Transcriptomic Analysis of Early Stages of Intestinal Regeneration in *Holothuria glaberrima*

David J. Quispe-Parra[1], Joshua Medina-Feliciano[1], Sebastián Cruz-González[1], Humberto Ortiz-Zuazaga[2], José E. García-Arrarás[1*]

[1]University of Puerto Rico, Biology Department, San Juan, 00925, Puerto Rico.

[2]University of Puerto Rico, Department of Computer Sciences, San Juan, 00925, Puerto Rico.

*jegarcia@hpcf.upr.edu

Abstract

Echinoderms comprise a group of animals with impressive regenerative capabilities. They can replace complex internal organs following injury or autotomy. In holothurians or sea cucumbers, cellular processes of intestinal regeneration have been extensively studied. The molecular machinery behind this faculty, however, remains to be understood. Here we assembled and annotated a de novo transcriptome using RNA-seq data consisting of regenerating and non-regenerating intestinal tissues from the sea cucumber *Holothuria glaberrima*. Comparisons of differential expression were made using the mesentery as a reference against 24 hour and 3 days regenerating intestine, revealing a large number of differentially expressed transcripts. Gene ontology and pathway enrichment analysis showed evidence of increasing transcriptional and translational activity. Further analysis of transcripts associated with transcription factors revealed diverse expression patterns with mechanisms involving developmental and cancer-related activity that could be related to the regenerative process. Our study demonstrates the broad and diversified gene expression profile during the early stages of the process using the mesentery as the focal point of intestinal regeneration. It also establishes the genes that are the most important candidates in the cellular processes that underlie regenerative responses.

Introduction

Regeneration is fundamental for renewal of cells and tissues in multicellular organisms, nevertheless the process of regenerating complex components, such as organs, is only possible in a few animal groups[1]. Understanding the limitations of this capability has become an important goal in improving health conditions related to tissue and organ functional maintenance[2]. However, discerning this process requires a clear understanding of the molecular mechanisms that drive the cellular functions responsible for regeneration.

Organisms with astonishing regenerative capacities are scattered among basal metazoan lineages and in various lophotrochozoan and deuterostome phyla. The latter encompasses a wide group of organisms that includes vertebrates along with tunicates, echinoderms, hemichordates, and cephalochordates. This phylogenetic placement makes highly

regenerative groups such as tunicates and echinoderms, attractive models for studying regeneration. Their potentially shared genetic mechanisms highlight their importance for biomedical studies[3,4].

Echinoderms comprise an important group of organisms with extraordinary regeneration capabilities. However, despite being among the most highly regenerative deuterostomes and their close relationship to vertebrates, the number of studies on their regenerative capacity is rather limited[5]. In recent years their use as regeneration model organisms has seen an upsurge with studies on spine regeneration in sea urchins [6], arm regeneration in crinoids, brittle stars and sea stars[7,8,9] and nervous system and viscera regeneration in sea cucumbers[10,11,12,13,14]. Sea cucumbers have been particularly attractive for intestinal regeneration studies because of their innate capacity of expelling their digestive tract in a process known as evisceration, which is then followed by its regeneration. Research relating regeneration of the whole digestive system is scarce among regenerating competent organisms, where in addition to sea cucumbers, other studies include planarians[15,16].

Efforts in understanding the regeneration of the holothurian digestive tract have provided important insights on the cellular processes involved in organogenesis. Experiments have identified the fundamental role of the cells in the mesentery during intestinal regeneration[17]. In normal, non-eviscerated animals, the mesentery extends along the interior body wall providing support to the intestine. Following evisceration, the torn edges at the distal part of the mesentery acquire a thick oval morphology forming the initial intestinal rudiment from the esophagus to the cloaca. This growth is formed by evisceration-activated signals that trigger dedifferentiation of peritoneocytes and muscle cells in the mesothelium of the mesentery. Myocyte dedifferentiation is associated with elimination of the contractile apparatus through condensation of myofilaments into membrane bound spindle-like structures (SLSs)[18]. The presence of these structures has been used to reveal a spatio-temporal gradient where, initially, dedifferentiation is localized near the free margin of the mesentery, while the rest of the mesentery remains mostly in a differentiated state. This pattern progresses until eventually different regions with increasing levels of de-differentiation can be observed in the mesentery. It has been hypothesized that dedifferentiated cells give rise to precursor cells that eventually divide and re-differentiate into cellular components of the regenerated organ[19].

In subsequent stages, mesothelial cells within the regenerating structure undergo an epithelial to mesenchymal transition (EMT) and ingress into the underlying connective tissue[20]. The increasing number of cells in the distal part of the mesentery together with the possible deposition of new extra cellular matrix (ECM), contribute to the rudiment growth. Therefore, about a week after evisceration, the intestinal rudiment can be clearly observed as a solid rod at the tip of the mesentery. In histological sections the rudiment appears as an elongated thickening of the mesentery with many mesenchymal cells in the inner connective tissue surrounded by a dedifferentiated coelomic epithelium. Within two weeks, the lumen forms and the reconstitution of mucosal epithelium is achieved by the intrusion

of tubular projections of the mucosal epithelium that migrate from anterior and posterior ends. It is important to highlight the role of the mesentery during all the intestinal regeneration process, since new cells originate from this structure as wells as the extracellular signals that modulate the whole process[17].

Although substantial advances have been achieved in dissecting the cellular mechanisms of intestinal regeneration, a similar analysis of the molecular basis is lagging. Initial methods to identify differentially expressed genes (DEGs) during intestinal regeneration consisted of differential screening (DS) and differential display (DD) to study specific genes that could be upregulated during regeneration[21,22,23]. Further studies with a broader approach targeted expressed sequence tags (ESTs) from cDNA libraries of regenerating intestines and the preparation of microarrays to reveal DEGs. In the last decade, these technologies have been used to uncover regeneration-associated genes in the transcriptomes of two other sea cucumbers, *Apostichopus japonicus*[24] and *Eupentacta fraudatrix*[25]. This has led to the identification of several DEGs linked with intestinal regeneration.

All previous intestinal comparative gene studies, however, share a major flaw. In their comparisons to determine the differences in gene expression profiles, they use the normal intestine as a baseline. This comparison is apt for late regenerative stages, but during the early events of regeneration, the regenerating rudiment shares more cells and tissues in common with the normal mesentery than with the normal intestine. Therefore, when using the normal intestine as a reference, genes expressed in the intestine luminal epithelium would appear as being under-expressed during the first week of regeneration since at that stage no lumen has formed. In contrast, genes that appeared as highly expressed in the early regenerate might be due to their preferential expression in the mesenterial tissues and not necessarily due to being activated by the regeneration process.

We have now applied RNA-seq technology to the analysis of intestinal regeneration in our model system, the sea cucumber *Holothuria glaberrima.* More importantly, to narrow down the genes involved in the early steps of the regenerative process we made two critical modifications in our experimental strategy. First, we focused on tissues at day 1 and day 3 of regeneration, stages that take place prior to, or concomitant with, the first observed cellular events. Second, regenerating tissues were compared to normal, uneviscerated mesentery. By using this approach, we obtained a real view concerning temporal differential gene expression during intestinal regeneration. Moreover, we were able to obtain a considerable amount of information regarding novel transcripts that appear to be activated during regeneration.

Results

Paired end reads obtained from the sequencing process had a satisfactory quality level after trimming. A phred score of 30 was obtained for all samples while N content was practically null in all samples as well (see Supplementary Fig. S1 online). De novo assembly was

generated using reads from this study and previous raw sequencing reads from the central nervous system of *H. glaberrima*[26]. The resulting transcriptome assessed with BUSCO had a 99.1% of complete core genes detected with 0.4% of fragmented genes and only 0.5% of missing core genes (see Supplementary Table S1 online).

Differential expression analysis

A total of 8460 differentially expressed transcripts (DETs) were found significant (adjusted P-value ≤ 0.05) at day 1 of regeneration (Fig. 1A). Upregulated transcripts (Log_2 Fold Change ≥ 2) comprised 3585, while downregulated (Log_2 Fold Change ≤ -2) were 4875. At day 3 of regeneration, the total number of transcripts with differential expression was 8216, of these 3663 were upregulated and 4553 downregulated (Fig. 1 B). When both 1 and 3 days regenerating intestines were compared to controls, a total of 3884 transcripts were differentially co-expressed on both days; 1723 were upregulated and 2161 downregulated (Fig. 1C).

To validate our data, we compared the expression of a series of annotated transcripts, to previous studies where the same genes had been analyzed using semiquantitative RT-PCR, RT-qPCR or *in situ* hybridization (Table 1). These transcripts correspond to annotated genes Myc, Orpin, Serum amyloid A, Mtf, WNT9, BMP, mortalin, β-catenin and survivin, and all of them shared the fact that their expression in the early intestinal rudiment (1-3 dpe) had been compared to that of the normal mesentery[21,27,28,29,30,31]. Our RNA-seq expression analysis for all these genes showed a similar expression pattern to their expression in prior studies. Thus, Myc, Orpin, Serum amyloid A, Mtf, WNT9, BMP, mortalin, were over-expressed while β-catenin and survivin showed no considerable difference between regenerating intestine and normal mesentery.

A wide range of annotated transcripts with differential expression were designated as DEGs. Here we provide a list of the top 10 upregulated and downregulated DEGs for days 1 and 3 as examples of some of the genes in our database (Tables 2-5). The ID column is the given name of the sequence after the clustering process to unify identical transcripts. Adjusted p-values were determined by a multiple test correction with the Benjamini and Hochberg method using p-values attained from the Wald test. Only the transcripts with adjusted p-value < 0.01 were chosen. At least half of upregulated DEGs were specific for each day, while others encoded proteins with similar descriptions between stages as in the case of actin, elongation factors, and cathepsin. Although annotated, some transcripts encode proteins not characterized in the databases as in the case of upregulated LOC592324 (accession: XM_030978239) and downregulated LOC575162 (accession: XM_030973217) at day 1. Downregulated DEGs shared the same characteristic with around half of their protein descriptions being specific for each stage and sharing proteins associated with transporters, aminopeptidases, lyases, homebox proteins and aquaporin channels. Additional comparison between day 1 and 3 of regeneration, gave interesting results concerning downregulated nervous system genes (see Supplementary Table S2 online).

Functional classification of DEGs

GO terms obtained from DEGs with a p-value less than 0.05 and considerable relation between the number of genes associated with the term (Gene Ratio) are shown in Figure 2. Most terms obtained from upregulated genes at both day 1 and 3, correlate with processes involved in transcription of DNA information into RNA as well as the translation of mRNA into proteins. Enriched terms of biological process (BP) include various protein transport mechanisms; cellular component (CC) terms are characterized for processes involving translation activity; molecular function (MF) terms comprise protein transport, translation, and RNA binding. Downregulated terms showed a wider variety of associations. In general, terms associated with signaling systems, cell adhesion, metabolic processes and the immune system were found. Similar terms were shown for CC, being the membrane components an enriched factor for downregulated genes. MF terms associated with downregulated genes in both days represented mostly cell signaling processes.

Pathways associated with intestinal regeneration were classified according to upregulated and downregulated genes for each day. Only resulting pathways with p-value < 0.05 were depicted in Figure 3. Enriched pathways for upregulated genes in day 1 and 3 were mostly associated with protein processing and transcriptional activity, being the pathway involved in proteasome activity the most enriched. Downregulated genes correlated mainly with various types of metabolic activity. Furthermore, enzymatic activity associated with peroxisomes were found to be enriched for both days. Intriguingly, phagosome activity was shown in pathways of upregulated genes for day 1, as wells as the downregulated genes of the same day.

Differentially Expressed Transcription Factors (DETFs)

One of the first cellular events associated with regeneration is the dedifferentiation of the mesentery cells. Such dedifferentiation is usually associated with cellular re-programming and the activation or inactivation of specific transcription factors (TFs). Thus, we identified the annotated TFs with expression changes (see Supplementary Table S3 online). The heatmap in Figure 4 shows DETF with an absolute value of Log_2 Fold Change (Log2FC) > 1 and an adjusted p-value < 0.05. Upregulation at day 1 was shown by the majority of the TFs in the heatmap, while a minority was downregulated at the same stage. The results show five main profile patterns. Group 1 (grey) corresponds to factors with moderate-to-low expression in normal tissues that increase during regeneration with a peak at day 1. Group 2 (light blue) corresponds to factors with high expression in normal tissues that decrease slightly in day 1 and 3. Group 3 (green) corresponds to factors with regular expression in normal mesentery that increase during the first three days of regeneration. Group 4 (yellow) correspond to factors that are moderately expressed in normal tissue and decrease their expression during regeneration. Finally, Group 5 (orange) includes factors that are weakly expressed in normal mesentery but increase their expression during regeneration.

Discussion

In the present study, using RNA-seq, we have constructed a large-scale differential gene expression profile of regenerating intestine in *H. glaberrima*. We validated our results by comparing the expression of genes found in our database with RT-PCR or in situ analyses where the expression of the same genes was done using the normal mesentery for comparison. Results showed an identical expression pattern in the early, 1- & 3-days post evisceration (dpe) stages. Among the genes showing parallel upregulated expression between RNA-seq and previous methods, we found the putative calcium-binding protein Orpin[28] and genes encoding proteins involved in cell differentiation and proliferation as in the case of Myc and Mtf[27,29]. Similarly, β-catenin showed no considerable change in expression in early stages in previous studies using RT-qPCR, and no changes were observed in our RNA-Seq results[27]. Other methods to detect gene expression were also found to give consistent results with our data. For instance, northern blot analysis of Serum amyloid A expression showed upregulation during day 3 of regeneration[21]. We found an increase in the expression of its mRNA at the same stage. Furthermore, *in situ* hybridization studies, although not quantitative, do show differences in mRNA expression that coincide with our data. In this case, WNT9, BMP, mortalin showed considerable increased labeling, in the early regenerate while survivin appeared to have scarce and dispersed labeling in the regenerating mesenteric region[30,31]. These results are in accordance with our RNA-seq findings, thus reassuring the reliability of the analysis obtained in our experimental approach to study early stages of intestinal regeneration in *H. glaberrima*.

The high number of DETs for 1 & 3 dpe (>8000), contrast with a previous study of intestinal regeneration in *A. japonicus* where less than 4000 DETs were found[24]. Notwithstanding, these differences can be attributed to the significance criteria established for each study. In *A. japonicus,* an adjusted p-value of < 0.001 was established as threshold for significance, while our study used a value of < 0.05. In fact, using the same significance criteria (see Supplementary Fig. S2 online), the number of DETs is similar (3587) to those reported in the *A. japonicus* study. Our results are also comparable with a study done in *E. fraudatrix* applying equivalent significance benchmarks (adjusted p-value < 0.05) documenting 17,227 DETs in regenerating intestine relative to intact gut[25].

Our interest, however, lies beyond simply quantifying the number of DETs but targets the expression trends during early stages of intestinal regeneration to determine genes involved in cellular events such as dedifferentiation, proliferation and migration. It is in this analysis where the major discrepancies are observed between this study and those of others.

Comparison of top DEGs at day 3 of regeneration between *A. japonicus*[24] and our data revealed no parallelism. All genes in up-regulated and down-regulated lists were different. Additional disparity was found in *E. fraudatrix* study, where most of their DETFs[25] were not found in our study, or if present, their differential expression showed no correlation. Examples of this expression discrepancy: EGR1, TCF24 and SNAI2, upregulated in *E. fraudatrix*, downregulated in *H. glaberrima*. There are even differences with previous

studies using the same species from our own group[32]. Diverging expressions were found mostly with ECM-related genes such as collagen alpha-1, laminin alpha 1, stromelysin-3 (MMP II) and developmental related Hox 9. In all cases, these genes were upregulated in the microarray study, whereas they were downregulated in our RNA-seq. As mentioned earlier we believe the differences between our results and previous studies stem from what each study used as a point of comparison to determine the gene expression profile. In all previous studies, the adult non-eviscerated intestine was used as the reference point to determine differential expression. However, as explained earlier, the ratio of the area of connective tissue to that of the mesothelium in the early intestinal rudiment is much smaller than in the adult intestine. Moreover, the adult intestine has a prominent mucosal layer that does not appear in the intestinal rudiment until the second week of regeneration. Therefore, the relative changes in cell/tissue distribution between the early regenerate and the fully formed organ are a clear cause for a skew in the resulting gene expression profile. Thus, the expression differences in previous studies might reflect differences in cell or tissue gene expression rather than differences in regeneration stages. On the other hand, our study takes these facts into account and focuses on a comparison with the normal mesentery which is a more trustworthy point of comparison because of similar distribution of cells and tissues.

Both functional and signal pathway analyses point out to certain activities that are central to the regeneration process. Overall upregulated transcripts were mostly involved with transcriptional processes, translation, and protein transport. This was expected as the regenerative process requires expression of abundant components for reconstitution of lost tissues. Post-translational protein processing and proteasome related pathways were also represented in the results. Both mechanisms appear to be evidently important for cell renewal specially the former which is needed for enzymatic modifications following synthesis. Proteasome-related transcripts have been previously observed to be upregulated during intestinal regeneration[33] and activity of the proteasome complex has been classified as indispensable for stem cell maintenance[34]. Proteasome activity is also important in cell signaling as the degradation of particular proteins can cause large shifts in cell processes[35]. Metabolic mechanisms were mostly reflected for downregulated transcripts, implying a halt in cell communication activity, catabolic and biosynthetic processes to concentrate biologic resources towards regeneration. Some pathways involved in metabolic activity are associated with decreased oxygen consumption rate (OCR) following evisceration[36]. This trend has been observed in induced pluripotent stem cells (iPSCs) that show inhibition of oxidative metabolism[37]. Regarding other downregulated GO terms: cell adhesion, included proteins known as members of broad families with diverse expression patterns such as cadherins[38]. A possible reduction in cell adhesion is consonant with mesenterial cell migration toward the forming blastema and with EMT events. Immune system at day 1 appears surprising given the fact that this activity is believed to be activated during regeneration and previous studies have shown representation of inflammatory response terms at day 3[39]. However, there are no studies analyzing expression pattern as early as day 1, which could indicate a late immune response in the beginning of the process. There is also evidence of positive regeneration response by suppression of the immune response in tadpole tail regeneration[40], which highlights the puzzling role of this complex network in

regeneration. The term peroxisome pathway in this downregulated group, suggests a decreased activity towards removal of reactive oxygen species (ROS). The importance of ROS has been documented in various regeneration model organisms for its positive effect during regeneration[41,42,43]. Finally, and worth highlighting, phagosome related pathway was represented in upregulated and downregulated transcripts for day 3. This tendency results from similar DEGs of actin and tubulin involved in phagosome motility and structure[44,45,46]. However, transcripts grouped in both expression trends were different, e.g. tubulin alpha-1A in downregulated and tubulin beta-4B in upregulated. Additionally, there were a greater number of transcripts promoting the phagocytic process in upregulated transcripts such as calnexin and calreticulin, two pro-phagocytic proteins with chaperone activity in endoplasmic reticulum[47].

Among the DEGs between normal mesentery and the early stages of intestinal regeneration, we have focused on TFs for obvious reasons: These are the genes that play central roles in most cellular processes by activating or inhibiting the transcription of many downstream genes involved in the process.

The DETFs can be classified depending on the genes that are being regulated and the effect they produce when expressed. Thus, it is not surprising that many of the DETFs regulate the transcription process itself. For example, TAF1A, MYBBP1A and PWP1 are involved in the assembly of RNA polymerase I preinitiation complex, processing of ribosomal RNA and ribosome assembly, respectively[48,49,50]. Similarly, EEF1A1 has been recognized as a translation elongation factor[51]. Interestingly all these genes appear to increase their expression at day 1 and are still upregulated, but at a lower level at day 3. This suggests that the injury triggers an early response to establish the transcription machinery necessary for the subsequent regenerative processes.

Far more interesting are those TF associated with cellular identity. In this respect the differential expression observed with MYC calls for particular attention. MYC is one of the Yamanaka factors used to induce pluripotency[52]. Moreover, previous studies from our lab have shown that it is overexpressed during both intestinal and radial nerve regeneration, two processes where cells undergo dedifferentiation[53,54]. RNAi performed in the radial nerve shows that downregulation of MYC decreases the dedifferentiation of the radial glia[27]. Thus, our study fully supports the view that MYC is important in the early regenerative process suggesting that it is important for the dedifferentiation of the mesenterial cells.

One of the most interesting findings is that most DETFs share one common characteristic: they have been associated with developmental processes involving mesodermal derivatives. Many of them have been shown to activate or inhibit processes leading to muscle, cartilage, or bone formation[55,56,57]. In many cases they are involved in keeping cells in an immature state or inhibiting their differentiation[58,59]. Furthermore, some are directly associated with the formation of digestive system organs, such as the liver, pancreas, cloaca, stomach and even with enterocyte identity[56,60,61,62]. Among these factors we found various transcripts associated with developmental-related homeobox (Hox) family of genes. Our results show

activation of MKX, HMX, TAP26 and DLX5 at day 1, while inhibition of HLX, EMX1 and CDX1 is observed for the same day. This contrasting expression trend is particular of Hox genes, which are characterized by their dynamic expression pattern as shown in sea urchin development[63]. Although Hox TFs appear to be involved in countless mechanisms, functions associated with MKX, HLX and CDX1 are worth mentioning. MKX has been shown to have a role as a regulator of collagen production[55], a structural protein previously shown to have important implications during intestine regeneration[56]. As for HLX and CDX1, their implication in enteric nervous system and intestine development[60,61] makes them interesting study targets. This activity could also be associated with downregulation of neurogenic associated transcripts observed when comparing day 3 vs 1 (see Supplementary Table S2 online). Considering previous studies of partial injury of the nervous system in regenerating mesentery[64], a stall in intestinal neuronal development can be hypothesized during dedifferentiation at early regeneration stages.

Other upregulated development-related factors include MYBBP1A, TEF, and MYNN. The first two have important roles in embryonic development[65,66], whereas MYNN has a localized expression in muscle cells during the same period[57]. On the other hand, downregulated transcripts identified as NRL/MAF (neural retina leucine zipper, Maf type), EGR1 and FOXA, have all been associated with differentiation mechanisms. The former has been shown to have an essential role in rod photoreceptor development[67]; EGR1 has a vast range of activities including differentiation and proliferation, most importantly its association with other factors that regulate cell growth[68]; FOXA has a role in competence establishment of the foregut endoderm and induction of liver specific genes for hepatogenesis[62]. Finally, it is worth mentioning that two transcripts annotated as KLF13 and ZNF260 were found, respectively upregulated and downregulated at day 1. Both belong to the Krüppel family of transcriptional regulators involved in modulation of numerous developmental features[58,59]. Downregulation of most of these transcripts appears reasonable since their differentiation role is contrary to the dedifferentiation processes taking place at the onset of regeneration.

Lastly are the DETFs associated with oncogenesis. SOX4 and LMO2 show upregulation after day 1 while HNF4G is downregulated in the same period. These cancer-related TFs, were recognized among the top considerable transcripts with differential expression. Similarities between regeneration and cancer-related events have been studied for their involvement in dedifferentiating mechanisms during cellular reprogramming[69]. This compelling process of returning to an undifferentiated state during regeneration, confers high proliferative and migratory capacities to cells, without the loss of cell growth control distinctive of cancer[70]. In the case of SOX4 and LMO2, their role has been established as regulators with aberrant expression leading to an oncogenic state[71,72]. As for HNF4G, its upregulation has been correlated as a prognostic marker for cancer development by promoting cell proliferation[73], which is interesting considering the opposite expression trend in our results. Other transcripts related to THAP6 and BTF3L4 that show upregulation at day 1, have been associated with an apoptotic regulatory role[74,75]. This

contrasting expression portrays regulatory differences of cancer related mechanisms being activated for dedifferentiation in early stages.

Diverse TF activity indicates the abundant network of mechanisms taking place during intestinal regeneration. Various TF were found to be directly or indirectly involved with regenerative processes. Although their primary role is rather aimed at developmental purposes in most cases, there are previously studied TF such as MYC whose DE can be proposed as landmarks of regeneration.

Conclusion

This is the first study with a mesentery directed approach for improving specificity in the characterization of differential gene expression in regenerating tissues. Furthermore, it is also the first one in sea cucumbers for analysis of differential expression during intestinal regeneration at a stage as early as 24 hours after the organism has expelled its internal organs. An expected high number of DETs were found using high-throughput sequencing methods and numerous functional processes associated with regeneration. What is more appealing, TF activity correlated with regenerative machinery was diverse in expression trends that not necessarily belonged to the same group of processes. Further studies can be considered to unravel the regulatory networks associated with TF activity and connect them with distinctive cellular processes.

Materials and Methods

Animals and treatment

Sea cucumbers were collected from the northeastern rocky coast of Puerto Rico[76]. Evisceration was carried out by an intracoelomic injection of 0.35M KCl. Subsequently, the organisms were left in salt-water aquaria for 1 and 3 dpe to undergo regeneration. Uneviscerated animals kept under the same conditions and for the same lengths of time as the eviscerated animals were used as controls. Prior to dissection, animals were anesthetized by immersion in ice-cold water for 45 minutes. A dorsal incision was made to expose the internal cavity of the organism and allow dissection of the mesentery separated from the body wall and intestine. In the case of non-eviscerated organisms, the intestine was separated from the mesentery that remained attached to the body wall and removed using small surgical scissors. The mesentery was obtained through out its length from the anterior end of the animal (next to the esophagus) to the posterior end (next to the cloaca). Dissected tissues were placed in RNAlater (Sigma, USA) solution and stored at 4ºC until the RNA extraction was performed.

RNA extraction

RNA extraction was done with a combination of the method established by Chomczynski (1993) using Tri-reagent® (N.93289, Sigma, USA) and the RNAeasy mini kit (Qiagen,

Germany). Three extractions were made for each stage (normal, day 1 and day 3) and each extraction consisted of pooled tissue samples from three organisms at the same stage. The concentration and quality of the RNA was determined by using the 2100 Bioanalyzer (Agilent Technologies, USA). Samples with a concentration greater than 200 ng/μL and an RNA Integrity Number (RIN) value of ≥ 8 were used for sequencing.

Sequencing of extracted RNA

RNA isolated from the mesentery of regenerating and non-regenerating sea cucumbers was sequenced at the Sequencing and Genotyping Facility of the University of Puerto Rico. Sequencing libraries were prepared using the TruSeq Stranded mRNA Library Prep Kit (Illumina, USA). Paired-end sequencing (2□×□150) was performed with the Illumina NextSeq® 500 system.

Transcriptome assembly

Raw reads were uploaded to the High-Performance Computing Facility of the University of Puerto Rico. Preprocessing steps of quality, trimming and filtering were done using FastQC[77] and Trimmomatic[78] tool packages to remove sequencing errors and adapters. A kmer trimming and digital normalization was applied previous to the assembly process using the khmer package[79]. A de novo transcriptome assembly was carried out using Trinity[80].

Differential gene expression

The differential expression analysis was carried out with Salmon[81] using the quality filtered data of RNA-seq and the assembled transcriptome generated with Trinity as reference. Count files generated with Salmon were grouped according to the type of sample (control, 24 hours and 3 days). Equivalence classes were also obtained to hierarchically cluster contigs by read similarity and expression with Corset[82]. Lastly, DESeq2 tool package[83] was used to extract differentially expressed cluster of contigs from the files generated with Salmon.

Functional annotation

The annotation pipeline Dammit[84] was used for annotating the transcriptome. Protein databases Pfam-A[85], Rfam[86], OrthoDB[87] and uniref90[88] were used for the annotation. The protein sequences from the genome of the sea urchin *Strongylocentrotus purpuratus* (Accession: GCA_000002235.4) were used as reference for the annotation process. Annotation completeness was evaluated with BUSCO[89].

Gene ontology and enrichment analysis

In order to have a broader understanding of the molecular processes that take place at different stages of gut regeneration we mapped Entrez Gene IDs of our transcripts using DAVID gene functional classification[90] tool to obtain a gene ontology (GO) analysis. Only DEGs with significant p-value (<0.05) and absolute expression level higher than 2 (Log2FC) were associated with GO terms. Results of ontology terms were summarized

with Revigo[91]. Pathway information was obtained by mapping KEGG ids from *S. purpuratus* datasets to *H. glaberrima* Entrez Gene IDs using kegga function from the Bioconductor package limma v3.11[92]. Gene ratio was manually calculated for each individual pathway using the number of DEGs for a term divided by the total number for that term. All transcripts used in the GO and pathway enrichment analysis were filtered based on their p value (<0.05).

Transcription factor analysis

Sequences annotated as transcription factors (TFs) were selected for expression analysis during the three stages of the regeneration process: non-regeneration stage, day 1 and day 3 of regeneration. A heatmap was generated with gplots package using the TF that had an absolute Log2FC higher than 2[93].

Data availability

The raw RNA-seq data can be found at the SRA NCBI Database under the BioProject accession PRJNA660762. Custom scripts and data for differential gene expression are available at https://github.com/davidjquispe/DGE_Analysis.

References

1. Alibardi, L. Review: Limb regeneration in humans: Dream or reality? *Annals of Anatomy - Anatomischer Anzeiger* **217**, 1–6 (2018).

2. Tsukamoto, A. *et al.* Challenging Regeneration to Transform Medicine. *STEM CELLS Translational Medicine* **5**, 1–7 (2016).

3. Bely, A. E. & Nyberg, K. G. Evolution of animal regeneration: re-emergence of a field. *Trends in Ecology & Evolution* **25**, 161–170 (2010).

4. Lai, A. G. & Aboobaker, A. A. EvoRegen in animals: Time to uncover deep conservation or convergence of adult stem cell evolution and regenerative processes. *Developmental Biology* **433**, 118–131 (2018).

5. García-Arrarás, J. E. & Dolmatov, I. Yu. Echinoderms: potential model systems for studies on muscle regeneration. *Curr Pharm Des* **16**, 942–955 (2010).

6. Reinardy, H. C., Emerson, C. E., Manley, J. M. & Bodnar, A. G. Tissue Regeneration and Biomineralization in Sea Urchins: Role of Notch Signaling and Presence of Stem Cell Markers. *PLoS One* **10**, (2015).

7. Hernroth, B. *et al.* Possibility of mixed progenitor cells in sea star arm regeneration. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* **314B**, 457–468 (2010).

8. Ferrario, C. *et al.* Fundamental aspects of arm repair phase in two echinoderm models. *Dev. Biol.* **433**, 297–309 (2018).

9. Mashanov, V. *et al.* Active Notch signaling is required for arm regeneration in a brittle star. *PLOS ONE* **15**, e0232981 (2020).

10. Shukalyuk, A. I. & Dolmatov, I. Yu. Regeneration of the Digestive Tube in the Holothurian Apostichopus japonicusafter Evisceration. *Russian Journal of Marine Biology* **27**, 168–173 (2001).

11. MASHANOV, V. S. & DOLMATOV, I. Y. Regeneration of digestive tract in the pentactulae of the far-eastern holothurian Eupentacta fraudatrix (Holothuroidea, Dendrochirota). *Invertebrate Reproduction & Development* **39**, 143–151 (2001).

12. San Miguel-Ruiz, J. E., Maldonado-Soto, A. R. & García-Arrarás, J. E. Regeneration of the radial nerve cord in the sea cucumber Holothuria glaberrima. *BMC Dev Biol* **9**, 3 (2009).

13. Mashanov, V. S. & García-Arrarás, J. E. Gut Regeneration in Holothurians: A Snapshot of Recent Developments. *The Biological Bulletin* (2016) doi:10.1086/BBLv221n1p93.

14. García-Arrarás, J. E. *et al.* Cell dedifferentiation and epithelial to mesenchymal transitions during intestinal regeneration in H. glaberrima. *BMC Developmental Biology* **11**, 61 (2011).

15. Forsthoefel, D. J., Park, A. E. & Newmark, P. A. Stem cell-based growth, regeneration, and remodeling of the planarian intestine. *Developmental Biology* **356**, 445–459 (2011).

16. Forsthoefel, D. J., Cejda, N. I., Khan, U. W. & Newmark, P. A. Cell-type diversity and regionalized gene expression in the planarian intestine. *eLife* **9**, e52613 (2020).

17. García-Arrarás, J. E., Bello, S. A. & Malavez, S. The mesentery as the epicenter for intestinal regeneration. *Seminars in Cell & Developmental Biology* **92**, 45–54 (2019).

18. Candelaria, A. G., Murray, G., File, S. K. & García-Arrarás, J. E. Contribution of mesenterial muscle dedifferentiation to intestine regeneration in the sea cucumber Holothuria glaberrima. *Cell Tissue Res* **325**, 55–65 (2006).

19. García-Arrarás, J. E. Dedifferentiation as a cell source for organ regeneration. in *Regenerative Engineering and Developmental Biology* (ed. Gardiner, D. M.) 373–394 (CRC Press, 2017). doi:10.1201/9781315120188-20.

20. García-Arrarás, J. E. *et al.* Cell dedifferentiation and epithelial to mesenchymal transitions during intestinal regeneration in H. glaberrima. *BMC Dev. Biol.* **11**, 61 (2011).

21. Santiago, P., Roig□López, J. L., Santiago, C. & García□Arrarás, J. E. Serum amyloid A protein in an echinoderm: Its primary structure and expression during intestinal regeneration in the sea cucumber Holothuria glaberrima. *Journal of Experimental Zoology* **288**, 335–344 (2000).

22. Méndez, A. T., Roig-López, J. L., Santiago, P., Santiago, C. & García-Arrarás, J. E. Identification of Hox Gene Sequences in the Sea Cucumber Holothuria glaberrima Selenka (Holothuroidea: Echinodermata). *Mar. Biotechnol.* **2**, 231–240 (2000).

23. Suárez-Castillo, E. C., Medina-Ortíz, W. E., Roig-López, J. L. & García-Arrarás, J. E. Ependymin, a gene involved in regeneration and neuroplasticity in vertebrates, is overexpressed during regeneration in the echinoderm Holothuria glaberrima. *Gene* **334**, 133–143 (2004).

24. Sun, L., Yang, H., Chen, M., Ma, D. & Lin, C. RNA-Seq Reveals Dynamic Changes of Gene Expression in Key Stages of Intestine Regeneration in the Sea Cucumber Apostichopus japonicas. *PLOS ONE* **8**, e69441 (2013).

25. Boyko, A. V., Girich, A. S., Tkacheva, E. S. & Dolmatov, I. Y. The Eupentacta fraudatrix transcriptome provides insights into regulation of cell transdifferentiation. *Scientific Reports* **10**, 1522 (2020).

26. Mashanov, V. S., Zueva, O. R. & García-Arrarás, J. E. Transcriptomic changes during regeneration of the central nervous system in an echinoderm. *BMC Genomics* **15**, 357 (2014).

27. Delgado, M. A. *Functional Role of B-catenin and Myc as Active Players in the Canonical Wnt Signaling Pathway During Intestinal Regeneration of the Sea Cucumber Holothuria Glaberrima*. (Universidad de Puerto Rico, 2019).

28. Soto-Acabá, A., Ortíz-Pineda, P. A. & García-Arrarás, J. E. Characterization of two novel EF-hand proteins identifies a clade of putative Ca2+-binding protein specific to the Ambulacraria. *bioRxiv* 2020.05.22.110411 (2020) doi:10.1101/2020.05.22.110411.

29. Hernández☐Pasos, J., Valentín☐Tirado, G. & García☐Arrarás, J. E. Melanotransferrin: New Homolog Genes and Their Differential Expression during Intestinal Regeneration in the Sea Cucumber Holothuria glaberrima. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* **328**, 259–274 (2017).

30. Mashanov, V. S., Zueva, O. R. & Garcia-Arraras, J. E. Expression of Wnt9, TCTP, and Bmp1/Tll in sea cucumber visceral regeneration. *Gene Expression Patterns* **12**, 24–35 (2012).

31. Mashanov, V. S., Zueva, O. R., Rojas-Catagena, C. & Garcia-Arraras, J. E. Visceral regeneration in a sea cucumber involves extensive expression of survivin and mortalin homologs in the mesothelium. *BMC Developmental Biology* **10**, 117 (2010).

32. Ortiz-Pineda, P. A. *et al.* Gene expression profiling of intestinal regeneration in the sea cucumber. *BMC Genomics* **10**, 262 (2009).

33. Pasten, C., Ortiz-Pineda, P. A. & E.García-Arrarás, J. Ubiquitin-proteasome system components are up-regulated during intestinal regeneration. *Genesis* **50**, 350–365 (2012).

34. Kitajima, Y. *et al.* The Ubiquitin-Proteasome System Is Indispensable for the Maintenance of Muscle Stem Cells. *Stem Cell Reports* **11**, 1523–1538 (2018).

35. Willis Monte S., Townley-Tilson W. H. Davin, Kang Eunice Y., Homeister Jonathon W. & Patterson Cam. Sent to Destroy: the ubiquitin proteasome system regulates cell signaling and protein quality control in cardiovascular development and disease. *Circulation Research* **106**, 463–478 (2010).

36. Zang, Y., Tian, X., Dong, S. & Dong, Y. Growth, metabolism and immune responses to evisceration and the regeneration of viscera in sea cucumber, Apostichopus japonicus. *Aquaculture* **358–359**, 50–60 (2012).

37. *Regenerative Biology of the Eye*. (Springer New York, 2014). doi:10.1007/978-1-4939-0787-8.

38. Hollnagel, A., Grund, C., Franke, W. W. & Arnold, H.-H. The Cell Adhesion Molecule M-Cadherin Is Not Essential for Muscle Development and Regeneration. *Molecular and Cellular Biology* **22**, 4760–4770 (2002).

39. Rojas-Cartagena, C. *et al.* Distinct profiles of expressed sequence tags during intestinal regeneration in the sea cucumber Holothuria glaberrima. *Physiological Genomics* **31**, 203–215 (2007).

40. Fukazawa, T., Naora, Y., Kunieda, T. & Kubo, T. Suppression of the immune response potentiates tadpole tail regeneration during the refractory period. *Development* **136**, 2323–2327 (2009).

41. Love, N. R. *et al.* Amputation-induced reactive oxygen species are required for successful Xenopus tadpole tail regeneration. *Nature Cell Biology* **15**, 222–228 (2013).

42. Pirotte, N. *et al.* Reactive Oxygen Species in Planarian Regeneration: An Upstream Necessity for Correct Patterning and Brain Formation. *Oxidative Medicine and Cellular Longevity* **2015**, 1–19 (2015).

43. Zhang, Q. *et al.* Reactive oxygen species generated from skeletal muscles are required for gecko tail regeneration. *Scientific Reports* **6**, 20752 (2016).

44. Zhang, F., Southwick, F. S. & Purich, D. L. Actin-based phagosome motility. *Cell Motil. Cytoskeleton* **53**, 81–88 (2002).

45. Jiang, M. *et al.* Microtubule motors transport phagosomes in the RPE, and lack of KLC1 leads to AMD-like pathogenesis. *J Cell Biol* **210**, 595–611 (2015).

46. Keller, S., Berghoff, K. & Kress, H. Phagosomal transport depends strongly on phagosome size. *Scientific Reports* **7**, 17068 (2017).

47. Müller-Taubenberger, A. *et al.* Calreticulin and calnexin in the endoplasmic reticulum are important for phagocytosis. *EMBO J* **20**, 6772–6782 (2001).

48. Friedrich, J. K., Panov, K. I., Cabart, P., Russell, J. & Zomerdijk, J. C. B. M. TBP-TAF Complex SL1 Directs RNA Polymerase I Pre-initiation Complex Formation and Stabilizes Upstream Binding Factor at the rDNA Promoter. *J Biol Chem* **280**, (2005).

49. Hochstatter, J. *et al.* Myb-binding Protein 1a (Mybbp1a) Regulates Levels and Processing of Pre-ribosomal RNA. *J. Biol. Chem.* **287**, 24365–24377 (2012).

50. Talkish, J., Campbell, I. W., Sahasranaman, A., Jakovljevic, J. & Woolford, J. L. Ribosome Assembly Factors Pwp1 and Nop12 Are Important for Folding of 5.8S rRNA during Ribosome Biogenesis in Saccharomyces cerevisiae. *Molecular and Cellular Biology* **34**, 1863–1877 (2014).

51. Vera, M. *et al.* The translation elongation factor eEF1A1 couples transcription to translation during heat shock response. *eLife* **3**, e03164 (2014).

52. Takahashi, K. & Yamanaka, S. Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors. *Cell* **126**, 663–676 (2006).

53. Mashanov, V. S., Zueva, O. R. & García-Arrarás, J. E. Expression of pluripotency factors in echinoderm regeneration. *Cell Tissue Res* **359**, 521–536 (2015).

54. Mashanov, V. S., Zueva, O. R. & García-Arrarás, J. E. Myc regulates programmed cell death and radial glia dedifferentiation after neural injury in an echinoderm. *BMC Developmental Biology* **15**, 24 (2015).

55. Liu, W. *et al.* The atypical homeodomain transcription factor Mohawk controls tendon morphogenesis. *Mol. Cell. Biol.* **30**, 4797–4807 (2010).

56. San Miguel-Ruiz, J. E. & García-Arrarás, J. E. Common cellular events occur during wound healing and organ regeneration in the sea cucumber Holothuria glaberrima. *BMC Developmental Biology* **7**, 115 (2007).

57. Cifuentes-Diaz, C. *et al.* Neuromuscular expression of the BTB/POZ and zinc finger protein myoneurin. *Muscle Nerve* **29**, 59–65 (2004).

58. Pearson, R., Fleetwood, J., Eaton, S., Crossley, M. & Bao, S. Krüppel-like transcription factors: A functional family. *The International Journal of Biochemistry & Cell Biology* **40**, 1996–2001 (2008).

59. Debrus, S. *et al.* The zinc finger-only protein Zfp260 is a novel cardiac regulator and a nuclear effector of alpha1-adrenergic signaling. *Mol. Cell. Biol.* **25**, 8669–8682 (2005).

60. Bates, M. D., Dunagan, D. T., Welch, L. C., Kaul, A. & Harvey, R. P. The Hlx homeobox transcription factor is required early in enteric nervous system development. *BMC Dev Biol* **6**, 33 (2006).

61. Silberg, D. G., Swain, G. P., Suh, E. R. & Traber, P. G. Cdx1 and Cdx2 expression during intestinal development. *Gastroenterology* **119**, 961–971 (2000).

62. Lee, C. S., Friedman, J. R., Fulmer, J. T. & Kaestner, K. H. The initiation of liver development is dependent on Foxa transcription factors. *Nature* **435**, 944–947 (2005).

63. Arenas-Mena, C., Cameron, A. R. & Davidson, E. H. Spatial expression of Hox cluster genes in the ontogeny of a sea urchin. *Development* **127**, 4631–4643 (2000).

64. Nieves-Ríos, C., Alvarez-Falcón, S., Malavez, S., Rodriguez-Otero, J. & García-Arrarás, J. E. The nervous system component of the mesentery of the sea cucumber Holothuria glaberrima in normal and regenerating animals. *Cell Tissue Res* **380**, 67–77 (2020).

65. Mori, S. *et al.* Myb-Binding Protein 1A (MYBBP1A) Is Essential for Early Embryonic Development, Controls Cell Cycle and Mitosis, and Acts as a Tumor Suppressor. *PLOS ONE* **7**, e39723 (2012).

66. Drolet, D. W. *et al.* TEF, a transcription factor expressed specifically in the anterior pituitary during embryogenesis, defines a new class of leucine zipper proteins. *Genes Dev.* **5**, 1739–1753 (1991).

67. Jayaram, H., Khaw, P.-T., MacLaren, R. E. & Limb, G. A. Focus on Molecules: Neural retina leucine zipper (NRL). *Experimental Eye Research* **104**, 99–100 (2012).

68. Ponti, D. *et al.* The Transcription Factor EGR1 Localizes to the Nucleolus and Is Linked to Suppression of Ribosomal Precursor Synthesis. *PLOS ONE* **9**, e96037 (2014).

69. Mosteiro, L. *et al.* Tissue damage and senescence provide critical signals for cellular reprogramming in vivo. *Science* **354**, (2016).

70. Oviedo, N. J. & Beane, W. S. Regeneration: The origin of cancer or a possible cure? *Seminars in Cell & Developmental Biology* **20**, 557–564 (2009).

71. Foronda, M. *et al.* Sox4 Links Tumor Suppression to Accelerated Aging in Mice by Modulating Stem Cell Activation. *Cell Reports* **8**, 487–500 (2014).

72. Meng Shu *et al.* LIM Domain Only 2 Regulates Endothelial Proliferation, Angiogenesis, and Tissue Regeneration. *Journal of the American Heart Association* **5**, e004117.

73. Wang, J. *et al.* Expression of HNF4G and its potential functions in lung cancer. *Oncotarget* **9**, 18018–18028 (2017).

74. Cm, B., Ca, B. & Jr, P. G. Structure of the C-terminal heme-binding domain of THAP domain containing protein 4 from Homo sapiens. *Proteins* **79**, 1337–1341 (2011).

75. Brockstedt, E., Otto, A., Rickers, A., Bommert, K. & Wittmann-Liebold, B. Preparative high-resolution two-dimensional electrophoresis enables the identification of RNA polymerase B

transcription factor 3 as an apoptosis-associated protein in the human BL60-2 Burkitt lymphoma cell line. *J. Protein Chem.* **18**, 225–231 (1999).

76. García☐Arrarás, J. E. *et al.* Cellular mechanisms of intestine regeneration in the sea cucumber, Holothuria glaberrima Selenka (Holothuroidea:Echinodermata). *Journal of Experimental Zoology* **281**, 288–304 (1998).

77. Andrews, S. FastQC: A Quality Control tool for High Throughput Sequence Data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (2010).

78. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

79. Crusoe, M. R. *et al.* The khmer software package: enabling efficient nucleotide sequence analysis. *F1000Res* **4**, (2015).

80. Grabherr, M. G. *et al.* Trinity: reconstructing a full-length transcriptome without a genome from RNA-Seq data. *Nat Biotechnol* **29**, 644–652 (2011).

81. Patro, R., Duggal, G. & Kingsford, C. Salmon: Accurate, Versatile and Ultrafast Quantification from RNA-seq Data using Lightweight-Alignment. *bioRxiv* 021592 (2015) doi:10.1101/021592.

82. Davidson, N. M. & Oshlack, A. Corset: enabling differential gene expression analysis for de novoassembled transcriptomes. *Genome Biology* **15**, 410 (2014).

83. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

84. Scott, C. dammit: an open and accessible de novo transcriptome annotator. (2016).

85. Finn, R. D. *et al.* Pfam: the protein families database. *Nucleic Acids Res* **42**, D222–D230 (2014).

86. Griffiths-Jones, S., Bateman, A., Marshall, M., Khanna, A. & Eddy, S. R. Rfam: an RNA family database. *Nucleic Acids Res* **31**, 439–441 (2003).

87. Waterhouse, R., Tegenfeldt, F., Li, J., M Zdobnov, E. & V Kriventseva, E. OrthoDB: A hierarchical catalog of animal, fungal and bacterial orthologs. *Nucleic acids research* **41**, (2012).

88. Suzek, B. E., Huang, H., McGarvey, P., Mazumder, R. & Wu, C. H. UniRef: comprehensive and non-redundant UniProt reference clusters. *Bioinformatics* **23**, 1282–1288 (2007).

89. Seppey, M., Manni, M. & Zdobnov, E. M. BUSCO: Assessing Genome Assembly and Annotation Completeness. *Methods Mol. Biol.* **1962**, 227–245 (2019).

90. Huang, D. W. *et al.* The DAVID Gene Functional Classification Tool: a novel biological module-centric algorithm to functionally analyze large gene lists. *Genome Biol* **8**, R183 (2007).

91. Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms. *PLOS ONE* **6**, e21800 (2011).

92. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

93. Warnes, G. R. *et al. gplots: Various R Programming Tools for Plotting Data*.

Acknowledgments

This project was funded by NIH (Grant R15NS01686 and R21AG057974) and the University of Puerto Rico. The authors acknowledge the help from the Sequencing and Genomics Facility of the UPR Río Piedras & MSRC/UPR (NIH/NIGMS-Award Number P20GM103475) that are supported by an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the NIH under grant number P20GM103475.

Author contributions statement

D.Q.P. performed preliminary work for data generation. J.M.F. built reference transcriptome and annotation with contribution from S.C.G. and input from H.O.Z. in computational proceedings. D.Q.P. and J.G.A carried out the analysis of results and drafting of the paper. All authors reviewed the manuscript.

Additional Information

Competing Interest

The authors declare no competing interests.

Figure 1. Gene expression patterns during early stages of intestinal regeneration. (A) Volcano plot of differentially expressed transcripts (DETs) at day 1 of regeneration. (B) Volcano plot of DETs at day 3 of regeneration. (C) Bar plots show upregulated and downregulated DETs in one or both days of regeneration.

Figure 2. Gene ontology enrichment analysis of differentially expressed genes (DEGs) at day 1 and day 3 of regeneration.

Figure 3. Pathway enrichment analysis of differentially expressed genes (DEGs) at day 1 and day 3 of regeneration.

Figure 4. Heatmap of differentially expressed transcription factors (DETFs) in non-regenerating stage mesentery (Normal), and regenerating (days 1 3) intestines. DETFs showing a similar expression profiles are grouped in the dendrogram and labeled by colors in the right part of the heatmap.

Table 1. Comparison of expressed transcripts in RNA-seq with previous studies

Gene	RNA-seq		RT-PCR/qPCR	Northern Blot/In situ
	Day 1 (Log2FC)	Day 3 (Log2FC)	Day 3	Days 2-3
Myc[27]	2.72 ↑	2.05 ↑	↑	NA
Orpin[28]	3.84 ↑	4.23 ↑	↑	NA
β-catenin[27]	0.88	0.46[NS]	-[NS]	NA
Melanotransferrin[29]	3.55 ↑	3.7 ↑	↑	NA
Serum amyloid A protein[21]	2.39 ↑	3.2 ↑	NA	↑

WNT9[30]	2.69 ↑	2.88 ↑	NA	↑
BMP[30]	3.53 ↑	3.31 ↑	NA	↑
survivin[31]	-0.90[NS]	-0.31[NS]	-[NS]	-
mortalin[31]	2.41 ↑	2.45 ↑	↑	↑

↑: Upregulated; ↓: Downregulated; -: No significant change; NS: Not significant (Pvalue > 0.05)

Table 2. Top 10 upregulated DEGs at day 1 of regeneration

ID	Description	Log2FC	Adjusted P-value
Cluster-36465.0	cathepsin Z	9.44↑	1.37e-10*
Cluster-29766.0	uncharacterized protein LOC592324	8.03↑	1.33e-07*
Cluster-27058.0	cubilin	7.78↑	9.56e-05*
Cluster-50639.0	serine/threonine-protein kinase NLK	7.43↑	3.73e-05*
Cluster-26489.0	actin	7.37↑	1.78e-04*
Cluster-40404.0	translation elongation factor 2	7.20↑	3.16e-04*
Cluster-34538.0	elongation factor 1 alpha	7.15↑	4.40e-05*
Cluster-26421.0	V-type proton ATPase 16 kDa proteolipid subunit	7.14↑	9.04e-04*
Cluster-33192.0	peptidyl-prolyl cis-trans isomerase	6.88↑	5.02e-05*
Cluster-50300.0	Histone H4	6.71↑	1.97e-04*

↑: Upregulated (Log2FC > 2); *: Significant (Pvalue < 0.05); Log2FC: Log_2 Fold Change

Table 3. Top 10 downregulated DEGs at day 1 of regeneration

ID	Description	Log2FC	Adjusted P-value
Cluster-9180.0	protein FAM166B	-19.48↓	1.92e-05*
Cluster-6039.0	pancreatic lipase-related protein 2	-11.15↓	4.67e-06*
Cluster-3567.3046	sodium-coupled monocarboxylate transporter 1	-10.09↓	2.33e-12*
Cluster-3567.56678	histidine ammonia-lyase	-9.94↓	1.30e-04*
Cluster-3567.37979	glutamyl aminopeptidase	-9.73↓	1.03e-11*
Cluster-4476.0	echinoidin-like	-9.46↓	3.23e-02*
Cluster-3825.2	homeobox protein CDX-2	-8.64↓	2.33e-03*
Cluster-1995.1	uncharacterized protein LOC575162	-8.20↓	7.09e-10*
Cluster-28066.0	sushi domain-containing protein 2	-7.92↓	4.74e-21*
Cluster-3567.59390	aquaporin-8	-7.86↓	2.44e-07*

↓: Downregulated (Log2FC < -2); *: Significant (Pvalue < 0.05); Log2FC: Log₂ Fold Change

Table 4. Top 10 upregulated DEGs at day 3 of regeneration

ID	Description	Log2FC	Adjusted P-value
Cluster-9244.0	uncharacterized protein LOC592324	9.12↑	1.01e-09*
Cluster-11178.0	ubiquitin	7.82↑	7.54e-06*
Cluster-51274.1	protein Wnt-6	7.68↑	3.64e-28*
Cluster-36666.1	actin related protein 1	7.47↑	3.31e-05*
Cluster-10976.0	major vault protein	7.35↑	1.26e-04*
Cluster-49832.1	actin CyI, cytoplasmic	7.22↑	1.79e-05*
Cluster-11042.0	ubiquitin-40S ribosomal protein S27a	7.04↑	3.05e-05*
Cluster-44753.0	transcription factor VBP	6.95↑	9.12e-10*
Cluster-29958.0	cathepsin L1	6.88↑	8.98e-04*
Cluster-10300.2	translation elongation factor 2	6.85↑	5.92e-04*

↑: Upregulated (Log2FC > 2); *: Significant (Pvalue < 0.05); Log2FC: Log₂ Fold Change

Table 5. Top 10 downregulated DEGs at day 3 of regeneration

ID	Description	Log2FC	Adjusted P-value
Cluster-3567.56219	aquaporin-8	-11.08↓	2.52e-14*
Cluster-3567.3046	sodium-coupled monocarboxylate transporter 1	-10.05↓	3.14e-12*
Cluster-3825.2	homeobox protein CDX-2	-9.57↓	6.50e-04*
Cluster-14925.2	tenascin-N	-9.31↓	1.87e-03*
Cluster-2933.0	b (0, +)-type amino acid transporter 1	-8.17↓	2.53e-07*
Cluster-3567.37979	glutamyl aminopeptidase	-7.97↓	2.87e-22*
Cluster-2754.1	aminopeptidase N	-7.83↓	9.95e-26*
Cluster-3567.56678	histidine ammonia-lyase	-7.82↓	1.19e-03*
Cluster-3567.63443	phospholipase B1, membrane-associated	-7.82↓	4.01e-06*
Cluster-3567.34581	tubulin alpha-1A chain	-7.72↓	1.15e-06*

↓: Downregulated (Log2FC < -2); *: Significant (Pvalue < 0.05); Log2FC: Log₂ Fold Change