The Phage-shock-protein (PSP) Envelope Stress Response: Discovery of Novel Partners and Evolutionary History

Janani Ravi[1,2]*, Vivek Anantharaman[3], Samuel Zorn Chen[4], Evan Pierce Brenner[1], Pratik Datta[2], L Aravind[3]*, Maria Laura Gennaro[2]*.

[1]Department of Biomedical Informatics, University of Colorado Anschutz Medical Campus, Aurora, CO, USA; [2]Public Health Research Institute, Rutgers University, Newark, NJ, USA; [3]National Center for Biotechnology Information, National Institutes of Health, Bethesda, MD, USA; [4]Computer Science Engineering Program, Michigan State University, East Lansing, MI, USA.

*Corresponding authors:

janani.ravi@cuanschutz.edu; aravind@nih.gov; marila.gennaro@rutgers.edu

Running title: *PSP evolution across the tree of life*

Abstract

Bacterial phage shock protein (PSP) systems stabilize the bacterial cell membrane and protect against envelope stress. These systems have been associated with virulence, but despite their critical roles, PSP components are not well-characterized outside proteobacteria. Using comparative genomics and protein sequence-structure-function analyses, we systematically identified PSP homologs, phyletic patterns, domain architectures, and gene neighborhoods. This approach underscored the evolutionary significance of the system, revealing that the core PspA gene (Snf7 in ESCRT outside bacteria) was present in the Last Universal Common Ancestor (LUCA), and that this ancestral functionality has since diversified into multiple novel, distinct PSP systems across life. Several novel partners of the PSP system were identified: (i) the Toastrack domain, likely facilitating assembly of sub-membrane stress-sensing and signaling complexes, (ii) the newly-defined HAAS-PadR-like transcriptional regulator pair system, and (iii) multiple independent associations with ATPase, CesT/Tir-like chaperone, and Band-7 domains in proteins thought to mediate sub-membrane dynamics. Our work also uncovered links between the PSP components and diverse SHOCT-like domains, suggesting roles in assembling membrane-associated complexes of proteins with disparate biochemical functions. Results are available at https://jravilab.shinyapps.io/psp_evolution.

Importance

Phage shock proteins (PSP) are virulence-associated, cell membrane stress-protective systems. They have mostly been characterized in proteobacteria and firmicutes. We now show that PSP systems present in the Last Universal Common Ancestor, and that homologs have evolved and diversified into newly identified functional contexts. Recognizing the conservation and evolution of PSP systems across bacterial phyla contributes new knowledge to our current understanding of

stress response mechanisms in prokaryotes. Moreover, the newly discovered PSP modularity will

likely prompt new studies of lineage-specific cell-envelope structures, lifestyles, and adaptation

mechanisms. Finally, our results validate use of domain architecture and genetic context for

discovery in comparative genomics.

Keywords

Phage shock protein response | Psp | Envelope stress response | Phylogeny and evolution | Domain

architectures | Genomic contexts and neighborhoods | Comparative Genomics

Introduction

Cell membranes are complex, dynamic structures composed of bilayer and non-bilayer lipids and

proteins (1). These membrane components engage in critical processes including membrane

biogenesis, cell division, cell shape maintenance, small molecule transport, cell-cell and cell-

environment signaling, maintenance of proton motive force (PMF), and motility through cytoskeletal

proteins (2). Cell membranes continuously adapt to external stresses, particularly in unicellular

organisms (3–7). Failure to adjust leads to membrane damage and cell death. A specialized suite of

lipids and proteins operate to maintain membrane form and function under stress (8, 9), including

the multiprotein bacterial Phage Shock Protein (PSP) system, and the archaeo-eukaryotic ESCRT

system (10–20).

The origins and divergence of membrane structure and function remain open questions in biology

regarding the very basis of life. Most membrane functions are conserved across the tree of life

despite compositional and structural variations (1), which provides insights into evolutionary history

and niche specialization. Where such functions differ can also be highly informative. Prior research

has validated that evolutionary processes across life are constrained, and the potential evolutionary

paths are not limitless (21, 22). Mapping out evolutionary history is therefore more attainable by

existence of constrained characteristics, which is also a basis for phenotypic convergence.

Membrane maintenance mechanisms obey these same rules, and characterizing membrane

homeostasis systems like PSP can provide functional and evolutionary insights across life.

PSP expression is regulated by the peripheral membrane protein PspA and the homologous plant

protein Vipp1 (11–16, 23–31), which both contain the coiled-coil structure PspA domain (16, 17, 19,

20, 32, 33) (**Table S1**). Bacterial PspA senses and responds to extracellular stress (14, 23, 33–35).

Its domain, Pfam PspA_IM30 (inner membrane 30KDa domain) is found in bacteria, photosynthetic

eukaryotes, and archaea (15–17, 33, 36, 37) (**Fig. 1A**). Proteins containing PspA_IM30 are known to

facilitate membrane fusion (16) and prevent leakage of protons through the membrane (2, 23) in

response to a wide range of surface stresses, including changes in membrane potential, ΔpH (2,

23) and PMF (38–41), (l)antibiotic stress (7), heat/osmotic shock (42) and mechanical stress via

stored curvature elastic stress (13, 43–45). Vipp1 is involved in thylakoid biogenesis, vesicular lipid

transport and storage, and membrane shaping (16, 26, 27, 32).

The classical PSP system first discovered in Proteobacteria (46) is not conserved across all bacterial

and archaeal clades (12, 33, 47). However, the PSP systems described in Firmicutes and

Actinobacteria (3, 11, 14, 15, 35, 36, 48, 49) include clade-specific PSP proteins with distinct

domain architectures and conserved functions (*e.g.,* transcriptional regulation, membrane tethering

and relocation). Several key questions remain, including how PspA evolved to function with distinct

partners across diverse lineages, about the identities of other integral components of the PSP

system beyond the classical systems and complete characterization of cognate partners, if any

homologs participate in novel functions, and whether common themes emerge from the phyletic

patterns of PSP systems.

We set out to evaluate the evolutionary and functional significance of the PSP system through a

comprehensive analysis of PspA occurrences, associated genomic contexts, and phyletic spreads

across the three superkingdoms or domains of life (50, 51). We resolved PSP proteins into

constituent domain architectures and queried each individual domain in our comparative genomics

and phylogenetic analyses. This domain-centric approach is ideal to address the questions posed

above. We identified several novel players in the PSP envelope stress response system and

described their putative function and evolution. The PSP repertoire — including new members,

homologs, domain architectures, genomic neighborhoods, phyletic spreads, and phylogenetic

analyses in all representative species — is available for researchers to systematically explore

through tabular and graphical summaries on our user-friendly, interactive companion [web app](#).

100 Results and Discussion

The PSP system presents an evolutionary and functional paradox: PspA appears across the tree of

life but exists in distinct genomic neighborhoods and with different functional partners (15, 30, 52).

We, therefore, sought to identify and characterize all homologs of PspA and cognate partner

domains across a dataset representative of the tree of life. We used comparative genomics and a

computational molecular evolutionary approach to comprehensively identify PSP homologs. We

discovered several novel conserved, lineage-specific domain architectures and genomic contexts

of PSP systems and delineated their phyletic spreads across the tree of life in the process (see

Methods for details). All our findings (data, results, and visual summaries) are available in an

interactive and searchable web application ([jravilab.shinyapps.io/psp_evolution](#)) referred to

henceforth as the 'web app.' The web app can be queried with the protein accession numbers cited

inline throughout the *Results* section.

113 Phyletic spread of classical Psp components across the tree of life

Describing the PSP network of proteins across the tree of life requires characterization of classical

PSP components in Proteobacteria, Actinobacteria, and Firmicutes. Among Proteobacteria, we

analyzed the *E. coli psp* operon, which is widely conserved across Gammaproteobacteria (10, 12,

39, 46, 53–55). PspA, PspB, and PspC are each composed of their namesake domains spanning

almost their entire protein lengths. The transcriptional regulator PspF is encoded on the opposite

strand and shares a central promoter region with the *pspABC* operon in most

Gammaproteobacteria, and only with *pspA* in other organisms. PspF contains an enhancer-binding,

specialized AAA-ATPase NtrC-AAA domain, in addition to the Fis-type HTH (helix-turn-helix) DNA-

binding domain. In Actinobacteria, the *M. tuberculosis psp* operon (14, 15, 36) includes the

namesake PspA protein; a transcription factor, ClgR, containing a cro-like-HTH (cHTH) DNA-

binding domain unique to actinobacteria; an integral membrane protein, Rv2743c, distinct from *E.*

coli PspC and PspB; and Rv2742c, an uncharacterized protein carrying a "Domain of Unknown

Function,'' DUF3046 (14, 15, 36). For consistency, we renamed Rv2743c and Rv2742c PspM and

PspN, respectively (Datta *et al.*, unpublished). *B. subtilis,* a member of Firmicutes, contains the *lia*

operon. LiaI is a small transmembrane (TM) protein (11, 49, 56), LiaF contains N-terminal TM helices

(DUF2157) and a C-terminal globular domain (DUF2154), and LiaG contains an uncharacterized

domain DUF4097. When we analyzed the domain architectures of these operons, we defined two

novel domains of the PSP system – the LiaI-LiaF-TM sensory domain that unifies four TM domains

(4TM) and DUF2157 from LiaI/LiaF, as well as Toastrack, unifying DUF4097/DUF2154/DUF2807

Pfam models. The *lia* operon is controlled by a two-component system LiaRS (11, 35). LiaR contains

a receiver domain, and a winged HTH (wHTH) DNA-binding domain (REC-wHTH), while LiaS is a

sensor kinase with a TM region, intracellular HAMP, and a histidine kinase (HISKIN) signaling

domain (57). Information on domains in this section is presented in **Text S1.1** and **Table S1**, and

graphically in **Fig. 1A** and **1B**.

PspA

From six distinct queries (Proteobacteria, Firmicutes, Actinobacteria, Cyanobacteria, Plants; see

Methods), homologs of PspA were sought comprehensively. Consistent with prior studies,

PspA_IM30 was identified in most bacterial and archaeal lineages (**Fig. 1C**; web app). Within

eukaryotes, only the SAR and Archaeplastida (greater plant lineage) carry PspA_Vipp1 homologs

(**Fig. 1C**). Analysis using multi-query sequence searches, we also identified Snf7 proteins from the

eukaryotic membrane-remodeling ESCRT systems **(18, 63–67)** as homologs (**Fig. 1B**). Snf7 was

then used for iterative searches, which yielded several pan-eukaryotic and archaeal homologs

distinct from Vipp1 (**Fig. 1B** and **1C**; **Table S3**), supporting a considerably wider evolutionary

distribution for PspA, corroborated by other recent reports (19, 20).

Toastrack and LiaI-LiaF-TM

Iterative domain-centric searches and multiple sequence alignments revealed that LiaI and LiaG

were similar to N-and C-terminal regions of LiaF. We define these two novel domains as **Toastrack**

(unifying DUF2154, DUF4097, DUF2807) and **LiaI-LiaF-TM**. Toastrack-containing proteins

characterized by a single-stranded right-handed beta-helix fold with a unique N-terminal 7-stranded

region featuring complex intertwining of the beta-helix strands (e.g., PDB: 4QRK, 4OPW). They are

pan-bacterial and sporadic in archaea and eukaryotes (**Fig. 1C**). Toastrack is frequently coupled to

PspC, LiaI-LiaF-TM, or both (**Fig. 4**; **Table S4**). LiaI-LiaF-TM contains a 4TM region, and is primarily

in bacteria, with transfers to Euryarchaeota (**Fig. 1B**). Search methodology is described in **Text**

S1.1, and visualizations of these novel domains provided on the 'MSA' tab of the web app.

PspB and PspC

PspB, often part of the proteobacterial *psp* operon, is another integral membrane protein implicated

in stress sensing and virulence (58–60). PspB has an N-terminal anchoring TM helix followed by an

α-helical cytoplasmic globular domain. While PspB is rarely found outside proteobacteria (**Fig. 1C**),

we discovered previously unrecognized divergent PspB domains fused to PspC (see section on

PspC architectures).

PspC was first identified in the proteobacterial PSP system; it is critical in sensing membrane stress

and restoring envelope integrity. Recent studies demonstrate that PspC in some bacterial species

may function independently of PspA in response to certain membrane stressors (59, 61–63).

Another integral membrane protein, our analysis showed PspC has two TM helices, the first being

a cryptic TM ('MSA' tab, web app). A conserved arginine (a positively charged residue) found

between the two TM regions is reminiscent of structures serving in membrane-associated stimulus

sensing, which may provide clues towards the biological roles of PspC. We observed that the PspC

domain is pan-bacterial and also present in a few archaeal clades (**Fig. 1C**). Our analysis showed

that PspC has two TM helices, the first being a cryptic TM. We also identified several novel domain

architectures and genomic contexts of PspC (see section on *PspC architectures*).

PspM and PspN

Domain-centric searches found no discernible homologs for PspMN outside Actinobacteria (**Fig.**

1C; web app), as with prior full protein searches (15, 36, 64). Two TM helices make up the

corynebacterial integral membrane partner **PspM** (encoded by *rv2743c*). Our analyses revealed

minimal PspM variation by MSA ('MSA' tab, web app) and a narrow phyletic spread restricted to

mycobacteria and corynebacteria (**Fig. 1B**) (15, 36, 64).

The fourth member in the Mycobacterial operon, PspN (Rv2742c), contains a C-terminal DUF3046

and an N-terminal domain we call **PspN_N** (**Table S1**; **Text S1.2**; 'MSA' tab in the web app) (15,

36, 64). **DUF3046** is prevalent only in actinobacteria (**Fig. 1C**). Moreover, the *M. tuberculosis*

genome carries a second copy of DUF3046 (*rv2738c*), located four genes downstream of *pspN*. We

infer that Rv2738c, rather than *pspN*, contains the ancestral copy of DUF3046 (**Text S1.2**). DUF3046

found as part of PspN is likely a duplicate translocated into the PspN open reading frame (ORF) of

mycobacteria, especially the *M. tuberculosis* complex. Further, unlike the pan-actinobacterial

DUF3046, the N-terminal region, PspN_N, is conserved only in *M. tuberculosis*, with remnants of

the coding region existing as potential pseudo-genes or degraded sequences in a few closely

related mycobacteria (*e.g., M. avium*).

Evolution of PspA

After identifying eukaryotic Snf7 as homologous, we sought to expand on the history of PspA. We

used a structure-informed MSA and phylogenetic tree ('MSA' tab, web app; **Fig. 2; Table S1**) to

investigate the PspA/Snf7 superfamily, and define its evolutionary origins.

Identifying clades carrying PspA and Snf7

PspA+ clades comprise all versions of the PspA domain phylogenetically similar to the 'classical'

PspA domain rather than the Snf7 domains. We found that most bacterial lineages contain PspA+

clade members (**Fig. 1C**), with a few transfers to archaea. Among eukaryotes, only those containing

plastids and two flagella (comprising the SAR/HA supergroup and excavata) have members of this

clade (**Fig. 1C**; **Fig. 2**; web app) (65, 66).

Similarly, we found that the curated set of PspA-like proteins contains a divergent cluster of proteins

belonging to what we now call the Snf7+ clade (**Fig. 2**; web app). Snf7 is part of the ESCRT-III

complex required for endosome-mediated trafficking via multivesicular body formation and sorting

and has a predominantly archaeo-eukaryotic phyletic pattern (**Fig. 1C**) (18, 67–71).

Tracing the roots of the PspA/Snf7 superfamily to the last universal common ancestor

(LUCA)

216
We explored the evolution of the PspA-Snf7 superfamily using an MSA of comprehensively selected

PspA/Snf7 homologs from distinct clades across the tree of life (50, 51) and different domain

architectures (**Fig. 2**; web app). The MSA revealed a unique insertion of heptad repeats in

actinobacterial PspA sequences, likely a lineage-specific adaptation. A few cyanobacterial PspA

homologs similar to the ancestral plant variant Vipp1 also contained a C-terminal extension (16, 32,

72). A maximum-likelihood phylogenetic tree was generated using this MSA (**Fig. 2**; web app), which

showed clear clustering into i) easily distinguishable clades of homologs from Actinobacteria,

Firmicutes, and Proteobacteria; ii) the eukaryotic Vipp1 clade grouped with cyanobacterial

homologs indicating plastid origins; and iii) Snf7 domain-containing homologs from archaea and

eukaryotes forming a well-defined, separate branch. Together, these observations support that the

PspA/Snf7 superfamily is ubiquitous across the superkingdoms/domains of life and originated from

a PspA-like gene present in the last universal common ancestor (LUCA), as independently

corroborated by recent studies (19, 20).

PspA: Novel architectures and neighborhoods

We used our comprehensive, domain-level search to delineate the domain architectures and

genomic contexts of the PspA homologs in organisms from diverse clades.

Domain Architectures

Our searches revealed that the underlying domain architectures of most PspA homologs (>98%;

~2500 homologs) show little variation (web app). In most lineages, PspA homologs contain only the

characteristic PspA_IM30 domain (**Fig. 3**; **Table S3;** web app). The remaining <2% of homologs

contain PspA_IM30 repeats in tandem or fusions with other domains, including NlpC/P60 and a

novel PspA-associated (PspAA) domain (**Fig. 3**; **Text S2.1**). The PspA_IM30–NlpC/P60 hydrolase

fusion is predicted to catalyze the modification of phosphatidylcholine, likely altering membrane

composition (**Fig. 3**; **Table S3**; AFZ52345.1) (73). Similarly, Snf7 showed minimal variation in domain

architecture, with occasional fusions (<5%) in eukaryotes, with some potentially attributable to

genome annotation errors. Some actinobacteria have an Snf7 homolog fused to an RND-family

lipid/fatty acid transporter, flanked by two genes encoding a *Mycobacterium*-specific TM protein

with a C-terminal cysteine-rich domain (74).

Paralogs

We found that genomes containing multiple copies of PspA do not maintain domain architecture

and genomic context among paralogs (**Fig. 3**; **Table S3**; 'Phylogeny –> Paralog' tab, web app). In

cyanobacteria, dyads/triads of PspA paralogs occur as adjacent, occasionally fused genes (**Fig. 3**;

Table S3; *e.g.,* BAG06015.1). Consistent with earlier reports (25, 27), we observed that these

neighboring PspA genes are part of two distinct clusters of homologs, one resembling the bacterial

PspA and the other the eukaryotic (plastid) Vipp1 (**Fig. 3**). Snf7 gene clusters also appear

occasionally in archaeal and eukaryotic species (**Table S3**; *e.g.* CBY21170.1). This suggests

tandem gene duplication and may relate to the role of these clusters in membrane stabilization as

oligomeric complexes, as in the eukaryotic ESCRT systems (75). Further expansion on PspA/Snf7

paralogs and their likely evolution, including inference of gene duplication *vs.* horizontal gene

transfer from domain architectures and genomic contexts, is available in the web app. Occasionally,

PspA and Snf7 co-occur in conserved gene neighborhoods (**Fig. 3**; **Table S3**) such as in

deltaproteobacteria (AKJ06548.1); in bacteroidetes (CBH24266.1), these neighborhoods also

contain a third gene encoding a coiled-coil protein with a structure reminiscent of the PspA coiled-

coils (**Table S3**).

Novel variations of classical Psp genomic contexts

We next explored the gene-neighborhood themes for each Psp member (**Fig. 3**; **Table S3**). Major

themes representing variations of classical Psp operons are summarized below. For the full list of

genomic contexts and phyletic spreads, see web app: 'Data,' 'Genomic Contexts' tabs.

Diversity of the pspFABC *operon*

In addition to proteobacteria, the classic *pspABC* occurs in nitrospirae and some spirochaetes (**Fig.**

3; **Table S3**). There are also variations to this theme (web app): 1) PspC is fused to a divergent C-

terminal PspB in addition to a single PspB in the operon (**Fig. 3**; **Table S3**; *e.g.,* ANW39986.1); 2)

PspB duplicates appear in the operon (**Fig. 3**; **Table S3**; *e.g.,* AOL22920.1); 3) PspD occurs along

with this operon only in gammaproteobacteria (**Fig. 3**; **Table S3**; *e.g.,* ANW39986.1). Key variations

in the transcription regulator PspF (NtrC-AAA and HTH) include: 1) an operon with additional genes

for PspB and PspC fusions and Toastrack-containing proteins in a few spirochaetes; 2) operons in

gammaproteobacteria with genes encoding NtrC-like transcription factors with N-terminal ligand-

binding ACT and PAS domains (**Fig. 3**; **Table S3**; *e.g.,* AEN65073.1) (76), and a further protein of

the DO-GTPase family, predicted to play a role in membrane-related stresses (77, 78), which

occasionally feature an additional PspF; and 3) in some alphaproteobacteria, the PspC in PspFABC

has been replaced by multiple Toastrack-containing proteins (**Fig. 3**; **Table S3**; *e.g.,* ANQ40502.1).

Thus, classic *pspABC* presents variation in structure and encoded function.

Associations with Vps4 and other classical AAA+-ATPases

The core of an ESCRT complex in archaea is defined by the co-occurrence of one or more copies

of Snf7 (*e.g.,* OLS27540.1; [**Table S3**]) and a gene encoding a **VPS4-like AAA+-ATPase** (with an

N-terminal MIT and C-terminal oligomerization domain [**Table S2**] (79). In our analysis, we observed

diversity in *vps4* gene neighborhoods (**Text S2.2**). ESCRT complexes are well-studied in

eukaryotes, but less so in archaea, where they were originally associated with cell division and

named for this role (Cdv system) (80). Crenarchaeota and Thaumarchaeota contain *cdvABC*. *cdvA*

has no eukaryotic homolog, while *cdvB* and *cdvC* encode homologs of ESCRT-III proteins and

Vps4, respectively. Recent studies have shown that eukaryotes originate within the archaeal clade

of Asgardarchaeota (80). Consistent with this, Asgardarchaeota show a conserved gene

neighborhood coupling Snf7, a VPS4-like AAA+ ATPase gene, and a gene coding for ESCRT-II

wHTH-domain (81) corresponding to the cognate eukaryotic ESCRT complex. These operons may

be further extended with additional copies of Snf7 genes and other genes coding for a TM protein

and an ABC ATPase. Additionally, examples of transfers of VPS4-like AAA+-ATPase from archaea

to several bacterial lineages were observed (**Table S3**; *e.g.,* ACB74714.1). In these cases, *snf7* is

displaced by an unrelated gene containing TPR repeat and 6TM, suggesting membrane proximity.

The bacterial PspA (**Table S3**; *e.g.* AEY64321.1) may occur with a distinct protein containing two

AAA+-ATPase domains – with the N-terminal copy inactive – in some bacterial clades (**Text S2.2**).

Both PspA and the membrane-associated Snf7, along with AAA+-ATPase, may occur in longer

operons with other genes encoding an ABC-ATPase, permease, and a solute- and peptidoglycan-

binding protein with PBPB-OmpA domains (**Table S3**; *e.g.,* OGG56892.1). These associations

between certain bacterial PspA and Snf7 clade members with AAA+-ATPases suggests a role in

ATP-dependent membrane remodeling (**Fig. 3**; **Table S3**) comparable to Snf7 and VPS4 in the

ESCRT-III-mediated membrane remodeling in the archaeo-eukaryotic lineage (19, 20, 67–69, 71,

75). The additional transporter components seen in some of these systems suggest a link between

this membrane remodeling and solute transport. Here the OmpA domain could interact with

peptidoglycan, while the PBPB domain binds the solute trafficked by the linked ABC transporter.

Finally, recent structural studies have reached similar evolutionary conclusions, i.e., that Vipp1,

PspA, and Snf7 proteins arose from a common ancestor and can play similar roles (19, 20).

Operons with CesT/Tir-like chaperones and Band-7 domain proteins

We discovered two novel overlapping genomic associations across various bacteria linking PspA

with the Band-7 domain and CesT/Tir-like chaperone domain proteins (**Fig. 3**; **Tables S2** and **S3**;

'MSA' tab, web app). Band-7 has been implicated in macromolecular complex assembly, playing a

chaperone-like role in binding peptides (82), and in both membrane-localization and membrane

dynamics (83). CesT/Tir chaperone domains have been shown to mediate protein-protein

interactions in the assembly and dynamics of the Type-III secretion systems of proteobacteria (84).

Using profile-profile searches, we identified a previously uncharacterized protein encoded by genes

linked to the yjfJ family of *pspA* genes (*e.g.,* ANH61663.1) as a novel member of the CesT/Tir

superfamily (**Fig. 3**; **Tables S1** and **S2**; *e.g.,* ANH61662.1). We also observed that the

proteobacterial proteins in the neighborhood of *pspA* (**Fig. 3;** *e.g.,* BAB38581.1), typified by YjfI

from *E. coli* (DUF2170; **Table S2**; *e.g.,* BAB38580.1), contained a CesT/Tir superfamily domain (85).

The dyad encoding CesT/Tir–PspA forms the core of one class of conserved gene neighborhood

(**Table S2**). We observed it as either a standalone unit or within larger operons, such as one group

combining CesT/Tir–PspA (**Fig. 3**; **Table S3**; ANH61663.1) with i) a gene encoding a membrane-

associated protein with the domain architecture TM+Band-7+coiled-coil+flotillin, ii) a novel AAA+-

ATPase fused to N-terminal coiled-coil and β-propeller repeats, and iii) a 3TM domain protein

prototyped by *E. coli* YqiJ and *B. subtilis* YuaF (previously implicated in resistance to cell-wall

targeting antibiotics) (86, 87). In some proteobacteria, this operon also contains a phospholipase D

(HKD) superfamily hydrolase (**Fig. 3**). Related abbreviated operons coding only for the Band-

7/flotillin-containing protein, the YqiJ/YuaF TM protein, and in some cases, the aforementioned

AAA+-ATPase protein is more widely distributed across bacteria and archaea. These might function

with PspA homologs encoded elsewhere in the genome.

The second major group incorporates the dyad into a 6–7 gene operon (**Fig. 3**; **Table S3**; *e.g.,*

AAN56746.1) featuring a gene encoding a spermine/spermidine-like synthase domain (86) fused to

an N-terminal 7TM transporter-like domain (**Fig. 3**; **Tables S2** and **S3**; *e.g.,* AAN56744.1). This

operon encodes three additional uncharacterized proteins, including the YjfL family 8TM protein

(DUF350), a novel lipoprotein (Ctha_1186 domain), and a β-strand rich protein that we predict to be

intracellularly localized (**Fig. 3**; **Tables S3** and **S4;** DUF4178; *e.g.,* AAN56747.1). In some cases, the

last gene in this operon codes for a ribbon-helix-helix (RHH) domain transcription factor (**Fig. 3**;

Tables S2 and **S3**).

The third group combines the CesT/Tir–PspA dyad (AMJ95269.1) with polyamine metabolism genes

encoding an ATP-grasp peptide ligase (AMJ95273.1) related to the glutathionyl spermidine

synthetase (**Fig. 3**; **Table S3**). This association was also recently noticed in a study of AdoMet

decarboxylase gene linkages (88). The operon also encodes a potassium channel with intracellular-

ligand-sensing, a YjfL family 4TM protein (**Tables S2** and **S3**; DUF350), a metal-chelating lipoprotein

(DUF1190), and another uncharacterized protein with predicted enzymatic activity (DUF2491).

Alternately, two variants of these operons lack the CesT/Tir–PspA dyad: one where its replacement

is a gene encoding a distinct protein occurring as either a secreted or lipoprotein version (DUF4247,

e.g., CCP45395.1) and a predicted beta-sandwich enzymatic domain with two highly conserved

histidines, as well as a conserved aspartic acid and glutamic acid (DUF2617, *e.g.,* CCP45394.1). In

the second variant, the dyad is replaced by a protein containing a Band-7 domain fused to C-

terminal 2TM and SHOCT domains (see below). These operons encode two further polyamine

metabolism genes, an AdoMet decarboxylase, and a polyamine oxidase.

Lastly, the *Bacillus subtilis ydjFGHL* operon typifies a coupling in Firmicutes of *pspA* (**Table S1**;

ANX09535.1) with Band-7 and C-terminal Zinc Ribbon (ZnR) domains in YdjL, and two ZnRs

followed by an uncharacterized domain related to YpeB featuring an NTF2-like fold and a TM

segment in protein YdjG. The conserved partner of PspA in these systems is the membrane-

associated YdjH, which is characterized by a so-called "**TPM phosphatase domain**" followed by

a low- complexity segment or a long coiled-coil (**Fig. 3**; **Table S3**) (88). Competing hypotheses

suggest that this domain either encodes a generic phosphatase based on its plant homolog (88) or

is involved in membrane-associated photosystem II repair and respiratory complex III assembly (89,

90). Our finding that it associates with PspA parallels its association with the chaperone CesT/Tir

and Band-7 involved in macromolecular assembly. Thus, our result supports the hypothesis that

the TPM domain with PspA plays a chaperone-like role in assembly of membrane-linked protein

complexes.

Membrane dynamics with Chaperone-like domains

The contextual association of PspA with AAA+-ATPase, two distinct families of CesT/Tir type

chaperones, Band-7 and TPM domains further supports functional roles of PspA homologs in

recruiting macromolecular assembly machinery to the inner leaf of the cell membrane. Polyamines

like spermine/spermidine have been implicated in membrane stability in a concentration-dependent

manner (91). The repeated coupling of polyamine metabolism genes we identified in PspA-CesT/Tir

operons implies that a PspA-based system of membrane dynamics monitors polyamine

concentration or aminopropylation of other substrates (92) to alter membrane structure and

membrane-associated protein complexes. We propose that these systems link extracellular stress

sensing with intracellular membrane dynamics based on the frequent coupling with cell-surface

lipoproteins. In particular, flotillin domain-containing proteins are likely to be recruited to lipid

subdomains with special components (*e.g.,* cardiolipin-rich inner membrane) to interface with PspA

(87).

The PspAA domain

Our *pspA* gene neighborhood analysis identified a unique partner in archaeal and bacterial phyla: a

protein containing a novel trihelical domain occurring in a two-gene cluster with *pspA* (**Fig. 3**; **Table**

S3; *e.g.,* AAM04874.1). This domain often occurs independently but is occasionally fused to an N-

terminal PspA in Actinobacteria and Chloroflexi (**Fig. 3**; **Table S3**; *e.g.,* ACU53894.1). This **PspA-**

associated domain A (PspAA) (**Fig. 3**; **Table S2**; **Text S1.3**) occurs in a PspA–PspAA dyad in a

two-component system (*e.g.*, CAB51252.1), and in another system consisting of a membrane-

associated metallopeptidase, another novel domain **PspA-associated domain B (PspAB)** (*e.g.*,

AAZ55047.1), and an occasional third gene encoding a SHOCT-like domain (**Fig. 3**; **Tables S2** and

S3; web app).

PspA with PspM (ClgRPspAMN) or Thioredoxin

Certain *pspA* homologs may occur in Cyanobacteria and Actinobacteria as an operonic dyad with

a gene encoding an active thioredoxin domain-containing protein (15). PspA in this dyad is typified

by corynebacterial RsmP (*e.g.,* AKK09942.1; **Fig. 3**; **Tables S2** and **S3**; **Text S2.3**), which is

predominantly found in rod-shaped actinobacteria and regulates cell shape via phosphorylation

(93). These PspA homologs either occur with thioredoxin (*e.g.,* AKK09942.1) or with a ClgR-like

cHTH and PspM genes (*e.g.,* CCP45543.1; **Fig. 3, Fig. S1**, **Table S3**). Corynebacterium spp. contain

two distinct PspA paralogs each displaying one of the above contextual associations ('Paralog' tab,

web app). The same thioredoxin family occurs with a different PspA clade (typically two copies) in

Cyanobacteria (*e.g.,* AFZ14666.1; Fig. 3; Table S3). The association of the ClgR-PspM and

thioredoxin and their mutual exclusion in the operon suggest that they act as alternative regulators

of the PspA homologs — repressors in the case of the ClgR, and a potential redox regulator in the

case of thioredoxin (**Fig. S1**; **Text S2.3**).

Operons with two-component systems

Two-component systems often associate with the PSP system. For instance, the *lialHGFSR* operon

from *B. subtilis* has been studied in the context of lantibiotic resistance (*e.g.*, ANX06812.1; **Fig. 3**;

Table S3) (11). In a few paenibacilli, genes encoding an additional PspA and PspC with a C-terminal

coiled-coil have been inserted into a *lialHGFSR*-like operon (*e.g.,* APB74393.1; **Fig. 3**; **Table S3**). In

Actinobacteria (*e.g.*, CAB51252.1; **Fig. 3**; **Table S3**), the PspA–PspAA dyad occurs with a firmicute-

like two-component system.

Classic two-component transcriptional signaling system

We identified two-component signaling system operons linked to either *pspA* or *PspC* in Firmicutes,

Actinobacteria, and other clades (**Fig. 3**; web app). In these operons, a membrane-bound histidine

kinase (HISKIN) communicates an external stress signal to the response regulator (REC–HTH)

protein, triggering a transcriptional response. It is very likely that PspA, LiaI-LiaF-TM, and Toastrack

tie into a two-component system regulating membrane integrity and permeability in response to the

stress signal. In Actinobacteria, where the HISKIN is fused to PspC, the signal is presumably sensed

by PspC. When *pspC/Toastrack* operons with two-component systems lack *pspA* in their

immediate operonic neighborhood, they likely recruit PspA proteins from elsewhere in the genome

to elicit the stress response function.

PspA-free variations of domain architectures and gene-neighborhood

We next investigated domain architectures and genomic contexts involving PSP components that

lack PspA, including those with the PspC and Toastrack domains.

PspC domain architectures and gene-neighborhood

PspC is present in most bacteria and the archaeal clades of Euryarchaeota and Asgardarchaeota.

Some orthologs of PspC are fused to an N-terminal ZnR domain (*e.g.,* ABC83427.1, **Table S4**),

while most other homologs occur either in predicted operons or fusions with multi-TM domains,

such as the LiaI-LiaF-TM and PspB (**Fig. 4**; **Table S4**). PspC also fuses to diverse signaling domains

such as i) HISKIN from two-component systems (see above); ii) a novel signaling domain we term

the <u>H</u>TH-<u>a</u>ssociated <u>α</u>-helical <u>s</u>ignaling domain (**HAAS**), which overlaps partly with the Pfam model

for DUF1700 (**Table S2**), and iii) the Toastrack domain (**Table S1**). Moreover, actinobacteria contain

a Lia-like system without PspA. The HISKIN domain in these systems is fused to N-terminal PspC

and LiaF-LiaI-TM domains (*e.g.,* AIJ13865.1) and accompanied by a REC–HTH-containing protein

(**Fig. 5**; **Table S4**). This core shares a promoter region with a second gene on the opposite strand

containing a PspC domain that is fused to additional TMs and, in some cases, a Toastrack domain

(*e.g.,* AIJ13866.1; **Fig. 5**; **Table S4**). These associations strongly imply that PspC acts as a sensor

domain to detect and communicate a membrane-proximal signal.

Contextual associations of the Toastrack Domain

The **Toastrack** domain repeatedly emerges as a partner to the PSP system. As noted, genes

encoding Toastrack-containing protein occur concurrently with *pspA* in several conserved

neighborhoods (**Fig. 4**; **Table S4**) and are fused to *pspC* in *pspA*-free settings. We found that the

Toastrack domains tend to be C-terminal to various TM domains such as PspC, LiaI-LiaF-TM, and

other multi-TM/single-TM domains unique to these systems (**Fig. 4**; **Text S2.4**). These fusions

strongly suggest that Toastrack is an intracellular domain. Further, the N-terminal regions of several

Toastrack architectures contain at least two divergent homologs of the bi-helical SHOCT domain

(*e.g.,* CAB15517.1; **Figs. 4** and **5**; **Tables S1** and **S4**), which we call **SHOCT-like** domains, which

are partly detected by the Pfam **DUF1707** model (**Text S2.4**) (94)**.** We observed these SHOCT-like

domains at both the N- and C-termini of various globular domains, including Toastrack. Therefore,

we predict that this SHOCT-like domain plays a role in anchoring disparate domains to the inner

leaf of the membrane.

The DNA-binding domain **LytTR** may also be encoded by operons carrying Toastrack fused to LiaI-

LiaF-TM in Firmicutes (*e.g.,* CAL82154.1; **Fig. 5**; **Table S4**). The Toastrack-containing protein (*e.g.,*

CAB15517.1) occurs in an operon with a *pspC* homolog (CAB15516.1) and has been implicated in

the activation of the membrane-associated *liaFSR* operon (**Table S1**) and protection against

membrane permeabilization (95–97). We also identified *pspA*-free variants of the classic *lia* operon

with Toastrack and a two-component system, typified by *vraT* of *Staphylococcus aureus*

(ABD31150.1; **Fig. 5**; **Table S4**). These systems carry LiaI-LiaF-TM and Toastrack with a two-

component system (*vraSR*) and are involved in methicillin and cell-wall-targeting antibiotic

resistance (3, 98–100). A similar organization of LiaI-LiaF-TM and Toastrack containing a protein

with a two-component system is found in a few bacterial species (*e.g.,* among Proteobacteria,

ABD83157.1, Ignavibacteriae, AFH48155.1; **Table S4**).

We discovered several other conserved gene neighborhoods centered on Toastrack-containing

genes likely to define novel functional analogs of the PSP system with potential roles in membrane-

linked stress response. One of these, a four-gene context found across diverse bacterial lineages,

includes a previously uncharacterized protein with similarity to the Pfam model DUF2089. The Pfam

model **DUF2089** can be divided into an N-terminal ZnR, central HTH, and C-terminal SHOCT-like

domains (ADE70705.1), variants of which appear with LiaI-LiaF-TM, B-box, or PspC domains (**Figs.**

4 and **5**; **Tables S2** and **S4**; **Text S2.4**). We propose that these Toastrack contexts function similarly

to the classical *lia* operon in transducing membrane-associated signals to a transcriptional output

affecting a wide range of genes, including through an operonically linked sigma factor.

Similarly, we observed operons coupling a membrane-anchored Toastrack-containing protein with

an ABC-ATPase, a permease subunit, a **GNTR-HTH** transcription factor with distinct C-terminal α-

helical domain (**Fig. 5**; **Tables S2** and **S4**; **Text S2.4**), and a DUF2884 membrane-associated

lipoprotein (*e.g.,* AJI31452.1). The latter might function as an extracellular solute-binding partner for

the ABC-ATPase and permease components. We found a similar operon in Actinobacteria with the

ribbon-helix-helix (**RHH**) domain transcription factor replacing **GNTR-HTH** (**Table S2**; **Text S2.4**).

In some Actinobacteria, the Toastrack domain is fused to a SHOCT-like domain or occurs in a

transport operon, likely coupling Toastrack to transcriptional regulation, membrane-proximal signal

sensing, and transport (**Fig. 5**; **Table S4**).

Based on these observations, we propose that the two extended β-sheets of the Toastrack domain

(formed by the β-helical repeats) facilitate the assembly of sub-membrane signaling complexes by

providing two expansive surfaces amenable to protein-protein interactions near the membrane.

Protein docking to the Toastrack domain may also activate transcription via fused or associated

HTH or LytTR domains, associated two-component systems, or the HAAS-HTH system.

The HAAS-PadR-like wHTH systems

We discovered a novel signaling system that frequently co-occurs with PspC and Toastrack

domains in gene neighborhoods that do not contain *pspA* (**Figs. 4** and **5**). This system consists of

two components, the HAAS domain and the PadR-like wHTH (**Table S2**). The HAAS domain is

partly similar to the Yip1 protein family (Pfam models DUF1700 and DUF1129; Pfam clan: Yip1). An

HHpred search further unifies this model with the Pfam profile DUF1048 (PDB: 2O3L). We therefore

refer to them collectively as the **HAAS** superfamily (HAAS under 'MSA' tab, web app). The core

HAAS fold has three consecutive α-helices, with the third helix displaying an unusual kink

corresponding to a conserved GxP motif, predicted to form part of a conserved groove that

mediates protein-protein interactions. The HAAS domain always occurs in gene-neighborhoods

coupled with a protein containing a standalone PadR-like wHTH DNA-binding domain (**Fig. 5**). This

co-migration suggests that HAAS transduces a signal from domains fused to it to this wHTH

transcription factor.

When coupled with PspC, the HAAS domain shows three broad themes: 1) it is found as part of a

multidomain TM protein with additional PspC, LiaI-LiaF-TM, and Toastrack domains; 2) it is fused

directly to a Toastrack domain in a gene neighborhood that also codes for a PspC domain protein,

occasionally fused to other TM domains; 3) constitutes a part of a TM protein fused to conserved

multi-TM domains other than the LiaI-LiaF-TM or PspC (*e.g.,* VanZ) (**Figs. 4** and **5**; **Table S4**). These

gene neighborhoods code for standalone *pspC* genes. Additionally, the HAAS domains occur in

PspC-independent contexts, but are typically coupled to the N-terminus of other multi-TM or

intracellular sensory domains. We found conserved fusions to at least 5 distinct multi-TM domains

(**Figs. 4** and **5**; **Table S4**), such as 1) the FtsX-like TM domains with extracellular solute-binding

MacB/PCD domains (*e.g.,* ACO32024.1); 2) FtsW/RodA/SpoVE family of TM domains (*e.g.,*

CAC98500.1) (100); 3) various uncharacterized 6TM, 4TM, and 2TM domains. In addition to

Toastrack, the HAAS domain may combine with other intracellular domains such as the

pentapeptide repeat domains mostly in firmicutes (*e.g.,* AOH56696.1; **Fig. 5**; **Table S4**). The

organisms in which these operons occur have Psp components elsewhere in the genome. These

proteins might be recruited to the stress response system, as suggested by the effects of PadR-

like wHTH deletion studies in the Listeria Lia systems (101).

In bacteria without the typical *lia* operon (carrying the LiaRS-like two-component system), we noted

that the HAAS–PadR-like-wHTH dyad replaces the histidine kinase-receiver domain transcription

factor pair. Therefore, we propose that the HAAS-PadR dyad constitutes an alternative to the

classical two-component systems. The HAAS domain may undergo a conformational change in

response to membrane-linked or intracellular signals detected by the sensor domains (*e.g.,* LiaI-

LiaF-TM or PspC) that fuse to or co-occur with the dyad. This conformational change is likely

transmitted to the associated PadR-like wHTH via the groove associated with the conserved GxP

motif; this change might lead to the release of the transcription factor to mediate a downstream

transcriptional response.

546   A unified view of PSP partners and evolution

We reconcile our findings on PSP-related genomic contexts and partner domains within the

framework of the tree of life to assemble the PSP puzzle. The above results include ~20,000

homologs of key PSP proteins, ~200 novel domain architectures (minimum 2 species) and ~500

genomic contexts (minimum 5 species) ([web app](#)). To visualize the broader context of PSP systems

across the superkingdoms, we first built a network of domain architectures for the extended PSP

system (Fig. 6A), with domains as nodes and co-occurrence within a protein as edges. This network

consolidates the PSP system with known and novel connections, including those between PspA

and PSP partner domains. Next, we summarized the phyletic spread and prevalence of all PSP

members and partners in a single heatmap (**Fig. 6B**). This view indicates that, of the PSP

components, (i) only PspA/Snf7 is present in all eukaryotes, whereas Toastrack has been

transferred from bacteria to certain eukaryotes (*e.g.,* human FAM185A with orthologs in animals

and other eukaryotes) ; (ii) PspC and PspA/Snf7 are present in most archaeal lineages, (iii)

occasional transfers of Toastrack, LiaI-LiaF-TM, and SHOCT domains are observed from bacteria

to Euryarchaeota; (iii) in bacteria, domains such as Toastrack, PspC, PspA, and LiaI-LiaF-TM, tend

to co-migrate. Finally, using an 'upset plot,' we unified: i) the relative occurrence of domain

architectures from PSP members and their most common neighbors, ii) combinations of these

domain architectures, and iii) their frequencies (**Fig. 6C**). These consolidated visualizations can be

explored in our [web app](#).

566   Conclusion

Recent work with the bacterial PSP stress response system suggests that while function is

maintained across phyla, the proteins, regulatory mechanisms, and membrane stress response

mechanisms vary widely among lineages (12, 15, 26, 36) (12, 15, 25, 32). These variations reflect

the utilization of a common peripheral inner membrane protein in lineage-specific envelope

dynamics and responses to unique environmental stresses. In this study, we systematically

analyzed PSP systems, their phyletic spread, and evolution (**Figs. 1** and **2**), discovering several

novel sequence and structural features (*e.g.,* domain architectures) and conserved gene-

neighborhoods of PSP components. We identified both classical and newly-predicted partners

(**Figs. 2** and **5**) and established their phyletic distributions across the tree of life (**Figs. 2** and **6**). We

also established that PspA/Snf7 traces to the LUCA (**Fig. 2**). Indeed, it has been independently

shown that the PspA/Vipp1 ESCRT system dates back to the LUCA (19, 20), further strengthening

the evolutionary significance of this finding. This confirms that the LUCA possessed a membrane

whose curvature and dynamics were mediated by an ancestral PspA/Snf7-like coiled-coil protein

assembled into polymeric structures adjacent to the inner-leaf of the membrane. This hypothesis is

in line with earlier work showing that LUCA contained some signal-recognition particle GTPases

(102). These proteins are unequivocal indicators of the presence of a secretory apparatus

translocating proteins across lipid bilayers. Hence, as a corollary they are also indicators of the

presence of a membrane. Similarly, the extant components of membrane linked bioenergetics

(F1/F0 ATPases or V-Type ATPases) are also indicative of a membrane in the LUCA (103, 104).

Thus, we infer that this ancient machinery was robust to the subsequent lineage-specific changes

in membrane chemistry, producing different membrane types (*e.g.,* ether-linked vs. ester-linked) in

the archaeal, bacterial, and eukaryotic lineages.

Our novel PSP findings make several new functional predictions: 1) Based on its occurrence with

membrane partners (*e.g.,* PspA, LiaI-LiaF-TM, PspC, HAAS, SHOCT-like domains) and transcription

regulators or two-component systems (*e.g.,* LiaRS, PadR-like-wHTH), we propose that Toastrack

is a membrane-proximal intracellular domain serving as a sub-membrane signaling complex

assembly platform; 2) The repeated association of PspA homologs with distinct AAA+-chaperones

and a non-ATPase chaperone (*e.g.,* CesT/Tir) or predicted chaperone-like proteins (*e.g.,* Band-7,

TPM) suggest that these associations are involved in the assembly of sub- and trans-membrane

complexes in response to specific envelope stresses. These observations imply a general role for

analogs of ESCRT-like complexes in these bacterial systems; 3) The association of PspA with

various transporters and polyamine metabolism systems indicates that the regulation of membrane

structure by PspA is associated with changes in solute concentrations and covalent modifications

that may affect membrane stability; 4) The discovery of several new versions of the SHOCT-like

domain suggests that these domains mediate membrane localization of disparate catalytic and non-

catalytic activities, which may subsequently interface with the PSP system; 5) A diverse array of

sub-membrane (*e.g.,* Toastrack), TM, and cell-surface domains (*e.g.,* PspC, LiaI-LiaF-TM) co-occur

with two-component or HAAS-PadR-like-wHTH systems, hinting at alternatives to conventional

two-component signaling that interface with the PSP system in responding to membrane stress.

Further research into these newly described systems will yield broadly applicable mechanistic

insights, given their conservation across swaths of life. For example, SHOCT domains are

understudied due to their believed absence in model organisms (94), but our approach identified

divergent homologs in the model organism *B. subtilis* ([CAB15517.1](#)), which may prompt new

studies. The extensive findings from applying gene neighborhood and domain architecture

searches support their viability, and such methods help reduce the space of annotated hypothetical

proteins without classified domains and add functional clues. Finally, we provide another pillar of

support to recent articles' conclusions that PspA/Snf7 homologs were already present in the last

universal common ancestor (19, 20), and deepen our understanding of the origins and diversification

of all life. All the findings (data, results, and visual summaries) from this work are found in an

interactive and queryable web application available at [jravilab.shinyapps.io/psp_evolution](#).

Methods

Query and subject selection

We queried classical PSP members — PspA (eight representatives across Proteobacteria, Actinobacteria, Firmicutes (two paralogs), Cyanobacteria, and Archaea); PspM (Rv2743c) and PspN (Rv2742c) (from *M. tuberculosis*); PspB and PspC (from *E. coli*); LiaI, LiaG, and LiaF (from *B. subtilis*) — against all sequenced genomes across the tree of life. We ran homology searches against custom databases of ~6500 completed representative or reference genomes with taxonomic lineages or the NCBI non-redundant database (**Tables S1** and **S2**) (105, 106). All PSP homologs are listed in the [web app](#), with representatives in **Tables S3** and **S4**. We obtained the phyletic order (sequence) from NCBI taxonomy and BV-BRC (105–109).

Identification and characterization of protein homologs

We analyzed the classical PSP proteins from the best-studied operons *pspF||ABCDE-G* (from *E. coli*), *liaIHGFSR* (from *B. subtilis*), and *clgRpspAMN* (from *M. tuberculosis*) and their phyletic spread to locate and stratify all PSP homologs. We first resolved these PSP proteins into their constituent domains and used each individual domain as a query for all our phylogenetic analyses to ensure an exhaustive search and identification of all related PSP proteins, including remote homologs. This approach allows us to find homologies beyond those from full-length proteins only. We then performed homology searches for each constituent domain across the tree of life (approximately 6500 representative or reference genomes). We used a combination of protein domain and orthology databases, iterative homology searches, and multiple sequence alignment to detect domains, signal peptides, and transmembrane (TM) regions to help construct the domain architecture of each query PSP protein. The ubiquitous presence of transcription factor HTH domains and histidine kinase-receiver domain two-component systems in bacterial phyla precluded

dedicated searches with these domains; instead, we noted their occurrence in predicted operons

alongside core PSP genes to identify transcriptional regulation partners.

We ensured the identification of a comprehensive set of homologs (close and remote) for each

queried protein via iterative searches using PSI-BLAST (110) with sequences of both full-length

proteins and the corresponding constituent domains. More distant relationships were determined

using profile-profile searches with the HHpred program (111). Protein searches were conducted

using homologous copies from multiple species as starting points. We aggregated search results

and recorded the numbers of homologs per species and genomes carrying each of the query

proteins ([web app](#)). These proteins were clustered into orthologous families using the similarity-

based clustering program BLASTCLUST (112). SignalP, TMHMM, Phobius, JPred, Pfam, and

custom profile databases (113–118) were used to identify signal peptides, TM regions, known

domains, and the secondary protein structures in every genome. Homolog information, including

domain architectures, is available in the [web app](#) ('Data,' 'Domain architectures' tabs).

Neighborhood search

Bacterial gene neighborhoods (±7 genes flanking each protein homolog) were retrieved using a Perl

script from GenBank (105, 106). Gene orientation, domains, and secondary structures of the

neighboring proteins were characterized using the same methods applied to query homologs

above. Genomic contexts are available in the [web app](#) ('Genomic contexts' tab). We note that

eukaryotic components (*e.g.,* Snf7) appearing alone are an artifact of the genomic context analysis

being restricted (and relevant) to bacteria and archaea (see [web app](#)).

Phylogenetic analysis

Multiple sequence alignment (MSA) of the identified homologs was performed using Kalign (119)

and MUSCLE (120) with default parameters. Domain/gene trees were constructed with FastTree

2.1 with default parameters (121) and visualized and labels colored in FigTree v1.4.4 (122). In

parallel, the PspA/Snf7 MSA was also passed to ModelTest-NG, which reported the best

performing substitution model (LG+I+G4+F) (123, 124), and then RAxML-NG-MPI (v1.2.0) (125) on

CU's Alpine HPC cluster and a maximum likelihood phylogenetic tree constructed (--all, –model

LG+I+G4+F, –tree rand{25} pars{25}, –bs-trees autoMRE, –threads 60). Bootstrapping was

performed until RAxML-NG assessed statistical convergence at 480 replicates [**Fig. S2**]. MSA and

phylogenetic trees are available in the web app ('Phylogeny' tab).

Network reconstruction

The PSP proximal neighborhood network was constructed on the domain architectures and

genomic contexts of PspA and its cognate partner proteins (**Tables S1** and **S2**). The nodes

represented the domains, and edges indicated a shared neighborhood (domains of the same

protein or neighboring proteins). Proximity networks are available in the web app ('Domain

architectures' tab).

Web application

We built our interactive and searchable web application using R Shiny (126). All data analyses and

visualizations were carried out using R or RStudio (127, 128) with R-packages (129–141). Data and

results from our study are available on the web application (jravilab.shinyapps.io/psp_evolution).

The web app can be queried with the protein accession numbers, domain names, or species cited

inline throughout the Results section.

Figures

Figure 1. The phyletic spread of classical Psp members across all major lineages.

A. The three classical PSP systems in *E. coli* (*pspF‖pspABC*), *M. tuberculosis* (*clgRpspAMN*), and *B. subtilis* (*lialHGFSR*) are shown. Proteins across three bacterial PSP systems are labeled and colors uniformly (across all figures that show domain architectures) indicate the nature of the protein; **black**, PspA homolog (*e.g.,* PspA, LiaH); **teal**, transcription factor/response regulator (*e.g.,* PspF, LiaR, MprA, ClgR) along with partner histidine kinases in **olive green** (*e.g.,* LiaS, MprB part of two component systems LiaRS, MprAB); **orange/yellow**, transmembrane protein (*e.g.,* PspB, PspC, LiaF, LiaI, PspM) embedded in the dashed blue membrane. Arrows represent interaction or activation, or inhibitory feedback in the case of LiaF and LiaS. A simple operon map is shown for each system at the bottom, but is expanded upon in the panel **B**. **B. Domain architectures of the classical Psp operons** in *E. coli*, *M. tuberculosis*, and *B. subtilis*. Domains are denoted by rectangular segments inside the block arrow representing each protein labeled below the arrow (*e.g.,* the protein LiaH contains only the PspA domain). The direction of the arrow indicates the direction of transcription. See *Results* for new domain definitions. **C. Phyletic spreads of Psp proteins.** Sunburst plots are shown to indicate the lineage distributions (as a fraction of the total number of homologs recorded across lineages) for the homologs of 'classical' domains/protein families of interest: PspA, Snf7, PspB, PspC, PspM, PspN (and DUF3046), LiaI-LiaF-TM, Toastrack. In each plot corresponding to a particular protein, the inner ring corresponds to the proportion of its homologs present in superkingdoms/domains of life, Bacteria, Archaea, and Eukaryota. The outer ring depicts the distribution of the homologs among key phyla. Interactive sunburst plots for each of the PSP proteins are available in the [web app](). The colors for each lineage (outer and inner rings) are shown separately for **C**.

Figure 2. PspA/Snf7 homologs across the tree of life.

Phylogenetic tree of PspA homologs across the tree of life. The phylogenetic tree was

constructed using FastTree and visualized with FigTree (parallel RAxML-NG generated PspA tree

with confidence values is shown in **Fig. S2**) from a multiple sequence alignment (using kalign3) of

representative PspA homologs across all major superkingdoms and phyla (see *Methods*; web app).

The key lineages are labeled next to distinct clusters of similar PspA proteins; leaf colors match the

lineage labels. The insets show the 3D structures for PspA (4WHE) and Snf7 (5FD7) from the Protein

Data Bank. The tree leaves are labeled by lineage, species, and accession numbers. A text-

searchable, scaled vector graphics version of this tree is available through the web app under the

Phylogeny tab. Representatives for PspA/Snf7 homologs are also available in **Table S3**. As a gene

tree, most homologs do cluster among evolutionarily similar species, though evidence of

widespread horizontal gene transfer is also evident. Two clusters of eukaryotic genes appear: Snf7

in Eukaryota and Asgardarchaeota group, mirroring recent research suggesting eukaryotic life arose

from this clade; and Vipp1 among Eukaryota and Cyanobacteria group, again reflecting the shared

origin of the protein in plants from Cyanobacteria.

Figure 3. PspA domain architectures and genomic contexts

The first row contains domain architectures of the most prevalent homologs of PspA (as in **1A**;

indicated in ***black*** throughout). The remaining rows show the predominant genomic contexts of

PspA homologs across multiple bacterial and archaeal lineages (identified by neighborhood

searches ±7 genes flanking each homolog; see *Methods*). Representative neighborhoods with

archetypal lineages and archetypal example proteins (with accession numbers and species) are

shown. The PspA contexts are grouped by neighboring domains such as (a) PspF, PspB/PspC; (b)

PspAA, PspAB; (c) ClgR, PspM/PspN, Thioredoxin (**Fig. S1**); (d) chaperones such as Band-7,

Flotillin, CesT_Tir, TPM_phosphatase, ZnR, SpermGS-ATPgrasp, and Spermine synthase (**Table**

S2); (e) two-component systems such as the Lia system and Toastrack, and other novel genomic

contexts; and (f) cyanobacterial variations. Key: *rectangles*, domains; *arrowheads*, the direction of

transcription; *domains enclosed by dotted lines*, absent in the genomic contexts in certain species;

white cross, substitution with protein(s) mentioned just below; *white triangle*, insertion with one or

more proteins; *two vertical lines* '‖', indicate a change in the direction of transcription; *small black*

boxes, domain repeats co-occurring within a protein/context. Archetypal accession numbers and

species are provided mostly on the left. Archetypal lineages are indicated in grey on the right for

each of the domain architectures and genomic contexts. Different domains are indicated by the

same color scheme as in **Figure 1**. Also, similar domains are given similar hues. For example,

membrane proteins are in different shades of orange (SIG, predicted signal peptide, dark orange,

PspC, orange, other transmembrane domain, light orange); transcription factors/regulators

(including HTH, helix-turn-helix domain) are in teal; DUFs, Domains of Unknown Function, and other

domains are in grey. Further details of the domain architectures and gene neighborhoods shown

are described in the text and **Table S3**, and the full list of PspA homologs, their domain

architectures, genomic contexts, and lineages are shown in the web app (under the 'Data,' 'Domain

architectures,' and 'Genomic contexts' tabs).

Figure 4. Lineage spread of PspA-free domain architectures.

The domain architectures of the most prevalent homologs of PspA partner domains (frequency of

occurrence >50 across lineages), including classical (Toastrack, LiaI-LiaF-TM, PspBC, PspMN,

DUF3046) and other novel neighbors (PspAA, PspAB, HAAS, SHOCT-bihelical, SHOCT-like, AAA[+]-

ATPase domains) are illustrated on the left. Broad distributions of domains across the tree of life

are indicated in **Fig. 1C**. The phyletic spread of the underlying domain architectures is shown here

along with their relative frequencies as a stacked barplot for all superkingdoms/domains of life (sub-

lineages or phyla are sorted; *e.g.,* all bacterial lineages appear together). Further details of the

domain architectures of all PspA partner domain homologs and their phyletic spreads are shown in

the web app, with representatives shown in **Table S4**.

Domain Architectures　　　**Number of Occurrences per Lineage**

Figure 5. The genomic contexts housing all Psp cognate partner domain homologs.

The genomic contexts and key lineage memberships are presented using the same schematic as

in **Figure 3**. The focus is on PSP partner domains such as Toastrack (blue), PspC, LiaI-LiaF-TM,

HAAS, SHOCT-bi-helical (in shades of orange since they are transmembrane domains), and the

various genomic neighborhoods with SHOCT-like proteins, transcription regulators (*e.g.,* PadR-

wHTH, SIGMA-HTH, GNTR-HTH), and two-component systems (**Table S2**). Further details of the

genomic contexts of all PspA-free partner domain homologs and their phyletic spreads are in the

web app, and representatives indicated in the figure are shown in **Table S4**.

Figure 6. PSP consolidated.

A. Domain proximity network. The network captures co-occurring domains within the top 97% of

the homologs of all the 'query' Psp members and their key partner domains (after sorting by

decreasing frequency of occurrence). The size of the nodes (domains) and width of edges (co-

occurrence of domains within a protein) are proportional to the frequency of their occurrence across

homologs. The query domains (original proteins/domains of interest; **Table S1**) and other commonly

co-occurring domains (**Table S2**) are *red* or *grey*. *Note*: A few connections may be absent in the

displayed network due to low occurrence (fewer than the threshold), *e.g.,* PspA and PspAA,

Betapropeller, and AAA-ATPase. The full network, and the domain-centric ones, can be accessed

on the [web app](). **B. Phyletic spread of the most common domains.** The heatmap shows the

presence/absence of homologs of PspA and partner domains across lineages. The color gradient

represents the number of homologs identified within each lineage (*e.g.,* the darkest shade indicates

the highest number of homologs in a particular lineage). *Rows*: Psp members and their most

frequent partners are queried against all sequenced and completed genomes across Bacteria,

Eukaryota, and Archaea. *Columns*: The major archaeal (*green*), bacterial (*orange*), eukaryotic

(*blue*), and viral (*grey*) lineages with representative sequenced genomes are considered. Details of

all homologs across the tree of life, their domain architectures, genomic contexts, and their lineage

distributions are shown in the [web app]() (representatives in **Table S3, S4**). **C. Predominant co-**

occurring domain architectures in genomic neighborhoods. UpSet plot of the most common

neighboring proteins (genomic contexts >100 occurrences are shown) among all Psp homologs.

Blue histogram: Distribution of the predominant domain architectures. *Dots and connections*:

Combinations in which these domain architectures come together in the genomic neighborhoods.

Red histogram: Frequency of occurrences of genomic neighborhoods comprising specific

combinations of the predominant domain architectures. Phyletic spreads and UpSet plots of the

domain architectures and genomic contexts for the homologs of all Psp member proteins are

available in the web app.

808 Declarations

809 Acknowledgments

We are incredibly grateful to Arjun Krishnan and Krishnan Raghunathan for detailed comments on

the manuscript and web application, guidance, and support; Emily Meyer and Jerome McKay for

critical edits of the manuscript. We also thank Srinand Sreevatsan, Beronda Montgomery, Deborah

Johnson, and the MSU Diversity Research Network for support and writing spaces. Finally, we

thank members of the JRaviLab for feedback on the web application, especially Jake Krol, for help

with the final deployment. This work utilized the Alpine high-performance computing resource at

the University of Colorado Boulder. Alpine is jointly funded by the University of Colorado Boulder,

the University of Colorado Anschutz, Colorado State University, and the National Science

Foundation (award: 2201538).

820 Funding

827 Author Contributions

J.R. and M.L.G. conceived the study; J.R., V.A., L.A., and M.L.G. designed the study; J.R. and V.A

acquired the data, performed all the analyses, and made the figures and tables. J.R., V.A., L.A., and

M.L.G. interpreted the results and wrote the manuscript. S.Z.C built the web application with all the

results (data summarization and visualization) with J.R.; S.Z.C. also contributed to making figures

and linking identifiers in the manuscript to reference databases. E.P.B. contributed to writing the

manuscript, performed phylogenetic tree construction, and mapped Pfam to Interpro domain IDs

throughout the manuscript. P.D. contributed to renaming some domains.

Data Availability and Reuse

All the data, analyses, and visualizations are available in our interactive and queryable web

application: https://jravilab.shinyapps.io/psp_evolution. Text, figures, and the web app are licensed

under Creative Commons Attribution CC BY 4.0.

References

1. van den Brink-van der Laan E, Killian JA, de Kruijff B. 2004. Nonbilayer lipids affect peripheral
and integral membrane proteins via changes in the lateral pressure profile. Biochim Biophys
Acta 1666:275–288.
2. Kobayashi R, Suzuki T, Yoshida M. 2007. Escherichia coli phage-shock protein A (PspA) binds
to membrane phospholipids and repairs proton leakage of the damaged membranes. Mol
Microbiol 66:100–109.
3. Jordan S, Hutchings MI, Mascher T. 2008. Cell envelope stress response in Gram-positive
bacteria. FEMS Microbiol Rev 32:107–146.
4. Raivio TL. 2005. Envelope stress responses and Gram-negative bacterial pathogenesis. Mol
Microbiol 56:1119–1128.
5. Rowley G, Spector M, Kormanec J, Roberts M. 2006. Pushing the envelope: extracytoplasmic
stress responses in bacterial pathogens. Nat Rev Microbiol 4:383–394.
6. Omardien S, Drijfhout JW, van Veen H, Schachtschabel S, Riool M, Hamoen LW, Brul S, Zaat
SAJ. 2018. Synthetic antimicrobial peptides delocalize membrane bound proteins thereby
inducing a cell envelope stress response. Biochim Biophys Acta Biomembr 1860:2416–2427.
7. Wenzel M, Kohl B, Münch D, Raatschen N, Albada HB, Hamoen L, Metzler-Nolte N, Sahl H-G,
Bandow JE. 2012. Proteomic response of Bacillus subtilis to lantibiotics reflects differences in
interaction with the cytoplasmic membrane. Antimicrob Agents Chemother 56:5749–5757.
8. Guest RL, Wang J, Wong JL, Raivio TL. 2017. A Bacterial Stress Response Regulates
Respiratory Protein Complexes To Control Envelope Stress Adaptation. J Bacteriol 199.
9. Gottesman S. 2017. Stress Reduction, Bacterial Style. J Bacteriol 199.
10. Brissette JL, Weiner L, Ripmaster TL, Model P. 1991. Characterization and sequence of the
Escherichia coli stress-induced psp operon. J Mol Biol 220:35–48.
11. Mascher T, Zimmer SL, Smith T-A, Helmann JD. 2004. Antibiotic-inducible promoter regulated
by the cell envelope stress-sensing two-component system LiaRS of Bacillus subtilis.
Antimicrob Agents Chemother 48:2888–2896.
12. Huvet M, Toni T, Sheng X, Thorne T, Jovanovic G, Engl C, Buck M, Pinney JW, Stumpf MPH.
2011. The evolution of the phage shock protein response system: interplay between protein
function, genomic organization, and system function. Mol Biol Evol 28:1141–1155.
13. McDonald C, Jovanovic G, Ces O, Buck M. 2015. Membrane Stored Curvature Elastic Stress
Modulates Recruitment of Maintenance Proteins PspA and Vipp1. mBio 6:e01188-01115.
14. Datta P, Ravi J, Guerrini V, Chauhan R, Neiditch MB, Shell SS, Fortune SM, Hancioglu B,
Igoshin OA, Gennaro ML. 2015. The Psp system of Mycobacterium tuberculosis integrates
envelope stress-sensing and envelope-preserving functions. Mol Microbiol 97:408–422.
15. Ravi J, Anantharaman V, Aravind L, Gennaro ML. 2018. Variations on a theme: evolution of the
phage-shock-protein system in Actinobacteria. Antonie Van Leeuwenhoek 111:753–760.
16. Heidrich J, Thurotte A, Schneider D. 2017. Specific interaction of IM30/Vipp1 with
cyanobacterial and chloroplast membranes results in membrane remodeling and eventually in
membrane fusion. Biochim Biophys Acta Biomembr 1859:537–549.
17. McDonald C, Jovanovic G, Wallace BA, Ces O, Buck M. 2017. Structure and function of PspA
and Vipp1 N-terminal peptides: Insights into the membrane stress sensing and mitigation.
Biochim Biophys Acta Biomembr 1859:28–39.

18. Lee I-H, Kai H, Carlson L-A, Groves JT, Hurley JH. 2015. Negative membrane curvature
catalyzes nucleation of endosomal sorting complex required for transport (ESCRT)-III assembly.
Proc Natl Acad Sci U S A 112:15892–15897.

19. Liu J, Tassinari M, Souza DP, Naskar S, Noel JK, Bohuszewicz O, Buck M, Williams TA, Baum
B, Low HH. 2021. Bacterial Vipp1 and PspA are members of the ancient ESCRT-III membrane-
remodeling superfamily. Cell 184:3660-3673.e18.

20. Junglas B, Huber ST, Heidler T, Schlösser L, Mann D, Hennig R, Clarke M, Hellmann N,
Schneider D, Sachse C. 2021. PspA adopts an ESCRT-III-like fold and remodels bacterial
membranes. Cell 184:3674-3688.e18.

21. Horinouchi T, Suzuki S, Hirasawa T, Ono N, Yomo T, Shimizu H, Furusawa C. 2015. Phenotypic
convergence in bacterial adaptive evolution to ethanol stress. BMC Evol Biol 15:180.

22. Furusawa C, Irie N. 2020. Toward understanding of evolutionary constraints: experimental and
theoretical approaches. Biophys Rev 12:1155–1161.

23. Kleerebezem M, Crielaard W, Tommassen J. 1996. Involvement of stress protein PspA (phage
shock protein A) of Escherichia coli in maintenance of the protonmotive force under stress
conditions. EMBO J 15:162–171.

24. Huvet M, Toni T, Tan H, Jovanovic G, Engl C, Buck M, Stumpf MPH. 2009. Model-based
evolutionary analysis: the natural history of phage-shock stress response. Biochem Soc Trans
37:762–767.

25. Vothknecht UC, Otters S, Hennig R, Schneider D. 2012. Vipp1: a very important protein in
plastids?! J Exp Bot 63:1699–1712.

26. Bultema JB, Fuhrmann E, Boekema EJ, Schneider D. 2010. Vipp1 and PspA: Related but not
twins. Commun Integr Biol 3:162–165.

27. Westphal S, Heins L, Soll J, Vothknecht UC. 2001. Vipp1 deletion mutant of Synechocystis: a
connection between bacterial phage shock and thylakoid biogenesis? Proc Natl Acad Sci U S A
98:4243–4248.

28. Heidrich J, Wulf V, Hennig R, Saur M, Markl J, Sönnichsen C, Schneider D. 2016. Organization
into Higher Ordered Ring Structures Counteracts Membrane Binding of IM30, a Protein
Associated with Inner Membranes in Chloroplasts and Cyanobacteria. J Biol Chem 291:14954–
14962.

29. Theis J, Gupta TK, Klingler J, Wan W, Albert S, Keller S, Engel BD, Schroda M. 2019. VIPP1
rods engulf membranes containing phosphatidylinositol phosphates. 1. Sci Rep 9:8725.

30. Zhang L, Kato Y, Otters S, Vothknecht UC, Sakamoto W. 2012. Essential role of VIPP1 in
chloroplast envelope maintenance in Arabidopsis. Plant Cell 24:3695–3707.

31. Armstrong RM, Adams KL, Zilisch JE, Bretl DJ, Sato H, Anderson DM, Zahrt TC. 2016. Rv2744c
Is a PspA Ortholog That Regulates Lipid Droplet Homeostasis and Nonreplicating Persistence in
Mycobacterium tuberculosis. J Bacteriol 198:1645–1661.

32. Gupta TK, Klumpe S, Gries K, Heinz S, Wietrzynski W, Ohnishi N, Niemeyer J, Spaniol B,
Schaffer M, Rast A, Ostermeier M, Strauss M, Plitzko JM, Baumeister W, Rudack T, Sakamoto
W, Nickelsen J, Schuller JM, Schroda M, Engel BD. 2021. Structural basis for VIPP1
oligomerization and maintenance of thylakoid membrane integrity. Cell 184:3643-3659.e23.

33. Joly N, Engl C, Jovanovic G, Huvet M, Toni T, Sheng X, Stumpf MPH, Buck M. 2010. Managing
membrane stress: the phage shock protein (Psp) response, from molecular mechanisms to
physiology. FEMS Microbiol Rev 34:797–827.

34. Eiamphungporn W, Helmann JD. 2008. The Bacillus subtilis sigma(M) regulon and its
contribution to cell envelope stress responses. Mol Microbiol 67:830–848.
35. Jordan S, Junker A, Helmann JD, Mascher T. 2006. Regulation of LiaRS-dependent gene
expression in bacillus subtilis: identification of inhibitor proteins, regulator binding sites, and
target genes of a conserved cell envelope stress-sensing two-component system. J Bacteriol
188:5153–5166.
36. Manganelli R, Gennaro ML. 2017. Protecting from Envelope Stress: Variations on the Phage-
Shock-Protein Theme. Trends Microbiol 25:205–216.
37. Popp PF, Gumerov VM, Andrianova EP, Bewersdorf L, Mascher T, Zhulin IB, Wolf D. 2022.
Phyletic Distribution and Diversification of the Phage Shock Protein Stress Response System in
Bacteria and Archaea. mSystems 7:e01348-21.
38. van der Laan M, Urbanus ML, Ten Hagen-Jongman CM, Nouwen N, Oudega B, Harms N,
Driessen AJM, Luirink J. 2003. A conserved function of YidC in the biogenesis of respiratory
chain complexes. Proc Natl Acad Sci U S A 100:5801–5806.
39. Maxson ME, Darwin AJ. 2006. Multiple promoters control expression of the Yersinia
enterocolitica phage-shock-protein A (pspA) operon. Microbiol Read Engl 152:1001–1010.
40. Weiner L, Model P. 1994. Role of an Escherichia coli stress-response operon in stationary-
phase survival. Proc Natl Acad Sci U S A 91:2191–2195.
41. Jovanovic G, Lloyd LJ, Stumpf MPH, Mayhew AJ, Buck M. 2006. Induction and function of the
phage shock protein extracytoplasmic stress response in Escherichia coli. J Biol Chem
281:21147–21161.
42. Mitchell AM, Silhavy TJ. 2019. Envelope stress responses: balancing damage repair and
toxicity. Nat Rev Microbiol 17:417–428.
43. Engl C, Beek AT, Bekker M, de Mattos JT, Jovanovic G, Buck M. 2011. Dissipation of proton
motive force is not sufficient to induce the phage shock protein response in Escherichia coli.
Curr Microbiol 62:1374–1385.
44. Weiner L, Brissette JL, Model P. 1991. Stress-induced expression of the Escherichia coli phage
shock protein operon is dependent on sigma 54 and modulated by positive and negative
feedback mechanisms. Genes Dev 5:1912–1923.
45. Jovanovic G, Engl C, Buck M. 2009. Physical, functional and conditional interactions between
ArcAB and phage shock proteins upon secretin-induced stress in Escherichia coli. Mol
Microbiol 74:16–28.
46. Brissette JL, Russel M, Weiner L, Model P. 1990. Phage shock protein, a stress protein of
Escherichia coli. Proc Natl Acad Sci U S A 87:862–866.
47. Yamaguchi S, Reid DA, Rothenberg E, Darwin AJ. 2013. Changes in Psp protein binding
partners, localization and behaviour upon activation of the Yersinia enterocolitica phage shock
protein response. Mol Microbiol 87:656–671.
48. Wolf D, Kalamorz F, Wecke T, Juszczak A, Mäder U, Homuth G, Jordan S, Kirstein J, Hoppert
M, Voigt B, Hecker M, Mascher T. 2010. In-depth profiling of the LiaR response of Bacillus
subtilis. J Bacteriol 192:4680–4693.
49. Jordan S, Rietkötter E, Strauch MA, Kalamorz F, Butcher BG, Helmann JD, Mascher T. 2007.
LiaRS-dependent gene expression is embedded in transition state regulation in Bacillus subtilis.
Microbiol Read Engl 153:2530–2540.
50. Castelle CJ, Banfield JF. 2018. Major New Microbial Groups Expand Diversity and Alter our
Understanding of the Tree of Life. Cell 172:1181–1197.

51. Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN,
Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman DA, Finstad KM, Amundson R,
Thomas BC, Banfield JF. 2016. A new view of the tree of life. Nat Microbiol 1:16048.

52. Flores-Kim J, Darwin AJ. 2016. The Phage Shock Protein Response. Annu Rev Microbiol 70:83–
101.

53. Darwin AJ, Miller VL. 2001. The psp locus of Yersinia enterocolitica is required for virulence and
for growth in vitro when the Ysc type III secretion system is produced. Mol Microbiol 39:429–
444.

54. Becker LA, Bang I-S, Crouch M-L, Fang FC. 2005. Compensatory role of PspA, a member of
the phage shock protein operon, in rpoE mutant Salmonella enterica serovar Typhimurium. Mol
Microbiol 56:1004–1016.

55. Elderkin S, Bordes P, Jones S, Rappas M, Buck M. 2005. Molecular determinants for PspA-
mediated repression of the AAA transcriptional activator PspF. J Bacteriol 187:3238–3248.

56. Suntharalingam P, Senadheera MD, Mair RW, Lévesque CM, Cvitkovitch DG. 2009. The LiaFSR
system regulates the cell envelope stress response in Streptococcus mutans. J Bacteriol
191:2973–2984.

57. Aravind L, Ponting CP. 1999. The cytoplasmic helical linker domain of receptor histidine kinase
and methyl-accepting proteins is common to many prokaryotic signalling proteins. FEMS
Microbiol Lett 176:111–116.

58. Maxson ME, Darwin AJ. 2006. PspB and PspC of Yersinia enterocolitica are dual function
proteins: regulators and effectors of the phage-shock-protein response. Mol Microbiol 59:1610–
1623.

59. Gueguen E, Savitzky DC, Darwin AJ. 2009. Analysis of the Yersinia enterocolitica PspBC
proteins defines functional domains, essential amino acids and new roles within the phage-
shock-protein response. Mol Microbiol 74:619–633.

60. Wallrodt I, Jelsbak L, Thomsen LE, Brix L, Lemire S, Gautier L, Nielsen DS, Jovanovic G, Buck
M, Olsen JE. 2014. Removal of the phage-shock protein PspB causes reduction of virulence in
Salmonella enterica serovar Typhimurium independently of NRAMP1. J Med Microbiol 63:788–
795.

61. Kleine B, Chattopadhyay A, Polen T, Pinto D, Mascher T, Bott M, Brocker M, Freudl R. 2017.
The three-component system EsrISR regulates a cell envelope stress response in
Corynebacterium glutamicum. Mol Microbiol 106:719–741.

62. Srivastava D, Moumene A, Flores-Kim J, Darwin AJ. 2017. Psp Stress Response Proteins Form
a Complex with Mislocalized Secretins in the Yersinia enterocolitica Cytoplasmic Membrane.
mBio 8.

63. Flores-Kim J, Darwin AJ. 2012. Phage shock protein C (PspC) of Yersinia enterocolitica is a
polytopic membrane protein with implications for regulation of the Psp stress response. J
Bacteriol 194:6548–6559.

64. Datta P, Ravi J, Guerrini V, Chauhan R, Neiditch MB, Shell SS, Fortune SM, Hancioglu B,
Igoshin OA, Gennaro ML. 2015. The Psp system of Mycobacterium tuberculosis integrates
envelope stress-sensing and envelope-preserving functions. Mol Microbiol 97:408–422.

65. Burki F, Shalchian-Tabrizi K, Pawlowski J. 2008. Phylogenomics reveals a new "megagroup"
including most photosynthetic eukaryotes. Biol Lett 4:366–369.

66.   Hampl V, Hug L, Leigh JW, Dacks JB, Lang BF, Simpson AGB, Roger AJ. 2009. Phylogenomic
1016   analyses support the monophyly of Excavata and resolve relationships among eukaryotic
1017   "supergroups." Proc Natl Acad Sci U S A 106:3859–3864.
1018   67.   Caillat C, Maity S, Miguet N, Roos WH, Weissenhorn W. 2019. The role of VPS4 in ESCRT-III
1019   polymer remodeling. Biochem Soc Trans 47:441–448.
1020   68.   Schöneberg J, Pavlin MR, Yan S, Righini M, Lee I-H, Carlson L-A, Bahrami AH, Goldman DH,
1021   Ren X, Hummer G, Bustamante C, Hurley JH. 2018. ATP-dependent force generation and
1022   membrane scission by ESCRT-III and Vps4. Science 362:1423–1428.
1023   69.   Weiss P, Huppert S, Kölling R. 2009. Analysis of the dual function of the ESCRT-III protein Snf7
1024   in endocytic trafficking and in gene expression. Biochem J 424:89–97.
1025   70.   Godinho RM da C, Crestani J, Kmetzsch L, Araujo G de S, Frases S, Staats CC, Schrank A,
1026   Vainstein MH, Rodrigues ML. 2014. The vacuolar-sorting protein Snf7 is required for export of
1027   virulence determinants in members of the Cryptococcus neoformans complex. Sci Rep 4:6198.
1028   71.   Webster BM, Colombi P, Jäger J, Lusk CP. 2014. Surveillance of nuclear pore complex
1029   assembly by ESCRT-III/Vps4. Cell 159:388–401.
1030   72.   Zhang L, Kondo H, Kamikubo H, Kataoka M, Sakamoto W. 2016. VIPP1 Has a Disordered C-
1031   Terminal Tail Necessary for Protecting Photosynthetic Membranes against Stress. Plant Physiol
1032   171:1983–1995.
1033   73.   Anantharaman V, Aravind L. 2003. Evolutionary history, structural features and biochemical
1034   diversity of the NlpC/P60 superfamily of enzymes. Genome Biol 4:R11.
1035   74.   Brosch R, Gordon SV, Marmiesse M, Brodin P, Buchrieser C, Eiglmeier K, Garnier T, Gutierrez
1036   C, Hewinson G, Kremer K, Parsons LM, Pym AS, Samper S, van Soolingen D, Cole ST. 2002. A
1037   new evolutionary scenario for the Mycobacterium tuberculosis complex. Proc Natl Acad Sci U S
1038   A 99:3684–3689.
1039   75.   Tang S, Henne WM, Borbat PP, Buchkovich NJ, Freed JH, Mao Y, Fromme JC, Emr SD. 2015.
1040   Structural basis for activation, assembly and membrane binding of ESCRT-III Snf7 filaments.
1041   eLife 4.
1042   76.   Anantharaman V, Koonin EV, Aravind L. 2001. Regulatory potential, phyletic distribution and
1043   evolution of ancient, intracellular small-molecule-binding domains. J Mol Biol 307:1271–1292.
1044   77.   Kaur G, Burroughs AM, Iyer LM, Aravind L. 2020. Highly regulated, diversifying NTP-dependent
1045   biological conflict systems with implications for the emergence of multicellularity. eLife 9.
1046   78.   Tsai JT, Sung N, Lee J, Chang C, Lee S, Tsai FTF. 2019. Crystal Structure of the YcjX Stress
1047   Protein Reveals a Ras-Like GTP-Binding Protein. J Mol Biol 431:3179–3190.
1048   79.   Wollert T, Wunder C, Lippincott-Schwartz J, Hurley JH. 2009. Membrane scission by the
1049   ESCRT-III complex. Nature 458:172–177.
1050   80.   Frohn BP, Härtel T, Cox J, Schwille P. 2022. Tracing back variations in archaeal ESCRT-based
1051   cell division to protein domain architectures. PloS One 17:e0266395.
1052   81.   Zaremba-Niedzwiedzka K, Caceres EF, Saw JH, Bäckström D, Juzokaite L, Vancaester E, Seitz
1053   KW, Anantharaman K, Starnawski P, Kjeldsen KU, Stott MB, Nunoura T, Banfield JF, Schramm
1054   A, Baker BJ, Spang A, Ettema TJG. 2017. Asgard archaea illuminate the origin of eukaryotic
1055   cellular complexity. Nature 541:353–358.
1056   82.   Burroughs AM, Aravind L. 2016. RNA damage in biological conflicts and the diversity of
1057   responding RNA repair systems. Nucleic Acids Res 44:8525–8555.
1058   83.   Gehl B, Sweetlove LJ. 2014. Mitochondrial Band-7 family proteins: scaffolds for respiratory
1059   chain assembly? Front Plant Sci 5:141.

84. Delahay RM, Shaw RK, Elliott SJ, Kaper JB, Knutton S, Frankel G. 2002. Functional analysis of
the enteropathogenic Escherichia coli type III secretion system chaperone CesT identifies
domains that mediate substrate interactions. Mol Microbiol 43:61–73.

85. Jovanovic G, Mehta P, McDonald C, Davidson AC, Uzdavinys P, Ying L, Buck M. 2014. The N-
terminal amphipathic helices determine regulatory and effector functions of phage shock protein
A (PspA) in Escherichia coli. J Mol Biol 426:1498–1511.

86. Lee M-J, Huang C-Y, Sun Y-J, Huang H. 2005. Cloning and characterization of spermidine
synthase and its implication in polyamine biosynthesis in Helicobacter pylori strain 26695.
Protein Expr Purif 43:140–148.

87. Jovanovic G, Mehta P, Ying L, Buck M. 2014. Anionic lipids and the cytoskeletal proteins MreB
and RodZ define the spatio-temporal distribution and function of membrane stress controller
PspA in Escherichia coli. Microbiol Read Engl 160:2374–2386.

88. Cerutti ML, Otero LH, Smal C, Pellizza L, Goldbaum FA, Klinke S, Aran M. 2017. Structural and
functional characterization of a cold adapted TPM-domain with ATPase/ADPase activity. J
Struct Biol 197:201–209.

89. Wegener KM, Bennewitz S, Oelmüller R, Pakrasi HB. 2011. The Psb32 protein aids in repairing
photodamaged photosystem II in the cyanobacterium Synechocystis 6803. Mol Plant 4:1052–
1061.

90. Sirpiö S, Allahverdiyeva Y, Suorsa M, Paakkarinen V, Vainonen J, Battchikova N, Aro E-M. 2007.
TLP18.3, a novel thylakoid lumen protein regulating photosystem II repair cycle. Biochem J
406:415–425.

91. Ballas SK, Mohandas N, Marton LJ, Shohet SB. 1983. Stabilization of erythrocyte membranes
by polyamines. Proc Natl Acad Sci U S A 80:1942–1946.

92. Li B, Kurihara S, Kim SH, Liang J, Michael AJ. 2019. A polyamine-independent role for S-
adenosylmethionine decarboxylase. Biochem J 476:2579–2594.

93. Fiuza M, Letek M, Leiba J, Villadangos AF, Vaquera J, Zanella-Cléon I, Mateos LM, Molle V, Gil
JA. 2010. Phosphorylation of a novel cytoskeletal protein (RsmP) regulates rod-shaped
morphology in Corynebacterium glutamicum. J Biol Chem 285:29387–29397.

94. Eberhardt RY, Bartholdson SJ, Punta M, Bateman A. 2013. The SHOCT domain: a widespread
domain under-represented in model organisms. PloS One 8:e57848.

95. Kingston AW, Liao X, Helmann JD. 2013. Contributions of the σ(W) , σ(M) and σ(X) regulons to
the lantibiotic resistome of Bacillus subtilis. Mol Microbiol 90:502–518.

96. Miller C, Kong J, Tran TT, Arias CA, Saxer G, Shamoo Y. 2013. Adaptation of Enterococcus
faecalis to daptomycin reveals an ordered progression to resistance. Antimicrob Agents
Chemother 57:5373–5383.

97. Wang L, Xiao S, Chen X, Chen S, Wang S, Wang C, Tang Y, Dong F. 2019. ytiB and ythA Genes
Reduce the Uranium Removal Capacity of Bacillus atrophaeus. Int J Mol Sci 20.

98. Boyle-Vavra S, Yin S, Jo DS, Montgomery CP, Daum RS. 2013. VraT/YvqF is required for
methicillin resistance and activation of the VraSR regulon in Staphylococcus aureus. Antimicrob
Agents Chemother 57:83–95.

99. Gardete S, Wu SW, Gill S, Tomasz A. 2006. Role of VraSR in antibiotic resistance and antibiotic-
induced stress response in Staphylococcus aureus. Antimicrob Agents Chemother 50:3424–
3434.

100. Wolf D, Domínguez-Cuevas P, Daniel RA, Mascher T. 2012. Cell envelope stress response in
cell wall-deficient L-forms of Bacillus subtilis. Antimicrob Agents Chemother 56:5907–5915.

101. Hauf S, Möller L, Fuchs S, Halbedel S. 2019. PadR-type repressors controlling production of a
non-canonical FtsW/RodA homologue and other trans-membrane proteins. Sci Rep 9:10023.
102. Leipe DD, Wolf YI, Koonin EV, Aravind L. 2002. Classification and evolution of P-loop GTPases
and related ATPases. J Mol Biol 317:41–72.
103. Lane N, Martin WF. 2012. The origin of membrane bioenergetics. Cell 151:1406–1416.
104. Sojo V, Pomiankowski A, Lane N. 2014. A bioenergetic basis for membrane divergence in
archaea and bacteria. PLoS Biol 12:e1001926.
105. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, Rajput B, Robbertse B,
Smith-White B, Ako-Adjei D, Astashyn A, Badretdin A, Bao Y, Blinkova O, Brover V, Chetvernin
V, Choi J, Cox E, Ermolaeva O, Farrell CM, Goldfarb T, Gupta T, Haft D, Hatcher E, Hlavina W,
Joardar VS, Kodali VK, Li W, Maglott D, Masterson P, McGarvey KM, Murphy MR, O'Neill K,
Pujar S, Rangwala SH, Rausch D, Riddick LD, Schoch C, Shkeda A, Storz SS, Sun H, Thibaud-
Nissen F, Tolstoy I, Tully RE, Vatsan AR, Wallin C, Webb D, Wu W, Landrum MJ, Kimchi A,
Tatusova T, DiCuccio M, Kitts P, Murphy TD, Pruitt KD. 2016. Reference sequence (RefSeq)
database at NCBI: current status, taxonomic expansion, and functional annotation. Nucleic
Acids Res 44:D733-745.
106. Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. 2016. GenBank. Nucleic Acids Res
44:D67–D72.
107. Olson RD, Assaf R, Brettin T, Conrad N, Cucinell C, Davis JJ, Dempsey DM, Dickerman A,
Dietrich EM, Kenyon RW, Kuscuoglu M, Lefkowitz EJ, Lu J, Machi D, Macken C, Mao C,
Niewiadomska A, Nguyen M, Olsen GJ, Overbeek JC, Parrello B, Parrello V, Porter JS, Pusch
GD, Shukla M, Singh I, Stewart L, Tan G, Thomas C, VanOeffelen M, Vonstein V, Wallace ZS,
Warren AS, Wattam AR, Xia F, Yoo H, Zhang Y, Zmasek CM, Scheuermann RH, Stevens RL.
2023. Introducing the Bacterial and Viral Bioinformatics Resource Center (BV-BRC): a resource
combining PATRIC, IRD and ViPR. Nucleic Acids Res 51:D678–D689.
108. Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz T,
Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen GJ,
Murphy-Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein V, Warren
A, Xia F, Yoo H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial Bioinformatics
Database and Analysis Resource Center. Nucleic Acids Res 45:D535–D542.
109. Federhen S. 2012. The NCBI Taxonomy database. Nucleic Acids Res 40:D136-143.
110. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped
BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids
Res 25:3389–3402.
111. Söding J, Biegert A, Lupas AN. 2005. The HHpred interactive server for protein homology
detection and structure prediction. Nucleic Acids Res 33:W244-248.
112. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. BMC Bioinformatics 10:421.
113. Nielsen H. 2017. Predicting Secretory Proteins with SignalP. Methods Mol Biol Clifton NJ
1611:59–73.
114. Käll L, Krogh A, Sonnhammer ELL. 2004. A combined transmembrane topology and signal
peptide prediction method. J Mol Biol 338:1027–1036.
115. Käll L, Krogh A, Sonnhammer ELL. 2007. Advantages of combined transmembrane topology
and signal peptide prediction--the Phobius web server. Nucleic Acids Res 35:W429-432.

116.   Cole C, Barber JD, Barton GJ. 2008. The Jpred 3 secondary structure prediction server. Nucleic
Acids Res 36:W197-201.
117.   El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson LJ,
Salazar GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L, Piovesan D, Tosatto SCE, Finn RD.
2019. The Pfam protein families database in 2019. Nucleic Acids Res 47:D427–D432.
118.   Finn RD, Clements J, Arndt W, Miller BL, Wheeler TJ, Schreiber F, Bateman A, Eddy SR. 2015.
HMMER web server: 2015 update. Nucleic Acids Res 43:W30-38.
119.   Lassmann T. 2019. Kalign 3: multiple sequence alignment of large data sets. Bioinforma Oxf
Engl https://doi.org/10.1093/bioinformatics/btz795.
120.   Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res 32:1792–1797.
121.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood trees for
large alignments. PloS One 5:e9490.
122.   Rambaut A. 2018. FigTree v1.4.4. GitHub. Repository.
https://github.com/rambaut/figtree/releases. Retrieved 13 July 2023.
123.   Flouri T, Izquierdo-Carrasco F, Darriba D, Aberer AJ, Nguyen L-T, Minh BQ, Von Haeseler A,
Stamatakis A. 2015. The Phylogenetic Likelihood Library. Syst Biol 64:356–362.
124.   Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T. 2020. ModelTest-NG: A New
and Scalable Tool for the Selection of DNA and Protein Evolutionary Models. Mol Biol Evol
37:291–294.
125.   Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019. RAxML-NG: a fast, scalable and
user-friendly tool for maximum likelihood phylogenetic inference. Bioinformatics 35:4453–4455.
126.   Chang W, Cheng J, Allaire JJ, Xie Y, McPherson J. 2019. shiny: Web Application Framework for
R. https://CRAN.R-project.org/package=shiny.
127.   R Core Team. 2020. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/.
128.   RStudio Team. 2020. RStudio: Integrated Development Environment for R. RStudio, PBC.,
Boston, MA. http://www.rstudio.com/.
129.   Hao ZhuThomas Travison DM Timothy Tsai, Will Beasley, Yihui Xie, GuangChuang Yu,
Stéphane Laurent, Rob Shepherd, Yoni Sidi, Brian Salzer, George Gui, Yeliang Fan. 2020.
Construct Complex Table with "kable" and Pipe Syntax. http://haozhu233.github.io/kableExtra/,
https://github.com/haozhu233/kableExtra.
130.   Mike Bostock CY Kerry Rodden, Kevin Warne, Kent Russell, Florian Breitwieser. 2020. Sunburst
"Htmlwidget." https://github.com/timelyportfolio/sunburstR.
131.   Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H. 2014. UpSet: Visualization of
Intersecting Sets. IEEE Trans Vis Comput Graph 20:1983–1992.
132.   Enrico Bonatesta UB Christoph Horejs-Kainrath. 2020. Multiple Sequence Alignment.
http://www.bioinf.jku.at/software/msa/.
133.   Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. Bioinforma Oxf Engl 35:526–528.
134.   Fellows I. 2018. Word Clouds. http://blog.fellstat.com/?cat=11.
135.   Jake Conway NG. 2019. A More Scalable Alternative to Venn and Euler Diagrams for Visualizing
Intersecting Sets. http://github.com/hms-dbmi/UpSetR.
136.   Müller K. 2017. here: A Simpler Way to Find Your Files. https://CRAN.R-
project.org/package=here.

137. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G, Hayes A,
Henry L, Hester J, Kuhn M, Pedersen TL, Miller E, Bache SM, Müller K, Ooms J, Robinson D,
Seidel DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H. 2019. Welcome to the
Tidyverse. J Open Source Softw 4:1686.
138. Martin Morgan MR. 2019. Access the Bioconductor Project Package Repository.
https://github.com/Bioconductor/BiocManager.
139. Perry M. 2016. Flexible Heatmaps for Functional Genomics and Sequence Features.
https://git.bioconductor.org/packages/heatmaps.
140. Xie Y, Allaire JJ, Grolemund G. 2018. R Markdown: The Definitive Guide. Chapman and
Hall/CRC, Boca Raton, Florida. https://bookdown.org/yihui/rmarkdown.
141. Allaire JJ, Xie, Y, McPherson J, Luraschi J, Ushey K, Atkins A, Wickham H, Cheng J, Chang W,
Iannone R. 2020. rmarkdown: Dynamic Documents for R.
https://github.com/rstudio/rmarkdown.
1207