Differential dopaminergic modulation of spontaneous cortico-subthalamic activity in Parkinson's disease

Short running title: Dopamine in cortico-subthalamic networks

Authors:

Abhinav Sharma[1], Diego Vidaurre[2,3], Jan Vesper[4], Alfons Schnitzler[1,5], Esther Florin[1]

[1]*Institute of Clinical Neuroscience and Medical Psychology, Medical Faculty, Heinrich-Heine University Düsseldorf, Germany*

[2]*Department of Psychiatry, University of Oxford, Oxford, UK*

[3]*Department of Clinical Health, Aarhus University, Aarhus, Denmark*

[4] *Department of Neurosurgery, University Hospital Düsseldorf, Germany*

[5] *Department of Neurology, Center for Movement Disorders and Neuromodulation, Medical Faculty, Heinrich-Heine University Düsseldorf, Germany.*

Correspondence to:
Prof. Dr. Esther Florin
Heinrich-Heine University Düsseldorf
Institute of Clinical Neuroscience and Medical Psychology
Universitätstr. 1
40225 Düsseldorf
Germany
Email: Esther.Florin@hhu.de
Tel: +49 211 81 08437

Abhinav Sharma
Heinrich-Heine University Düsseldorf
Institute of Clinical Neuroscience and Medical Psychology
Universitätstr. 1
40225 Düsseldorf
Germany
Email: Abhinav.Sharma@med.uni-duesseldorf.de

Abstract

Pathological oscillations including elevated beta activity in the subthalamic nucleus (STN) and between STN and cortical areas are a hallmark of neural activity in Parkinson's disease (PD). Oscillations also play an important role in normal physiological processes and serve distinct functional roles at different points in time. We characterised the effect of dopaminergic medication on oscillatory whole-brain networks in PD in a time-resolved manner by employing a hidden Markov model on combined STN local field potentials and magnetoencephalography (MEG) recordings from 17 PD patients. Dopaminergic medication led to communication within the medial and orbitofrontal cortex in the delta/theta frequency range. This is in line with deteriorated frontal executive functioning as a side effect of dopamine treatment in Parkinson's disease. In addition, dopamine caused the beta band activity to switch from an STN-mediated motor network to a frontoparietal-mediated one. In contrast, dopamine did not modify locally-originating STN oscillations in PD. STN–STN synchrony emerged both on and off medication. By providing electrophysiological evidence for the differential effects of dopaminergic medication on the discovered networks, our findings open further avenues for electrical and pharmacological interventions in PD.

Keywords: whole-brain, time-resolved analysis, local field potentials, Magnetoencephalography

List of abbreviations:

comms - Communication state; DBS - Deep brain stimulation; EEG – Electroencephalography; ECG – Electrocardiography; EOG – Electrooculography; FO - Fractional occupancy; GMM - Gaussian mixture model; HMM - Hidden Markov model; hyper –DA Hyper-dopaminergic state; intraMED - intra-medication; interMED – inter-medication; L-DOPA – Levodopa; LFP - Local field potential; MEG – Magnetoencephalography; NNMF - Non-negative matrix factorization; PD - Parkinson's disease; PFC - Prefrontal cortex; PCA - Principal component analysis; STN - Subthalamic nucleus; TDE -Time-delay embedding

1 Introduction

Oscillatory activity serves crucial cognitive roles in the brain[1,2] and alterations of oscillatory activity have been linked to neurological and psychiatric diseases.[3] Different large-scale brain networks operate with their own oscillatory fingerprint and carry out specific functions.[4–6] Given the dynamics of cognition, different brain networks need to be recruited and deployed flexibly. Hence, the duration for which a network is active, its overall temporal presence, and even the interval between the different activations of a specific network might provide a unique window to understanding brain functions. Crucially, alterations of these temporal properties or networks might be related to neurological disorders.

In Parkinson's disease (PD), beta oscillations within the subthalamic nucleus (STN) and motor cortex (13–30 Hz) correlate with the motor symptoms of PD.[7–9] Beta oscillations also play a critical role in synchronising communication in a healthy brain.[10] At the cellular level, loss of nigral dopamine neurons in PD leads to widespread changes in brain networks, to varying degrees across different patients. Dopamine loss is managed in patients via dopaminergic medication. As dopamine is a widespread neuromodulator in the brain,[11] this raises the question of whether all medication-induced changes can restore physiological oscillatory networks.[12]

We characterised the differential effects of dopaminergic medication and delineated pathological versus physiologically-relevant spectral connectivity in PD. In our present work, we studied PD brain activity via a hidden Markov model (HMM), a data-driven learning algorithm.[6,13] Due to the importance of cortico–subcortical interactions in PD, we investigated these interactions with combined spontaneous whole-brain magnetoencephalography (MEG) and STN local field potentials (LFPs) recordings from PD patients. By investigating whole-brain connectivity, including the STN, we extended previous work that was limited to specific connectivity pairs.[14,15] Moreover, we identified the temporal properties of the networks both ON and OFF medication. The temporal properties provide an encompassing view of network alterations in PD and the effect of dopamine on these networks.

We found that cortico–cortical, cortico–STN, and STN–STN networks were differentially modulated by dopaminergic medication. Within those networks, the effect of medication was disruptive for the cortico–cortical networks, physiologically restorative for the cortico–STN networks, and limited for the STN–STN interactions. These results provide novel information on the oscillatory network connectivity occurring in PD and the differential changes caused by dopaminergic intervention. These whole-brain networks, along with their electrophysiological

signatures, open up new potential targets for both electric and pharmacological interventions in PD.

2 Materials and Methods

2.1 Subjects

In total, 17 (4 female) right-handed PD patients (age: 55.2 ± 9.3 years) undergoing surgery for therapeutic STN deep brain stimulation (DBS) were recruited for this study. Patients had been selected for DBS treatment according to the guidelines of the German Society for Neurology. The experimental procedure was explained to all participants and they gave written consent. The study was approved by the local ethics committee (study number 5608R) and conducted in accordance with the Declaration of Helsinki. Bilateral DBS electrodes were implanted in the dorsal part of the STN at the Department of Functional Neurosurgery and Stereotaxy in Düsseldorf. The implanted DBS electrodes used were the St. Jude Medical directional lead 6172 (Abbott Laboratories, Lake Bluff, USA) and in one case the Boston Scientific Vercise segmented lead (Boston Scientific Corporation, Marlborough, USA). These electrodes have four contact heights and the two middle heights are segmented into three equally-spaced contacts.

The DBS leads were externalised and we measured the patients after 1 to 3 days. To simultaneously acquire MEG and LFP signals, we connected the externalised leads to an electroencephalogram (EEG) amplifier integrated with the MEG system. We used a whole-head MEG system with 306 channels (Elekta Vectorview, Elekta Neuromag, Finland) housed within a magnetically-shielded chamber. All patients were requested to sit still and awake during data acquisition. To ensure that patients did not fall asleep, we tracked patients' pupil diameter with an eye tracker. To remove eye blink and cardiac artefacts, electrooculography (EOG) and electrocardiography (ECG) were recorded along with the LFP and MEG signals. In order to co-register the MEG recording with the individual magnetic resonance imaging (MRI), four head position indicator coils were placed on the patient's head. Their position as well as additional head points were digitised using the Polhemus Isotrack system (Polhemus, Colchester, USA). The data were recorded with a sampling rate of 2400 Hz and a low-pass filter of 800 Hz was applied. An electrode was placed at the mastoid and all LFP signals were referenced to it.

For the clinical OFF medication state, oral PD medication was withdrawn overnight for at least 12 hours. If a patient had an apomorphine pump, this pump was stopped at least 1 hour before the measurement. First, we recorded resting-state activity in the medication OFF condition. The patients were then given their morning dose of L-DOPA in the form of fast-acting levodopa. Data were acquired in three runs of 10 minutes, for a total of 30 minutes for each medication condition. We started the ON medication measurement at least half an hour after the administration of the dose and after clinical improvement was seen. The same procedure as for the OFF medication state was followed for the ON medication measurement.

2.2   Pre-processing

All data processing and analyses were performed using Matlab (version R 2016b; Math Works, Natick, USA). Custom-written Matlab scripts and the Brainstorm toolbox (http://neuroimage.usc.edu/brainstorm/Introduction) were used.[16] To ensure artefact-free data, two people independently inspected the data visually, cleaned artefacts, and compared the cleaning output. The final cleaned data included changes agreed upon by both the people involved in cleaning. The Neuromag system provides signal-space projection (SSP) vectors for the cleaning of external artefacts from the MEG channels, which were applied. The line noise was removed from all channels with a notch filter at 50, 100, 150, …, 550 and 600 Hz with a 3 dB bandwidth of 1 Hz. The LFP recordings from the DBS electrode were re-referenced against the mean of all LFP channels. Very noisy and flat MEG/LFP channels were excluded from further analysis. Time segments containing artefacts were removed from the time series. However, if artefacts regularly occurred only in one single channel, this whole channel was removed instead. Frequently-arising artefacts following the same basic pattern, such as eye blinks or cardiac artefacts, were removed via SSP. All data were high-pass filtered with 1 Hz to remove movement-related low-frequency artefacts. Finally, the data were down-sampled to 1,000 Hz.

Source estimation was performed on these recordings at an individual level using each individual's anatomy. Therefore, using Freesurfer (https://surfer.nmr.mgh.harvard.edu/, v.5.3.0), the individual cortical surfaces were extracted from the individual T1-weighted MRI scans (3T scanner and 1 mm³ voxel size). We used the overlapping spheres method with 306 spheres for the forward model. As the inverse model, we used a linearly constrained minimum variance (LCMV) beamformer. The data covariance matrix for the LCMV beamformer was

computed directly from each 10-minute recording. The data covariance was regularised using the median eigenvalue of the data covariance matrix. The noise covariance was obtained from an empty room recording on the same day as the actual measurement.

For each subject, the invasive entry point of the STN was identified based on intraoperative microelectrode recordings.[17,18] Subsequently, the first recording height after the entry into the STN was selected to obtain the three directional LFP recordings from the respective hemisphere. The source-reconstructed MEG data were projected to the default cortical anatomy (MNI 152 with 15002 vertices) and then down-sampled temporally to 250 Hz for each medication condition for every subject. We used the Mindboggle atlas to spatially reduce the data dimensions. For each of the 42 cortical regions in the atlas, a multidimensional time series consisting of the vertices within that anatomical region was extracted. To reduce the multivariate times series for each region to a single one, we employed the first principal component explaining the highest variance share in each region. The first principal component row vectors from all 42 anatomical regions were stacked into a MEG cortical time series matrix. To correct for volume conduction in the signal, symmetric orthogonalisation[19] was applied to each subject's resulting cortical time series matrix. The row vectors of this orthogonalised matrix and the six LFPs (three each for left and right STN) were z-scored. Subsequently, they were stacked into one multidimensional time series (N by T) matrix. Here, N = 48 is the total number of nodes/regions (42 regions from the cortex and 6 LFP electrode contacts) and T denotes the length of the time dimension. This 48 by T data matrix obtained from each subject was concatenated along the temporal dimension across all subjects for each specific medication condition. Finally, to resolve sign ambiguity inherent in source-reconstructed MEG data as well as resolve polarity of LFP channels across subjects, a sign-flip correction[13] procedure was applied to this final 48 by (T by number of subjects) dataset within a medication condition. The pre-processing steps were performed for OFF and ON medication separately.

2.3   Hidden Markov model analysis

The HMM is a data-driven probabilistic algorithm which finds recurrent network patterns in multivariate time series.[13,20] Each network pattern is referred to as a 'state' in the HMM framework, such that these networks can activate or deactivate at various points in time. Here onwards, 'state' or 'network' is used interchangeably. We used a specific variety of the HMM, the time-delay embedded HMM (TDE-HMM), where whole-brain networks are defined in terms of both spectral power and phase-coupling.[6] Hence, for every time point, the HMM

algorithm provided the probability that a network is active. Here onwards, a contiguous block of time for which the probability of a particular network being active remained higher than all the other networks is referred to as a 'state visit'. Hence, the HMM produced temporally-resolved spatial networks for the underlying time series. In our approach we also performed spectral analyses of these state visits, leading to a complete spatio–spectral connectivity profile across the cortex and the STN. By applying the HMM analysis to the combined MEG-LFP dataset we were able to temporally, spatially and spectrally separate cortico–cortical, cortico–STN and STN–STN networks.

2.3.1 Estimation of the hidden Markov model

Since we were interested in recovering phase-related networks, the TDE-HMM was fit directly on the time series obtained after pre-processing steps described previously (Section 2.2), as opposed to its power envelope. This preserved the cross-covariance within and across the underlying raw time series of the cortical regions and the STN. The model estimation finds recurrent patterns of covariance between regions (42 cortical regions and 6 STN contacts) and segregates them into 'states' or 'networks'. Based on these covariance patterns, for each state, the power spectra of each cortical region and the coherence amongst regions can be extracted.

We opted for six different states as a reasonable trade-off between the spectral quality of the results and their redundancy. The HMM-MAR toolbox[13] was used for fitting the TDE-HMM model. We employed the TDE version of the HMM where the embedding took place in a 60 ms window (i.e. a 15 time point window for a sampling frequency of 250 Hz). Since time embedding would increase the number of rows of the data from 48 to 48 times the window length (also referred to as number of lags), an additional PCA (reduction across 48 by number of lags) step was performed after time embedding. The number of components retained was 96 (48 x 2). This approach follows Vidaurre *et al.*.[6] To characterise each state, a full covariance matrix with an inverse Wishart prior was used. The diagonal of the prior for the transition probability matrix was set as 10. To ensure that the mean of the time series did not take part in driving the states, the 'zero mean' option in HMM toolbox was set to 1. To speed up the process of fitting we used the stochastic version of variational inference for the HMM. In order to start the optimisation process, the 'HMM-MAR'-type initialisation was used (for details see (Vidaurre *et al.*[13]). The HMM was fit separately OFF and ON medication.

2.3.2 Statistical analysis of the states

After the six states were obtained for HMM OFF and HMM ON medication, these states were statistically compared within each medication condition as well as between medication conditions. In addition, the temporal properties of these states were compared.

2.3.2.1 Intra-medication analysis

We investigated the spectral connectivity patterns across the different states within a medication condition (intra-medication or IntraMed). The objective was to uncover significant coherent connectivity standing out from the background within each frequency band (delta/theta [1-8 Hz], alpha [8-12 Hz], and beta [13-30 Hz]) in the respective states. The HMM output included the state time courses (i.e. when the states activated) for the entire concatenated data time series. The state time courses allowed the extraction of state- and subject-specific data for further state- and subject-level analysis. For each HMM state, we filtered the state-specific data[1] for all the subjects between 1 and 45 Hz. Then we calculated the Fourier transform of the data using a multitaper approach to extract the frequency components from the short segments of each state visit.[2] Seven Slepian tapers with a time-bandwidth product of 4 were used, resulting in a frequency resolution of 0.5Hz and therefore binned frequency domain values. Subsequently, we calculated the coherence and power spectral density of this binned (frequency bins obtained during the multitaper step) data for every subject and every state. The coherence and the power spectral density obtained were three-dimensional matrices of size f (number of frequency bins) by N (42 cortical locations + 6 STN contacts) by N.

Based on the coherence matrices, we performed a frequency band-specific analysis. Canonical definitions of frequency bands assign equal weight to each frequency bin within a band for every subject. This might not be suitable when considering analyses of brain signals across a large dataset. For example, the beta peak varies between individual subjects. Assigning the same weight to each bin in the beta range might reduce the beta effect at the group level. To allow for inter-subject variability in each frequency bin's contribution to a frequency band, we determined the frequency modes in a data-driven manner.[6] Because we focused on interactions

---

[1] For state-wise data extraction please refer the HMM toolbox wiki (https://github.com/OHBA-analysis/HMM-MAR/wiki/User-Guide).

[2] see (Vidaurre *et al.*, 2018*b*) for discussion on multitaper for short time data segments.

that are important to establish the STN–cortex communication, the identification of the relevant frequency modes was restricted to the cross-coherence between the STN-LFPs and cortical signals; in other words, the block matrix consisting of rows 1 to 6 (STN) and columns 7 to 48 (cortex). For each subject, this extracted submatrix was then vectorised across columns. This gave us a (number of frequency bins by 252 [6 STN locations by 42 cortical locations]) matrix for each state. For every subject, this matrix was concatenated along the spatial dimension across all states producing a (number of frequency bins by [252 by 6 (number of states)]) matrix. We called this the subject-level coherence matrix. We averaged these matrices across all subjects along the spectral dimension (number of frequency bins) to yield a (number of frequency bins by [252 by 6]) group-level coherence matrix. We factorised the group-level coherence matrix into four frequency modes using a non-negative matrix factorisation (NNMF) (Lee and Seung, 2001). Each of the resulting four frequency modes obtained was of size (1 by number of frequency bins). Three of them resembled the canonical delta/theta (delta and theta frequencies were combined into one band), alpha, and beta bands whereas the last one represented noise. Since NNMF does not guarantee a unique solution, we performed multiple instances of the factorisation. At each instance, we visualised the output to ensure frequency specificity of the frequency modes. The stability of the output was ensured by using 'Robust NNMF', which is a variant of the NNMF algorithm.[6] We then computed the inner product between the subject- and group-level coherence matrix and the frequency modes obtained above. We called these the subject-level and group-level projection results, respectively.

To separate background noise from the strongest coherent connections, a Gaussian mixture model (GMM) approach was used.[6] For the group-level projection results, we normalised the activity in each state for each spectral band by subtracting the mean coherence within each frequency mode across all states. As a prior for the mixture model, we used two single-dimensional Gaussian distributions with unit variance: one mixture component to capture noise and the other to capture the significant connections. This GMM with two mixtures was applied to the coherence values (absolute value) of each state. Connections were considered significant if their p-value after correction for multiple comparisons was smaller than 0.05.

2.3.2.2 Inter-medication analysis

To test for differences in coherence across medication conditions (inter-medication or InterMed), the first step was to objectively establish a comparison between the states found in

the two HMMs fit separately for each condition. There is no a priori reason for the states detected in each condition to resemble each other. To find OFF and ON medication states that may resemble each other, we calculated the Riemannian distance[21] the state covariance matrices of the OFF and ON HMM. This yielded an OFF states by ON states (6 x 6) distance matrix. Subsequently, finding the appropriately matched OFF and ON states reduced to a standard linear assignment problem. We found an ON state counterpart to each OFF state by minimising the total sum of distances using the Munkres linear assignment algorithm.[20] This approach yielded a one-to-one pairing of OFF and ON medication states, and all further analysis was conducted on these pairs. For ease of reading, we gave each pair its own label. For example, when we refer to a 'communication' state in the following sections, then such a state was discovered OFF medication and its corresponding state ON medication is its distance-matched partner. In subsequent sections, all mentions of ON or OFF medication refer to these state pairs unless mentioned otherwise.

We used the subject-level projection results obtained during IntraMed analysis to perform InterMed analyses. We performed two-sided independent sample t-tests between the matched states to compare the coherence, which was calculated between different regions of interest (see Dataset preparation). We grouped individual atlas regions into canonical cortical regions like frontal, sensorimotor, parietal, visual, medial PFC, and STN contacts. For example, in the beta band, STN (contacts)-sensorimotor coherence in the OFF condition was compared to the STN (contacts)-sensorimotor coherence in the ON condition. The p-values obtained were corrected for multiple comparisons for a total number of possible combinations.

2.3.3 Temporal properties of hidden Markov model states

To test for changes in the temporal properties OFF versus ON medication, we compared the lifetimes, interval between visits, and fractional occupancy (FO) for each state both within and across HMMs using two-sided independent sample t-tests. Lifetime/dwell time of a state refers to the time spent by the neural activity in that state. Interval of visit was defined as the time between successive visits of the same state. Finally, the FO of a state was defined as the fraction of time spent in each state. Extremely short state visits might not reflect neural processes, hence we only used values that were greater than 100 ms for lifetime comparisons.

2.4 Data availability

Because of privacy law, data are available upon personal request. Enquiries can be sent to the corresponding author.

3 Results

Under resting-state conditions in PD patients, we simultaneously recorded whole-brain MEG activity with LFPs from the STN using directional electrodes implanted for DBS. Using an HMM we identified recurrent patterns of transient network connectivity between the cortex and the STN, which we henceforth refer to as an 'HMM state'. Each HMM state was characterised by coherence calculated between different pairs of regions. Additionally, the temporal evolution of the HMM states was determined. The PD data were acquired under medication (L-DOPA) OFF and ON conditions, which allowed us to delineate the physiological versus pathological spatio–spectral and temporal changes observed in PD. OFF medication, patients had on average a Unified Parkinson's Disease Rating Scale (UPDRS) part III of 29.24 ± 10.74. This was reduced by L-DOPA (176.5 ± 56.2 mg) to 19.47 ± 8.52, indicating an improvement in motor symptoms. HMMs model discrete-state stochastic dynamical systems[22] that bring a dynamical systems perspective to the understanding of pathological versus physiological connectivity in PD.

3.1 Spontaneous brain activity in Parkinson's disease can be resolved into distinct states

Motor or tremor network-based studies have evaluated specific cortico-subthalamic spectral connectivity[23] but did not account for the whole-brain changes affecting them. Conversely, resting-state whole-brain MEG studies have identified network changes related to both motor and non-motor symptoms of PD but did not account for STN-STN or cortico–STN activity affecting these networks.[24,25] Using an HMM, we delineated cortico-subthalamic spectral changes from both global source-level cortical interactions as well as local STN–STN interactions. Three of the six HMM states could be attributed to physiologically interpretable connectivity patterns. We could not interpret the other three states within the current physiological frameworks both OFF and ON medication and they are therefore not considered in the following (see Supplementary Figures S1-S3). The connectivity between different brain regions for each state was visualised for the frequency modes shown in Figure 1. Figures 2-4 show the connectivity patterns for the three physiologically meaningful states in both the OFF

(top row) and ON medication condition (bottom row). We refer to the state obtained in Figure 2 as the *hyper-dopaminergic state (hyper-DA)*. The frontal cortex changes observed in this state, ON medication, are in line with the dopamine overdose hypothesis in PD (see below),[26,27] and therefore this state was labelled as hyper-DA. This state was characterised mostly by local coherence within segregated networks OFF medication in the alpha and beta band. In contrast, there was a widespread increase in coherence across the brain from OFF to ON medication. Therefore, ON medication, the connectivity strength in the alpha and beta band was not significantly different from the mean noise level. Figure 3 displays the second state. A large proportion of spectral connections in this state enable cortico–STN communication[8,14,15,28,29] and thus we labelled this as the *communication state (comms)*. This state was characterised by connectivity between multiple cortical regions and the STN OFF medication, but increased specificity of cortical–STN connectivity ON medication. Finally, Figure 4 shows the third state. Within this state, highly synchronous STN–STN spectral connectivity emerged, both OFF and ON medication and therefore we named it the *local state (local)*. The spectral characteristics of this state largely remain unaffected under the influence of dopaminergic medication. In the following sections, we describe these three states in detail.

3.2 Hyper-dopaminergic state is characterised by increased frontal coherence due to elevated dopamine levels

Dopaminergic medication is known to produce cognitive side effects in PD patients.[12] According to the dopamine overdose hypothesis, a reason for these effects is the presence of excess dopamine in brain regions not affected in PD.[27,30] Previous task-based and neuroimaging studies in PD demonstrated frontal cognitive impairment due to dopaminergic medication.[30–32] Supporting the dopamine overdose hypothesis in PD, we identified a delta/theta oscillatory network involving lateral and medial orbitofrontal cortical regions. To the best of our knowledge, we report these results for the first time under resting-state conditions.

The delta/theta network emerged between the lateral and medial orbitofrontal cortex ON medication ($p < 0.05$, Figure 2C delta). On the contrary, OFF medication no significant connectivity was detected in the delta/theta band. In the alpha and beta band OFF medication there was significant connectivity within the frontal regions, STN, and to a limited extent in the posterior parietal regions ($p < 0.05$, Figure 2A).

Another effect of excess dopamine was significantly increased connectivity of frontal cortex and temporal cortex both with the STN and multiple cortical regions across all frequency modes

(p < 0.01, Figure 2 delta, alpha, and beta). The change in sensorimotor–STN connectivity primarily took place in the alpha band with an increase ON medication. Sensorimotor–cortical connectivity was increased ON medication across multiple cortical regions in both the alpha and beta band (p < 0.01, Figure 2 alpha and beta). However, STN–STN coherence remained unchanged OFF versus ON medication across all frequency modes.

Viewed together, the hyper-dopaminergic state captured increased coherence across the cortex ON medication within the alpha and beta band. This, however, implies that ON medication no connectivity strength was significantly higher than the mean noise level within the alpha or beta band. ON medication significant coherence emerged in the delta/theta band primarily between different regions of the orbitofrontal cortex. Increased frontal cortical coherence in the orbitofrontal cortex points towards an overdosing effect of tonic dopamine levels, which might explain the cognitive deficits observed in PD ON medication.

3.3   Dopaminergic medication selectively reduced connectivity in the communication state

Previous MEG–LFP studies in PD involving dopaminergic intervention have reported changes in beta and alpha band connectivity between specific cortical regions and the STN.[14,15,33] However, these studies did not report simultaneous cortico–cortical as well as STN–STN changes within different frequency modes. Our analysis revealed that the communication state ON medication was characterised by selective cortico–STN spectral connectivity and an overall shift in cortex-wide activity towards physiologically-relevant network connectivity.

In particular, ON medication, connectivity between STN and cortex became more selective in the alpha and beta band. OFF medication, significant coherence was present between multiple cortical regions and STN in the alpha and beta band (p < 0.05, Figure 3A). Importantly, coherence OFF medication was significantly larger than ON medication between STN and sensorimotor, STN and temporal and STN and frontal cortices (p < 0.05 for all connections, Figure 3B alpha and beta). Furthermore, ON medication in the alpha band only the connectivity between temporal and parietal cortical regions and the STN was preserved (Figure 3C alpha; p < 0.05), consistent with previous findings.[14] In contrast, in the beta band only STN-medial orbitofrontal connectivity remained intact (p < 0.05, Figure 3C beta).

Most previous PD studies report a decrease in the motor–STN coherence ON medication in the beta band[14,15,34–36] but do not indicate any changes that the sensorimotor regions might experience at the whole-brain level. In the communication state, OFF medication, STN–premotor (sensory), STN–frontal, and STN–parietal connectivity was present ($p < 0.05$, Figure 3A alpha and beta). STN–cortical coherence was then significantly reduced ON compared to OFF medication ($p < 0.05$, Figure 3B alpha and beta). Finally, ON medication, a sensorimotor–frontoparietal network emerged ($p < 0.05$, Figure 3C beta) where sensorimotor, frontal and parietal regions were no longer connected to the STN, but instead directly communicated with each other in the beta band. Hence, there was a transition from STN-mediated sensorimotor connectivity to the cortex OFF medication to a more direct cortico–cortical connectivity ON medication.

In the communication state, we found STN–cortico and cortico–cortical changes and simultaneously a change in STN–STN connectivity. In the ON condition, STN–STN connectivity was significantly different from the mean noise level across all three frequency modes ($p < 0.05$, Figure 3C). But on the other hand, there was no significant change in the STN–STN connectivity OFF versus ON medication ($p=0.21$ - delta/theta; $p=0.25$ - alpha; $p=0.10$ - beta; Figure 3B).

To summarise, coherence decreased ON medication across a wide range of cortical regions both at the cortico–cortical and cortico–STN level. Still, significant connectivity was selectively preserved in a spectrally-specific manner ON medication both at the cortico–cortical level and the cortico–STN level. Furthermore, a sensorimotor–frontoparietal network emerged ON medication. The most surprising aspect of this state was the emergence of bilateral STN–STN coherence ON medication across all frequency modes. The results provide evidence for the plausible role of dopamine to increase the signal-to-noise ratio across the brain, as well as the importance of beta-band activity in enabling cortical control of motor activity ON medication.

3.4 Dopamine selectively modifies delta/theta oscillations within the local state

STN–STN intra-hemispheric oscillations positively correlate to motor symptom severity in PD OFF medication, whereas dopamine-dependent nonlinear phase relationships exist between inter-hemispheric STN–STN activity.[37] Crucially, previous studies could not rule out the influence of cortico–STN connectivity on these inter-hemispheric STN–STN interactions.

Using our whole-brain framework, we found a state exhibiting STN–STN connectivity only, both OFF and ON medication.

In this local state, dopaminergic intervention had only a limited effect on local STN–STN connectivity. OFF medication, STN–STN coherence was present across all three frequency modes ($p < 0.05$, Figure 4A), while ON medication, significant STN–STN coherence emerged only in the alpha and beta bands ($p < 0.05$, Figure 4C alpha and beta). ON medication, STN–STN delta/theta connectivity strength was not significantly different from the mean noise level ($p < 0.05$, Figure 4C delta).

InterMed analysis also revealed that OFF compared to ON medication coherence was reduced across the entire cortex both at the inter-cortical and the STN–cortex level across all frequency modes. The most affected areas were similar to the ones in the communication state, in other words, the sensorimotor, frontal, and temporal regions. Their coherence with the STN was also significantly reduced, ON compared to OFF medication (STN–sensorimotor, $p < 0.01$ - delta/theta, beta; $p < 0.05$ - alpha; STN–temporal, $p < 0.01$ - delta/theta, alpha, beta; and STN–frontal, $p < 0.01$ - delta/theta, alpha and beta; Figure 4B).

In summary, STN–STN connectivity was not significantly altered OFF to ON medication. At the same time, coherence decreased from OFF to ON medication at both the cortico–cortical and the cortico–STN level. Therefore, only significant STN–STN connectivity existed both OFF and ON medication, while cortico–STN or cortico–cortical connectivity changes remained at the mean noise level.

3.5   States with a generic coherence decrease have longer lifetimes

We have thus far identified three distinct HMM states, which were differentially modified, both spatially and spectrally, under the effect of dopaminergic medication. Previous research has shown that ON medication, spectrally-specific cortico–STN connectivity remains preserved in PD compared to OFF medication.[14,15] This indicates the existence of functionally relevant cortico–STN loops. A decrease in coherence between the cortex and the STN has also been observed ON medication,[38] which was correlated with improved motor functions in PD. All the connectivity effects were observed in our results for the communication state. Furthermore, in the communication state, we showed the existence of a frontoparietal sensorimotor network in the beta band ON medication. Recent evidence indicates that with the loss of dopamine and the

start of PD symptoms, delta/theta oscillations emerge within the basal ganglia.[39] In line with these findings, STN–STN delta/theta oscillations in the local state were reduced ON medication. Using the temporal properties of the identified networks, we investigated whether states showing physiological connectivity patterns lasted longer ON medication. A state that is physiological should exhibit increased lifetime and/or should occur more often ON medication.

Figure 5A–C shows the temporal properties for the three states for both the OFF and ON medication conditions. OFF medication, the local state was the one with the longest lifetime (local > hyper-DA, $p < 0.01$; local > comms, $p < 0.01$). The communication state OFF medication had the shortest lifetime among all three states (comms < hyper-DA, $p < 0.01$; comms < local, $p < 0.01$) and the shortest interval between visits (interval of visit comms < hyper-DA, $p < 0.01$; comms < local, $p < 0.01$). The largest interval between visits was for the hyper-dopaminergic state OFF medication (hyper-DA > local, $p < 0.01$; hyper-DA > comms, $p < 0.01$). The FO for the local and communication state was similar, but significantly higher than for the hyper-dopaminergic state (local > hyper-DA, $p < 0.01$; local $\approx$ comms, $p = 0.65$; comms > hyper-DA, $p < 0.01$). ON medication, the comparison between temporal properties of all three states retained the same significance levels as OFF medication, except for the lifetime of the communication state, which was no longer significantly different from that of the hyper-dopaminergic state ($p = 0.75$). Within each medication condition, the states retained their temporal characteristics relative to each other.

Across medication conditions, significant changes were present in the temporal properties of the states. The lifetimes for both the local and communication state were significantly increased by medication (ON > OFF: local, $p < 0.01$; comms, $p \leq 0.01$) but the lifetime for the hyper-dopaminergic state was not significantly influenced by medication. The hyper-dopaminergic state was visited even less often ON medication (Interval: ON > OFF hyper-DA, $p < 0.01$). The interval between visits remained unchanged for the local and communication states. The FO for all three states was not significantly changed from OFF to ON medication. According to the temporal properties analysed, the hyper-dopaminergic state was visited least often compared to the other two states both OFF and ON medication. Both the local and communication state tended to last longer ON medication. This indicates that the states showing physiologically-relevant spectral connectivity lasted longer ON medication.

4 Discussion

In this study, we parsed simultaneously-recorded MEG-STN LFP signals into discrete time-resolved states to reveal distinct spectral communication patterns. We identified three states exhibiting distinct coherence patterns ON and OFF medication: a hyper-dopaminergic, a communication, and a local state. Our results indicate a tendency of neural activity to engage in connectivity patterns in which coherence decreases under the effect of dopaminergic medication and which maintain selective cortico–STN connectivity (communication and local states). Only within the hyper-dopaminergic state did coherence increase under dopaminergic medication. These results are in line with the multiple effects of dopaminergic medication reported in resting and task-based PD studies.[37,40,41]

The differential effect of dopamine allowed us to delineate pathological and physiological spectral connectivity. The hyper-DA state provided electrophysiological evidence in the delta/theta band for the overdose effect of dopaminergic medication in PD. Prior to the electrophysiological evidence in our study, there was only evidence through task-based or fMRI studies.[30–32] The communication state revealed that simultaneous cortico–cortical and STN–STN interactions emerge ON medication, with spectrally- and spatially-specific cortico–STN interactions. In addition, ON medication, a frontoparietal motor network was present, indicating a shift from STN-mediated motor connectivity to a cortical one. These findings have not been reported in previous studies. The local state exhibited the limited ability of dopaminergic medication to modify local STN–STN delta oscillations. Our analysis also revealed significant changes in the temporal properties of the connectivity profiles, including lifetime and FO, under the effect of dopaminergic medication. This insight might in the future prove important for modifying medication as well as DBS-based strategies for therapeutic purposes.

4.1 Increased tonic dopamine causes excessive frontal cortical activity

The hyper-dopaminergic state showed significant coherent connectivity between the orbitofrontal cortical regions in the delta/theta band ON medication. According to the dopamine overdose hypothesis[26,27,42,43] in PD, the commonly used doses of dopaminergic medication to mitigate the motor symptoms cause the ventral frontostriatal cortical circuits to experience an excessive increase in tonic dopamine levels. This medication-induced increase is due to excessive compensation of dopamine in the ventral striatal circuitry, which experiences a lower

loss of dopamine than its dorsal counterpart. The reason is that in PD dopaminergic neurons in the substantia nigra are primarily lost and therefore the dopamine depletion within the dorsal circuitry is higher than within the ventral one.[26,27] Frontal regions involved with the ventral striatal circuitry include the orbitofrontal cortex, anterior cingulate, and the inferior temporal cortex.[27,44] Increased frontal cortex connectivity potentially explains the cognitive deficits observed in PD.[38,45] Our detected emergence of frontal cortico–cortical coherence (between orbitofrontal and medial orbitofrontal regions) specifically in the delta/theta band could explain the cognitive deficits observed in PD due to dopaminergic medication, given the role of frontal delta/theta oscillations in cognition.[46,47]

A comparison of temporal properties of the hyper-dopaminergic state OFF versus ON medication revealed that the interval between visits was significantly increased ON medication, while the FO of this state was significantly reduced. In fact, the FO of the hyper-dopaminergic state was the lowest among the three states. The temporal results indicate that the hyper-dopaminergic state is least visited. Neural activity ON medication is not likely to visit this state, but whenever it does, its visits are of the same duration as OFF medication. Hence, the hyper-dopaminergic state's presence could explain the cognitive side effects observed ON medication in PD.

4.2   Selective spectral connectivity remains preserved with increased dopamine levels

In line with previous research,[14,15] within the communication state, selective spectral connectivity between the cortex and STN was maintained ON versus OFF medication. Moreover, coherent ensembles emerged primarily in the alpha and beta band within frontal and parietal regions. This is in line with previous HMM results obtained for healthy subjects that also detect anterior and posterior cognitive networks.[6]

An interesting feature of the communication state was the emergence of local STN–STN coherence in all three frequency modes. Bilateral STN–STN coherence in the alpha and beta band did not change in the communication state ON versus OFF medication. This activity became significant only ON medication (IntraMed analysis). Since synchrony limits information transfer,[48–50] local oscillations are a potential mechanism to prevent excessive communication with the cortex. Another possibility is that a loss of cortical afferents causes

local basal ganglia oscillations to become more pronounced. The causal nature of the cortico-basal ganglia interaction is an endeavour for future research.

An important finding in the communication state was the emergence of a frontoparietal sensorimotor network in the beta band ON medication. This was in contrast to an STN-mediated network OFF medication, where sensorimotor frontal and parietal regions were coherent with the STN but not directly with each other. Previous research has shown that critical information regarding sensorimotor decision-making involves frontoparietal regions.[51–54] Hence, the emergence of frontoparietal connectivity with motor regions points towards the physiological relevance of the communication state. Moreover, neural activity ON medication remained longer in the communication state as the lifetime of this state significantly increased compared to OFF medication. The finding is in line with our hypothesis that a state showing physiologically-relevant spectral connectivity lasts longer ON medication.

4.3 Tonic dopamine has a limited effect on local STN–STN interactions

In the communication state, STN–STN coherence accompanied network changes affecting cortico–STN communication ON medication, thereby likely having a functional role. In contrast, in the local state, STN–STN coherence emerged without the presence of any significant cortico–STN coherence either OFF or ON medication. This may indicate that the observed STN–STN activity in the local state emerged due to local basal ganglia circuitry. No changes were observed in the alpha and beta band in the local state ON versus OFF medication, which may indicate the inability of tonic dopamine to modify basal ganglia circuit activity. These results provide more evidence that the changes in STN–STN coherence observed in previous studies[29,35,55] reflect cortical interaction affecting STN activity. Future studies should analyse changes occurring within the STN. To the best of our knowledge, we are the first to uncover modulation of STN–STN delta/theta oscillations by dopaminergic medication. Studies have shown that local basal ganglia delta oscillations, which do not require input from the motor cortex, are robust biomarkers of dopamine depletion.[39] Hence, selective elimination of delta/theta oscillations under dopaminergic medication in the local state points towards restoration of physiologically-relevant network activity.

4.4 Limitations of the study

In the present study, we employed a data-driven approach based on an HMM. In order to find the appropriate model, we had to specify the number of states a priori. We selected the number of HMM model states based on a compromise between spectral quality of results and their redundancy. The number of states could also be determined by selecting the one with the highest negative free energy. However, model selection based on free energy often does not yield concrete results.[56] Another limitation is the use of multivariate Gaussian distributions to characterise the state covariance matrices. Although it improves the tractability of the HMM estimation process, it is by construction unable to capture higher-order statistics of the data beyond the first two moments. For example, burst activity might also be a relevant property of brain networks.[57]

Besides these limitations inherent in the analysis approach, there are also some related to the experimental design. As this is a study containing invasive LFP recordings, we can never have a healthy control group. The medication led to a marked motor improvement in these patients based on the UPDRS, but the patients still have impairments. To differentiate pathological and physiological network activity, we had to rely on the temporal properties of the networks. A further limitation was that all our recordings were made under resting conditions, preventing us from discerning the functional role of oscillations within the discovered networks. We opted for the current design because resting-state data allows the study of networks independent of a task and because using a specific task bears the risk that the patients are not able to properly perform it. Nevertheless, future studies should analyse the behaviour of specific networks using tasks to probe them.

5 Summary

Using a data-driven machine learning approach, we identified three distinct networks (states) that captured differential effects of dopaminergic medication on spectral connectivity in PD. Our findings uncovered a hyper-dopaminergic state that captured the potentially adverse effects of increased dopamine levels due to dopaminergic medication. Furthermore, a communication state was identified that maintained spatio-spectrally selective cortico–STN connectivity ON medication. We also found a local STN–STN coherent state, pointing towards the limited effect of dopaminergic medication to modify local basal ganglia activity. Our findings bring forth a

dynamical systems perspective for differentiating pathological versus physiologically-relevant spectral connectivity in PD. Furthermore, we were able to uncover differential changes induced by altered levels of a neuromodulator such as dopamine in a completely data-driven manner without providing detailed information about large-scale dopaminergic networks to the HMM. This shows another advantage of our dynamical systems-level approach. Furthermore, our whole brain–STN approach provides novel electrophysiological evidence of distributed changes due to dopaminergic medication in brain connectivity, extending previous pairwise connectivity results reported in PD.

Acknowledgements

EF gratefully acknowledges support by the Volkswagen Foundation (Lichtenberg program 89387). Computational support and infrastructure was provided by the "Centre for Information and Media Technology" (ZIM) at the University of Düsseldorf (Germany). We would like to thank Johannes Pfeifer for his valuable feedback on the manuscript.

Funding

This study was funded by the Volkswagen Foundation (Lichtenberg program 89387).

Competing interests

AS has been serving as a consultant for Medtronic Inc, Boston Scientific, St. Jude Medical, Grünenthal, and has received lecture fees from Abbvie, Boston Scientific, St. Jude Medical, Medtronic Inc, UCB

6   References

1.   Akam T, Kullmann DM. Oscillatory multiplexing of population codes for selective communication in the mammalian brain. *Nat Rev Neurosci*. 2014;15(2):111-122.

2.   Akam T, Kullmann DM. Oscillations and filtering networks support flexible routing of

information. *Neuron*. 2010;67(2):308-320.

3.   Schnitzler A, Gross J. Normal and pathological oscillatory communication in the brain. *Nat Rev Neurosci*. 2005;6(4):285-296.

4.   Keitel A, Gross J. Individual human brain areas can be identified from their characteristic spectral activation fingerprints. Engel AK, ed. *PLoS Biol*. 2016;14(6):e1002498.

5.   Mellem MS, Wohltjen S, Gotts SJ, Ghuman AS, Martin A. Intrinsic frequency biases and profiles across human cortex. *J Neurophysiol*. 2017;118(5):2853-2864.

6.   Vidaurre D, Hunt LT, Quinn AJ, et al. Spontaneous cortical activity transiently organises into frequency specific phase-coupling networks. *Nat Commun*. 2018;9(1):1-13.

7.   Marreiros AC, Cagnan H, Moran RJ, Friston KJ, Brown P. Basal ganglia-cortical interactions in Parkinsonian patients. *Neuroimage*. 2013;66:301-310.

8.   van Wijk BCM, Beudel M, Jha A, et al. Subthalamic nucleus phase-amplitude coupling correlates with motor impairment in Parkinson's disease. *Clin Neurophysiol*. 2016;127(4):2010-2019.

9.   West TO, Berthouze L, Halliday DM, et al. Propagation of beta/gamma rhythms in the cortico-basal ganglia circuits of the parkinsonian rat. *J Neurophysiol*. 2018;119(5):1608-1628.

10.   Engel AK, Fries P. Beta-band oscillations—signalling the status quo? *Curr Opin Neurobiol*. 2010;20(2):156-165.

11.   Gershman SJ, Uchida N. Believing in dopamine. *Nat Rev Neurosci*. 2019;20(11):703-714.

12.   Voon V, Fernagut PO, Wickens J, et al. Chronic dopaminergic stimulation in Parkinson's disease: from dyskinesias to impulse control disorders. *Lancet Neurol*. 2009;8(12):1140-1149.

13.   Vidaurre D, Quinn AJ, Baker AP, Dupret D, Tejero-Cantero A, Woolrich MW. Spectrally resolved fast transient brain states in electrophysiological data. *Neuroimage*. 2016;126:81-95.

14. Litvak V, Jha A, Eusebio A, et al. Resting oscillatory cortico-subthalamic connectivity in patients with Parkinson's disease. *Brain*. 2011;134(2):359-374.

15. Hirschmann J, Özkurt TE, Butz M, et al. Differential modulation of STN-cortical and cortico-muscular coherence by movement and levodopa in Parkinson's disease. *Neuroimage*. 2013;68:203-213.

16. Tadel F, Baillet S, Mosher JC, Pantazis D, Leahy RM. Brainstorm: A user-friendly application for MEG/EEG analysis. *Comput Intell Neurosci*. 2011;2011.

17. Gross RE, Krack P, Rodriguez-Oroz MC, Rezai AR, Benabid AL. Electrophysiological mapping for the implantation of deep brain stimulators for Parkinson's disease and tremor. *Mov Disord*. 2006;21(SUPPL. 14):259-283.

18. Moran A, Bar-Gad I, Bergman H, Israel Z. Real-time refinement of subthalamic nucleus targeting using Bayesian decision-making on the root mean square measure. *Mov Disord*. 2006;21(9):1425-1431.

19. Colclough GL, Brookes MJ, Smith SM, Woolrich MW. A symmetric multivariate leakage correction for MEG connectomes. *Neuroimage*. 2015;117:439-448.

20. Vidaurre D, Abeysuriya R, Becker R, et al. Discovering dynamic brain networks from big data in rest and task. *Neuroimage*. 2018;180(Pt B):646-656.

21. Förstner W, Moonen B. A metric for covariance matrices. In: *Geodesy-The Challenge of the 3rd Millennium*. Springer Berlin Heidelberg; 2003:299-309.

22. Fraser AM. *Hidden Markov Models and Dynamical Systems*. SIAM; 2008.

23. Boon LI, Geraedts VJ, Hillebrand A, et al. A systematic review of MEG-based studies in Parkinson's disease: The motor system and beyond. *Hum Brain Mapp*. 2019;40(9):2827-2848.

24. Olde Dubbelink KTE, Stoffers D, Deijen JB, Twisk JWR, Stam CJ, Berendse HW. Cognitive decline in Parkinson's disease is associated with slowing of resting-state brain activity: A longitudinal study. *Neurobiol Aging*. 2013;34(2):408-418.

25. Olde Dubbelink KTE, Stoffers D, Deijen JB, et al. Resting-state functional connectivity as a marker of disease progression in Parkinson's disease: A longitudinal MEG study. *NeuroImage Clin*. 2013;2(1):612-619.

26. Kelly C, De Zubicaray G, Di Martino A, et al. L-dopa modulates functional connectivity in striatal cognitive and motor networks: A double-blind placebo-controlled study. *J Neurosci*. 2009;29(22):7364-7378.

27. MacDonald PA, Monchi O. Differential effects of dopaminergic therapies on dorsal and ventral striatum in Parkinson's disease: Implications for cognitive function. *Parkinsons Dis*. 2011;2011:1-18.

28. Lalo E, Thobois S, Sharott A, et al. Patterns of bidirectional communication between cortex and basal ganglia during movement in patients with Parkinson disease. *J Neurosci*. 2008;28(12):3008-3016.

29. Oswal A, Brown P, Litvak V. Movement related dynamics of subthalmo-cortical alpha connectivity in Parkinson's disease. *Neuroimage*. 2013;70:132-142.

30. MacDonald PA, MacDonald AA, Seergobin KN, et al. The effect of dopamine therapy on ventral and dorsal striatum-mediated cognition in Parkinson's disease: Support from functional MRI. *Brain*. 2011;134(5):1447-1463.

31. Cools R, Clark L, Owen AM, Robbins TW. Defining the neural mechanisms of probabilistic reversal learning using event-related functional magnetic resonance imaging. *J Neurosci*. 2002;22(11):4563-4567.

32. Ray N, Strafella AP. Dopamine, reward, and frontostriatal circuitry in impulse control disorders in Parkinson's disease: Insights from functional imaging. *Clin EEG Neurosci*. 2010;41(2):87-93.

33. Oswal A, Beudel M, Zrinzo L, et al. Deep brain stimulation modulates synchrony within spatially and spectrally distinct resting state networks in Parkinson's disease. *Brain*. 2016;139(5):1482-1496.

34. Hammond C, Bergman H, Brown P. Pathological synchronization in Parkinson's disease: Networks, models and treatments. *Trends Neurosci*. 2007;30(7):357-364.

35. Little S, Tan H, Anzak A, Pogosyan A, Kühn A, Brown P. Bilateral functional connectivity of the basal ganglia in patients with Parkinson's disease and its modulation by dopaminergic treatment. *PLoS One*. 2013;8(12):e82762.

36. Marinelli L, Quartarone A, Hallett M, Frazzitta G, Felice Ghilardi M, Ghilardi MF. *The Many Facets of Motor Learning and Their Relevance for Parkinson's Disease*. Vol

128. Elsevier Ireland Ltd; 2017:1127-1141.

37.   West T, Farmer S, Berthouze L, et al. The parkinsonian subthalamic network: Measures of power, linear, and non-linear synchronization and their relationship to l-dopa treatment and off state motor severity. *Front Hum Neurosci*. 2016;10(OCT2016):517.

38.   George JS, Strunk J, Mak-Mccully R, Houser M, Poizner H, Aron AR. Dopaminergic therapy in Parkinson's disease decreases cortical beta band coherence in the resting state and increases cortical beta band power during executive control. *NeuroImage Clin*. 2013;3:261-270.

39.   Whalen XTC, Willard AM, Rubin JE, Gittis AH. Delta oscillations are a robust biomarker of dopamine depletion severity and motor dysfunction in awake mice. *J Neurophysiol*. 2020;124(2):312-359.

40.   Jubault T, Monetta L, Strafella AP, Lafontaine AL, Monchi O. L-dopa medication in Parkinson's disease restores activity in the motor cortico-striatal loop but does not modify the cognitive network. Gribble PL, ed. *PLoS One*. 2009;4(7):e6154.

41.   Tinkhauser G, Pogosyan A, Tan H, Herz DM, Kühn AA, Brown P. Beta burst dynamics in Parkinson's disease OFF and ON dopaminergic medication. *Brain*. 2017;140(11):2968-2981.

42.   Cools R. Enhanced or impaired cognitive function in Parkinson's disease as a function of dopaminergic medication and task demands. *Cereb Cortex*. 2001;11(12):1136-1143.

43.   Vaillancourt DE, Schonfeld D, Kwak Y, Bohnen NI, Seidler R. Dopamine overdose hypothesis: Evidence and clinical implications. *Mov Disord*. 2013;28(14):1920-1929.

44.   Cools R. Dopaminergic modulation of cognitive function-implications for L-DOPA treatment in Parkinson's disease. *Neurosci Biobehav Rev*. 2006;30(1):1-23.

45.   Shohamy D, Myers CE, Grossman S, Sage J, Gluck MA. The role of dopamine in cognitive sequence learning: Evidence from Parkinson's disease. *Behav Brain Res*. 2005;156(2):191-199.

46.   Harmony T. The functional significance of delta oscillations in cognitive processing. *Front Integr Neurosci*. 2013;7(DEC):83.

47.   Zavala BA, Tan H, Little S, et al. Midline frontal cortex low-frequency activity drives subthalamic nucleus oscillations during conflict. *J Neurosci*. 2014;34(21):7322-7333.

48.   Cruz A V., Mallet N, Magill PJ, Brown P, Averbeck BB. Effects of dopamine depletion on network entropy in the external globus pallidus. *J Neurophysiol*. 2009;102(2):1092-1102.

49.   Cagnan H, Duff EP, Brown P. The relative phases of basal ganglia activities dynamically shape effective connectivity in Parkinson's disease. *Brain*. 2015;138(6):1667-1678.

50.   Holt AB, Kormann E, Gulberti A, et al. Phase-dependent suppression of beta oscillations in parkinson's disease patients. *J Neurosci*. 2019;39(6):1119-1134.

51.   Gertz H, Fiehler K. Human posterior parietal cortex encodes the movement goal in a pro-/anti-reach task. *J Neurophysiol*. 2015;114(1):170-183.

52.   Siegel M, Buschman TJ, Miller EK. Cortical information flow during flexible sensorimotor decisions. *Science (80-)*. 2015;348(6241):1352-1355.

53.   Gallivan JP, Chapman CS, Wolpert DM, Flanagan JR. Decision-making in sensorimotor control. *Nat Rev Neurosci*. 2018;19(9):519-534.

54.   Martínez-Vázquez P, Gail A. Directed interaction between monkey premotor and posterior parietal cortex during motor-goal retrieval from working memory. *Cereb Cortex*. 2018;28(5):1866-1881.

55.   Shimamoto SA, Ryapolova-Webb ES, Ostrem JL, Galifianakis NB, Miller KJ, Starr PA. Subthalamic nucleus neurons are synchronized to primary motor cortex local field potentials in Parkinson's disease. *J Neurosci*. 2013;33(17):7220-7233.

56.   Baker AP, Brookes MJ, Rezek IA, et al. Fast transient networks in spontaneous human brain activity. *Elife*. 2014;2014(3).

57.   Florin E, Watanabe M, Logothetis NK. The role of sub-second neural events in spontaneous brain activity. *Curr Opin Neurobiol*. 2015;32:24-30.

FIGURE DESCRIPTIONS

Figure 1:

Title: Data-driven frequency modes derived from the OFF and ON medication conditions.

Description: Each plotted curve shows a different spectral band. The x-axis represents frequency in Hz and the y-axis represents the weights obtained from the non-negative matrix factorisation (NNMF) factorisation in arbitrary units. The weights can be considered as amplitudes corresponding to the frequency bins. The frequency resolution of the modes is 0.5 Hz. Panels A and B show the OFF and ON medication frequency modes, respectively.

Figure 2:

Title: The hyper-dopaminergic state

The hyper-dopaminergic state was characterised by a significant increase in coherence ON compared to OFF medication (see Panel B). Due to this, no connections within the alpha and beta band ON medication were significantly higher than the mean (Panel C). However, in the delta band, ON medication medial prefrontal-orbitofrontal connectivity emerged.

Description: A and C: Each node in the circular graph represents a brain region based on the Mindboggle atlas. The regions from the atlas are listed in Table 1 along with their corresponding numbers that are used in the circular graph. The colour code in the circular graph represents a group of regions clustered according to the atlas (starting from node number 1) STN contacts (contacts 1, 2, 3 = right STN and contacts 4, 5, 6 = left STN), frontal, medial frontal, temporal, sensorimotor, parietal, and visual cortices.

In the circular graph, only the significant connections (p < 0.05; corrected for multiple comparisons, IntraMed analysis) are displayed as black curves connecting the nodes. The circles from left to right represent the delta/theta, alpha, and beta bands. Panel A shows results for OFF medication data and Panel C for the ON medication condition.

Panel B shows the result for inter-medication analysis (InterMed) for the hyper-dopaminergic state. In each symmetric matrix, every row and column corresponds to a specific atlas cluster denoted by the dot colour on the side of the matrix. Each matrix entry is the result of the InterMed analysis where OFF medication connectivity between ith row and jth column was compared to the ON medication connectivity between the same connections. A cell is white if

the comparison mentioned on top of the matrix (either ON > OFF or OFF > ON) was significant at a threshold of $p < 0.01$.

Figure 3:

Title: The communication state

Description: For the general description, see the note to Figure 2.

The communication state was characterised by preservation of spectrally selective cortico–STN connectivity ON medication. Also, ON medication, a sensorimotor–frontoparietal network emerged.

Figure 4:

Title: The local state

Description: For the general description, see the note to Figure 2.

The local state was characterised by preservation of STN–STN coherence in the alpha and beta band OFF versus ON medication. STN–STN theta/delta coherence was no longer significant ON medication.

Figure 5:

Title: Temporal properties of states

Description: Panel A shows the fractional occupancy for the three states for the hyper-dopaminergic (hyper-DA), communication, and the local state. Each point represents the mean for a state and the error bar represents standard error. Orange denotes ON medication data and blue OFF medication data.

Panel B shows the mean interval of visits (in milliseconds) of the three states ON and OFF medication.

Panel C shows the lifetime (in milliseconds) for the three states. Figure insets are used for clarity in case error bars are not clearly visible. The y-axis of each figure inset has the same units as the main figure.

Tables:

Table 1 Regions of the Mindboggle atlas used

STN- Subthalamic nucleus, Vis- Visual, Par-Parietal, Smtr-Sensory motor, Tmp-Temporal, Mpf-Medial prefrontal, Frnt-Frontal, Ctx- Cortex
The colour code is for the ring figures presented as part of the results.

STN	1	Contact 1 right	Smtr-Ctx	12	Postcentral
	2	Contact 2 right		13	Precentral
	3	Contact 3 right	Tmp-Ctx	14	Middle temporal
	1	Contact 4 left		15	Superior temporal
	2	Contact 5 left	Mpf-Ctx	16	Caudal middle frontal
	3	Contact 6 left		17	Medial orbitofrontal
Vis-Ctx	4	Cuneus		18	Medial orbitofrontal
	5	Lateral occipital		19	Lateral orbitofrontal
	6	Lingual	Frnt-Ctx	20	Pars opercularis
Par-Ctx	7	Inferior parietal		21	Pars orbitalis
	8	Para central		22	Pars triangularis
	9	Precuneus		23	Rostral middlefrontal
	10	Superior parietal		24	Superior frontal
	11	Supramarginal			

Supplementary Material

Figure S1-S3:

S1-S3 show the three additional states that were found in the OFF and ON conditions. S1-S3 A show the OFF states and S1-S3 B show their distance-matched ON states. To the best of our knowledge, we could not interpret these states within our current physiological understanding of Parkinson's disease.

A Fractional occupancy

B Interval of visit

C Lifetime