Human cerebral cortex networks use expanding and contracting state dynamic to shape cortical functions

Alex Willumsen[1], Jens Midgaard[1], Bo Jespersen[2], Christoffer K.K. Hansen[1], Salina N. Lam[1], Sabine Hansen[1], Ron Kupers[1,3], Martin E. Fabricius[4], Minna Litman[5], Lars Pinborg[5,6], José D.Tascón-Vidarte[3], Anne Sabers[5], Per E. Roland[1*]

[1] Dept. Neuroscience, Panum Institute, University of Copenhagen, Denmark
[2] Department of Neurosurgery, Rigshospitalet, University Hospital of Copenhagen, Denmark
[3] DIKU, Dept. Computer Sciences, University of Copenhagen, Denmark
[4] Department of Clinical Neurophysiology, Rigshospitalet, University Hospital of Copenhagen, Denmark
[5] Epilepsy Clinic, Department of Neurology, Rigshospitalet, University Hospital of Copenhagen, Denmark
[6] Neurobiology Research Unit, Department of Neurology, Rigshospitalet, University Hospital of Copenhagen, Denmark

Short title: Fluctuating attractors in the human brain

* Correspondence to: Per E. Roland, Department of Neuroscience, Panum Institute, University of Copenhagen, Blegdamsvej 3B, 2200 Copenhagen, Denmark. Email: per.ebbe.roland@sund.ku.dk

Summary

We lack viable explanations of how collective activities of neurons in networks produce brain functions. We recorded field potentials from many local networks in the human cerebral cortex during a wide variety of brain functions. The network dynamics showed that each local cortical network produced fluctuating attractor states. The state trajectories continuously stretched and contracted during all brain functions, leaving no stable patterns. Different local networks all produced this dynamic, despite different architectures. Single trial stimuli and tasks modified the stretching and contractions. These modified fluctuations cross-correlated among particular networks during specific brain functions. Spontaneous activity, rest, sensory, motor and cognitive functions all emerged from this dynamic. Its mathematical structure provides a general theoretical model of cortical dynamics that can be tested experimentally. This universal dynamic is a simple functional organizing principle for brain functions at the mm^3 scale that is distinct from existing frameworks.

Keywords: cortical dynamics, brain theory, single trials, mental imagery, perception, attractor volume, excitation-inhibition, fluctuations and oscillations, entrainment, spontaneous activity

Human cerebral cortex networks use expanding and contracting state dynamic to shape cortical functions

Graphical abstract

In brief

Willumsen et al. developed a method to show that local cortical networks contribute to sensory, motor and cognitive functions by stretching and contracting the trajectory of the multidimensional field potential. In single trials the networks communicate by cross-correlating the stretching and contracting. This ubiquitous attractor dynamic forms a departure from existing models of how postsynaptic dynamics contribute to sensory, motor and cognitive brain functions.

Highlights

- Cortical fluctuating expanding and contacting attractor dynamics (FECAT) drive collective postsynaptic operations at the mm^3 scale

- FECAT dynamic accounted for all behavioral conditions and all tested cortical areas

- Cortical states show no stationary patterns, but continuously expand and contract with a stable attractor dynamic

- Our method reveals multi-dimensional cortical dynamics in field potentials, also useful for EEG and MEG

Despite numerous efforts over the past decades, there is no consensus theory or working hypothesis yet, which adequately explains how the collective action of neurons in local networks can create brain functions (Breakspear, 2017; Buzsaki, 2019; McCormick et al., 2020; Singer et al., 2019). Here we examine whether there is any order or systematic rules in the collective postsynaptic actions at local cortical network scale. We chose this spatial scale because brain functions are assumed to emerge from collective actions of large populations of neurons (Breakspear, 2017; Buzsaki, 2019; McCormick et al., 2020; Pitkow and Angelaki, 2017; Rabinovich et al., 2012; Roland, 2017; Singer et al., 2019).

In each local cortical network, the activities of millions of synapses create equally many postsynaptic currents (Iacaruso et al., 2017). These give orders of magnitude larger number of combinatorial possibilities of the network. Yet, local cortical networks reliably create perception in a few hundred ms (Thorpe et al., 1996). This makes one suspect that there are organizing principles or dynamic rules controlling how the postsynaptic currents efficiently combine in the network. Here we search for such rules. The postsynaptic membrane currents collectively generate field potentials (Einevoll et al., 2014; Pesaran et al., 2018). The analysis of field potentials, often separated into different frequency bandwidths, has provided important new insights into the functions of hippocampal networks and some large-scale cortical operations (Buzsaki, 2019; Muller et al., 2018; Pesaran et al., 2018; Singer et al. 2019). However, this type of analysis cannot capture the dynamics of the cortical states hidden in the field potential, for example in single trials (Babloyantz and Destexhe, 1986; Breakspear, 2017; Rabinovich et al., 2012; Roschke and Basar,1990; Stam, 1996). First, the field potential is multi-dimensional, because it is created by many currents. Each measurement of the field potential therefore corresponds to one point in a multidimensional space (scaled in μV) that we call a local cortical state of the local network. Second, in this multi-dimensional space, the local cortical states evolve one after another, and most likely in a non-linear manner (Babloyantz and Destexhe, 1986; Breakspear, 2017; Rabinovich et al., 2012).

We measured field potentials in many cortical regions while patients performed sensory, motor and cognitive tasks, and also during spontaneous activity. Thereafter, we extracted multi-dimensional local cortical states from each local cortical network to capture the dynamic and non-linear progress of states in each network (Takens, 1981). We looked for possible dynamic rules driving the temporal evolution of local cortical states in the ms to s time scales. Dynamical rules, in this context, are succinct, exhaustive, often mathematical descriptions of how cortical states in local networks evolve (Babloyantz and Destexhe, 1986; Breakspear, 2017; Rabinovich et al., 2012; Roschke and Basar, 1990; Stam, 1996; Takens, 1981). In case there are several rules driving cortical states, we use the term dynamics and reserve the term dynamic to mean one rule only.

We did find systematic order in the way cortical states from local networks evolved. All probed local networks used only one dynamic rule to create local cortical states. The succession of states evolved continuously, irregularly stretching and contracting their trajectory. Trajectories were located to an attractor that consequently expanded and contracted continuously. That is, fluctuating expanding and contracting attractor

dynamic determined how local cortical states evolved. This rule has implications for theoretical and experimental neuroscience. For example, it explains the nature of the continuous flow of states under all conditions, how spontaneous ongoing cortical states and sensory-evoked cortical states can arise from the same dynamic, and how different brain functions may emerge. It has mathematical structure and could serve as a theoretical model for studies of larger scale brain functions.

Figure 1. Logic of the analysis and definition of fluctuating attractor. A. Electrode lead with field potential *f*(*t*) sampled at 512 Hz. To find the dynamics of the local networks we must first know the *f*(*t*) dimensionality. Although the field potential is created mainly by currents from multiple dendrites of many neurons, the collective result of these many currents may be of lower dimensionality. We used embedding to estimate the dimensionality. Thereto, we constructed a vector *F*(*t*) out of 20 copies of *f*(*t*) delayed by successive multiples of τ ms. The dimensionality of *F*(*t*) is the number of dimensions needed to achieve an exhaustive description of *F*(*t*). We then calculated the correlation dimensionality from this 20-dimensional vector *F*(*t*). **B.** If, for example, the estimated dimensionality of *F*(*t*) is 3.6 we can get a good picture of the evolution of local cortical states in a 3-dimensional state space (in reality the dimensionality ranged from 3.5 to 9). In state space, each point is one local cortical state. We followed the progress of local cortical states every 1.95 ms (1/512 s) forming the blue and red parts of the trajectory. In each local network, one attractor confines the states for each behavioral condition to one hyper-elliptic part of state space. **C.** The hyper-ellipsoid expands and contracts, making the volume to enlarge and diminish. We call this a fluctuating expanding and contracting attractor (FECAT).

RESULTS

We recorded field potentials from multiple sites in the cerebral cortex of 13 patients with medical intractable epilepsy (**STAR Methods**). In a local cortical network, the field potential is dominated by transmembrane currents from more than 10^5 dendrites and their subsequent extracellular currents (Einevoll et al., 2014; Pesaran et al., 2018; **STAR Methods**).

Each field potential time-series reflects the underlying dynamics of the local network in the immediate neighborhood of the electrode lead at which it is recorded. Each analyzed time series was 60s long (**STAR Methods; Figure S1**). All epileptic activity was removed and the analysis was restricted to those leads and time series that remained without any abnormality throughout the experiments. This left 3476 60s files of normal field potential recordings, distributed over 290 electrode leads under 40 different conditions. These conditions comprised visual, auditory, gustatory, olfactory and somatosensory stimulations, as well as tests with voluntary movements in which the dynamics of the brain operations was driven partly by external cues or stimuli. We also included memory retrieval and mental navigation tasks that did not involve external stimuli and overt behavior (**STAR Methods**).

Dimensionality of field potential dynamics

The dimensionality is the number of dimensions we need to get an exhaustive account of the information about cortical states in the field potential (Roschke and Basar, 1990; Stam, 1996). Although the field potential is multidimensional, its true dimensionality is unknown. We therefore need first to determine the number of dimensions needed to create the multidimensional space, in which we can study how the cortical states produced by each local network evolve. First, we embedded the field potential (Takens, 1981) (**Figures 1 and S1; STAR Methods**). Then, we estimated its correlation dimensionality. To find the dynamics driving the evolution of the multi-dimensional field potential in one local network implies to demonstrate how the local cortical states produced by the local network evolve in a multidimensional state space of sufficient dimensions. A local cortical state is one point in this multidimensional state space. In the state spaces in this study, one point represents the state of the local network over 1.95 ms (**Figure 1**). Despite the intrinsic high dimensionality of the sources of the field potential, the estimated dimensionality was always comparatively low, ranging from 3 to 9 (**Figures 2 and S2**).

The local correlation dimensionalities in awake conditions fell into two clusters, one with relatively high means and low standard deviations, the other with lower means and higher standard deviations (**Figures 2A and S2**). The validity of the clustering is apparent from the silhouette plots (**STAR Methods**) in **Figure S3.**
The difference between the two clusters was also tested by traditional statistical tests (**STAR Methods**) and was significant (p< 0.02 or less, except for patient 11; **Table S1).** High dimensionality is associated with high complexity of dynamics and low dimensionality with lower complexity (Roschke and Basar, 1990; Stam, 1996). The estimated dimensionality for each condition and electrode lead was used to build a state space of that dimensionality, in which we studied the cortical states evolving from the local cortical network (**Figure 1; STAR Methods**).

Figure 2. Correlation dimensionalities of field potentials. A. Correlation dimensions as function of anatomical location for all tests of patient 17. Red star: median. Red or blue dots: mean, depending on membership of cluster (**Figures S2 and S3)**. Standard deviation shown as error bars. Abbreviations: amy: amygdala; hip: hippocampus; phg: parahippocampal gyrus; fus: fusiform gyrus; ang: angular gyrus; calc: calcarine cortex; prec: precuneus; stg: superior temporal gyrus; mtg: middle temporal gyrus; itg: inferior temporal gyrus; ins: insula; postc: postcentral gyrus; smg: supramarginal gyrus. **B.** Axial slices of the standard brain, showing positions of electrode leads. Green spots for leads from patient 17; other colors refer to other patients.

Fluctuations of the volumes of cortical states

Figure 3 shows how the local cortical networks confine their evolving cortical states to a single region, shaped like a hyper-ellipsoid in state space. All cortical states in all patients (3476 60s-epochs, distributed over 290 leads, giving a total of 106782720 states) were located inside the enclosing volume of the hyper-ellipsoid in **Figure 3A.** We found qualitatively identical, albeit smaller, hyper-ellipsoids at all electrode leads and under all experimental conditions (**STAR Methods**). Inside each hyper-ellipsoid, the temporal succession of states formed a trajectory (**Figures 3B and S4**). Thus, the local dynamics measured at every cortical location and under every condition was restricted to a single hyper-elliptic region of its state space.

Data for 60 s or more are necessary to compute a reliable estimate of the correlation dimensionality (**Figure S1**), while the field potential cannot be regarded as stationary for so long. There is then a need for evaluating the dynamics at shorter, physiologically and behaviorally relevant, timescales. Consequently, we made a principal component analysis of the embedded trajectory of appropriate dimensionality. Then we estimated the volume of successive local states accumulated over 50 ms with a sliding time window of 250 ms (for choice of parameters see **STAR Methods**). The cortical states in such short time intervals form short trajectories, rather than hyper-ellipsoids (**Figures 3 and S4**). To follow fast changes in the local states at each lead and condition, we estimated these state space changes as volumes of hyper-rectangles (**STAR Methods**).

Figure 3. Each local cortical network constrains dynamics to one fluctuating expanding and contracting attractor. **A.** State space containing all normal states (n= 106,782,720) in all patients and conditions giving 3476 enclosing volumes (one color for each electrode-condition combination, but only the largest is visible; **STAR Methods**). Together, all enclosing volumes occupy a single, limited part of state space in shape of a hyper-ellipsoid. **B.** Single condition hyper-ellipsoid. Six embedding dimensions of **F**(t) from the posterior part of the superior temporal sulcus, reconstructed from 60 s trajectory during identification of emotional faces (patient 5, lead 24). Left: projections of the first 3 dimensions (dim). Right: dimensions 4,5,6. Red: trajectory temporarily exploiting nearby state space. Blue: trajectory inside the state space defined by the hyper-ellipsoid enclosing volume $V_{95\%}$ (**STAR Methods**). **C.** Fluctuations of the volume calculated by equation 4 (**STAR Methods**) of local cortical states (attractor states) during 60 s of rest. Note $\log_{10}$ scale on the y-axis. **D.** Distribution of volumes (equation 4) of fluctuating attractor. X-axis $\log_{10}V$ (orders of magnitude) aligned to the median.

Left: patient 12 during 20 randomly selected 60 s episodes of awake spontaneous activities. Right: all experimental conditions in awake patients (n=3363). **E.** Projections of first 3 dimensions of 5s trajectories from the parietal operculum during classifying nouns (patient 17). Fluctuating hyper-elliptic attractor of big, medium and small volume (left, center and right, respectively). Sections of trajectory exceeding Mahalanobis distances > 90% from the center of gravity are shown in red. Start and end of 5 s episode trajectory are shown in dark blue and yellow, respectively. **F.** Projections of first 3 dimensions of 1 s trajectories from fusiform gyrus during recall of visual events at admission day (patient 12). Left: big volume trajectory of dimensions 1-3 (upper part) and 4-6 (lower part); Right: small volume trajectory. Conventions as in E. **G.** Distributions (log normal, **Figure S4**) of times to return for outside states to the enclosing $V_{90\%}$, when $V_{90\%}$ is big (top), medium (middle) and small (bottom).

Note that we use two complementary methods to estimate two different volumes of the local cortical states. The hyper-ellipsoid method was used for successions of cortical states 1s and longer (**Figure 3A, B, E, F**). The hyper-rectangle method was used for shorter successions of cortical states (equation 4, **STAR Methods**; **Figures 3C and 3D**). During ongoing autonomous activity only (rest condition), the enclosing hyper-rectangle volume continuously expanded and contracted (**Figure 3C**). In all (n=110) realizations of rest conditions, the enclosing volume continuously expanded and contracted irregularly. These expansions and contractions were also the dominant qualitative feature at all cortical sites and under all tests and conditions (n=3476) (**Figures 3D and S5; database)**. For comparison, we also plotted the similar volume fluctuations from 20 randomly selected 60 s episodes during spontaneous behavior of one awake patient when not engaged in any test. The ranges of the enclosing volume (from minimum to maximum) were at least four orders of magnitude, with no difference in the ranges between spontaneous behavior and experimental conditions (**Figure 3D right and left).** Fluctuations of the (one-dimensional) cortical field potential are well known (Einevoll et al., 2014; Pesaran et al., 2018). However, that these vigorous fluctuations should be the dominant feature of the dynamics of evolving cortical states at full dimensionality was not foreseen.

Over shorter (1s, 5s) time scales, the flow of states can be localized to well-defined trajectories, as shown in the projections in **Figures 3E and 3F.** The volume fluctuations were irregular (**Figure S5; database**) meaning that in each cortical network, a shorter succession of cortical states expanded and contracted irregularly. The short succession of states stretched the trajectory, thereby increasing the volume enclosing the multi-dimensional trajectory, and contracted the multi-dimensional trajectory of the states, thereby decreasing the enclosing volume. These volume changes were estimated as the volumes of the enclosing hyper-ellipsoid or hyper-rectangle containing a short succession of states. Equivalently, one can describe the expansion and contraction as stretching and contraction of the multidimensional trajectory forming the succession of states. These expansions and contractions continued under all conditions **(database)**. These volume fluctuations were not very fast (50 ms to a few hundreds of ms). When the volume increases, the multi-variance of the cortical states and their trajectory increases. When the volume decreases, the multi-variance of the states and their trajectory decreases (for details and flow of states, see **STAR Methods**).

Attractor properties of the dynamics

Fast volume fluctuations of four orders of magnitude indicate that the local networks they originate from produce unstable dynamics. For this reason, we examined the stability of the evolving cortical states. In **Figure 3B** we set the enclosing volume as the volume of the hyper-ellipsoid enclosing all states whose Mahalanobis distances from the center lie inside the 95% quantile (**STAR Methods**). As the outside states quickly returned to the 95% enclosing volume, we suspected that these observed dynamics could be attractor-like. Therefore, we examined the fate of outside states at all time scales up to 60 s. In addition, we examined outside states when the enclosing volume, because of its fluctuations, was small, medium and large (for definitions see **STAR Methods**). We defined the putative attractor as the set of states whose Mahalanobis distances from the center of the enclosing hyper-elliptic volume were less than 90% of the maximum measured value. Thus, large volumes, medium-sized volumes and small volumes all had by definition 10% of states as outliers (**Figures 3E and 3F**). Most of these states returned to the putative attractor within 50 ms, at a different position than that from which they had departed, as is typical for a dynamical system (**Figures 3G and S4**). Changing the definition of the outside states to 20% or 5% did not change the distribution of the return times. The states evolved within a hyper-elliptic region in state space. States which occasionally escaped this region were swiftly pulled back by the attractor, irrespective of the size of the hyper-ellipse.

To sum up, each local cortical network produces cortical states. The succession of states forms a trajectory which for 90% is contained inside a hyper-elliptic object. This object attracts states outside and pulls the outside states back. Therefore, it is a kind of attractor. This attractor expands and contracts continuously in state space, thereby expanding and compressing the states it contains. Or, phrased equivalently, stretches and contracts the trajectory it contains (**Figures 3E and 3F**).

These results are compatible with single attractor dynamics. This analysis, however, does not distinguish whether what we defined as outside states in reality were parts of the attractor itself - or departures from the attractor that were rapidly pulled back. None of our experimental conditions succeeded to perturb the attractor systematically, indicating overall stability in cortex under normal conditions. Moreover, the state flows were fast in all dimensions within the state space enclosed by the attractor. This reduces the likelihood that the data supports other, for example multi-stable, dynamics (**Figure S6**). Furthermore, an attractor, as normally defined, has a single invariant dimensionality and occupies an invariant part of state space. What we observed is not an attractor in the normal sense, because of the expansions and contractions in state space**.** To describe this lower-dimensional fluctuating expanding and contracting attracting object, we coined the term fluctuating expanding and contracting attractor (FECAT).

Fluctuating expanding contracting attractor dynamic is a universal large-scale cortical state dynamic

In these 3476 episodes, comprising all tested conditions and all 290 leads we could not find any exception from fluctuating attractor dynamic (**Figures 3A, 3B, 3G, S4** and **S6; database**).

Figure 4. Different cortical network architectures create cortical states with FECAT-dynamic. Projections of attractor trajectories over 60 s; Red curves: trajectories temporarily exploiting nearby state space. Blue curves: trajectories inside the state space defined by the enclosing hyper-elliptic volume $V_{95\%}$. Below: fluctuations of the volume of cortical states (equation 4, **STAR Methods**) during the same 60 s. **A.** Stereognosis (patient 6). **B.** Conversation (patient 4). **C.** Ortho-nasal smell (patient 16). **D.** Motor test 6 (patient 17). **E.** Visual recognition (patient 17). **F.** Memory retrieval, mental navigation (patient 5). **G.** Watching TV (patient 12). **H.** Eating (patient 14). **I.** Facial emotion identification (patient 4).

This means that local networks from different cortical locations and within different cortical areas used the same type of dynamic to drive their cortical states. Indeed, local cortical networks in prefrontal cortex, primary visual cortex, supramarginal gyrus, and hippocampus, have widely different anatomical and neurotransmitter receptor architectures (Elston, 2003; Zilles and Amounts 2010). Notwithstanding these architectonic differences, the local networks created cortical states using FECAT-dynamic (**Figure 4; database).**

Figure 5. Different brain functions have cortical states emerging from FECAT dynamic. Projections of attractor trajectories over 60s; Red curves: trajectories temporarily exploiting nearby state space. Blue curves: trajectories inside the state space defined by the enclosing hyper-elliptic volume $V_{95\%}$ (**STAR Methods**). Below: fluctuations of the volume of cortical states (equation 4, **STAR Methods**) during the same 60 s. **A.** Control (rest) condition, superior temporal gyrus, anterior part (patient 4). **B.** Mental navigation (memory retrieval test 8), posterior hippocampus (patient 6). **C.** Memory retrieval test 10 (admission day), frontal operculum (patient 12). **D.** Classification of nouns, parahippocampal gyrus (patient 8). **E.** Somatosensory attention, anterior insula (patient 16). **F.** Watching TV, superior frontal gyrus,

anterior part (patient 9). **G.** Visual motion to right (visual test 7), posterior hippocampus (patient 15). **H.** Visual test 4, middle part of inferior frontal gyrus (patient 14). Note entrainment. **I.** Recognition (visual test 6), subiculum (patient 10). **J.** Taste, amygdala (patient 16).

Similarly, during all tested conditions, cortical states evolved under the rule of FECAT-dynamic. For example, cortical states during spontaneous behavior, rest condition and during different brain functions such as mental navigation, classification of nouns, somatosensory attention, tasting sweets etc., were all products of FECAT-dynamic (**Figure 5; database**). The different conditions and brain functions produced somewhat different patterns of expansions and contractions (**Figures 5 and S5, database).** However, single trial tests were associated with a single expansion and contraction.

Attractor size fluctuations reshape to single trial input

Single trial tasks reshaped the ongoing expansions and contractions, but did not perturb the dynamic. The volume fluctuations of the attractor are thus sensitive to single trial input, for example, asking to imagine the face of a well-known person. This elicited a small expansion followed by a slightly longer contraction modifying the ongoing fluctuations (**Figure 6A and 6B).** Similarly, a small expansion followed by a contraction was also observed in single trial tasks in which patients were asked to identify facial emotional expressions, imagine well-known landmarks or tourist attractions, judge whether spoken nouns were concrete or abstract, look at geometrical shapes, and recognize natural scenes (**Figures 5, 6F and S5**). Each patient did from 2 to 7 such single trial tests. This suggests that a single trial engagement is associated with an expansion followed by a contraction of the volume of the fluctuating attractor. This is compatible also with the stretching and compression of the trajectories in **Figures 3E and 3F**.

But how do the cortical networks work together to perform these brain functions? The face imagery test (**Figures 6A and 6B**) was associated with modified expansions and contractions in many other local networks. In these local networks, the modified expansions and contractions of the fluctuating attractors cross-correlated mutually and cross-correlated with the modified expansion and contraction in **Figure 6A.** These pairwise cross-correlations appeared with different lags (**Figures 6C and 6D**). However, not all local cortical networks had significant cross-correlations of the fluctuations in a given test (**Figure S5**). Thus, only a subset of the many local networks modified and correlated their fluctuating attractor fluctuations during the tests**,** while the remaining networks continued their un-correlated fluctuations (**Figure S5**). These results also held for the other single-trial tests. With few exceptions, each of these tests was associated with a unique combination of cross-correlated cortical local networks (**Table S2**). On average, 43.5±24.8 % (SD) of the local networks engaged with correlated fluctuations in these tests (n=43). This demonstrates the sensitivity of the attractor volume fluctuations to single transients and stimuli. At this macroscopic scale, the single trial tests seem to be operated by different combinations of cortical local networks. The local networks engaged by cross- correlating their single expansion-contraction with different lags.

Figure 6. Single trial expansion and contraction of the fluctuating attractor. A. 4 s average of 11 z-scores of attractor volume (V) showing one expansion and contraction. Average of the sound track below. Each trial aligned to the peak of the sound. **B.** Fluctuations in 11 trials of V of the attractor in $\log_{10}$ scale, recall of faces (memory retrieval test 1, **STAR Methods**). Patient 17, lead 50, cortex in posterior part of the superior temporal sulcus; below:

sound track with peaks indicated by the red stippled lines for each of the names of the 11 well-known persons to imagine (**STAR Methods**). **C.** Statistically significant cross-correlations of average z-scores between leads engaged in the task. Green stippled lines: 95% confidence limits. **D.** z-score averages, with mean in red, of expansions and contractions from the leads engaged in the task. **E.** Pairwise cross-correlations (cc) of the leads in D. **F.** Top: z-score average of V from 9 trials; below: average photodiode signal showing end of last screen presentation, 1 s black screen interruption, and first 4 s of emotional expression to judge. Bottom: fluctuating volume of the attractor in $\log_{10}$ scale with photodiode signal of 9 single trial screen presentations below. Patient. 5, lead 6, 6 dimensions, middle temporal gyrus.

We also observed expansions and contractions comparable to those in **Figure 6** in the absence of sensory stimulation (example in **Figure 3F)**. All expansions and contractions of attractor volumes during all conditions can be found in the **database** (https://github.com/ChristofferKKHansen/RolandLab).

Entrainment of cortical states

Rhythmic visual stimulation entrains the volume fluctuations of the attractors. Entrainment of field potentials in visual cortex of monkeys has been reported (Lakatos et al., 2008). In visual tests 4,5 and 11, patients were instructed to watch the items shown on the monitor, without responding. This elicited an entrainment of the expansions and contractions of the attractor, with a frequency identical to (**Figures 7A and 7F**) or (occasionally) double that of the stimulus presentation rate (**Figure 7G**). Rhythmic visual presentations with no intermission of a black screen (visual test 3), and stimulus durations of 6 s (facial emotions) did not produce entrainment. In contrast, regular presentation of visual stimuli of 125 or 1000 ms duration at 0.5 or 1 Hz did produce entrainment (including visual test 6).
As for the other single trial tests, one stimulus elicited one expansion followed by one contraction (**Figures 7A, 7B, 7E and 7F**). The phase of the expansion-contraction oscillation however varied with cortical location. In some local networks, the expansion led the stimulus presentation (example in **Figures 7A and 7E**), while at other sites it trailed it (**Figures 7B and 7F**). The frequency of these entrained oscillations was strictly locked to the presentation frequency (n= 41 out of 41 tests) (**Figures 7A, 7B, 7E and 7F**) with lags ranging from -500 to + 500 ms (**Figures 7C and 7D).** In these strictly rhythmic trials, anticipation of the next stimulus as well as evoked activity may explain the large range in lags (see also **Figure 6F**) (Gillies et al., 2019; Staresina et al., 2019).

The proportion of leads showing attractor volume entrainment in the four visual tests (4, 5, 6 and 11; **STAR Methods**) ranged from 0% to 100% (**Table S2**). Natural pictures presented for 125 ms at 1 Hz engaged from 91 % to 100% of the leads (mean 97%; n=8). The entrainment was not obvious in the raw data, but it was evident in the stimulus onset time-locked averaged data (**Figures 7A, 7B, 7E and 7F**). This indicates that the system is not simply undergoing oscillatory (limit-cycle) dynamics; the time-locked oscillations here are just another property of the attractor fluctuations.

Figure 7. Entrainment of fluctuating attractors. A. Average z-scores of volume fluctuations over trials with average stimulus presentation. Bottom: 60 s episode of fluctuating size V of the attractor in $\log_{10}$ scale with onset of trials marked with red stippled lines. Natural pictures presented 125 ms with 1 s black screen intermissions (visual test 4). Patient 11 lead 2, anterior part of cingulate gyrus. **B.** Same patient, same test, but lead 101, middle temporal gyrus, posterior part; phase shift of the fluctuating expansions-contractions of the attractor (compared to A). **C.** Pairwise cross-correlations above 70% quantile of average z-scores during the task for the time segment defined by the 2 vertical lines in A. Green stippled lines: 95% confidence limits. **D.** Matrix of the pairwise lags (in ms) for all leads. **E.** Visual recognition test, patient 6, lead 2, anterior part of cingulate cortex. Average of z-scores (top), 60 s episode of recognition of fluctuating size V of the attractor in log10 scale with photodiode signal of picture presentation below. **F.** Fluctuating attractor entrainment of double frequency. Geometrical pictures (visual test 11) patient 5, lead 25, cortex lining posterior part of superior temporal sulcus.

DISCUSSION

Our results provide an experimentally-based overall explanation of how cortical states contained in field potentials from the human cerebral cortex evolve during a wide spectrum of behavioral conditions. All examined local cortical networks produced successions of local cortical states forming trajectories that stretched and contracted

irregularly and unceasingly. The trajectories predominantly resided inside a single multidimensional object, an attractor that continuously expanded and contracted in state space. The states inside this attractor pulled states that temporarily escaped back into the attractor, irrespective of the volume of the attractor. Stimuli, single trial tasks, and other behavioral conditions temporarily reshaped the expansions and contractions and altered the progress velocity of cortical states, but did not radically alter or change this dynamic into other dynamical regimen. The reshaped expansions and contractions cross-correlated between selected subsets of cortical networks, depending on the test. This hitherto unreported fluctuating expanding and contracting attractor dynamic gives a simple general explanation of larger scale (postsynaptic) dynamics of the cerebral cortex at behaviorally relevant time-scales.

The state space dimensionalities in our data at all recorded sites were larger than 3 but less than 10. Previous studies on cortical dynamics reported dimensionalities up to 3 (Bruno et al., 2017; Hennequin et al., 2018; Knierim and Zhang, 2012; Oprisan et al., 2015; Priesemann et al., 2013; Rabinovich et al., 2012; Tognioli and Scott Kelso, 2011; Wimmer et al., 2014). One reason for this discrepancy might be that FECAT-dynamic was derived from the actual cortical field potential signal, instead of being a theoretical attractor model borrowed from physics or other disciplines.

Methodological considerations

Field potentials recorded from the cerebral cortex will always be multi-dimensional in practice and hence best comprehensible in a multi-dimensional state-space (McCormick et al., 2020). To get a comprehensive account of the function of the neural structures producing the field potential, the dynamics should be studied in a state space of sufficient dimensionality. Bandwidth and power spectra analyses cannot reveal the underlying dynamics in field potentials (Babloyantz and Destexhe, 1986; Breakspear, 2017; Knierim and Zhang, 2012; Oprisan et al., 2015; Rabinovich et al., 2012; Roschke and Basar, 1990; Stam, 1996,). In contrast, by extracting non-linear field potential dynamics in state space of sufficiently high dimensionality (Takens, 1981) and by avoiding methods requiring stationarity, we discovered the fluctuating attractors. A low-dimensional, slowly decaying, oscillating spiral attractor was recently described (Bruno et al., 2017). However, we were unable to find attractors similar to FECAT in the literature. Attractors of similar dimensionalities to FECAT derived from the human EEG have been proposed (Babloyantz and Destexhe,1986; Roschke and Basar, 1990; Stam, 1996) based on a working assumption of stationarity. It is possible that these putative EEG attractors are in fact FECAT attractors, which could be tested, for example by the strategy outlined in **Fig. 1** and **STAR Methods.**

We studied the volume expansions and contractions of local cortical states in state space over successive 50 ms intervals. Since we used only 25 states for the calculation of the volume, it made no sense to narrow the observation intervals. This does not preclude studies of faster fluctuations of the expansions and contractions, if the sampling rate of the data is sufficiently high and if the autocorrelation in the time series is rapidly declining. The volume of the states varied significantly with the dimension of state space. Most likely, the dimensionality of the dynamics changes faster than over 60 s, the time needed to get a reasonably accurate estimate. However, the volume of the cortical states may reflect smaller changes in dimensionality. If the distances between states increase, the product of the square

roots of variances and hence the volume of the states increases (equation 4, **STAR Methods**). This is roughly equivalent with an increase in dimensionality. Similarly, if the distances between states decrease, the variances and hence the volume of the states decrease. With our data we could only observe volume fluctuations slower than the average flow of states. However, the flow of the states did not seem to fluctuate faster than the volume (see **Figure S6** values).

The electrode lead sites were not uniformly distributed over the cortex in the whole material, and were sparsely scattered in individual patients. This precluded a coherent anatomical reconstruction of the lead positions for examining spatio-temporal cortical dynamics *(Arieli et al., 1995; Ayzenshtat et al., 2010; Ferezou et al., 2007; Muller et al., 2018; Roland et al., 2017)*.

Evolution of postsynaptic cortical states and local excitation and inhibition

Local networks in prefrontal, parietal and temporal cortices, hippocampus and amygdala all operated with FECAT-dynamic. These networks have well-described different anatomical and receptor architecture (Zilles and Amunts, 2010). The common FECAT-dynamic thus should have a common cause. One explanation of this consistent result is that delays and timing of excitation and inhibition are preserved in the field potentials from these different networks and expressed in the FECAT-dynamic.

The field potential is a weighted sum of the extracellular currents in the neighborhood of the electrode lead. The main sources of these extracellular currents are the underlying multiple postsynaptic membrane currents (Einevoll et al., 2014; Pesaran et al., 2018). The field potential is measured at a relatively coarse mm^3 scale, causing many details of the biophysical computations of inward and outward currents to get lost (Iacaruso et al., 2017). Only closely timed and dominant excitations and inhibitions in the network may remain in the multidimensional field potential. So, it might well be that architectural differences in the local networks are responsible for important biophysical computations that are not captured at the mm^3 scale. At this scale, the expansions, contractions, progress of states and attraction could be caused by multidimensional action of excitation and inhibition in the local network. Without excitation, states will not progress and expand in state space. Without inhibition, evolving states will not contract, and states escaping from the attractor will not be pulled back (Roland et al., 2017).

The field potential is closely correlated and in phase with the voltage sensitive dye signal in the cerebral cortex of rodents, carnivores, and awake primates (Arieli et al., 1995; Ayzenshtat et al., 2010; Ferezou et al., 2007; Lippert, et al., 2007; Petersen et al., 2003; Xu et al., 2007). The visual cortical areas in the ferret have similar short stretching of trajectories in state space when excitation dominates the local network, followed by a slower 1 s contraction when inhibition dominates (Roland et al., 2017). Thus, the stimulus-induced expansions and contractions of cortical states we find in human local networks may relate to local dominance and timing of excitation, followed by local dominance and timing of inhibition. The ongoing expansions and contractions in state space observed in the absence of experimental stimuli may also reflect such shifts.

An advantage of such an organization of common network dynamics is that it facilitates the communication between different local networks. If networks operated with different dynamics it would take time to mutually adjust the dynamics. Indeed, in the single trial tests, the reshaped expansion and contraction of cortical states cross-correlated between a subset of cortical networks. What is communicated between the networks must be reflected in the timing and dominance of excitation and inhibition in the target networks. These excitation-inhibition shifts exist at multiple (local spatial) scales. Finer details in perception, imagery and motor control which cannot be resolved at this mm^3 scale could be resolved by mechanisms also affecting the delayed timing of excitation and inhibition at microscopic scales (Dehghani et al., 2016; Harvey and Roland, 2013; McCormick et al., 2020). The consistent FECAT-dynamic is the overriding principle at the mm^3 scale. However, whether FECAT-dynamic is preserved at finer scales and how the state expansions and contractions relate to local excitation and inhibition are questions that can be tested more readily in animals.

If the multi-dimensional timing and dominance between excitation and inhibition is expressed in the FECAT-dynamic, this would also explain how all the different brain functions we examined could emerge from FECAT-dynamic. Having only one (type of) fluctuating attractor dynamic operating under all conditions means that the cortex does not need to change dynamics to react to new transients and to shift to another brain function.

Implications of FECAT-dynamic

Changes in field potentials that are not locked to stimulus presentation or motor behavior are often regarded as background activity, obscure ongoing activity, default mode, resting, "dark activity" or noise (Buzsaki, 2019; Singer et al., 2019; but see McCormick et al., 2020). However, cortical states in all the different behavioral conditions, spontaneous and resting states included, evolved under FECAT-dynamic. Also, the expansions and contractions of cortical states during experimental conditions were within the same ranges as during spontaneous activities and rest (**Figures 3C, 3D and 3G**). These results show that there is no reason to maintain that spontaneous attractor state fluctuations during rest and under spontaneous activity in the awake conditions are noise or driven by other types of dynamics than the dynamic driving perception, retrieval of memories, thinking, and preparation of motor activity. The FECAT-dynamic accounting for all these conditions ensures that cortical networks easily switch from internally initiated functions to externally initiated functions and vice versa. Thus, spontaneous ongoing and resting activity, sensory evoked and preparatory motor activity is unified by the FECAT-dynamic creating local cortical states.

It is therefore unlikely that FECAT-dynamic is caused by stimuli, test demands and other external causes. The progress, the expansions and contractions, and attractor dynamics continued without external signals or demands. This result, and the finding that all conditions evolved under FECAT-dynamic, indicate that FECAT is an ongoing continuous intrinsic dynamic of cortical local networks that can be modified, but not changed or replaced by another dynamic in response to external demands or stimuli. Having ever ongoing progress, expansions and contractions of cortical attractor states that is modifiable in reaction to intrinsic as well as extrinsic sources has important

biological advantages. It eliminates needs for starting mechanisms initiating evoked and intrinsic activities.

Cortical networks must remain permanently stable, preventing runaway excitation (Buzsaki, 2019; Rabinoivich et al., 2012; Singer et al., 2019). On the other hand, most known stable attractors prevent the network to react to small transients (Rabinovich et al., 2012; Tognioli and Scott Kelso, 2011; Wimmer et al., 2014). The attractor property of FECAT-dynamic explains how cortical states under all (physiological) conditions can evolve with stability (i.e. do not split (bifurcate), change into another dynamic or run off). Yet, the modulations of the fluctuations demonstrate the sensitivity of this dynamic to small single trial inputs, as expected from a complex dynamical system. This unifies cortical stability with sensitivity to small transient inputs, and resolves a long-lasting conundrum in theoretical and computational neuroscience (Breakspear, 2017; Hennequin et al., 2018; Rabinoivich et al., 2012; Tognioli and Scott Kelso, 2011).

One principal result is that irregular fluctuations of the attractor volumes are the rule. However, FECAT-dynamic demonstrates that both fluctuating states and entrained oscillatory cortical states can arise from the same dynamic. This resolves a longstanding controversy between scientists who regard asynchronous irregular cortical states as characteristic for the cerebral cortex, and others who emphasize oscillations as fundamental for cortical operations (Breakspear, 2017; Buzsaki, 2019; Dehghani et al., 2016; Einevoll et al., 2014; McCormick et al., 2020; Mulller et al., 2018; Pesaran et al., 2018; Singer et al., 2019).

Like almost all other dynamics, FECAT-dynamic accounts for the *temporal* evolution of states in local cortical networks. Local cortical networks are often incapable of distinguishing clear differences in stimuli, especially in single trials (Forsberg et al., 2016; Ganguli et al., 2008). Different conditions might differ by the detailed nature of the attractor fluctuations and by their *spatio*-temporal cross-correlations in cortical space. Cross-correlations of attractor volume fluctuations captured the cooperation between local cortical networks across the cortex in the examples provided. This makes single trial cortical operations robust. Notably, the expansions and contractions were cross-correlated, not the field potentials themselves as in previous reports (Muller et al., 2018). The results also indicated that cross-correlated expansions and contractions of cortical states among a certain combination of local networks characterized a particular single trial test. This suggests that engagements in different tasks can may be distinguished by *spatio*-temporal differences in the participations of local networks (Arieli et al. 1995; Ayzenshtat et al., 2010; Ferezou et al. 2007; Iacaruso et al., 2017; Roland, 2017; Singer et al., 2019).

The trajectories inside the attractor show how cortical states evolve in state space and time. Since the trajectories of the multidimensional field potential in state space continuously stretched and contracted, our results are incompatible with dynamics temporarily settling into particular stable states (equilibria) or representations. This applied even to conditions in which patients were exposed to stationary pictures or imagined stationary objects. These conditions were associated with a short expansion, stretching the trajectory, followed by a 1 to 2 s contraction of the trajectory (**Figures 4, 5 and 6**). When the trajectory stretches, the state-space-time sequence (making the trajectory) gets stretched (evolve faster). When the trajectory contracts, the state-

space-time shortens (evolve more slowly). This is under the assumption that the local networks have no absolute clock-time. The contractions, as argued above, may relate to an overall inhibition in the local networks coinciding with visual perception (Lee, 2003; Harvey and Roland, 2013; Ozeki et al., 2009; Roland et al., 2017). Thus, cross-correlated contractions in state-space-time between networks may relate to the subjective experience of scenes or objects, since the experience concurred with the contractions.

Cortical states derived from field potentials are traditionally associated with more global conditions such as sleep and awake states, attention and alertness, in time scales ranging from seconds to hours (Buzsaki, 2019; Rheinberger and Jasper, 1937; Singer et al., 2019; Stam, 1996). However, the postsynaptic currents in a local cortical network most likely also reflect the collective computations contributing to brain functions at the mm^3 spatial scale and the ms time scale (Arieli et al., 1995; Ayzenshtat et al., 2010; Ferezou et al., 2007; Harvey and Roland, 2013; Iacaruso et al., 2017; McCormick et al., 2020; Petersen et al., 2003; Roland et al., 2017). The cortical states extracted from the field potentials should therefore also reflect contributions to specific brain functions being performed. This makes FECAT-dynamic a rule explaining larger scale cortical operations within and between local cortical networks.

The fluctuating expanding and contracting attractor is a mathematical object. This makes it a theoretically overall and simple description of how cortical states evolve that is easy to test experimentally, also at finer scales of observation and in other species. However, FECAT dynamic does not deal with single-cell spiking dynamics.

Perspectives

Rather than having millions of individual postsynaptic currents driving each local network, consistent fluctuating expanding and contracting attractor dynamic was used as an overall collective mechanism to drive evolving cortical states at mm^3 spatial scales. Although the cortical states were vigorously changing, the attractor property makes the state evolution well controlled and robust. Our data left no evidence of any stationarity, stable patterns, stable states or equilibria. In contrast, the ever-continuing expansions and contractions and flow of cortical states were the main features of cortical functional dynamics under normal physiological conditions. Cortical functions are likely to emerge from modifications of the flow, expansions and contractions spreading among the local networks. However, the outcomes - i.e. percepts, emotions, mental images etc.- are also based on continuously changing cortical states.

Conceptually, fluctuating expanding and contracting attractor dynamic resolve and replace current dichotomies in systems neuroscience between stable brain dynamics versus sensitive transient brain dynamics, spontaneous versus evoked cortical activity, asynchronous irregular fluctuations versus oscillatory activity. FECAT-dynamic also unifies the dynamics created by different local cortical networks and the dynamics of the collective cooperation of local networks.

FECAT-dynamic is neither a "top-down" or "bottom-up" description of cortical operations, but its mathematical structure makes it a theoretical model of cortical dynamics that can be tested experimentally also at finer scales of observation in animals. This dynamic may also be hidden in MEG and EEG signals in humans and non-human primates. We expect fluctuating expanding and contracting attractor

dynamic also in other mammalian brains. Future studies should examine the extent to which the modifications of the expansions and contractions and the flow of states predict behavioral conditions in detail.

Acknowledgments: We gratefully acknowledge the expert technical help of Lennart Derm and Thomas Kirkeskov Olsson, Dept. Clinical Neurophysiology University Hospital Copenhagen and the editorial help of Paul Cumming. We are greatly indebted to John Hertz for his thoughtful discussions and important input on earlier versions of the manuscript, to Aasa Feragen for advice on multidimensional volumes, to Laurence Dricot for help with the anatomical localization of the electrode leads, and to Viktor Jirsa for his critical reading and comments.

Funding: This study was supported by the Velux Foundation, Denmark.

Author contributions: PER conceived and designed the study. PER and AS got the funding. BJ did surgery implantation and postoperative survey. MF was responsible for field potential acquisition and wrote the ethics application with AS and PER. AW created all data analysis pipelines and wrote all software together with SNL SH, CKKH. JM and AW developed hardware. Data acquisition: JM, AW, CKKH. Experimental testing of patients: JM, RK, PER. PER, AW, SNL developed the attractor volume model. AW, PER, JM, CKKH did the data analysis. SNL did statistics. RK, JTD did the anatomical analysis. RK made all Figures. ML, LP, AS managed and controlled the clinical conditions of the patients. PER wrote the earlier versions. PER, JM and RK edited the final version of the manuscript. **Competing interests:** Authors declare no competing interests. **Data and materials availability:** Data are available at **github** https://github.com/ChristofferKKHansen/RolandLab.

Supplementary Materials:

Figures S1-S7

Tables S1-S2

References

Andrillon, T., Nir, Y, Staba, R., Ferrarelli, F., Cirelli, C. and Tononi, G. (2011). Sleep spindles in humans: insight from intracranial EEG and unit recordings. J. Neurosci. 31, 17821-17834.

Arieli, A., Shoham, D., Hildesheim, D.R., and Grinvald, A. (1995). Coherent spatiotemporal patterns of ongoing activity revealed by real-time optical imaging coupled with single-unit recording in the cat visual cortex. J. Neurophysiol. 73, 2072-2093.

Avants, B.B., Epstein, C.L., Grossman, M. and Gee, J.C. (2008). Symmetric diffeomorphic image registration with cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. Medical image analysis, 12, 26-41.

Ayzenshtat, I., Meirovithz, E., Edelman, H., Werner-Reiss, U., Bienenstock, U., Abeles, M., and Slovin, H. (2010). Precise spatiotemporal patterns among visual cortical areas and their relation to visual stimulus processing. J. Neurosci. 30, 11232-11245.

Babloyantz, A., and Destexhe, A. (1986). Low-dimensional chaos in an instance of epilepsy. Proc. Natl. Acad. Sci. U. S. A. 83, 35613-3517.

Breakspear, M. (2017). Dynamic models of large-scale brain activity. Nat. Neurosci. 20, 340-352.

Bruno, A.M., Frost, W., and Humphries M.D. (2017). A spiral attractor network drives rhythmic locomotion. eLIfe 2017; 6e27342, DOI:10.7554/eLife 27342.

Buzsaki, G. (2019). The brain inside out, MIT Press, Cambridge, Mass.

Camastra, F. (2003). Data dimensionality estimation methods: A survey. Pattern Recogn. 36, 2945-2954.

Churchland, M.M., Yu, B.M., Cunningham, J.P., Sugrue, L.P., Cohen, M.R., Corrado, G.S., Newsome, W.T., Clark, A.M., Hosseini, P., Scott, B.B., et al. (2010). Stimulus onset quenches neural variability: a widespread cortical phenomenon. Nat. Neurosci. 13, 369–378.

Dehghani, N., Peyrache, A., Telenczuk, B., Le Van Quyen, M., Halgren, E., Cash, S.S., Hatsopoulos, N.G., and Destexhe, A. (2016). Dynamic balance of excitation and inhibition in human and monkey neocortex. Sci. Rep. 6, 23176. doi: 10.1038/srep23176.

Einevoll, G.T., Kayser, C., Logothetis, N.K., and Panzieri, S. (2014). Modelling and analysis of local field potentials for studying the function of cortical circuits. Nat. Neurosci. 14, 770-785.

Elston, G.N. (2003). Cortex, cognition and the cell: New insights into the pyramidal neuron and prefrontal function. Cereb. Cortex 13, 1124-118.

Ferezou, I., Haiss, F., Gentet, L.J., Aronoff, R., Weber, B., and Petersen, C.C.H. (2007). Spatiotemporal dynamics of cortical sensorimotor integration in behaving mice. Neuron 56, 907-923.

Fonov, V.S., Evans, A.C., McKinstry, R.C., Almli, C. and Collins, D. (2009). Unbiased nonlinear average age-appropriate brain templates from birth to adulthood. NeuroImage 47, S102.

Forsberg, L.E., Bonde, L.H., Harvey, M.A., and Roland, P.E. (2016). The second spiking threshold: Dynamics of laminar network spiking in the visual cortex. Front. Syst. Neurosci. 10, 65. doi: 10.3389/fnsys.2016.00065.

Gallego, J.A., Perich, M.G., Miller, L.E., and Solla, S.A. (2017). Neural manifolds for the control of movement. Neuron 94, 978-984.

Ganguli, S., Huh, D., and Sompolinsky, H. (2008). Memory traces in dynamical systems. Proc. Natl. Acad. Sci. 105, 18970-18975.

Gillies M.J., Huang, Y., Hyam, J.A., Aziz, T.Z., Green, A.L. (2019) Direct evidence for a role of the human anterior cingulate cortex in central command. Auton. Neurosci. 216, 61-68.

Grassberger, P. and Procaccia, I. (1983). Characterization of strange attractors. Phys. Rev. Lett. 50, 346-349.

Harvey, M.A., and Roland, P.E. (2013). Laminar firing and membrane dynamics in four visual areas exposed to two objects moving to occlusion. Front. Syst. Neurosci. 7: 23. doi: 103389/fnsys.2013.00023.

Hastie, T., Tibshirani, R. and Friedman, J. (2009). The elements of statistical learning, Springer, New York.

Hennequin, G., Ahmadian, Y., Rubin, D.B., Lengyel, M., and Miller, K.D. (2018). The dynamical regime of sensory cortex: Stable dynamics around a single Stimulus-tuned attractor account for patterns of noise variability. Neuron 16, 846-860.

Hori, T., Sugita, I.Y., Koga, E., Shirakawa, S., Inoue, K., Uchida, S., Kuwahara, H., Kousaka, M., Kobayashi, T., Tsuji, Y., Terashima, M., Fukuda, K. and Fukuda, N. (2001). Sleep Computing Committee of the Japanese Society of Sleep Research Society Proposed supplements and amendments to "A manual of standardized terminology, techniques, and scoring system for sleep stages of human subjects", the Rechtschaffen and Kales (1968) standard. Psychiatry Clin. Neurosci. 55, 305-310.

Iacaruso, F. M., Gasler, I.T., and Hofer S.B. (2017). Synaptic organization of visual space in primary visual cortex. Nature 547,449-452.

Iglesias, J.E., Liu, C.-Y., Thompson, P.M. and Tu, Z. (2011). Robust brain extraction across datasets and comparison with publicly available methods. IEEE transactions on medical imaging, 30, 1617-1634.

ITK. Insight segmentation and registration toolkit. Web Page: https://itk.org/, 2019. [Consulted in June 2019].

Kennel, M.B., Brown, R., and Abarbane, I.H.D.I. (1992). Determining embedding dimension for phase-space reconstruction using a geometrical construction. Phys. Rev. A 45, 3403-3411.

Knierim, J.J., and Zhang, K. (2012). Attractor dynamics of spatially correlated neural activity in the limbic system. Ann. Rev. Neurosci. 35, 267-285.

Lakatos, P., Karmos, G., Mehta, A.D., Ulbert, I., and Schroeder, C.E. (2008). Entrainment of neuronal oscillations as a mechanism of attentional selection. Science 320,110-113.

Lee, T.S. (2003) Computations in early visual cortex. J. Physiol. Paris, 97, 121-139.

Lindén, H., Pettersen, K.H. and Einevoll, G.T. (2010). Intrinsic dendritic filtering gives low-pass power spectra of local field potentials. J. Comput. Neurosci. 29, 423-444.

Lippert, M.T., Takagaki, K., Xu, W., Huang, X., and Wu, J.-Y. (2007). Methods for voltage-sensitive dye imaging of rat cortical activity with high signal-to-noise ratio. J. Neurophysiol. 98, 502-512.

Lundqvist, D., Flykt, A. and Öhman, A. (1998). The Karolinska Directed Emotional Faces (KDEF). Stockholm: Department of Neurosciences Karolinska Hospital.

McCormick, D.A., Nestvogel, D.B., He, B.J. (2020) Neuromodulation of brain state and behavior. Annu. Rev. Neurosci. 43, 391-415.

Muller, L., Chavane, F., Reynolds, J., and Sejnowsky T. (2018). Cortical travelling waves: mechanisms and computational principles. Nat. Rev. Neurosci. 19, 255-268.

Oprisan, S.A., Lynn, P.E., Tompa, T., and Lavin, A. (2015). Low-dimensional attractor for neural activity from local field potential in optogenetic mice. Front. Comput. Neurosci. 9, 125 (2015). doi: 10389/fncom.201500125.

Ozeki, H., Finn, I.M., Schaffer, E.S., Miller, K.D., and Ferster, D. (2009). Inhibitory stabilization of the cortical network underlies visual surround suppression. Neuron 62, 578-592.

Pesaran, B., Vinck, M., Einevoll, G.T., Sirota, A., Fries, P., Siegel, M., Truccolo, W., Schroeder, C.E., and Srinivasan, R. (2018). Investigating large-scale brain dynamics using field potential recordings: analysis and interpretation. Nat. Neurosci. 21, 903-919.

Pettersen, K.H. and Einevoll, G.T. (2008). Amplitude variability and extracellular low-pass filtering of neuronal spikes. Biophys. J. 94, 784-802.

Petersen, C.C.H., Hahn, T.T.G., Mehta, M., Grinvald, A., and Sakmann, B. (2003). Interaction of sensory responses with spontaneous depolarization in layer 2/3 barrel cortex. Proc. Natl. Acad. Sci. U. S. A. 100, 13638-13643.

Priesemann, V., Valderrama, M., Wibral, M., and Le Van Quyen, M. (2013). Neuronal avalanches differ from wakefulness to deep sleep – Evidence from intracranial depth recordings in humans. PLoS Comput. Biol. 9, e1002985. doi: 10.1371/journal.pcbi.1002985.

Pritchard, W.S. and Duke, D.W. (1995). Measuring chaos in the brain: A tutorial review of EEG dimensions estimation. Brain Cogn. 27, 353-397.

Rabinovich, M.I., Friston, K.J., and Varona, P. Editors (2012). Principles of Brain Dynamics (MIT Press, Cambridge, Mass., 2012).

Rheinberger, M. and Jasper, H. (1937). The electrical activity of the cerebral cortex in the unanesthetized cat. Am. J. Physiol. 119, 186-196.

Roland, P.E. (1981). Somatotopical tuning on the postcentral gyrus during focal attention in man. A regional cerebral blood flow study. J. Neurophysiol., 46:744-754.

Roland, P.E. (2017). Space-time dynamics of membrane currents evolve to shape excitation, spiking, and inhibition in the cortex at small and large scales. Neuron 94, 934-942.

Roland, P.E., Bonde, L.H., Forsberg, L., and Harvey, M. (2017). Breaking the excitation-inhibition balance makes the cortical network's space-time dynamics distinguish simple visual scenes. Front. Syst. Neurosci. 11:14. doi: 10.3389/fnsys.2017.00014.

Roland, P.E. and Friberg, L. (1985). Localization of cortical areas activated by thinking. J. Neurophysiol., 53:1219-1243.

Roschke, J., and Basar, E. (1990). "The EEG is not simple noise: strange attractors in intracranial structures" in Chaos in brain function, E. Basar, Ed. (Springer, Berlin), pp. 49-62.

Singer, W., Sejnowski, T.J., and Rakic, P. Editors (2019) The Neocortex, MIT Press, Cambridge, Mass.

Stam, C.J. (1996). Non-linear dynamic analysis of EEG and MEG: Review of an emerging field. Clin. Neurophysiol. 116, 2266-2301.

Staresina, B.P., Reber, T.P., Niediek, J., Boström, J., Elger, C.E. (2019) Recollection in the human hippocampal-entorhinal cell circuitry. Nat Commun. 10(1):1503. doi:10.1038/s41467-019-09558-3.

Takens, F. (1981). "Detecting strange attractors in turbulence". In Dynamical Systems and Turbulence, Lecture Notes in Mathematics, D.A. Rand, L.-S. Young, Eds. (Springer-Verlag). vol. 898, pp. 366–38.

Thévenaz, P. and Unser, M (2000). Optimization of mutual information for multiresolution image registration. IEEE transactions on image processing, 9, 2083-2099.

Thorpe, S., Fize, D., and Marlot, C. (1996). Speed of processing in the human visual system. Nature 381,520-522.

Tognioli, E., and Scott Kelso, J.A. (2011). The metastable brain. Neuron 81, 35-48.

Wimmer, K., Nykamp, D.Q., Constantinidis, C., and Compte, A. (2014). Bump attractor dynamics in prefrontal cortex explains behavioral precision in spatial working memory. Nat. Neurosci. 17, 431-439.

Xu, W., Huang, X., Takagaki, K., and Wu, J.Y. (2007). Compression and reflection of visually evoked cortical waves. Neuron 55, 119-129.

Zilles, K., and Amunts, K. (2010). Centenary of Brodmann's map- conception and fate. Nat. Rev. Neurosci. 11, 139-145.

STAR Methods

A total of 13 patients (6 females, 7 males; age range: 19 to 55 years; mean: 38 years, SD: 12 years) participated in this study. The patients had drug resistant epilepsy, were candidates for surgical resection and were implanted with depth electrodes, at locations in cerebral cortex determined to reveal the location of an epileptic focus and spread of epileptic discharges. The study was approved by the Human Ethics Committee for the Greater Copenhagen region (permission 56441). The patients spent from 5 to 13 days with the electrodes implanted.

The aim of this study was to obtain field potentials, without any abnormalities, from as many cortical sites as possible. The experimental testing conditions were designed to be as diverse as possible in the clinical ward. The purpose of this diversity was to examine whether the dynamics driving the evolution of the field potential were different for different tests, or explored different parts of state space. The experimental conditions were not designed to map or functionally dissect sensory, motor and cognitive brain functions. The analysis design is summarized in **Figure 1.**

After removal of all epileptic activity (see below), the material comprised 290 electrode leads. The anatomical locations of the electrode leads are listed in **Table 1**.

Experimental Conditions

Typically, 1 to 3 days after the implantation, the patient participated in different tests aimed to induce behavioral brain states as diverse as possible. In addition, we selected episodes of sleep, conversation with the nurse, TV watching, and eating. The purpose of the many behavioral conditions was to let the field potential dynamics explore state space over a wide range. For different practical reasons (medication, post-ictal sleep, early dismissal) not all patients performed all tests. Subjects were sitting or lying in bed. All behavioral conditions were collected several hours away from ictal and post-ictal activity, typically in the first two days in the ward.

Visual tests

We collected a series of 200 color pictures of landscapes and outdoor environments, devoid of images of humans (96 dpi, 1060 x 1920 pixels) downloaded from freeware sources on the internet. The pictures were presented on a screen measuring 53x30 cm (refresh rate: 144Hz) which was placed approximately 120 cm from the eyes. Patients were instructed just to watch; with the exception of visual test 6, no behavioral responses were required. A total of 11 visual tests was conducted.
Visual test 1: 200 pictures appeared, 1 frame each (6.94 ms) with no intermission.
Visual test 2: 100 pictures were randomly selected and each was presented for 49 ms with no intermission.
Visual test 3: the same 100 pictures were presented each for 125 ms with no intermission.
Visual test 4: the same 100 pictures were presented each for 125 ms, followed by a 999-ms lasting black screen
Visual test 5: the same 100 pictures were presented each for 999 ms, followed by a 999-ms lasting black screen.

Visual test 6: The next day, the 100 pictures were mixed with 100 new ones that the patient had not seen before. The 200 pictures were randomly presented each for 999 ms, followed by a 1999 ms lasting black screen. Following each picture, the patient had to indicate whether he/she had seen it earlier (recognition test).

Table 1: anatomical localisation of the electrode leads

Anatomical site	№ leads
Hippocampus	38
head (20)	
body & tail (18)	
Middle temporal gyrus	35
anterior part (8)	
middle part (19)	
posterior part (8)	
Cingulate gyrus	29
anterior part (19)	
middle part (5)	
posterrior part (5)	
Insula	27
anterior part (13)	
posterior part (14)	
Superior temporal sulcus	18
anterior part (3)	
middle part (7)	
posterior part (8)	
Superior temporal gyrus	12
anterior + middle part (10)	
posterior part (2)	
Superior frontal gyrus	12
Parietal operculum	11
Temporal pole	10
Inferior temporal gyrus	10
Supramarginal gyrus	9
Fusiform gyrus	9
Superior frontal sulcus	9
anterior part (1)	
posterior part (8)	
Parahippocampal gyrus	8
Orbitofrontal cortex	7
lateral part (4)	
medial part (3)	
Middle frontal gyrus	5
Frontal operculum	5
Lingual gyrus	4
Entorhinal cortex	3
Inferior frontal gyrus	2
Precuneus	2
Gyrus precentralis	1
Gyrus postcentralis	1
Calcarine cortex	1
Angular gyrus	1
Subiculum	1

Visual test 7: apparent motion. A 3°x 3° black square moved on a white background in steps of 2° from the left to the right border of the screen, in 583 ms. The test was repeated 100 times.

Visual test 8: same as visual test 7, but here the square moved from right to left.

Visual test 9: one stationary 3°x 3° white square was shown for 1 minute.

Visual test 10: one 3°x 3° red square, flanked by four 3°x 3° green squares, was shown for 2 minutes.

Visual test 11: one white square, 3 white squares in a row, one red square flanked by 4 green squares, or one green square flanked by 4 red squares were shown for 250 ms. Presentation was random and balanced such that each alternative was shown 50 times. All squares measured 3°x3°.

Control conditions

In the Ganzfeld condition, the patients' eyes were occluded for 10 minutes by white ping pong balls, cut in halves, and covered by surgical gauze pads secured by tape, preventing any patterned visual input. The patients were asked to keep eyes open during the test. In the rest condition, the patient had eyes closed, was encouraged to relax, concentrating on having it black in front of the mind's eye, but stay awake. Patients were supine in both control conditions.

Emotions

A total of 35 pictures (72 dpi) were taken from the Karolinska Directed Emotional Faces (KDEF) test (Lundquist et al.,1998), showing 7 different emotional expressions (angry, disgusted, sad, fearful, happy, surprised and neutral). Each image was shown for 6 s, followed by a 2s dark screen. Following each picture, patients were asked to indicate the type of emotional expression.

Memory retrieval tests

During these tests, the patient had eyes closed.

Memory test 1: recall faces. With 5 s interval, the experimenter read the names of 29 well-known persons or cartoon characters (e.g. queen Margrethe, Donald Duck, Angela Merkel). The patient responded by yes or no whether he/she could clearly visualize the face.

Memory test 2: recall places. With 5 s interval, the experimenter read the names of 15 well-known landmarks or tourist attractions (e.g. the little mermaid, golden gate bridge, the Chinese wall). The patient responded by yes or no whether he/she could clearly visualize the place.

Memory test 3: mental navigation. The patient was asked to imagine going out of the front door of his/her residence, going to the right down the street, then walking down the first street appearing to the left and then alternatively to the right and left as in (Roland and Friberg, 1985). After 65 s the patient was asked to tell where she/he is.

Memory test 4: same as 3, but starting taking the home street to the left (Roland and Friberg, 1985).

Memory test 5: a known target at some distance from the patient's house was chosen. The patient was asked to imagine walking/cycling slowly to the target and say "now" upon arrival.

Memory tests 6 and 7: same as memory test 5 but each with a new target.

Memory test 8: patient was asked to imagine standing at target (condition 5) and walking/ cycling back home. Upon arrival, patient says "now".

Memory test 9: The patient imagines standing at target in condition 6, otherwise as in condition 8.

Memory test 10: patient was asked to recall all events in temporal order during the hospital admission day, from getting out of bed in the morning until arriving at the hospital. Upon arrival, patient says "now".

Classification of nouns

A list of 19 Danish nouns all starting with "hoved" (head or main in English) was read (6 s inter-stimulus interval) and the patient was asked to classify each noun as being "concrete" or "abstract".

Motor tests

An EMG disc electrode was fixed to the skin over the forearm extensors, contralateral to the brain hemisphere carrying the largest number of implanted electrodes. A reference electrode was fixed near the wrist, and the EMG signal was sampled by an EMG amplifier (10 Hz-5kHz band-pass filter; amplified and finally recorded at 30 kHz) and used to define the motor epochs.

Motor test 1: the patient was asked to move the arm with the electrode such that the hand touches the left side of the chest, to move it back to the original rest position, and then to touch the right side of the chest. Then in similar order, the same hand moves to touch right and left shoulder, right and left foot, right and left knee, right and left abdomen, forehead, right and left heel, right and left thigh, opposite elbow, and opposite hand, in total lasting 60 s.

Motor test 2: the patient was asked to quickly bend the index-finger and thumb to touch, then quickly to extend the fingers again and to continue this as quickly as possible for 60s in total.

Motor test 3: as in motor test 2, but synchronously with the index and thumbs of both hands.

Motor test 4: as in motor test 3, but counterstroke.

Motor test 5: rhythm. With the hand contralateral to the hemisphere with the majority of implanted electrodes, the index touches the thumb for approximately one s, followed by 3 short touches and so on in the same rhythm for total of 60 s.

Motor test 6: With the same hand as condition 5, the thumb touches shortly the index, middle finger, ring finger and little finger, the little finger, ring finger, middle finger and index and so back and forth for total of 60 s.

Attention tests

Somatosensory attention: the patient expected a touch stimulus close to detection threshold with a thin calibrated filament (pressure of 10 mg to 20 mg) on the tip of the index finger. The stimulus was first presented after 60 s had elapsed (Roland, 1981).

Auditory attention: The patient was asked to listen to a list of random numbers between 1 and 100, pronounced close to his/her auditory threshold. The patient had to repeat each detected number. In the ensuing 60-s period, no numbers were spoken. After 60 s have elapsed, a faint but detectable number was spoken, as a control.

Stereognosis

A small object (e.g. a cap of a USB-stick) was placed in the palm of the patient's hand. The patient had eyes closed and was prompted to manipulate the object with the fingertips. The testing lasted up to 30 s, or until the object was recognized. The test was then repeated with the other hand with another object.

Smell and taste

Ortho-nasal smell was tested according to conventional clinical bedside tests using mild odorants such as tea and chocolate. The patient was instructed to sniff deeply, with eyes closed, while the odorant was held close to patient's nose. Five odorants were used consecutively. Each odorant was tested for 20 s or until the odor was identified, whichever came first.

Retro-nasal smell and taste were tested by the patient pinching the nose with one hand, while he/she placed a sweet in the mouth with the other hand. Mouth-breathing was allowed. The patient was instructed to swallow only saliva, while chewing gently on the sweet. The patient was asked to report the taste of the sweet. After 20 s, the patient was instructed to exhale through the nose, and report any changes in the "taste" (flavor) of the sweet.

Sleep

1. Light sleep. Visual confirmation of sleep spindles, 9-16 Hz, minimum duration 0.5 s, maximum 2 s, appearing more frequently than 0.3 Hz in the field potential time series. In addition, visual confirmation of K-complexes > 200 μV, duration >200 ms, diphasic configuration. Finally, the selected 60 s episode should have maximum 20% delta activity. Light sleep was defined as a period of 60 seconds with at least 2 sleep spindles and 1 K-complex. Sleep spindles and K-complexes were detected by an algorithm and afterwards visually confirmed (Andrillon et al., 2011). The criteria for sleep spindles was an increase in 9-16 Hz activity of at least 3 standard deviations, with a minimum duration 0.5 s, and a maximum 2 s.
2. Slow wave sleep. The episode selected had more than 60% slow waves,1-2 Hz > 150 μV peak-to-peak. The amplitudes of the slow waves were more than 3 standard deviations of 0.1-2 Hz activity and had peak-to-valley of at least 150μV occupying more than 20% of the 60 second time period (Hori et al., 2001). All episodes selected were visually confirmed as slow wave sleep.

Field potentials

Field potentials were sampled at 512 Hz with a band-pass filter of 0.1 to 256 Hz from the implanted electrodes (SEEG electrodes; Ad-tech Medical Instruments, 1.28 mm in diameter, lead length 1.57 mm) in 13 patients. The electrode locations, based exclusively on clinical criteria, were determined from the post-operative CT-scan (1mm^3 voxels) co-registered to the patient's T1-weighted magnetic resonance Image (MRI) of the brain. After rejecting electrode leads located in white matter, artifacts and leads with epileptic activity (see below), the field potential from 290 leads located in the cerebral cortex were further analyzed.

The sources of the field potential are the transmembrane currents $I_n(t)$.

From (Lindén et al., 2010):

$$f(\boldsymbol{r}, t) = \frac{1}{4\pi\sigma} \sum_{n=1}^{N} \frac{I_n(t)}{|\boldsymbol{r} - \boldsymbol{r}_n|} \qquad (1)$$

$f(t)$ is the field potential at a particular instant for a population of neuron membranes close to the electrode lead (Pettersen and Einevoll, 2008). $I_n(t)$ is the net transmembrane current at a particular distance $\boldsymbol{r}_n$ from the lead. Under the assumption that the extracellular conductivity, σ, is constant over the whole range of the spatial summation in equation 1, $f(t)$ at each electrode lead, is the weighted (by their inverse distance from the lead) sum of the membrane and extracellular currents in the sampling space.

The field potential may show net increase of inward currents similar to a net decrease at another electrode, because polarity depends on the reference and might be reversed in recordings. Similarly, inward currents and outward currents in the space sampled by the electrode might cancel, despite lively synaptic activity (Einevoll et al., 2014; Pesaran et al., 2018). Artifacts (synchronous in all leads) and episodes with noise were removed. Thereafter, we used a common reference for each of the gray matter leads. The common reference was the median of the signal in all intra-cranial leads. This efficiently reduces the influence of remote sources on the local field potential. Each field potential then is a time series of fluctuations around an overall median (**Figure1**).

Detection and removing inter-ictal epileptic activity from data

To identify epileptic transients in the raw field potential, we performed a wavelet transform on the raw field potential and applied a threshold to the transform to identify epileptic transients. Transient epileptic events were identified using a Mexican hat wavelet (from Matlab® version 2018b). We used three wavelets with half widths of 0.02, 0.04 and 0.1 seconds). These wavelets were then convolved with the field potential signal to generate a time series where the amplitude correlates with transients in data with temporal characteristics of inter-ictal activity. We squared the convolved signal and defined a threshold of $0.5 \cdot 10^6$ μV as the limit of normal activity. Thereafter all files with detected epileptic activity were removed from the original set of 12949 $f(t)$ files at 476 cortical locations, leaving 3476 field potential files from 290 cortical locations each of length 60 s.

Reconstruction of the dynamics (embedding)

We assume that the dynamics of a local network in the cerebral cortex can be captured from a field potential recording. The dynamics are the set of rules by which the field potential evolves. For a given behavioral condition, the sampled field potential is a time series one can examine in state space to reveal its dynamics. We reconstructed the behavior in state space from the field potentials from the 290 cortical electrode leads for each of the behavioral conditions. In order to find the true dimensionality of the dynamics, we employed the delay embedding method (Takens, 1981). The state of the system at time t was represented by a d-dimensional vector:

$$\mathbf{F}(t) = [f(t), f(t+\tau), f(t+2\tau), \dots f(t+(d-1)\tau)]. \quad (2)$$

Here, *f(t)* is the field potential. The delay parameter τ is taken equal to the empirical autocorrelation time of *f*(*t*) (Pritchard and Duke, 1995). The embedding dimensionality was 20, i.e. d=20, judged enough to capture the underlying dynamics. This method is standard and has a long history in nonlinear system dynamics. The delay-embedded vector signal captures all generic information about the dynamics (Takens, 1981; Pritchard and Duke, 1995). We take this as a working hypothesis here as well.

From each *f(t)*, 20 delayed embedded versions were generated, with delays, τ = 59.9 ms. The delay parameter τ was chosen as the mean of 12949 original field potential time series as the value where the autocorrelation function of these time series = 1/e (Pritchard and Duke, 1995) (**Fig S1**). For each behavioral condition, the first 60 s of the *f*(*t*) was chosen as the basis of the embeddings.

Dimensionality of state space

The dimensionality of the dynamics was estimated with the Grassberger and Procaccia (1983) method by an automatic algorithm developed in-house. Specifically, for each point of the embedded time series, the distances to all other points in the state space were calculated. The number of points within a radius of ε was then counted for different values of ε. The correlation integral is given as:

$$C_\varepsilon = \frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^{n} \mathrm{H}(\varepsilon - \|x_j - x_i\|_2) \,, (3)$$

where n is the number of points, x_i and x_j are d-dimensional coordinate vectors, ε is the radius threshold and H is the Heaviside step function. The $\vert\,\vert$ indicates the norm. The calculation of C_ε is computationally intensive but parallelizing the code made it possible to reduce the computation time using the Danish Super Computer center Computerome (http://www.computerome.dtu.dk). The algorithm arrives at the state space dimension by considering the dependence between the correlation integral and the distance thresholds for a given embedding dimension. Specifically, the slope of each curve produced by a log(C_E) and log(ε) plot is automatically examined to determine the correlation dimension of the upper linear part of the curve **(Figure S1)**. The automated program for this identification has a manual counterpart since, in a small number of cases, it was difficult to determine the slope automatically. Finally, to determine the dimension, the dependence between the correlation dimension and embedding dimension is considered. If the correlation dimension is constant for 3 or more successive embedding dimensions, the algorithm concludes that the associated correlation dimension is the state space dimension, in short correlation dimensionality (**Figure S1)**. The constant part of this dependence is determined by a least squares approach (**Figure S1**). Both the automatic detection of the linear slope and the automated horizontal line determination of the dimension have multiple tuning parameters that can be more or less restrictive depending on the amount of noise in the data. For 28 records it was not possible to find a horizontal segment.

The correlation integral estimate depends on the number of sample points, here 30720 (60 s). Increasing the number of samples will increase the dimensionality slightly (**Figure S1**), but further decreases the likelihood that the dynamics, and hence the dimensionality, remains (quasi) stationary. So, the 60 s is a compromise.

The correlation integral algorithm also determined the amount of false nearest neighbors, and the inequality $R_{d+1}(n)/R_A > A_{tol}$ (see Kennel et al., 1992 for all details). If the embedding dimension chosen, in our case 20, is too low, the trajectories will intersect and overlap in state space. Such intersections and overlaps are called false nearest neighbors. False nearest neighbors thus tell that the embedding dimension chosen to reveal the dynamics is too low. R_A is an estimate of the size of the set containing all trajectories, $R_{d+1}(n)$ the sum of distances between each of the n points and neighbors, and the A_{tol} set to = 1 (Kennel et al., 1992). When the number of dimensions increase the distances between the points in state space increase; this is especially true for time series with a limited number of samples. The above inequality limits the possibility that false neighbors at high dimensionality become true neighbors. At embedding dimension 4 the number of false nearest neighbors was practically zero and the A_{tol} criterion fulfilled up to embedding dimension 20 (**Figure S1**). This means that the **F**(t) unfolds as expected from a dynamical system at embedding dimension 4 such that there are no crossings (intersections) of the trajectories in the attracting object (Kennel et al., 1992). For comparison, we shuffled the original time series to produced surrogate data with the random method (Matlab® randperm) (**Figure S1**).

The correlation dimensionality estimate seems superior to the alternatives (Camastra, 2003). The extraction of the dynamics from the embedded times series of the field potential was done by using integer dimensionality close to the estimated correlation dimensionality. With the absence of false nearest neighbors and the A_{tol} criterion fulfilled, these factors together should give a close to exhaustive description of the dynamics contained in the field potential.

Statistical analysis of dimensionality

When the distribution of estimated correlation dimensions over all behavioral conditions was plotted for each electrode lead, it appeared that in some cortical locations, the mean correlation dimension was high and the square-root of the correlation dimension variance relatively low; whereas in other locations mean correlation dimension was lower and the square-root of the variance relatively high (**Figures 2A, S2**). For each patient, we checked whether the distributions of the mean correlation dimension and square-root of the variance were approximately Gaussian. Under the assumption that the correlation dimensions distributed in two groups, one with high mean and low standard deviation, standard deviation, and one with lower means and higher standard deviation we subjected the correlation dimensions for each patient separately to a k-means cluster analysis (k=2). Two different centers were selected and a distance from the center to each point is measured. The algorithm then aims to have the lowest possible sum of distances. The algorithm is given as:

1. Initialize: select *k* points as medoids
2. Compute distances of each data point to each medoid
3. Assign each observation to the cluster with the closest medoid.
4. Compute average of the observations in each cluster to get a new medoid location
5. Repeat step 2-4 until cluster centers does not change

The medoid method implies that the algorithm finds one experimental co-ordinate, with minimum distances to all members of that cluster, as the center point for each cluster

(which reduces the influences of outliers). **Figure S2** shows the results. The quality of the clustering is apparent from the silhouette plots (Hastie et al., 2009). The silhouette value ranging from –1 to 1 shows how similar a cluster point is to its own cluster. A value a bit above 0 is usual in accepting the clustering. Silhouette values are shown in **Figure S3**.

To further statistically characterize the separation of the two groups we made a linear regression of the points of each cluster and compared the regressions lines (F-test) and verified that the residuals were approximately distributed as a Gaussian distribution. **Table S1** shows the results.

Dynamics confined to one part of state space

Figure 3A shows all 106,782,720 normal states in one state space. These states were distributed in 3476 sets. As the set of points occupying state space were multidimensional, their occupancy of state space was approximated. If the dimensionality was 6.6, for example, the set of points were located to a 6-dimensional state space, because the states do not exist in dimension 7. Each of the 3476 sets including all states was covered in a convex hull, the smallest convex set that contains all 106,782,720 states. The convex hull was computed for all relevant embedding dimensions. This was done using the Matlab® function convhulln using tetrahedra forms as facets. For the illustration, each of the 3476 sets was assigned one color; dimensions 1 to 3 and 4 to 6 were plotted separately in the figure (**Figure 3A**).

Fluctuating dynamics

With data sampled over 60s, the sets plotted in 3,4,5,6, or 7 dimensions will be unable to reveal fast dynamics relevant for perception, thinking, and preparation and execution of behavior. To examine faster changes in state space occupancy, we approximated location of the states as a hyper-ellipsoid (see **Figure 3A**). Mahalanobis distance is a measure of distances between data points in a multidimensional space, that is scale invariant and independent of Gaussian assumptions. Each state has a Mahalanobis distance in a multidimensional space from the center of the hyper-ellipsoid of the integer dimension just lower than the correlation dimensionality. We calculated the enclosing hyper-volume, of each of the 3476 sets.

We also used the *embedded time series* **F**(t) in the appropriate integer dimension to estimate the volume of the attractor (attracting set of states). If the correlation dimensionality of the set was 5.8, we calculated the first 5 principal components of the 5.8 dimensionality set. In this case, the estimated hyper-volume of the set in 5-dimensional state space was calculated as:

$$V = \sqrt{var1} \; \sqrt{var2} \; \sqrt{var3} \; \sqrt{var4} \; \sqrt{var5} \quad (4)$$

in which the product of the square-roots of the variances in each of the 5 orthogonal principal components dimensions is a hyper-rectangle giving the estimate V of the set in state space. The algorithm determined the V every 50 ms with a certain sliding window (was set to 128 samples = 250ms). The 50 ms are less than τ (60 ms). The sliding window was less than 5 τ, because most data ranged between 5 and 7 dimensions (see Flow of states below). This means that each value of V is an estimate of the volume of 25 consecutive states. Note that this method estimates V as the volume of an orthotope (hyper-rectangle) and not of a hyper-ellipsoid. In equation 4,

the orthotope is 5 dimensional. In addition, the succession of the 128 states was confined to a trajectory, that however could take various shapes (as also seen for the 1 s and 5 s trajectories in **Figure 3**). The states, however over some 10s to 20 s become well bounded by a hyper-ellipsoid as the convex hull. V fluctuated roughly from 10^7 to 10^{17} μV.

The amplitude variations of the fluctuations of the log $_{10}$ V from the median in all 60 s test episodes in all awake states (n= 3526) in all patients (n=13) were measured and shown in **Figure 3D.** For comparison, the similar fluctuations from 20 randomly selected 60 s episodes in awake states, i.e. spontaneous behavior not belonging to any test episode, were also calculated (**Figure 3D)**. A coefficient of variation of the z-score values of V (see below) was also calculated in steps of 50 ms with a sliding window of 250 ms (**Figure S5**).

Attractor properties

The set of states as points occupying state space were located to one part of state space. Neither the states themselves nor the trajectories they formed did spread to parts of state space outside the convex hull in **Figure 3A**. The reason for this could be that the network in which the states evolved had attracting properties, i.e. attracted states nearby. If we defined the hyper-ellipsoid as being within 90% limits of the Mahalanobis distances, the 10% external states (outliers) should sooner or later end up in this hyper-ellipsoid if the local network had attracting forces. We used the Matlab® function mahal. The Mahal$_{Threshold}$ = 90% percentile of Mahalanobis distances of all states from the center of the hyper-ellipsoid. The states included in the hyper-ellipsoid < Mahal$_{Treshold.}$
Figure S4G shows the probability densities of all states. We defined the putative attractor as the enclosed volume V$_{90\%}$ containing 90% of all Mahalanobis distances. Thus 10% by definition were outliers.

Since the hyper-volume, V, itself fluctuated over a scale from 0.2 to 10 Hz, during which V ranged from small to maximal, the attracting property should be universal, present when V was small, medium and large. As the distribution of fluctuations in log$_{10}$V was near Gaussian (**Figure 3D**) we defined small logV$_{90\%}$ as the lower 16 % of the distribution of the 90% Mahalanobis distances in each patient, medium as the mid 68% logV$_{90\%}$, and large as the top 16% of logV$_{90\%}$.
The algorithm then located small medium and big states of logV$_{90\%}$ and listed the length of the outlying states, i.e. the time in ms from departure to the return to the hyper-ellipsoid. This was done for all conditions and electrode leads (n=3476) **(Figure 3E, F, G)**.

Flow of states. The **F**(t) was visualized in the integer dimension just lower than the correlation dimensionality. If the correlation dimensionality was 6.6, for example, we chose 6 dimensions for visualization (**Figure 3**). Six dimensions gives 6 orthogonal planes (**Figure S4**). The planes were defined by their normal vector and origin. The set contains only one, in this case, 6-dimensional trajectory, describing the evolution of states. Note that each state is identified in state space as, in this example, a 6-dimensional coordinate [μV$_{ti}$, μV$_{ti+1\tau}$, μV$_{ti+2\tau}$, μV$_{ti+3\tau}$, μV$_{ti+4\tau}$, μV$_{ti+5\tau}$]. The states in state space progress every 1/512 s from one 6-dimensional μV value to the next. The 6-dimensional μV value however is composed from 6 samples spread in time over 360

ms. This is how the flow of states must be interpreted. If the Euclidian distances between successive 6-dimensional μV values increase, the flow of states increases. When Euclidian distances between successive 6-dimensional μV values decrease, the flow of states decreases.

A relatively detailed description of how the trajectories behave can be obtained from calculations of how long time it takes for states to cross one of the symmetry planes (**Figure S4**). As an example, depending on where in this 6-dimensional space a continuous set of states are, they may first cross for example the 3rd symmetry plane (from one side or the other), after a while they may cross the 1st symmetry plane and after yet a while the 5th and so on. The cross time is the number of states positioned on one side of the plane until they cross that plane (**Figure S4C, E**, **Figure S6**). Cross times were calculated for the whole material of 60 s files (n=3476) (**Figure S6).**

To further prove the presence of attracting forces, we calculated flow vectors (Forsberg et al., 2016) as the average velocity over 4 samples (7.81 ms). In **Figure S4**, only the flow vectors after the time point where the Mahalanobis distance was maximal are shown. We calculated also the Euclidian distance in μV between the point, where they exceeded 95% limit and the point of return to the enclosing volume in state space for all 3476 files (**Figure S4**).

Single trial analysis of the fluctuating attractor

For the conditions in which the patients were stimulated by visual stimuli (visual tests 4, 5, 6, 11; emotions) the start of the stimulus is shown in **Figure 6.** For the conditions in which the patients were verbally prompted to recall faces and places (memory retrieval tests 1, 2) and the classification of nouns, the trials are marked at the peak in the analog sound track (sampled with 30 KHz). For the visual stimuli conditions, the single trials were aligned to the start of the stimulus from the photodiode analog track (sampled with 30 KHz). An epoch from -4000 ms to + 4000 ms was chosen. For each of these epochs, the log V signal was converted to z-scores by the Matlab® function z-score. Using the peak in the sound track (providing the most well-defined trigger signal) to align all single trials in the 60 s episodes, segments, 8000 ms long, of z-scored V were calculated. Thereafter all 8000 ms segments, one for each trial, were averaged and the average V plotted (**Figures 6 and 7).** For the conditions in which the patients were verbally prompted to recall faces and places (memory retrieval tests 1,2) and the classification of nouns, the single trials were aligned to the peak in the sound track and similarly treated. This was done for each lead.

For each behavioral condition, the cross correlation between all leads of their individual average z-score was calculated by the Matlab® function xcorr using the option coef, for the time interval -1000 ms to + 2000 ms. Thereafter the pairwise lag between any 2 leads was calculated. A 95% confidence interval of the pairwise cross correlations served to separate significant pairs from random fluctuations.

Anatomical Registration

Image registration was applied for each patient's images in order to compare all the subjects in a unique reference space. The applied procedure is depicted in **Figure S7.** The MNI152 atlas is used as the global reference standard brain (Fonov et al., 2009).

Each individual MRI image with all electrode leads marked was transformed into standard format and merged with the MRIs in standard format of the other patients.