Macrophage-derived insulin/IGF antagonist ImpL2 regulates systemic metabolism for

mounting an effective acute immune response in *Drosophila*

Gabriela Krejčová[1#], Adam Bajgar[1#*], Pavla Nedbalová[1], Julie Kovářová[2], Nick Kamps-Hughes[3], Helena

Zemanová[1], Lukáš Strych[1], Tomáš Doležal[1*]

[1]University of South Bohemia, Czech Republic, [2]Biology Centre CAS, Czech Republic, [3]University of Oregon,

United States

[#]equal contribution

[*]For correspondence: AB - bajgaa00@prf.jcu.cz TD - tomas.dolezal@prf.jcu.cz

Abstract

In response to invading pathogens, macrophages metabolically polarize towards Hif1α-induced

aerobic glycolysis, requiring increased supply of nutrients. Here, we show that in order to obtain

sufficient resources, *Drosophila* macrophages release the insulin/IGF antagonist ImpL2, whose

expression is regulated by Hif1α. ImpL2 remotely induces the release of lipids and carbohydrates

from adipose tissue by reducing insulin signaling, followed by increased nutrient accumulation in

activated immune cells. ImpL2 thus translates the metabolic requirements of immune cells into a

systemic metabolic switch. Although these ImpL2 effects are essential during the acute immune

response to streptococcal infection, they become maladaptive upon chronic infection by an

intracellular pathogen. The relevance of our model to mammalian immunometabolism is

demonstrated by the increased expression of the ImpL2 homolog IGFBP7 in human macrophages

exposed to *Streptococcus*.

Keywords:

macrophage, immuno-metabolism, Drosophila, immunity, bacterial infection, Streptococcus, ImpL2,

IGFBP7, energy mobilization, Hif1α, aerobic glycolysis, macrophage polarization, insulin resistance,

selfish immune system, adipose tissue remodeling, Foxo, infection-induced insulin resistance,

Listeria, Insulin/IGF antagonist, wasting, cachexia

Introduction

Macrophages represent the front line of defence against invading pathogens. Although the

effectiveness of the immune response correlates with the number of immune cells (Nicholson and

Nicholson 2008), their maintenance requires energy and their excessive activation can lead to a

myriad of pathologies and metabolic disorders (Shattuck-Heidorn et al. 2016; Zmora et al. 2017).

Animals have therefore evolved a strategy that allows them to maintain sufficient numbers of

quiescent immune cells that can be rapidly activated in response to the detection of pathogen- or

danger-associated molecular patterns (Kelly and O'Neill 2015). Numerous populations of sentinel

macrophages wait for activating stimuli without presenting a substantial energy burden. As a

consequence, macrophages depend on rapid and sufficient supply from external sources, making the

acute phase of the immune response challenging for the whole organism (Newsholme et al. 1986).

In response to the recognition of an invading pathogen, macrophages must rapidly alter their

metabolism to generate enough energy and precursors to support their bactericidal function.

Bactericidal (M1) macrophages therefore substantially increase the rate of glycolysis and the pentose

phosphate pathway and rewire their mitochondrial metabolism in a Hif1α-dependent manner (Van

den Bossche, O'Neill, and Menon 2017). Such a metabolic setup resembles the Warburg effect, which

was originally described as a unique metabolic program for cancer cells (Warburg 1925). We have

recently shown that the metabolic polarization of macrophages is an ancient and evolutionarily

conserved process, as *Drosophila* macrophages also undergo a Hif1α-triggered metabolic switch that

is essential for their bactericidal function (Krejčová et al. 2019).

An adverse aspect of M1 polarization is that these cells require more energy and become functionally

dependent on external sources of glucose, glutamine, and lipids. Therefore, macrophages release

signals in response to their metabolic activation that regulate systemic metabolism, thus securing

nutrient supply at the expense of other organs (Straub 2014). Such privileged behavior, in which

macrophages usurp nutrients from other processes, is crucial for an effective immune response

(Bajgar and Dolezal 2018). One factor, mediating such behavior, is extracellular adenosine, which

links the current metabolic state of activated immune cells to the systemic mobilization of

carbohydrates that serve as a resource for immune defense (Bajgar et al. 2015; Bajgar and Dolezal

2018). However, this response is very complex and we assume the existence of other signaling

factors with an analogous function.

To discover other signaling factors that are released by activated immune cells and regulate systemic

metabolism, we sought inspiration from neoplastic tumor research. This idea is based on the notion

that tumors and activated immune cells share common features of their cellular metabolism, as both

utilize aerobic glycolysis triggered by Hif1α (Biswas and Mantovani 2012; Nagao et al. 2019). They

also share an impact on systemic metabolism, as both cancer and sepsis patients exhibit a phenotype

similar to the wasting caused by cytokine-induced insulin resistance (Dev, Bruera, and Dalal 2018).

Although insulin resistance is mostly studied as a pathological condition, its evolutionary

conservation indicates that it must carry an adaptive physiological function (Soeters and Soeters

2012; Odegaard and Chawla 2013). We therefore hypothesized that activated immune cells could

release the same factors as tumor cells, but with a beneficial role for the acute response as opposed

to cancer-induced cachexia.

In this study, we focus on the insulin/IGF antagonist ImpL2 (Imaginal morphogenesis protein-Late 2),

which is released by neoplastic tumor cells, to suppress insulin signaling via binding to Drosophila

insulin-like peptides, thereby causing energy wasting (Alee 2011; Arquier et al. 2006; Honegger et al.

2008; Kwon et al. 2015; Figueroa-Clarevega and Bilder 2015). In addition to its production by

neoplastic tumors, ImpL2 is known to be released by lipid-overloaded macrophages (Morgantini et al.

2019), as well as by other cells employing Hif1α activity-dependent metabolic programs in *Drosophila*

(Alee 2011; Owusu-Ansah, Song, and Perrimon 2013; Kwon et al. 2015; Figueroa-Clarevega and Bilder

2015). We therefore decided to test the role of ImpL2 as a macrophage-derived signaling factor that

may be responsible for nutrient mobilization during the acute phase of immune response to bacterial

infection.

Here we show that activated macrophages produce ImpL2 in a Hif1-dependent manner during the

acute phase of infection, resulting in Foxo-mediated changes in adipose tissue metabolism. As an

outcome of ImpL2 action, we observed increased titers of circulating carbohydrates and lipids and

their accumulation in macrophages. ImpL2 release by macrophages is necessary for resistance to

streptococcal infection. In contrast to this beneficial role in fighting extracellular pathogens, the

effects of ImpL2 are maladaptive in response to intracellularly growing *Listeria*. Conservation of

ImpL2 function between insects and mammals is indicated by increased expression of the ImpL2

homolog IGFBP7 in human macrophages exposed to streptococci.

Results

ImpL2 expression increases in immune-activated macrophages in a Hif1α-dependent manner

To test the potential role of ImpL2 during infection, we first monitored its expression profile during

the acute phase of *Streptococcus pneumoniae* infection. ImpL2 expression increased significantly in

infected flies compared to PBS-injected controls, as early as 3 hours post-infection (hpi), reaching up

to a threefold increase in expression 21 hpi on an organismal level (Figure 1A). To identify the tissues

responsible for the infection-induced rise in ImpL2 expression, we employed the Gal4 driver specific

for the *ImpL2-RA* transcriptional variant (Bader et al. 2013) to drive expression of the fluorescent

marker UAS-mCherry. The pattern of ImpL2-positive cells resembled the characteristic distribution of

hemocytes in adult flies (Figure 1B), and their number increased substantially upon infection (Figure

1B and C). The ImpL2-RA>mCherry marker clearly colocalized with hemocyte-specific antibody

against the scavenger receptor Nimrod C1 (NimC1; Figure 1D). To verify that cells expressing ImpL2-

RA>mCherry are macrophages, we injected flies with the phagocytic marker *S. aureus*-pHrodo-Green,

and it indeed colocalized with these cells (Figure 1-figure supplement 1). Cells expressing ImpL2-

RA>mCherry also actively recognized and engulfed *S. pneumoniae ex vivo* (Figure 1E). The evidence

that ImpL2 is produced by macrophages during infection is further supported by the more than

sixfold increase in expression in CrqGal4>UAS-GFP labeled (Clark et al. 2011) FACS sorted

macrophages (Figure 1F).

As there are three alternative transcriptional start sites for ImpL2 (Figure 1I), we analyzed the

expression pattern of each isoform at 24 hpi. Among all analyzed isoforms, the expression of *ImpL2-*

RA was the highest of all transcriptional variants in macrophages and was hardly detectable in fat

body or muscles (Figure 1-figure supplement 2). While the *ImpL2-RB+D* forms were the most

abundant variants in adipose tissue, their expression was five times weaker than *ImpL2-RA*

expression in macrophages (see different y-axis scales in Figure 1-figure supplement 2). In addition,

expression of the *ImpL2-RA* isoform increased more than sixfold in macrophages, the most significant

increase in expression of the transcripts in response to infection (Figure 1F and Figure 1-figure

supplement 2). Thus, the *ImpL2-RA* transcriptional variant expressed in macrophages contributes

significantly to the overall increase in ImpL2 expression observed at the onset of infection.

The metabolic switch in activated macrophages is regulated by the transcription factor Hif1α

(Krejčová et al. 2019), which is known to be a potent regulator of ImpL2 expression in cells, which

rely on anaerobic metabolism (Allee 2011; Li et al. 2013; Owusu-Ansah, Song, and Perrimon 2013).

We therefore hypothesized that this same regulation could underlie the infection-induced increase in

ImpL2 in activated macrophages. Macrophage-specific knockdown of Hif1α (for efficiency, see Figure

1-figure supplement 3) resulted in the inability of these cells to trigger the characteristic infection-

induced expression of ImpL2, and this effect was particularly evident for the *ImpL2-RA* isoform

(Figure 1G). We identified a cluster of four Hif1α binding sites upstream of the transcription start site

of *ImpL2-RA* (Figure 1I). Moreover, this enhancer region shows the strongest hypoxic induction of all

sequence surrounding the *ImpL2* gene (Kamps-Hughes et al. 2015 and Figure 1I), suggesting that

Hif1α may directly drive *ImpL2-RA* transcription in macrophages. In-depth analysis of this 500bp

region revealed the presence of eight immune and stress response elements clustered together (four

hypoxia response elements, two Relish bindings sites, and two heat shock factor binding sites; Figure

1I and Figure 1-figure supplement 4). Chip-qPCR analysis revealed direct binding of Hif1α to this part

of the *ImpL2-RA* promoter, and this interaction was further enhanced after infection (in Figure 1H

denoted as region 5).

These experiments identified activated macrophages as prominent producers of ImpL2 in infected

adult flies. The increase of ImpL2 production was associated with a Hif1α-induced metabolic switch

in macrophages, through direct binding of Hif1α to the regulatory sequence of the *ImpL2-RA* isoform.

136   Macrophage-derived ImpL2 is required for Foxo-mediated mobilization of fat body reserves

Activated macrophages release signaling factors that mobilize reserves to provide sufficient nutrients

for the activated immune system (Bajgar and Dolezal 2018; Dolezal et al. 2019). Here we show that

macrophages increase ImpL2 production during infection. As ImpL2 has previously been associated

with reserve mobilization, leading to wasting in flies with experimentally induced neoplastic growth,

we further tested the impact of macrophage-derived ImpL2 on systemic metabolism. To do so we

employed *Drosophila* genetic tools to manipulate ImpL2 expression specifically in macrophages

(Crq>Gal4) or specifically in macrophages temporally restricted to the adult stage (Hml>Gal4,

Gal80^{TS}). Using conventional ImpL2 knockdown (ImpL2^{RNAi}) and overexpression (ImpL2^{CDS}) constructs,

we achieved significant changes in ImpL2 expression (Figure 1F). These experimental manipulations

allowed us to either prevent infection-induced upregulation of ImpL2 in macrophages or to simulate

the increase in ImpL2 expression in uninfected individuals (Figure 1F).

Both infection and overexpression of ImpL2 in macrophages significantly reduced triglyceride content

in whole flies (Figure 2A), which was accompanied by dramatic changes in adipose tissue morphology

(Figure 2B-E). The number of lipid droplets increased as their average size decreased (Figure 2B, C,

and E), making the lipids more accessible for lipases. The overall area of adipose tissue occupied by

lipid droplets was markedly smaller (Figure 2D). It should be emphasized that these infection-induced

effects were suppressed by macrophage-specific ImpL2 knockdown (Figure 2A-E). Detailed lipidomic

analysis of adipose tissue by mass spectrometry revealed that both infection and macrophage-

specific overexpression of ImpL2 caused a proportional shift in lipid content from storage lipids

(triglycerides) to polar lipid species (phosphatidylethanolamine,   phosphatidylinositol), which

participate in lipid mobilization and transport (Figure 2F). These effects were reduced by ImpL2

knockdown (Figure 2F). Similar effects were also observed for glycogen stores as both infection and

overexpression of ImpL2 significantly reduced glycogen content in whole flies, while knockdown of

ImpL2 abolished this effect (Figure 2G). This data thus indicate that macrophage-derived ImpL2 may

serve as a mediator of cross-talk between the immune system and lipid metabolism in the adipose

tissue.

In addition to a substantially increased lipolytic and glycogenolytic programs, we also observed

enhanced autophagy in the adipose tissue of infected flies bearing the Atg8a-mCherry reporter

(Figure 3A), further implying an increase in catabolic metabolism. These infection-induced metabolic

changes are consistent with the expression of many metabolic genes in adipose tissue. The

expression of *MTP*, *apoLPP*, *apoLTP,* and *Bmm* genes associated with lipid mobilization, as well as

Atg1 and *Atg6* genes associated with autophagy, were upregulated in adipose tissue during infection

(Figure 3B and Figure 3-figure supplement 1). Furthermore, their expression is under the control of

macrophage-derived ImpL2. Indeed, knockdown of ImpL2 suppressed this effect, whereas

overexpression of ImpL2 mimicked the response induced by infection (Figure 3B and Figure 3-figure

supplement 1).

Many of the aforementioned metabolic genes whose expression is altered by ImpL2 during infection

(Figure 3B) are known Foxo targets, therefore we analyzed the impact of macrophage-derived ImpL2

on the subcellular localization of Foxo in adipose tissue. We found that Foxo displays nuclear

localization upon infection, whereas it remained predominantly in the cytoplasm of adipocytes in

uninfected flies (Figure 4). Macrophage-specific overexpression of ImpL2 recapitulated the infection-

induced effect on Foxo nuclear localization, and conversely, knockdown of ImpL2 reversed this effect

(Figure 4). These effects of ImpL2 on Foxo localization are in agreement with the expression of Foxo

target genes (Figure 3B and Figure 3-figure supplement 1). To verify that the observed effects of

macrophage-derived ImpL2 on the fat body are indeed mediated by Foxo, we tried to rescue the

effect of ImpL2 overexpression in hemocytes by hypomorphic mutation of *Foxo*. The lipid droplet

phenotype induced by ImpL2 overexpression was completely reversed by the heterozygous

foxo^BG01018 mutation (Figure 5A), demonstrating that the effects of macrophage-derived ImpL2 on fat

body metabolism are indeed mediated by Foxo.

ImpL2 is known to antagonize insulin signaling by binding to *Drosophila* insulin-like peptides

(Honegger et al. 2008), which is here supported by effects on Foxo in the fat body, a known target of

insulin signaling. Therefore, we tried to check the state of insulin signaling in the fat body by the

commonly used PI3K reporter tGPH and phosphorylation of Akt. In response to infection, the tGPH

reporter showed increased cytosolic localization in comparison to control flies when analyzed by

confocal microscopy (Figure 5B). This effect was phenocopied by overexpression of ImpL2 in

macrophages even in the absence of infection (Figure 5B), demonstrating the ability of macrophage-

derived ImpL2 to antagonize insulin signaling in the fat body. Unfortunatelly, the effect of infection

on tGPH localization was not strong enough to clearly see a difference after knocking down Impl2 in a

double-blind evaluation. pAkt also appeared to be too variable during *S. pneumoniae* infection,

precluding the reasonable use of Akt phosphorylation in our model.

Macrophage-derived ImpL2 increases lipids and carbohydrates both in circulation and in

macrophages

Stimulated glycogen and triglyceride catabolism is manifested by hyperglycemia and hyperlipidemia

in the circulation of infected flies. Elevated titers of glucose, trehalose, glycerides and free fatty acids

were detected in the hemolymph of these flies (Figure 6A and Figure6-figure supplement 1).

Increased levels of circulating lipids and carbohydrates were accompanied by increased accumulation

of these energy-rich compounds in infection-activated macrophages (Figure 6B). These results are

supported by the higher occurrence of lipid droplets in the cytosol of infection-activated

macrophages after staining with the neutral lipid dye OilRedO (Figure 6C and D). The ability of

macrophages to uptake mobilized lipids was verified by injection of fluorescently labeled lipoproteins

(LDL-pHrodo; Figure 6E and F). These lipoproteins can be endocytosed via recognition by the

scavenger receptor Croquemort (fly homolog of mammalian CD36), which is abundantly expressed in

Drosophila macrophages. Injection of different concentrations of LDL-pHrodo showed that

lipoprotein uptake by macrophages correlates with the amount of lipoproteins in the circulation,

even beyond the physiological concentrations commonly occuring in the hemolymph (Gilbert and

Chino 1974). Reserve mobilization, which leads to an increase in circulating nutrients and their

subsequent accumulation in activated macrophages, depends on the production of ImpL2 by

macrophages. Indeed, knockdown of macrophage-specific ImpL2 suppresses all observed effects

stimulated by infection, whereas overexpression of ImpL2 mimicked these effects even in uninfected

individuals (Figure 6A-D).

Taken together, the above results suggest that ImpL2 produced by macrophages during infection

affects insulin signaling in adipose tissue, thereby triggering a Foxo-mediated transcriptional program

that provides macrophages with the nutrients they require.

ImpL2 is required for an effective immune response but can be detrimental during chronic

infection

The efficiency of the immune response depends on an adequate supply of energy and essential

precursors to activated immune cells. Therefore, we decided to investigate whether ImpL2-mediated

release of reserves is necessary for an effective immune response. Lack of ImpL2 production by

macrophages significantly reduced survival of flies infected with *S. pneumoniae* (Figure 7A and Figure

7-figure supplement 1). This infection is associated with elevated pathogen load (Figure 7B),

indicating decreased resistance in these individuals. Elimination of bacteria and survival of *S.*

pneumonia infection is critically dependent on efficient phagocytosis, otherwise flies succumb to

infection within two days (Bajgar and Dolezal 2018). Indeed, reduced resistance in flies with knocked-

down ImpL2 is associated with reduced phagocytic rates (Figure 7E and F). Infection-induced

increases in antimicrobial peptide production by macrophages were reduced by ImpL2 knockdown

for two of the three peptides analyzed (Figure 7-figure supplement 2). On the other hand,

overexpression of ImpL2 24 hours prior to infection improved resistance to streptococcal infection,

as evidenced by lower pathogen load and improved survival (Figure 6C and D). However, we did not

detect a difference in phagocytosis and expression of the two antimicrobial peptides during infection

is rather reduced in flies overexpressing ImpL2 (Figure 7-figure supplement 2), leaving the reasons for

improved survival unclear.

Although our results demonstrate a beneficial role of ImpL2 during the acute phase of infection,

ImpL2 has previously been associated with detrimental effects via induction of chronic cachexia-like

wasting in a model of neoplastic tumor in *Drosophila* (Kwon et al. 2015). This suggests that the

beneficial role of ImpL2 may be restricted to the short period of the acute phase of the immune

response. Therefore, we tested the effects of ImpL2 manipulations in flies challenged with chronic

infection caused by the intracellular pathogen *Listeria monocytogenes*. Flies injected with Listeria are

unable to eliminate these bacteria and the length of their survival is determined by disease

tolerance (Louie et al. 2016) whereas virulence and intracellular growth of *Listeria* depend on the

availability of nutrients in the cytosol of the host cell (Chen, Pensinger, and Sauer 2017).

Concordantly, the overexpression of ImpL2, which led to increased nutrient supplementation in

immune cells, resulted in chronically increased *Listeria* burden (Figure 8A and B). This indicates that

the originally beneficial effects of ImpL2 observed during the acute response to streptococcal

infection may become detrimental during chronic immune challenge. Consistent with this, silencing

of ImpL2 production by macrophage significantly reduced the intracellular load of *L. monocytogenes*

at both 24 hpi and 12 days post-infection (Figure 8C and D), which may be explained by either a

decreased ability of macrophages to engulf *Listeria* or a reduction of nutrients in their cytosol.

In conclusion, ImpL2-mediated resource mobilization is essential for an adequate antibacterial

immune response and resistance to infection. On the other hand, chronic or excessive production of

ImpL2 tends to intensify the deleterious effects of chronic intracellular infections.

Expression of *IGFBP7,* a mammalian homolog of ImpL2, is upregulated in immune-activated THP-1

cells

It has previously been shown that IGFBP7, the mammalian homologue of ImpL2, is produced by lipid-

loaded liver macrophages in obese mice and human patients, with consequent effects on systemic

metabolism through the regulation of insulin signaling in hepatocytes. ImpL2 released from lipid-

loaded macrophages appears to play an analogous role in *Drosophila* fed a haigh-fat diet (Morgantini

et al. 2019). To test the potential importance of the role of IGFBP7 in humans during infection, we

measured the expression of IGFBPs in the PMA-activated THP-1 human monocytic cell line 24 hours

after exposure to *S. pneumoniae* bacteria (Figure 9A). Expression of several IGFBP family members,

namely *IGFBP1, IGFBP2, IGFBP5,* and *IGFBP6,* was not detected, indicating that these genes are not

abundantly expressed in these cells (data not shown). Interestingly, *IGFBP7* gene expression was

significantly increased in inoculated culture compared to PBS-treated controls (Figure 9B). In

addition, the expression of the *IGFBP3* gene, which inhibits insulin resistance (Mohanraj et al. 2013),

was reduced fourfold in activated THP-1 cells (Figure 9B), which is in agreement with the expected

pattern associated with the adoption of bactericidal polarization by mammalian macrophages. This

data therefore suggest that the increase in ImpL2 / IGFBP7 production by macrophages in response

to infection is evolutionarily conserved in mammals.

Discussion

In this work, we show that infection-induced Hif1α transcriptional activity connects an intracellular

metabolic switch to ImpL2 production in activated macrophages (Figre 10). ImpL2 subsequently

induces the mobilization of resources from adipose tissue into the circulation, which may then

become available to activated immune cells. This metabolic rearrangement of adipose tissue is most

likely mediated by suppression of insulin signaling and subsequent transcriptional activity of Foxo.

Such a program results in accelerated lipolysis and carbohydrate mobilization and the increased titer

of circulating lipids and carbohydrates facilitates their utilization by macrophages. Although not

directly tested in this work, these macromolecules are known to be important for bactericidal

function of macrophages (Remmerie and Scott 2018). Indeed, here we show that such ImpL2-

induced metabolic adaptation is essential for the acute immune response, but becomes maladaptive

in the case of infection by intracellular pathogens that metabolically exploit the host cell. This is a

remarkable example of how a beneficial metabolic program can become maladaptive in chronic

diseases. This mechanism may not be limited to insects, indicated by an experiment showing that

human macrophages activated by the same bacteria increase the expression of IGFBP7, the human

homolog of *Drosophila* ImpL2.

We have previously shown that *Drosophila* macrophages undergo an evolutionarily conserved

polarization to the M1 phenotype and thus their metabolism adjusts to Hif1α-induced aerobic

glycolysis in response to bacterial challenge. This cellular metabolic shift is accompanied by systemic

metabolic changes necessary to supplement the sudden nutritional needs of macrophages (Krejčová

et al. 2019). Coordination of cellular metabolism in macrophages with systemic metabolism is crucial

for resistance to infection and implies the existence of circulating factors mediating this interorgan

crosstalk (Dolezal et al. 2019). Inspired by research on neoplastic cancer cells (Kwon et al. 2015;

Figueroa-Clarevega and Bilder 2015; Bunker et al. 2015), in this study we focus on the cachectic

factor ImpL2 as a candidate that has previously been linked to Hif1α transcriptional activity while

having the potential to regulate systemic metabolism (Alee 2011; Li et al. 2013). To test whether the

link between Hif1α activity and ImpL2 expression during hypoxia also applies to infection-activated

macrophages, we performed a genomic *in silico* meta-analysis of hypoxic enhancer activity and Hif1α

binding sites followed by Chip-qPCR to confirm the binding of Hif1α to previously identified genomic

loci (Kamps-Hughes et al. 2015). This approach revealed the induction of a hypoxic enhancer in close

proximity to the transcription start site of the *ImpL2-RA* transcript variant, which is further supported

by the presence of four hypoxia response elements at this locus. *ImpL2-RA* is the most strongly

expressed transcriptional variant in macrophages and its expression is further increased after

infection. This infection-induced increase is Hif1α dependent, indicating a connection between ImpL2

expression and an internal metabolic switch in infection-activated macrophages. We have previously

identified extracellular adenosine as a systemic factor derived from immune cells, with an analogous

function to ImpL2, mobilizing carbohydrate stores to be available to the immune system (Bajgar et al.

2015; Bajgar and Dolezal 2018). While adenosine is a signaling molecule whose generation depends

on the rate of cellular metabolism, the production of ImpL2 protein is linked to the transcriptional

activity of the central metabolic regulator Hif1α. Thus, it is noteworthy that regulation of systemic

metabolism can be mediated by factors with different mechanisms of production that reflect cellular

metabolism of immune cells.

In addition to the regulation of carbohydrate metabolism, a prominent effect of ImpL2 is its

regulation of systemic lipid metabolism. The metabolic changes observed in infection-challenged flies

resemble lipemia as a characteristic symptom of sepsis in severely ill patients (Harris, Gosnell, and

Kumwenda 2000). In sepsis-induced lipemia, insulin-resistant hepatocytes elevate the level of

circulating lipids, which are preferentially utilized by activated immune cells in the periphery

(Khovidhunkit et al. 2004; Aspichueta et al. 2012). Scavenger receptor-mediated endocytosis of

serum lipids together with attenuated reverse cholesterol transport contribute substantially to

cholesterol accumulation in bactericidal immune cells (Podrez et al. 2000). A number of immune-

related macrophage functions depend on sufficient delivery of lipids, ranging from phagocytosis and

phagolysosome maturation, to catecholamine production and immune memory formation via the

mevalonate pathway (summarized in Remmerie and Scott 2018). Although providing macrophages

with substantial amounts of lipids is essential for their proper function during the acute phase of

infection, chronic exposure of macrophages to excessive lipids can lead to the adoption of a foam-

cell phenotype, which promotes metabolic syndrome and atherosclerosis (Chistiakov et al. 2017;

Febbraio, Guy, and Silverstein 2004).

Although we have not yet fully elucidate the mechanism of infection-induced lipid catabolism, we

found that nuclear translocation of Foxo, followed by increased expression of several Foxo-target

genes involved in this process, is regulated by macrophage-derived ImpL2. Foxo-driven mobilization

of lipid stores requires suppression of insulin signaling in adipose tissue (Molaei, Vandehoef, and

Karpac 2019; Luong et al. 2006), which is in agreemnet with the well-documented ability of ImpL2 to

bind to *Drosophila* insulin-like peptides (DILPs) and thereby reduce insulin signaling in adipose tissue.

This is further supported by our observations of reduced adipocyte insulin signaling in infected flies

and flies with experimentally enhanced ImpL2 production in macrophages (Alee 2011; Alic et al.

2011; Arquier et al. 2006; Figueroa-Clarevega and Bilder 2015; Honegger et al. 2008; Kwon et al.

2015; Okamoto et al. 2013). Metabolic adjustments and the switch from anabolism to catabolism in

adipose tissue during infection are regulated by multiple mechanisms (reviewed in Dolezal et al.

2019). Specifically, Foxo has previously been linked to wasting in flies during chronic infection

(Dionne et al. 2006), suggesting its pathological effects during a prolonged immune response. In

contrast, the remote effect of ImpL2 on reserve mobilization described in this work represents a

rather beneficial role for ImpL2 during the acute immune response. Suppression of insulin signaling is

often associated with chronic inflammation and can become detrimental to the organism by

disrupting metabolic balance. However, the concept of selfish immunity (Straub 2014) considers

insulin resistance as an evolutionary adaptive mechanism for rerouting nutrients toward the immune

system during the acute immune response. Our results showing that ImpL2 is required for an

effective immune response through changes in lipid and carbohydrate metabolism can be considered

as experimental evidence for insulin resistance as an adaptive mechanism for resource mobilization

in the acute phase of the immune response.

Neoplastic tumors share a characteristic cellular metabolism (Warburg 1925) with bactericidal

macrophages (Andrejeva and Rathmell 2017). We suggest that they also share the production of

ImpL2, which was originally perceived as a cancer-derived cachectic factor (Kwon et al. 2015;

Figueroa-Clarevega and Bilder 2015), released from these cells to usurp energy from other tissues via

cytokine-induced insulin resistance. The perception of ImpL2 expression as a consequence of cellular

metabolic settings is supported by other conditions when ATP generation is independent of

mitochondria, such as experimentally induced mitochondrial dysfunction, hypoxia, and neoplastic

tumor growth (Alee 2011; Figueroa-Clarevega and Bilder 2015; Kwon et al. 2015; Li et al. 2013).

Although this mechanism of metabolic regulation is beneficial to the organism during acute infection,

it becomes detrimental in the case of tumor growth or chronic infection. Elimination of *S. pneumonia*

requires effective phagocytosis – the flies either clear the infection or die. When phagocytosis is

blocked, flies succumb to infection within two days (Bajgar and Dolezal 2018), so this immune

response can be regarded as acute. ImpL2-mediated metabolic changes appear to be important for

such a response. On the other hand, the harmful effect of ImpL2 on the survival of infection caused

by the intracellular pathogen *L. monocytogenes* is demonstrated here. The flies are unable to

eliminate these bacteria, which escape from the phagosome and establish a chronic intracellular

infection from which the flies sooner or later die. In this case, reducing phagocytosis by knocking

down ImpL2 lowered the short- and long-term intracellular load of the pathogen, leading to longer

host survival. Various intracellular bacteria are known to take advantage of host cell supply and

hijack their metabolic cascades to literarily nourish themselves (Péan et al. 2017; Teng, Ang, and

Guan 2017). We do not know whether the observed increased *L. monocytogenes* burden in flies

overexpressing ImpL2 is due to increased nutrients available to the pathogen, but these flies die

more rapidly, indicating a rather negative role for ImpL2 during chronic infection.

In this study, we revealed that human macrophages activated by *S. pneumoniae* upregulate the

expression of IGFBP7, the human homolog of *Drosophila* ImpL2, suggesting that a similar mechanism

of metabolic regulation may operate in humans. Remarkably, not only infection but also

experimentally induced metabolic syndrome in both *Drosophila* and mammals led to increased

ImpL2/IGFBP7 expression in macrophages (Morgantini et al. 2019). While ImpL2-producing

macrophages affect metabolism in the *Drosophila* fat body, which integrates functions of both

mammalian adipose tissue and liver, IGFBP7-producing Kupffer cells act directly on hepatocytes to

induce systemic metabolic changes (Morgantini et al. 2019; Akiel et al. 2017). The ability of both

ImpL2 and IGFBP7 to bind extracellular insulin, thereby reducing systemic insulin signaling, further

documents their functional homology, suggesting that their mechanism of action may also be

conserved (Honegger et al. 2008; Arquier et al. 2006; Oh et al. 1996). Although this mechanism of

resource mobilization may be important for the acute phase response, when prolonged it contributes

to the progression of metabolic syndrome and atherosclerosis in mammals (Tomkin 2012). It is

therefore not surprising that circulating IGFBP7 levels are considered a marker of systemic metabolic

imbalance accompanying several human diseases such as obesity, acute kidney injury, liver fibrosis,

and chronic obstructive pulmonary disease (Liu et al. 2015; Gunnerson et al. 2016; Martínez-Castillo

et al. 2020; Ruan et al. 2017).

Combining our data with previous work, we conclude that the insulin/IGF antagonist ImpL2, which is

released from activated macrophages as a reflection of their metabolic polarization, mediates

nutrient mobilization from adipose tissue by reducing insulin signaling to liberate sufficient resources

for the activated immune system. Despite the fact that both ImpL2 and insulin resistance are mostly

studied in the context of pathological conditions such as cancer-induced cachexia, obesity, and

chronic inflammatory conditions, our data revealed that they may play a beneficial role in the acute

phase of bacterial infection. The relevance of our model is further supported by the production of

IGFBP7, a homologue of ImpL2, in response to streptococcal infection in human macrophages. This

leads us to the hypothesis that analogous mechanisms may also apply to macrophage-induced

nutrient mobilization from hepatocytes during the acute phase of the immune response; however,

this relationship remains to be experimentally verified.

Funding

Tomáš Doležal - Czech Science Foundation (Project 20-09103S)

Adam Bajgar - Czech Science Foundation (Project 20-14030S)

Gabriela Krejčová - USB Grant Agency (Project 050/2019/P)

Acknowledgment

The authors acknowledge funding from the Grant Agency of the Czech Republic to TD (Project 20-

09103S; www.gacr.cz) and to AB (Project 20-14030S; www.gacr.cz). GK was supported by USB Grant

Agency (Project 050/2019/P). We thank to Lucie Hrádková for laboratory services, enthusiasm and

support, Alena Krejčí-Bruce and Lenka Chodáková for critical comments and inspiring discussions,

Pavel Branný and Linda Doubravová for help with the preparation of the S.p.-GFP strain and Hana

Sehadová for help with scanning electron microscopy. We thank to Hugo Stocker for the ImpL2^{RNAi},

ImpL2^{CDS}, and ImpL2-RA-Gal4 fly lines, Gabor Juhasz for the Atg8amCherry fly, and Marc Dionne for

Crq>GFP fly line. Other fly stocks were obtained from the Bloomington Center (Bloomington, IN) and

the VDRC (Vienna, Austria). The *S. pneumoniae* and *L. monocytogenes* bacterial strains were obtained

from Dr. David Schneider. We also thank Martin Moss and Petr Šimek for the lipidomics service, the

Department of Medical Biology (USB) for allowing us to use the S3eBioRad sorter, Biology Centre CAS

for allowing us to use a confocal microscope, and a laboratory equipped to maintain human tissue

cultures. We are also grateful to developers of Fiji: an open-source platform for biological-image

analysis (doi:10.1038/nmeth.2019)

Materials and methods

***Drosophila melanogaster* strains and culture**

The flies were raised on a diet containing cornmeal (80 g/l), sucrose (50 g/l), yeast (40 g/l), agar (10

g/l), and 10%-methylparaben (16.7 mL/l) and maintained in a humidity-controlled environment with

a natural 12 h/12 h light/dark cycle at 25°C. Flies carrying Gal80 protein were raised at 18°C and

transferred to 29°C 24 h prior to infection in order to degrade temperature-sensitive Gal80. Prior to

the experiments, flies were kept in plastic vials on a sucrose-free cornmeal diet (cornmeal 53.5 g/l,

yeast 28.2 g/l, agar 6.2 g/l and 10%-methylparaben 16.7 mL/l) for 7 days. Flies infected with *S.*

pneumoniae were kept on a sucrose-free cornmeal diet in incubators at 29°C due to the temperature

sensitivity of *S. pneumoniae*. They were transferred to fresh vials every other day without the use of

CO_2 to ensure good food condition. Flies infected with *L. monocytogenes* were kept on a sucrose-free

cornmeal diet at 25°C. The Drosophila Stock Center in Bloomington provided *ImpL2^{RNAi}* (*y1 sc* v1;*

P{TRiP.HMC03863}attP40; FBst0055855) *TRiP^{control}* (*y[1] v[1] ; P{y[+t7.7]=CaryP}attP40*; FBst0036304)

and *20xUAS-6xmCherry* (*P{20XUAS-6XmCherry-HA}attP2*; FBtp0094992) flies. *ImpL2-RA-Gal4*

(FBal0290965) and *UAS-ImpL2^{cds}* (*UAS-s.ImpL2*; FBal0249386) were kind gifts from Hugo Stocker.

CrqGal4>2xeGFP were obtained from Marc Dionne. The *Atg8a-mCherry* strain was kindly provided by

Gabor Juhasz. The *w^{1118}* strain has a genetic background based on *CantonS*.

Genotypes of experimental models

Figure 1

ImpL2-RA>mCherry corresponds to *w^{1118}/w^{1118}; 20xUAS-6xmCherry/+; ImpL2-RA-Gal4/+*

HmlGal4>GFP refers to *w^{1118}/w^{1118}; HmlΔ-Gal4 UAS-2xeGFP/HmlΔ-Gal4 UAS-2xeGFP; +/+*

HmlGal4>GFP; Hif1α^{RNAi} corresponds to *w^{1118}/+; HmlΔ-Gal4 UAS-2xeGFP/+; UAS-Hif1α^{RNAi}/+*

ImpL2^{RNAi} refers to *w^{1118}/+; HmlΔ-Gal4/ UAS-ImpL2^{RNAi}; P{tubPGal80ts}/+*

ImpL2^{cds} corresponds to *w^{1118}/w^{1118}; HmlΔ-Gal4/+; P{tubPGal80ts}/UAS-ImpL2^{cds}*

TRiP^control refers to *w^1118/+; HmlΔ-Gal4/+; P{tubPGal80ts}/TRiP^control*

w^1118 corresponds to *w^1118/ w^1118; HmlΔ-Gal4/+; P{tubPGal80ts}/+*

Figure 1-figure supplement 3

Hml>Gal4, UAS-GFP x TRiP^control corresponds to *w^1118/+; HmlΔ-Gal4 UAS-2xeGFP/+; TRiP^control/+*

Hml>Gal4, UAS-GFP x Hif1α^RNAi corresponds to *w^1118/+; HmlΔ-Gal4 UAS-2xeGFP/+; UAS-Hif1α^RNAi/+*

Figure 2

ImpL2^RNAi refers to *w^1118/+; HmlΔ-Gal4/ UAS-ImpL2^RNAi; P{tubPGal80ts}/+*

ImpL2^cds corresponds to *w^1118/w^1118; HmlΔ-Gal4/+; P{tubPGal80ts}/UAS-ImpL2^cds*

TRiP^control refers to *w^1118/+; HmlΔ-Gal4/+; P{tubPGal80ts}/TRiP^control*

w^1118 corresponds to *w^1118/ w^1118; HmlΔ-Gal4/+; P{tubPGal80ts}/+*

Figure 3

ImpL2^RNAi refers to *w^1118/+; HmlΔ-Gal4/ UAS-ImpL2^RNAi; P{tubPGal80ts}/+*

ImpL2^cds corresponds to *w^1118/w^1118; HmlΔ-Gal4/+; P{tubPGal80ts}/UAS-ImpL2^cds*

Atg8a-mCherry refers to gen *+/+; Atg8a-mCherry/Atg8a-mCherry*

Figure 3-figure supplement 1

ImpL2^RNAi refers to *w^1118/+; HmlΔ-Gal4/ UAS-ImpL2^RNAi; P{tubPGal80ts}/+*

ImpL2^{cds} corresponds to *w¹¹¹⁸/w¹¹¹⁸; HmlΔ-Gal4/+; P{tubPGal80ts}/UAS-ImpL2^{cds}*

TRiP^{control} refers to *w¹¹¹⁸/+; HmlΔ-Gal4/+; P{tubPGal80ts}/TRiP^{control}*

w¹¹¹⁸ corresponds to *w¹¹¹⁸/ w¹¹¹⁸; HmlΔ-Gal4/+; P{tubPGal80ts}/+*

Figure 4

ImpL2^{RNAi} refers to *w¹¹¹⁸/+; HmlΔ-Gal4/ UAS-ImpL2^{RNAi}; P{tubPGal80ts}/+*

ImpL2^{cds} corresponds to *w¹¹¹⁸/w¹¹¹⁸; HmlΔ-Gal4/+; P{tubPGal80ts}/UAS-ImpL2^{cds}*

TRiP^{control} refers to *w¹¹¹⁸/+; HmlΔ-Gal4/+; P{tubPGal80ts}/TRiP^{control}*

w¹¹¹⁸ corresponds to *w¹¹¹⁸/ w¹¹¹⁸; HmlΔ-Gal4/+; P{tubPGal80ts}/+*

Figure 5

Hml>Gal4 TubGal80 x w¹¹¹⁸ corresponds to *w¹¹¹⁸/w¹¹¹⁸; HmlΔ-Gal4 P{tubPGal80ts}/+; +/+*

Hml>Gal4 TubGal80 x ImpL2^{cds} corresponds to *w¹¹¹⁸/w¹¹¹⁸; HmlΔ-Gal4 P{tubPGal80ts}/+; UAS-*
ImpL2^{cd}/+

Hml>Gal4 TubGal80 x foxo^{BG01018} ImpL2^{cds} refers to *w¹¹¹⁸/w¹¹¹⁸; HmlΔ-Gal4 P{tubPGal80ts}/+;*
P{w[+mGT]=GT1}foxo^{BG01018} UAS-ImpL2^{cds}/+

Hml>Gal4 TubGal80; tGPH-GFP x w¹¹¹⁸ corresponds to *w¹¹¹⁸/ w¹¹¹⁸; HmlΔ-Gal4 P{tubPGal80ts}/+;*
tGPH/+

Hml>Gal4 TubGal80; tGPH-GFP x ImpL2^{cds} corresponds to *w¹¹¹⁸/ w¹¹¹⁸; HmlΔ-Gal4 P{tubPGal80ts}/+;*
tGPH/UAS-ImpL2^{cds}

Figure 6

ImpL2^{RNAi} refers to *w¹¹¹⁸/+; HmlΔ-Gal4/ UAS-ImpL2^{RNAi}; P{tubPGal80ts}/+*

ImpL2^{cds} corresponds to *w^{1118}/w^{1118}; HmlΔ-Gal4/+; P{tubPGal80ts}/UAS-ImpL2^{cds}*

TRiP^{control} refers to *w^{1118}/+; HmlΔ-Gal4/+; P{tubPGal80ts}/TRiP^{control}*

w^{1118} corresponds to *w^{1118}/ w^{1118}; HmlΔ-Gal4/+; P{tubPGal80ts}/+*

Crq>Gal4; UAS2xGFP x ImpL2^{RNAi} refers to *w^{1118}/+; UAS-ImpL2^{RNAi}/+; Crq-Gal4, UAS-2xeGFP/+*

Crq>Gal4; UAS2xGFP x ImpL2^{cds} corresponds to *w^{1118}/ w^{1118}; +/+; Crq-Gal4, UAS-2xeGFP/UAS-*

ImpL2^{cds}

Crq>Gal4; UAS2xGFP x TRiP^{control} refers to *w^{1118}/+; +/+; Crq-Gal4, UAS-2xeGFP/TRiP^{control}*

Crq>Gal4; UAS2xGFP x W^{1118} corresponds to *w^{1118}/ w^{1118}; Crq-Gal4, UAS-2xeGFP/+*

CrqGal4>GFP refers to *w^{1118}/ w^{1118}; Crq-Gal4, UAS-2xeGFP/Crq-Gal4, UAS-2xeGFP*

Figure 7

ImpL2^{RNAi} refers to *w^{1118}/+; HmlΔ-Gal4/ UAS-ImpL2^{RNAi}; P{tubPGal80ts}/+*

ImpL2^{cds} corresponds to *w^{1118}/w^{1118}; HmlΔ-Gal4/+; P{tubPGal80ts}/UAS-ImpL2^{cds}*

TRiP^{control} refers to *w^{1118}/+; HmlΔ-Gal4/+; P{tubPGal80ts}/TRiP^{control}*

w^{1118} corresponds to *w^{1118}/ w^{1118}; HmlΔ-Gal4/+; P{tubPGal80ts}/+*

Crq>Gal4 x ImpL2^{cds} corresponds to *w^{1118}/ w^{1118}; +/+; Crq-Gal4, UAS-2xeGFP/UAS-ImpL2^{cds}*

Figure 7-figure supplement 1

Hml-Gal4; TubGal80^{TS} refers to *w^{1118}/ w^{1118}; HmlΔ-Gal4/+; P{tubPGal80ts}/+*

Hml-Gal4; TubGal80^{TS} > ImpL2^{cds} corresponds to *w^{1118}/w^{1118}; HmlΔ-Gal4/+; P{tubPGal80ts}/UAS-*

ImpL2^{cds}

ImpL2^{cds} x w¹¹¹⁸ corresponds to *w¹¹¹⁸/w¹¹¹⁸; +/UAS-ImpL2^{cds}*

Hml-Gal4; TubGal80^{TS} > ImpL2^{RNAi} corresponds to *w¹¹¹⁸/+; HmlΔ-Gal4/ UAS-ImpL2^{RNAi};*

P{tubPGal80ts}/+

ImpL2^{RNAi} corresponds to *w¹¹¹⁸/+; +/ UAS-ImpL2^{RNAi}*

Figure 7-figure supplement 2

ImpL2^{RNAi} refers to *w¹¹¹⁸/+; HmlΔ-Gal4/ UAS-ImpL2^{RNAi}; P{tubPGal80ts}/+*

ImpL2^{cds} corresponds to *w¹¹¹⁸/w¹¹¹⁸; HmlΔ-Gal4/+; P{tubPGal80ts}/UAS-ImpL2^{cds}*

TRiP^{control} refers to *w¹¹¹⁸/+; HmlΔ-Gal4/+; P{tubPGal80ts}/TRiP^{control}*

w¹¹¹⁸ corresponds to *w¹¹¹⁸/ w¹¹¹⁸; HmlΔ-Gal4/+; P{tubPGal80ts}/+*

Figure 8

ImpL2^{RNAi} refers to *w¹¹¹⁸/+; HmlΔ-Gal4/ UAS-ImpL2^{RNAi}; P{tubPGal80ts}/+*

ImpL2^{cds} corresponds to *w¹¹¹⁸/w¹¹¹⁸; HmlΔ-Gal4/+; P{tubPGal80ts}/UAS-ImpL2^{cds}*

TRiP^{control} refers to *w¹¹¹⁸/+; HmlΔ-Gal4/+; P{tubPGal80ts}/TRiP^{control}*

w¹¹¹⁸ corresponds to *w¹¹¹⁸/ w¹¹¹⁸; HmlΔ-Gal4/+; P{tubPGal80ts}/+*

Hypoxic enhancer activity analysis

Genome-wide experimental data characterizing hypoxia-induced transcriptional enhancer activity

(Kamps-Hughes et al. 2015) was used to analyze the ImpL2 region for hypoxic enhancers. Hypoxic

transcriptional induction is defined as the ratio of expressed randomer tag counts in hypoxic versus

normoxic conditions and was binned by 500-bp regions across the *ImpL2* locus. Each 500-bp bin was

then analyzed for transcription factor binding sites corresponding to hypoxic (Hif1α), immune (Rel)

and stress (HSF) response transcription factors. Transcription factor position frequency matrices

were downloaded from the JASPAR database

(https://academic.oup.com/nar/article/32/suppl_1/D91/2505159) and queried against the 500-bp

bin sequences using BOBRO software (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3074163/).

BOBRO generates a p-value of enrichment for the position frequency matrix within the search

sequence and also identifies the location of the predicted binding sites.

Chip-qPCR assay

The Pro-A Drosophila CHIP Seq Kit (Chromatrap) was used to co-immunoprecipitate genomic regions

specifically bound by the transcription factor HIF1α. A transgenic fly strain carrying the Hif1α protein

fused to GFP (BDSC: 42672) was used for this purpose. The procedure was performed according to

the supplier's instructions. Briefly, the slurry was prepared by homogenizing ten infected or PBS

injected males in three biological replicates. The Rabbit Anti-GFP antibody (ABfinity) was bound to

the chromatographic column. Genomic DNA was fragmented to an approximate size of 500 bp by

three cycles of 60-second sonication. The fragment size was verified by gel electrophoresis. All

samples were tested with positive and negative controls. The amount of precipitated genomic

fragments was normalized to the amount of fragments in slurry before precipitation. The ImpL2-RA

promoter sequence was covered with seven primer pairs corresponding to the 500-bp bins upstream

of the transcription start site previously assessed in the *in silico* analysis. The genomic region of S-

adenosylmethionine synthetase was used as a negative control since it does not contain any

sequences of hypoxia response elements. Primer sequences are listed in the Key Resources Table.

The amount of HIF1α-bound regions of the ImpL2 promoter was quantified on a 96CFX 1000 Touch

Real-Time Cycler (BioRad) using TP 2x SYBR Master Mix (Top-Bio) in three technical replicates with

the following protocol: initial denaturation - 3 min at 95°C, amplification – 15 s at 94°C, 20 s at 56°C,

25 s at 72°C for 40 cycles. Melting curve analysis was performed at 65–85°C/step 0.5°C. The qPCR

data were analyzed using double delta Ct analysis

Gene expression analysis

Gene expression analyzes were performed on several types of samples: six whole flies, six thoraxes,

six fat bodies, or 20,000 isolated macrophages. Macrophages were isolated by a cell sorter (S3e Cell

Sorter, BioRad) as described in the section Isolation of macrophages, while dissections were made in

PBS, transferred to TRIzol Reagent (Invitrogen) and homogenized using a DEPC-treated pestle.

Subsequently, RNA was extracted by TRIzol Reagent (Invitrogen) according to the manufacturer's

protocol. Superscript III Reverse Transcriptase (Invitrogen) primed by oligo(dT)20 primer was used for

reverse transcription. Relative expression rates for particular genes were quantified on a 384CFX

1000 Touch Real-Time Cycler (BioRad) using the TP 2x SYBR Master Mix (Top-Bio) in three technical

replicates with the following protocol: initial denaturation - 3 min at 95°C, amplification – 15 s at

94°C, 20 s at 56°C, 25 s at 72°C for 40 cycles. Melting curve analysis was performed at 65–85°C/step

0.5°C. The primer sequences are listed in the Key Resources Table. The qPCR data were analyzed

using double delta Ct analysis, and the expressions or specific genes were normalized to the

expression of Ribosomal protein 49 (Rp49) in the corresponding sample. The relative values (fold

change) to control are shown in the graphs. Samples for gene expression analysis were collected

from three independent experiments.

Metabolite measurement

To measure metabolite concentration, isolated macrophages, whole flies or hemolymph were used.

Hemolymph was isolated from 25 adult males by centrifugation (14,000 RPM, 5 min) through a

silicagel filter into 50 μL of PBS. For measurement of metabolites from whole flies, five flies were

homogenized in 200 μL of PBS and centrifuged (3 min, 4°C, 8,000 RPM) to discard insoluble debris.

50,000 macrophages were isolated by cell sorter (S3e Cell Sorter, BioRad) as described in the section

Isolation of macrophages. Half of all samples were used for the quantification of proteins. Samples

for glucose, glycogen, trehalose, and triglyceride measurement were denatured at 75°C for 10 min,

whereas samples for protein quantification were frozen immediately in -80°C. Glucose was measured

using a Glucose (GO) Assay (GAGO-20) Kit (Sigma) according to the manufacturer's protocol.

Colorimetric reaction was measured at 540 nm. For glycogen quantification, sample was mixed with

amyloglucosidase (Sigma) and incubated at 37°C for 30 min. A Bicinchoninic Acid Assay (BCA) Kit

(Sigma) was used for protein quantification according to the supplier's protocol and the absorbance

was measured at 595 nm. Cholesterol and cholesteryl esters were measured on isolated lipid fraction

by using Cholesterol/Cholesteryl Ester Quantitation Kit (Sigma) according to the supplier's protocol.

Triglycerides were measured using Triglyceride quantification Colorimetric/Fluorometric Kit (Sigma).

For trehalose quantification, sample was mixed with trehalase (Sigma) and incubated at 37°C for 30

min. Samples for metabolite concentration were collected from three independent experiments.

Staining of lipid droplets

Flies were dissected in Grace's Insect Medium (Sigma) and subsequently stained with DAPI and Cell

Brite Fix Membrane Stain 488 (Biotium) diluted with Grace's Insect Medium according to the

manufacturer's protocol at 37°C. Tissues were washed in PBS and then fixed with 4% PFA

(Polysciences). After 20 min, the tissues were washed in PBS and pre-washed with 60% isopropanol.

Dissected abdomens were then stained with OilRedO dissolved in 60% isopropanol for 10 min. The

tissues were then washed with 60% isopropanol and mounted in an Aqua Polymount (Polysciences).

Tissues were imaged using an Olympus FluoView 3000 confocal microscope (Olympus). Content of

lipids in adipose tissue and size of lipid droplets were analyzed using Fiji software. Flies for the

analysis of lipid droplets in the fat body were collected from three independent experiments and

representative images are shown.

Lipidomic analysis

Adipose tissues from six flies from each group were dissected in ice-cold PBS for lipidomics analysis.

The removed tissues were stored in PBS buffer in Eppendorf tubes at − 80 °C. Immediately after

thawing, they were extracted by 500 µl of cold chloroform: methanol solution (v/v; 1:1). The samples

were then homogenized by a Tissue Lyser II (Qiagen, Prague, Czech Republic) at 50 Hz, -18°C for 5

min and kept further in an ultrasonic bath (0°C, 5 min). Further, the mixture was centrifuged at

10,000 RPM at 4°C for 10 min followed by the removal of the supernatant. The extraction step was

repeated at the same conditions. The lower layer of pooled supernatant was evaporated to dryness

under a gentle stream of Argon. The dry total lipid extract was re-dissolved in 500 µl of chloroform:

methanol solution (v/v; 1:1) and directly measured using previously described methods (Bayley et al.

2020). Briefly, high performance liquid chromatography (Accela 600 pump, Accela AS autosampler)

combined with mass spectrometry LTQ-XL (all Thermo Fisher Scientific, San Jose, CA, USA) were used.

The chromatographic conditions were as follows: Injection volume 5 µl; column Gemini 3 µM C18

HPLC column (150 × 2 mm ID, Phenomenex, Torrance, CA, USA) at 35°C; the mobile phase (A) 5 mM

ammonium acetate in methanol with ammonia (0.025%), (B) water and (C) isopropanol: MeOH (8:2);

gradient change of A:B:C as follows: 0 min: 92:8:0, 7 min: 97:3:0, 12 min: 100:0:0, 19 min: 93:0:7, 20-

29 min: 90:0:10, 40-45 min: 40:0:60, 48 min: 100:0:0, and 50-65 min: 92:8:0 with flow rate 200

µl/min. The mass spectrometry condition: positive (3 kV) and negative (-2,5 kV) ion detection mode;

capillary temperature 200°C. Eluted ions were detected with full scan mode from 200 to 1000 Da

with the collisionally induced MS2 fragmentation (NCE 35). Data were acquired and processed by

means of XCalibur 4.0 software (Thermo Fisher). The corrected areas under individual analytical

peaks were expressed in percentages assuming that the total area of all detected is 100%.

Autophagy visualization

To visualize autophagy in adipose tissue, Atg8a-mCherry-bearing flies were dissected in PBS, fixed

with 4% PFA, and then washed with PBS and stained with DAPI. Fat bodies were imaged using an

Olympus FluoView 3000 confocal microscope (Olympus).

Immunostaining

Flies were dissected in ice-cold PBS and fixed with 4% PFA in PBS (Polysciences) for 20 minutes. After

three washes in PBS-Tween (0.1%), nonspecific binding was blocked by 10% NGS in PBS for 1 hour at

RT. Tissues were then incubated with primary antibodis (for NimC1: Mouse anti-NimC1 antibody

P1a+b, 1:100, kindly provided by István Andó); for Foxo: Rabbit anti-Foxo, CosmoBio, 1:1,000; for

tGPH: Rabbit anti-GFP, ABfinity, 1:100) at 4°C overnight. After washing the unbound primary

antibody (three times for 10 min in PBS-Tween), secondary antibody was applied at a dilution of

1:250 for 2 hours at RT (Goat anti-Mouse IgG (H+L) Alexa 555, Invitogen or Goat anti-Rabbit IgG (H+L)

Cy2, Jackson-Immunoresearch). Nuclei were stained with DAPI. Tissues were mounted with Aqua

Polymount (Polysciences). Tissues were imaged using an Olympus FluoView 3000 confocal

microscope (Olympus) and images were reconstructed using Fiji software. Foxo localization was

detected by Plot-Profile analysis using Fiji software.

Isolation of macrophages

GFP-labeled macrophages were isolated from *Crq-Gal4, UAS-eGFP* male flies using fluorescence-

activated cell sorting (FACS). Approximately 200 flies were anaesthetized with CO_2, washed in PBS

and homogenized in 600 mL of PBS using a pestle. The homogenate was sieved through a nylon cell

strainer (40 μm). This strainer was then additionally washed with 200 μL of PBS, which was added to

the homogenate subsequently. The samples were centrifuged (3 min, 4°C, 3,500 RPM) and the

supernatant was washed with ice-cold PBS after each centrifugation (three times). Prior to sorting,

samples were transferred to FACS polystyrene tubes using a disposable bacterial filter (50 μm,

Sysmex) and macrophages were sorted into 100 μL of PBS using a S3^{TM} Cell Sorter (BioRad). Isolated

cells were verified by fluorescence microscopy and differential interference contrast.

Phagocytic activity

To analyze phagocytic rate, flies were infected with 20,000 of *S. pneumoniae* and after 24 h, they

were injected with 50 nl of pHrodo^{TM} Red *S. aureus* (Thermo Fischer Scientific). After 1 h, abdomens

of flies were dissected in PBS and then fixed for 20 min with 4% PFA. Aqua Polymount (Polysciences)

was used to mount the sample. Macrophages were imaged using an Olympus FluoView 3000

confocal microscope and red dots depicting phagocytic events were manually counted per cell.

Lipoprotein uptake

To analyze lipoprotein uptake by *Drosophila* macrophages, Crq>GFP flies were injected with 50 nl of

1x (corresponding to physiological concentration), 5x or 10x concentrated pHrodo^{TM} Red LDL

(Invitrogen) into the ventrolateral side of the abdomen using an Eppendorf Femtojet microinjector.

After 1 hour, the fly abdomens were opened in PBS and subsequently fixed for 20 min with 4% PFA in

PBS (Polysciences). Aqua Polymount (Polysciences) was used to mount the sample. Macrophages

were imaged using an Olympus FluoView 3000 confocal microscope.

Bacterial strain and fly injection

The *Streptococcus pneumoniae* strain EJ1 was stored at -80°C in Tryptic Soy Broth (TSB) media

containing 10% glycerol. For the experiments, bacteria were streaked onto agar plates containing 3%

TSB and 100 mg/mL streptomycin and subsequently incubated at 37°C in 5% CO_2 overnight. Single

colonies were inoculated into 3 mL of TSB liquid media with 100 mg/mL of streptomycin and 100,000

units of catalase and incubated at 37°C + 5% CO2 overnight. Bacterial density was measured after an

additional 4 h so that it reached an approximate 0.4 OD600. Final bacterial cultures were centrifuged

and dissolved in PBS so the final OD reached 2.4. The *S. pneumoniae* culture was kept on ice prior to

injection and during the injection itself. Seven-day-old males were anaesthetized with CO_2 and

injected with 50 nL culture containing 20,000 *S. pneumoniae* bacteria or 50 nL of mock buffer (PBS)

into the ventrolateral side of the abdomen using an Eppendorf Femtojet microinjector. The *Listeria*

monocytogenes strain 10403S was stored at -80°C in brain and heart infusion (BHI) broth containing

25% glycerol. For the experiments bacteria were streaked onto Luria Bertani (LB) agar plates

containing 100 μg/mL streptomycin and incubated at 37°C overnight; plates were stored at 4°C and

used for inoculation for a period of two weeks. Single colonies were used to inoculate 3 mL of BHI

and incubated overnight at 37°C without shaking to obtain a morning 600 nm optical density (OD600)

of approx. 0.4. Further, *L. monocytogenes* cultures were diluted to OD600 0.01 in phosphate

buffered saline (PBS) and stored on ice prior to loading into an injection needle. Approximately 1,000

Listeria per fly were injected.

Survival analysis

Streptococcus-injected flies were kept at 29°C in vials with approximately 30 individuals per vial and

were transferred to fresh food every other day. *Listeria*-injected flies were kept at 25°C. Dead flies

were counted daily. At least three independent experiments were performed and combined into a

single survival curve generated in Graphpad Prism software; individual experiments showed

comparable results. The average number of individuals was more than 500 for each genotype and

replicate.

Pathogen load measurement

Single flies were homogenized in PBS using a motorized plastic pestle in 1.5 mL tubes. Bacteria were

plated in spots onto LB (*L. monocytogenes*) or TSB (*S. pneumoniae*) agar plates containing

streptomycin in serial dilutions and incubated overnight at 37°C before manual counting. To

determine intracellular *L. monocytogenes* loads, flies were injected with 50 nL of gentamycin solution

(1 mg/mL in PBS) 3 h prior to fly homogenization. Pathogen loads of 16 flies were determined for

each genotype and treatment in each experiment; at least three independent infection experiments

were conducted and the results were combined into one graph (in all presented cases, individual

experiments showed comparable results).

THP-1 cell lines

THP-1 cells were cultured at 37°C, 5% CO_2, in RPMI-1640 medium (Sigma), supplemented with 2 mM

glutamine (Applichem), 2 g/L sodium bicarbonate (J&K Scientific), 10% FBS (Biosera) and 100 mg/L

streptomycin (Sigma). Prior to the experiment, cells were transferred to 24-well plates at 10^5

cells/well in four biological replicates. THP-1 cells were activated with phorbol-12-myristate-13-

acetate (200 ng/mL, MedChemExpress). After 24 hours, *S. pneumoniae* bacteria were added (MOI 50

bacteria/macrophage) and the plate was centrifuged briefly (2 min, 200xg). Following 6-h incubation,

the cells were washed with RPMI-1640 medium, and fresh RPMI-1640 supplemented with

gentamycin (0.1 mg/mL, Sigma) was added. After 1 h incubation, the medium was replaced with

RPMI-1640 supplemented with penicillin-streptomycin (1%, Biosera). After another 17 h, the cells

were harvested into TRIzol Reagent (Invitrogen) followed by RNA isolation.

Statistics

All data were analyzed using Graphpad Prism software; specific tests are listed for each graph in the

figure legend.

Key resources table

Reagent type (species) or resource	Designation	Source reference	Identifier	Additional information
Strain, strain background (*Streptococcus pneumoniae*)	*EJ1 strain*	Provided by David Schneider		Dilution 20,000 units
Chemical compound, drug	TRIzol Reagent	Invitrogen	Cat# 15-596-018	
Chemical compound, drug	Superscript III Reverse Transcriptase	Invitrogen	Cat# 18080044	
Chemical compound, drug	2x SYBR Master Mix	Top-Bio	Cat# T607	
Chemical compound, drug	Grace´s Insect Medium	Sigma	Cat# G9771-1L	
Chemical compound, drug	OilRedO	Sigma	Cat# O0625-25G	
Chemical compound, drug	DAPI	Sigma	Cat# D9542	
Chemical compound,	Cell Brite Fix Membrane Stain 488	Biotium	Cat# 30090-T	

Reagent type (species) or resource	Designation	Source reference	Identifier	Additional information
drug				
Chemical compound, drug	Alexa Fluor™ Plus 405 Phalloidin	Invitrogen	Cat# A30104	
Chemical compound, drug	Pro-A Drosophila CHIP Seq Kit	Chromatrap	Cat# 500274	
Chemical compound, drug	pHrodo™ Red S. aureus Bioparticles™ Conjugate for Phagocytosis	ThermoFisher	Cat# A10010	
Chemical compound, drug	pHrodo™ Red LDL	Invitrogen	Cat# L34356	
Chemical compound, drug	Phorbol-12-myristate-13-acetate	MedChemExpress	Cat# HY-18739	
Chemical compound, drug	RPMI-1640 Medium	Sigma	Cat# R7755	
Chemical compound, drug	Glutamine	Applichem	Cat# A3704	
Chemical compound, drug	Gentamycin	Sigma	Cat# 1405-41-0	
Chemical compound, drug	FBS	Biosera	Cat# FB-1001	
Chemical compound, drug	Penicilin-Streptomycin	Biosera	Cat# XC-A4122/100	
Antibody	Rabbit Anti-GFP	ABfinity	Cat# G10362	

Reagent type (species) or resource	Designation	Source reference	Identifier	Additional information
Antibody	Anti-NimC1 (P1a+b)	Kindly gifted by István Andó		
Antibody	Goat anti-Mouse IgG (H+L) Alexa 555	Invitrogen	A28180	
Antibody	Rabbit Drosophila anti-Foxo	CosmoBio	Cat# THU-A-DFoxo	
Antibody	Goat anti-Rabbit IgG (H+L) Cy2	Jackson-Immunoresearch	AB_2338021	
Commercial assay, kit	Glucose (GO) Assay Kit	Sigma	Cat# GAGO20-1KT	
Commercial assay, kit	Bicinchoninic Acid Assay Kit	Sigma	Cat# BCA1	
Commercial assay, kit	Cholesterol/Cholesteryl Ester Quantitation Kit	Sigma	Cat# 428901-1KIT	
Commercial assay, kit	Triglyceride quantification Colorimetric/Fluorometric Kit	Sigma	Cat# MAK266-1KT	
Chemical compound, drug	Trehalase from porcine kidney	Sigma	Cat# T8778-1UN	
Genetic reagent (*Drosophila melanogaster*)	*HmlG4G80: w^{1118}; HmlΔ-Gal4; P{tubPGal80ts}*	Cross made in our laboratory by Tomas Dolezal		
Genetic reagent (*Drosophila melanogaster*)	*w^{1118}; HmlΔ-Gal4 P{tubPGal80ts}/+; P{w[+mGT]=GT1}foxo^{BG01018} /UAS-ImpL2^{cds}*	Cross made in our laboratory by Tomas Dolezal		
Genetic reagent (*Drosophila melanogaster*)	*w^{1118}; HmlΔ-Gal4 P{tubPGal80ts}/+; tGPH-GFP/TRiP^{control}*	Cross made in our laboratory by Tomas Dolezal		
Genetic reagent (*Drosophila*	*w^{1118}; PBac{sima-*	Bloomington	BDSC:	

Reagent type (species) or resource	Designation	Source reference	Identifier	Additional information
melanogaster)	GFP.AC.FPTB}VK00037	Drosophila Stock Center	42672	
Genetic reagent (Drosophila melanogaster)	ImpL2^{RNAi}	Bloomington Drosophila Stock Center		
Genetic reagent (Drosophila melanogaster)	ImpL2-RA-Gal4	Provided by Hugo Stocker		
Genetic reagent (Drosophila melanogaster)	Atg8a-mCherry	Provided by Gabor Juhasz		
Genetic reagent (Drosophila melanogaster)	ImpL2^{cds}	Provided by Hugo Stocker		
Genetic reagent (Drosophila melanogaster)	20xUAS-6xmCherry	Bloomington Drosophila Stock Center		
Genetic reagent (Drosophila melanogaster)	CrqGal4>2xeGFP	Provided by Marc C. Dionne		
Genetic reagent (D. melanogaster)	Hif1α[RNAi]: P{KK110834}VIE-260B	Vienna Drosophila Resource Center	VDRC: v106504	FBst0478328
Genetic reagent (D. melanogaster)	TRiP ^{control}: y(1) v(1); P{y[+t7.7]=CaryP}attP2	Bloomington Drosophila Stock Center	BDSC: 36303	FBst0036303
Genetic reagent (D. melanogaster)	w: w^{1118}	Genetic background based on CantonS		
Sequence-based reagent	Foxo forward: CAATGCAAGAGATGGTCTCTC	KRD	CG3143	FBgn0038197
Sequence-based reagent	Foxo reverse: TGACCAGGTTCGTGTTGACC	KRD	CG3143	FBgn0038197
Sequence-based	4EBP Forward: CCATGATCACCAGGAAGGTTGTC	KRD	CG8846	FBgn0261560

Reagent type (species) or resource	Designation	Source reference	Identifier	Additional information
reagent	A			
Sequence-based reagent	4EBP Reverse: AGCCCGCTCGTAGATAAGTTTGGT	KRD	CG8846	FBgn0261560
Sequence-based reagent	EIF-4E Forward: AAGAAGAACATTCGCCCATG T	KRD	CG4035	FBgn0015218
Sequence-based reagent	EIF-4E Reverse: GGACTGCCGACGGAAACAAC	KRD	CG4035	FBgn0015218
Sequence-based reagent	Glut-CoA deh Forward: CTGATTGGTGCCTTTGGCC	KRD	CG9547	FBgn0031824
Sequence-based reagent	Glut-CoA deh Reverse: ATAGCGAGCGTTGTTCAGAC	KRD	CG9547	FBgn0031824
Sequence-based reagent	Lip4 Forward: CCACTTGGGTGATGATGGGAC	KRD	CG6113	FBgn0032264
Sequence-based reagent	Lip4 Reverse: ACGTATTGCCACGAACATTGG	KRD	CG6113	FBgn0032264
Sequence-based reagent	MTP Forward: GAATCGAAATGCCCAGACG	KRD	CG9342	FBgn0266369
Sequence-based reagent	MTP Reverse: AACGTTGGTTTGTTTGAGAAGC	KRD	CG9342	FBgn0266369
Sequence-based reagent	HSL Forward: CAGTCCTACGAGATTCACGG	KRD	CG11055	FBgn0034491
Sequence-based reagent	HSL Reverse: GGCTTCGTTGGATAACATTGTG	KRD	CG11055	FBgn0034491
Sequence-based reagent	Bmm Forward: CACCGCGCCGCAATGAATGTATA A	KRD	CG5295	FBgn0036449
Sequence-based reagent	Bmm Reverse: TTCAATCACTGTTTGTCGGTCGG C	KRD	CG5295	FBgn0036449
Sequence-based	GlyP Forward:	KRD	CG7254	FBgn0004507

Reagent type (species) or resource	Designation	Source reference	Identifier	Additional information
reagent	TCCACCCTGAGGGACTACTAC			
Sequence-based reagent	GlyP Reverse: GGTGTTGGTCAGTGAGCGAC	KRD	CG7254	FBgn0004507
Sequence-based reagent	GlyS Forward: CGATGAAGAAGTACAGGGTC	KRD	CG6904	FBgn0266064
Sequence-based reagent	GlyS Reverse: TATCATCGTTATTGCCTGGAG	KRD	CG6904	FBgn0266064
Sequence-based reagent	apoLTP Forward: TGGGTGGATTGAAGCCACAG	KRD	CG15828	FBgn0032136
Sequence-based reagent	apoLTP Reverse: TGAGTGGATTTTCTCTCACTGC	KRD	CG15828	FBgn0032136
Sequence-based reagent	apoLPP Forward: TTGGAATCCTAGCTTCTGTGCT	KRD	CG11064	FBgn0087002
Sequence-based reagent	apoLPP Reverse: AGTCATAGTAGTTGCCGGGTAT	KRD	CG11064	FBgn0087002
Sequence-based reagent	Atg1 Forward: CTAAAGCCGTCGTCCAATGT	KRD	CG10967	FBgn0260945
Sequence-based reagent	Atg1 Reverse: GAACAGCATGCTCCGGTATT	KRD	CG10967	FBgn0260945
Sequence-based reagent	Atg6 Forward: CGGAGTTATCTTTGCCCATCT	KRD	CG5429	FBgn0264325
Sequence-based reagent	Atg6 Reverse: GTGCTACACGCTGTTGCTC	KRD	CG5429	FBgn0264325
Sequence-based reagent	Rp49 Forward: AAGCTGTCGCACAAATGGCG	KRD	CG7939	FBgn0002626
Sequence-based reagent	Rp49 Reverse: GCACGTTGTGCACCAGGAAC	KRD	CG7939	FBgn0002626
Sequence-based reagent	ImpL2 Forward: TTCGCGGTTTCTGGGCACCC	KRD	CG15009	FBgn0001257

Reagent type (species) or resource	Designation	Source reference	Identifier	Additional information
Sequence-based reagent	ImpL2 Reverse: GCGCGTCCGATCGTCGCATA	KRD	CG15009	FBgn0001257
Sequence-based reagent	ImpL2-RA Forward: GTGCCAACGAAGCTTCGAGTG	KRD	CG15009	FBgn0001257
Sequence-based reagent	ImpL2-RA Reverse: GCGTGGCTTCTCCTCCTCC	KRD	CG15009	FBgn0001257
Sequence-based reagent	ImpL2-RB, D Forward: GTCGTCGGAAAGGATACCGC	KRD	CG15009	FBgn0001257
Sequence-based reagent	ImpL2-RB, D Reverse: TTCCGTCGTCAATCCAATAGCC	KRD	CG15009	FBgn0001257
Sequence-based reagent	ImpL2-RC Forward: ATAGACTCGGAGACCTCCCC	KRD	CG15009	FBgn0001257
Sequence-based reagent	ImpL2-RC Reverse: GCGTGGCTTCTCCTCCTCC	KRD	CG15009	FBgn0001257
Sequence-based reagent	IGFBP7 Forward: GCCATCACCCAGGTCAGCAAG	KRD	GC04M057030	
Sequence-based reagent	IGFBP7 Reverse: GGATTCCGATGACCTCACAGCT	KRD	GC04M057030	
Sequence-based reagent	ACTB Forward: ATTGCCGACAGGATGCAGAA	KRD	GC07M005527	
Sequence-based reagent	ACTB Reverse: GCTGATCCACATCTGCTGGAA	KRD	GC07M005527	
Sequence-based reagent	IGFBP1 Forward: TCCTTTGGGACGCCATCAGTAC	KRD	GC07P046552	
Sequence-based reagent	IGFBP1 Reverse: GATGTCTCCTGTGCCTTGGCTA	KRD	GC07P046552	

Reagent type (species) or resource	Designation	Source reference	Identifier	Additional information
Sequence-based reagent	IGFBP2 Forward: CGAGGGCACTTGTGAGAAGCG	KRD	GC02P216632	
Sequence-based reagent	IGFBP2 Reverse: TGTTCATGGTGCTGTCCACGTG	KRD	GC02P216632	
Sequence-based reagent	IGFBP3 Forward: CGCTACAAAGTTGACTACGAGTC	KRD	GC07M045912	
Sequence-based reagent	IGFBP3 Reverse: GTCTTCCATTTCTCTACGGCAGG	KRD	GC07M045912	
Sequence-based reagent	IGFBP4 Forward: ACCCACGAGGACCTCTACATCA	KRD	GC17P040443	
Sequence-based reagent	IGFBP4 Reverse: CACACCAGCACTTGCCACGCT	KRD	GC17P040443	
Sequence-based reagent	IGFBP5 Forward: CGTGCTGTGTACCTGCCCAATT	KRD	GC02M216672	
Sequence-based reagent	IGFBP5 Reverse: ACTTGTCCACGCACCAGCAGAT	KRD	GC02M216672	
Sequence-based reagent	IGFBP6 Forward: CACAGGATGTGAACCGCAGAGA	KRD	GC12P053097	
Sequence-based reagent	IGFBP6 Reverse: CACTGAGTCCAGATGTCTACGG	KRD	GC12P053097	
Sequence-based reagent	Defensin Forward: GTTCTTCGTTCTCGTGG	KRD	CG1385	FBgn0010385
Sequence-based reagent	Defensin Reverse: CTTTGAACCCCTTGGC	KRD	CG1385	FBgn0010385
Sequence-based reagent	Metchnikowin Forward: AACTTAATCTTGGAGCGA	KRD	CG8175	FBgn0014865

Reagent type (species) or resource	Designation	Source reference	Identifier	Additional information
Sequence-based reagent	Metchnikowin Reverse: CGGTCTTGGTTGGTTAG	KRD	CG8175	FBgn0014865
Sequence-based reagent	Drosocin Forward: CCATCGTTTTCCTGCT	KRD	CG10816	FBgn0010388
Sequence-based reagent	Drosocin Reverse: CCATCGTTTTCCTGCT	KRD	CG10816	FBgn0010388
Sequence-based reagent	Reg 1 Forward: GAGCAAATTGGACTCTACAGG	KRD		
Sequence-based reagent	Reg 1 Reverse: GGGGAGCAACAAGTAACTCG	KRD		
Sequence-based reagent	Reg 2 Forward: CTTTGGGCTGATAATTCCGG	KRD		
Sequence-based reagent	Reg 2 Reverse: TACATATATCCATAGAACCACG	KRD		
Sequence-based reagent	Reg 5 Forward: AGCCATCCATCTATGTGCC	KRD		
Sequence-based reagent	Reg 5 Reverse: TCAAACGCCACGAGACGAC	KRD		
Sequence-based reagent	Reg 7 Forward: GCAACTCAAATTCTTCAAACTCG	KRD		
Sequence-based reagent	Reg 7 Reverse: TCGGACCACTTGCTTTGTGT	KRD		
Sequence-based reagent	Reg 9 Forward: GAACCGTCGCCTTCCAG	KRD		

Reagent type (species) or resource	Designation	Source reference	Identifier	Additional information
Sequence-based reagent	Reg 9 Reverse: TGCCATGCCATTTGTTTCGC	KRD		
Sequence-based reagent	Sam-S Forward: CAAATCAGCGACGCTATCTTGG	KRD	CG2674	FBgn0005278
Sequence-based reagent	Sam-S Reverse: TGTCTCACGAACAACCTTCTGG	KRD	CG2674	FBgn0005278
Software, algorithm	Graphpad Prism	https://www.graphpad.com/	Graphpad Prism	RRID:SCR_002798
Software, algorithm	Microsoft Excel	https://www.microsoft.com/	Microsoft Excel	
Software, algorithm	Fiji	ImageJ - https://fiji.sc	ImageJ	RRID:SCR_002285
Other	S3e Cell Sorter	BioRad -http://www.bio-rad.com/	BioRad	
Other	Olympus FluoView 3000	Olympus - https://www.olympus-global.com/	Olympus	RRID:SCR_017015 RRID:SCR_014215
Other	Olympus IX71	Olympus - https://www.olympus-global.com/	Olympus	

References

Alee. 2011. "ImpL2 Represses Insulin Signaling in Response to Hypoxia." Dissertation Thesis. University of Oregon, Eugene, Oregon, USA.

Akiel, Maaged et al. 2017. "IGFBP7 Deletion Promotes Hepatocellular Carcinoma." *Cancer Research* 77(15): 4014–25. http://cancerres.aacrjournals.org/lookup/doi/10.1158/0008-5472.CAN-16-2885.

Alic, Nazif, Matthew P. Hoddinott, Giovanna Vinti, and Linda Partridge. 2011. "Lifespan Extension by Increased Expression of the Drosophila Homologue of the IGFBP7 Tumour Suppressor." *Aging Cell* 10(1): 137–47. http://doi.wiley.com/10.1111/j.1474-9726.2010.00653.x.

Andrejeva, Gabriela, and Jeffrey C. Rathmell. 2017. "Similarities and Distinctions of Cancer and
Immune Metabolism in Inflammation and Tumors." *Cell Metabolism* 26(1): 49–70.
https://linkinghub.elsevier.com/retrieve/pii/S1550413117303467.

Arquier, Nathalie et al. 2006. "Analysis of the Hypoxia-Sensing Pathway in Drosophila Melanogaster."
Biochemical Journal 393(2): 471–80.
https://portlandpress.com/biochemj/article/393/2/471/78893/Analysis-of-the-hypoxiasensing-
pathway-in.

Aspichueta, Patricia et al. 2012. "Disrupted VLDL Features and Lipoprotein Metabolism in Sepsis." In
Dyslipidemia - From Prevention to Treatment, InTech.
http://www.intechopen.com/books/dyslipidemia-from-prevention-to-treatment/disrupted-
vldl-features-and-lipoprotein-metabolism-in-sepsis.

Bader, R. et al. 2013. "The IGFBP7 Homolog Imp-L2 Promotes Insulin Signaling in Distinct Neurons of
the Drosophila Brain." *Journal of Cell Science* 126(12): 2571–76.
http://jcs.biologists.org/cgi/doi/10.1242/jcs.120261.

Bajgar, Adam et al. 2015. "Extracellular Adenosine Mediates a Systemic Metabolic Switch during
Immune Response" ed. Marc S. Dionne. *PLOS Biology* 13(4): e1002135.
https://dx.plos.org/10.1371/journal.pbio.1002135.

Bajgar, Adam, and Tomas Dolezal. 2018. "Extracellular Adenosine Modulates Host-Pathogen
Interactions through Regulation of Systemic Metabolism during Immune Response in
Drosophila" ed. Brian P. Lazzaro. *PLOS Pathogens* 14(4): e1007022.
https://dx.plos.org/10.1371/journal.ppat.1007022.

Bayley, Jeppe Seamus et al. 2020. "Cold-Acclimation Increases Depolarization Resistance and
Tolerance in Muscle Fibers from a Chill-Susceptible Insect, Locusta Migratoria." *American
Journal of Physiology-Regulatory, Integrative and Comparative Physiology*: ajpregu.00068.2020.
https://journals.physiology.org/doi/10.1152/ajpregu.00068.2020.

Biswas, Subhra K., and Alberto Mantovani. 2012. "Orchestration of Metabolism by Macrophages."
Cell Metabolism 15(4): 432–37.
https://linkinghub.elsevier.com/retrieve/pii/S1550413112000150.

Van den Bossche, Jan, Luke A. O'Neill, and Deepthi Menon. 2017. "Macrophage Immunometabolism:
Where Are We (Going)?" *Trends in Immunology* 38(6): 395–406.
https://linkinghub.elsevier.com/retrieve/pii/S147149061730042X.

Bunker, Brandon D et al. 2015. "The Transcriptional Response to Tumorigenic Polarity Loss in
Drosophila." *eLife* 4. http://www.ncbi.nlm.nih.gov/pubmed/25719210.

Chen, Grischa Y., Daniel A. Pensinger, and John-Demian Sauer. 2017. "Listeria Monocytogenes
Cytosolic Metabolism Promotes Replication, Survival, and Evasion of Innate Immunity." *Cellular
Microbiology* 19(10): e12762. https://onlinelibrary.wiley.com/doi/10.1111/cmi.12762.

Chistiakov, Dimitry A. et al. 2017. "Mechanisms of Foam Cell Formation in Atherosclerosis." *Journal
of Molecular Medicine* 95(11): 1153–65. http://link.springer.com/10.1007/s00109-017-1575-8.

Clark, Rebecca I. et al. 2011. "Multiple TGF-β Superfamily Signals Modulate the Adult Drosophila
Immune Response." *Current Biology* 21(19): 1672–77.
https://linkinghub.elsevier.com/retrieve/pii/S0960982211009547.

Dev, R, E. Bruera, and S. Dalal. 2018. "Insulin Resistance and Body Composition in Cancer Patients."
Annals of Oncology 29: ii18–26.
https://linkinghub.elsevier.com/retrieve/pii/S0923753419316801.

795 Dionne, Marc S., Linh N. Pham, Mimi Shirasu-Hiza, and David S. Schneider. 2006. "Akt and Foxo
796   Dysregulation Contribute to Infection-Induced Wasting in Drosophila." *Current Biology* 16(20):
797   1977–85. https://linkinghub.elsevier.com/retrieve/pii/S0960982206020641.

798 Dolezal, Tomas et al. 2019. "Molecular Regulations of Metabolism during Immune Response in
799   Insects." *Insect Biochemistry and Molecular Biology* 109: 31–42.
800   https://linkinghub.elsevier.com/retrieve/pii/S0965174818304600.

801 Febbraio, Maria, Ella Guy, and Roy L. Silverstein. 2004. "Stem Cell Transplantation Reveals That
802   Absence of Macrophage CD36 Is Protective Against Atherosclerosis." *Arteriosclerosis,*
803   *Thrombosis, and Vascular Biology* 24(12): 2333–38.
804   https://www.ahajournals.org/doi/10.1161/01.ATV.0000148007.06370.68.

805 Figueroa-Clarevega, Alejandra, and David Bilder. 2015. "Malignant Drosophila Tumors Interrupt
806   Insulin Signaling to Induce Cachexia-like Wasting." *Developmental Cell* 33(1): 47–55.
807   https://linkinghub.elsevier.com/retrieve/pii/S1534580715001434.

808 Gilbert, Lawrence I., and Haruo Chino. 1974. "Transport of Lipids in Insects." *Journal of Lipid Research*
809   15(5): 439–56. https://linkinghub.elsevier.com/retrieve/pii/S002222752036764X.

810 Gunnerson, Kyle J. et al. 2016. "TIMP2•IGFBP7 Biomarker Panel Accurately Predicts Acute Kidney
811   Injury in High-Risk Surgical Patients." *Journal of Trauma and Acute Care Surgery* 80(2): 243–49.
812   http://journals.lww.com/01586154-201602000-00009.

813 H. Tomkin, Gerald. 2012. "LDL as a Cause of Atherosclerosis." *The Open Atherosclerosis & Thrombosis*
814   *Journal* 5(1): 13–21. http://benthamopen.com/ABSTRACT/TOATHERTJ-5-13.

815 Harris, H W, J E Gosnell, and Z L Kumwenda. 2000. "The Lipemia of Sepsis: Triglyceride-Rich
816   Lipoproteins as Agents of Innate Immunity." *Journal of endotoxin research* 6(6): 421–30.
817   http://www.ncbi.nlm.nih.gov/pubmed/11521066.

818 Honegger, Basil et al. 2008. "Imp-L2, a Putative Homolog of Vertebrate IGF-Binding Protein 7,
819   Counteracts Insulin Signaling in Drosophila and Is Essential for Starvation Resistance." *Journal of*
820   *Biology* 7(3): 10. http://jbiol.biomedcentral.com/articles/10.1186/jbiol72.

821 Kamps-Hughes, Nick, Jessica L. Preston, Melissa A. Randel, and Eric A. Johnson. 2015. "Genome-Wide
822   Identification of Hypoxia-Induced Enhancer Regions." *PeerJ* 3: e1527.
823   https://peerj.com/articles/1527.

824 Kelly, Beth, and Luke AJ O'Neill. 2015. "Metabolic Reprogramming in Macrophages and Dendritic
825   Cells in Innate Immunity." *Cell Research* 25(7): 771–84.
826   http://www.nature.com/articles/cr201568.

827 Khovidhunkit, Weerapan et al. 2004. "Thematic Review Series: The Pathogenesis of Atherosclerosis.
828   Effects of Infection and Inflammation on Lipid and Lipoprotein Metabolism Mechanisms and
829   Consequences to the Host." *Journal of Lipid Research* 45(7): 1169–96.
830   http://www.jlr.org/lookup/doi/10.1194/jlr.R300019-JLR200.

831 Krejčová, Gabriela et al. 2019. "Drosophila Macrophages Switch to Aerobic Glycolysis to Mount
832   Effective Antibacterial Defense." *eLife* 8. https://elifesciences.org/articles/50414.

833 Kwon, Young et al. 2015. "Systemic Organ Wasting Induced by Localized Expression of the Secreted
834   Insulin/IGF Antagonist ImpL2." *Developmental Cell* 33(1): 36–46.
835   https://linkinghub.elsevier.com/retrieve/pii/S1534580715001148.

836 Li, Yan et al. 2013. "HIF- and Non-HIF-Regulated Hypoxic Responses Require the Estrogen-Related
837   Receptor in Drosophila Melanogaster" ed. Eric Rulifson. *PLoS Genetics* 9(1): e1003230.

https://dx.plos.org/10.1371/journal.pgen.1003230.

Liu, Yi et al. 2015. "Serum IGFBP7 Levels Associate with Insulin Resistance and the Risk of Metabolic
Syndrome in a Chinese Population." *Scientific Reports* 5(1): 10227.
http://www.nature.com/articles/srep10227.

Louie, Alexander et al. 2016. "How Many Parameters Does It Take to Describe Disease Tolerance?"
ed. Andy P. Dobson. *PLOS Biology* 14(4): e1002435.
https://dx.plos.org/10.1371/journal.pbio.1002435.

Luong, Nancy et al. 2006. "Activated FOXO-Mediated Insulin Resistance Is Blocked by Reduction of
TOR Activity." *Cell Metabolism* 4(2): 133–42.
https://linkinghub.elsevier.com/retrieve/pii/S1550413106002348.

Martínez-Castillo, Moisés et al. 2020. "Differential Production of Insulin-like Growth Factor-Binding
Proteins in Liver Fibrosis Progression." *Molecular and Cellular Biochemistry* 469(1–2): 65–75.
http://link.springer.com/10.1007/s11010-020-03728-4.

Mohanraj L, Kim H-S, Li W, Cai Q, Kim KE, Shin H-J, et al. 2013. IGFBP-3 Inhibits Cytokine-Induced
Insulin Resistance and Early Manifestations of Atherosclerosis. *PLoS ONE* 8(1): e55084.
https://doi.org/10.1371/journal.pone.0055084

Molaei, Maral, Crissie Vandehoef, and Jason Karpac. 2019. "NF-ΚB Shapes Metabolic Adaptation by
Attenuating Foxo-Mediated Lipolysis in Drosophila." *Developmental Cell* 49(5): 802-810.e6.
https://linkinghub.elsevier.com/retrieve/pii/S1534580719302795.

Morgantini, Cecilia et al. 2019. "Liver Macrophages Regulate Systemic Metabolism through Non-
Inflammatory Factors." *Nature Metabolism* 1(4): 445–59.
http://www.nature.com/articles/s42255-019-0044-9.

Nagao, Ayako et al. 2019. "HIF-1-Dependent Reprogramming of Glucose Metabolic Pathway of
Cancer Cells and Its Therapeutic Significance." *International Journal of Molecular Sciences* 20(2):
238. https://www.mdpi.com/1422-0067/20/2/238.

Newsholme, P, R Curi, S Gordon, and E A Newsholme. 1986. "Metabolism of Glucose, Glutamine,
Long-Chain Fatty Acids and Ketone Bodies by Murine Macrophages." *Biochemical Journal*
239(1): 121–25. https://portlandpress.com/biochemj/article/239/1/121/21648/Metabolism-of-
glucose-glutamine-longchain-fatty.

Nicholson, David, and Lindsay B. Nicholson. 2008. "A Simple Immune System Simulation Reveals
Optimal Movement and Cell Density Parameters for Successful Target Clearance." *Immunology*
123(4): 519–27. http://doi.wiley.com/10.1111/j.1365-2567.2007.02721.x.

Odegaard, J. I., and A. Chawla. 2013. "Pleiotropic Actions of Insulin Resistance and Inflammation in
Metabolic Homeostasis." *Science* 339(6116): 172–77.
https://www.sciencemag.org/lookup/doi/10.1126/science.1230721.

Oh, Youngman et al. 1996. "Synthesis and Characterization of Insulin-like Growth Factor-Binding
Protein (IGFBP)-7." *Journal of Biological Chemistry* 271(48): 30322–25.
http://www.jbc.org/lookup/doi/10.1074/jbc.271.48.30322.

Okamoto, N. et al. 2013. "A Secreted Decoy of InR Antagonizes Insulin/IGF Signaling to Restrict Body
Growth in Drosophila." *Genes & Development* 27(1): 87–97.
http://genesdev.cshlp.org/cgi/doi/10.1101/gad.204479.112.

Owusu-Ansah, Edward, Wei Song, and Norbert Perrimon. 2013. "Muscle Mitohormesis Promotes
Longevity via Systemic Repression of Insulin Signaling." *Cell* 155(3): 699–712.
https://linkinghub.elsevier.com/retrieve/pii/S0092867413011586.

Péan, Claire B. et al. 2017. "Regulation of Phagocyte Triglyceride by a STAT-ATG2 Pathway Controls
Mycobacterial Infection." *Nature Communications* 8(1): 14642.
http://www.nature.com/articles/ncomms14642.

Podrez, Eugene A. et al. 2000. "Macrophage Scavenger Receptor CD36 Is the Major Receptor for LDL
Modified by Monocyte-Generated Reactive Nitrogen Species." *Journal of Clinical Investigation*
105(8): 1095–1108. http://www.jci.org/articles/view/8574.

Remmerie, Anneleen, and Charlotte L. Scott. 2018. "Macrophages and Lipid Metabolism." *Cellular*
Immunology 330: 27–42. https://linkinghub.elsevier.com/retrieve/pii/S0008874918300327.

Ruan, Wenjing et al. 2017. "Serum Levels of IGFBP7 Are Elevated during Acute Exacerbation in COPD
Patients." *International Journal of Chronic Obstructive Pulmonary Disease* Volume 12: 1775–80.
https://www.dovepress.com/serum-levels-of-igfbp7-are-elevated-during-acute-exacerbation-
in-copd--peer-reviewed-article-COPD.

Shattuck-Heidorn, Heather et al. 2016. "Energetics and the Immune System: Trade-Offs Associated
with Non-Acute Levels of CRP in Adolescent Gambian Girls." *Evolution, Medicine, and Public*
Health: eow034. https://academic.oup.com/emph/article-lookup/doi/10.1093/emph/eow034.

Soeters, Maarten R., and Peter B. Soeters. 2012. "The Evolutionary Benefit of Insulin Resistance."
Clinical Nutrition 31(6): 1002–7.
https://linkinghub.elsevier.com/retrieve/pii/S0261561412001112.

Straub, Rainer H. 2014. "Insulin Resistance, Selfish Brain, and Selfish Immune System: An
Evolutionarily Positively Selected Program Used in Chronic Inflammatory Diseases." *Arthritis*
Research & Therapy 16(Suppl 2): S4. http://arthritis-
research.biomedcentral.com/articles/10.1186/ar4688.

Teng, Ooiean, Candice Ke En Ang, and Xue Li Guan. 2017. "Macrophage–Bacteria Interactions—A
Lipid-Centric Relationship." *Frontiers in Immunology* 8.
http://journal.frontiersin.org/article/10.3389/fimmu.2017.01836/full.

Warburg, Otto. 1925. "Über Den Stoffwechsel Der Carcinomzelle." *Klinische Wochenschrift* 4(12):
534–36. http://link.springer.com/10.1007/BF01726151.

Zmora, Niv, Stavros Bashiardes, Maayan Levy, and Eran Elinav. 2017. "The Role of the Immune
System in Metabolic Health and Disease." *Cell Metabolism* 25(3): 506–21.
https://linkinghub.elsevier.com/retrieve/pii/S1550413117300967.

 Figure legends

 Figure 1. **Streptococcal infection increases ImpL2 expression in a Hif1α-dependent manner in**

 activated macrophages

 (**A**) Expression of ImpL2 mRNA at whole-body level in flies (CrqGal4>GFP) infected with *S.*

 pneumoniae (S.p. infected) and control flies (Buff. injected) at various time points; results ompared

 by unpaired t test with Holm-Sidak method for multiple comparisons. (**B**) Representative confocal

 microscopy images of control (left) and infected (right) ImpL2-RA>mCherry individuals imaged at 24

 hpi, from Z stack of 7 layers, autofluorescence in the green channel was used to visualize the fly's

 body (see Figure 1-figure supplement 5 for images in color blind friendly pallete). (**C**) Number of

 ImpL2-positive macrophages in the thorax of control and infected adult flies (ImpL2-RA>mCherry) at

 24 hpi; results compared by unpaired t test. (**D**) Confocal microscopy image of infected and

 dissected ImpL2-RA>mCherry bearing fly stained with anti-NimC1 antibody (green) depicting the

 expression of ImpL2 in hemocytes, from Z stack of 11 layers. The scale bar represents 10 μm. (**E**)

 Confocal microscopy image of ImpL2-RA>mCherry hemocyte actively interacting with GFP-labeled *S.*

 pneumoniae (S.p.-GFP) *ex vivo.* The scale bar represents 13 μm. (**F**) Expression of ImpL2 mRNA (using

 primers targeting all transcript variants) in hemocytes isolated from Buff. injected and S.p. infected

 flies with macrophage-specific ImpL2 knockdown (ImpL2^{RNAi}), overexpression (ImpL2^{CDS}) and their

 respective controls (TRiP^{control}, w^{1118}) at 24 hpi (right) and gene expression of ImpL2-RA transcript in

 hemocytes isolated from Buff. injected and S.p. infected flies with hemocyte-specific ImpL2

 knockdown (ImpL2^{RNAi}) and its respective control (TRiP^{control}) at 24 hpi (left), documenting the

 efficiency of RNAi fly line used in this manuscript. Results compared by 2way ANOVA Tukey's multiple

 comparisons test. (**G**) Gene expression of ImpL2-RA (left) and all ImpL2 transcript variants (right) in

 hemocytes isolated from Buff. injected and S.p. infected control (HmlGal4>GFP) flies and flies with

 hemocyte-specific Hif1α knockdown at 24 hpi. Results compared by 2way ANOVA Tukey's multiple

comparisons test. (**H**) Relative amount of selected ImpL2 genomic regions (visualized in I) bound by

the transcription factor Hif1α in Buff. injected ans S.p. infected flies. Selected genomic region of S-

adenosylmethionin synthetase was used as a negative control. Results compared by 2way ANOVA

Tukey's multiple comparisons test. (**I**) *In silico* analysis of hypoxic enhancer activity by 500 bp bins at

the ImpL2 locus and visualization of ImpL2 transcript variants (Flybase.org). Each black dot plots a

log2 fold change (y-axis) of the difference in the randomer tag counts mapped to the 500-bp bin

between normoxic and hypoxic conditions. The triangles show the position and negative log10 p-

value (y-axis) of multiple hypoxic (Hif1α), immune (Rel) and stress (HSF) response transcription factor

binding sites. The close up shows distribution and clustering of individual response elements for the

most significant bin. In **A, F** and **G**, expression levels normalized against rp49 are reported as fold

change relative to levels of ImpL2 (using primers targeting all transcript variants) and ImpL2-RA

respectively, in Buff. injected controls, which were arbitrarily set to 1. The individual dots represent

biological replicates with line/bar showing mean ± SD, asterisks mark statistically significant

differences (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$). Hif1α, hypoxia-inducible factor 1 α; Rel, relish; HSF,

heat shock factor.

Figure 1-figure supplement 1. **ImpL2-RA>mCherry-positive cells display hemocyte morphology and**

actively phagocytose *Staphylococcus aureus*

Repesentative confocal microscopy image of ImpL2-RA>mCherry hemocyte depicting its phagocytic

ability visualized by *S. aureus*-pHrodo (green). The scale bar represents 10 μm.

Figure 1-figure supplement 2. **Gene expression of ImpL2 transcipt variants in distinct tissues**

Gene expression of ImpL2-RA (left), ImpL2-RB and ImpL2-RD (middle) and ImpL2-RC (right) transcript

variants in either hemocytes, fat body or thoracic muscles dissected from control and infected

Crq>GFP flies at 24 hpi. Expression levels normalized against rp49 are presented as a fold change

relative to levels of ImpL2-RA in hemocytes of Buff. injected control flies, which were arbitrarily set to

1. Individual dots represent biological replicates. Results were compared by unpaired t test with

Holm-Sidak method for multiple comparisons. Values are mean ± SD, asterisks mark statistically

significant differences (*p<0.05; **p<0.01; ***p<0.001).

Figure 1-figure supplement 3. **Efficiency of genetic manipulation of Hif1α**

Gene expression of Hif1α in hemocytes of Buff. injected and S.p. infected flies with or without

hemocyte-specific Hif1α knockdown. Expression levels normalized against rp49 are presented as a

fold change relative to levels of Hif1α in Buff. injected HmlGal>GFP flies, which were arbitrarily set to

1. Individual dots represent biological replicates. Results compared by 2way ANOVA Tukey's multiple

comparisons test. Values are mean ± SD, asterisks mark statistically significant differences (*p<0.05;

**p<0.01; ***p<0.001).

Figure 1-figure supplement 4. **ImpL2-RA regulatory sequence with transcription factor binding sites**

500-bp long sequence upstream of ImpL2-RA transcriptional start site (as shown on map) with

marked binding sites for Hif1α, Rel and HSF1. Hif1α, hypoxia-inducible factor 1 α; Rel, relish; HSF1,

heat shock factor 1.

Figure 1-figure supplement 5. **Copy of Figure 1B and 1D in color blind friendly pallete.**

(**A, B**) Representative confocal microscopy images of control (left) and infected (right) ImpL2-

RA>mCherry individuals imaged at 24 hpi, from Z stack of 7 layers, autofluorescence in the green

channel was used to visualize the fly's body. (**C**) Confocal microscopy image of infected and dissected

ImpL2-RA>mCherry bearing fly stained with anti-NimC1 antibody (green) depicting the expression of

ImpL2 in hemocytes, from Z stack of 11 layers. The scale bar represents 10 μm.

Figure 2. **Macrophage-derived ImpL2 regulates changes in adipose tissue metabolism induced by**

infection

(**A**) Triglyceride concentration in Buff. injected and S.p. infected flies with macrophage-specific ImpL2

knockdown (ImpL2^{RNAi}), overexpression (ImpL2^{CDS}) and their respective controls (TRiP^{control}, w^{1118}) at

whole-body level at 24 hpi. (**B-D**) Average size (**B**) and number (**C**) of lipid droplets (values log10-

transformed) and percentage of area occupied (**D**) in the fat body of Buff. injected and S.p. infected

flies with macrophage-specific ImpL2 knockdown (ImpL2^{RNAi}), overexpression (ImpL2^{CDS}) and their

respective controls (TRiP^{control}, w^{1118}) at 24 hpi. (**E**) Representative confocal microscopy images of

dissected fat body of Buff. injected and S.p. infected flies with macrophage-specific ImpL2

knockdown (ImpL2^{RNAi}), overexpression (ImpL2^{CDS}) and their respective controls (TRiP^{control}, w^{1118}) at

24 hpi, stained with OilRedO (red), DAPI (cyan) and CellBrite (white). The scale bar represents 20 μm.

(**F**) Relative proportions of different lipid species in the fat body of Buff. injected and S.p. infected

flies with macrophage-specific ImpL2 knockdown (ImpL2^{RNAi}), overexpression (ImpL2^{CDS}) and their

respective controls (TRiP^{control}, w^{1118}) at 24 hpi. (**G**) Glycogen concentration in Buff. injected and S.p.

infected flies with -specific ImpL2 knockdown (ImpL2^{RNAi}), overexpression (ImpL2^{CDS}) and their

respective controls (TRiP^{control}, w^{1118}) at whole-body level at 24 hpi. Metabolite concentrations were

normalized to the amount of proteins in each sample. Values are mean ± SD, asterisks mark

statistically significant differences (*$p<0.05$; **$p<0.01$; ***$p<0.001$). Results in (**A-D** and **G**) were

compared by 2way ANOVA Tukey's multiple comparisons test. (**B-D**) were quantified from 8 confocal

microscopy images for each genotype and treatment. Lipidomic analysis in (**F**) was performed in

three biological replicates for each genotype and treatment. PC, phosphatidylcholine; PE,

phosphatidylethanolamine; TG, triglycerides; DG, diglycerides; PG, phosphatidylglycerol; PI,

phosphatidylinositol; LysoPC, Lyso-phosphatidylcholine; LysoPE, Lyso-phosphatidylethanolamine.

1012

Figure 3. **Infection-induced autophagy and changes in gene expression in adipose tissue**

(**A**) Representative confocal microscopy images of fat body of control and infected flies with Atg8a-

mCherry autophagy reporter (Atg8a, red; DAPI, cyan). The scale bar represents 20 μm. (**B**) Graphical

representation of log2-fold change in mRNA expression of genes involved in insulin signaling,

lipoprotein and lipid metabolism, glycogenolysis and autophagy in the fat body of Buff.-injected and

S.p.-infected flies with macrophage-specific ImpL2 knockdown (ImpL2^{RNAi}), overexpression (ImpL2^{CDS})

and their respective controls (TRiP^{control}, w^{1118}) at 24 hpi. The first column (Infection) shows the log2-

fold change in mRNA expression in S.p-infected flies compared to Buff.-injected control flies (average

change for TRiP^{control} and w^{1118}). The second to fourth columns represent the log2-fold change in

mRNA expression compared to the corresponding control for knockdown (ImpL^{RNAi} to TRiP^{control}) and

overexpression (ImpL2^{CDS} to w^{1118}) in either uninfected (Buff.) or S.p. infected (S.p.) flies. Foxo,

forkhead box O; 4EBP, Thor; EIF-4E1, eukaryotic translation initiation factor 4E1; MTP, microsomal

triacylglycerol transfer protein; apoLPP, apolipophorin; apoLTP, apolipoprotein lipid transfer particle;

Bmm, brummer; Lip4, lipase 4; Hsl, hormone sensitive lipase; Glutaryl-CoA deh., Glutaryl-CoA

dehydrogenase; GlyS, glycogen synthase; GlyP, glycogen phosphorylase; Atg1, autophagy-related

gene 1; Atg6, autophagy-related gene 6.

1029

Figure 3-figure supplement 1. **Expression levels of genes regulating metabolism in adipose tissue**

Gene expression of *Foxo, 4EBP, EIF4E1, Glutaryl-CoA deh., Lip4, HSL, Bmm, GlyS, GlyP, apoLTP,*

apoLPP, MTP, Atg1, and *Atg6* genes in adipose tissue of Buff. injected (black bars) and S.p. infected

(grey bars) flies with macrophage-specific ImpL2 knockdown, overexpression and their respective

controls; all genotypes depicted below the x-axis were crossed with *Hml>Gal4; TubGal80^{TS}*.

Expression levels normalized against rp49 are presented as a fold-change relative to the levels in the

Buff. injected TRiP^{control}, which was arbitrarily set to 1. The individual dots represent biological

replicates. Results were compared by 2way ANOVA Tukey's multiple comparisons test. Bars show

mean ± SD, asterisks mark statistically significant differences (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$).

1039

Figure 4. **Macrophage-derived ImpL2 affects Foxo activity in adipose tissue**

Representative confocal microscopy images of Foxo immunolocalization in the fat body of Buff.

injected and S.p. infected flies with macrophage-specific ImpL2 knockdown (ImpL2^{RNAi}),

overexpression (ImpL2^{CDS}) and their respective controls (TRiP^{control}, w^{1118}) at 24 hpi; anti-dFoxo

antibody, green; DAPI, cyan. Histograms of Foxo cellular localization represent Foxo and DAPI signal

intensity in sections indicated by a white freehand line. The scale bar represents 20 μm.

1046

Figure 5. **Macrophage-derived ImpL2 affects adipose tissue insulin sensitivity**

(**A**) Representative confocal images of dissected adipose tissue of control flies (w^{1118}), flies with

macrophage-specific ImpL2 overexpression (ImpL2^{CDS}), and flies with macrophage-specific ImpL2

overexpression (ImpL2^{CDS}) and ubiquitous expression of hypomorphic variant of Foxo (foxoBG01018)

stained by OilRedO (red), CellBrite (white) and DAPI (cyan) at 24 hpi. The scale bar represents 20 μm.

(**B**) Representative confocal images of localization of the PI3K reporter tGPH in adipose tissue of Buff.

injected and S.p. infected control flies (w^{1118}) and flies with specific ImpL2 overexpression (ImpL2^{CDS})

at 24 hpi. tGPH, green; DAPI, cyan. The scale bar represents 20 μm.

1055

Figure 6. **Macrophage-derived ImpL2 increases circulating carbohydrates and lipids to be available**

for activated macrophages

(**A**) Concentrations of circulating glucose, trehalose, glycerides, and free fatty acids in the

hemolymph of Buff. injected and S.p. infected flies with macrophage-specific ImpL2 knockdown

(ImpL2^{RNAi}), overexpression (ImpL2^{CDS}) and their respective controls (TRiP^{control}, w^{1118}) at 24 hpi. (**B**)

Concentrations of glucose, trehalose, glycerides, and free fatty acids (**H**) in macrophages of Buff.

injected and S.p. infected flies with macrophage-specific ImpL2 knockdown (ImpL2^{RNAi}),

overexpression (ImpL2^{CDS}), and their respective controls (TRiP^{control}, w^{1118}) at 24 hpi. (**C**)

Representative confocal microscopy image of macrophages (Crq>GFP) containing lipid droplets;

neutral lipids were stained with OilRedO (red). The scale bar represents 10 μm. The image represents

a Z-stack consisting of a maximum projection of 5 layers. (**D**) Proportional occurrence of

macrophages containing at least one lipid droplet (stained by OilRedO) at 24 hpi, data combined

from fifty Crq>GFP positive cells. (**E**) Representative confocal microscopy images depicting the ability

of macrophage (Crq>GFP) to endocytose lipoproteins (human LDL-pHrodo, red). The scale bar

represents 10 μm. (**F**) Number of endocytic events in macrophages of flies, which were injected with

various concentrations (1x – dose corresponding to physiological concentration, 5x, and 10x) of

lipoproteins (human LDL-pHrodo). The results are combined from three biological replicates. (**A-B**)

Metabolite concentrations were normalized to the amount of proteins in each sample. Carbohydrate,

resp. lipid measurements were performed in eight, resp. four replicates represented by individual

dots. Results in (**A,B** and **D**) were compared by 2way ANOVA Tukey's multiple comparisons test and

in (**F**) one-way ANOVA Tukey's multiple comparisons test. Bars/lines show means ± SD, asterisks mark

statistically significant differences (*p<0.05; **p<0.01; ***p<0.001).

1078

Figure 6-figure supplement 1. **Concentration of carbohydrates and lipids in control fly lines**

(**A-E**) Concentrations of circulating glucose (**A**), trehalose (**B**) and glycerides (**C**) in hemolymph and

triglycerides (**D**) and glycogen (**E**) at a whole body level of Buff. injected and S.p. infected Hml-Gal4

TubGal80^{TS} x TRiP^{control}, Hml-Gal4 TubGal80^{TS} x w^{1118}, ImpL2^{RNAi} x TRiP^{control}, and ImpL2^{CDS} x w^{1118}

control flies at 24 hpi. Metabolite concentrations were normalized to the amount of proteins in each

sample. Results were compared by 2way ANOVA Tukey's multiple comparisons test. Bars show

means ± SD, asterisks mark statistically significant differences (*p<0.05; **p<0.01; ***p<0.001).

1086

Figure 7. **Macrophage-derived ImpL2 is important for resistance to bacterial infection**

(**A, C**) Survival rate of *S. pneumoniae* (S.p.) infected and Buff. injected flies with macrophage-specific

ImpL2 knockdown (ImpL2^{RNAi}) (**A**), overexpression (ImpL2^{CDS}) (**C**) and their respective controls

(TRiP^{control}, w^{1118}). Three independent experiments were performed and combined into one survival

curve; the number of individuals per replicate was at least 500 for each genotype. (**B, D**) Pathogen

load in S.p. infected and Buff. injected flies with knockdown (ImpL2^{RNAi}) (**B**), overexpression

(ImpL2^{CDS}) (**D**) and their respective controls (TRiP^{control}, w^{1118}) at 24 and 48 hpi. The individual dots in

the plot represent the number of bacteria (colony forming units-CFUs) in one fly. The data show

results combined from three independent biological replicates. (**E**) Confocal microscopy image

depicting the ability of a macrophage (Crq>GFP) to phagocytose an invading pathogen (*S. aureus*-

pHrodo-red). The scale bar represents 10 μm. The image represents a Z-stack consisting of a

maximum projection of 5 layers. (**F**) Phagocytic rate calculated for flies with macrophage-specific

ImpL2 knockdown (ImpL2^{RNAi}), overexpression (ImpL2^{CDS}), and their respective controls (TRiP^{control},

w^{1118}); each dot in the plot represents the log10-transformed number of phagocytic events per

macrophage (Crq>GFP); results are combined from three independent experiments. Survival data (**A**

and **C**) were analyzed by Log-rank and Grehan-Breslow Wilcoxon tests, pathogen load (**B** and **D**) by

Mann-Whitney test and phagocytic capability (**F**) by one-way ANOVA Tukey's multiple comparisons

test. Lines are means ± SD, asterisks mark statistically significant differences (*$p<0.05$; **$p<0.01$;

***$p<0.001$).

1106

Figure 7-figure supplement 1. **Survival of additional control genotypes of *S. pneumoniae* infection.**

Survival rate of S.p. infected and Buff. injected flies with macrophage-specific ImpL2 knockdown

(ImpL2^{RNAi}) (upper graph) and overexpression (ImpL2^{CDS}) (lower graph) and and their respective

controls (ImpL2^{RNAi} x TRiP^{control}, HmlG4G80 x TRiP^{control}, ImpL2^{cds} x w^{1118} and HmlG4G80 x w^{1118}). Three

independent experiments were performed and combined into one survival curve; the number of

individuals per replicate was at least 500 for each genotype. Survival data were analyzed by Log-rank

and Grehan-Breslow Wilcoxon tests.

1114

Figure 7-figure supplement 2. **Expression of antimicrobial peptides in hemocytes**

Gene expression of Drosocin (upper graph), Metchnikowin (middle graph) and Defensin (lower

graph) in macrophages of flies with macrophage-specific ImpL2 knockdown (ImpL2^{RNAi}),

overexpression (ImpL2^{CDS}), and their respective controls (TRiP^{control}, w^{1118}) at 24 hpi. Expression levels

normalized against rp49 are reported as fold change relative to levels of Drosocin, Metchnikowin,

and Defensin, respectively, in Buff. injected TRiP^{control}, which were arbitrarily set to 1.The individual

dots represent biological replicates. Results were compared by 2way ANOVA Tukey's multiple

comparisons test. Bars show means ± SD, asterisks mark statistically significant differences (*$p<0.05$;

**$p<0.01$; ***$p<0.001$).

1124

Figure 8. **ImpL2 enhances deleterious effects of chronic infection**

(**A, C**) Survival rate of *L. monocytogenes* (L.m.) infected and Buff. injected flies with macrophage-

specific ImpL2 overexpression (ImpL2^{CDS}) (**A**), knockdown (ImpL2^{RNAi}) (**C**), and their respective controls

(TRiP^{control}, w^{1118}). Three independent experiments were performed and combined into each survival

curve; the number of individuals per replicate was at least 500 for each genotype. (**B, D**) Pathogen

load of *L. monocytogenes* in flies with macrophage-specific ImpL2 overexpression (ImpL2^{CDS}) (B),

knockdown (ImpL2^{RNAi}) (**D**), (**E**) and their respective controls (TRiP^{control}, w^{1118}) shown as either total

load or intracellular *Listeria* subpopulation at 24 hpi and at 12 days post-infection. The individual dots

in the plot represent the number of bacteria (colony forming units-CFUs) in one fly. The data show

results combined from three independent biological replicates. Survival data (**A** and **C**) were analyzed

by Log-rank and Grehan-Breslow Wilcoxon tests and pathogen load (**B** and **D**) by Mann-Whitney test.

Values are mean ± SD, asterisks mark statistically significant differences (*$p<0.05$; **$p<0.01$;

***$p<0.001$).

1138

Figure 9. **IGFBP7, the mammalian homolog of ImpL2, is produced in human macrophages in**

response to *S. pneumoniae* in vitro

(**A**) Confocal microscopy images of PMA-activated control THP-1 cells (left) and PMA-activated THP-1

cells actively interacting with GFP-labeled *S. pneumoniae* (green) *in vitro* (right). The scale bar

represents 10 µm. The image on the right is a maximum projection of 3 layers. (**B**) Gene expression of

IGFBP7, IGFBP3 and IGFBP4 in PMA-activated THP-1 cells 24 hours after exposure to *S. pneumoniae*

in vitro. Expression levels normalized against ACTB are shown as a fold change relative to levels of

IGFBP7 in controls that were arbitrarily set to 1. Measured data were compared in Graphpad Prism

using 2way ANOVA Sidak's multiple comparison test. Values are mean ± SD, asterisks mark

statistically significant differences (*$p<0.05$; **$p<0.01$; ***$p<0.001$).

1149

Figure 9. **Schematic representation of the proposed model of ImpL2 function during infection**

Upon recognition of the bacteria, the macrophages switch to Hif1α-driven aerobic glycolysis. Hif1α

also induces the expression of the ImpL2-RA isoform by binding to hypoxia response elements in the

promotor of the *ImpL2* gene. Macrophage-derived ImpL2 then remotely activates Foxo re-

localization to the nucleus in the fat body, leading to the expression of metabolic Foxo-target genes.

Mobilized sources, manifested by hyperlipidemia and hyperglycemia in the hemolymph, are

subsequently used by activated macrophages. These transient changes in ImpL2-regulated

metabolism are essential for efficient bacterial killing and resistance to infection. However, if

prolonged, for example during chronic infection, macrophage-derived ImpL2 can harm the host by

potential wasting. Hif1α, hypoxia-inducible factor 1 α; HREs, hypoxia response elements; ImpL2,

imaginal morphogenesis protein-late 2; Foxo, forkhead box O.

1161

1162   **Source Data Files**

1163   All source data files are in Graphpad Prism format, which was used for all data processing. The free

1164   Graphpad viewer is available at www.graphpad.com

1165

Fig1.pzfx – Source data for Figure 1., Figure 1.-figure supplement 2. and Figure 1.-figure supplement

3.

Fig2 .pzfx – Source data for Figure 2.

Fig3.pzfx – Source data for Figure 3. and Figure 3.-figure supplement 1.

Fig6.pzfx – Source data for Figure 6.

Fig6suppl1.pzfx – Source data for Figure 6.-figure supplement 1.

Fig7 – Source Data 1.pzfx – Source data for Figure 7.

Fig7suppl1.pzfx – Source data for Figure 7.-figure supplement 1.

Fig6suppl2.pzfx – Source data for Figure 7.-figure supplement 2.

Fig8.pzfx – Source data for Figure 8.

Fig9.pzfx – Source data for Figure 9.

A ImpL2 expression in whole flies

B ImpL2-RA>mCherry Buff. injected | ImpL2-RA>mCherry S.p. infected

C Number of ImpL2-positive macrophages

D ImpL2-RA>mCherry NimC1 S.p. infected

E ImpL2-RA>mCherry S.p.-GFP phalloidin

F Expression of ImpL2 in macrophages

G ImpL2 expression upon Hif1α^RNAi in macrophages

H Chip-qPCR of Hif1α-ImpL2

I

A *Hml>Gal4 TubGal80^{TS}* crossed to:

w^{1118} | *ImpL2^{CDS}* | *foxo^{BG01018} ImpL2^{CDS}*

OilRedO
CellBrite
DARI

B

Hml>Gal4 TubGal80^{TS}; tGPH-GFP
crossed to:

Buff. injected | *S.p.* infected

tGPH DAPI

w^{1118}

ImpL2^{CDS}

A HEMOLYMPH

B HEMOCYTES

C CrqGal4>GFP OilRedO

D Percentage of OilRedO positive macrophages

E CrqGal4>GFP 1x LDL-pHrodo — CrqGal4>GFP 10x LDL-pHrodo

F LDL-pHrodo endocytosis

A ImpL2 knockdown - *S. pneumoniae* infection

Percent survival vs Days post infection

Hml>Gal4; TubGal80^TS x

— ImpL2^RNAi / S.p. infected
⋯ ImpL2^RNAi / Buff. injected
— TRiP^control / S.p. infected
⋯ TRiP^control / Buff. injected

***

B Pathogen load of *S. pneumoniae*

CFUs (in thousands) at 24 hpi and 48 hpi

Hml>Gal4; TubGal80^TS x

● TRiP^control
● ImpL2^RNAi

**

C ImpL2 overexpression - *S.pneumoniae* infection

Percent survival vs Days post infection

Hml>Gal4; TubGal80^TS x

— ImpL2^CDS / S.p. infected
⋯ ImpL2^CDS / Buff. injected
— w^1118 / S.p. infected
⋯ w^1118 / Buff. injected

***

D Pathogen load of *S. pneumoniae*

CFUs (in thousands) at 24 hpi and 48 hpi

Hml>Gal4; TubGal80^TS x

● w^1118
● ImpL2^CDS

***

E CrqGal4>GFP S.a.-pHrodo

F Phagocytic capability

No. of phagocytic events (log)

TRiP^control, ImpL2^RNAi, w^1118, ImpL2^CDS

x CrqGal4>GFP

***

A   ImpL2 overexpression - *L. monocytogenes* infection

Hml>Gal4; TubGal80^{TS} x

- ImpL2^{CDS} / L.m. infected
- ImpL2^{CDS} / Buff. injected
- W^{1118} / L.m. infected
- W^{1118} / Buff. injected

B   Pathogen load of *L. monocytogenes*

Hml>Gal4; TubGal80^{TS} x

- w^{1118}
- ImpL2^{CDS}

C   ImpL2 knockdown - *L. monocytogenes* infection

Hml>Gal4; TubGal80^{TS} x

- ImpL2^{RNAi} / L.m. infected
- ImpL2^{RNAi} / Buff. injected
- TRiP^{control} / L.m. infected
- TRiP^{control} / Buff. injected

D   Pathogen load of *L. monocytogenes*

Hml>Gal4; TubGal80^{TS} x

- TRiP^{control}
- ImpL2^{RNAi}

A

control

S.p. infected

THP-1 cells

DAPI S.p.-GFP phalloidin

B

Expression of IGFBPs in THP-1 cells