# Altered basal lipid metabolism underlies the functional impairment of naive CD8[+] T cells in elderly humans

**Short title:** Metabolic properties of naive CD8[+] T cells

Francesco Nicoli,[1,2,*] Mariela P. Cabral-Piccin,[1,10] Laura Papagno,[1,10] Eleonora Gallerani,[2] Victor Folcher,[1] Marion Dubois,[1] Emmanuel Clave,[3] Hélène Vallet,[1,4] Justin J. Frere,[5] Emma Gostick,[6] Sian Llewellyn-Lacey,[6] David A. Price,[6,7] Antoine Toubert,[3,8] Jacques Boddaert,[1,4] Antonella Caputo,[2] Riccardo Gavioli,[2] and Victor Appay[1,9,*]

[1] Centre d'Immunologie et des Maladies Infectieuses (CIMI-Paris), Sorbonne Université, INSERM U1135, 75013 Paris, France

[2] Department of Chemical and Pharmaceutical Sciences, University of Ferrara, Ferrara 44121, Italy

[4] Institut de Recherche Saint Louis, EMiLy, Université de Paris, INSERM U1160, 75010 Paris, France

[4] Service de Gériatrie, Hôpital Pitié-Salpêtrière, AP-HP, 75013 Paris, France

[5] Department of Immunobiology and the Arizona Center on Aging, University of Arizona College of Medicine Tucson, Tucson, AZ 85724, USA

[6] Division of Infection and Immunity, Cardiff University School of Medicine, Cardiff CF14 4XN, UK

[7] Systems Immunity Research Institute, Cardiff University School of Medicine, Cardiff CF14 4XN, UK

[8] Laboratoire d'Immunologie et d'Histocompatibilité, Hôpital Saint-Louis, AP-HP, 75010 Paris, France

[9] International Research Center of Medical Sciences, Kumamoto University, Kumamoto 860-0811, Japan

[10] These authors contributed equally

[*] Correspondence: nclfnc1@unife.it (F.N.) or victor.appay@upmc.fr (V.A.)

35 **ABSTRACT**
36

Aging is associated with functional deficits in the naive T cell compartment, which compromise

the generation of *de novo* immune responses against previously unencountered antigens. The

mechanisms that underlie this phenomenon have nonetheless remained unclear. We identified an

age-related link between altered basal lipid metabolism and impaired antigen responsiveness in

the naive $CD8^+$ T cell compartment. These abnormalities were associated with an enhanced

susceptibility to activation-induced apoptosis and could be recapitulated *in vitro* by exposure to

the homeostatic cytokine interleukin (IL)-7. Importantly, reversal of the bioenergetic anomalies

with lipid-altering drugs, such as rosiglitazone, almost completely restored the functional

capabilities of naive $CD8^+$ T cells. Interventions that favor lipid catabolism may therefore find

utility as adjunctive therapies in the elderly to promote vaccine-induced immunity against

emerging pathogens, such as seasonal influenza viruses and SARS-CoV-2.

**Keywords**

Fatty acids, immunometabolism, immunosenescence, naive T cells.

**INTRODUCTION**
53
Life expectancy has increased considerably over the last century as a consequence of advances in

medicine and improved public health systems. However, old age is associated with a high

prevalence of chronic diseases and an increased susceptibility to cancer and emerging pathogens,

such as SARS-CoV-2 (F. Nicoli et al., 2020). Age-related deficits in the immune system are

thought to play a key role in the development of many pathological conditions (Dorshkind &

Swain, 2009; Fulop et al., 2017; Ventura, Casciaro, Gangemi, & Buquicchio, 2017). Immune

aging is characterized by a progressive erosion of the naive $CD8^+$ T cell compartment, which

impairs *de novo* immune responses against newly encountered antigens (Briceno et al., 2016;

Nikolich-Zugich, 2014). In addition to a decline in absolute numbers (Wertheimer et al., 2014),

naive $CD8^+$ T cells in elderly individuals exhibit increased rates of homeostatic proliferation

(Sauce et al., 2012) and impaired differentiation in response to T cell receptor (TCR)-mediated

activation (Briceno et al., 2016).

A growing body of evidence indicates that lymphocyte metabolism is a key determinant of

immune functionality (Almeida, Lochner, Berod, & Sparwasser, 2016; Francesco Nicoli, 2020; F.

Nicoli, L. Papagno, et al., 2018; Palmer, Ostrowski, Balderson, Christian, & Crowe, 2015).

Systemic metabolic disturbances are common in elderly individuals, and increased levels of

adipokines and proinflammatory lipid species in particular have been implicated as critical

mediators of inflammaging, which is thought to exacerbate many age-related diseases (Baylis,

Bartlett, Patel, & Roberts, 2013). In this study, we investigated the bioenergetic features of naive

$CD8^+$ T cells in middle-aged and elderly humans, aiming to establish a link between metabolic

disturbances and immunosenescence. Naive $CD8^+$ T cells displayed a constellation of functional

and metabolic abnormalities in elderly people, including enhanced lipid influx and storage

accompanied by a marked susceptibility to apoptosis and impaired proliferation in response to

activation via the TCR. Importantly, these deficits were rectified in the presence of lipid-altering

drugs, opening potential therapeutic avenues to slow the process of immunosenescence.

**RESULTS**

**Naive CD8⁺ T cells in the elderly are susceptible to apoptosis and exhibit impaired**

**proliferation after stimulation via the TCR**

In preliminary experiments, we compared the activation profiles of naive CD8$^+$ T cells in middle-

aged and elderly individuals, mimicking antigen-driven signals with plate-bound anti-CD3. No

age-related differences in activation per se were detected 24 hr after stimulation, as determined by

measuring the upregulation of CD69 and CD134 (Figure 1A). Signals transduced via the TCR

elicit an mTOR-driven metabolic switch that supports the function and the viability of activated

naive CD8$^+$ T cells (F. Nicoli, L. Papagno, et al., 2018). We therefore assessed mTOR activity in

parallel by quantifying phospho-S6 (pS6). In line with the comparable activation profiles, naive

CD8$^+$ T cells in middle-aged and elderly individuals upregulated mTOR activity to a similar

extent after stimulation with plate-bound anti-CD3 (Figure 1B).

To extend these findings, we measured the expression of metabolism-related genes, comparing

unstimulated and activated naive CD8$^+$ T cells (Figure 1C). Genes encoding various enzymes

involved in glycolysis were upregulated similarly in flow-sorted naive CD8$^+$ T cells from middle-

aged and elderly individuals after stimulation with plate-bound anti-CD3. In contrast, genes

associated with lipid metabolism or signaling pathways involved in metabolic regulation were not

generally upregulated in response to stimulation, with the exception of *MYC*, which was

overexpressed in activated naive CD8$^+$ T cells, irrespective of age. Genes that play a critical role

in the metabolic switch were also overexpressed in activated naive CD8$^+$ T cells, irrespective of

age, with the exception of *HIF1* and *RPS6KB1*, which were upregulated to a greater extent in

activated naive CD8$^+$ T cells from middle-aged *versus* elderly individuals.

**Figure 1. Activation and gene expression in the naive CD8$^+$ T cell compartment.** (**A & B**) PBMCs from middle-aged (Mid) and elderly individuals (Old) were incubated in the absence or presence of plate-bound anti-CD3. Surface expression of the activation markers CD69 and CD134 was measured after 24 hr (A), and intracellular expression of mTOR activity marker pS6 was measured after 3 hr (B). Data are shown for naive CD8$^+$ T cells. Each dot represents one donor. Horizontal lines indicate median values. * p < 0.05 (Mann-Whitney *U* test). (**C**) Flow-sorted naive CD8$^+$ T cells from middle-aged (n = 4; Mid) and elderly individuals (n = 5; Old) were incubated in the absence or presence of plate-bound anti-CD3. Gene expression levels were measured after 3 hr. Data are shown relative to the unstimulated condition. Bars indicate mean ± SEM. * p < 0.05 (Mann-Whitney *U* test).

Despite these phenotypic and transcriptional similarities, naive CD8$^+$ T cells from middle-aged individuals proliferated to a greater extent than naive CD8$^+$ T cells from elderly individuals in response to stimulation with plate-bound anti-CD3 (Figure 2A), consistent with a previous report (Briceno et al., 2016). Early apoptosis was also less commonly observed among activated naive CD8$^+$ T cells from middle-aged *versus* elderly individuals, as determined by measuring the

expression of active caspase-3 (Figure 2B). In addition, we noted a strong inverse correlation

between the frequency of naive CD8[+] T cells that proliferated after stimulation and the frequency

of naive CD8[+] T cells that expressed active caspase-3 after stimulation, irrespective of age (Figure

2C).

Collectively, these data revealed that enhanced susceptibility to apoptosis was associated with an

age-related deficit in the proliferative capabilities of naive CD8[+] T cells, despite a largely

unaltered response to activation via the TCR.

**Figure 2. Apoptosis and proliferation in the naive CD8[+] T cell compartment.** (**A** & **B**)
PBMCs from middle-aged (Mid) and elderly individuals (Old) were stimulated with plate-bound
anti-CD3. Proliferation was measured after 4 days (A), and active caspase-3 expression was
measured after 1 day (B). Left panels: representative flow cytometry profiles. Right panels: data
summaries. Data are shown for naive CD8[+] T cells. Each dot represents one donor. Horizontal
lines indicate median values. * $p < 0.05$ (Mann-Whitney U test). (**C**) Correlation between the
frequency of naive CD8[+] T cells that proliferated after stimulation and the frequency of naive
CD8[+] T cells that expressed active caspase-3 after stimulation. Each dot represents one donor.
Significance was determined using Spearman's rank correlation.

**Naive CD8[+] T cells in the elderly display elevated levels of basal activation**

Unstimulated naive CD8[+] T cells from middle-aged individuals expressed CD134 less commonly

than unstimulated naive CD8[+] T cells from elderly individuals (Figure 1A). To consolidate this

observation, which suggested increased levels of basal activation in the elderly, we measured the

expression of T-bet, which is classically upregulated in response to activation via the TCR. The

basal expression frequencies of T-bet mirrored the basal expression frequencies of CD134 (Figure

3A). Equivalent results were obtained using a more stringent definition of naive CD8[+] T cells

(Figures S1A and S1B), which excluded phenotypically similar memory CD8[+] T cells (Pulko et

al., 2016). The basal expression frequency of T-bet also correlated directly with the activation-

induced expression frequency of active caspase-3 among naive CD8[+] T cells (Figure 3B).

Glycolysis is the main metabolic pathway that supports the activation of naive CD8[+] T cells (F.

Nicoli, L. Papagno, et al., 2018; L. Zhang & Romero, 2018). We found no significant differences

in basal glucose uptake between unstimulated naive CD8[+] T cells from middle-aged individuals

and unstimulated naive CD8[+] T cells from elderly individuals (Figure 3C). Moreover, we found

similar basal expression levels of glycolysis-related genes, with the exception of *HK2*, which was

overexpressed in unstimulated naive CD8[+] T cells from middle-aged *versus* elderly individuals

(Figure S2A). This gene encodes a selectively regulated isoform of hexokinase (H. Tan et al.,

2017), which catalyzes glucose phosphorylation, and is usually induced in response to stimulation

via the TCR (Lis et al., 2016; H. Tan et al., 2017).

**Figure 3. Basal activation in the naive CD8[+] T cell compartment.** (**A**) T-bet expression was
measured in unstimulated naive CD8[+] T cells from middle-aged (Mid) and elderly individuals
(Old). Left panel: representative flow cytometry profiles. Right panel: data summary. Each dot
represents one donor. Horizontal lines indicate median values. *** p < 0.001 (Mann-Whitney *U*
test). (**B**) Correlation between the basal expression frequency of T-bet and the activation-induced
expression frequency of active caspase-3 among naive CD8[+] T cells. Each dot represents one
donor. Significance was determined using Spearman's rank correlation. (**C**) Glucose uptake was
measured in unstimulated naive CD8[+] T cells from middle-aged (Mid) and elderly individuals
(Old) by determining the mean fluorescence intensity (MFI) of 2-NBDG. Left panel:
representative flow cytometry profiles. Right panel: data summary. Each dot represents one
donor. Horizontal lines indicate median values. Significance was determined using the Mann-
Whitney *U* test.

Collectively, these data revealed an age-related increase in the basal activation status of

unstimulated naive $CD8^+$ T cells, which appeared to be independent of the glycolysis pathway or

triggering via the TCR.

**Naive $CD8^+$ T cells in the elderly are metabolically distinct**

To explore these age-related homeostatic changes in more detail, we investigated the potential

roles of other metabolic pathways. Interestingly, fatty acid (FA) uptake was decreased among

unstimulated naive $CD8^+$ T cells from middle-aged *versus* elderly individuals (Figure 4A), but

this difference was not associated with significant changes in the expression levels of genes

encoding various enzymes involved in FA oxidation (FAO) or FA synthesis (FAS). However, we

noted that *DGAT1*, which encodes diacylglycerol O-acyltransferase 1, a key enzyme involved in

the storage of FAs as triacylglycerol (TAG), was expressed at lower levels in unstimulated naive

$CD8^+$ T cells from middle-aged *versus* elderly individuals (Figure S2B). In line with these results,

we also found that unstimulated naive $CD8^+$ T cells from middle-aged individuals stored lower

amounts of neutral lipids (NLs) than unstimulated naive $CD8^+$ T cells from elderly individuals

(Figure 4B).

**Figure 4. Basal metabolism in the naive CD8+ T cell compartment.** (**A & B**) FA uptake (A) and NL content (B) were measured in unstimulated naive CD8+ T cells from middle-aged (Mid) and elderly individuals (Old) by determining the mean fluorescence intensity (MFI) of BODIPY FL C16 and BODIPY 493/503, respectively. Left panels: representative flow cytometry profiles.

Right panels: data summaries. Each dot represents one donor. Horizontal lines indicate median
values. * p < 0.05 (Mann-Whitney *U* test). (**C**) Expression levels of genes related to signaling
pathways involved in metabolic regulation were measured in unstimulated naive CD8$^+$ T cells
flow-sorted from middle-aged (n = 5; Mid) and elderly individuals (n = 5; Old). Data are shown
relative to the housekeeping gene 18S. Bars indicate mean ± SEM. * p < 0.05 (Mann-Whitney *U*
test). (**D** & **E**) Mitochondrial membrane potential (D) and mass (E) were measured in
unstimulated naive CD8$^+$ T cells from middle-aged (Mid) and elderly individuals (Old) by
determining the mean fluorescence intensity (MFI) of TMRM and Mitotracker Deep Red,
respectively. Left panels: representative flow cytometry profiles. Right panels: data summaries.
Each dot represents one donor. Horizontal lines indicate median values. ** p < 0.01 (Mann-
Whitney *U* test).
205

In further experiments, we assessed the basal expression levels of various genes encoding

transcription factors involved in metabolic regulation. Similar patterns of expression were

observed in unstimulated naive CD8$^+$ T cells, irrespective of age, with the exception of *ID2*,

which was expressed at lower levels in unstimulated naive CD8$^+$ T cells from middle-aged *versus*

elderly individuals (Figure 4C). *ID2* is involved in metabolic adaptation (Hou et al., 2009; Z.

Zhang, Rahme, Chatterjee, Havrda, & Israel, 2017) and promotes lipid storage via the

downmodulation of *PGC-1α* (Hou et al., 2009), which enhances FAO and inhibits TAG synthesis

(Morris et al., 2012). Moreover, *ID2* promotes an overall increase in mitochondrial membrane

potential (ΔΨM), without affecting mitochondrial biogenesis or, by extension, mitochondrial

mass (Z. Zhang et al., 2017). In line with these known functions, ΔΨM was lower in unstimulated

naive CD8$^+$ T cells from middle-aged *versus* elderly individuals (Figure 4D), whereas

mitochondrial mass was largely unaffected by age (Figure 4E). We also noted a direct correlation

between ΔΨM and the frequency of unstimulated naive CD8$^+$ T cells that expressed T-bet,

suggesting a link with the loss of quiescence (Figure S3).

Collectively, these data revealed an age-related shift in the basal metabolic properties of naive

CD8$^+$ T cells, typified by high levels of FA uptake and NL storage and a supranormal ΔΨM.

**Naive CD8$^+$ T cells in the elderly display high levels of lipid uptake and storage and**

**impaired priming in the presence of IL-7**

Naive CD8$^+$ T cells are maintained primarily via homeostatic proliferation in the elderly,

counteracting an age-related decrease in the rate of *de novo* production (Cicin-Sain et al., 2007;

Sauce et al., 2012). In turn, homeostatic proliferation is driven by the common cytokine receptor γ

chain family of cytokines, particularly interleukin (IL)-7 (Guimond, Fry, & Mackall, 2005),

which regulates the proliferation of naive and memory T cells under lymphopenic conditions (Fry

& Mackall, 2005; Surh & Sprent, 2005). We hypothesized that homeostatic proliferation might

alter the basal metabolic properties of naive CD8$^+$ T cells in elderly individuals, who typically

harbor increased plasma concentrations of IL-7 (Sauce et al., 2012). In line with this notion, we

found that *in vitro* treatment with IL-7 triggered FA uptake and NL storage in the naive CD8$^+$ T

cell compartment (Figure 5A and 5B).

To explore the biological relevance of this observation, we used an *in vitro* model to prime naive

CD8$^+$ T cells specific for the melanoma-associated epitope EV10 (Melan-A$_{26–35/A27L}$), which is

restricted by the common allotype HLA-A*02:01 (Gutjahr et al., 2019; F. Nicoli, L. Papagno, et

al., 2018). Pretreatment with IL-7 significantly inhibited the expansion of EV10-specific CD8$^+$ T

cells over a period of 10 days (Figure 5C). In contrast, no inhibition was observed if IL-7 was

added to the cultures on day 4 or day 7, after the initial encounter with cognate antigen (Fig.

S4A). Moreover, the addition of IL-7 on day 10 enhanced the recovery of EV10-specific CD8$^+$ T

cells on day 30, consistent with a survival effect in the memory compartment (Figure S4B).

Collectively, these data revealed an association between altered basal lipid metabolism and

impaired antigen responsiveness as a function of age in the naive CD8$^+$ T cell compartment, both

of which could be recapitulated with exogenous IL-7.

**Figure 5. Role of IL-7 in the naive CD8$^+$ T cell compartment.** (**A & B**) PBMCs from middle-aged individuals were incubated in the absence or presence of IL-7 for 24 (A) or 48 hr (B). FA uptake (A) and NL content (B) were measured in naive CD8$^+$ T cells by determining the mean fluorescence intensity (MFI) of BODIPY FL C16 and BODIPY 493/503, respectively. Left panels: representative flow cytometry profiles. Right panels: data summaries. Each dot represents one donor. ** $p < 0.01$ (Wilcoxon signed rank test). (**C**) Percentage of tetramer$^+$ EV10-specific CD8$^+$ T cells expanded for 10 days in the presence of Flt3 ligand and TLR8L after preincubation for 24 hr in the absence or presence of IL-7. Left panel: representative flow cytometry profiles. Right panel: data summary. Each dot represents one donor. ** $p < 0.01$ (Wilcoxon signed rank test). NT: not treated.

**Naive CD8$^+$ T cells in the elderly can be reinvigorated with lipid-altering drugs**

T cell homeostasis and viability can be affected by high levels of FAs (de Jong, Kloppenburg, Toes, & Ioan-Facsinay, 2014; Francesco Nicoli, Paul, & Appay, 2018). We therefore assessed the

potential link between apoptosis and lipid metabolism. The frequency of unstimulated naive CD8[+]

T cells that expressed active caspase-3 correlated directly with basal levels of FA uptake (Figure

6A) and NL content (Figure 6B). To determine the biological relevance of these associations, we

treated naive CD8[+] T cells with rosiglitazone, a drug known to foster lipid catabolism by

activating triglyceride lipase (Kershaw et al., 2007) and preventing the conversion of FAs into

NLs (Askari et al., 2007). Pretreatment with rosiglitazone inhibited activation-induced apoptosis

among naive CD8[+] T cells, irrespective of age (Figure 6C). Similar results were obtained using

fenofibrate, which induces lipid catabolism by enhancing FAO (Figure S5).

**Figure 6. Effects of lipid-altering drugs in the naive CD8[+] T cell compartment.** (**A & B**)
Correlations between the frequency of unstimulated naive CD8[+] T cells that expressed active
caspase-3 and basal levels of FA uptake (A) and NL content (B) measured by determining the
mean fluorescence intensity (MFI) of BODIPY FL C16 and BODIPY 493/503, respectively. Each
dot represents one donor. Significance was determined using Spearman's rank correlation. (**C**)
PBMCs from middle-aged (n = 9; Mid) and elderly individuals (n = 8; Old) were stimulated with
plate-bound anti-CD3 in the absence or presence of rosiglitazone (Ros). Active caspase-3
expression was measured after 24 hr. Data are shown for naive CD8[+] T cells. Left panel:
representative flow cytometry profiles. Right panel: data summary. Bars indicate mean ± SEM. *
$p < 0.05$ (Wilcoxon signed rank test).

In line with a previous study (Angela et al., 2016), we found that NLs accumulated in naive CD8[+]

T cells as a function of proliferation in response to stimulation with plate-bound anti-CD3 (Figure

7A). We also found that proliferation under otherwise identical conditions was inhibited in the

presence of hexadecyltrimethylammonium bromide (CTAB), which disrupts FA uptake (Figure

7B). This result was expected in light of the fact that FAs are essential components of cell

membranes. However, preincubation with palmitic acid, a common saturated FA, similarly

inhibited the proliferation of naive CD8$^+$ T cells in response to stimulation with plate-bound anti-

CD3 (Figure 7C). To explore these opposing effects in more detail, we tested our system using

two interventions, namely serum starvation, which inhibits FA uptake, and exposure to

rosiglitazone, which boosts NL catabolism (Figure 7D). Naive CD8$^+$ T cells from elderly

individuals proliferated to a greater extent after serum starvation, and the addition of rosiglitazone

further enhanced these activation-induced proliferative responses to levels approaching those

observed in direct comparisons with untreated naive CD8$^+$ T cells from middle-aged individuals

(Figure 7E). Of note, these interventions suppressed the activation-induced proliferation of naive

CD8$^+$ T cells from middle-aged individuals, highlighting the importance of basal activation and

lipid metabolism as determinants of therapeutic efficacy.

To assess the potential relevance of these findings in the context of antigen-driven immune

responses, we used our *in vitro* model to prime naive CD8$^+$ T cells specific for EV10 (Briceno et

al., 2016). As expected from the proliferation data, EV10-specific CD8$^+$ T cells from middle-aged

individuals expanded to a greater extent than EV10-specific CD8$^+$ T cells from elderly individuals

(Figure 7F), and preincubation with rosiglitazone consistently enhanced the expansion of EV10-

specific CD8$^+$ T cells from elderly individuals (Figure 7G),

Collectively, these data revealed that age-related functional deficits associated with abnormal

lipid metabolism and greater levels of basal activation in the naive CD8$^+$ T cell compartment were

reversed in the presence of rosiglitazone, offering a potentially useful therapeutic option to

enhance immune reactivity against newly encountered antigens in the elderly.

**Figure 7. Potential therapeutic properties of lipid-altering drugs in the naive CD8+ T cell compartment.** (**A–C**) PBMCs from middle-aged individuals were stimulated with plate-bound anti-CD3. Proliferation was measured after 4 days. Data are shown for naive CD8+ T cells. (A) NL content was measured by determining the mean fluorescence intensity (MFI) of BODIPY 493/503 and plotted against the dilution of Cell Proliferation Dye (CPD). Flow cytometry profiles are representative of three independent experiments. (B) PBMCs were stimulated in the absence or presence of hexadecyltrimethylammonium bromide (CTAB). Flow cytometry profiles are representative of three independent experiments. (C) PBMCs were preincubated for 1 day in serum-free medium in the absence or presence of palmitic acid (PA). Each dot represents one donor. Horizontal lines indicate median values. * $p < 0.05$ (Wilcoxon signed rank test). (**D**) PBMCs were preincubated for 2 days in serum-free medium in the absence or presence of rosiglitazone (Ros). NL content was measured in naive CD8+ T cells by determining the mean fluorescence intensity (MFI) of BODIPY 493/503. Flow cytometry profiles are representative of five independent experiments. (**E**) PBMCs from middle-aged adults (Mid) and elderly individuals (Old) were preincubated for 2 days in serum-free medium in the absence or presence of rosiglitazone (Ros) and stimulated with plate-bound anti-CD3. Proliferation was measured after 4 days. Data are shown for naive CD8+ T cells. Left panel: representative flow cytometry profiles. Right panel: data summary. Each dot represents one donor. Horizontal lines indicate median

values. * p < 0.05, ** p < 0.01 (Mann-Whitney *U* test). (**F**) Percentage of tetramer[+] EV10-specific
CD8[+] T cells expanded for 10 days in the presence of Flt3 ligand and a cocktail of inflammatory
cytokines. Each dot represents one donor. Horizontal lines indicate median values. * p < 0.05
(Mann-Whitney *U* test). (**G**) Percentage of tetramer[+] EV10-specific CD8[+] T cells expanded for 10
days in the presence of Flt3 ligand and a cocktail of inflammatory cytokines after preincubation
for 2 days in the absence or presence of rosiglitazone (Ros). Left panel: representative flow
cytometry profiles. Right panel: data summary. Each dot represents one donor. Horizontal lines
indicate median values. * p < 0.05 (Wilcoxon signed rank test). NT: not treated.
341

**DISCUSSION**

A detailed understanding of age-related deficits in the naive T cell compartment is essential for

the rational development of immunotherapies and vaccines that protect elderly individuals from

emerging threats, such as COVID-19. We found that naive CD8[+] T cells in elderly individuals

were susceptible to apoptosis and proliferated suboptimally in response to stimulation via the

TCR. These abnormalities were associated with enhanced levels of basal activation, measured in

terms of $\Delta\Psi M$ and the *ex vivo* expression frequencies of T-bet and CD134.

Recent studies have shown that metabolic processes govern the behavior of T cells (Gubser et al.,

2013; F. Nicoli, L. Papagno, et al., 2018; O'Neill, Kishton, & Rathmell, 2016; L. Zhang &

Romero, 2018). In the naive CD8[+] T cell compartment, autophagy and glycolysis are typically

upregulated in response to activation (Arnold et al., 2014; F. Nicoli, L. Papagno, et al., 2018;

Whang et al., 2017), whereas homeostatic energy requirements are fulfilled primarily via FAO

(Almeida et al., 2016; Green & Beck, 2017; O'Sullivan et al., 2014; Pearce et al., 2009; Raud,

McGuire, Jones, Sparwasser, & Berod, 2018). This metabolic switch was largely unaffected by

age in our study, but at rest, naive CD8[+] T cells in the elderly displayed abnormally high levels of

FA uptake and stored abnormally high amounts of NLs.

In line with previous reports showing that high levels of intracellular lipids may be toxic (de Jong

et al., 2014; F. Nicoli, L. Papagno, et al., 2018; Zurier, Rossetti, Seiler, & Laposata, 1999), we

found that active caspase-3 expression correlated directly with FA uptake and NL content in the

naive CD8[+] T cell compartment. Lipids are essential for T cell activation and proliferation

(Angela et al., 2016). Excessively high amounts of intracellular lipids can nonetheless impair T

cell proliferation and viability (Fernanda Cury-Boaventura, Cristine Kanunfre, Gorjao, Martins de

Lima, & Curi, 2006; Howie et al., 2017; Takahashi et al., 2012). Accordingly, we found that

activation-induced apoptosis was reduced by interventions that enhanced lipid clearance in naive

CD8[+] T cells. Of note, NLs *per se* are not toxic. The conversion of FAs into NLs therefore most

likely protects against lipotoxicity under homeostatic conditions (de Jong et al., 2014).

The heightened basal activation status of naive CD8[+] T cells in elderly individuals seemed to be

sustained energetically by increased mitochondrial activity, given that T-bet expression correlated

directly with $\Delta\Psi M$. Inflammation is closely linked with metabolic dysregulation in the elderly

(Pararasa et al., 2016). High levels of circulating proinflammatory cytokines and lipids are

common features of advanced age and may contribute to the disruption of cellular quiescence.

Moreover, hematopoietic progenitor cells in elderly individuals are often metabolically active,

and this characteristic may be heritable (Fali et al., 2018). Increased rates of homeostatic

proliferation are required to maintain naive CD8[+] T cell numbers in the elderly (Sauce et al.,

2012), and the predominant energetic pathway that supports this process is thought to be FAO

(Chang et al., 2013; Ibitokou et al., 2018). It is therefore plausible that high basal levels of FA

uptake and NL storage constitute a bioenergetic pattern that favors homeostatic proliferation. IL-7

is the main driver of homeostatic proliferation in this context (J. T. Tan et al., 2001), and plasma

levels of IL-7 are typically increased in the elderly (Sauce et al., 2012). We found that IL-7

promoted FA uptake and NL storage in the naive CD8[+] T cell compartment, as reported

previously for memory CD8[+] T cells (Cui et al., 2015). In line with the findings of this latter

study, IL-7 enhanced the *in vitro* survival of antigen-experienced CD8[+] T cells. However, IL-7

also inhibited the *in vitro* expansion of naive CD8$^+$ T cells in response to antigen encounter,

suggesting that homeostatic proliferation may compromise *de novo* adaptive immunity.

Aging is characterized by profound metabolic perturbations (Bonomini, Rodella, & Rezzani,

2015), including increased lipogenesis (Kuhla, Blei, Jaster, & Vollmar, 2011) and reduced

lipolysis (Toth & Tchernof, 2000), leading to higher systemic levels of free FAs and TAG (Mc

Auley & Mooney, 2015; Pararasa et al., 2016). The combination a homeostatic environment and

high systemic levels of proinflammatory cytokines and lipids may therefore underlie the altered

metabolism and functional deficits that characterize naive CD8$^+$ T cells in the elderly. A key

finding of our study was the observation that rosiglitazone, a drug already licensed for the

treatment of diabetes mellitus, can reverse these abnormalities. This is an encouraging result from

a translational perspective, akin to recent data showing that increased FAO can improve the

development and the quality of effector CD8$^+$ T cells (Chowdhury, Chamoto, & Honjo, 2018;

Chowdhury, Chamoto, Kumar, & Honjo, 2018; F. Nicoli, L. Papagno, et al., 2018; Y. Zhang et

al., 2017). Accordingly, we propose that lipid-altering drugs may be useful adjunctive therapies to

enhance adaptive immune responses against previously unencountered antigens, particularly in

elderly individuals, who often respond poorly to vaccination and remain vulnerable to emerging

pathogens, such as seasonal influenza viruses and SARS-CoV-2.

**MATERIAL AND METHODS**

**Human subjects and samples**

Two groups of healthy volunteers were enrolled in this study: (i) middle-aged Caucasians (19–55 years); and (ii) elderly Caucasians (>65 years). Individuals with malignancies, acute diseases, or severe chronic diseases, such as atherosclerosis, congestive heart failure, poorly controlled diabetes mellitus, renal or hepatic disease, various inflammatory conditions, or chronic obstructive pulmonary disease, as well as individuals on immunosuppressive therapy, were excluded from the study. Ethical approval was granted by the Comité de Protection des Personnes of the Pitié Salpétrière Hospital (Paris, France). All volunteers provided written informed consent in accordance with the Declaration of Helsinki. PBMCs were isolated from venous blood samples via density gradient centrifugation according to standard protocols and cryopreserved in complete medium supplemented with dimethyl sulfoxide (DMSO; 10% v/v; Sigma-Aldrich) and fetal calf serum (FCS; 20% v/v; Sigma-Aldrich). Complete medium (R+) consisted of RPMI 1640 supplemented with non-essential amino acids (1% v/v), penicillin/streptomycin (100 U/mL), L-glutamine (2 mM), and sodium pyruvate (1 mM) (all from Thermo Fisher Scientific).

**Flow cytometry and cell sorting**

PBMCs were stained for surface markers using combinations of the following directly conjugated monoclonal antibodies: anti-CCR7–BV650 (clone 3D12; BD Biosciences), anti-CCR7–PE-Cy7 (clone 3D12; BD Biosciences), anti-CD3–BV605 (clone SK7; BD Biosciences), anti-CD8–APC (clone RPA-T8; BD Biosciences), anti-CD8–APC-Cy7 (clone SK1; BD Biosciences), anti-CD8–FITC (clone RPA-T8; BD Biosciences), anti-CD27–AF700 (clone O323; BioLegend), anti-CD27–PE (clone M-T271; BD Biosciences), anti-CD45RA–ECD (clone 2H4LDH11LDB9; Beckman Coulter), anti-CD45RA–PerCP-Cy5.5 (clone HI100; eBioscience), anti-CD45RA–V450

(clone HI100; BD Biosciences), anti-CD49b–PE-Cy7 (clone 9F10; BioLegend), anti-CD57–

Pacific Blue (clone HCD57; BioLegend), and anti-CD95–FITC (clone DX2; BD Biosciences).

Naive CD8$^+$ T cells were defined as CD3$^+$ CD8$^+$ CD27$^+$ CD45RA$^+$ CCR7$^+$ in most experiments

and further identified as CD49b$^-$ CD57$^-$ CD95$^-$ for gene expression studies and intracellular

measurements of T-bet. Non-viable cells were eliminated from the analysis using LIVE/DEAD

Fixable Aqua (Thermo Fisher Scientific). Intracellular stains were performed using anti-T-bet–

eFluor660 (clone 4B10; eBioscience) in conjunction with a Transcription Factor Buffer Set (BD

Biosciences). Samples were acquired using an LSR Fortessa or a FACSCanto II (BD

Biosciences). Naive CD8$^+$ T-cells were flow-sorted using a FACSAria II (BD Biosciences). Data

were analyzed using FACSDiva software version 7 (BD Biosciences) and/or FlowJo software

version 10 (FlowJo LLC).

**Activation and apoptosis assays**

PBMCs were stimulated for 24 hr with plate-bound anti-CD3 (clone OKT3; Thermo Fisher

Scientific) in the absence or presence of fenofibrate (50 µM; Sigma-Aldrich) or rosiglitazone (40

µM; Sigma-Aldrich). Activation markers were quantified on the cell surface using anti-CD69–

FITC (clone L78; BD Biosciences) and anti-CD134–BV711 (clone ACT35; BD Biosciences).

Intracellular stains were performed using anti-active caspase-3–PE (clone C92-605; BD

Biosciences) in conjunction with a Cytofix/Cytoperm Fixation/Permeabilization Solution Kit (BD

Biosciences).

**Metabolism assays**

To determine glucose uptake, PBMCs were incubated in phosphate-buffered saline (PBS)

containing 2′-(N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)amino)-2-deoxyglucose (2-NBDG; 50 µM;

Thermo Fisher Scientific). To determine FA uptake, PBMCs were incubated for 20 min at 37°C

in PBS containing 4,4-difluoro-5,7-dimethyl-4-bora-3a,4a-diaza-*s*-indacene-3-hexadecanoic acid

(BODIPY FL C16; 1 μM; Thermo Fisher Scientific). To determine neutral lipid content, PBMCs

were incubated for 20 min at 37°C in PBS containing 4,4-difluoro-1,3,5,7,8-pentamethyl-4-bora-

3a,4a-diaza-*s*-indacene (BODIPY 493/503; 10 μM; Thermo Fisher Scientific). To determine

mitochondrial mass, PBMCs were incubated for 30 min at 37°C in R+ containing Mitotracker

Deep Red (500 nM; Thermo Fisher Scientific). To determine mitochondrial membrane potential,

PBMCs were incubated for 30 min at 37°C in R+ containing tetramethylrhodamine, methyl ester,

perchlorate (TMRM; 25 nM; Thermo Fisher Scientific). To determine mTOR activity, PBMCs

were incubated for 10 min at 37°C in Cytofix Fixation Buffer (BD Biosciences), washed,

incubated for 30 min at 4°C in Phosflow Perm Buffer III (BD Biosciences), washed again, and

stained for 1 hr at room temperature with anti-pS6–Pacific Blue (clone D57.2.2E; Cell Signaling

Technology).

**Proliferation assays**

PBMCs were labeled with Cell Proliferation Dye (CPD) eFluor450 (Thermo Fisher Scientific)

and stimulated for 4 days with plate-bound anti-CD3 (clone OKT3; Thermo Fisher Scientific) in

the absence or presence of CTAB (1 μM; Sigma-Aldrich). In some experiments, cells were

precultured in AIM-V medium (Thermo Fisher Scientific) supplemented with bovine serum

albumin (BSA; 10% v/v; Sigma-Aldrich) for 1 day in the absence or presence of palmitic acid

(300 μM; Sigma-Aldrich), and in other experiments, cells were precultured in AIM-V medium

(Thermo Fisher Scientific) without BSA supplementation for 2 days in the absence or presence of

rosiglitazone (40 μM; Sigma-Aldrich). Proliferation was measured using flow cytometry to

quantify the dilution of CPD.

**Peptides and tetramers**

All peptides were synthesized at >95% purity (BioSynthesis Inc.). The EV20 peptide

(YTAAEELAGIGILTVILGVL, Melan-A$_{21–40/A27L}$) was used for *in vitro* priming studies.

Fluorochrome-labeled tetrameric complexes of HLA-A*02:01–EV10 (ELAGIGILTV, Melan-

A$_{26–35/A27L}$) were generated in-house as described previously (Price et al., 2005).

***In vitro* priming of antigen-specific CD8$^+$ T cells**

Naive precursors specific for HLA-A2–EV10 were primed *in vitro* using an accelerated dendritic

cell (DC) coculture protocol as described previously (Alanio et al., 2015; Lissina et al., 2016; F.

Nicoli, L. Papagno, et al., 2018). Briefly, thawed PBMCs were resuspended at $5 \times 10^6$ cells/well

in AIM-V medium (Thermo Fisher Scientific) supplemented with Flt3 ligand (Flt3L; 50 ng/ml;

R&D Systems) in the absence or presence of rosiglitazone (40 μM; Sigma-Aldrich) or IL-7 (20

ng/mL; R&D Systems). After 24 hr (day 1), the Melan-A peptide EV20 (1 μM) was added to the

cultures, and DC maturation was induced using a standard cocktail of inflammatory cytokines,

incorporating IL-1β (10 ng/mL), IL-7 (0.5 ng/mL), prostaglandin E2 (PGE2; 1 μM), and TNF

(1,000 U/mL) (all from R&D Systems), or ssRNA40 (TLR8L; 0.5 μg/mL; InvivoGen). The

cultures were supplemented on day 2 with FCS (10% v/v; Sigma-Aldrich). Medium was replaced

every 3 days thereafter with fresh R+ containing FCS (10% v/v; Sigma-Aldrich). In some

experiments, the cultures were further supplemented on day 4 or day 7 with IL-7 (10 ng/mL;

R&D Systems). Antigen-specific CD8$^+$ T cells were characterized via flow cytometry on day 10.

Long-term cultures were maintained similarly until day 30 in the absence or presence of IL-7 (10

mg/mL; R&D Systems).

**RNA extraction and qPCR analysis**

PBMCs were activated for 5 hr with plate-bound anti-CD3 (clone OKT3; Thermo Fisher

Scientific). RNA was extracted from flow-sorted naive CD8$^+$ T cells (n = 300 per condition)

using a NucleoSpin RNA XS Kit (Macherey-Nagel), and cDNA was synthesized using Reverse

Transcription Master Mix (Fluidigm). Specific targets were amplified using PreAmp Master Mix

(Fluidigm). Gene expression was assessed using a BioMark HD System (Fluidigm) with

EvaGreen Supermix (Bio-Rad). RNA expression levels were calculated using the $2^{-\Delta\Delta CT}$ method

with reference to a housekeeping gene (human 18S) (F. Nicoli, E. Gallerani, et al., 2018).

**General statistics**

Univariate statistical analyses were performed using nonparametric tests in Prism software

version 8 (GraphPad Software Inc.). Significance was assigned at $p < 0.05$.

**ACKNOWLEDGEMENTS**

We thank Simone Candioli and Silvio Spatocco (University of Ferrara, Italy), Irene Bonazzi and

Ilaria Signoretto (University of Padua, Italy), and Alain Savenay (AP-HP, Paris, France) for

technical assistance, Mario Pende (Institut Necker-Enfants Malades, Paris, France) for helpful

discussions, and Veronique Morin and Rima Zoorob (INSERM U1135, Paris, France) and Silvia

Menegatti and Lars Rogge (Institut Pasteur, Paris, France) for assistance with gene expression

analyses. This work was supported by the Agence National de la Recherche (ANR-14-CE14-

0030-01) and the Université Franco-Italienne/Università Italo-Francese (Galileo Project G10-718;

PHC Galilee Project 39582TJ). Core provision of the FACSCanto II was funded by the University

of Ferrara (Bando per l'acquisizione di strumenti per la ricerca di ateneo-2015). D.A.P. was

supported by a Wellcome Trust Senior Investigator Award (100326/Z/12/Z).

**CONFLICT OF INTERESTS**

The authors declare no conflict of interest.

**AUTHORS' CONTRIBUTIONS**

F.N., R.G., and V.A. conceptualized the project; F.N., M.P.C.P., L.P., E.Ga., V.F., and J.J.F.

performed experiments and analyzed data; M.D., E.C., H.V., E.Go., S.L.L., D.A.P., A.T., and J.B.

provided critical resources; F.N., J.F.F., R.G., and V.A. drafted the manuscript; F.N., D.A.P.,

A.C., R.G., and V.A. edited the manuscript; D.A.P., A.T., A.C., and V.A. acquired funds to

support the work. All authors contributed intellectually and approved the manuscript.

**REFERENCES**

Alanio, C., Nicoli, F., Sultanik, P., Flecken, T., Perot, B., Duffy, D., . . . Albert, M. L. (2015).
Bystander hyperactivation of preimmune CD8+ T cells in chronic HCV patients. *Elife, 4*, 1-
20. doi:10.7554/eLife.07916
Almeida, L., Lochner, M., Berod, L., & Sparwasser, T. (2016). Metabolic pathways in T cell
activation and lineage differentiation. *Semin Immunol, 28*(5), 514-524.
doi:10.1016/j.smim.2016.10.009
Angela, M., Endo, Y., Asou, H. K., Yamamoto, T., Tumes, D. J., Tokuyama, H., . . . Nakayama, T.
(2016). Fatty acid metabolic reprogramming via mTOR-mediated inductions of
PPARgamma directs early activation of T cells. *Nat Commun, 7*, 13683.
doi:10.1038/ncomms13683
Arnold, C. R., Pritz, T., Brunner, S., Knabb, C., Salvenmoser, W., Holzwarth, B., . . . Grubeck-
Loebenstein, B. (2014). T cell receptor-mediated activation is a potent inducer of
macroautophagy in human CD8(+)CD28(+) T cells but not in CD8(+)CD28(-) T cells. *Exp
Gerontol, 54*, 75-83. doi:10.1016/j.exger.2014.01.018
Askari, B., Kanter, J. E., Sherrid, A. M., Golej, D. L., Bender, A. T., Liu, J., . . . Bornfeldt, K. E. (2007).
Rosiglitazone inhibits acyl-CoA synthetase activity and fatty acid partitioning to
diacylglycerol and triacylglycerol via a peroxisome proliferator-activated receptor-
gamma-independent mechanism in human arterial smooth muscle cells and
macrophages. *Diabetes, 56*(4), 1143-1152. doi:10.2337/db06-0267
Baylis, D., Bartlett, D. B., Patel, H. P., & Roberts, H. C. (2013). Understanding how we age:
insights into inflammaging. *Longev Healthspan, 2*(1), 8. doi:10.1186/2046-2395-2-8
Bonomini, F., Rodella, L. F., & Rezzani, R. (2015). Metabolic syndrome, aging and involvement of
oxidative stress. *Aging Dis, 6*(2), 109-120. doi:10.14336/AD.2014.0305
Briceno, O., Lissina, A., Wanke, K., Afonso, G., von Braun, A., Ragon, K., . . . Appay, V. (2016).
Reduced naive CD8(+) T-cell priming efficacy in elderly adults. *Aging Cell, 15*(1), 14-21.
doi:10.1111/acel.12384
Chang, C. H., Curtis, J. D., Maggi, L. B., Jr., Faubert, B., Villarino, A. V., O'Sullivan, D., . . . Pearce, E.
L. (2013). Posttranscriptional control of T cell effector function by aerobic glycolysis. *Cell,
153*(6), 1239-1251. doi:10.1016/j.cell.2013.05.016
Chowdhury, P. S., Chamoto, K., & Honjo, T. (2018). Combination therapy strategies for improving
PD-1 blockade efficacy: a new era in cancer immunotherapy. *J Intern Med, 283*(2), 110-
120. doi:10.1111/joim.12708
Chowdhury, P. S., Chamoto, K., Kumar, A., & Honjo, T. (2018). PPAR-Induced Fatty Acid Oxidation
in T Cells Increases the Number of Tumor-Reactive CD8(+) T Cells and Facilitates Anti-PD-
1 Therapy. *Cancer Immunol Res*. doi:10.1158/2326-6066.CIR-18-0095
Cicin-Sain, L., Messaoudi, I., Park, B., Currier, N., Planer, S., Fischer, M., . . . Nikolich-Zugich, J.
(2007). Dramatic increase in naive T cell turnover is linked to loss of naive T cells from old
primates. *Proc Natl Acad Sci U S A, 104*(50), 19960-19965. doi:10.1073/pnas.0705905104
Cui, G., Staron, M. M., Gray, S. M., Ho, P. C., Amezquita, R. A., Wu, J., & Kaech, S. M. (2015). IL-7-
Induced Glycerol Transport and TAG Synthesis Promotes Memory CD8+ T Cell Longevity.
*Cell, 161*(4), 750-761. doi:10.1016/j.cell.2015.03.021
de Jong, A. J., Kloppenburg, M., Toes, R. E., & Ioan-Facsinay, A. (2014). Fatty acids, lipid
mediators, and T-cell function. *Front Immunol, 5*, 483. doi:10.3389/fimmu.2014.00483

Dorshkind, K., & Swain, S. (2009). Age-associated declines in immune system development and
function: causes, consequences, and reversal. *Curr Opin Immunol, 21*(4), 404-407.
doi:10.1016/j.coi.2009.07.001
Fali, T., Fabre-Mersseman, V., Yamamoto, T., Bayard, C., Papagno, L., Fastenackels, S., . . . Appay,
V. (2018). Elderly human hematopoietic progenitor cells express cellular senescence
markers and are more susceptible to pyroptosis. *JCI Insight, 3*(13).
doi:10.1172/jci.insight.95319
Fernanda Cury-Boaventura, M., Cristine Kanunfre, C., Gorjao, R., Martins de Lima, T., & Curi, R.
(2006). Mechanisms involved in Jurkat cell death induced by oleic and linoleic acids. *Clin
Nutr, 25*(6), 1004-1014. doi:10.1016/j.clnu.2006.05.008
Fry, T. J., & Mackall, C. L. (2005). The many faces of IL-7: from lymphopoiesis to peripheral T cell
maintenance. *J Immunol, 174*(11), 6571-6576.
Fulop, T., Larbi, A., Dupuis, G., Le Page, A., Frost, E. H., Cohen, A. A., . . . Franceschi, C. (2017).
Immunosenescence and Inflamm-Aging As Two Sides of the Same Coin: Friends or Foes?
*Front Immunol, 8*, 1960. doi:10.3389/fimmu.2017.01960
Green, W. D., & Beck, M. A. (2017). Obesity altered T cell metabolism and the response to
infection. *Curr Opin Immunol, 46*, 1-7. doi:10.1016/j.coi.2017.03.008
Gubser, P. M., Bantug, G. R., Razik, L., Fischer, M., Dimeloe, S., Hoenger, G., . . . Hess, C. (2013).
Rapid effector function of memory CD8+ T cells requires an immediate-early glycolytic
switch. *Nat Immunol, 14*(10), 1064-1072. doi:10.1038/ni.2687
Guimond, M., Fry, T. J., & Mackall, C. L. (2005). Cytokine signals in T-cell homeostasis. *J
Immunother, 28*(4), 289-294.
Gutjahr, A., Papagno, L., Nicoli, F., Kanuma, T., Kuse, N., Cabral-Piccin, M. P., . . . Appay, V.
(2019). The STING ligand cGAMP potentiates the efficacy of vaccine-induced CD8+ T cells.
*JCI Insight, 4*(7). doi:10.1172/jci.insight.125107
Hou, T. Y., Ward, S. M., Murad, J. M., Watson, N. P., Israel, M. A., & Duffield, G. E. (2009). ID2
(inhibitor of DNA binding 2) is a rhythmically expressed transcriptional repressor required
for circadian clock output in mouse liver. *J Biol Chem, 284*(46), 31735-31745.
doi:10.1074/jbc.M109.013961
Howie, D., Cobbold, S. P., Adams, E., Ten Bokum, A., Necula, A. S., Zhang, W., . . . Waldmann, H.
(2017). Foxp3 drives oxidative phosphorylation and protection from lipotoxicity. *JCI
Insight, 2*(3), e89160. doi:10.1172/jci.insight.89160
Ibitokou, S. A., Dillon, B. E., Sinha, M., Szczesny, B., Delgadillo, A., Reda Abdelrahman, D., . . .
Stephens, R. (2018). Early Inhibition of Fatty Acid Synthesis Reduces Generation of
Memory Precursor Effector T Cells in Chronic Infection. *J Immunol, 200*(2), 643-656.
doi:10.4049/jimmunol.1602110
Kershaw, E. E., Schupp, M., Guan, H. P., Gardner, N. P., Lazar, M. A., & Flier, J. S. (2007).
PPARgamma regulates adipose triglyceride lipase in adipocytes in vitro and in vivo. *Am J
Physiol Endocrinol Metab, 293*(6), E1736-1745. doi:10.1152/ajpendo.00122.2007
Kuhla, A., Blei, T., Jaster, R., & Vollmar, B. (2011). Aging is associated with a shift of fatty
metabolism toward lipogenesis. *J Gerontol A Biol Sci Med Sci, 66*(11), 1192-1200.
doi:10.1093/gerona/glr124
Lis, P., Dylag, M., Niedzwiecka, K., Ko, Y. H., Pedersen, P. L., Goffeau, A., & Ulaszewski, S. (2016).
The HK2 Dependent "Warburg Effect" and Mitochondrial Oxidative Phosphorylation in
Cancer: Targets for Effective Therapy with 3-Bromopyruvate. *Molecules, 21*(12).
doi:10.3390/molecules21121730

Lissina, A., Briceno, O., Afonso, G., Larsen, M., Gostick, E., Price, D. A., . . . Appay, V. (2016).
Priming of Qualitatively Superior Human Effector CD8+ T Cells Using TLR8 Ligand
Combined with FLT3 Ligand. *J Immunol, 196*(1), 256-263. doi:10.4049/jimmunol.1501140
Mc Auley, M. T., & Mooney, K. M. (2015). Computationally Modeling Lipid Metabolism and
Aging: A Mini-review. *Comput Struct Biotechnol J, 13*, 38-46.
doi:10.1016/j.csbj.2014.11.006
Morris, E. M., Meers, G. M., Booth, F. W., Fritsche, K. L., Hardin, C. D., Thyfault, J. P., & Ibdah, J.
A. (2012). PGC-1alpha overexpression results in increased hepatic fatty acid oxidation
with reduced triacylglycerol accumulation and secretion. *Am J Physiol Gastrointest Liver
Physiol, 303*(8), G979-992. doi:10.1152/ajpgi.00169.2012
Nicoli, F. (2020). Angry, Hungry T-Cells: How Are T-Cell Responses Induced in Low Nutrient
Conditions? *Immunometabolism, 2*(1), e200004. doi:10.20900/immunometab20200004
Nicoli, F., Gallerani, E., Sforza, F., Finessi, V., Chachage, M., Geldmacher, C., . . . Gavioli, R. (2018).
The HIV-1 Tat protein affects human CD4+ T-cell programing and activation, and favors
the differentiation of naive CD4+ T cells. *AIDS, 32*(5), 575-581.
doi:10.1097/QAD.0000000000001734
Nicoli, F., Papagno, L., Frere, J. J., Cabral-Piccin, M. P., Clave, E., Gostick, E., . . . Appay, V. (2018).
Naive CD8(+) T-Cells Engage a Versatile Metabolic Program Upon Activation in Humans
and Differ Energetically From Memory CD8(+) T-Cells. *Front Immunol, 9*, 2736.
doi:10.3389/fimmu.2018.02736
Nicoli, F., Paul, S., & Appay, V. (2018). Harnessing the Induction of CD8+ T-Cell Responses
Through Metabolic Regulation by Pathogen-Recognition-Receptor Triggering in Antigen
Presenting Cells. *Frontiers in Immunology, 9*(2372). doi:10.3389/fimmu.2018.02372
Nicoli, F., Solis-Soto, M. T., Paudel, D., Marconi, P., Gavioli, R., Appay, V., & Caputo, A. (2020).
Age-related decline of de novo T cell responsiveness as a cause of COVID-19 severity.
*Geroscience*. doi:10.1007/s11357-020-00217-w
Nikolich-Zugich, J. (2014). Aging of the T cell compartment in mice and humans: from no naive
expectations to foggy memories. *J Immunol, 193*(6), 2622-2629.
doi:10.4049/jimmunol.1401174
O'Neill, L. A., Kishton, R. J., & Rathmell, J. (2016). A guide to immunometabolism for
immunologists. *Nat Rev Immunol, 16*(9), 553-565. doi:10.1038/nri.2016.70
O'Sullivan, D., van der Windt, G. J., Huang, S. C., Curtis, J. D., Chang, C. H., Buck, M. D., . . .
Pearce, E. L. (2014). Memory CD8(+) T cells use cell-intrinsic lipolysis to support the
metabolic programming necessary for development. *Immunity, 41*(1), 75-88.
doi:10.1016/j.immuni.2014.06.005
Palmer, C. S., Ostrowski, M., Balderson, B., Christian, N., & Crowe, S. M. (2015). Glucose
metabolism regulates T cell activation, differentiation, and functions. *Front Immunol, 6*,
1. doi:10.3389/fimmu.2015.00001
Pararasa, C., Ikwuobe, J., Shigdar, S., Boukouvalas, A., Nabney, I. T., Brown, J. E., . . . Griffiths, H.
R. (2016). Age-associated changes in long-chain fatty acid profile during healthy aging
promote pro-inflammatory monocyte polarization via PPARgamma. *Aging Cell, 15*(1),
128-139. doi:10.1111/acel.12416
Pearce, E. L., Walsh, M. C., Cejas, P. J., Harms, G. M., Shen, H., Wang, L. S., . . . Choi, Y. (2009).
Enhancing CD8 T-cell memory by modulating fatty acid metabolism. *Nature, 460*(7251),
103-107. doi:10.1038/nature08097

Price, D. A., Brenchley, J. M., Ruff, L. E., Betts, M. R., Hill, B. J., Roederer, M., . . . Douek, D. C.
(2005). Avidity for antigen shapes clonal dominance in CD8+ T cell populations specific
for persistent DNA viruses. *J Exp Med, 202*(10), 1349-1361. doi:10.1084/jem.20051357
Pulko, V., Davies, J. S., Martinez, C., Lanteri, M. C., Busch, M. P., Diamond, M. S., . . . Nikolich-
Zugich, J. (2016). Human memory T cells with a naive phenotype accumulate with aging
and respond to persistent viruses. *Nat Immunol, 17*(8), 966-975. doi:10.1038/ni.3483
Raud, B., McGuire, P. J., Jones, R. G., Sparwasser, T., & Berod, L. (2018). Fatty acid metabolism in
CD8(+) T cell memory: Challenging current concepts. *Immunol Rev, 283*(1), 213-231.
doi:10.1111/imr.12655
Sauce, D., Larsen, M., Fastenackels, S., Roux, A., Gorochov, G., Katlama, C., . . . Appay, V. (2012).
Lymphopenia-driven homeostatic regulation of naive T cells in elderly and
thymectomized young adults. *J Immunol, 189*(12), 5541-5548.
doi:10.4049/jimmunol.1201235
Surh, C. D., & Sprent, J. (2005). Regulation of mature T cell homeostasis. *Semin Immunol, 17*(3),
183-191. doi:10.1016/j.smim.2005.02.007
Takahashi, H. K., Cambiaghi, T. D., Luchessi, A. D., Hirabara, S. M., Vinolo, M. A., Newsholme, P.,
& Curi, R. (2012). Activation of survival and apoptotic signaling pathways in lymphocytes
exposed to palmitic acid. *J Cell Physiol, 227*(1), 339-350. doi:10.1002/jcp.22740
Tan, H., Yang, K., Li, Y., Shaw, T. I., Wang, Y., Blanco, D. B., . . . Chi, H. (2017). Integrative
Proteomics and Phosphoproteomics Profiling Reveals Dynamic Signaling Networks and
Bioenergetics Pathways Underlying T Cell Activation. *Immunity, 46*(3), 488-503.
doi:10.1016/j.immuni.2017.02.010
Tan, J. T., Dudl, E., LeRoy, E., Murray, R., Sprent, J., Weinberg, K. I., & Surh, C. D. (2001). IL-7 is
critical for homeostatic proliferation and survival of naive T cells. *Proc Natl Acad Sci U S
A, 98*(15), 8732-8737. doi:10.1073/pnas.161126098
Toth, M. J., & Tchernof, A. (2000). Lipid metabolism in the elderly. *Eur J Clin Nutr, 54 Suppl 3*,
S121-125.
Ventura, M. T., Casciaro, M., Gangemi, S., & Buquicchio, R. (2017). Immunosenescence in aging:
between immune cells depletion and cytokines up-regulation. *Clin Mol Allergy, 15*, 21.
doi:10.1186/s12948-017-0077-0
Wertheimer, A. M., Bennett, M. S., Park, B., Uhrlaub, J. L., Martinez, C., Pulko, V., . . . Nikolich-
Zugich, J. (2014). Aging and cytomegalovirus infection differentially and jointly affect
distinct circulating T cell subsets in humans. *J Immunol, 192*(5), 2143-2155.
doi:10.4049/jimmunol.1301721
Whang, M. I., Tavares, R. M., Benjamin, D. I., Kattah, M. G., Advincula, R., Nomura, D. K., . . . Ma,
A. (2017). The Ubiquitin Binding Protein TAX1BP1 Mediates Autophagasome Induction
and the Metabolic Transition of Activated T Cells. *Immunity, 46*(3), 405-420.
doi:10.1016/j.immuni.2017.02.018
Zhang, L., & Romero, P. (2018). Metabolic Control of CD8(+) T Cell Fate Decisions and Antitumor
Immunity. *Trends Mol Med, 24*(1), 30-48. doi:10.1016/j.molmed.2017.11.005
Zhang, Y., Kurupati, R., Liu, L., Zhou, X. Y., Zhang, G., Hudaihed, A., . . . Ertl, H. C. J. (2017).
Enhancing CD8+ T Cell Fatty Acid Catabolism within a Metabolically Challenging Tumor
Microenvironment Increases the Efficacy of Melanoma Immunotherapy. *Cancer Cell,
32*(3), 377-391 e379. doi:10.1016/j.ccell.2017.08.004
Zhang, Z., Rahme, G. J., Chatterjee, P. D., Havrda, M. C., & Israel, M. A. (2017). ID2 promotes
survival of glioblastoma cells during metabolic stress by regulating mitochondrial
function. *Cell Death Dis, 8*(2), e2615. doi:10.1038/cddis.2017.14

Zurier, R. B., Rossetti, R. G., Seiler, C. M., & Laposata, M. (1999). Human peripheral blood T
lymphocyte proliferation after activation of the T cell receptor: effects of unsaturated
fatty acids. *Prostaglandins Leukot Essent Fatty Acids, 60*(5-6), 371-375.

**SUPPLEMENTAL INFORMATION**

**Table S1. Antibodies used in flow cytometry experiments.**

| ANTIBODY | FLUOROCHROME | SUPPLIER AND REFERENCE |
|---|---|---|
| Active caspase-3 | PE | BD Biosciences - 561011 |
| CCR7 | BV650 | BD Biosciences - 563407 |
| CCR7 | PE-Cy7 | BD Biosciences - 557648 |
| CD3 | BV605 | BD Biosciences - 563219 |
| CD8 | APC | BD Biosciences - 555369 |
| CD8 | APC-Cy7 | BD Biosciences - 557834 |
| CD8 | FITC | BD Biosciences - 555366 |
| CD27 | AF700 | BioLegend - 302814 |
| CD27 | BUV395 | BD Biosciences - 563815 |
| CD27 | PE | BD Biosciences - 555441 |
| CD45RA | ECD | Beckman Coulter - IM2711U |
| CD45RA | PerCP-Cy5.5 | eBioscience - 45-0458-42 |
| CD45RA | V450 | BD Biosciences - 560362 |
| CD49d | PE-Cy7 | BioLegend - 304314 |
| CD57 | Pacific Blue | BioLegend - 322316 |
| CD69 | FITC | BD Biosciences - 347823 |
| CD95 | FITC | BD Biosciences - 555673 |
| CD134 | BV711 | BD Biosciences - 563664 |
| pS6 | Pacific Blue | Cell Signaling Technology - 8520S |
| T-bet | eFluor660 | eBioscience - 50-5825-80 |

**Figure S1. T-bet expression in naive CD8$^+$ T cells.** (**A**) Flow cytometric gating strategy used to

exclude memory CD8$^+$ T cells with a naive phenotype (MNP). (B) T-bet expression in naive

CD8$^+$ T cells (CD3$^+$ CD8$^+$ CD27$^+$ CD45RA$^+$ CCR7$^+$ CD49d$^-$ CD57$^-$ CD95$^-$) from middle-aged

(Mid) and elderly individuals (Old). Each dot represents one donor. Horizontal lines indicate

median values. * p < 0.05 (Mann-Whitney *U* test).

**Figure S2. Expression of metabolism-related genes in naive CD8<sup>+</sup> T cells. (A & B)** Expression

levels of genes related to glucose (A) and fatty acid metabolism (B) were measured in

unstimulated naive CD8<sup>+</sup> T cells flow-sorted from middle-aged (n = 5; Mid) and elderly

individuals (n = 5; Old). Bars indicate mean ± SEM. * $p < 0.05$ (Mann-Whitney *U* test).

**Figure S3. Correlation between T-bet expression and mitochondrial membrane potential in naive CD8+ T cells.** Correlation between the frequency of unstimulated naive CD8+ T cells that expressed T-bet and basal mitochondrial membrane potential, measured by determining the mean fluorescence intensity (MFI) of TMRM. Each dot represents one donor. Significance was determined using Spearman's rank correlation.

**Figure S4. Effect of IL-7 on primed epitope-specific CD8[+] T cells. (A & B)** EV10-specific

CD8[+] T cells were primed and expanded for 10 days in the presence of Flt3 ligand and TLR8L.

(A) Cells were left unmanipulated or treated with IL-7 on day 4 or day 7 and tetramer-stained on

day 10. Each dot represents one donor. (B) Cells were left unmanipulated or treated with IL-7

from day 11 to day 30 and tetramer-stained on day 30. Each dot represents one donor. * $p < 0.05$

(Wilcoxon signed rank test). NT: not treated.

775 **Figure S5. Effects of fenofibrate on naive CD8[+] T cells.** PBMCs from middle-aged (n = 9; Mid)

776 and elderly individuals (n = 8; Old) were stimulated with plate-bound anti-CD3 in the absence or

777 presence of fenofibrate (Fen). Active caspase-3 expression was measured after 24 hr. Data are

778 shown for naive CD8[+] T cells. Left panel: representative flow cytometry profiles. Right panel:

779 data summary. Bars indicate mean ± SEM. * $p < 0.05$ (Wilcoxon signed rank test).

780
781