Molecular characterization revealed the role of thaumatin-like proteins in stress response in bread wheat

Alok Sharma[1], Himanshu Sharma[1,2], Ruchika Rajput[3], Ashutosh Pandey[3] and Santosh Kumar Upadhyay[*1]

1. Department of Botany, Panjab University, Chandigarh, India-160014

2. I.K. Gujral Punjab Technical University, Kapurthala, Jalandhar, India-144603

3. National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi, India-110067

*Corresponding Author

*Corresponding Author: Dr. Santosh Kumar Upadhyay, Department of Botany, Panjab University, Chandigarh, India-160014. Email: skupadhyay@pu.ac.in, Tel: +91-172-2534001, Fax: +91-172-2779510

ABSTRACT

Thaumatin-like proteins (TLPs) are related to the pathogenesis-related-5 (PR-5) family and involved in stress response. Herein, a total of 93 *TLP* genes were identified in the genome of *Triticum aestivum*. Further, we identified 26, 27, 39 and 37 *TLP* genes in the *Brachypodium distachyon*, *Oryza sativa*, *Sorghum bicolor* and *Zea mays* genomes for comparative characterization, respectively. They could be grouped into small and long TLPs with conserved thaumatin signature motif. Tightly clustered genes exhibited conserved gene and protein structure. The physicochemical analyses suggested significant differences between small and long TLPs. Evolutionary analyses suggested the role of duplication events and purifying selection in the expansion of the TLP gene family. Expression analyses revealed the possible roles of *TLPs* in plant development and abiotic and fungal stress response. Recombinant expression of *TaTLP2-B* in *Saccharomyces cerevisiae* provided significant tolerance against cold, heat, drought and salt stresses. The results depicted the importance of TLPs in cereal crops that would be highly useful in future crop improvement programs.

Keywords: Abiotic stress, Biotic stress, Cereals, Thaumatin-like proteins, *TaTLP2-B*, *Triticum aestivum*

1. Introduction

The growth and development of plants have always been affected by various abiotic and biotic stress conditions. In response to these stress conditions, plants produce numerous molecules including pathogenesis-related (PR) proteins [1,2]. PR proteins belong to a superfamily of defence-related proteins consisting of PR1-PR17 protein families, that have been classified based on amino acid sequences, serological reaction, enzymatic activity etc. [3]

Thaumatin-like proteins (TLPs) are related to the PR5 family of the PR superfamily [3]. These are called TLPs due to their significant similarity with a ~23 kDa sweet-tasting protein known as thaumatin [4]. Thaumatin protein was first isolated from an African shrub known as *Thaumatococcus danielli* [5]. Over the period, TLPs have been reported in a diverse group of organisms including insects, nematodes, fungi and plants [6,7]. In plants, TLPs are reported from algae to angiosperms [6-9].

Based on the molecular weight (MW), TLP proteins are grouped as long (L-type) and small (S-type) TLPs, having the MW ~21-26 kDa and ~16-17 kDa, respectively. Long TLPs have been identified from each group of plants, while small TLPs are confined to only gymnospermous and monocot plants [6,9]. A total of 16 and 10 conserved cysteine residues, forming eight and five disulfide linkages, are known to be present in long and small TLPs, respectively. These disulfide bonds provide resistance against extreme pH, heat and protease degradation [6,10]. Each TLP consists of a conserved REDDD motif and a thaumatin signature motif G-X-[GF]-X-C-X-T-[GA]-D-C-X- (1,2)-G-X-(2,3)-C. The REDDD motif is involved in receptor binding for their antifungal action [6]. Numerous studies reported that the overexpression of TLPs provides significant resistance against various fungi in both dicot and monocot plants [11-13]. The mechanism of action of TLPs in fungal resistance is ambiguous; however, it has been presumed

that they work by degradation and permeabilization of the fungal cell walls [6,14]. Besides,

TLPs are also known to be involved in antifreeze action and abiotic stress resistance in plants

[15]. Further, they are also reported to be involved in flowering, fruit ripening and seed

germination [16-18].

Despite utmost importance, a detailed characterization of TLPs is still limited in various cereal

crops. Therefore, the current study aims at an inclusive characterization of TLPs in bread wheat,

an important cereal crop. Besides, we have also performed a comparative analysis of various

features of TLPs in other cereals including *B. distachyon*, *O. sativa*, *S. bicolor* and *Z. mays*.

Chromosomal distribution, phylogeny, physicochemical properties, gene duplication events, *cis*-

regulatory elements, and gene and protein structural analyses were performed. To decipher their

roles in plant development and under abiotic and biotic stresses, their expression analysis was

performed using high throughput RNA-seq data. The expression of eight selected *TaTLP* genes

was validated using qRT-PCR. An abiotic stress-responsive gene, *TaTLP2-B*, of *T. aestivum* was

cloned and used for functional characterization in *Saccharomyces cerevisiae* (yeast) cells.

Recombinant expression of TaTLP2-B provided significant tolerance to the yeast in spot assay

against various abiotic stresses. The study will increase the knowledge about numerous TLP

proteins and provide the base for the functional characterization of identified genes in future

studies.

2. Material and methods

2.1. Plant materials, growth conditions and stress treatments

Surface sterilization of bread wheat seeds (cv. Chinese Spring) was done with 1.2% sodium

hypochlorite in 10% ethanol followed by the washing with double autoclaved water. The

sterilized seeds were kept overnight on moist Whatman filter papers at 4°C for stratification and

were further kept at room temperature for germination. The seedlings were then transferred to

fresh phytaboxes and were allowed to grow in plant growth chambers under 60% relative

humidity, 22°C temperature and a 16 and 8 h light and dark period, respectively [19]. Seven-

days old seedlings were kept under heat (40°C), drought (20% PEG) and combined heat and

drought stress treatments for 6, 12, 24 and 48 hours (h). For salinity stress, plants were subjected

to 150 mM NaCl for 6, 12, 24 and 48 h. The plant samples including roots and shoots were

collected and stored at -80°C till further experiments.

2.2. Identification and nomenclature of the TLP genes

Extensive BLAST searches were used to identify the TLP proteins in five different cereals

including *B. distachyon*, *O. sativa*, *S. bicolor*, *T. aestivum* and *Z. mays*. Arabidopsis TLP

sequences were used as a query against the protein model sequences of each crop downloaded

from the Ensembl Plants [9,20]. The presence of the thaumatin (PF00314) domain was analyzed

using the Hidden Markov Model (HMM) and Pfam BLAST searches at e-value 10^{-10}. Identified

TLP sequences were further subjected to the NCBI Conserved Domain Database (CDD) BLAST

to further confirm the presence of the thaumatin domain. The *TLP* genes, having a complete

thaumatin family signature, were selected for further analysis. Identified *TLPs* were named as

per their sequence of occurrence at various chromosomes in each crop, except *T. aestivum*. The

international   rules   for   gene   symbolization   of   *T.   aestivum*

(http://wheat.pw.usda.gov/ggpages/wgc/98/Intro.htm) were followed for the nomenclature of

TaTLPs.

2.3. Chromosomal localization and duplication events

Plant Ensemble (http://plants.ensembl.org/) was used for gathering the chromosomal and sub-

genomic location of *TLPs* of all five crops. The homeologous *TLP* genes in *T. aestivum* were

identified based on ≥90% sequence similarity and their occurrence at the related chromosomes.

The MapInspect software was used for the graphical representation of *TLP* genes on their

respective chromosomes (http://www.plantbreeding.wur .nl/uk/software_mapinspect.html.2012).

Duplication events (DEs) were identified using a bidirectional blast hit approach with sequence

similarity of ≥80%, while tandem and segmental DEs were segregated based on their distance

and occurrence at respective chromosomes as per the previous studies [21].

2.4. Multiple sequence alignments, phylogeny and structural analysis

The Muscle and Multalin tools were used for the multiple sequence alignments of all the TLP*s*

with a known thaumatin protein (P02883.2|THM1_THADA) to find the conserved residues

[22,23]. The phylogenetic tree was constructed by the maximum likelihood method with 1000

bootstraps using the MEGA X software [24].

2.5. Ka/Ks and Tajima's relativity test

The alignment of the protein and nucleotide sequences of the paralogous gene pairs was done

using the ClustalOmega server (https://www.ebi.ac.uk/Tools/msa/clustalo). The synonymous

substitution per synonymous site (Ks), the non-synonymous substitution per non-synonymous

site (Ka), and their ratio (Ka/Ks) were calculated using the PAL2NAL programme [25,26]. The

calculation of the divergence time of each pair of duplicated genes was done using the formula

T=Ks/2r, where T represents the divergence time and r represents the divergence rate. The

divergence rate was assumed to be 6.5×10^{-9} for cereals [27]. The Tajima's relativity test was

performed to find out the evolutionary rate between paralogous genes [28].

6

2.6. Gene structure organization

The genomic and coding DNA sequence (CDS) of identified *TLP* genes were used for the

analysis of gene structure in terms of exon-intron organization and intron phases using the GSDS

2.0 server as done in earlier studies [29,30].

2.7. Physicochemical properties of the TLPs

Various physicochemical characteristics such as peptide length, molecular weight (MW) and

isoelectric point (pI) were analyzed using the ExPasy tool, which were further confirmed from

the Ensemble plants and Sequence Manipulation Suite [31-33]. Tools including CELLO v.2.5,

ngLOC, ProtComp9 and WoLF PSORT were used for the prediction of subcellular localization

of TLP proteins [34-37]. Tools such as Phobius and DAS-Tmfilter were used for the prediction

of transmembrane regions [38,39]. The signal peptides were detected using the tools Phobius and

SignalP [38,40]. SMART server was used for the domain analysis, whereas, motifs were

analyzed using MEME v.4.11.4 [41,42].

2.8. Cis-regulatory element analysis

For *cis*-regulatory elements analysis, 1.5 kb upstream genomic sequences from the initiation

codon were retrieved for each *TLP* gene. These promoter sequences were analyzed using the

PLACE software [43]. Identified *cis*-regulatory elements were categorized based on their

functions.

2.9. Expression profiling using RNA-seq data

Expression analysis in various tissues of each crop was done using the high throughput RNA-seq

data retrieved from the URGI database (wheaturgi.versailles.inra.fr/files/RNASeqWheat/) and

Expression ATLAS [44-46]. In *T. aestivum*, the expression data generated in replicates for

various tissue developmental stages, under biotic (fungal pathogen) and abiotic (heat, drought

7

and salt) conditions were used [47-49]. Data generated for three developmental stages of root,

stem, leaf, spike and grain was used to analyze the tissue-specific expression in wheat. Further,

the RNA-seq data (PRJNA243835) generated in triplicates after the 24, 48 and 72 h of

inoculation of *Blumeria graminis* f. sp. *tritici* (Bgt) and *Puccinia striiformis* f. sp. *tritici* (Pst) in

7-d-old seedlings were used for the *TaTLP* expression analysis under biotic stress [47]. Under

abiotic stress conditions, duplicate RNA-seq data (SRP045409) for 1 and 6 h of treatments of

heat (40º C), drought (20% PEG 6000) and a combination of both heat and drought stresses were

used to study the *TaTLP* expression [48]. Root RNA-seq data (SRP062745) available in

triplicates for the treatment of 150 mM NaCl at 6, 12, 24 and 48 h treatment were used to

analyze the expression of *TaTLPs* under salt stress [49]. Trinity package was used to calculate

the expression value in terms of FPKM value [50]. Hierarchical Clustering Explorer 3.5 was

used to generate heat maps of differentially expressed genes, which were clustered using the

Euclidean distance method [51].

2.11. RNA isolation, cDNA synthesis and qRT-PCR

The total RNA of each sample was isolated using the Spectrum™ Plant Total RNA kit (Sigma,

USA). Seedlings (shoot and root) grown under normal conditions were used as a control.

TURBO DNA-free™ Kit (Invitrogen, USA) was used to remove the genomic DNA

contamination. The qualitative and quantitative analysis of RNA was done using agarose gel

electrophoresis and nanodrop quantification. The Superscript III First-Strand Synthesis Super-

mix (Invitrogen, USA) was used to synthesize the cDNA from one microgram of total RNA. A

real-time qRT-PCR was performed with the gene-specific primers of selected genes using SYBR

Green at the 7900 HT Fast Real-Time PCR System (Applied Biosystems) following the method

established in our laboratory [21]. The fold expression change was calculated using the delta-

delta CT method ($2^{-\Delta\Delta CT}$) using the expression of ADP-ribosylation factor (*TaARF*) as an

internal control [52]. All the experiments were carried out in triplicates and expressed as mean ±

SD. A significant difference between the control and treatments was examined by using the two-

tailed student's t-test.

2.12. Cloning and functional characterization

The full-length open reading frame (ORF) of the *TaTLP2-B* gene was amplified from cDNA

using the *TaTLP2-B* forward (5' GTAATGGCTCTTCTTCCTCCTCTGCTTCTG 3') and

TaTLP2-B reverse (5' AATCTGGGCCACACGATCGCCCC 3') primers. The amplified DNA

was cloned into the pJET1.2 cloning vector and sequenced. *TaTLP2-B* gene was re-amplified

from the confirmed clone using the same primers and ligated into the pYES2.1/V5-His-Topo

vector (Invitrogen, USA). The recombinant plasmid (pYES2.1-*TaTLP2-B*) was transformed into

S. cerevisiae (W303) (yeast) cells for further characterization. For control, *lacz* gene containing

pYES2.1 vector (pYES2.1/V5-His/lacZ) was used during the studies. Spot assays using

recombinant yeast cells were performed under various abiotic stress conditions. Recombinant

yeast cells were grown overnight in SD/−ura (with 2% dextrose) medium at 30°C and 200 rpm as

primary culture and further inoculated for secondary culture (1:100 dilutions). As soon as OD_{600}

reached 0.4, 2% galactose was added to induce the expression of recombinant protein in yeast

and kept at 30°C and 200 rpm for 6 h. The OD_{600} was adjusted to 0.4 and an equal volume (500

µl) of each induced culture was further diluted in 10 ml medium. The diluted cultures were

treated with heat (37°C), cold (4°C), drought (30% PEG), combined heat- drought and salt (1 M

NaCl) stresses for 24 h, separately. After the treatments, serial dilutions (10^{0}, 10^{-2}, 10^{-4} and 10^{-6})

were prepared, 5 µl of each dilution was spotted on SD/-ura agar plates, and incubated at 30°C

for 2–3 d [21]. All the experiments were performed in triplicates and results were compared

visually.

2.13. Subcellular localization of the TaTLP2-B

Subcellular localization of TaTLP2-B was analyzed using CaMV35S-driven C-terminal YFP

fusion construct generated by Gateway LR-recombination into binary vector PEG101. Plasmids

were then transformed into *Agrobacterium tumefaciens* GV3101 strain. The cells were

resuspended in a freshly prepared infiltration medium (10 mM $MgCl^2$, 10 mM MES/KOH, pH

5.7 and 150 μM acetosyringone). The resulting constructs were infiltrated onto the abaxial

surface of *Nicotiana benthamiana* leaf and kept at 22°C for 48 h. The YFP fluorescence was

observed using a Leica TCS SP8 (Leica Microsystems, Wetzlar, Germany) laser-scanning

confocal microscope at 514-527 nm.

3. Results

3.1. Identification and chromosomal localization of the TLP genes

Numerous properties of TLPs ascribed to defence and development pathways and lack of such

studies in numerous cereals lead us to perform a comprehensive analysis of the TLP family in

five major cereals. An extensive BLAST search identified 26, 27, 39, 93 and 37 *TLP* genes in *B*.

distachyon, O. sativa, S. *bicolor, T. aestivum* and *Z. mays*, respectively. The genes lacking or

having incomplete thaumatin signature motifs were excluded from the study. Four *TLPs* were

identified as small *TLPs* (sTLPs) in each *B. distachyon, O. sativa* and *Z. mays*, while 10 and 20

sTLPs were detected in *S. bicolor* and *T. aestivum,* respectively, (**supplementary file S1**). In the

case of *T. aestivum*, identified *TLP* genes from the A, B and D sub-genomes formed 32

homeologous groups based on their sequence homology ($\geq$ 90%). All of the identified proteins

exhibited a complete thaumatin signature motif (**supplementary fig. S1**).

The chromosomal localization suggested the scattered distribution of the *TLP* genes at the

majority of chromosomes in each crop (**Fig. 1**). In *B. distachyon,* chromosome 4 consisted of a

maximum of 10 *TLP* (both long and small) genes, while in the case of *O. sativa, S. bicolor, T.*

aestivum and *Z. mays*, the majority of genes (6, 11, 18 and 11) were localized onchromosomes

12, 8, 5A and 1, respectively.

3.2. Multiple sequence alignments

TLP proteins consist of various characteristic and conserved residues, which are important for

their functional activities. Multiple sequence alignments of TLP proteins with a known

thaumatin protein (P02883.2|THM1_THADA) revealed the identification of 16 and 10 conserved

cysteine residues in the majority of long and small TLP proteins of the studied cereal crops,

respectively. Moreover, 7-15 cysteine residues had also been observed in a few TLPs of *O.*

sativa, S. bicolor, T. aestivum and *Z. mays*. The REDDD motif was also found conserved in the

majority of TLP proteins. However, in certain TLP proteins, a few amino acid residues (AAs)

were replaced by other AAs having either similar or different properties. For instance, Arginine

(R) was replaced by Lysine (K) or Asparagine (N). This could be responsible for the differential

evolvement of various characteristic features of TLPs during evolution. The thaumatin signature

motif, which is a characteristic feature of TLPs [6], was found to be highly conserved in all the

identified TLP proteins. Some other regions like FF hydrophobic motif and bottom of acidic cleft

forming amino acids were also found well conserved (**Fig. 2**, **supplementary fig. S1**).

3.3. Evolutionary analyses

3.3.1. Phylogeny

To decipher the evolutionary relationship, a phylogenetic tree was built using the full-length TLP

protein sequences of the five cereals and *A. thaliana*. These were clustered into 11 different

clades based on their phylogenetic relatedness, named as groups I to XI (**Fig. 3**). The highest

number of TLPs were found in group XI, followed by group II and group X, while group IV was

the smallest with only five genes. All the sTLPs were tightly clustered into group XI, which

could be due to their smaller size. Further, the majority of groups consisted of TLPs from all the

five cereals, except groups III, V and VI that lacked members from one or more plant species.

Besides, group IV comprised only three TaTLP and two AtTLP proteins. Besides, the

homeologous TaTLPs of *T. aestivum* were tightly clustered in proximity.

3.3.2. Duplication events investigation

Duplication event (DE) analysis was carried out to understand their roles in the evolution and

expansion of the TLP gene family in the studied crop species. A total of one, four, three, six and

eight DEs were predicted in *B. distachyon, O. sativa, S. bicolor*, *T. aestivum* and *Z mays*,

respectively (**Fig. 1**, **supplementary file S2**). Only tandem duplication events (TDEs) were

found in the case of paralogous genes of *B. distachyon* (*BdTLP1-BdTLP2), O. sativa* (*OsTLP5-

OsTLP7, OsTLP22-OsTLP24, etc.*) and *S. bicolor* (*SbTLP28-SbTLP29, SbTLP31-SbTLP34,

etc.*). However, in the case of *T. aestivum*, five DEs (*TaTLP1-A-TaTLP24-D, TaTLP1-B-

TaTLP24-B, etc.*) were segmental and one was TDE (*TaTLP16-A-TaTLP17-A*). In *Z. mays*,

seven segmental (*ZmTLP1-ZmTLP35, ZmTLP2-ZmTLP34* etc.) and one TDE (*ZmTLP6-

ZmTLP7) were found. All the duplicated gene pairs of each cereal crop were tightly clustered in

proximity in the phylogenetic tree.

3.3.3. Ka/Ks and Tajima's relative rate test

Over the course of evolution, various evolutionary forces and natural pressures affected the

duplicated genes [53]. To understand the evolutionary divergence between the paralogous gene

pairs, the Ka/Ks analysis was carried out. The Ka/Ks ratio of more than one suggests the positive

(non-purifying) and less than one indicates negative (purifying) selection pressure. All the

paralogous genes showed the Ka/Ks value lesser than one, which suggested the negative or

purifying selection on duplicated *TLP* genes (**Table 1**). However, Ka/Ks analysis for *ZmTLP6*

and *ZmTLP7* was not performed due to their 100% similarity. Additionally, the divergence time

of DEs was also calculated using the Ks value and previously described methods [54,55]. The

divergence time of duplicated genes was calculated as 83 million years ago (MYA) in *BdTLPs*,

while the range varied from 52-108, 6-137, 63-99 and 1-63 MYA in *O. sativa, S. bicolor, T.*

aestivum and *Z. mays*, respectively (**Table 1**). Tajima's relative rate test found the insignificant

$\chi 2$ value for all the duplicated gene pairs at P > 0.05 (**Table 2**). The P-value of more than 0.05

depicted their acceptance of the molecular evolutionary clock hypothesis [28].

3.4. Gene architecture, domain and motif analyses

The number of exons varied from one to three in *O. sativa,* while one to four in *B. distachyon, S.*

bicolor and *T. aestivum.* In the case of *Z. mays,* most of the *TLPs* consisted of one to three exons,

while *ZmTLP16, ZmTLP20, ZmTLP25* and *ZmTLP33* exhibited eleven, seven, eight and ten

exons, respectively. Moreover, a total of 44% (98/222) identified *TLPs* were intronless.

13

Intriguingly, all the *sTLPs* were intronless, except *TaTLP6-A* and *TaTLP20-A1*. The intron phase

analysis revealed that the occurrence of a maximum number of introns in phase 1, followed by

phase 2, while the least number of introns were in phase 0 (**supplementary fig. S2**).

The functional nature of a protein depends upon the occurrence of domain composition. All of

the identified cereals' TLPs consisted of a thaumatin domain (PF00314), which confirmed that

they are thaumatin-like proteins. The size of the thaumatin domain ranged from 202-217 AAs

to134-154 AAs in the long and small TLPs, respectively. In addition to the thaumatin domain,

ZmTLP16, ZmTLP20, ZmTLP25 and ZmTLP33 also consisted of a nuclear protein 96 (NUP96)

domain of ~211 AAs at the C-terminus of these proteins (**supplementary file S3**).

Motif investigation revealed the occurrence of 10 highly conserved motifs in the TLP proteins.

Motifs 1-9 were parts of the thaumatin domain, while motif 10 was unknown. Motifs 5, 6 and 8

were the most conserved motifs in TLP proteins, in which the thaumatin signature sequence was

present in motif 8 (**supplementary fig. S2**). The occurrence of conserved motifs across the

cereal species suggested the conserved nature of TLP proteins in related plant species.

3.5. Physicochemical properties

To understand the various important features, several physicochemical properties of identified

TLPs were studied. The TLPs were analyzed for molecular weight (MW), peptide length,

isoelectric point (pI), subcellular localization, transmembrane (TM) helix and signal peptide

(**supplementary file S4**). The average length of the long and small TLPs ranged from284-334

AAs to 175-185 AAs in the studied cereal crops, respectively. Similarly, the average MWs

ranged from ~29-35 kDa to ~17-19 kDa, respectively. However, the average pI ranged from 5.89

14

to 6.95 and from 4.90 to 6.82 for the long and small TLPs, respectively (**Table 3**,

supplementary file S4).

The majority of TLP proteins lacked TM helices, which suggested their soluble/cytoplasmic

nature. However, five TLPs of *B. distachyon* and *O. sativa*, seven of *T. aestivum* and *Z. mays*,

and 10 TLPs of *S. bicolor* consisted of a single TM region, and OsTLP18, TaTLP4-D and

TaTLP5-D comprised of two TM helices, which suggested their membrane-bound nature

(**supplementary file S4**).

A total of 23, 21, 36, 86 and 33 TLPs of *B. distachyon, O. sativa, S. bicolor*, *T. aestivum* and *Z.*

mays, consisted of the N-terminal signal peptide, respectively. Further, the majority of TLP

proteins were predicted to be localized in the extracellular region, while three TLPs from *O.*

sativa (OsTLP10, OsTLP15 and OsTLP16) showed nuclear, and BdTLP16, TaTLP21-A,

ZmTLP20, ZmTLP25 and ZmTLP33 showed plasma membrane localization (**supplementary**

file S4). To validate the subcellular localization, a *TLP* gene (*TaTLP2-B*) was cloned and

expressed with YFP as a translational fusion protein. Analysis of YFP fluorescence of fusion

protein confirmed their extracellular localization (**Fig. 4**). Moreover, we could also see some

fluorescence in the cytoplasmic region. The results suggested that the majority of TaTLP2-B was

localized in the extracellular region, where they could be involved in various defence related

functions (such as anti-fungal etc.). The cytoplasmic localization could be due to their translation

in the cytoplasm, or it might also function inside the cytoplasm.

3.6. Promoter region analyses

Promoter elements are necessary for the regulation of gene expression under different conditions.

Therefore, we performed the *cis*-regulatory analysis of *TLPs* to foresee their regulatory

mechanisms. Based on their putative functions, the identified *cis*-regulatory elements were

segregated into four groups; growth and development, light, hormone and stress-responsive

elements (**supplementary table S1**). The promoter regions of *TLP* genes comprised of several

growth-related elements including TE2F2NTPCNA, EBOX, RYREPEAT4, RAV1AAT,

POLLEN1LEAAT52 etc. In the case of light-responsive elements, GT1CONSENSUS, GATA

box, GBOXLERBCS, BOXII, IBOXCORE were some common *cis*-regulatory elements.

However, under the hormonal responsive category, DPBFCOREDCD3, ARFAT, ARR1,

ERELEE4 and GARE10OSREP1 were abscisic acid, auxin, cytokinin, ethylene and gibberellic

acid-responsive elements, respectively.

3.7. Expression profiling of the TLP genes under different tissues and developmental stages

Expression analysis of genes is a significant way to understand their involvement in various

developmental and physiological processes. Previously, TLPs are reported to be involved in

plant growth and developmental processes [56,57]. Therefore, we performed the expression

analysis of TLPs under various tissues and their developmental stages. A total of 23, 27, 37, 84

and 29 *TLP* genes showed expression in one or more tissue developmental stages in *B.*

distachyon, O. sativa, S. bicolor, T. aestivum and *Z. mays*, respectively (**Fig. 5A-E**). Based on

expression profile, these genes were clustered into 3-5 groups in various crop species. *BdTLPs* of

group 1 were highly expressed in early and emerging inflorescence stages, while *BdTLP9* and

BdTLP12, and *BdTLP20* and *BdTLP23* were upregulated in endosperm and embryo,

respectively. Group 2 genes were highly expressed in the leaf, pistil, embryo and endosperm

tissues. However, group 3 genes were highly expressed at 10 days after pollination (DAP) of

seeds; moreover, *BdTLP8* and *BdTLP18* were upregulated in the pistil, as well (**Fig. 5A**). In *O.*

sativa, the majority of group 1 genes were highly expressed in seed, group 2 in callus, group 3

genes in the shoot, group 4 in inflorescence and group 5 genes in emerging inflorescence and anther (**Fig. 5B**). In *S. bicolor,* group 1 and group 4 genes showed higher expression in root and leaf tissues, respectively. However, groups 2 and 3 *TLP* genes were highly expressed in various reproductive tissues like anther, pistil, endosperm, embryo and flower (**Fig. 5C**).

In *T. aestivum*, the majority of group 1 genes exhibited higher expression in root tissue followed by the shoot. However, most of the groups 2, 3 and 4 *TLP* genes were highly expressed in various developmental stages of spike and grain tissues (**Fig. 5D**).

In the case of *Z. mays,* group 1 genes showed significant expression in unpollinated silk, pericarp and aleurone layer. While, the majority of group 2 genes were specifically upregulated in the root tissues, and group 4 genes in the leaf and internode tissues. Group 3 genes were specifically upregulated in the embryo (**Fig. 5E**). These results suggested the role of *TLP* genes in both vegetative and reproductive tissues development in all the studied cereal crops. However, the higher expression of a few genes in a specific tissue or developmental stages suggested their precise role in particular tissue development.

3.8. Expression profiling of the TaTLP genes under biotic and abiotic stress conditions

Since TLPs belong to the defence-related protein family and are well known for their stress-responsive behaviour. The expression analysis of *TaTLP* genes was also performed under various biotic and abiotic stress conditions to reveal their roles in stress resistance [47-49]. For biotic stress, the available RNA-seq data generated at 24, 48 and 72 h of infestation of two important fungal pathogens, namely, *Blumeria graminis* f. sp. *tritici* (Bgt) and *Puccinia striiformis* f. sp. *tritici* (Pst) was used for expression analysis [47]. In *T. aestivum*, a total of 50 *TaTLP* genes exhibited differential expression (≥2 fold) in these stress conditions, which could

17

be clustered into four groups (**Fig. 6A**). The majority of group 1 *TaTLP* genes were upregulated

at 24 and 48 h of Bgt infestation. However, all the group 3 and 4 *TaTLP* genes were significantly

upregulated at the late (72 h) and early (24 h) periods of Pst infestations, respectively. *TaTLP3-*

A2, *TaTLP12-A,* and *TaTLP32-A* were the most upregulated genes after Bgt infection with 37, 36

and 35 folds, respectively. However, in the case of Pst infestation, *TaTLP10-A* (51 fold up) and

TaTLP27-D (17 fold up) were highly upregulated.

Abiotic stresses are another major threat to plants, which affect various physiological and

biological processes. To understand the putative role of *TaTLPs* under abiotic stress conditions,

expression analysis was carried out under drought (DS), heat (HS) and combined heat and

drought (HD) stress using online available RNA-seq data [48]. A total of 52 *TaTLPs* exhibited

significant differential expression in these stresses, which formed four clusters in the heatmap

(**Fig. 6B**). Almost all the genes of group 1 were downregulated in the DS, HS and HD stresses,

except *TaTLP12-A* that was slightly upregulated in DS treatment. In the case of group 2, the

majority of genes were upregulated at 6 h of DS, HS and HD treatments, while either unaffected

or downregulated at 1 h of treatment. These might be the late responsive *TaTLP* genes. Most of

the group 3 and group 4 *TaTLP* genes were found to be DS and HS responsive, respectively.

Expression analysis of the *TaTLP* genes was also performed at 6, 12, 24 and 48 h of salt stress

using available RNA-seq data [49]. Out of 93 genes, 63 *TaTLP* genes were found to be

differentially expressed under salt stress, which were clustered into 4 groups, based on their

expression patterns (**Fig. 6C**). All the group 1 genes were downregulated at all the stages of salt

stress treatment. In contrast, all the group 3 genes were found significantly upregulated at 48 h of

treatment. However, group 2 genes were upregulated at the early stage (6 h) of treatment, while

they get normalized at later stages. The results suggested that the group 2 and group 3 genes are

early and late responsive *TaTLP* genes, respectively.

To validate the RNA-seq expression data, qRT-PCR analysis of eight *TaTLP* genes was

performed using gene-specific primers at various hours of heat, drought and salt treatments (**Fig.**

7A-P, supplementary file S5). Overall the result was in agreement with the expression observed

using RNA-seq data, with a few exceptions. In the case of heat stress, *TaTLP2-B*, *TaTLP7-D*,

TaTLP14-B1 and *TaTLP25-B* were more upregulated at 1 h, while other genes were highly

upregulated at 6 h of treatment. In the case of drought stress, *TaTLP-7D* was exclusively more

upregulated at 1 h, whereas other genes were upregulated at 6 h. However, *TaTLP2-B* and

TaTLP10-D were downregulated at DS 1 h. The majority of genes were highly upregulated at 1 h

HD treatment. In the case of salt stress, seven genes were highly upregulated at 12 h of salt

stress, except *TaTLP14-B1* that showed maximum expression at 48 h. The results further

confirmed that a few *TaTLP* genes are early while others are late responsive.

3.10. Comparative expression profiling of the TaTLP paralogous genes

To understand the function of duplicated genes, a comparative expression profiling of each

paralogous pair was performed. Generally, based on expression pattern, duplicated genes could

categorize into the retention of function, pseudo-functionalization and neo-functionalization. Out

of six paralogous gene pairs, three pairs (*TaTLP1-A-TaTLP24-D, TaTLP1-B-TaTLP24-B,*

TaTLP15-A-TaTLP18-B) showed a similar trend of expression, suggested retention of function in

these duplicated genes (**Fig. 8A-C**). Two duplicated pairs (*TaTLP12-D-TaTLP13-D* and

TaTLP16-A-TaTLP17-A) showed insignificant expression of one of the gene suggested pseudo-

functionalization (**Fig. 8D, E**). Retention of function in the majority of duplicated genes and the

19

absence of neo-functionalization suggested functional conservation in *TLP* genes during

evolution.

3.11. Cloning and functional characterization of the TaTLP2-B in Saccharomyces cerevisiae

In our study, since the *TaTLP2-B* exhibited significant differential expression under abiotic stress

conditions, it was selected for functional characterization. *TaTLP2-B* open reading frame was

amplified using end primers and cloned into pYES2.1 yeast expression vector. The *LacZ* gene-

containing vector was used as control (**Fig. 9A-F**). In the spot assay, we observed similar growth

of both control (LacZ) and TaTLP2-B expressing yeast cells in the controlled conditions (**Fig.**

9A). However, in the case of cold, drought, heat, combined heat drought and salt stress

conditions, TaTLP2-B recombinant cells exhibited higher growth than the control cells (**Fig. 9B-**

F). The results revealed that the overexpression of the TaTLP2-B provided abiotic stress

tolerance to the recombinant yeast cells. This gene can be used for the development of abiotic

stress-resistant transgenic crops in future studies.

4. Discussion

TLPs are important proteins involved in the tissue development and stress resistance pathways of

plants [6]. Therefore, we have studied the TLPs in five major cereals including *B. distachyon*, *O.*

sativa, *S. bicolor*, *T. aestivum* and *Z. mays*. In the present study, we have identified a total of 26,

27, 39, 93 and 37 *TLPs* in the genome of *B. distachyon*, *O. sativa*, *S. bicolor*, *T. aestivum* and *Z.*

mays, respectively. Variable numbers of the *TLP* genes in different plants have been reported in

earlier studies. For instance, 24, 33 and 49 *TLP* genes have been reported in diploid *A. thaliana*,

Vitis vinifera and *Populus trichocarpa*, respectively [7,58]. In the case of *O. sativa* and *Z. mays*,

49 *TLP* genes in each are reported in the earlier study [7], because probably they had not

removed the genes with incomplete thaumatin signature motifs, as done in the present study. The

occurrence of the highest number of *TLP* genes in *T. aestivum* could be attributed to their

complex allohexaploid genome configuration (AABBDD) [20]. Further, differential expansion

of the *TLP* genes in various plant species might be attributed to the number of duplication events.

The duplication events are responsible for the expansion of gene families and these are the major

forces to attain neo-functionality by causing genetic variability [59]. In DEs analysis, our results

suggested that tandem DEs are a major factor in *B. distachyon*, *O. sativa* and *S. bicolor* and

segmental DEs in *T. aestivum* and *Z. mays* behind the gene expansion of TLP family. The role of

tandem and segmental duplications in the evolution of TLP gene family has been earlier reported

in various plant species including *A. thaliana*, *O. sativa*, and *Z. mays* [7]. The higher number of

paralogous genes in *T. aestivum* and *Z. mays* might be attributed to their larger genome size and

the presence of more transposable elements [53,60].

Furthermore, the divergence time of monocots from eudicots was estimated as ~170-235 Mya,

which were further diverged into grasses around ~77 Mya [60,61]. Therefore, the paralogous

TLP genes in *B. distachyon* might have evolved before the divergence of grasses. The main

duplication incidence in *S. bicolor* and *O. sativa* genome was estimated at around ~70 Mya [62].

However, our results indicated that the majority of paralogous *TLP* genes of *O. sativa* and *S.*

bicolor are evolved after the main DE occurrence. In *Z. mays*, paralogous *TLP* genes were

probably formed close to the divergence time of maize and sorghum, i.e. ~11-28 Mya [62].

However, in the case of *T. aestivum*, paralogous *TLP* genes were probably evolved earlier than

21

the hybridization event of A, B and D subgenomes [63]. However, two duplicated pairs, having

NUP 96 domain showed their recent incidence of DEs. Duplicated genes face various

environmental forces and natural pressures during the process of evolution [53]. Our results

suggested the purifying selection of *TLP* genes as a major force of selection, as reported in the

various other plant species [7,64]. Moreover, positive selection has also been reported for *TLP*

genes in poplar, which suggested varied natural selection procedures during TLPs' evolution [8].

Since the TLPs are paraphyletic in origin, they are supposed to be derived from multiple

ancestral genes [65]. Clustering of identified TLPs in multiple clades is in agreement with the

paraphyletic nature of origin. Further, a variable number of TLPs in each clade is due to the

differential duplication of *TLP* genes. Similarly, the uneven distribution and paraphyletic nature

of TLPs have also been reported by previous studies [7,64]. The tight clustering of homeologous

and paralogous genes could be due to the high sequence homology among the respective

sequences.

The organization of introns and exons of *TLPs* have been reported to be in the range of one to ten

exons [6,7,9,53]. On similar patterns, our results suggest comparable findings with one to four

exons in most of the TLPs. However, most of the monocots' TLPs have intron-less nature, which

was also found in our analysis [7]. Additionally, our results with respect to the protein lengths,

molecular weights and isoelectric points of long and small TLP proteins, found in accordance

with previously studied TLPs [6,7,9]. In our analysis, the TLP2-B was found to be localized in

extracellular region. Similarly, the extracellular localization of a wheat TLP (TaLr19TLP1) is

also reported in an earlier study [66].

In addition, the presence of thaumatin-like domain in all the TLP proteins, make our studies

similar to previously reported findings. However, four ZmTLPs (ZmTLP16, ZmTLP20,

22

ZmTLP25, and ZmTLP33) have an additional NUP 96 domain. The NUP 96 domain is reported

to be involved in plant development and stress resistance against pathogens [67]. Various abiotic

and biotic stress-responsive elements were also found in most of the studied *TLPs*, for instance,

DRE2COREZMRAB17, ASF1, BIHD1, CBFHV, CGG-box, GCCCORE,

LTRECOREATCOR15, MYB, MYC, WRKY1, W-box, T/G-box etc. Various plant hormone-

related and stress-related *cis*-regulatory elements have also been reported in four cotton species

[56]. Moreover, in previous studies, the elements like ERE, W-box, WRKY, were found to be

involved in plant growth and stress resistance [68-70]. The occurrence of numerous *cis*-

regulatory elements suggested diverse functions of *TLP* genes in plants. Further, the expression

profiling of *TLPs* under various tissues and developmental stages also advocated the putative

involvement of these genes in development of vegetative and reproductive tissues. Similar

expression trend in various plant tissues and their role in reproductive tissues like flowers and

seeds has also been reported in earlier studies, for instance, *TLP* genes in *Gossypium hirsutum*,

also showed their varied expression pattern in vegetative as well as in reproductive organs [18,

56, 71]. The role of *TLP* genes in fungal resistance has been reported in earlier studies against

numerous pathogens including *Fusarium* sp. [72], *Microdochium nivale* [73], *Rhizoctonia solani*

[74] and *Verticillium dahlia* [56] in wheat and other plant species. The differential expression of

TaTLP genes in response to Pst and Bgt infestation further established their roles in fungal stress

responses. Moreover, similar to our finding's differential expression and upregulation of *TLP*

genes at various hours of drought, salt and cold stress treatment has also been reported in other

plants like *Brassica rapa* and *Gossypium* sp. [56,75]. Further, the over-expression of *TLP* genes

provided increased tolerance against various abiotic stresses in tobacco, cotton, Arabidopsis etc.

[11,56,76]. In a recently conducted study, the important role of a cotton *TLP* (*GhTLP19*) gene

509 was studied in combating drought stress [56]. Moreover, the GbTLP1 and ObTLP1 of *G.*

510 *barbadense* and *Ocimum basilicum*, respectively, were found effective against drought and salt

511 stresses [57,76]. Collectively, current findings depict that the TLP proteins could be utilized to

512 develop the stress resistance transgenic cereal crops, which will be a boost for the agriculture.

5. Conclusions

Thaumatin-like proteins (TLPs), a part of the PR5 protein family, are known to be involved in

various biotic and abiotic stresses. The curiosity to unravel the various possibilities led us to a

detailed characterization of a total of 222 *TLP* genes in five cereal crops. Phylogenetic analysis

revealed the paraphyletic origin of TLPs in cereals, while the occurrence of duplication events

suggested the role of paralogous genes in the expansion of the TLP gene family. Gene expression

analysis using RNA-seq-data and qRT-PCR suggested the important roles of *TLP* genes in plant

growth and development and abiotic and biotic stress responses. Significant tolerance in the

TaTLP2-B expressing yeast cells further established their role in stress response. Overall the

study revealed that the *TLP* genes can be used for stress-resistant transgenic crop development.

Since the identified genes are directly derived from the food crops, they would be easily de-

regulated from the regulatory authority of the transgenic crops. Further, the current study will

facilitate the detailed functional characterization of *TLP* genes of cereal crops in future studies.

Competing interests

The authors declare that they have no competing interests

Author's contribution

SKU conceived the idea. AS and SKU designed the experiments. AS, HS and RR performed the experiments. AS, HS and AP analyzed the data. AS, HS and SKU wrote the manuscript. All the authors have read and finalized the manuscript.

Acknowledgements

The authors are grateful to the Panjab University, Chandigarh, and National Institute of Plant Genome Research, New Delhi, India for research facilities, Ensembl Plants, URGI and NCBI for data availability. AS and HS is grateful to CSIR for the senior research fellowship. HS is also thankful to the IKGPTU, Jalandhar for the Ph.D registration. SKU is grateful to the Council of Scientific and Industrial Research (CSIR) and the Department of Science and Technology (DST), Government of India for a research grant (No. 38(1489)/19/EMR-II) and partial financial support under the Promotion of University Research and Scientific Excellence (PURSE) grant scheme, respectively.

References

545   1.   E. Kombrink, I.E. Somssich, Pathogenesis-Related Proteins and Plant Defense, in: Plant

546   Relationships, Springer Berlin Heidelberg, Berlin, Heidelberg, 1997: pp. 107–128.

547   https://doi.org/10.1007/978-3-662-10370-8_7.

548   2.   P. Chadha, R.H. Das, A pathogenesis-related protein, AhPR10 from peanut: an insight of

549   its mode of antifungal activity, Planta. 225 (2006) 213–222.

550   https://doi.org/10.1007/s00425-006-0344-7.

551   3.   L.C. van Loon, M. Rep, C.M.J. Pieterse, Significance of inducible defense-related

552   proteins in infected plants, Annu. Rev. Phytopathol. 44 (2006) 135–162.

553   https://doi.org/10.1146/annurev.phyto.44.070505.143425.

554   4.   R. Velazhahan, S.K. Datta, S. Muthukrishnan, The PR-5 family: Thaumatin-like proteins,

555   in: Pathog. Proteins Plants, CRC Press, 1999: pp. 107–129.

556   https://doi.org/10.1201/9781420049299.

557   5.   H. van der Wel, K. Loeve, Isolation and characterization of thaumatin I and II, the sweet-

558   tasting proteins from *Thaumatococcus daniellii* Benth., Eur. J. Biochem. 31 (1972) 221–

559   5. https://doi.org/10.1111/j.1432-1033.1972.tb02522.x.

560   6.   J.-J. Liu, R. Sturrock, A.K.M. Ekramoddoullah, The superfamily of thaumatin-like

561   proteins: its origin, evolution, and expression towards biological function, Plant Cell Rep.

562   29 (2010) 419–436. https://doi.org/10.1007/s00299-010-0826-8.

563   7.   J. Cao, Y. Lv, Z. Hou, X. Li, L. Ding, Expansion and evolution of thaumatin-like protein

564   (TLP) gene family in six plants, Plant Growth Regul. 79 (2016) 299–307.

565   https://doi.org/10.1007/s10725-015-0134-y.

8. J.-J. Liu, A. Zamani, A.K.M. Ekramoddoullah, Expression profiling of a complex

thaumatin-like protein family in western white pine, Planta. 231 (2010) 637–651.

https://doi.org/10.1007/s00425-009-1068-2.

9. B. Petre, I. Major, N. Rouhier, S. Duplessis, Genome-wide analysis of eukaryote

thaumatin-like proteins (TLPs) with an emphasis on poplar, BMC Plant Biol. 11 (2011)

33. https://doi.org/10.1186/1471-2229-11-33.

10. R. Ghosh, C. Chakrabarti, Crystal structure analysis of NP24-I: a thaumatin-like protein,

Planta. 228 (2008) 883–890. https://doi.org/10.1007/s00425-008-0790-5.

11. M. V. Rajam, N. Chandola, P. Saiprasad Goud, D. Singh, V. Kashyap, M.L. Choudhary,

D. Sihachakr, Thaumatin gene confers resistance to fungal pathogens as well as tolerance

to abiotic stresses in transgenic tobacco plants, Biol. Plant. 51 (2007) 135–141.

https://doi.org/10.1007/s10535-007-0026-8.

12. K. Datta, R. Velazhahan, N. Oliva, I. Ona, T. Mew, G.S. Khush, S. Muthukrishnan, S.K.

Datta, Over-expression of the cloned rice thaumatin-like protein (*PR-5*) gene in

transgenic rice plants enhances environmental friendly resistance to *Rhizoctonia solani*

causing sheath blight disease, Theor. Appl. Genet. 98 (1999) 1138–1145.

https://doi.org/10.1007/s001220051178.

13. X. Wang, C. Tang, L. Deng, G. Cai, X. Liu, B. Liu, Q. Han, H. Buchenauer, G. Wei, D.

Han, L. Huang, Z. Kang, Characterization of a pathogenesis-related thaumatin-like

protein gene *TaPR5* from wheat induced by stripe rust fungus, Physiol. Plant. 139 (2010)

27–38. https://doi.org/10.1111/j.1399-3054.2009.01338.x.

14. L.R. Abad, M.P. D'Urzo, D. Liu, M.L. Narasimhan, M. Reuveni, J.K. Zhu, X. Niu, N.K.

Singh, P.M. Hasegawa, R.A. Bressan, Antifungal activity of tobacco osmotin has

27

specificity and involves plasma membrane permeabilization, Plant Sci. 118 (1996) 11–

23. https://doi.org/10.1016/0168-9452(96)04420-2.

15. Y. Zhang, D.S. Shih, Isolation of an osmotin-like protein gene from strawberry and

analysis of the response of this gene to abiotic stresses, J. Plant Physiol. 164 (2007) 68–

77. https://doi.org/10.1016/j.jplph.2006.02.002.

16. A.D. Neale, J.A. Wahleithner, M. Lund, H.T. Bonnett, A. Kelly, D.R. Meeks-Wagner,

W.J. Peacock, E.S. Dennis, chitinase, [beta]-1,3-glucanase, osmotin, and extensin are

expressed in tobacco explants during flower formation, PLANT CELL ONLINE. 2

(1990) 673–684. https://doi.org/10.1105/tpc.2.7.673.

17. R.A. Salzman, I. Tikhonova, B.P. Bordelon, P.M. Hasegawa, R.A. Bressan, Coordinate

accumulation of antifungal proteins and hexoses constitutes a developmentally controlled

defense response during fruit ripening in grape., Plant Physiol. 117 (1998) 465–72.

http://www.ncbi.nlm.nih.gov/pubmed/9625699.

18. P.J. Seo, A.-K. Lee, F. Xiang, C.-M. Park, Molecular and functional profiling of

Arabidopsis pathogenesis-related genes: Insights into their roles in salt response of seed

germination, Plant Cell Physiol. 49 (2008) 334–344. https://doi.org/10.1093/pcp/pcn011.

19. P. Lei, X. Wei, R. Gao, F. Huo, X. Nie, W. Tong, W. Song, Genome-wide identification

of PYL gene family in wheat: Evolution, expression and 3D structure analysis,

Genomics. 113 (2021) 854–866. https://doi.org/10.1016/j.ygeno.2020.12.017.

20. A. Sharma, Shumayla, S. Tyagi, A. Alok, K. Singh, S.K. Upadhyay, Thaumatin-like

protein kinases: Molecular characterization and transcriptional profiling in five cereal

crops, Plant Sci. 290 (2020). https://doi.org/10.1016/j.plantsci.2019.110317.

21. Shumayla, S. Tyagi, A. Sharma, K. Singh, S.K. Upadhyay, Genomic dissection and
transcriptional profiling of Cysteine-rich receptor-like kinases in five cereals and
functional characterization of *TaCRK68-A*, Int. J. Biol. Macromol. 134 (2019) 316–329.
https://doi.org/10.1016/j.ijbiomac.2019.05.016.

22. R.C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high
throughput, Nucleic Acids Res. 32 (2004) 1792–1797.
https://doi.org/10.1093/nar/gkh340.

23. F. Corpet, Multiple sequence alignment with hierarchical clustering., Nucleic Acids Res.
16 (1988) 10881–90. http://www.ncbi.nlm.nih.gov/pubmed/2849754 (accessed May 27,
2017).

24. S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, MEGA X: Molecular Evolutionary
Genetics Analysis across Computing Platforms, (n.d.).
https://doi.org/10.1093/molbev/msy096.

25. M. Suyama, D. Torrents, P. Bork, PAL2NAL: Robust conversion of protein sequence
alignments into the corresponding codon alignments, Nucleic Acids Res. 34 (2006).
https://doi.org/10.1093/nar/gkl315.

26. M. Nagaraju, P.S. Reddy, S.A. Kumar, A. Kumar, G. Rajasheker, D.M. Rao, P.B. Kavi
Kishor, Genome-wide identification and transcriptional profiling of small heat shock
protein gene family under diverse abiotic stress conditions in *Sorghum bicolor* (L.), Int. J.
Biol. Macromol. 142 (2020) 822–834. https://doi.org/10.1016/j.ijbiomac.2019.10.023.

27. B.S. Gaut, B.R. Morton, B.C. McCaig, M.T. Clegg, Substitution rate comparisons
between grasses and palms: synonymous rate differences at the nuclear gene Adh parallel

rate differences at the plastid gene rbcL., Proc. Natl. Acad. Sci. U. S. A. 93 (1996) 10274.

https://doi.org/10.1073/PNAS.93.19.10274.

28. F. Tajima, Simple methods for testing the molecular evolutionary clock hypothesis.,

Genetics. 135 (1993).

29. B. Hu, J. Jin, A.-Y. Guo, H. Zhang, J. Luo, G. Gao, GSDS 2.0: an upgraded gene feature

visualization   server.,   Oxford   University   Press,   2015.

https://doi.org/10.1093/bioinformatics/btu817.

30. H. Sharma, M. Taneja, S.K. Upadhyay, Identification, characterization and expression

profiling of cation-proton antiporter superfamily in *Triticum aestivum* L. and functional

analysis   of   *TaNHX4-B*,   Genomics.   112   (2020)   356–370.

https://doi.org/10.1016/j.ygeno.2019.02.015.

31. E. Gasteiger, C. Hoogland, A. Gattiker, S. Duvaud, M.R. Wilkins, R.D. Appel, A.

Bairoch, Protein Identification and Analysis Tools on the ExPASy Server, in: Proteomics

Protoc.   Handb.,   Humana   Press,   Totowa,   NJ,   2005:   pp.   571–607.

https://doi.org/10.1385/1-59259-890-0:571.

32. T. Nussbaumer, M.M. Martis, S.K. Roessner, M. Pfeifer, K.C. Bader, S. Sharma, H.

Gundlach, M. Spannagl, MIPS PlantsDB: a database framework for comparative plant

genome   research.,   Nucleic   Acids   Res.   41   (2013)   D1144-51.

https://doi.org/10.1093/nar/gks1153.

33. P. Stothard, The sequence manipulation suite: JavaScript programs for analyzing and

formatting   protein   and   DNA   sequences.,   Biotechniques.   28   (2000)   1102,   1104.

https://doi.org/10.2144/00286ir01.

34. C.-S. Yu, Y.-C. Chen, C.-H. Lu, J.-K. Hwang, Prediction of protein subcellular localization, Proteins Struct. Funct. Bioinforma. 64 (2006) 643–651. https://doi.org/10.1002/prot.21018.

35. B.R. King, C. Guda, ngLOC: an n-gram-based Bayesian method for estimating the subcellular proteomes of eukaryotes, Genome Biol. 8 (2007) R68. https://doi.org/10.1186/gb-2007-8-5-r68.

36. O. Emanuelsson, H. Nielsen, S. Brunak, G. Von Heijne, Predicting subcellular localization of proteins based on their N-terminal amino acid sequence, J. Mol. Biol. 300 (2000) 1005–1016. https://doi.org/10.1006/jmbi.2000.3903.

37. P. Horton, K.-J. Park, T. Obayashi, N. Fujita, H. Harada, C.J. Adams-Collier, K. Nakai, WoLF PSORT: protein localization predictor, Nucleic Acids Res. 35 (2007) W585–W587. https://doi.org/10.1093/nar/gkm259.

38. L. Käll, A. Krogh, E.L.. Sonnhammer, A combined transmembrane topology and signal peptide prediction method, J. Mol. Biol. 338 (2004) 1027–1036. https://doi.org/10.1016/j.jmb.2004.03.016.

39. M. Cserzo, F. Eisenhaber, B. Eisenhaber, I. Simon, TM or not TM: transmembrane protein prediction with low false positive rate using DAS-TMfilter, Bioinformatics. 20 (2004) 136–137. https://doi.org/10.1093/bioinformatics/btg394.

40. T.N. Petersen, S. Brunak, G. von Heijne, H. Nielsen, SignalP 4.0: discriminating signal peptides from transmembrane regions, Nat. Methods. 8 (2011) 785–786. https://doi.org/10.1038/nmeth.1701.

41. I. Letunic, T. Doerks, P. Bork, SMART: recent updates, new developments and status in 2015., Nucleic Acids Res. 43 (2015) D257-60. https://doi.org/10.1093/nar/gku949.

42. T.L. Bailey, M. Boden, F.A. Buske, M. Frith, C.E. Grant, L. Clementi, J. Ren, W.W. Li,
W.S. Noble, MEME SUITE: tools for motif discovery and searching, 2009.
https://doi.org/10.1093/nar/gkp335.

43. K. Higo, Y. Ugawa, M. Iwamoto, H. Higo, PLACE: a database of plant cis -acting
regulatory DNA elements, Nucleic Acids Res. 26 (1998) 358–359.

44. F. Choulet, A. Alberti, S. Theil, N.M. Glover, V. Barbe, J. Daron, L. Pingault, P.
Sourdille, A. Couloux, E. Paux, Structural and Functional Partitioning of Bread Wheat
Chromosome 3B, (n.d.). https://doi.org/10.1126/science.1249721ï.

45. L. Pingault, F. Choulet, A. Alberti, N. Glover, P. Wincker, C. Feuillet, E. Paux, Deep
transcriptome sequencing provides new insights into the structural and functional
organization of the wheat genome, Genome Biol. 16 (2015) 29.
https://doi.org/10.1186/s13059-015-0601-9.

46. I. Papatheodorou, N.A. Fonseca, M. Keays, Y.A. Tang, E. Barrera, W. Bazant, M. Burke,
A. Füllgrabe, A.M.P. Fuentes, N. George, L. Huerta, S. Koskinen, S. Mohammed, M.
Geniza, J. Preece, P. Jaiswal, A.F. Jarnuczak, W. Huber, O. Stegle, J.A. Vizcaino, A.
Brazma, R. Petryszak, Expression Atlas: Gene and protein expression across multiple
studies and organisms, Nucleic Acids Res. 46 (2018) D246–D251.
https://doi.org/10.1093/nar/gkx1158.

47. H. Zhang, Y. Yang, C. Wang, M. Liu, H. Li, Y. Fu, Y. Wang, Y. Nie, X. Liu, W. Ji,
Large-scale transcriptome comparison reveals distinct gene activations in wheat
responding to stripe rust and powdery mildew, BMC Genomics. 15 (2014) 898.
https://doi.org/10.1186/1471-2164-15-898.

48. Z. Liu, M. Xin, J. Qin, H. Peng, Z. Ni, Y. Yao, Q. Sun, Temporal transcriptome profiling reveals expression partitioning of homeologous genes contributing to heat and drought acclimation in wheat (*Triticum aestivum* L.), BMC Plant Biol. 15 (2015) 152. https://doi.org/10.1186/s12870-015-0511-8.

49. Y. Zhang, Z. Liu, A.A. Khan, Q. Lin, Y. Han, P. Mu, Y. Liu, H. Zhang, L. Li, X. Meng, Z. Ni, M. Xin, Expression partitioning of homeologs and tandem duplications contribute to salt tolerance in wheat (*Triticum aestivum* L.)., Sci. Rep. 6 (2016) 21476. https://doi.org/10.1038/srep21476.

50. B.J. Haas, A. Papanicolaou, M. Yassour, M. Grabherr, P.D. Blood, J. Bowden, M.B. Couger, D. Eccles, B. Li, M. Lieber, M.D. MacManes, M. Ott, J. Orvis, N. Pochet, F. Strozzi, N. Weeks, R. Westerman, T. William, C.N. Dewey, R. Henschel, R.D. LeDuc, N. Friedman, A. Regev, De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis, Nat. Protoc. 8 (2013) 1494–1512. https://doi.org/10.1038/nprot.2013.084.

51. J. Seo, H. Gordish-Dressman, E.P. Hoffman, An interactive power analysis tool for microarray hypothesis testing and generation, Bioinformatics. 22 (2006) 808–814. https://doi.org/10.1093/bioinformatics/btk052.

52. K.J. Livak, T.D. Schmittgen, Analysis of relative gene expression data using Real-Time quantitative PCR and the 2−ΔΔCT method, Methods. 25 (2001) 402–408. https://doi.org/10.1006/meth.2001.1262.

53. H. Ellegren, Comparative genomics and the study of evolution by natural selection, Mol. Ecol. 17 (2008) 4586–4596. https://doi.org/10.1111/j.1365-294X.2008.03954.x.

54. S. Huang, A. Sirikhachornkit, X. Su, J. Faris, B. Gill, R. Haselkorn, P. Gornicki, Genes encoding plastid acetyl-CoA carboxylase and 3-phosphoglycerate kinase of the Triticum/Aegilops complex and the evolutionary history of polyploid wheat, Proc. Natl. Acad. Sci. U. S. A. 99 (2002) 8133–8138. https://doi.org/10.1073/pnas.072223799.

55. M. Sharma, A. Singh, A. Shankar, A. Pandey, V. Baranwal, S. Kapoor, A.K. Tyagi, G.K. Pandey, Comprehensive expression analysis of rice armadillo gene family during abiotic stress and development, DNA Res. 21 (2014) 267–283. https://doi.org/10.1093/dnares/dst056.

56. Z. Li, X. Wang, Y. Cui, K. Qiao, L. Zhu, S. Fan, Q. Ma, Comprehensive genome-wide analysis of thaumatin-like gene family in four cotton species and functional identification of ghtlp19 involved in regulating tolerance to verticillium dahlia and drought, Front. Plant Sci. 11 (2020) 1597. https://doi.org/10.3389/fpls.2020.575015.

57. M.F.H. Munis, L. Tu, F. Deng, J. Tan, L. Xu, S. Xu, L. Long, X. Zhang, A thaumatin-like protein gene involved in cotton fiber secondary cell wall development enhances resistance against *Verticillium dahliae* and other stresses in transgenic tobacco, Biochem. Biophys. Res. Commun. 393 (2010) 38–44. https://doi.org/10.1016/j.bbrc.2010.01.069.

58. X. Yan, H. Qiao, X. Zhang, C. Guo, M. Wang, Y. Wang, X. Wang, Analysis of the grape (*Vitis vinifera* L.) thaumatin-like protein (*TLP*) gene family and demonstration that *TLP29* contributes to disease resistance, Sci. Rep. 7 (2017) 4269. https://doi.org/10.1038/s41598-017-04105-w.

59. R. Appels, K. Eversole, N. Stein, C. Feuillet, B. Keller, J. Rogers, et al., Shifting the limits in wheat research and breeding using a fully annotated reference genome, Science (80-.). 361 (2018) eaar7191. https://doi.org/10.1126/science.aar7191.

60. B.S. Gaut, Evolutionary dynamics of grass genomes, New Phytol. 154 (2002) 15–28.

https://doi.org/10.1046/j.1469-8137.2002.00352.x.

61. Y.W. Yang, K.N. Lai, P.Y. Tai, W.H. Li, Rates of nucleotide substitution in angiosperm

mitochondrial DNA sequences and dates of divergence between Brassica and other

angiosperm   lineages,   J.   Mol.   Evol.   48   (1999)   597–604.

https://doi.org/10.1007/PL00006502.

62. A.H. Paterson, J.E. Bowers, B.A. Chapman, Ancient polyploidization predating divergence

of the cereals, and its consequences for comparative genomics, Proc. Natl. Acad. Sci. U. S.

A. 101 (2004) 9903–9908. https://doi.org/10.1073/pnas.0307901101.

63. T. Marcussen, S.R. Sandve, L. Heier, M. Spannagl, M. Pfeifer, K.S. Jakobsen, et al.,

Ancient hybridizations among the ancestral genomes of bread wheat, Science (80-.). 345

(2014) 1250092–1250092. https://doi.org/10.1126/science.1250092.

64. Y. Liu, J. Cui, X. Zhou, Y. Luan, F. Luan, Genome-wide identification, characterization

and expression analysis of the TLP gene family in melon (*Cucumis melo* L.), Genomics.

112 (2020) 2499–2509. https://doi.org/10.1016/j.ygeno.2020.02.001.

65. R.G. Shatters, L.M. Boykin, S.L. Lapointe, W.B. Hunter, A.A. Weathersbee, Phylogenetic

and structural relationships of the *PR5* Gene family reveal an ancient multigene family

conserved   in   plants   and   select   animal   taxa,   J.   Mol.   Evol.   63   (2006)   12–29.

https://doi.org/10.1007/s00239-005-0053-z.

66. J. Zhang, F. Wang, F. Liang, Y. Zhang, L. Ma, H. Wang, D. Liu, Functional analysis of a

pathogenesis-related thaumatin-like protein gene *TaLr35PR5* from wheat induced by leaf

rust fungus, BMC Plant Biol. 18 (2018) 1–11. https://doi.org/10.1186/s12870-018-1297-2.

67. Y. Zhang, X. Li, A putative nucleoporin 96 is required for both basal defense and constitutive resistance responses mediated by suppressor of npr1-1, constitutive 1, Plant Cell. 17 (2005) 1306–1316. https://doi.org/10.1105/tpc.104.029926.

68. K. Hiroyuki and R. Terauchi, Regulation of expression of rice thaumatin-like protein: inducibility by elicitor requires promoter W-box elements, Plant Cell Rep. 27 (2008) 1521–1528. doi: 10.1007/s00299-008-0536-7

69. P.J. Rushton, I.E. Somssich, P. Ringler, Q.J. Shen, WRKY transcription factors, Trends Plant Sci. 15 (2010) 247–258. https://doi.org/10.1016/j.tplants.2010.02.006.

70. J.Y. Gou, L.J. Wang, S.P. Chen, W.L. Hu, X.Y. Chen, Gene expression and metabolite profiles of cotton fiber during cell elongation and secondary cell wall synthesis, Cell Res. 17 (2007) 422–434. https://doi.org/10.1038/sj.cr.7310150.

71. B.R. Fils-Lycaon, P.A. Wiersma, K.C. Eastwell, P. Sautiere, A cherry protein and its gene, abundantly expressed in ripening fruit, have been identified as thaumatin-like., Plant Physiol. 111 (1996) 269–73. http://www.ncbi.nlm.nih.gov/pubmed/8685266.

72. C.A. Mackintosh, J. Lewis, L.E. Radmer, S. Shin, S.J. Heinen, L.A. Smith, M.N. Wyckoff, R. Dill-Macky, C.K. Evans, S. Kravchenko, G.D. Baldridge, R.J. Zeyen, G.J. Muehlbauer, Overexpression of defense response genes in transgenic wheat enhances resistance to Fusarium head blight, Plant Cell Rep. 26 (2007) 479–488. https://doi.org/10.1007/s00299-006-0265-8.

73. C. Kuwabara, D. Takezawa, T. Shimada, T. Hamada, S. Fujikawa, K. Arakawa, Abscisic acid- and cold-induced thaumatin-like protein in winter wheat has an antifungal activity against snow mould, *Microdochium nivale*, Physiol. Plant. 115 (2002) 101–110. https://doi.org/10.1034/j.1399-3054.2002.1150112.x.

74. G. Naseri, M.M. Sohani, A. Pourmassalehgou, S. Allahi, In planta transformation of rice

(Oryza sativa) using thaumatin-like protein gene for enhancing resistance to sheath blight,

African J. Biotechnol. 11 (2012) 7885–7893. https://doi.org/10.5897/AJB11.3331.

75. N.U. Ahmed, J.-I. Park, H.-J. Jung, K.-K. Kang, Y.-P. Lim, Y. Hur, I.-S. Nou, Molecular

characterization of thaumatin family genes related to stresses in *Brassica rapa*, Sci. Hortic.

(Amsterdam). 152 (2013) 26–34. https://doi.org/10.1016/j.scienta.2013.01.007.

76. R.C. Misra, Sandeep, M. Kamthan, S. Kumar, S. Ghosh, A thaumatin-like protein of

Ocimum basilicum confers tolerance to fungal pathogen and abiotic stress in transgenic

Arabidopsis, Sci. Rep. 6 (2016) 25340. https://doi.org/10.1038/srep25340.

Figure legends

Fig. 1 Chromosomal distribution and paralogous gene analysis of *TLPs* in five cereal crops. Image depicts the scattered localization of *TLP* genes of **(A)** *B. distachyon*, **(B)** *O. sativa*, **(C)** *S. bicolor*, **(D)** *T. aestivum* and **(E)** *Z. mays* at their respective chromosomes. Red lines and blue lines denote the tandem and segmental duplication events, respectively.

Fig. 2 Multiple sequence alignment of selected long and small TLPs of *B. distachyon, O. sativa, S. bicolor, T. aestivum* and *Z. mays* with a known thaumatin protein sequence. Figure shows the alignment of long (OsTLP5, SbTLP2, BdTLP13, ZmTLP13 and TaTLP1-A) and small (BdTLP10, OsTLP25, ZmTLP6, SbTLP29 and TaTLP6-B) TLPs with a known thaumatin (P02883.2|THM1_THADA) protein sequence. The characteristic thaumatin signature motif is shown in the sky blue box. All the conserved cysteine residues were marked with an asterisk. The REDDD motif was marked with a red circle. The FF hydrophobic motif and bottom of acidic cleft forming regions are marked with a green triangle and brown box, respectively. The conserved regions of the thaumatin domain are marked with a black line.

Fig. 3 Phylogenetic tree analysis of *A. thaliana, B. distachyon, O. sativa. S, bicolor, T. aestivum and Z. mays* using full-length peptide of TLPs. The figure shows the phylogenetic tree of TLPs, built by the maximum likelihood approach with 1000 bootstrap values using MEGAX software. The phylogenetic tree displays the clustering of TLPs of six plant species in I-XI different groups, each group is coloured differently. The TLPs from each plant are marked with a different colour.

Fig. 4 Subcellular localization analysis of TaTLP2-B by YFP fusion. The YFP fluorescence depicts the localization of TaTLP2-B protein majorly in the extracellular region.

Fig. 5 Expression analysis of *TLP* genes of *B. distachyon* (*BdTLP*), *O. sativa* (*OsTLP*). *S. bicolor* (*SbTLP*), *T. aestivum* (*TaTLP*) and *Z. mays* (*ZmTLP*) under various tissue and developmental stages. Heatmaps **(A-E)** shows the clustering and expression profiling of *TLP* genes in (**A**) *B. distachyon*, (**B**) *O. sativa*, (**C**) *S. bicolor*, (**D**) *T. aestivum* and (**E**) *Z. mays*. The different clusters are formed based on expression values of *TLP* genes in the respective crop which are represented with different colours. The colour bar depicts the upregulated and downregulated expression with red and green colours, respectively.

Fig. 6 Expression profiling of *TLP* genes of *T. aestivum* under biotic and abiotic stress conditions. Heat maps **(A-C)** shows the clustering and expression profiling of *TaTLP* genes under **(A)** *Blumeria graminis* (Bgt) and *Puccinia striiformis* (Pst), **(B)** heat (HS), drought (DS) and combined heat-drought (HD) and **(C)** salt stress conditions. The clustering of *TaTLP* genes in each heat map is based on their expression pattern in respective stress conditions. The colour bar shows high and low expressions with red and green colours, respectively.

Fig. 7 qRT-PCR analysis of eight *TaTLP* genes. Expression patterns in the form of fold changes have been shown in the bar graphs. Figures **7 A, C, E, G, I, K, M, and O** show expression of *TaTLP2-B, TaTLP5-D, TaTLP7-D, TaTLP9-D, TaTLP10-D, TaTLP14-B1, TaTLP25-B and TaTLP29-A* genes, under HS, DS and HD stress conditions, while figure **7 B, D, F, H, J, L, N, and P** shows the expression pattern of these genes under salt stress, respectively. Significance between the control and treated conditions is carried using a two-tailed student's t-test. The ns, *, ** and *** markings represent the significance at p-value >0.05, <=0.05, <=0.01, <=0.001, respectively.

Fig. 8 Comparative expression analysis of duplicated gene pairs in *T. aestivum*. Graphs **(A-E)**

represents the comparative expression pattern of paralogous *TaTLPs* under various tissue

and developmental stages. Based on their expression trend, *TaTLPs* are categorized into

the retention of function **(A-C)** and pseudo functionalization types **(D, E)**.

Fig. 9. Spot assay of recombinant *Saccharomyces cerevisiae* (yeast) cells under various stress

conditions. Figure **(A)** shows similar growth of both TaTLP2-B (pYES2.1-TaTLP2-B)

and control vector (pYES2.1/V5-His/lacZ) containing yeast cells under control

conditions. Figure **(B-F)** shows the spot assay analysis of *TaTLP2-B* and control vector

containing yeast cells in **(B)** cold (4°C), **(C)** drought (30% PEG), **(D)** heat (37°C), **(E)**

combined heat and drought (37°C and 30% PEG), and **(F)** salt (1M NaCl) stress

conditions, respectively.

Supplementary files

Figure S1. Multiple sequence alignment of 222 TLPs of *B. distachyon, O. sativa, S. bicolor*, *T.*

aestivum and *Z. mays*. Conserved cysteine residues are marked with an asterisk. The

red dot denotes the REDDD motif. The FF hydrophobic motif is highlighted with a

green triangle. The thaumatin signature motif, conserved domain, amino acids forming

the bottom of the acidic cleft are marked with sky blue, red and brown lines,

respectively.

Figure S2. Intron/exon organization and conserved motifs of TLPs of *B. distachyon*, *O. sativa*, *S.*

bicolor, T. aestivum and *Z. mays*.

File S1. List of identified *TLP* genes with their proposed nomenclature in *B. distachyon, O. sativa*

S, bicolor, T. aestivum and *Z. mays*. The Red colour represents the *sTLPs* of each crop.

File S2. List of paralogous *TLP* genes in *B. distachyon*, *O. sativa*, *S. bicolor*, *T. aestivum* and *Z.*

mays.

File S3. Domain organization of TLP proteins of *B. distachyon*, *O. sativa*, *S. bicolor*, *T. aestivum*

and *Z. mays*.

File S4. The characterization table of *B. distachyon, O. sativa*, *S. bicolor*, *T. aestivum* and *Z. mays*

TLPs.

File S5. List of qRT-PCR primers.

Table S1. List of predicted *cis*-regulatory elements in the promoter region of *TLP* genes of *B.*

distachyon, O. sativa, S. bicolor, T. aestivum and *Z, mays*.

 Table 1 The Ka/Ks ratio and divergence time of duplicated *TLP* gene pairs.

Gene A	Gene B	Ka	Ks	Ka/Ks	Selection pressure	T = Ks/2r
BdTLPK1	BdTLPK2	0.1102	1.0810	0.1020	Purifying	83.2
OsTLP5	OsTLP7	0.1014	0.6788	0.1494	Purifying	52.2
OsTLP22	OsTLP24	0.0697	0.4134	0.3190	Purifying	27.7
OsTLP23	OsTLP25	0.1149	0.3603	0.31646	Purifying	29.4
OsTLP26	OsTLP27	0.1569	1.3980	0.1122	Purifying	107.6
SbTLP28	SbTLP29	0.0911	0.6808	0.1338	Purifying	52.4
SbTLP31	SbTLP34	0.1388	1.7816	0.0779	Purifying	137
SbTLP32	SbTLP33	0.0055	0.0716	0.0771	Purifying	5.5
TaTLP1-A	TaTLP24-D	0.1224	0.8739	0.1401	Purifying	67.2
TaTLP1-B	TaTLP24-B	0.0937	0.8163	0.1148	Purifying	62.8
TaTLP12-B	TaTLP13-D	0.1193	1.1836	0.1008	Purifying	91
TaTLP15-A	TaTLP18-B	0.1368	0.9879	0.1384	Purifying	76
TaTLP15-B2	TaTLP18-A	0.1423	0.9582	0.1486	Purifying	73.7
TaTLP16-A	TaTLP17-A	0.1456	1.2825	0.1135	Purifying	98.7
ZmTLP1	ZmTLP35	0.0955	0.3929	0.2431	Purifying	30.2
ZmTLP2	ZmTLP34	0.1044	0.8149	0.1281	Purifying	62.7
ZmTLP3	ZmTLP22	0.0495	0.3036	0.1630	Purifying	23.4
ZmTLP9	ZmTLP12	0.0767	0.3783	0.2027	Purifying	29.1

ZmTLP13	*ZmTLP37*	0.0523	0.3200	0.1634	Purifying	24.7
ZmTLP16	*ZmTLP33*	0.0133	0.0141	0.9483	Purifying	1.1
ZmTLP20	*ZmTLP25*	0.0797	0.0889	0.8965	Purifying	6.9

882   **Ka**- non-synonymous substitutions per non-synonymous site, **Ks**- synonymous substitutions

883   per synonymous site, **T**- Divergence time

 Table 2 Tajima's relative rate test of paralogous genes.

Group A	Group B	Outgroup	Nt	Na	Nb	χ^2	P
BdTLP1	*BdTLP2*	*BdTLP11*	403	47	42	0.28	0.59611
OsTLP5	*OsTLP7*	*OsTLP18*	401	28	29	0.02	0.89463
OsTLP22	*OsTLP24*	*OsTLP18*	333	15	17	0.13	0.72367
OsTLP23	*OsTLP25*	*OsTLP18*	313	20	27	1.04	0.30723
OsTLP26	*OsTLP27*	*OsTLP18*	411	34	34	0	1
SbTLP28	*SbTLP29*	*SbTLP5*	316	19	30	2.47	0.11608
SbTLP31	*SbTLP34*	*SbTLP5*	312	27	32	0.42	0.51508
SbTLP32	*SbTLP33*	*SbTLP5*	347	2	2	0	1
TaTLP1-A	*TaTLP24-D*	*TaTLP4-D*	483	37	30	0.73	0.39245
TaTLP1-B	*TaTLP24-B*	*TaTLP4-D*	486	36	31	0.37	0.5413
TaTLP12-B	*TaTLP13-D*	*TaTLP4-D*	321	24	33	1.42	0.23323
TaTLP15-A	*TaTLP18-B*	*TaTLP4-D*	391	36	38	0.05	0.81615
TaTLP15-B2	*TaTLP18-A*	*TaTLP4-D*	260	28	27	0.02	0.89274
TaTLP16-A	*TaTLP17-A*	*TaTLP4-D*	291	36	31	0.37	0.5413
ZmTLP1	*ZmTLP35*	*ZmTLP28*	551	36	33	0.13	0.71798
ZmTLP2	*ZmTLP34*	*ZmTLP28*	541	34	25	1.37	0.24132
ZmTLP3	*ZmTLP22*	*ZmTLP28*	560	11	14	0.36	0.54851
ZmTLP9	*ZmTLP12*	*ZmTLP28*	372	6	12	2	0.1573
ZmTLP13	*ZmTLP37*	*ZmTLP28*	464	19	22	0.22	0.63941

ZmTLP16	ZmTLP33	ZmTLP28	551	5	7	0.33	0.5637
ZmTLP20	ZmTLP25	ZmTLP28	525	17	23	0.9	0.34278

890   **Nt-** Identical sites in all three sequences**, Na**-Unique differences in Sequence A, **Nb-** Unique

891   differences in Sequence B

Table 3. Physicochemical properties of long and small TLP proteins.

Property	Type	*Brachypodium distachyon*	*Oryza sativa*	*Sorghum bicolor*	*Triticum aestivum*	*Zea mays*
Average Protein length	Long TLPs	299	289	294	284	334
	Small TLPs	185	178	178	177	175
Average Molecular weight (kDa)	Long TLPs	30.8	29.8	30.0	29.1	35.0
	Small TLPs	19.5	18.1	18.3	18.22	17.7
Average pI	Long TLPs	6.18	6.95	6.05	6.18	5.89
	Small TLPs	6.82	6.06	5.74	6.17	4.90

Cysteine residue · REDDD motif · FF hydrophobic motif · Bottom of acidic cleft forming amino acids · Signature motif · Thaumatin conserverd domain

YFP Bright field Merged

TaTLP2-B-YFP