Nitazoxanide and JIB-04 have broad-spectrum antiviral activity and inhibit SARS-

CoV-2 replication in cell culture and coronavirus pathogenesis in a pig model

Juhee Son[1,2,*], Shimeng Huang[3,*], Qiru Zeng[1,*], Traci L. Bricker[4], James Brett Case[4],

Jinzhu Zhou[3], Ruochen Zang[1,5], Zhuoming Liu[1], Xinjian Chang[3], Houda H. Harastani[4], Lu

Chen[6], Maria Florencia Gomez Castro[1], Yongxiang Zhao[3], Hinissan P. Kohio[1], Gaopeng

Hou[1], Baochao Fan[3], Beibei Niu[3], Rongli Guo[3], Paul W. Rothlauf[1,7], Adam L. Bailey[8], Xin

Wang[5], Pei-Yong Shi[9], Sean P.J. Whelan[1], Michael S. Diamond[1,4,8], Adrianus C.M.

Boon[1,4,8], Bin Li[3,#], Siyuan Ding[1,#]

[1]Department of Molecular Microbiology, Washington University School of Medicine, St.

Louis, MO, USA. [2]Program in Molecular Cell Biology, Washington University School of

Medicine, St. Louis, MO, USA. [3]Institute of Veterinary Medicine, Jiangsu Academy of

Agricultural Sciences, Nanjing, China. [4]Department of Medicine, Division of Infectious

Diseases, Washington University School of Medicine, St. Louis, MO, USA. [5]Key

Laboratory of Marine Drugs, Ministry of Education, Ocean University of China, Qingdao,

China. [6]National Center for Advancing Translational Sciences, National Institutes of

Health, Rockville, MD, USA. [7]Program in Virology, Harvard Medical School, Boston, MA,

USA. [8]Department of Pathology and Immunology, Washington University School of

Medicine, St. Louis, MO, USA. [9]Department of Biochemistry and Molecular Biology,

University of Texas Medical Branch, Galveston TX, USA.

* These authors contributed equally: Juhee Son, Shimeng Huang, Qiru Zeng

Corresponding authors: Bin Li, libinana@126.com, Siyuan Ding, siyuan.ding@wustl.edu

Abstract

Pathogenic coronaviruses represent a major threat to global public health. Here, using a

recombinant reporter virus-based compound screening approach, we identified several

small-molecule inhibitors that potently block the replication of the newly emerged severe

acute respiratory syndrome virus 2 (SARS-CoV-2). Two compounds, nitazoxanide and

JIB-04 inhibited SARS-CoV-2 replication in Vero E6 cells with an EC_{50} of 4.90 μM and

0.69 μM, respectively, with specificity indices of greater than 150. Both inhibitors had *in*

vitro antiviral activity in multiple cell types against some DNA and RNA viruses, including

porcine transmissible gastroenteritis virus. In an *in vivo* porcine model of coronavirus

infection, administration of JIB-04 reduced virus infection and associated tissue pathology,

which resulted in improved body weight gain and survival. These results highlight the

potential utility of nitazoxanide and JIB-04 as antiviral agents against SARS-CoV-2 and

other viral pathogens.

INTRODUCTION

The coronavirus disease 2019 (COVID-19) pandemic has caused unprecedented global

morbidity, mortality, and socioeconomic destabilization. Thus, there is an urgent unmet

need to develop safe and effective countermeasures to combat the disease and provide

immediate treatment. Multiple efforts are underway to identify candidate drugs that inhibit

the replication of the novel severe acute respiratory syndrome virus 2 (SARS-CoV-2) (*1-*

5), the cause of COVID-19 (*6, 7*). So far, several small-molecule inhibitors that interfere

with SARS-CoV-2 cell entry have been identified, including transmembrane serine

protease inhibitors camostat (*8*) and nafamostat (*9*), and endosomal inhibitors including

chloroquine and derivatives (*9*), E-64d (*8*), apilimod (*10*), and 25-hydroxycholesterol (*11*).

Drug screens and structural studies also revealed compounds that target SARS-CoV-2

viral enzymes such as the viral RNA-dependent RNA polymerase (RdRp) (*12-14*) and the

main protease (M^{pro}, also known as $3CL^{pro}$) (*15-17*). Here, utilizing a fluorescent SARS-

CoV-2 virus and an imaging-based screen approach, we identified several known and

previously unknown broadly antiviral compounds that successfully inhibit SARS-CoV-2

replication.

RESULTS

To identify small molecules with anti-SARS-CoV-2 activity, we performed a screen with a

recombinant SARS-CoV-2 that encodes mNeonGreen as a reporter of infection (*18*) and

an in-house collection of ~200 compounds that comprise FDA-approved drugs, well-

defined broad-spectrum antivirals, and investigational new drugs. We identified

molecules with stronger antiviral efficacy (>44.8% inhibition) than the positive controls

chloroquine and remdesivir against SARS-CoV-2 replication in Vero E6 cells (**Fig. 1A** and

Dataset S1).

We selected two small molecules for detailed characterization: [2-[(5-nitro-1,3-thiazol-2-

yl)carbamoyl]phenyl] acetate (nitazoxanide, abbreviated as NTZ), an FDA-approved drug

for treatment of diarrhea caused by *Giardia lamblia* and *Cryptosporidium parvum*

infections (*19*), and 5-chloro-N-[(E)-[phenyl(pyridin-2-yl)methylidene]amino]pyridin-2-

amine (JIB-04 E-isomer), a pan-Jumonji histone demethylase inhibitor (*20*) (**Fig. S1A**).

NTZ was chosen based on its broad-spectrum antiviral activity against several respiratory

viruses (rhinovirus A, influenza A virus, human metapneumovirus and respiratory

syncytial virus) (https://drugvirus.info/) and safety profile in humans. NTZ is currently

being tested in clinical trials for COVID-19 in Mexico and Brazil (ClinicalTrials.gov

identifiers: NCT04341493; NCT04348409). However, no pre-clinical data is published to

support an *in vitro* inhibitory function. JIB-04 was picked for its stronger anti-SARS-CoV-

2 activity than NTZ (**Fig. S1B**). Moreover, it has not been reported as an antiviral molecule

before and several histone demethylases were recently uncovered in a CRISPR/Cas9

genetic knockout screen of SARS-CoV-2 host dependency factors (*21*).

We confirmed that pre-treatment of cells with either NTZ or JIB-04 for 1 hr inhibited the

replication of a clinical isolate of SARS-CoV-2 (2019-nCoV/USA-WA1/2020 strain) by

viral antigen staining in Vero E6 cells (**Fig. 1B**). In Vero E6 cells, NTZ and JIB-04 had

half-maximal effective concentration (EC_{50}) values of 4.90 µM (with 95% confidence

interval of 3.05-7.25 µM) and 695 nM (with 95% confidence interval of 567-822 nM),

respectively. Both drugs did not induce cytotoxicity at 300 µM and thus had excellent

selectivity indices (SI > 150). Intracellular SARS-CoV-2 RNA levels also were reduced

significantly by NTZ and JIB-04 but not by camostat treatment (**Fig. 1C**).

To examine whether the lead compounds target cellular pathways (e.g., entry, translation, replication, or assembly) shared by different viruses, we counter-screened 15 molecules with a vesicular stomatitis virus (VSV) eGFP virus (*22*) and a replication-competent chimeric VSV eGFP virus that encodes SARS-CoV-2 spike (S) protein in place of the native glycoprotein (*23*). Concanamycin A and MG132 preferentially (>50-fold) inhibited VSV-SARS-CoV-2 than VSV, implying that at least part of the antiviral activity was specific to the S protein (**Fig. 2A**). In contrast, the majority of the top hits from the drug screen (thapsigargin, AG-1478, FTY720, 3-isobutyl-1-methylxanthine (IBMX), arbidol, nigericin, JIB-04, and nitazoxanide) suppressed both VSV and VSV-SARS-CoV-2 replication in MA104 cells (**Fig. 2A**). Using flow cytometry, we observed that NTZ or JIB-04 reduced the infectivity of VSV and VSV-SARS-CoV-2 at 6 hr post infection (hpi), within a single replication cycle (**Fig. 2B**). Less GFP intensity was detected in infected cells (**Fig. S2A**), suggesting that the inhibitors might act on a common post-entry pathway. NTZ or JIB-04 reduced VSV-SARS-CoV-2 replication in a dose-dependent manner without apparent cytotoxicity (**Fig. 2C** and **S2B**). The inhibitory effects were more apparent when cells were infected with VSV-SARS-CoV-2 at a low multiplicity of infection (MOI) (**Fig. S2C**). At 30 µM, JIB-04 treatment led to 100-fold reduction of intracellular VSV-SARS-CoV-2 RNA levels, comparable to the efficacy of chloroquine (**Fig. S2D**).

Unlike NTZ, with reported broad-spectrum antiviral activity (*24-27*), JIB-04 is not known to have an antiviral function. We found that besides VSV (**Fig. 2A**), JIB-04 had antiviral property against both DNA and RNA viruses, including vaccinia virus and two strains of

rotaviruses (RVs) (**Fig. 2D**). JIB-04 also inhibited the replication of enteric viruses, i.e. a

strain of porcine RV and transmissible gastroenteritis virus (TGEV) (**Fig. 2E-F** and **Fig.**

S2E), a porcine CoV that infects the small intestine of pigs and causes severe diarrheal

disease (*28*).

Ciliated airway epithelial cells are a target for SARS-CoV-2 *in vivo* (*29*), and the human

lung epithelial cell line, Calu-3, has been used to evaluate SARS-CoV-2 infection as an

in vitro surrogate (*8, 30*). NTZ inhibition of VSV-SARS-CoV-2 replication, as measured by

viral RNA levels, was ineffective in HEK293-hACE2-TMPRSS2 and Calu-3 cells, in

contrast to JIB-04 (**Fig. 2G**), suggesting distinct antiviral mechanisms.

Although NTZ and JIB-04 have been previously connected to interferon (IFN) and

autophagy activation (*20, 31, 32*), the antiviral activity that we observed was independent

of these pathways. Neither treatment led to the induction of IFN and IFN-stimulated gene

expression or the formation of LC3-positive punctate structures (**Fig. S3A-B**). Using a

dose-response drug matrix, we found that JIB-04 had synergistic antiviral effects with

chloroquine and NTZ in MA104 cells (**Fig. 3A**). Synergism was calculated by the highest

single agent (HSA) synergy model using SynergyFinder 2.0 (*33*). JIB-04 at 0.1-3 µM had

the most synergy with either chloroquine or NTZ at 1-10 µM (**Fig. 3A**). A combinatorial

effect of JIB-04 and camostat also was observed in Calu-3 cells (**Fig. S3C**), further

highlighting a unique antiviral mode of action of JIB-04.

Time of addition experiments also supported that both NTZ and JIB-04 primarily acted at

a post-entry step (**Fig. 3B**). Pre-incubation of the cells with JIB-04 for 1 hr significantly

reduced SARS-CoV-2 S mRNA transcription post VSV-SARS-CoV-2 infection (**Fig. 3C**).

Consistently, the newly synthesized full-length S protein levels were strongly decreased

(**Fig. 3D**). We obtained similar results with SARS-CoV-2 plasmid transfection. JIB-04

treatment suppressed S mRNA transcription (**Fig. S3D**) and almost completely prevented

the production of full-length and protease-processed SARS-CoV-2 S proteins (**Fig. S3E**).

In parallel, we examined the cellular signaling pathways modulated by JIB-04. Small

interfering RNA (siRNA)-mediated knockdown of JIB-04 cellular targets (*i.e.*, histone

demethylases KDM4B, KDM4C, KDM5A, or KDM5B (*20*)) recapitulated the antiviral effect

of JIB-04 (**Fig. 3E** and **Fig. S3F-G**). We hypothesized that JIB-04 treatment led to H3K9

and H3K27 methylation and silenced expression of a subset of genes. To evaluate this

idea, we performed RNA-sequencing on mock and JIB-04 treated cells with or without

virus infection to identify the potential JIB-04 targets. Pathway analysis revealed

dampened metabolic signaling such as the cytochrome P450 system in JIB-04 treated

cells (**Fig. S3H**). Specifically, JIB-04 downregulated two cytochrome P450 enzymes,

CYP1A1 and *CYP1B1*, and aryl hydrocarbon receptor repressor (*AHRR*), product of

which represses transactivator of CYP1A1 and CYP1B1 (*34*) (**Fig. 3F**). We validated by

quantitative PCR that JIB-04 treatment reduced *CYP1A1*, *CYP1B1*, and *AHRR* mRNA

levels by 4-6-fold (**Fig. 3G**). To explore the pharmacological utility of this finding, we tested

the antiviral activity of cytochrome P450 enzyme inhibitors Fluoxetine and Fluvoxamine

(*35*). Both compounds inhibited the replication of VSV-SARS-CoV-2 (**Fig. 3H**) and wild-

type SARS-CoV-2 (**Fig. 3I**).

Next, we used a neonatal pig TGEV infection model (*36*) to assess the antiviral efficacy

of JIB-04 against coronavirus *in vivo*. Two-day old piglets were injected via intraperitoneal

route with JIB-04 at 24 hr and 6 hr before oral inoculation of TGEV (**Fig. 4A**). We

monitored body weight on a daily basis and recorded diarrhea development and mortality

every 6 hr. The animals in the control group lost more weight and had more severe

diarrhea than those receiving JIB-04 (**Fig. S4A-B**). At 2 days post infection, 3 out of 5

piglets in the DMSO group succumbed to infection as compared to 1 out of 5 animals in

the JIB-04 group (**Fig. 4B**). Consistent with our *in vitro* results (**Fig. 2E-F**), the TGEV viral

loads throughout the gastrointestinal tract, measured by viral RNA and titer levels, were

substantially lower in the JIB-04 treated piglets (**Fig. 4C-D**). The number of viral antigen

positive intestinal epithelial cells also was decreased following JIB-04 treatment (**Fig.**

S4C). Independent of virus inhibition, JIB-04 enhanced the expression of tight-junction

genes such as claudin-1 and ZO-1 and pro-inflammatory cytokines including IFN-β and

TNF-α (**Fig. S4D-E**). Furthermore, we observed less severe enteropathy in the JIB-04

treated animals (**Fig. 4E**). Taken together, our data demonstrate *in vivo* antiviral activity

of JIB-04 against a major porcine CoV pathogen.

DISCUSSION

Using a repurposed compound screening approach, we identified drugs with reported

inhibitory activity against SARS-CoV-2, such as tetrandrine (*37*) and arbidol (*38*), and

characterized a number of previously unreported small-molecule (JIB-04, AG-1478,

nigericin, etc.) that reduced SARS-CoV-2 infection. Several compounds converge on the

endosomal pathway: brefeldin A, concanamycin A, tetrandrine, and U18666A.

Furthermore, FTY720 induced formation of enlarged endosome/lysosome structure,

similar to that triggered by apilimod treatment (*39*), highlighting an important role of

endosomal trafficking in SARS-CoV-2 entry and infection, at least in cell culture.

Our results highlight NTZ and JIB-04 as potential therapeutics for SARS-CoV-2 and the

need for further evaluation. Compared to NTZ, JIB-04 does not have reported antiviral

properties. Another compound in our screen, GSK-J4, also is a histone demethylase

inhibitor, which targets JMJD3/KDM6B. However, it did not inhibit SARS-CoV-2

replication. Thus, a definitive involvement of certain KDM family members in SARS-CoV-

2 host interactions has not been established.

Although NTZ has been used to treat protozoan infections since 1984 (*40*), the

mechanisms underlying its anti-microbial activities are not fully elucidated. Our *in vitro*

results support a post-entry inhibitory mode of action, which was previously proposed for

hepatitis C virus (*27*). In a non-placebo-controlled setting, NTZ, given at the dosage of

500 mg twice a day for 5 days, was used to treat 150 severe COVID-19 patients in a

public hospital in Guayaquil, Ecuador (Villacís-Bermeo BA, Personal Communication,

May 15, 2020). Besides antiviral activity, NTZ is reported to reduce plasma IL-6 level in

mice (*41*), which suggests that it also might reduce inflammatory responses. We propose

that studies are needed to determine possible combinatorial efficacy of these antivirals in

combination with remdesivir in COVID-19 patients.

MATERIALS AND METHODS

Reagents, cells, and viruses

Reagents: NTZ used in *in vitro* assays (N0290, Sigma-Aldrich, > 98% purity), JIB-04 E-

isomer used in *in vitro* assays (S7281, Selleckchem, 99.8% purity), JIB-04 E-isomer used

in *in vivo* experiments (HY-13953, Med-ChemExpress), Fluvoxamine maleate (S1336,

Selleckchem), Fluoxetine HCl (S1333, Selleckchem), low molecular weight poly(I:C)

complexed with LyoVec (tlrl-picwlv, InvivoGen). EGFP-LC3 plasmid was a gift from

Christina Stallings at Washington University School of Medicine. SARS-CoV-2 S plasmid

was used as previously described (*42*). pUC19 empty plasmid was used as mock in all

transfection experiments.

Cells: Vero E6 cells (CRL-1586, ATCC) and Vero cells (CCL81, ATCC) were cultured in

DMEM supplemented with 10% fetal bovine serum (FBS), 10 mM HEPES, 1 mM sodium

pyruvate, 0.1 mM non-essential amino acids, and 1 X Penicillin-Streptomycin-

Glutamine. Calu-3 cells (HTB-55, ATCC) and swine ST cells (CRL-1746, ATCC) were

DMEM supplemented with 10% FBS and 1 X Penicillin-Streptomycin-Glutamine. HEK293,

HEK293-hACE2, and HEK293-hACE2-TMPRSS2 cells were cultured in complete DMEM

containing G418 and/or blasticidin and used as previously described (*11*). MA104 and

Vero E6-TMPRSS2 cells were cultured as before (*42*).

Viruses: Rhesus RV RRV strain, bovine RV UK strain, and porcine RV NJ2012 strain

(Genbank: MT874983-MT874993) were propagated and titrated as before (*43*). Vaccinia

virus MVA strain was used as before (*44*). HSV-1 syn17+ strain was a gift from Ann Arvin

at Stanford University. TGEV JS2012 strain was propagated as before (*45*). TGEV was

titrated by serial dilutions in cells in 96-well plates that were seeded at a density of 1 X

10^4 cells per well. Cytopathic effects were observed at 3-7 dpi and the $TCID_{50}$ values

were calculated and converted to PFU/ml. A clinical isolate of SARS-CoV-2 (2019-

nCoV/USA-WA1/2020 strain) was obtained from the Centers for Disease Control and

Prevention (gift of Natalie Thornburg). A SARS-CoV-2 mNeonGreen reporter virus was

used as previously reported (*18*). Both the clinical isolate and the mNeonGreen SARS-

CoV-2 viruses were propagated in Vero CCL81 cells and titrated by focus-forming assays

on Vero E6 cells. Recombinant VSV-eGFP (*22*) and VSV-eGFP-SARS-CoV-2 were

previously described (*23*). Cells infected with GFP-positive virus were imaged with ECHO

REVOLVE 4 fluorescence microscope. Plaque assays were performed in MA104 cells

seeded in 6-well plates using an adapted version of the rotavirus plaque assay protocol

(*43*).

Inhibitor screen

The small-molecule inhibitors used in this study are from in-house collection and the

COVID Box (Medicines for Malaria Venture; www.mmv.org/mmv-open/covid-box).

Compound names, vendors, and catalog numbers are listed in **Table S1**. At 24 hpi, cells

were fixed in 4% paraformaldehyde (PFA) in PBS and stained with Hoechst 33342. The

levels of viral antigens, reflected by mNeonGreen signals, were scanned by Amersham

Typhoon 5 (GE). Image background was subtracted using rolling ball algorithm (radius =

5 pixels). To further remove the imaging artifacts and well-to-well variation, we removed

the region which fell below the threshold calculated by Moments algorithm. Number of

positive pixels and total intensity (after background subtraction) were computed for each

well and log10 transformed. The number of cells in each well was quantified based on

Hoechst 33342 staining detected by Cytation 3 imaging reader (BioTek). Image analysis

was performed using ImageJ and customized R scripts. The quantification of

mNeonGreen and Hoechst 33342 is provided in **Dataset S1**.

Cell cytotoxicity assay

The viability of Vero E6 and MA104 cells after drug treatment was determined using the

Cell Counting Kit 8 (ab228554, Abcam). Briefly, cells in 96-well plates were treated with

NTZ or JIB-04 at desired concentrations at 37°C. After 25 hr of incubation, the inhibitor-

containing medium was replaced with fresh complete medium with 10 µl of WST-8

solution in each well. The cells were incubated at 37 °C for 2 hr with protection from light.

Absorbance at 460nm was measured using Gen5 software and BioTek ELx800

Microplate Reader.

RNA extraction and quantitative PCR

Total RNA was extracted from cells using RNeasy Mini kit (Qiagen). For spike plasmid

transfection experiments, total RNA was extracted using Aurum Total RNA Mini Kit

(Biorad) with DNase digestion. Complementary DNA was synthesized with High Capacity

cDNA Reverse Transcription kit (Thermo Fisher) as previously described (*46*).

Quantitative PCR was performed using AriaMX (Agilent) with 12.5 μl of either Power

SYBR Green master mix or Taqman master mix (Applied Biosystems) in a 25 μl reaction.

Gene expression was normalized to the housekeeping gene GAPDH. All SYBR Green

primers and Taqman probes used in this study are listed in **Table S2**.

Western blotting

Cells were lysed in RIPA buffer supplemented with protease inhibitor cocktail and

phosphatase inhibitor. Lysates were boiled for 5 min in 1 x Laemmli Sample Buffer (Bio-

Rad) containing 5% β-mercaptoethanol. Proteins were resolved in SDS-PAGE and

detected as described (*47*) using the following antibodies: GAPDH (631402, Biolegend),

rotavirus VP6 (rabbit polyclonal, ABclonal technology), and SARS-CoV-2 S2 (40592-T62,

Sino Biological). Secondary antibodies were anti-rabbit (7074, Cell Signaling) or anti-

mouse (7076, Cell Signaling) immunoglobulin G horseradish peroxidase-linked

antibodies. Protein bands were visualized with Clarity ECL Substrate (Bio-rad) and a

Biorad Gel Doc XR system.

Small interfering RNA transfection

HEK293 cells were transfected using Lipofectamine RNAiMAX Transfection Reagent

(Thermo Fisher). Cells were harvested at 48 hr post transfection and knockdown

efficiency was determined by RT-qPCR. siRNA transfected cells were infected with PoRV

(MOI=0.01) for 12 hr and viral RNA copy numbers were examined by RT-qPCR. All siRNA

used in this study were designed and synthesized by GenePharma (Shanghai, China)

and their sequences of siRNAs are listed in **Table S2**.

Flow cytometry

Vero E6-TMPRSS2 cells were inoculated with VSV-GFP or VSV-SARS-CoV-2 at an MOI

of 3 for 1 hr at 37°C. At 6 hpi, cells were harvested and fixed in 4% PFA in PBS.

Percentage of GFP positive cells and GFP intensity were determined by BD

LSRFortessa™ X-20 cell analyzer and analyzed by FlowJo v10.6.2 (BD).

RNA-sequencing

HEK293 cells were pre-treated with JIB-04 (10 µM) for 12 hr, and mock or porcine RV-

infected (MOI=0.01) for another 12 hr. Total RNA from cells in triplicate was extracted

using RNeasy Mini kit (Qiagen). RNA sample quality was measured by both NanoDrop

spectrophotometer (Thermo Fisher) and Bioanalyzer 2100 (Agilent). Libraries were

sequenced on the Illumina NovaSeq 6000 platform. Differential gene expression analysis

was performed using DESeq2. The RNA-seq raw and processed datasets were deposited

onto NCBI Gene Expression Omnibus database (GSE156219).

TGEV piglet infection

Newborn piglets (Landrace × Yorkshire) were spontaneously delivered from sows, and

their body weights were recorded at birth. Fifteen neonatal male pigs at birth were

obtained from a TGEV-free farm in Nanjing without suckling. All piglets were confirmed

negative for TGEV by RT-PCR and ELISA (IDEXX, USA). The pigs were randomly

separated into three groups, housed in separate rooms, and fed the same artificial milk

substitutes that meet the nutrient and energy recommendations of the National Research

Council [NRC, 2012] at the animal facility of the Institute of Veterinary Medicine, Jiangsu

Academy of Agricultural Sciences, Nanjing, Jiangsu Province. The experiments were

divided into three groups: a DMSO inoculation control group (control, n=5); a DMSO

inoculation and TGEV infection group (TGEV-DMSO, n=5); a JIB-04 inoculation and

TGEV infection group (TGEV-JIB-04, n=5)). Neonatal pigs were intraperitoneally injected

twice with JIB-04 (75 mg/kg) or DMSO at 24 hr and 6 hr prior to TGEV infection. TGEV-

DMSO and TGEV-JIB-04 groups were orally infected with $1\times10^{7.25}$ (1.778×10^7) $TCID_{50}$

(equivalent to 1.245×10^7 PFU) of TGEV in 1.5 ml of DMEM per pig. Neonatal pigs were

weighed and observed for clinical signs every 8 hr throughout the study. Serum samples

were collected from each pig at 24 and 48 hpi to detect specific anti-TGEV antibodies.

The occurrence of diarrhea was monitored, and its severity was recorded based on an

established scoring system (*44*). In brief, diarrhea was scored on the basis of color,

consistency, and amount, and numbered as follows: 0 = normal; 1 = pasty; 2 = semi-liquid;

3 = liquid, and score ≥ 2 considered as diarrhea. At 48 hpi, all pigs were euthanized, and

intestinal tissues were collected for pathological examination and viral load analysis using

RT-qPCR and primers in **Table S2**.

Histopathological and immunofluorescence analysis

Intestinal tissues harvested from pigs were fixed in 4% PFA in PBS and incubated in 50%

ethanol overnight. After fixation, tissues were embedded in paraffin, sectioned, and

subjected to hematoxylin and eosin staining by standard procedures. For

immunofluorescence analysis, samples were incubated with rabbit anti-TGEV-N antibody

(1:500, DA0224, Shanghai YouLong Biotech) for 30 min at 37 ºC. After three washes,

samples were stained with Cy3-conjugated goat anti-rabbit secondary antibody

(Beyotime) and DAPI (Invitrogen). Images were obtained using a fluorescence

microscope (Carl Zeiss).

Ethics statement

Animal experiments were approved by the Committee on the Ethics of Animal Care and

Use of the Science and Technology Agency of Jiangsu Province. The approval ID is

NKYVET 2014-63, granted by the Jiangsu Academy of Agricultural Sciences

Experimental Animal Ethics Committee. All efforts were made to minimize animal

suffering. The virus challenge and tissue collection were performed in strict accordance

with the guidelines of Jiangsu Province Animal Regulations (Decree No. 2020-18).

Statistical analysis

All bar graphs were displayed as means ± standard error of mean (SEM). Statistical

significance in data Fig. 2E, 2F, 3C, S3D, S3F, 4C, and 4D was calculated by Student's *t*

test using Prism 8.4.3 (GraphPad). Statistical significance in data Fig. 1C, 2D, 2G, 3B,

3E, 3G, S3A, S3C, and S4B was calculated by pairwise ANOVA using Prism 8.4.3. Non-

linear regression (curve fit) was performed to calculate EC_{50} and CC_{50} values for Fig. 1B,

2A, and 2C using Prism 8.4.3. HSA synergy model was used to calculate the synergy

scores of dose-response data in Fig. 3A. Gehan-Breslow-Wilcoxon test was used to

compare the survival curves in Fig. 4B. All data were presented as asterisks (*p≤0.05;

**p≤0.01; ***p≤0.001). All experiments other than Fig. 1A, 3I, and 4 were repeated at least

twice. The raw data and p value calculations are included in Dataset S1.

SUPPLEMENTARY MATERIALS

Table S1. List of chemicals used in the anti-SARS-CoV-2 compound screen

Table S2. List of quantitative PCR primers and siRNA

Table S3. Raw data and statistics

Figure 1

375
376

Fig. 1. Nitazoxanide and JIB-04 inhibit SARS-CoV-2 replication

(A) Small molecule inhibitor screen. Vero E6 cells were treated with individual

compounds (listed in Table S1) at 10 µM for 1 hour (hr) and infected with SARS-

CoV-2-mNeonGreen (MOI=0.5). At 24 hr post infection (hpi), cells were fixed and

nuclei were stained by Hoechst 33342. The intensities of mNeonGreen and

Hoechst were quantified by the Typhoon biomolecular imager and Cytation plate

reader, respectively. The ratio of mNeonGreen and Hoechst is plotted as

percentage of inhibition.

(B) Dose-response analysis of wild-type SARS-CoV-2 replication with NTZ or JIB-04

treatment. Vero E6 cells were treated with compounds for 1 hr and infected with a

clinical isolate of SARS-CoV-2 (MOI=0.5). S protein levels were quantified at 24

hpi based on immunofluorescence. For CC_{50} measurement, cells were treated with

inhibitors at 0.3 µM to 1 mM for 25 hr. SI: selectivity index.

(C) Dose-response analysis of intracellular viral RNA levels with compounds. Vero E6

cells were treated with NTZ (10 µM), JIB-04 (10 µM), chloroquine (10 µM),

remdesivir (3 µM), or camostat (10 µM) for 1 hr and infected with a clinical isolate

of SARS-CoV-2 (MOI=0.5). SARS-CoV-2 RNA levels at 24 hpi were measured by

RT-qPCR.

For all panels except A, experiments were repeated at least three times with similar

results. Fig. 1A was performed once with raw data included in Dataset S1. Data

are represented as mean ± SEM. Statistical significance is from pooled data of the

multiple independent experiments (*$p \leq 0.05$).

Figure 2

400
401

Fig. 2. Nitazoxanide and JIB-04 broadly inhibit DNA and RNA viruses in different cell types

(A) Dose-response analysis of VSV and VSV-SARS-CoV-2 replication with 15 compounds. MA104 cells were treated with indicated compounds at 0.01 to 30 µM for 1 hr and infected with VSV or VSV-SARS-CoV-2 (MOI=3). GFP signals at 24 hpi were quantified to calculate the percentage of inhibition. EC_{50} values for VSV and VSV-SARS-CoV-2 are shown in each graph in red and blue, respectively.

(B) Virus infectivity with NTZ or JIB-04 treatment. Vero E6-TMPRSS2 cells were treated with compounds (10 µM) for 1 hr and infected with VSV or VSV-SARS-CoV-2 (MOI=3). At 6 hpi, percentages of GFP positive cells were quantified by flow cytometry.

(C) Dose-response analysis of VSV-SARS-CoV-2 replication and cytotoxicity with NTZ or JIB-04 treatment. For EC_{50} measurement, MA104 cells were treated with compounds at 0.01 to 30 µM for 1 hr and infected with VSV-SARS-CoV-2 (MOI=3) for 24 hr. For CC_{50} measurement, cells were treated with compounds at 0.1 µM to 3 mM for 25 hr. SI: selectivity index.

(D) Intracellular viral RNA levels with NTZ or JIB-04 treatment. MA104 cells were treated with compounds (10 µM) for 1 hr and infected with vaccinia virus (VACV), herpes simplex virus-1 (HSV-1), or rotavirus (RV, RRV and UK strains) (MOI=1). Viral RNA levels at 24 hpi were measured by RT-qPCR for VACV B10R, HSV-1 ICP-27, and RV NSP5, respectively.

(E) Viral RNA copy numbers with JIB-04 treatment. HEK293 cells were treated with JIB-04 (10 µM) for 6 hr and infected with porcine rotavirus (MOI=0.01) for 6 hr. ST

cells were treated with JIB-04 (10 µM) for 12 hr and infected with transmissible

gastroenteritis virus (TGEV) (MOI=0.01) for 12 hr. Viral RNA copy numbers were

measured by RT-qPCR.

(F) TGEV titers in the cell supernatant with JIB-04 treatment. ST cells were treated

with JIB-04 (10 µM) for 12 hr and infected with TGEV (MOI=0.01). Virus titers at 6

and 12 hpi were measured by plaque assays.

(G)Intracellular viral RNA levels with NTZ or JIB-04 treatment in different cell types.

HEK293-hACE2, HEK293-hACE2-TMPRSS2, and Calu-3 cells were treated with

compounds (10 µM) for 1 hr and infected with VSV-SARS-CoV-2 (MOI=1). VSV

RNA levels at 24 hpi were measured by RT-qPCR.

For all panels except A, experiments were repeated at least three times with similar

results. Fig. 2A was performed once. Data are represented as mean ± SEM.

Statistical significance is from pooled data of the multiple independent experiments

(*p≤0.05; **p≤0.01; ***p≤0.001).

Figure 3

Fig. 3. JIB-04 exhibits distinct post-entry antiviral mechanisms

(A) Drug combination dose-response matrix and VSV-SARS-CoV-2 replication. MA104 cells were treated with JIB-04 and chloroquine or JIB-04 and NTZ for 1 hr and infected with VSV-SARS-CoV-2 (MOI=3). GFP signals at 24 hpi were quantified to calculate the percentage of inhibition.

(B) Time of compound addition and VSV-SARS-CoV-2 replication. MA104 cells were treated with NTZ or JIB-04 (10 µM) at indicated time points relative to VSV-SARS-CoV-2 infection (MOI=3, 0 hpi). GFP signals at 8 hpi were quantified to calculate the percentage of inhibition.

(C) Intracellular SARS-CoV-2 S RNA levels with JIB-04 treatment. MA104 cells were treated with JIB-04 (10 µM) for 1 hr and infected with VSV-SARS-CoV-2 (MOI=1) for 1, 3, 5, and 7 hr. S RNA levels were measured by RT-qPCR.

(D) Western blot analysis of SARS-CoV-2 S protein levels with JIB-04 treatment. MA104 cells were treated with JIB-04 (10 µM) for 1 hr and infected with VSV-SARS-CoV-2 (MOI=1) for 1, 3, 5, and 7 hr. FL: full-length. S2: cleaved S2 fragment. (* non-specific band)

(E) Histone demethylase siRNA knockdown and RV replication. HEK293 cells were transfected with scrambled siRNA or siRNA targeting indicated histone demethylases for 48 hr and infected with porcine RV (MOI=0.01). Viral RNA copy numbers at 12 hpi were quantified by RT-qPCR.

(F) Volcano plot of differentially expressed transcripts with JIB-04 treatment and RV infection. HEK293 cells were treated with DMSO or JIB-04 (10 µM) for 12 hr, and mock-infected (left panel) or infected with porcine RV (MOI=0.01, right panel) for

another 12 hr. Red dots represent upregulated genes and green dots represent

downregulated genes in JIB-04 treated cells.

(G)Expression of three top genes in (F) with JIB-04 treatment. HEK293 cells were

treated with JIB-04 (10 µM) for 12 hr and mock-infected or infected porcine RV

(MOI=0.01) for 12 hr. mRNA levels of CYP1A1, CYP1B1, and AHRR at 12 hpi

were measured by RT-qPCR.

(H)Dose-response analysis of VSV-SARS-CoV-2 replication with fluoxetine or

fluvoxamine treatment. MA104 cells were treated with compounds at 0.01 to 30

µM for 1 hr and infected with VSV-SARS-CoV-2 (MOI=3). GFP signals at 24 hpi

were quantified to calculate the percentage of inhibition. For CC_{50} measurement,

cells were treated with compounds at 0.1 µM to 300 µM for 25 hr.

(I) Dose-response analysis of wild-type SARS-CoV-2 replication with fluoxetine or

fluvoxamine treatment. Vero E6 cells were treated with compounds for 1 hr and

infected with a clinical isolate of SARS-CoV-2 (MOI=0.5). S protein levels at 24 hpi

were quantified based on immunofluorescence. For CC_{50} measurement, cells were

treated with compounds at 0.1 µM to 300 µM for 25 hr.

For all panels except A and I, experiments were repeated at least three times with

similar results. Fig. 3A was performed twice. Inhibition assay in Fig. 3I was

performed once and cytotoxicity assay was performed in triplicates. Data are

represented as mean ± SEM. Statistical significance is from pooled data of the

multiple independent experiments (*p≤0.05; **p≤0.01; ***p≤0.001).

Figure 4

487
488
Fig. 4. JIB-04 suppresses TGEV replication and pathogenesis in pigs

(A) Experimental schemes for testing the protective efficacy of JIB-04 treatment

against TGEV challenge in three groups of neonatal pigs. Control: DMSO injection,

mock infection; TGEV-DMSO: DMSO injection, TGEV infection; TGEV-JIB-04:

JIB-04 injection, TGEV infection.

(B) Survival curve of TGEV infected pigs with JIB-04 treatment. Neonatal pigs were

intraperitoneally injected with vehicle control DMSO or JIB-04 and mock-infected

or infected with 1.2×10^7 PFU of TGEV. Survival was monitored every 8 hr with

data censored at 48 hpi, when all pigs were sacrificed.

(C) TGEV RNA copy numbers in the gastrointestinal (GI) tract of infected pigs. TGEV

infected piglets were sacrificed at 48 hpi. For the DMSO group, two animals

sacrificed at 48 hpi and one that died at 40 hpi (colored in blue) were examined.

For the JIB-04 groups, four animals sacrificed at 48 hpi were examined. TGEV

genome copy numbers at 48 hpi were quantified by RT-qPCR.

(D) Same as (C) except that virus titers were measured by plaque assays.

(E) Hematoxylin and eosin staining of different GI tract sections from pigs sacrificed at

48 hpi. Representative images of 3 animals. Scale bar, 100 μm.

Data are represented as mean ± SEM. Statistical significance is from pooled data

of the multiple independent experiments (*p≤0.05).

Supplemental Figures and Legends

Figure S1

510
511
512 **Fig. S1. Nitazoxanide and JIB-04 inhibit SARS-CoV-2 replication**

513 (A) Chemical structures of NTZ and JIB-04 E-isomer from ChemSpider database.

514 (B) Representative images of Vero E6 cells infected by SARS-CoV-2-mNeonGreen

515 (MOI=0.5) at 24 hpi in Fig. 1A.

516 Experiments were repeated at least three times with similar results.

Figure S2

Fig. S2. Nitazoxanide and JIB-04 inhibit the replication of multiple viruses

(A) Mean fluorescence intensity of GFP positive cells in Fig. 2B was quantified by flow

cytometry.

(B) Dose-response analysis of VSV-SARS-CoV-2 replication with NTZ or JIB-04

treatment. MA104 cells were treated with compounds at indicated concentrations

for 1 hr and infected with VSV-SARS-CoV-2 (MOI=3). At 24 hpi, images of GFP

positive infected cells were acquired by the ECHO fluorescence microscope.

(C) Same as (B) except that cells were infected with an MOI of 0.1.

(D) Dose-response analysis of intracellular viral RNA levels with NTZ, JIB-04, or

chloroquine treatment. MA104 cells were treated with compounds at 0.1 to 30 µM

for 1 hr and infected with VSV-SARS-CoV-2 (MOI=3). VSV RNA levels at 24 hpi

were measured by RT-qPCR.

(E) Western blot analysis of RV antigen VP6 levels with JIB-04 treatment. HEK293

cells were treated with JIB-04 at 1, 5, or 10 µM for 6 hr and infected with porcine

RV (MOI=0.01) for 12 hr. GAPDH was used as a loading control.

All experiments were repeated at least three times with similar results. Data are

represented as mean ± SEM.

539
540

Figure S3

Fig. S3. Inhibition or knockdown of specific KDM histone demethylases inhibits

virus replication

(A) Expression of IFN and IFN-stimulated genes with NTZ or JIB-04 treatment. HEK293 cells were treated with NTZ (10 µM), JIB-04 (3 µM), or transfected with low-molecular-weight poly(I:C) (100 ng/ml) for 24 hr. mRNA levels of IFNL3 and CXCL10 were measured by RT-qPCR.

(B) Autophagy formation with compound treatment. HEK293 cells were transfected with EGFP-LC3 plasmid for 24 hr and treated with rapamycin (100 nM), NTZ (10 µM), or JIB-04 (3 µM) for another 18 hr. GFP positive punctate structures indicate autophagy activation. Scale bar, 20 µm.

(C) Intracellular viral RNA levels with JIB-04 and camostat treatment. Calu-3 cells were treated with compounds (10 µM) for 1 hr and infected with VSV-SARS-CoV (MOI=3). VSV RNA levels at 24 hpi were measured by RT-qPCR.

(D) Intracellular SARS-CoV-2 S RNA levels with JIB-04 treatment. HEK293 cells were transfected with SARS-CoV-2 S plasmid for 2 hr and treated with JIB-04 (3 µM) for 24 hr. S RNA levels were measured by RT-qPCR.

(E) Western blot analysis of SARS-CoV-2 S protein levels with JIB-04 treatment. HEK293 cells were transfected with SARS-CoV-2 S plasmid for 2 hr and treated with JIB-04 (3 µM) for 24 hr. FL: full-length. S2: cleaved S2 fragment.

(F) siRNA-mediated knockdown of JIB-04 target histone demethylases. HEK293 cells were transfected with scrambled siRNA or siRNA targeting indicated histone demethylases for 48 hr. mRNA levels of indicated histone demethylases were measured by RT-qPCR.

(G)Western blot analysis of RV antigen VP6 levels in cells with histone demethylase

siRNA knockdown. HEK293 cells were transfected with scrambled siRNA or siRNA

targeting indicated histone demethylases for 48 hr and infected with porcine RV

(MOI=0.01) for 12 hr.

(H)Pathway enrichment analysis of gene expression regulated by JIB-04 treatment.

Downregulated genes in Fig. 3F with p values < 1e-10 were analyzed by DAVID

functional annotation.

All experiments were repeated at least three times with similar results. Data are

represented as mean ± SEM. Statistical significance is from pooled data of the

multiple independent experiments (*$p \leq 0.05$; **$p \leq 0.01$; ***$p \leq 0.001$).

Figure S4

Fig. S4. JIB-04 reduces TGEV induced weight loss and pathogenesis

(A) Weight of TGEV infected pigs with JIB-04 treatment in Fig. 4B. The body weight of

individual animals was monitored every 24 hr.

(B) Diarrhea occurrence in TGEV infected pigs with JIB-04 treatment in Fig. 4B.

Diarrhea severity was scored for the fecal specimens of DMSO or JIB-04 treated,

mock or TGEV infected animals every 8 hr.

(C) Immunofluorescence staining of TGEV antigen in different GI tract sections from

pigs sacrificed at 48 hpi. Blue: cell nuclei; red: TGEV nucleocapsid protein.

Representative images of 3 animals. Scale bar, 100 μm.

(D) Expression of tight junction associated genes in the GI tract of TGEV infected pigs

with JIB-04 treatment. For the DMSO group, two animals sacrificed at 48 hpi and

one that died at 40 hpi (colored in blue) were examined. For the JIB-04 groups,

four animals sacrificed at 48 hpi were examined. CLDN1 and TJP1 mRNA levels

at 48 hpi were measured by RT-qPCR.

(E) Same as (D) except that IFNB and TNF mRNA levels were measured instead.

Data are represented as mean ± SEM. Statistical significance is from pooled data

of the multiple independent experiments (*$p \leq 0.05$; **$p \leq 0.01$).

597
598
599

REFERENCES AND NOTES

1. F. Touret, Gilles, M., Barral, K., Nougairede, A., Decroly, E., de Lamballerie, X., Coutard, B., In vitro screening of a FDA approved chemical library reveals potential inhibitors of SARS-CoV-2 replication. *bioRxiv*, (2020).
2. L. Riva, Yuan, S., Yin, X., Martin-Sancho, L., Matsunaga, N., et al., , A Large-scale Drug Repositioning Survey for SARS-CoV-2 Antivirals. *bioRxiv*, (2020).
3. C. Mirabelli, Wotring, J.W., Zhang, C.J., McCarty, S.M., Fursmidt, R., et al., , Morphological Cell Profiling of SARS-CoV-2 Infection Identifies Drug Repurposing Candidates for COVID-19. *bioRxiv*, (2020).
4. M. Dittmar, Lee, J.S., Whig, K., Segrist, E., Li, M., Jurado, K., Samby, K., Ramage, H., Schultz, D., Cherry, S., Drug repurposing screens reveal FDA approved drugs active against SARS-Cov-2. *bioRxiv*, (2020).
5. K. Heiser, McLean, P.F., Davis, C.T., et al., , Identification of potential treatments for COVID-19 through artificial intelligence-enabled phenomic analysis of human cells infected with SARS-CoV-2. *bioRxiv*, (2020).
6. F. Wu *et al.*, A new coronavirus associated with human respiratory disease in China. *Nature* **579**, 265-269 (2020).
7. P. Zhou *et al.*, A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273 (2020).
8. M. Hoffmann *et al.*, SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*, (2020).
9. M. Wang *et al.*, Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell research* **30**, 269-271 (2020).
10. Y. L. Kang, Chou, Y.Y., Rothlauf, P.W., Liu, Z., Soh, T.K., Cureton, D., Case, J.B., Chen, R.E., Diamond, M.S., Whelan, S.P.J., Kirchhausen, T., Inhibition of PIKfyve kinase prevents infection by EBOV and SARS-CoV-2. *bioRxiv*, (2020).
11. R. Zang, Case, J.B., Gomez Castro, M.F., Liu, Z., Zeng, Q., Zhao, H., Son, J., Rothlauf, P.W., Hou, G., Bose, S., Wang, X., Vahey, M.D., Kirchhausen, T., Fremond, D.H., Diamond, M.S., Whelan, S.P.J., Ding, S., Cholesterol 25-hydroxylase suppresses SARS-CoV-2 replication by blocking membrane fusion. *bioRxiv*, (2020).
12. W. Yin *et al.*, Structural basis for inhibition of the RNA-dependent RNA polymerase from SARS-CoV-2 by remdesivir. *Science*, (2020).
13. Y. Gao *et al.*, Structure of the RNA-dependent RNA polymerase from COVID-19 virus. *Science* **368**, 779-782 (2020).
14. R. N. Kirchdoerfer, A. B. Ward, Structure of the SARS-CoV nsp12 polymerase bound to nsp7 and nsp8 co-factors. *Nature communications* **10**, 2342 (2019).
15. L. Zhang *et al.*, Crystal structure of SARS-CoV-2 main protease provides a basis for design of improved alpha-ketoamide inhibitors. *Science* **368**, 409-412 (2020).
16. W. Dai *et al.*, Structure-based design of antiviral drug candidates targeting the SARS-CoV-2 main protease. *Science*, (2020).
17. Z. Jin *et al.*, Structure of M(pro) from SARS-CoV-2 and discovery of its inhibitors. *Nature* **582**, 289-293 (2020).
18. X. Xie *et al.*, An Infectious cDNA Clone of SARS-CoV-2. *Cell host & microbe*, (2020).

19. L. M. Fox, L. D. Saravolatz, Nitazoxanide: a new thiazolide antiparasitic agent. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **40**, 1173-1180 (2005).

20. L. Wang *et al.*, A small molecule modulates Jumonji histone demethylase activity and selectively inhibits cancer growth. *Nature communications* **4**, 2035 (2013).

21. J. Wei, Alfajaro, M.M., Hanna, R.E., et al., , Genome-wide CRISPR screen reveals host genes that regulate SARS-CoV-2 infection. *bioRxiv*, (2020).

22. S. Cherry *et al.*, Genome-wide RNAi screen reveals a specific sensitivity of IRES-containing RNA viruses to host translation inhibition. *Genes & development* **19**, 445-452 (2005).

23. J. B. Case, Rothlauf, P.W., Chen, R.E., Liu, Z., Zhao, H., Kim, A.S., Bloyet, L., Zeng, Q., Tahan, S., Droit, L., et al.,, Neutralizing antibody and soluble ACE2 inhibition of a replication-competent VSV-SARS-CoV-2 and a clinical isolate of SARS-CoV-2. *Cell host & microbe*, (2020).

24. W. Dang *et al.*, Nitazoxanide Inhibits Human Norovirus Replication and Synergizes with Ribavirin by Activation of Cellular Antiviral Response. *Antimicrobial agents and chemotherapy* **62**, (2018).

25. V. Hargest, B. Sharp, B. Livingston, V. Cortez, S. Schultz-Cherry, Astrovirus Replication Is Inhibited by Nitazoxanide In Vitro and In Vivo. *Journal of virology* **94**, (2020).

26. S. La Frazia *et al.*, Thiazolides, a new class of antiviral agents effective against rotavirus infection, target viral morphogenesis, inhibiting viroplasm formation. *Journal of virology* **87**, 11096-11106 (2013).

27. M. Elazar *et al.*, The anti-hepatitis C agent nitazoxanide induces phosphorylation of eukaryotic initiation factor 2alpha via protein kinase activated by double-stranded RNA activation. *Gastroenterology* **137**, 1827-1835 (2009).

28. L. J. Saif, Animal coronavirus vaccines: lessons for SARS. *Dev Biol (Basel)* **119**, 129-140 (2004).

29. Y. J. Hou *et al.*, SARS-CoV-2 Reverse Genetics Reveals a Variable Infection Gradient in the Respiratory Tract. *Cell*, (2020).

30. T. P. Sheahan *et al.*, An orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in human airway epithelial cell cultures and multiple coronaviruses in mice. *Science translational medicine* **12**, (2020).

31. M. Xu *et al.*, SHMT2 and the BRCC36/BRISC deubiquitinase regulate HIV-1 Tat K63-ubiquitylation and destruction by autophagy. *PLoS pathogens* **14**, e1007071 (2018).

32. K. K. Lam *et al.*, Nitazoxanide stimulates autophagy and inhibits mTORC1 signaling and intracellular proliferation of Mycobacterium tuberculosis. *PLoS pathogens* **8**, e1002691 (2012).

33. A. Ianevski, L. He, T. Aittokallio, J. Tang, SynergyFinder: a web application for analyzing drug combination dose-response matrix data. *Bioinformatics* **36**, 2645 (2020).

34. S. I. Karchner, D. G. Franks, W. H. Powell, M. E. Hahn, Regulatory interactions among three members of the vertebrate aryl hydrocarbon receptor family: AHR repressor, AHR1, and AHR2. *The Journal of biological chemistry* **277**, 6949-6959 (2002).

35. A. Hemeryck, F. M. Belpaire, Selective serotonin reuptake inhibitors and
cytochrome P-450 mediated drug-drug interactions: an update. *Curr Drug Metab*
3, 13-37 (2002).
36. L. Luo *et al.*, Aminopeptidase N-null neonatal piglets are protected from
transmissible gastroenteritis virus but not porcine epidemic diarrhea virus.
Scientific reports **9**, 13186 (2019).
37. X. Ou *et al.*, Characterization of spike glycoprotein of SARS-CoV-2 on virus entry
and its immune cross-reactivity with SARS-CoV. *Nature communications* **11**, 1620
(2020).
38. X. Wang *et al.*, The anti-influenza virus drug, arbidol is an efficient inhibitor of
SARS-CoV-2 in vitro. *Cell Discov* **6**, 28 (2020).
39. Y. L. Kang *et al.*, Inhibition of PIKfyve kinase prevents infection by EBOV and
SARS-CoV-2. *bioRxiv*, (2020).
40. J. F. Rossignol, H. Maisonneuve, Nitazoxanide in the treatment of Taenia saginata
and Hymenolepis nana infections. *The American journal of tropical medicine and*
hygiene **33**, 511-512 (1984).
41. S. K. Hong *et al.*, Nitazoxanide suppresses IL-6 production in LPS-stimulated
mouse macrophages and TG-injected mice. *Int Immunopharmacol* **13**, 23-27
(2012).
42. R. Zang *et al.*, TMPRSS2 and TMPRSS4 promote SARS-CoV-2 infection of
human small intestinal enterocytes. *Sci Immunol* **5**, (2020).
43. S. Ding *et al.*, Rotavirus VP3 targets MAVS for degradation to inhibit type III
interferon expression in intestinal epithelial cells. *eLife* **7**, (2018).
44. B. Li *et al.*, Drebrin restricts rotavirus entry by inhibiting dynamin-mediated
endocytosis. *Proceedings of the National Academy of Sciences of the United*
States of America **114**, E3642-E3651 (2017).
45. R. Guo *et al.*, Characterization and evaluation of the pathogenicity of a natural
recombinant transmissible gastroenteritis virus in China. *Virology* **545**, 24-32
(2020).
46. C. R. Bolen, S. Ding, M. D. Robek, S. H. Kleinstein, Dynamic expression profiling
of Type I and Type III Interferon-stimulated hepatocytes reveals a stable hierarchy
of gene expression. *Hepatology*, (2013).
47. S. Ding, W. Khoury-Hanold, A. Iwasaki, M. D. Robek, Epigenetic reprogramming
of the type III interferon response potentiates antiviral activity and suppresses
tumor growth. *PLoS biology* **12**, e1001758 (2014).
727
728

Funding

This study is supported by the National Institutes of Health (NIH) DDRCC grant P30 DK052574, NIH grants K99/R00 AI135031 and R01 AI150796 and the COVID-19 Fast Grants Funding to S.D., NIH contracts and grants (75N93019C00062 and R01 AI127828) and the Defense Advanced Research Project Agency (HR001117S0019) to M.S.D., NIH grant U01 AI151810 to A.C.M.B., and unrestricted funds from Washington University School of Medicine and NIH grant R37 AI059371 to S.P.W. The *in vivo* pig studies are supported by the Jiangsu Province Natural Sciences Foundation (BK20190003) and National Natural Science Foundation of China (31872481) to B.L. J.B.C. is supported by a Helen Hay Whitney Foundation postdoctoral fellowship.

Author contributions

J.S., Q.Z., R.Z., and S.D. designed, executed, and analyzed *in vitro* efficacy studies. M.F.G.C., H.P.K., and G.H. assisted with the RNA extraction and RT-qPCR analysis. Y.Z. performed the *in vitro* TGEV inhibition studies. Z.L. performed the flow cytometry analysis. L.C. wrote the algorithm that quantifies inhibitor screen results. P.W.R. and S.P.J.W. constructed the VSV-SARS-CoV-2 virus. Q.Z., J.Z., and R.G. propagated and titrated viruses. J.B.C. propagated and infected the clinical isolate of SARS-CoV-2. P.Y.S. provided the recombinant SARS-CoV-2 mNeonGreen virus. A.L.B propagated the mNeonGreen virus and designed the SARS-CoV-2 Taqman probe. S.H., B.L., and S.D. designed the *in vivo* efficacy studies. S.H., J.Z., X.C., B.F., and B.N., performed the *in*

vivo TGEV infection experiments, dissected the animals and harvested tissues, and

measured viral titers and cytokine mRNA levels. X.W., S.P.J.W., M.S.D., A.C.M.B., B.L.,

and S.D. provided supervision and funding for the study. J.S. and S.D. wrote the

manuscript with the input and edits from S.H., Q.Z., J.B.C., J.Z., Z.L., M.F.G.C., H.P.K.,

G.H., S.P.J.W., M.S.D., A.C.M.B., and B.L.

Competing interests

Acknowledgements

The authors thank Dr. Harry Greenberg (Stanford University, USA) and Dr. Bolívar A.

Villacís-Bermeo (Guayaquil, Guayas, Ecuador) for constructive comments and

arguments. We appreciate the assistance from Matthew Williams (Department of

Molecular Microbiology Media and Glassware Facility) and Erica Lantelme (Department

of Pathology and Immunology, Flow Cytometry Core Facility). Cytation plate scanning

was assisted by Zhou Huang (Department of Molecular Microbiology).

Data and materials availability

All raw data and statistical analysis in the current study are available in Table S3 and

Dataset S1. RNA-seq dataset has been deposited onto NCBI GEO (GSE156219).