Overview of the expression of key regeneration genes in embryo development and matured tissues in Axolotl (*Ambystoma mexicanum*)

Authors:
- Harshil Shah, No Affiliation, harshilsemail1@gmail.com
- Juan Caballero, Universidad Autonoma de Queretaro, jcaballero@uaq.mx

Date
09/25/2020

ABSTRACT

The axolotl is a Mexican endangered species with the capability to perform tissue regeneration in amputated extremities. Being an important research model, the complete genome of the Axolotl was sequenced in 2018 for the first time, revealing an enormous genome: the largest of any animal ever sequenced, and about 10 times larger than the human genome, a new landmark achievement in biology research.

In this report, we collected 70 known genes that play an important role in tissue regeneration and searched for their expression during embryo development, regeneration, and in 6 adult tissues: the heart, liver, gills, front leg, rear leg, and tail. We observe those 70 genes expression levels in the 3 conditions and approximately 3 genes seem to be expressed exclusively in regeneration.

This report displays a few insights on how this marvelous species can modulate its regenerative capabilities. There is still much more to explore with the Axolotl. Further research into their regenerative capabilities could provide researchers with the perspicacity to possibly accomplish human limb regeneration.

Keywords:
Axolotl, tissue regeneration, expression analysis, RNA-seq

INTRODUCTION

The axolotl (*Ambystoma mexicanum*) is famous for its lifelong youthfulness, more scientifically described as neoteny, the fact that it does not undergo the metamorphosis from larval, aquatic stage to adult, terrestrial salamander-like related amphibian species would, but reaches maturity while in the larval form (Gross 2018). The Axolotl is also paedomorphic, retaining juvenile features throughout their lifespan and into adulthood, including the external gills, a body fin, and moveable eyelids (https://animals.sandiegozoo.org/animals/axolotl).

Diverse molecular analysis has been performed on this species, including the complete genome of the Axolotl which was sequenced in 2018 for the first time. The genome revealed enormous chromosomes: it is the largest of any animal ever sequenced, and about 10 times larger than the human genome (Dockrill 2020). The full sequencing marked a landmark in Biology because the data available could be researched for potential awareness into their regenerative ability as well as other useful traits the Axolotl possesses.

The axolotl's capability allows them to regenerate their limbs, lungs, heart, jaws, spines, and even parts of their brain (https://animals.sandiegozoo.org/animals/axolotl). Studies have found that they can regrow a new limb five times perfectly in a few weeks. Other than being regenerative, wild Axolotl's can change the hue of their skin a few shades for camouflage.

The stages of regeneration for a salamander leg are visible in the composite images Figure 1, Figure 2, and Figure 3. The intact limb is at the left; the series to the right shows the gradual regeneration of the limb over a couple of months. The steps include: first the intact limb receives trauma and we see the wound without the limb.

The first step of regeneration involves epithelialization, a process of covering excoriated or traumatized epithelial surfaces. First, a clot of blood cells rapidly stops bleeding at the wounded site. Then, a layer of cells begins to form an epithelial and cover the wound completely (Dunlap 2018). The cellular and molecular processes involved in initiation, maintenance, and completion of epithelialization are essential for successful wound closure.

Next, in innervation, nerves are replaced into the wounded areas and go into muscle fibers, innervating the muscle fiber. The wounded epidermis transforms into a layer of signaling cells called the Apical Epithelial Cap, which has a vital role in regeneration. Interesting research conducted previously further analyzes how cells at the stump of the wound do not revert to a completely embryonic state as previously thought, but rather the parts that were muscle remembers that it needs to grow muscle, whereas the part that was nerve remembers that it needs to grow nerve. They are still able to grow into tissues, but only certain kinds of tissue (Farrow 2017).

The cells underneath the epidermis also begin to rapidly divide, forming a cone-shaped structure known as a blastema. The cells that make up the blastema are thought to be bone, cartilage, muscle, or other cells that de-differentiate to become like stem cells.

Dedifferentiation is an important biological phenomenon whereby cells regress from a specialized function to a simpler state reminiscent of stem cells. Stem cells are self-renewing cells capable of giving rise to differentiated cells when supplied with the appropriate factors, and these stem cells can then begin migrating to necessary areas.

These de-differentiated stem cells in the blastema then grow and multiply, eventually regaining their identity as fully developed cells to form necessary tissues and organs. As the blastema and its cells continue to divide, the growing structure flattens and eventually resembles a perfect copy of the lost limb, including nerves and blood vessels that are connected to the rest of the body.

Figure 1. Photographs of limb regeneration in Axolotl (Farrow 2017)

Figure 2. Graphic representation of the limb regeneration in Axolotl (Mccusker 2015)

Figure 3. Bone and muscle regeneration in Axolotl limb amputation (Dunlap 2018)

Further research from James Godwin, the lead study author, from the Australian Regenerative Medicine Institute (ARMI) at Monash University in Melbourne, concludes that macrophages play an essential role in the regenerative process. They experimented with salamanders by injecting the animals with a chemical substance that depleted the macrophage-levels of the animals and then amputating limbs to see the body's reaction. They found that those salamanders could not regenerate the limbs and had severe scarring build-up. However, after they reverse-engineered this and re-introduced the macrophages, the animal was able to regenerate those same lost limbs, thus demonstrating the importance of macrophages in the process (Lewis 2013). Macrophages have already been known to play crucial roles in mouse embryo tissues and development for many animals. Godwin believes that the ability to regenerate was turned off due to evolution, but it may be possible to reevaluate that and reverse the process (Lewis 2013).

Axolotl genome and research

Learning more about the Axolotl's anatomy, geographic density, and lifestyle/cycle could contribute to the overall understanding of the animal's abilities and tendencies. Studying the major organs of the Axolotl could provide information on the roles of different genes In molecular processes and metabolism. Since our research and experiment focused on the Axolotl's Rear and Front Leg, Gills, Heart, Liver, and Tail, Figure 4 shows a diagram of the Axolotl body labeled with major organs.

Figure 4. Axolotl main organs.

The axolotl's capability allows them to regenerate their limbs, lungs, heart, jaws, spines, and even parts of their brain (https://animals.sandiegozoo.org/animals/axolotl). Studies have

found that they can regrow a new limb five times perfectly in a few weeks. Other than being regenerative, wild Axolotl's can change the hue of their skin a few shades for camouflage.

The final superpower is that the Axolotl is 1,000 times more resistant to cancers than mammals, and researchers from the University of Nottingham used an extract from axolotl eggs to stimulate tumor suppressor genes and prevent breast cancer growth (Travers 2018). Scientists from the University brought the Breast Cancer Cells back under control by reactivating their tumor suppressor genes (University of Nottingham 2011). Treating the cells with Axolotl oocyte extract reactivated the suppressor genes and stopped the cancer from growing, and within 60 days, there was no cancerous growth. Cancers are caused by mutations within the cell division cycle and uncontrollable growth of mutated cells, and among the most important of these genes are tumor suppressor genes. These genes repress the development of cancers and normally act as a control point in the cell division cycle. Therefore, the switching off tumor suppressor genes is a common cause of cancers, including breast cancer" (University of Nottingham 2011). Using novel technology that makes use of the eggs of the axolotl salamander, cancer cells were treated with extracts made from axolotl oocytes that could reverse the epigenetic marks on tumor suppressor genes, causing these genes to reactivate, and thereby stopping the cancerous cell growth. Axolotl oocytes are packed with molecules that have very powerful epigenetic modifying activity. Previously, the Johnson's lab at the University of Nottingham showed that extracts prepared from these oocytes have a powerful capacity to change epigenetic marks on the DNA of human cells (University of Nottingham 2011).

For this research, the three major functions we focused on were limb regeneration, development in the embryo, and major organs, as mentioned before. Focusing on each, first, we took data from experiments when Axolotls were amputated limbs in a laboratory and studied over a 40-period study. The key stages of regeneration, as the data reflected, were the: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 16, 19, 24, and 40 stages. For the embryo development, Figure 5 shows the 40 stages of development of the Axolotl from an egg to larva. Finally, we included data of the expression levels of proteins in the Axolotl in major organs, as listed above: Front and Rear Leg, Tail, Gills, Heart, and Liver.

METHODS

Sequences

We used the sequences of all Axolotl assembled transcriptome reported in (Caballero-Perez 2018), in brief, such database was built using the assembly combination of regeneration transcriptome (Bryant 2017), embryo development (Jiang 2017) and the RNA-seq of the adult tissues. After merging the assemblies, the transcriptome was quantified for each stage with Kallisto and normalized in transcript per million to enable cross-comparisons.

Selection of key genes in regeneration

We used the list of 70 key genes involved in Axolotl regeneration as reported in (Sibai), in the initial screening we noted many genes were incomplete, therefore we collected the orthologue gene searching the main human RefSeq protein for each gene in NCBI GenBank.

Transcript identification in Axolotl transcriptome

We utilized NCBI Blast+ to search the transcriptome database using "tblastn" to use the human protein as a query, and the transcriptome nucleotide database, so all transcripts are translated to the equivalent proteins for the comparison. With the specific Axolotl genes that were most similar to the human gene found (E-value < 1e-6), we collected those transcripts identifiers and searched the database for their expression levels in the three major functions: regeneration, development, and tissues.

RESULTS

Expression patterns

Expression of Axolotl Genes Table

	Regeneration	Embryo	Tissues
Extremely High Expression	2	22	26
High Expression	2	8	8
Medium Expression	7	8	12
Low Expression	54	31	16
No Expression	5	1	8

All 70 genes were identified in Axolotl transcriptome. The genes were classified into different levels of expression based on the greatest number of stages/time-periods in which the transcript was expressed at that level. No expression was considered for 0 tpm, low expression 0-2 tpm, medium expression 2-5 tpm, high expression 5-20 tpm, and extreme high expression was >20 tpm.

Table 1. Global expression of genes in the 3 transcriptomes.

As shown in Table 1, in regeneration, most genes had low expression while a few had high and extremely high expression. The few genes that are left are expressed in extremely high quantities. In the embryo/development, more genes were expressed in extremely high quantities as well as many genes that were expressed in lower quantities. Finally, in the major organs of the Axolotl, most genes were extremely high expressed out of all the three functions

Table 2 displays the expression for each of the genes. The first column lists the gene symbol that was being inputted into the transcriptome database. The next column lists the Axolotl gene identifier that was shown to be most similar to the human gene, and it is listed as in the Amex transcriptome database. The next three columns list the expression levels of the Axolotl genes in regeneration, development, and major organ tissues in detail, with all the different expression levels present throughout the experiment.

Gene Symbol	Axolotl Gene	Expression in Regeneration	Expression in Embryo / Development	Expression in Axolotl Tissues
AURKA	Amex3_3159454	Low expression throughout process	Extremely high expression, gradually decreasing	No expression
AURKB	Amex3_3260416	Medium expression in regeneration, increasing gradually towards later stages	Very high expression, gradually decreasing	High expression in tissues (Especially front and rear legs, gills, and tail)
CCNJ	Amex3_3258442	Low expression	High expression, decreases substantially at end	Low expression l
CDC20	Amex3_3534199	Low expression in the 3rd, 4th, and 7th stage(6hrs, 12hrs, 5d respectively)	Low expression, with medium expression in the 7th and 11th stage	Low expression in the heart, rear leg, and tail
CDCA8	Amex3_2115615	Low expression	Extremely high expression in all stages of development	High expression (14+ in Front and Rear legs, Tail, and Gills)
KIF11	Amex3_208660	No expression	Low expression only in the 1st, 2nd, 9th, 12th, 14th, 16th, and 19th stages	Low expression only in Front leg and tail
FLNA	Amex3_208660	No expression	Low expression only in the 1, 2, 9, 12, 14, 16, 19, and 24 stages	Low expression only in Front leg and Tail
KIF22	Amex3_3258744	Low expression	Extremely high expression, gradually declining	High expression in Gills and Rear leg, Low expression in rest
KIF23	Amex3_2629094	Low expression	Low expression in only 2, 4, 9, 12, 16, 19, 24, 40 stages	No expression
MAD2L1	Amex3_3532025	Low expression	Extremely high expression, decreases to	Low expression

			medium expression after 11 stage	
MAD2L1BP	Amex3_3533474	Low expression	Extremely high expression, decreases to medium expression after 24 stage	Medium expression
MTBP	Amex3_3531074	Low expression at 12hrs, 3d, 10d, 21d, 28d	Medium expression in 1, 2, 3, 4 stages, low expression in all other stages	Low expression
NEK2	Amex3_3259832	Low expression	Extremely high expression	Low expression in Heart and Liver, Medium expression in the rest
NIPBL	Amex3_208253	Low expression only at 6hrs, 1d, 14d, 21d, 28d	Medium expression in middle stages, Low expression in rest	Low expression
PLK1	Amex3_402983	Low expression	Extremely high expression, decreasing gradually	Low expression in Heart and Liver, High expression in rest
PRC1	Amex3_1857349	None	High expression in the first and last stages, extremely high expression in the middle	Low expression
RCC1	Amex3_819365	Low expression 10d, 21d	Medium expression	Low expression in only Front leg
TRIP13	Amex3_3159454	Low expression at 3hrs, 6hrs, 1d, 3d, 5d, 14d, and 28d	Extremely high expression, decreases substantially at end to low expression	No expression
APEX1	Amex3_2503633	Low expression	Extremely high expression	High expression
FEN1	Amex3_3533277	Low expression	Medium to High expression	High expression in tail, medium expression in rest
HCFC1	Amex3_2112644	Low expression	Low to Medium expression	Low expression
PGGT1B	Amex3_3156215	Low expression	Medium expression in first two stages, then high expression	Low expression only in Front and Rear leg, Gills, and Heart
RPS6KB1	Amex3_3106233	Low expression	Low expression in first two stages, medium expression in next three stages, then high expression	Medium expression
SFPQ	Amex3_1628	Low expression at 0hrs, 12hrs, 1d, 3d, 7d, 10d, 21d, 28d	Medium to High expression	Low expression in Heart, High expression in Rear leg, Medium expression in rest
ASF1B	Amex3_2564976	Low expression	Extremely high expression, decreases to medium expression at end	Medium expression in Rear leg, Low expression in rest
HELLS	Amex3_374305	Low expression	Medium expression at stage 3, Low expression	No expression

			only at 1, 2,4,14,19,24 stages	
HIST3H2BB	Amex3_161534	High to Extremely high expression	Low expression only at 1,2,5,8,9,16 stage	No expression
HMGB2	Amex3_403585	No expression till 12hrs, then Low expression	Low expression till 8 stage, then Extremely high expression	Low to Medium expression
NASP	Amex3_2503343	No expression till 6hrs, then Low expression	Extremely high expression	Medium expression in Liver, High expression in rest
RPA2	Amex3_2565539	Low expression	Extremely high to High expression	Low expression in Heart and Liver, Medium expression in rest
SART3	Amex3_3258344	Low expression	Extremely high expression	Medium expression
TOP1	Amex3_1383108	Low expression	Extremely high expression	Medium expression
TOP2B	Amex3_99453	Low expression	Low expression till 7d, then medium expression till 10d, then Extremely high expression	High expression in Gills and Rear leg, Medium expression in rest
ACTA1	Amex3_288061	Medium expression at 3hrs,12hrs, Low expression in rest	Low expression till 21d, then High expression	Medium expression in Gills and Rear leg, Low expression in Heart
ACTN3	Amex3_2626973	No expression	Low expression at 8,9,24,40 stage	Extremely high expression in Tail, High expression in Front and Rear leg, Low expression in Heart
DES	Amex3_334965	Medium expression till 1d, then Low expression	Low expression only at 1,3,14,16,19,24,40 stages	Low expression in Liver, Extremely high expression in rest
MYBPC2	Amex3_77332	Medium expression till 12hrs, then Low expression, gradually decreasing trend	Low expression only at 1,5,14,16,24,40 stages	Low expression in Heart and Liver, Extremely high expression in rest
MYBPC3	Amex3_68648	Low expression only at 0hrs,12hrs,1d,3d,5d,14d, 21d,28d	Low expression only at 19 stage	Extremely high expression in Heart, Low expression in Front leg and Gills, no expression in rest
MYH2	Amex3_1761	Low expression	Low expression only at 40 stage	No expression in Heart or Liver, Medium expression in Gills, Extremely high expression in rest
MYH4	Amex3_1761	Low expression	Low expression only at 40 stage	No expression in Heart or Liver, Medium expression in Gills, Extremely high expression in rest
MYH7	Amex3_105660	Low expression	Low expression only after 7 stage	Low expression in Liver, Medium expression in Gills, High expression in Tail, Extremely high

				expression in rest
MYL1	Amex3_2633364	High expression till 12hrs, then Medium expression till 5d, then Low expression	Low expression	No expression in Liver, Low expression in Heart, High expression in Gills, Extremely high expression in rest
MYL2	Amex3_22809	Low expression at 7d, 10d, 14d, Medium expression for rest	Low expression	No expression in Liver, Low expression in Heart, Medium expression in Gills, Extremely high expression in rest
MYL3	Amex3_329288	Extremely high expression till 12hrs, then High expression till 5d, then Low expression	Low expression only at 1,14 stage, Medium expression at 40 stage	Medium expression in Liver, Extremely high expression in rest
MYL4	Amex3_2639977	Low expression	Low expression	No expression in Front leg and Gills, Low expression in Liver and Tail, Medium expression in Rear leg, Extremely high expression in Heart
TNNC1	Amex3_339993	High expression till 12hrs, then Medium expression till 3d, then Low expression	High expression at 40 stage, Medium expression at 24 stage, low expression for rest	Low expression in Liver, Extremely high expression in rest
TNNC2	Amex3_361070	Extremely high expression, decreases to Low expression	High expression at 1 stage, then Low expression for rest	Low expression in Liver, High expression in Heart, Extremely high expression in rest
TNNI1	Amex3_115364	Extremely high expression till 12hrs, then High expression till 3d, then Low expression for rest	Low expression only at 2,5,10,12,16,19,24,40 stage	No expression in Liver, Medium expression in Heart, High expression in Gills, Extremely high expression for rest
TNNI2	Amex3_2638627	Extremely high expression till 1d, then high expression till 5d, then Low expression for rest	Extremely high expression for 1 stage, then Low expression for rest	No expression in Liver, Low expression in Heart, Extremely high expression in rest
TNNT1	Amex3_198319	High expression till 3d, then Low expression	Low expression only at 2,9,10,11,12,14,16,19,24,40 stage	No expression in Liver, Low expression in Heart, Medium expression in Gills, Extremely high expression in rest
TNNT3	Amex3_823025	Extremely high expression till 3d, then Medium and Low expression	High expression at 1 stage, then Low expression for rest	No expression in Liver, Medium expression in Heart, Extremely high expression in rest
TPM1	Amex3_190914	Extremely high expression till 5d, then Low expression	Medium expression	Medium expression in Liver, Extremely high expression in rest
TPM3	Amex3_345240	Medium expression till 3d, then Low expression	Low expression only at 9,10,11,12,19,24,40 stage	No expression in Liver, Extremely high expression in rest
ABAT	Amex3_2629439	Low expression	Low / Medium expression	Extremely high expression in Liver,Medium expression in Heart, Low

				expression in rest
ANXA6	Amex3_42541	Low expression	Low expression	High expression in Gills, Extremely high expression in rest
BIN1	Amex3_656102	Low expression only at 0hrs,10d,21d,28d	Medium / High expression	Low expression only in Tail, no expression in rest
MYH10	Amex3_3530194	Low expression	Extremely high / High expression	Extremely high expression in Heart, Medium / High expression in rest
MYOZ1	Amex3_2633493	Medium / High expression till 7d, then Low expression	Low expression	No expression in Heart or Liver, High expression in Gills, Extremely high expression in rest
GAPDH	Amex3_128063	Low expression only at 12hrs,1d	Medium expression at 40 stage, Low expression 1,2,6 stage, No expression in other stages	No expression
GPD1	Amex3_132015	Low expression	Low expression	Medium expression in Gills, Extremely high expression in liver, High expression in rest
GPI	Amex3_3534586	High expression at 6hrs and 1d, Low expression at 10d, Medium expression in rest	High expression in 2,24 stage, Extremely high expression in rest	Extremely high expression
MDH2	Amex3_124638	Low expression	Low expression	Extremely high expression
PFKM	Amex3_2632336	Low expression	Low expresssion	No expression in Liver, Low expression in Gills, High expression in Heart and Tail, Extremely high expression in rest
PGK1	Amex3_375656	High expression at 6hrs, Low expression at 10d, Medium expression for rest	Extremely high / High expression	Extremely high expression
SLC25A12	Amex3_349153	Low expression	Low expression	Low expression in Liver, High expression in Gills and Tail, Extremely high expression in rest
TPI1	Amex3_2565813	Low expression only at 0hrs, 6hrs,12hrs,1d,3d,7d,28d	Extremely high expression at 1,7,9,10,11,12,40 stage, High expression in rest	Extremely high expression
PGM1	Amex3_3531432	Low expression	Low expression	Low expression
PRKAG2	Amex3_604	Low expression only at 6hrs,1d,10d,14d	Medium expression till 12 stage, then Low expression for rest	Low expression in Tail and Heart, Medium expression for rest
ATP1B2	Amex3_110410	Medium expression	Low expression till 8 stage, High expression at 9 stage, then Extremely high expression	Low expression in Liver, Medium expression in Heart, Extremely high expression in rest

GABARAPL2	Amex3_3107823	Low expression	High expression at 2,40 stage, Medium expression at 12,24 stage, Low expression at 14,16,19 stage, Extremely high expression for rest	High expression in Liver, Extremely high expression in rest

Table 2. Detailed gene expression in the 3 main datasets, expression levels are classified as: Low expression: 0 - 2 tpm, Medium expression: 2-5 tpm, High expression: 5-20 tpm, and Extremely High: >20 tpm.

Figure 5. Heatmap for gene expression in the 3 datasets.

Figure 5 displays the overall results of all the genes, which are listed at the top of the graph, of each other the three data retrievals/comparisons we made: the regeneration process, development process, and the protein concentration in major tissues/organs. The colors show the relative expression levels. We see that regeneration, as observed above, has most genes were expressed at a low or no expression. We also observe that in embryo and major tissues more genes are expressed at higher quantities.

DISCUSSION

Regeneration Genes

We were able to find that there are orthologues genes conserved between humans and Axolotl gene sequences. Considering the functions of many genes and the proteins they produce in both organisms, especially for major organs and tissues, it can be observed that many of the same functions need to be carried out by the organisms for survival. That implies that they utilize similar if not the same proteins adapted best for the organism.

The gene expression observed in their respective transcripts showed that genes were expressed in a variety of levels and each condition displayed gene expression in different quantities.

In regeneration, most genes had low expression while a few had high and extremely high expression, displaying how fewer proteins are required for the regeneration process, and most genes are expressed in moderately lower levels but still play instrumental roles in the regeneration process. The few genes expressed in higher numbers are probably expressed in much higher quantities to make up and are required in large quantities to aid in the regenerative process.

In the embryo/development, our observation concludes that more proteins are required for the developmental process and more genes are required in large numbers, and this could be explained because development takes longer and requires more resources to complete correctly, compared to regeneration, which only is re-developing one part of the body at the time.

Finally, gene expression in the major organs demonstrated the vast amount of genes and proteins required for the Axolotl's regular body functions. There was also a considerable amount of genes that were expressed in lower and medium quantities, showing that some functions only require small quantities to complete the necessary processes.

Conservation in Other Species

Figure 8 shows the evolutionary relationship of different species of the p53 cancer suppressor gene. The numbers on the branches of the phylogenetic trees are bootstrapping values.

Figure 9 is a phylogenetic tree of BH3-only proteins. Sequences found for zebrafish, Xenopus, human and axolotl were translated into amino acid sequences and aligned using ClustalW2 and T-coffee. A tree was generated and visualized. The scale bar indicates amino acid changes" (Vesna 2018). The Axolotl and Humans are shown to have evolutionary connections from the Bad, Bid, and Bim gene, where they are the closest relatives of those genes compared to the five organisms in the study.

Figure 8. Phylogenetic analysis for the P53 gene (Villiard 2007).

Figure 9. Phylogenetic analysis for BH3 protein family (Vesna 2018)

The Axolotl also shares characteristics with other species, as portrayed in both Figures. In Figure 9, the Eastern Newt (*Notophthalmus viridescens*) and Sub-Saharan Frogs (*Xenopus* sp) are most closely related with the Axolotl's p53 tumor suppressor gene. In Figure 9, the Danio species are most closely evolutionarily related to the Axolotl in the Beclin and Bim genes. Like this, as the Axolotl's characteristics are more studied and their similarities are discovered with other organisms, more evolutionary relationships can be drawn and eventually the entire Axolotl genome will be linked with other organisms in a detailed and specific phylogenetic tree.

CONCLUSION

Future Investigation
The Axolotl genome is still relatively recent and there is still much to explore, we have only begun to scratch the surface of the regenerative capabilities Axolotls possess and future research is required to obtain more information on that ability.

The next steps for our research are to continue research into Axolotl's and Human shared gene similarities and how genes can be manipulated to yield desirable results for regeneration.

For further research, topics can include: using comparisons for different human gene and finding the similarities between the human and Axolotl genome, studying the regenerative ability of other animals such as other salamanders as well, and uncovering the mysteries behind the other "superpowers" Axolotls have and utilizing them to help humans, such as Axolotl's being more resistant to cancer.

Funding

This study was supported by the SkoolMentor program.

Acknowledgments

The authors acknowledge the Molecular and Developmental Complexity Group (Alfredo Cruz Lab) at UGA-LANGEBIO for generating and providing the data.

REFERENCES

1. "15 Limb Loss Statistics that May Surprise You." *AccessProsthetics*, 18 Oct 2017. https://accessprosthetics.com/15-limb-loss-statistics-may-surprise/.

2. Aryal, Sagar. "Western Blotting - Introduction, Principle, and Applications." *Online Microbiology and Biology Notes*, 23 Oct 2017, *Microbe Notes*. https://microbenotes .com/western-blotting -introduction-principle-and-applications/.

3. "Axolotl". *San Diego Zoo*. https://animals.sandiegozoo.org/animals/axolotl

4. "Biology of Axolotls." *Axolotls*, http://www.axolotl.org/biology.htm.

5. Bryant DM, Johnson K, DiTommaso T, et al. A Tissue-Mapped Axolotl De Novo Transcriptome Enables Identification of Limb Regeneration Factors. *Cell Rep*. 2017;18(3):762-776. doi:10.1016/j.celrep.2016.12.063.

6. Caballero-Pérez J, Espinal-Centeno A, Falcon F, García-Ortega LF, Curiel-Quesada E, Cruz-Hernández A, Bako L, et al. Transcriptional landscapes of Axolotl (*Ambystoma mexicanum*). Dev Biol 2018. 433(2): 227-239

7. Dockrill, Peter. "Scientists Pinpoint Axolotl Genes Involved in Their Unique Body Regeneration." *ScienceAlert*, 30 Jan 2020. https://www.sciencealert.com/scientists-discover-axolotl-genes-involved-in-their-amazing-limb-regeneration.

8. Dunlap, Garrett. "Regeneration: What the Axolotl Can Teach us About Regrowing Human Limbs." *Harvard University Graduate School of Arts and Sciences*, 12 Jan 2018, *Science In The News*. http://sitn.hms.harvard.edu/flash/2018/regeneration - axolotl-can-teach-us-regrowing-human-limbs/

9. Farrow, Jonathan. "A is For Axolotl." *Rising Ape Collective*, 27 Feb 2017. https:// rising-ape.com/2015/02/27/a-is-for-axolotl/.

10. Gross, Michael. "Regenerative Lessons from the Axolotl." *Current Biology*, vol. 28, no. 5, 5 March 2018, Pages R187-R189, *ScienceDirect*. https://www.sciencedirect. com/science/article/pii/S0960982218302185

11. Jiang P, Nelson JD, Leng N, et al. Analysis of embryonic development in the unsequenced axolotl: Waves of transcriptomic upheaval and stability. *Dev Biol*. 2017;426(2):143-154. doi:10.1016/j.ydbio.2016.05.024.

12. Lewis, Tanya. "Missing Parts? Salamander Regenerative Secret Revealed." *LiveScience*, 20 May 2013. https://www.livescience.com/34513-how-salamanders-regenerate-lost-limbs.html.

13. Mahmood, Tahrin, and Ping-Chang Yang. "Western blot: technique, theory, and trouble shooting." *North American Journal of Medical Sciences* vol. 4,9 (2012): 429-34. doi:10.4103/1947-2714.100998

14. Mccusker, Catherine & Bryant, Susan & Gardiner, David. "The Axolotl Limb Blastema: Cellular and Molecular Mechanisms Driving Blastema Formation and Limb Regeneration in Tetrapods: The Axolotl Limb Blastema." *ResearchGate*, 2015. https://www.researchgate.net/publication/272755143_The_axolotl_limb_blastema_cel lular_and_molecular_mechanisms_driving_blastema_formation_and_limb_regenerati on_in_tetrapods_The_Axolotl_Limb_Blastema.

15. Preston, Elizabeth. "Salamander's Genome Guards Secrets of Limb Regrowth." *Quantamagazine*, 2 July 2018. https://www.quantamagazine.org/axolotl-genome - slowly-yields-secrets-of-limb-regrowth-20180702/.

16. Robertson, Sally. "Gene Expression Techniques." *News Medical LifeSciences*, 24 May 2019, https://www.news-medical.net/life-sciences/Gene-Expression-Techniques .aspx

17. Sibai, M., Parlayan, C., Tuğlu, P. *et al.* Integrative Analysis of Axolotl Gene Expression Data from Regenerative and Wound Healing Limb Tissues. *Sci Rep* 9, 20280 (2019). https://doi.org/10.1038/s41598-019-56829-6.
18. Travers, Julia. "This Salamander Resists Cancer and Regrows Limbs." *Labroots*, 8 Jan 2018. https://www.labroots.com/trending/videos/11340/this-salamander -resists-cancer-and-regrows-limbs
19. University of Nottingham. "Scientists bring cancer cells back under control." *ScienceDaily*. *ScienceDaily*, 18 January 2011. https://www.sciencedaily.com/releases/2011/01/110118122554.htm
20. Vesna, Bucan & Peck, Claas-Tido & Nasser, Inas & Liebsch, Christina & Vogt, Peter & Strauß, Sarah. (2018). "Identification of axolotl BH3-only proteins and expression in axolotl organs and apoptotic limb regeneration tissue." *ResearchGate*, August 2018, https://www.researchgate.net/figure/Phylogenetic-tree-of-BH3-only- proteins-Sequences-found-for-zebrafish-Xenopus-human-and_fig1_327263450.
21. Villiard, Eric & Brinkmann, Henner & Moiseeva, Olga & Mallette, Frederick & Ferbeyre, Gerardo & Roy, Stephane. "Urodele p53 tolerates amino acid changes found in p53 variants linked to human cancer." *ResearchGate*, 2007, https://www.researchgate.net/figure/Phylogenetic-tree-of-p53-protein-sequences-in-vertebrates-with-bootstrap-values-Maximum_fig3_5942083.