**SARS-CoV-2 induces double-stranded RNA-mediated innate immune responses in**

**respiratory epithelial derived cells and cardiomyocytes**

Yize Li [a,z,*,#], David M Renner[a,z,*], Courtney E Comar [a,z,*], Jillian N Whelan[a,z,*], Hanako M Reyes[a,z]

Fabian Leonardo Cardenas-Diaz[b,y], Rachel Truitt[b,x], Li Hui Tan[c], Beihua Dong[d], Konstantinos

Dionysios Alysandratos[e], Jessie Huang[e], James N. Palmer[c], Nithin D. Adappa[c], Michael A.

Kohanski[c], Darrell N. Kotton[e], Robert H Silverman[d],  Wenli Yang[b], Edward Morrisey[b,y],  Noam A.

Cohen[c,f,g],  Susan R Weiss [a,z,#, ^]

Departments of Microbiology[a], Medicine[b], Otorhinolaryngology[c], Institute for Regenerative

Medicine[x], Penn Center for Research on Coronaviruses and Other Emerging Pathogens[z], Penn-

CHOP Lung Biology Institute[y], Perlman School of Medicine at the University of Pennsylvania,

Philadelphia, PA, 19104

[d]Department of Cancer Biology, Lerner Research Institute, Cleveland Clinic, Cleveland, OH

44195 USA

[e]Department of Medicine, the Pulmonary Center, Center for Regenerative Medicine Boston

University School of Medicine, Boston, MA 02118, USA

[f]Corporal Michael J. Crescenz VA Medical Center, Philadelphia PA, 19104

[g]Monell Chemical Senses Center, Philadelphia, PA, 19104

#Address correspondence to: Susan R. Weiss, weisssr@pennmedicine.upenn.edu or Yize Li

yizeli@pennmedicine.upenn.edu

^lead contact: Susan R. Weiss, weisssr@pennmedicine.upenn.edu

     [*]These authors contributed equally to this work

**Summary**

Coronaviruses are adept at evading and/or antagonizing double-stranded RNA-induced host

antiviral pathways, including interferon signaling, OAS-RNase L and PKR while robust cytokine

responses characterize severe coronavirus disease. Knowledge of how newly emerged SARS-

CoV-2 interacts with these pathways is minimal. SARS-CoV-2 readily infects patient-derived nasal

epithelial cells and induced pluripotent stem cell-derived alveolar type 2 cells(iAT2) and

cardiomyocytes(iCM). Robust activation of interferons or RNase L is not observed, while PKR

activation is evident in iAT2 and iCM. In SARS-CoV-2 infected Calu-3 and A549$^{ACE2}$ lung derived

cell lines, activation of all pathways is observed, similar to a mutant MERS-CoV lacking innate

immune antagonists. Moreover, increased replication in *RNASEL* knockout A549$^{ACE2}$ cells,

implicates RNase L in restricting SARS-CoV-2. Finally, while SARS-CoV-2 is less adept at

antagonizing these host defense pathways compared to other coronaviruses, the innate immune

response is still generally weak. These host-virus interactions may contribute to the unique

pathogenesis of SARS-CoV-2.

49 **Keywords:** SARS-CoV-2; interferon, OAS-RNase L, PKR, viral accessory proteins

**Introduction**

Severe acute respiratory syndrome coronavirus (SARS-CoV)-2 emerged in China in late 2019, causing the COVID-19 pandemic with extensive morbidity and mortality, leading to major changes in day-to-day life in many parts of the world. This was the third lethal respiratory coronavirus, after SARS-CoV in 2002 and Middle East respiratory coronavirus (MERS-CoV) in 2012, to emerge from bats in the twenty first century. Although these viruses are all members of the *Betacoronavirus* genus (Llanes et al., 2020), each has caused a somewhat different pattern of pathogenesis and spread in humans (Fehr and Perlman, 2015); therefore it is important to understand how these viruses interact with their host. Additionally, recent reports indicate that ancestors of SARS-CoV-2 as well as other SARS-CoV-like viruses have been circulating in bats for decades, suggesting potential future instances of direct spread to humans (Boni et al., 2020).

Coronaviruses are enveloped viruses with large, positive-sense single-stranded (ss)RNA genomes of around 30kb that can infect a diverse range of mammals and other species. Coronaviruses use much of their genomes, including their approximately 20 kb Orf1ab replicase locus comprising the 5' two thirds of the genome, to encode proteins that antagonize host cell responses (Perlman and Netland, 2009). As a result they are remarkably adept at antagonizing host responses, in particular the double-stranded RNA (dsRNA)-induced pathways that are essential components of the host innate immune response (Cruz et al., 2011; Dedeurwaerder et al., 2014; Koetzner et al., 2010; Kopecky-Bromberg et al., 2007; Weiss and Navas-Martin, 2005). In addition, interspersed among the structural genes encoded in the 3' third of the genome are lineage-specific genes encoding accessory proteins, which are non-essential for RNA replication and variable among CoV lineages that further divide the *Betacoronavirus* genus (Cui et al., 2019). These accessory proteins often have functions in antagonizing host cell responses and thus likely contribute to discrepancies in pathogenesis and tropism observed among the different lineages (Comar et al., 2019; Kikkert, 2020; Zhao et al., 2012).

Like other RNA viruses, coronaviruses produce double-stranded RNA (dsRNA) early on during

the infection cycle as a result of genome replication and mRNA transcription (Sola et al., 2015).

Host cell pattern recognition receptors (PRRs) sense viral dsRNA as pathogenic non-self and

respond by activating several antiviral pathways critical for early defense against viral invasion.

DsRNA sensing by cytosolic PRRs can be divided into three key pathways – interferon (IFN)

production, ribonuclease L (RNase L) activation, and protein kinase R (PKR) activation (**Fig 1**)

(Hur, 2019). Detection of dsRNA by MDA5 during coronavirus infection (Roth-Cross et al., 2008),

leads to the production of type I ($\alpha/\beta$) and type III ($\lambda$) IFN. Upon binding to its specific cell surface

receptor, IFN triggers phosphorylation of STAT1 and STAT2 transcription factors, which then

induce expression of interferon stimulated genes (ISGs) with antiviral activities (Lopusna et al.,

2013; Platanias, 2005). In parallel, dsRNA is also sensed by oligoadenylate synthetases (OASs),

primarily OAS3, which synthesize 2',5'-linked oligoadenylates (2-5A)(Li et al., 2016; Whelan et

al., 2019). Generation of 2-5A induces dimerization and activation of ribonuclease L (RNase L),

leading to degradation of viral and host ssRNA. Finally, dsRNA sensing by protein kinase R (PKR)

will induce PKR autophosphorylation, permitting PKR to then phosphorylate the translation

initiation factor eIF2$\alpha$, which results in protein synthesis shutdown and restriction of viral

replication (Sadler and Williams, 2008). While RNase L and PKR antiviral activity are not

dependent on IFN production (Whelan et al., 2019), the genes encoding OASs and PKR are ISGs,

therefore these pathways can be activated and/or reinforced by IFN production. Similarly, RNase

L and PKR activation can promote IFN production, cellular stress, inflammation, and/or apoptotic

death (Banerjee et al., 2014; Castelli et al., 1997; Chakrabarti et al., 2015; Kang and Tang, 2012;

Malathi et al., 2007; Zhou et al., 1997), thus further reducing host cell viability.

Induction and inhibition of innate immune responses during infection with SARS-CoV-2 have yet

to be fully characterized. Furthermore, while it is well understood that SARS-CoV-2 enters the

human body through the upper respiratory tract, it is unclear which cell types of the upper and

lower respiratory system contribute to sustained infection and resulting disease in the airways

and elsewhere. We have performed SARS-CoV-2 infections of primary nasal epithelial cells,

induced pluripotent stem cell (iPSC)-derived alveolar type 2 cells (iAT2), and iPSC-derived

cardiomyocytes (iCM), which together represent the host tissues likely affected by clinical SARS-

CoV-2 infection (Hou et al., 2020; Sharma et al., 2020). We assessed viral replication in these

cell types as well as the degree of ensuing dsRNA-sensing responses. We also employed two

respiratory tract derived immune-competent cells lines, Calu-3 cells and A549 cells, to investigate

dsRNA-induced pathway activation during SARS-CoV-2 infection. In addition, we compared host

responses to SARS-CoV-2 with those of MERS-CoV and MERS-CoV-ΔNS4ab, a mutant lacking

expression of two dsRNA-induced innate immune pathway antagonists that we have

characterized previously (Comar et al., 2019).

115
116    **Results**
117
**SARS-CoV-2 replicates efficiently in cells derived from upper and lower respiratory tract.**

We compared the replication of SARS-CoV-2 and MERS-CoV in nasal epithelia-derived cells, a

relevant site of infection *in vivo*, from four different donors (**Fig 2A**). For each virus, replication

was similar in cells from all four individuals, although the extent of replication was somewhat

variable. The trends in replication kinetics, however, were significantly different between SARS-

CoV-2 and MERS-CoV infections. Replication of SARS-CoV-2 increased until 96hpi, but then

plateaued at nearly $10^6$ plaque-forming units (PFU)/ml. MERS-CoV replication peaked at 96hpi,

at a lower titer than SARS-CoV-2. Nasal epithelial cell cultures were stained with antibodies to

identify ciliated cells (anti-type IV β-tubulin), a key feature of this cell type, and either SARS-CoV-

2 or MERS-CoV nucleocapsid expression (anti-N protein) (**Fig 2B**). We detected abundant N

expressed in both SARS-CoV-2 and MERS-CoV infected cells, indicating that these cells were

sufficiently infected at 48 hours post infection (hpi). Interestingly, robust replication occurred in

these cultures, despite a very low level of ACE2 protein expression in cells from the three

individuals examined (**Fig 2C**).

We measured dsRNA-induced host responses to SARS-CoV-2 infection, including type I and type

III IFN mRNA induction, RNase L activation, and PKR activation, in the nasal cells. For RT-qPCR

analysis, we extracted RNA from SARS-CoV-2 infected cultures from four different donors at

120hpi. We verified that virus was replicating by quantifying viral genome copies from intracellular

RNA (**Fig S1A**). We then quantified mRNA expression of IFN-β (type I IFN), IFN-λ (type III IFN),

select ISGs induced downstream of IFN production (*OAS2*, *IFIT1*, *IFIH1*), and the neutrophil

attracting chemokine IL-8 (*CXCL8*), which has been implicated in nasal inflammation during viral

infection (Mukaida, 2003; Turner et al., 1998) (**Fig 2D**)**.** The data shown represents the average

of cells from three donors. There was some induction of type I and to a lesser extent type III IFN

mRNA, and minimal induction of the ISG mRNAs examined. Similarly, *CXCL8* encoding IL-8  was

barely induced. Interestingly, this may be at least partially due to high basal levels of IFN (notably

IFN-λ) and ISG (notably OAS2) mRNAs compared with other cell types examined below, which

would result in reporting of weak fold changes in mRNA levels compared with mock infected cells

(**Fig S2**). To further investigate this very weak ISG induction, using cells from the same donors

as the IFN/ISG mRNA quantification, we assessed the phosphorylation of STAT1, a transcription

factor that is itself encoded by an ISG, which is primarily a key mediator of type I and type III IFN

signaling. Upon infection, STAT1 is phosphorylated before complexing with STAT2 and IRF9 to

form ISGF3, which translocates to the nucleus where it mediates the activation of ISG

transcription (Stark and Darnell, 2012). Consistent with the weak activation of ISGs, there was no

evidence of phosphorylation of STAT1, likely due to low levels or transient phosphorylation (**Fig**

**2C**). In addition, we did not detect PKR activation, as indicated by the absence of phosphorylated

PKR and eIF2$\alpha$ in SARS-CoV-2 infected cells, while PKR and eIF2$\alpha$ were clearly expressed (**Fig**

**2C**). We also assessed activation of the OAS-RNase L pathway during SARS-CoV-2 infection of

cells from two of the same four donors. Since 28S and 18S ribosomal RNAs (rRNAs) are targeted

for degradation by activated RNase L, we can evaluate 28S and 18S rRNA integrity using a

Bioanalyzer as a readout for RNase L activation. The absence of any rRNA degradation in SARS-

CoV-2 infected cells (**Fig 2E**) indicated that RNase L was not activated despite abundant RNase

L protein expression (**Fig 2C**).

161
162
Next, we sought to examine host innate immune responses during infection of alveolar type 2

cells (AT2), a major target of SARS-CoV-2 infection in humans (Hou et al., 2020; Ng et al., 2016;

Qian et al., 2013). Initially, we attempted infections in lung organoid cultures, however they were

not sufficiently permissive to SARS-CoV-2. Thus, we employed induced pluripotent stem cell

(iPSC)-derived iAT2 cells (SPC2 line), expressing tdTomato from the endogenous locus of

surfactant protein-C (SFTPC), an AT2 cell specific marker  (Jacob et al., 2019). As in nasal cells,

virus replicated efficiently, reaching a titer of $10^6$ PFU/ml by 48hpi (**Fig 3A**). Staining of cultures

with an anti-N antibody showed that most of the iAT2 cells were infected, without obvious

cytopathic effect (CPE) during infection (**Fig 3B**). Notably, SARS-CoV-2 infection of iAT2 cells

was robust despite ACE2 expression being below the level of detection by immunoblotting (**Fig**

**3C**). We observed activation of the PKR pathway as indicated by both PKR and eIF2$\alpha$

phosphorylation by immunoblotting (**Fig 3C**). We extracted RNA from infected iAT2 cells for RT-

qPCR analysis, verified these cells were replicating virus by quantifying genome RNA copies (**Fig**

**S1B**), and assessed IFN/ISG induction. As with the nasal cells, we observed weak induction of

IFN-$\beta$ and IFN-$\lambda$ mRNA from mock infected and infected cells (**Fig 3D**), while MDA5 (Roth-Cross

et al., 2008) a dsRNA sensor in the pathway leading to IFN production during coronavirus

expression, is not detected (**Fig 3C**). We used the alphavirus Sindbis virus (SINV) as a positive

control, which we have previously shown induces robust activation of all dsRNA-induced

pathways (Comar et al., 2019). Surprisingly, we observed greater increases in OAS2 and IFIT

mRNA expression by SARS-CoV-2 compared with SINV (**Fig 3D**), but with minimal induction of

IFIH1 mRNA or protein (MDA5) (**Fig 3C&D**). However, we did not observe phosphorylation of

STAT1 (**Fig 3C**), as in the nasal cells above. Additionally, we did not observe any degradation of

rRNA in SARS-CoV-2 infected cells, and only slight degradation by SINV despite ample

expression of RNase L (**Fig 3E**), suggesting minimal activation of RNase L in iAT2 cells in general.

187
**SARS-CoV-2 replicates and induces innate immune responses in iPSC-derived**

**cardiomyocytes.** Since many COVID-19 patients experience cardiovascular symptoms and

pathology (Lindner et al., 2020; Shi et al., 2020), we investigated SARS-CoV-2 infection of iPSC

derived-cardiomyocytes (iCM). SARS-CoV-2 replicated robustly in these cells reaching titers of

approximately $10^6$ PFU/ml by 48hpi (**Fig 4A**), similar to replication in nasal and iAT2 cells. Cells

were stained with an antibody against cardiac troponin-T (cTnT) as a marker for cardiomyocytes,

and an antibody against the viral N protein to identify infected cells (**Fig 4B**). In addition, we

detected clear cytopathic effect (CPE), in iCM, which differed from infected nasal and iAT2 cells.

This CPE included syncytia resulting from cell-to-cell fusion, which is typical of coronaviruses and

a result of intracellular cleavage of the viral spike protein by the host enzyme furin (Belouzard et

al., 2009; de Haan et al., 2004; Gombold et al., 1993; Qiu et al., 2006; Yamada and Liu, 2009).

Interestingly, while we observed detectable ACE2 protein expression in mock infected cells in two

independent experiments, we observed loss of ACE2 expression upon SARS-CoV-2 infection,

consistent with a recent study (Sharma et al., 2020) (**Fig 4C**). We extracted RNA from mock

infected cells and cells infected with SARS-CoV-2 or SINV, verified that virus was replicating by

quantifying viral genome in intracellular RNA (**Fig S1C**), and quantified expression of mRNAs for

IFNs and select ISGs. We found low levels of IFN/ISGs transcript in iCM similar to the nasal and

iAT2 cells (**Fig D**), perhaps due to the undetectable levels of MDA5 and MAVS protein expression

in these cells (**Fig 4C**). SINV also induced host mRNAs weakly, with the exception of IFN-$\lambda$, in

these cells (**Fig 4D**). We observed no degradation of rRNA, suggesting an absence of RNase L

activation in iCM with SARS-CoV-2 or SINV (**Fig 4E**), despite clear infection with either virus (**Fig**

**S1C**). This was not surprising as there was no RNase L detectable by immunoblot in these cells

(**Fig 4C**). Finally, as in iAT2 cells, we observed phosphorylation of PKR and eIF2$\alpha$, indicating that

the PKR antiviral pathway is activated (**Fig 4C**).

212
213
214    **SARS-CoV-2 replicates in respiratory epithelial cell lines and induces dsRNA responsive**
215    **pathways.**
216
We aimed to further characterize the relationship between SARS-CoV-2 and dsRNA-induced host

response pathways, which were activated to variable levels in the different primary and iPSC

derived cells evaluated. We chose two respiratory epithelium-derived human cell lines, A549 and

Calu-3, both of which are immune competent and have been used for studies of SARS-CoV

(Blanco-Melo et al., 2020) and MERS-CoV (Comar et al., 2019; Thornbrough et al., 2016). A549

cells were not permissive to SARS-CoV-2, which correlated with undetectable levels of the SARS-

CoV-2 receptor ACE2 (**Fig S3**). Therefore, we generated A549 cells expressing the ACE2

receptor (A549$^{ACE2}$) by lentiviral transduction, and used two single cell clones, C44 and C34, for

all experiments (**Fig S3**). Both A549$^{ACE2}$ clones express high levels of ACE2 greater than the

endogenously expressed ACE2 in Calu-3 cells (**Fig S3**) and in the primary cells discussed above

(**Fig 2-4**).

We performed single step growth curves to measure replication of SARS-CoV-2 over the course

of one infectious cycle in A549$^{ACE2}$ cells, simian Vero-E6 cells, which are commonly used to

prepare SARS-CoV-2 stocks, and Calu-3 cells (clone HTB-55). SARS-CoV-2 replicated robustly

in A549$^{ACE2}$ and Vero-E6 cells (**Fig 5A**) but in comparison viral yields were lower in Calu-3 cells

(**Fig 5B**). Since Calu-3 cells also support MERS-CoV infection, we compared SARS-CoV-2

replication to that of MERS-CoV (WT) and MERS-CoV-ΔNS4ab, a mutant deleted in host cell

antagonists NS4a, a dsRNA-binding protein, and NS4b, a 2'5'-phosphodiesterase that prevents

RNase L activation and nuclear translocation of NF-κB (Canton et al., 2018; Comar et al., 2019).

Consistent with our previous work (Comar et al., 2019), MERS-CoV-ΔNS4ab reduced viral titers

from WT MERS-CoV levels, although they remained higher than SARS-CoV-2 titers (**Fig 5B**). To

further understand the replication of SARS-CoV-2, we stained A549, Calu-3 and Vero-E6 cells at

24 hpi with antibodies against viral N protein and viral dsRNA, including additional Calu-3 staining

at 48 hpi since replication kinetics are slower (**Fig 5C**). We observed cytopathic effect in all three

cell types, with N staining in the cytoplasm. Syncytia were observed in A549$^{ACE2}$ and Calu-3 cells,

but not in Vero cells (**Fig 5C**). We also observed viral dsRNA localized to perinuclear foci as we

and others have described during infection with other coronaviruses (Comar et al., 2019; Knoops

et al., 2008; Lundin et al., 2014; Rabouw et al., 2016).

We used RT-qPCR to quantify the induction of type I and type III IFNs and select ISGs at 24 and

48 hpi (**Fig 6A**), as well as the intracellular viral genome copies to verify replication (**Fig 6B**) in

A549$^{ACE2}$ cells. Using SINV as a positive control, we found relatively low levels of both IFNβ and

IFNλ mRNA at 24 and 48 hpi by SARS-CoV-2, compared to SINV (**Fig 6A**). Notably, IFN induction

was greater than observed in the nasal, iAT2, or iCM cells, possibly due to lower basal levels of

IFNβ, but not IFNλ, mRNA in the A549$^{ACE2}$ cells, which allow for greater fold changes over mock

infected cells (**Fig S2**). As observed previously with SINV and MERS-CoV, SARS-CoV-2 induced

more IFNλ than IFNβ, typical of A549 cells (Comar et al., 2019). Levels of ISG mRNAs were

variable, with SARS-CoV-2 inducing moderate levels of OAS2 and IFIT1 mRNAs, but only late in

infection (48 hpi), similar to those induced by SINV at 24 hpi **(Fig 6A)**. We observed minimal

effects on mRNA levels of IFIH1 and the cytokine CXCL8 at both timepoints (**Fig 6A**).

Furthermore, we did not detect any STAT1 phosphorylation at 24 hpi (**Fig 6C**), which correlates

with weak ISG expression, suggesting defective IFN signaling downstream of IFN production. The

data shown in Fig 6 were derived from A549$^{ACE2}$ clone C44; similar data were obtained for a

second clone, C33 (**Fig S4**).

We evaluated IFN/ISG responses in Calu-3 cells, which provided a second lung-derived cell line

that additionally supports both SARS-CoV-2 and MERS-CoV infection, allowing us to compare

host responses between the two lethal CoVs. We compared SARS-CoV-2 responses to both WT

MERS-CoV and mutant MERS-CoV-ΔNS4ab (**Fig 7A**). Although we observed reduced MERS-

CoV-ΔNS4ab infectious virus production compared with WT MERS-CoV (**Fig 5B**), we detected

similar intracellular viral genome levels of all three viruses (**Fig 7B**). We found previously that

MERS-CoV-ΔNS4ab induces higher levels of IFNs and ISGs compared to WT MERS-CoV, and

also activates RNase L and PKR (Comar et al., 2019). Herein, in Calu-3 cells, we observed

greater SARS-CoV-2 induction of IFN mRNAs as compared to A549$^{ACE2}$ cells (**Fig 6A&S4B**).

Interestingly, SARS-CoV-2 induced higher IFN mRNA levels than WT MERS-CoV at 24 and 48

hpi, which more closely resemble those induced by the IFN antagonist-deficient mutant MERS-

CoV-ΔNS4ab by 48hpi (**Fig 7A**). Similarly, SARS-CoV-2 generally induced more ISG mRNA than

WT MERS-CoV, and even more OAS2 mRNA than MERS-ΔNS4ab (**Fig 7A**). Induction of CXCL8

was weak for all viruses (**Fig 7A**). Notably, SARS-CoV-2 induced ISG mRNAs in Calu-3 (24hpi)

without the delay observed in A549$^{ACE2}$ cells. Consistent with earlier ISG mRNA induction during

infection, SARS-CoV-2 infection promotes phosphorylation of STAT1 in Calu-3 cells (**Fig 7C**) as

recently reported (Lokugamage et al., 2020). SARS-CoV-2 induction of pSTAT1 as well as rapid

IFIT1 and OAS2 mRNA induction suggests a similar host response to SARS-CoV-2 as that

observed during mutant MERS-CoV-ΔNS4ab infection, and not that of WT MERS-CoV infection.

282
**SARS-CoV-2 infection activates RNase L and PKR.** We assessed activation of the RNase L

pathway by analyzing intracellular rRNA from infected cells for RNA integrity, as described above.

We found that SARS-CoV-2 promotes rRNA degradation minimally at 24 hpi and more clearly at

48 hpi in A549$^{ACE2}$, using SINV as a positive control (**Fig 8A**). Evaluation of RNase L activation in

SARS-CoV-2, WT MERS-CoV, and MERS-CoV-ΔNS4ab infected Calu-3 cells showed SARS-

CoV-2 activation of RNase L to a similar extent as MERS-CoV-ΔNS4ab, which lacks both a

dsRNA binding protein and a RNase L-specific antagonist (Comar et al., 2019; Siu et al., 2014)

(**Fig 8B**). In contrast, as we previously reported, WT MERS-CoV failed to activate RNase L

(Comar et al., 2019; Thornbrough et al., 2016) (**Fig 8B**). We also observed activation of PKR as

indicated by phosphorylation of PKR and downstream eIF2α, in both A549$^{ACE2}$ cells (**Fig 8C**) and

Calu-3 cells (**Fig 8D**) infected with SARS-CoV-2. In Calu-3 cells, SARS-CoV-2 induced PKR

phosphorylation to a similar extent as MERS-CoV-ΔNS4ab, while WT MERS-CoV failed to induce

a response. These data are consistent with IFN/ISG induction data described above, suggesting

that SARS-CoV-2 may not antagonize dsRNA pathways as efficiently as MERS-CoV, but instead

induces host responses similar to those observed during MERS-CoV-ΔNS4ab infection.

The A549$^{ACE2}$ cells were valuable in that they provided a system with intact innate immune

responses that was also amenable to CRISPR-Cas9 engineering. Thus, we used the A549$^{ACE2}$

cells to construct additional cell lines with targeted deletions of *MAVS, RNASEL*, or *PKR*, as we

have done previously for parental A549 cells (Li et al., 2017; Li et al., 2016). We could then use

these cells to determine whether activation of IFN, RNase L, and/or PKR resulted in attenuation

of SARS-CoV-2 replication (Li et al., 2017; Li et al., 2016). We validated the knockout (KO)

A549$^{ACE2}$ cell lines by western blot (**Fig 9A**) and compared replication of SARS-CoV-2 in *MAVS*

KO, *RNASEL* KO and *PKR* KO cells with levels in WT A549$^{ACE2}$ cells (**Fig 9B**). Interestingly, there

was little effect on SARS-CoV-2 replication with MAVS or PKR expression absent. In *RNASEL*

KO cells at 48 hpi, virus replication was two- to four-fold higher compared to WT A549$^{ACE2}$ cells

(**Fig 9B**). While the difference in replication between *RNASEL* KO and WT was not extensive, it

was statistically significant in three independent experiments. Additionally, infected *RNASEL* KO

cells exhibited strikingly more CPE as compared with WT, *PKR* KO, or *MAVS* KO cells, as

demonstrated by crystal violet-staining of infected cells (**Fig 9C**). We also assessed rRNA

degradation and, as expected, found that rRNA remained intact in the *RNASEL* KO A549$^{ACE2}$

cells, which further validated these cells. However, rRNA was degraded in *PKR* or *MAVS* KO

cells, indicating RNase L activation in both of these cell types (**Fig 9D**). These data are consistent

with our previous findings that activation of the RNase L pathway does not depend on MAVS

signaling in A549 cells infected with SINV or Zika virus (ZIKV) (Li et al., 2019; Whelan et al.,

2019). RNase L activation instead occurs in parallel with IFN production (**Fig 1**), and is likely

dependent on basal levels of OAS gene expression (Birdwell et al., 2016).

320
321
322
323  **Discussion**
324
SARS-CoV-2 is the causative agent of the severe acute respiratory, COVID-19. SARS-CoV-2

infection begins in the ciliated cells of the nasal epithelium and spreads to the lower respiratory

tract, possibly by aspiration of infectious nasal secretions, infecting alveolar type 2 (AT2) cells

and causing acute respiratory distress syndrome (ARDS) (Hou et al., 2020). Despite this typical

axis of infection, many infected individuals exhibit symptoms and pathology in other organ

systems, including the cardiovascular system (Puelles et al., 2020; Shi et al., 2020). Here, we

present data expanding our understanding of cellular responses from each one of these critical

areas. We focused on understanding the early host-viral interactions that trigger activation of the

dsRNA-sensing pathways, including the RNase L, PKR, and type I and III IFN pathways. We

evaluated responses to SARS-CoV-2 infection in primary nasal epithelia-derived upper airway

cells and iPSC-derived type II airway (iAT2) cells, which represent major sites of infection during

COVID-19 disease, as well as iPSC-derived cardiomyocytes (iCM), another likely target of

infection (Sharma et al., 2020). To complement these studies, we used two lung derived

transformed cell lines, Calu-3 cells and two different A549$^{ACE2}$ clones, to more mechanistically

dissect activation and antagonism of these pathways by SARS-CoV-2. We found that the extent

of IFN induction and signaling is variable among the primary cell types and cell lines used, but is

consistently only poorly induced. Interestingly, we show that SARS-CoV-2 infection results in

more IFN signaling (phosphorylation of STAT1 and IFN/ISG expression) when compared to

MERS-CoV in Calu-3 cells. We also found that SARS-CoV-2 activates RNase L and PKR in both

cell lines used and PKR in iAT2 cells and iCM, but not in primary nasal cells. Using KO cell lines,

we demonstrate that RNase L expression significantly impacts SARS-CoV-2 viral titers and CPE

observed during infection. These data suggest that while SARS-CoV-2 is generally a weak

activator of dsRNA-induced responses in cells of the respiratory and cardiovascular systems,

SARS-CoV-2 is less adept at antagonizing host responses than MERS-CoV.

Nasal cells are the initial replication site of SARS-CoV-2 and MERS-CoV. In quantifying virus

replication in infected nasal cell culture, we found that SARS-CoV-2 replicates to higher titer than

MERS-CoV, and the time period for shedding of virus is much longer (**Fig 2A**). We suggest that

this longer period of replication in nasal cells and stronger immune responses in Calu-3 cells may

in part explain why SARS-CoV-2 is less virulent, yet more contagious than MERS-CoV. Indeed

for SARS-CoV-2, $R_0$=5.7 (Sanche et al., 2020) while for MERS-CoV, ($R_0$=0.45)(Oh et al., 2016).

Infection of all three primary cell types – nasal cells, iAT2 cells, and iCM – resulted in high levels

of SARS-CoV-2 replication, while only cardiomyocytes exhibited obvious CPE (**Figs 2-4**). It was

easily observed with syncytia formation in both lines, but the kinetics of viral spread were slower

in Calu-3 (**Fig 5**). IFA staining with viral dsRNA-specific antibody (J2) showed SARS-CoV-2

dsRNA localized to perinuclear areas in A549$^{ACE2}$ and Calu-3 cells, which is typical of coronavirus

infection (**Fig 5**). The protein expression level of the SARS-CoV-2 host receptor ACE2 (Lu et al.,

2020; Wan et al., 2020; Zhou et al., 2020) in primary cells and Calu-3 cells was either low or

undetectable, indicating that high levels of receptor are not necessary for productive infection (**Fig**

**2-4&S3**). This is similar to previous observations in the murine coronavirus (MHV) system where

viral receptor CEACAM1a is very weakly expressed in the mouse brain, a major site of infection,

and particularly in neurons, the most frequently infected cells, as measured by RT-qPCR for

mRNA and antibody staining for protein (Bender et al., 2010).

The canonical IFN production and signaling pathways activated by the sensing of dsRNA, an

obligate intermediate in viral genome replication and mRNA transcription, provide a crucial early

antiviral response (**Fig 1**). The role of IFN responses during coronavirus infection is complex and

at times contradictory. For example, SARS-CoV induces poor IFN signaling during early infection

in mice, and treatment with IFN improves outcome. However, IFN production later in infection has

a pathogenic effect (Channappanavar and Perlman, 2017). Similarly, weak IFN responses have

been observed during initial stages of SARS-CoV-2 infection, but IFN produced later may

contribute to the strong inflammatory responses and resulting immunopathology observed during

SARS-CoV-2 infection cytokine storms (Giamarellos-Bourboulis et al., 2020; Hadjadj et al., 2020).

Furthermore, murine coronavirus (MHV) induces minimal levels of IFN in cell culture and *in vivo*,

yet IFN$\alpha/\beta$ receptor (IFNAR) KO mice lacking type I IFN signaling responses experience ectopic

virus spread and rapid death (Roth-Cross et al., 2008).

While IFNs may contribute to pathogenesis later on in infection, coronaviruses, including the lethal

human viruses, often prevent these responses early on during infection in both animal models

and humans (Arabi et al., 2017; Channappanavar et al., 2016; Channappanavar et al., 2019).

Antagonism of dsRNA-induced antiviral pathways has been well characterized for lineage a

betacoronaviruses (subgenera embeco; for example, murine coronavirus MHV) (Zhao et al.,

2012) and lineage c betacoronaviruses (merbeco; MERS-CoV and other related bat viruses)

(Comar et al., 2019), however there is less known about lineage b (sarbeco) betacoronaviruses,

including SARS-CoV (2002) and SARS-CoV-2. A recent report described a cohort of COVID-19

patients with impaired IFN and innate immune responses that correlated with enhanced disease

severity, suggesting that subversion of these host responses is important for severe SARS-CoV-

2-mediated clinical outcomes (Hadjadj et al., 2020). Thus, it is important to understand the basic

biology of SARS-CoV-2 interaction with the host innate immune system.

We and others have previously found that both MHV (lineage a) and MERS-CoV (lineage c)

betacoronaviruses induce only minimal type I and type III IFNs, and fail to activate RNase L or

PKR pathways (Rabouw et al., 2016; Thornbrough et al., 2016; Ye et al., 2007; Zhao et al., 2012;

Zhao et al., 2011). We found that SARS-CoV-2, like other betacoronaviruses, induced limited

amounts of type I and type III IFN mRNAs, although this was somewhat variable among the cell

types examined. Using SINV as a control for robust activation of IFN, we detected low levels of

type I and type III IFN mRNA in nasal cell, iAT2 cells, and iCM (**Fig 2-4**). However, we observed

higher levels of OAS2, an ISG, relative to SINV in iAT2 cells (**Fig 3D**). As we have observed

among murine cells, we saw vastly different levels of basal expression of both IFN and ISG

mRNAs among the cell types infected (**Fig S2**) (Li and Weiss, 2016; Zhao et al., 2013; Zhao et

al., 2011). It is understood that higher basal levels of innate immune response mRNAs typically

result in a lower threshold for activation of corresponding responses. Interestingly, we observed

significantly higher basal levels, especially IFNλ, in (uninfected) nasal cells as compared to iAT2

cells and iCM (**Fig S2A**). As major barrier cells, we speculate that this may be important for

protection as these cells are more exposed to infectious agents in the environment. Indeed, it is

well documented that type III IFNs serve as an added defense for epithelial cells, which may

perhaps explain some of the differences observed in basal gene expression between nasal cells

and iCM (Ank et al., 2008; Forero et al., 2019; Galani et al., 2017).  As previously reported in heart

tissue, the iCM expressed undetectable levels of both MAVS and RNase L, (Uhlén et al., 2015;

Zhou et al., 1997), which is possibly to protect the heart from excessive inflammation.

In A549$^{ACE2}$ cells, SARS-CoV-2 induced low levels of IFNλ and IFNβ mRNAs and somewhat

higher ISG mRNA by 48 hpi, as compared with SINV (**Fig 6A**). We observed greater increases

in IFN induction in Calu-3 compared to A549$^{ACE2}$ (**Fig 7A**), which may be at least partially due to

higher basal levels of IFNs in the Calu-3 cells (**Fig S2**). Calu-3 cells were employed to directly

compare the host response to SARS-CoV-2 infection with that of MERS-CoV and mutant MERS-

CoV-ΔNS4ab, which lacks the NS4a and NS4b proteins that inhibit IFN production and signaling

(Canton et al., 2018; Comar et al., 2019; Rabouw et al., 2016). In Calu-3 cells, SARS-CoV-2

induced more IFN mRNA than WT MERS-CoV, closer to the level of MERS-CoV-ΔNS4ab (**Fig**

**7A**). Furthermore, SARS-CoV-2 induced higher levels of ISG mRNAs than MERS-CoV and, in

the case of OAS2, higher than MERS-CoV-ΔNS4ab as well.  Consistent with this, in Calu-3 cells

SARS-CoV-2  and  MERS-CoV-ΔNS4ab,  but  not  WT  MERS-CoV,  promoted  STAT1

phosphorylation (**Fig 7C**), which leads to ISG transcription and antiviral responses. Overall, our

results displayed a trend of relatively weak IFN responses induced by SARS-CoV-2 in airway

epithelial cells with limited ISG induction, when compared with host responses to viruses from

other families. Additionally, our data show that enhanced IFN/ISG responses in Calu-3 cells

restrict virus production, while lower host responses in A549$^{ACE2}$ cells correlate with higher viral

titers (**Fig 5**). Considering how robust ACE2 expression appears dispensable for infection of

some cell types (nasal, iAT2, Calu-3), these data also indicate that stronger innate immune

responses may be more effective at restricting SARS-CoV-2 replication than low ACE2

expression level.

We found that SARS-CoV-2 was unable to prevent activation of RNase L and PKR, although to

different extents among the cell types, unlike two other betacoronaviruses, MHV and MERS-CoV,

which shut down these pathways (Comar et al., 2019; Ye et al., 2007; Zhao et al., 2012). We

observed PKR activation as indicated by phosphorylation of PKR and eIF2$\alpha$ in SARS-CoV-2

infected iAT2 (**Fig 3C**) and iCM (one/two experiments) (**Fig 4C**), but not in nasal cells (**Fig 2C**).

However, we did not detect rRNA degradation indicative of RNase L activation in these cell types

(**Fig 2E, 3E, 4E**). Activation of both RNase L and PKR were observed in A549[ACE2] and Calu-3

cells during infection with SARS-CoV-2 (**Fig 8**). In Calu-3 cells, this contrasted MERS-CoV and

was more similar to MERS-CoV-ΔNS4ab. Previous studies have shown that MERS-CoV NS4a

restricts phosphorylation of PKR by binding dsRNA, reducing its accessibility to PKR (Comar et

al., 2019; Rabouw et al., 2016). Additionally, MERS-CoV NS4b, a 2'-5' phosphodiesterase,

prevents RNase L activation by degrading 2-5A, the small molecular activator of RNase L (Comar

et al., 2019; Thornbrough et al., 2016). Current understanding of SARS-CoV-2 protein function

infers an absence of these types of protein antagonists, therefore it is not surprising that both of

these pathways are activated during infection of both A549[ACE2] and Calu-3. Indeed, MERS-CoV-

ΔNS4ab attenuation compared to WT MERS-CoV, as well as lower SARS-CoV-2 titers than those

of MERS-CoV (**Fig 5B**), may be at least in part due to RNase L and PKR activation in addition to

IFN/ISG induction in Calu-3 cells.

Consistent with previous studies involving the alphavirus SINV, we found that neither MERS-CoV-

ΔNS4ab nor SARS-CoV-2 activated dsRNA-induced pathway responses as robustly as SINV (Li

et al., 2016; Whelan et al.). We suggest this may be due to CoV antagonists encoded by the nsp

genes of the replicase locus. For CoVs, conserved replicase proteins have been identified as

important modulators of innate immune invasion, including nsp1, nsp14, nsp15, and nsp16

(Gordon et al., 2020; Kindler et al., 2017; Perlman and Netland, 2009; Volk et al., 2020). Most

notably, nsp15 encodes an endoribonuclease (EndoU) that has been shown in the MHV system

to restrict dsRNA accumulation and thus limit activation of both RNase L and PKR (Deng et al.,

2017; Kindler et al., 2017). A very recent report using a replicon system concludes that SARS-

CoV-2 nsp1 and nsp6 are more effective antagonists than the homologous proteins of SARS-CoV

or MERS-CoV (Xia et al., 2020). Nevertheless, increased, albeit modest, replication and

enhanced cell death in SARS-CoV-2 infected *RNASEL* KO cells indicates that this pathway is

activated and indeed restricts replication and downstream cell death caused by SARS-CoV-2

infection (**Fig 9B&C**). In contrast, we found that *PKR* KO had no effect on viral titer. This is

consistent with our previous finding that MERS-CoV infection inhibits host protein synthesis

independent of PKR, so that PKR phosphorylation during MERS-CoV-ΔNS4ab infection did not

lead to further reduction (Comar et al., 2019). Finally, KO of *MAVS* and the consequent loss of

IFN production had no significant effect on viral titer or cell death. This is similar to our previous

findings demonstrating that RNase L activation can occur independent of virus-induced IFN

production with other viruses (SINV) (Li et al., 2019) and ZIKV (Whelan et al., 2019) in A549

cells, as well as for MHV in murine bone marrow-derived macrophages (Birdwell et al., 2016).

This underscores the importance of the RNase L and PKR antiviral pathways, which can be

activated early in infection upon concurrent dsRNA sensing by OAS, PKR, and MDA5 receptors

before IFN is produced. Alternatively, these pathways can be activated in cells infected by virus

that produce low levels of IFN only late in infection, as we observe here with SARS-CoV-2. Further

studies are required to determine whether activation of PKR or RNase L during  SARS-CoV-2

infection results in functional outcomes characteristic of these pathways, including inhibition of

protein synthesis, induction of apoptosis, cleavage of viral RNA, or induction of inflammatory

responses (**Fig 1**). Interestingly, we observed possible RNase L-induced apoptosis in the SARS-

CoV-2 infected A549$^{ACE2}$ WT, *MAVS* KO, and *PKR* KO cells, when compared with mock infected

counterparts (**Fig 9C**). However, *RNASEL* KO cells displayed the most cell death among the four

cell lines, suggesting that virus-induced cell lysis in the *RNASEL* KO cells where viral titers are

highest (**Fig 9B**) is more detrimental to cells than RNase L-induced programmed cell death.

There have been few studies on the interaction of the prototype lineage b betacoronavirus,

SARS-CoV (2002), with these dsRNA-induced pathways, with most of the studies describing IFN

antagonists involving overexpressed protein and overexpressed artificial IFN promoter

sequences for readouts. In this way, it was reported that Orf3b, Orf6 and nucleocapsid (N) protein

are antagonists of IFN production and signaling (Kopecky-Bromberg et al., 2007). The role of the

SARS-CoV Orf6 encoded protein as an IFN antagonist was subsequently confirmed in the context

of virus infection, and further defined as Orf6 binding to host nuclear importins to prevent

translocation of STAT1 and other transcription factors, thus impairing induction of antiviral ISGs

(Frieman et al., 2007; Sims et al., 2013). SARS-CoV-2 encodes an Orf6 protein with 69%

homology to SARS-CoV Orf6 (Lokugamage et al., 2020), as well as a similar, truncated Orf3b

which has been shown by overexpression studies to antagonize IFN mRNA transcription (Konno

et al., 2020). Strikingly, recent work using overexpression systems of SARS-CoV-2 Orf6 found

that it interacts with host KPNA2, the same target of the SARS-CoV Orf6 protein (Xia et al., 2020).

However, further investigation is required to determine how similarly the proteins from the two

viruses function in the context of infection.

We have shown that SARS-CoV-2 activates dsRNA-induced innate immune responses to levels

similar to those of a MERS-CoV mutant lacking two accessory proteins that antagonize these

pathways, which highlights the distinctions among coronaviruses in interacting with these

pathways. However, like MERS-CoV and MHV, SARS-CoV-2 induces limited and late IFN/ISG

responses, indicating that proteins antagonizing innate immune responses are likely encoded.

Our future studies will focus on identifying specific innate immunity antagonists among lineage b

betacoronavirus accessory proteins as well as conserved nsp proteins. The recent development

of reverse genetics systems for SARS-CoV-2 (Hou et al., 2020; Thi Nhu Thao et al., 2020; Xie et

al., 2020) will allow us to study these proteins during viral infection. This will provide a system

complete with coronavirus-mediated alterations of cellular architecture typically nonexistent in

overexpression systems, which is critical to fully appreciate the functions of these proteins

(Knoops et al., 2008; Lundin et al., 2014; Sola et al., 2015). Our future studies will reveal key

features of SARS-CoV-2 biology, as well as those of other currently circulating SARS-related

viruses with pandemic potential.

**Materials and Methods**

**Viruses.** SARS-CoV-2 (USA-WA1/2020 strain)  was obtained from BEI and propagated in Vero-

E6 cells. The genome RNA was sequenced was found to be identical to GenBank: MN985325.1**.**

Recombinant MERS-CoV and MERS-CoV-ΔNS4ab were described previously (Comar et al.,

2019) and were propagated in Vero-CCL81 cells. Sindbis virus Girdwood (G100) was obtained

from Dr. Mark Heise, University of North Carolina, Chapel Hill, and was prepared as previously

described (Suthar et al., 2005). Sendai virus (SeV) strain Cantell (Basler et al., 2003)  was

obtained from Dr. Carolina B. Lopez (University of Pennsylvania).

**Cell lines**. African green monkey kidney Vero cells (E6)  or (CCL81) (obtained from ATCC) were

cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco catalog no. 11965),

supplemented with 10% fetal bovine serum (FBS), 100 U/ml of penicillin, 100 μg/ml streptomycin,

50 μg/ml gentamicin, 1mM sodium pyruvate, and 10mM HEPES. Human A549 cells(Verified by

ATCC) were cultured in RPMI 1640 (Gibco catalog no. 11875) supplemented with 10% FBS, 100

U/ml of penicillin, and 100 μg/ml streptomycin. Human HEK 293T cells were cultured in DMEM

supplemented with 10% FBS and 1 mM sodium pyruvate. Human Calu-3 cells (clone HTB-55)

were cultured in MEM supplemented with 20% FBS without antibiotics.

540
541  **Primary cell cultures**
542

**Human sinonasal air liquid interface** (**ALI) cultures.** Sinonasal mucosal specimens were acquired from residual clinical material obtained during sinonasal surgery subsequent to approval fromm The University of Pennsylvania Institutional Review Board. ALI cultures were established from enzymatically dissociated human sinonasal epithelial cells (HSEC) as previously described (Lee et al., 2017; Lee et al., 2014) and grown to confluence with bronchial epithelial basal medium (BEBM; Lonza, Alpharetta, GA) supplemented with BEGM Singlequots (Lonza), 100 U/ml penicillin and 0.25 µg /ml amphotericin B for 7 days. Cells were then trypsinized and seeded on porous polyester membranes (2-3× $10^4$ cells per membrane) in cell culture inserts (Transwell-clear, diameter 12 mm, 0.4 µm pores; Corning, Acton, MA). Five days later the culture medium was removed from the upper compartment and the epithelium was allowed to differentiate by using the differentiation medium consisting of 1:1 DMEM (Invitrogen, Grand Island, NY) and BEBM (Lonza), supplemented with BEGM Singlequots (Lonza) with 0.1 nM retinoic acid (Sigma-Aldrich), 100 UI/ml penicillin, 0.25 µg /ml amphotericin B and 2% Nu serum (Corning) in the basal compartment. Cultures were fed every three days for 6 weeks prior to infection with SARS-CoV-2. The day prior infection, the cells were fed and the apical side of the cultures were washed with 100µl of warm PBS X 3.

**Alveolar organoids and 2D cultures**. iPSC (SPC2 iPSC line, clone SPC2-ST-B2, Boston University) derived alveolar epithelial type 2 cells (iAT2s) were differentiated and maintained as alveolospheres embedded in 3D Matrigel in CK+DCI media, as previously described (Jacob et al., 2019). iAT2s were passaged approximately every two weeks by dissociation into single cells via the sequential application of dispase (2mg/ml, Thermo Fisher Scientific, 17105-04) for 1h at 37°C and 0.05% trypsin (Invitrogen, 25300054) for 15min at 37°C and re-plated at a density of 400 cells/µl of Matrigel (Corning, 356231) in CK+DCI media supplemented with ROCK inhibitor for the first 48h, as previously described (Jacob et al., 2019). For generation of 2D alveolar cells for viral infection, alveolospheres were dispersed into single cells, then plated on pre-coated 1/30

Matrigel plates at a cell density of 125,000 cells/cm2 using CK+DCI media with ROCK inhibitor

for the first 48h and then the medium was changed to CK+DCI media at day 3 and infected with

SARS-CoV-2 virus.

**Cardiomyocytes**. Experiments involving the use of human iPSCs were approved by the

University of Pennsylvania Embryonic Stem Cell Research Oversight Committee.  The iPSC line

(PENN123i-SV20) used for cardiomyocyte generation was derived by the UPenn iPSC core as

previously described (Pashos et al., 2017; Yang et al., 2015).  This line has been deposited at the

WiCell repository (Wicell.org).   iPSCs were maintained on Geltrex (Thermofisher Scientific)-

coated plates in iPS-Brew XF (Miltenyi Biotec) media at 37$^{o}$C in 5% $CO_2$/5% $O_2$/90% air humidified

atmosphere.    Cells were passaged every 5-7 days using Stem-MACS Passaging Solution

(Miltenyi Biotec).     Differentiation of SV20 into cardiomyocytes (iCMs) was performed using

previously described protocols (Laflamme et al., 2007; Palpant et al., 2017).  In general, iCMs

were >95% positive for cardiac Troponin T staining by FACS.  Day 18-25 differentiated cells were

replated and used for viral infection experiments.

**Generation of A549$^{ACE2}$ cells.** A549$^{ACE2}$ cells were constructed by lentivirus transduction of

*hACE2*. The plasmid encoding the cDNA of *hACE2* was purchased from Addgene. The cDNA

was amplified using forward primer 5′-ACTCTAGAATGTCAAGCTCTTCCTGGCTCCTTC-3′ and

reverse                                         primer                                         5′-

TTGTCGACTTACGTAGAATCGAGACCGAGGAGAGGGTTAGGGATAGGCTTACCAAAGGAG

GTCTGAAC ′-3(Contained V5 tag sequences). The fragment containing hACE2-V5 was digested

by the XbaI and SalI restriction enzymes from the hACE2 cDNA and was cloned into pLenti-GFP

(Addgene) in place of green fluorescent protein (GFP), generating pLenti-hACE2-V5. The

resulting plasmids were packaged in lentiviruses pseudotyped with vesicular stomatitis virus

glycoprotein G (VSV-G) to establish the gene knock-in cells. Supernatants harvested 48 hours

post-transfection were used for transduction into A549 cells. Forty-eight hours after transduction,

cells were subjected to hygromycin (1 mg/ml) selection for 3 days and single-cell cloned. Clones

were screened for ACE2 expression and susceptibility to SARS-CoV-2 replication.

**CRISPR/Cas9 engineered cells**. *RNASEL*, *PKR* and *MAVS* KO A549$^{ACE2}$ cells were constructed

using the same Lenti-CRISPR system and guide RNA sequences as previously described  (Li et

al., 2017; Li et al., 2016).

**Viral growth kinetics**. The nasal ALI cultures were apically infected with SARS-CoV-2(MOI=5),

MERS-CoV(MOI=5)**.** Briefly, viral stocks were diluted in nasal cell media, 50μl was added to each

well, the cells were incubated in 37°C for one hour, then the virus was removed and the cells were

wash three times with 200μl of PBS. For viral growth curves, at indicated time points, 200 μl of

PBS was added to the apical surface, collected 5 minutes later and frozen for subsequent analysis

of shed virus by plaque assay. The inserts were transferred to new 24-well plates with fresh media

after each collection. For iAT2 or iCM, cells were plated in 12 or 6-well plates, 4X10$^5$ cells (iAT2)

or 6.25X10$^5$ cells per well (iCM), cells were infected with SARS-CoV-2 at MOI=5 (iAT2) or

MOI=1(iCM).  At 6, 24, 48 hours postinfection, 200μl of supernatant were harvested and stored

in -80°C for infectious virus titration. For infections, Cell lines were plated in 12-well plates, A549

and Vero-E6 at 5X10$^5$ cells per well) and Calu-3 at 3X10$^5$ cells per well). Viruses were diluted in

serum-free RPMI (A549 infections) or serum-free DMEM (Vero infections) or serum-free MEM

(Calu-3 infections) and added to cells for absorption for 1 hour at 37°C. Cells were washed three

times with PBS and fed with DMEM or RPMI +2% FBS for Vero and RPMI infections, respectively,

or 4% FBS in MEM for Calu-3 infections (Thornbrough et al., 2016). For virus titration 200 μl of

supernatant was collected at the times indicated and stored at -80°C for plaque assay on Vero-

E6 (SARS-CoV-2) or Vero-CCL81 (MERS-CoV) cells as previously described (Scobey et al.,

2013).

**Plaque assay.** Briefly virus supernatant was 10-fold serial diluted and inoculum was absorbed on

Vero cells for 1 hour at 37°C. Inoculum was overlaid with DMEM plus agarose (either 0.7% or

0.1%) and incubated for 3 days at 37°C. Cells were fixed with 4% paraformaldehyde and stained

with 1% crystal violet for counting plaques. All infections and virus manipulations were conducted

in a biosafety level 3 (BSL-3) laboratory using appropriate and approved personal protective

equipment and protocols.

**Immunofluorescent staining.** For nasal ALI culture,  following 48 hours of infection, the cultures

were fixed in 4% paraformaldehyde at room temperature for 30 minutes.  The transwell supports

were washed 3 times with PBS prior to excision of the membrane containing the cells. The cells

were permeabilized with 0.2% Triton X-100 in PBS and then immersed in PBS with 0.2% Triton

X-100, 10% normal donkey serum, and 1% BSA for 60 min at room temperature.  Primary

antibody incubation was incubated overnight at 4 ºC (Type IV tubulin, Abcam ab11315, rabbit anti

SARS-CoV-2 Nucleocapsid protein, GeneTex, Irvine, CA).  Visualization was carried out with Alex

Fluor®-conjugated donkey anti-mouse or anti-rabbit IgGs(Thermo-Fisher) (1:1000; 60 min

incubation at room temperature).  Confocal images were acquired with an Olympus Fluoview

System (Z-axis step 0.5µm; sequential scanning).  For iAT2, the cell monolayer was fixed using

4% paraformaldehyde (PFA) for 30min, 1X PBS was used to removed PFA and proceed with

antibody staining. Fixed cells were treated with a blocking solution containing 0.1% Triton X-100

and 5% donkey serum in 1X PBS for 30min. Immunostaining was performed for SARS-CoV-2

nucleocapsid protein expression using the SARS-CoV-2 nucleocapsid antibody at 1:1000 dilution

in blocking solution incubated for 30min. After washing primary antibody away, a secondary Alexa

fluor 488 donkey anti-rabbit IgG(H+L) antibody(Thermo-Fisher) was used at 1:400 dilution in

blocking solution and incubated for 30min. Secondary antibody was washed away with 1X PBS

and DAPI was used for nuclear staining at 2.5μg/ml. iCM were fixed in 4% paraformaldehyde and

permeabilized with 0.1% Triton X-100 for 15min. Cells were blocked with 10% normal donkey

serum (Sigma D9663) in 0.2% Tween 20 (Biorad 170-6531) for 1hr. Antibodies against cardiac

troponin T (cTnT, Abcam ab8295; 1:100 mouse) and SARS-CoV-2 nucleocapsid were incubated

with cells in blocking solution overnight at 4 °C. Donkey anti-mouse Alexa Fluor 647 (Invitrogen

A31571) and Donkey anti-rabbit Alexa Fluor 488 (Invitrogen A21206) were diluted 1:250 in

blocking solution and incubated with cells for 2hr at RT. Slides were mounted in Slowfade Gold

anti-fade reagent with DAPI (Invitrogen S36939). Images were acquired with BZ-X710 all-in-one

fluorescence microscope equipped with BZ-X Viewer software (Keyence Corporation). At the

indicated times post-infection cells were fixed onto glass coverslips (Calu-3 coverslips were

coated with rat tail collagen type-1: Cell Applications, Inc. Cat. # 122-20) with 4%

paraformaldehyde for 30 minutes at room temperature. Cells were then washed three times with

PBS and permeabilized for 10 minutes with PBS+0.1% Triton-X100. Cells were then blocked in

PBS and 3% BSA for 30-60 minutes at room temperature. Primary antibodies were diluted in

block buffer and incubated on a rocker at room temperature for one hour. Cells were washed

three times with block buffer and then incubated rocking at room temperature for 60 minutes with

secondary antibodies diluted in block buffer. Finally, cells were washed twice with block buffer

and once with PBS, and nuclei stained with DAPI diluted in PBS (2ng/uL final concentration).

SARS-CoV-2 nucleoprotein and dsRNA (J2,1:1000, Scions) were detected. Secondary

antibodies were from Invitrogen: goat anti-mouse IgG AlexaFluor594 (A-11005) for J2 and goat

anti-rabbit IgG AlexaFluor488 (A-11070) for nucleocapsid. Coverslips were mounted onto slides

for analysis by widefield microscopy with Nikon Eclipse Ti2 using a Nikon 40x/0.95NA Plan APO

objective and NikonDS-Qi1Mc-U3 12 bit camera. Images were processed using Fiji/Image J

software.

**Western immunoblotting.** Cells were washed once with ice-cold PBS and lysates harvested at

the indicated times post infection with lysis buffer (1% NP40, 2mM EDTA, 10% glycerol, 150mM

NaCl, 50mM Tris HCl) supplemented with protease inhibitors (Roche – complete mini EDTA-free

protease inhibitor) and phosphatase inhibitors (Roche – PhosStop easy pack). After 5 minutes

lysates were harvested, incubated on ice for 20 minutes, centrifuged for 20 minutes at 4°C and

supernatants mixed 3:1 with 4x Laemmli sample buffer. Samples were heated at 95°C for 5

minutes, then separated on 4-15% SDS-PAGE, and transferred to polyvinylidene difluoride

(PVDF) membranes. Blots were blocked with 5% nonfat milk or 5% BSA and probed with

antibodies (table below) diluted in the same block buffer. Primary antibodies were incubated

overnight at 4°C or for 1 hour at room temperature. All secondary antibody incubation steps were

done for 1 hour at room temperature. Blots were visualized using Thermo Scientific SuperSignal

west chemiluminescent substrates (Cat #: 34095 or 34080). Blots were probed sequentially with

antibodies and in between antibody treatments stripped using Thermo scientific Restore western

blot stripping buffer (Cat #: 21059).

| Primary Antibody | Antibody species | Blocking buffer | Dilution | Catalog number |
|---|---|---|---|---|
| p-PKR (phospho-T446) [E120] | rabbit | 5% milk/TBST | 1 : 1000 | Abcam 32036 |
| PKR (D7F7) | rabbit | 5% milk/TBST | 1:1000 | Cell Signaling Technology 12297S |
| p-eif2α (S51) | rabbit | 5% BSA/TBST | 1:1000 | Cell Signaling Technology 9721S |
| eif2α | rabbit | 5% BSA/TBST | 1:1000 | Cell Signaling Technology 9722S |
| GAPDH (14C10) | rabbit | 5% milk/TBST | 1:2000 | Cell Signaling Technology 2118S |
| SARS-CoV-2 N | rabbit | 5% milk/TBST | 1:2000 | GTX135357 (Gentex) |
| MERS-CoV N | mouse | 5% milk/TBST | 1:2000 | 40068-MM10 (Sino Biological) |
| pSTAT1 (Tyr701) | rabbit | 5% BSA/TBST | 1:1000 | Cell Signaling Technology 7649 |

| | | | | |
|---|---|---|---|---|
| STAT1 | mouse | 5% BSA/TBST | 1:1000 | Santa Cruz (C136): SC-464 |
| ACE2 | rabbit | 5% milk/TBST | 1:1000 | Cell Signaling Technology 4355S |
| MAVS | rabbit | 5% milk/TBST | 1:1000 | Cell Signaling Technology 24930S |
| V5 | rabbit | 5% milk/TBST | 1:1000 | Cell Signaling Technology 13202S |
| RNase L | mouse | 5% milk/TBST | 1:1000 | gift from Dr. Robert Silverman (Cleveland Clinic) |
| MDA5 | rabbit | 5% milk/TBST | 1:1000 | Cell Signaling Technology 5321S |
| **Secondary Antibody** | | | | |
| goat anti-rabbit IgG | HRP linked | same as primary | 1:3000 | Cell Signaling Technology 7074S |
| goat anti-mouse IgG | HRP linked | same as primary | 1:3000 | Cell Signaling Technology 7076S |

**Quantitative PCR (RT-qPCR).**

A549, Calu-3, and iAT2 cells were lysed at indicated times post infection in RLT buffer and DNase-

treated before total RNA was extracted using the RNeasy Plus Mini Kit (Qiagen). RNA from iCM

and nasal cells was extracted using TRIzol-LS (Ambion), and DNase-treated using the DNA-

*free*$^{TM}$ Kit (Invitrogen). RNA was reverse transcribed into cDNA with a High Capacity cDNA

Reverse Transcriptase Kit (Applied Biosystems). cDNA was amplified using specific RT-qPCR

primers (see Table below), iQ$^{TM}$ SYBR$^{®}$ Green Supermix (Bio-Rad), and the QuantStudio$^{TM}$ 3

PCR system (Thermo Fisher). Host gene expression displayed as fold change over mock-infected

samples was generated by first normalizing cycle threshold ($C_T$) values to 18S rRNA to generate

$\Delta C_T$ values ($\Delta C_T$ = $C_T$ gene of interest - $C_T$ 18S rRNA). Next, $\Delta(\Delta C_T)$ values were determined by

subtracting the mock-infected $\Delta C_T$ values from the virus-infected samples. Technical triplicates

were averaged and means displayed using the equation $2^{-\Delta(\Delta CT)}$. For basal expression levels, $C_T$

values were normalized to 18S rRNA to generate $\Delta C_T$ values ($\Delta C_T$ = $C_T$ gene of interest - $C_T$ 18S

rRNA), and displayed as $2^{-\Delta Ct}$. Basal expression levels were also calculated as fold change over

A549$^{ACE2}$ clone 44 using the equation $2^{-\Delta(\Delta CT)}$. $\Delta(\Delta C_T)$ values were calculated by subtracting $\Delta C_T$

values from each cell type from the $\Delta C_T$ value of A549$^{ACE2}$ clone 44. Absolute quantification of

SARS-CoV-2 and MERS-CoV genomes was calculated using a standard curve generated from

serially diluted known concentrations of a digested plasmid containing the region of interest. For

SARS-CoV-2, construct pcDNA6B-nCoV-NSP12-FLAG encoding the RDRP gene (gift from Dr.

George Stark, Cleveland Clinic) was digested with Xho1 and purified by Qiagen QIAquick PCR

Purification Kit to be used as a standard in the RT-qPCR reaction. For MERS-CoV, cDNA MERS-

D1 (Scobey et al., 2013) containing basepairs 12259–15470 of the MERS-CoV genome was

digested with BgII and purified by Qiagen QIAquick PCR Purification Kit to be used as a standard

in the RT-PCR reaction. Copy numbers were generated by standard curve analysis in the

QuantStudio™ 3 software, and copy numbers per ug RNA were calculated based on the volume

of cDNA used in the qPCR reaction, and concentration of RNA used to generated cDNA. Primer

sequences are as follows:

| | Forward primer (5' to 3') | Reverse primer (5' to 3') |
|---|---|---|
| *IFNL1* | CGCCTTGGAAGAGTCACTCA | GAAGCCTCAGGTCCCAATTC |
| *OAS2* | TTCTGCCTGCACCACTCTTCACGAC | GCCAGTCTTCAGAGCTGTGCCTTTG |
| *IFIT1* | 5'-TGGTGACCTGGGGCAACTTT | AGGCCTTGGCCCGTTCATAA |
| *IFNB* | GTCAGAGTGGAAATCCTAAG | ACAGCATCTGCTGGTTGAAG |
| GAPDH | GCAAATTCCATGGCACCGT | TCGCCCCACTTGATTTTGG |
| *IFIH1* | GCACAGAGCGGTAGACCCTGCTT | AGGCCTTGGCCCGTTCATAA |
| *CXCL8* | GAGAGTGATTGAGAGTGGACCAC | CACAACCCTCTGCACCCAGTTT |
| 18S rRNA | TTCGATGGTAGTCGCTGTGC | CTGCTGCCTTCCTTGAATGTGGTA |
| SARS-CoV-2 genome (nsp12/RdRp) | GGTAACTGGTATGATTTCG | CTGGTCAAGGTTAATATAGG |

| MERS-CoV genome (nsp7) | GCACATCTGTGGTTCTCCTCTCT | AAGCCCAGGCCCTACTATTAGC |
|---|---|---|

**Analyses of RNase L-mediated rRNA degradation.** RNA was harvested with buffer RLT

(Qiagen RNeasy #74106) or Trizol-LS (Ambion) and analyzed on an RNA chip with an Agilent

Bioanalyzer using the Agilent RNA 6000 Nano Kit and its prescribed protocol as we have

described previously (Cat #: 5067-1511).

**Statistical analysis.** All statistical analyses and plotting of data were performed using GraphPad

Prism software (GraphPad Software, Inc., CA). SARS-CoV-2 and MERS-CoV replication trends

in nasal cells were analyzed by two-way ANOVA comparing averaged titers from all four donor

cells for each virus at each timepoint. MERS-CoV and MERS-CoV-ΔNS4ab viral replication and

primary cell RT-qPCR gene expression between SARS-CoV-2 and SINV were analyzed by paired

Student $t$ test. RT-qPCR analysis in A549$^{ACE2}$ cells was analyzed by one-way ANOVA, comparing

SARS-CoV-2 at each timepoint to SINV. RT-qPCR analysis in Calu-3 cells was analyzed by two-

way ANOVA, comparing SARS-CoV-2 at each timepoint to MERS-CoV and MERS-CoV-ΔNS4ab.

SARS-CoV-2 replication in A549$^{ACE2}$ WT cells compared with A549$^{ACE2}$ KO cells was analyzed by

two-way ANOVA. Displayed significance is determined by p-value (P), where * = P < 0.05; ** = P

< 0.01; *** = P < 0.001; **** = P < 0.0001; ns = not significant.

**Acknowledgements**

We thank Nicholas Parenti for technical help and Dr. Nikki Tanneti for reading the manuscript.

This work was supported by NIH grants AI140442 and supplement for SARS-CoV-2 (SRW),

AI104887 (SRW and RHS); funds from Penn Center for Coronavirus Research and Other

Emerging Pathogens (SRW, YL); NIH grants U01HL148857, R01HL087825, U01HL134745 and

R01HL132999 (EM); VA administration grant CX001617 (NAC);  NIH grants U01TR001810, N01

75N92020C00005, R01HL095993, and an Evergrande MassCPR award (DNK, JH, and KDA).

RT and WY were supported in part by institutional funds from the University of Pennsylania

Perelman School of Medicine to the iPSC Core and by NIH grant U01TR001810. DMR was

supported in part by T32-AI055400 and CEC was supported in part by T32 NS-007180,

**Author Contributions**

Conceptualization: YL, CEC, DMR, SRW
Methodology: YL CEC, DMR, JNW, HMR, WY, NAC, JNP, NDA, MAK, EM, RHS, SRW.
Investigation, Performed experiments: YL, CEC, DMR, JNW, HMR, FLC-D, RT, LHT, BD
Writing –Original Draft: SRW, JNW
Writing –Review & Editing: YL, CEC, DMR, JNW, HMR, SRW, EM, NAC, WY, DNK
Funding Acquisition: SRW, WY, EM, NC, RHS, DNK
Resources: WY, NC, EM, RHS, SRW
Supervision: WY, NC, EM, RHS, SRW

**Declaration of Interests**

756    The authors declare no competing interests.

**References**

Ank, N., Iversen, M.B., Bartholdy, C., Staeheli, P., Hartmann, R., Jensen, U.B., Dagnaes-Hansen,
F., Thomsen, A.R., Chen, Z., Haugen, H.*, et al.* (2008). An important role for type III interferon
(IFN-lambda/IL-28) in TLR-induced antiviral activity. J Immunol *180*, 2474-2485.

Arabi, Y.M., Balkhy, H.H., Hayden, F.G., Bouchama, A., Luke, T., Baillie, J.K., Al-Omari, A., Hajeer,
A.H., Senga, M., Denison, M.R.*, et al.* (2017). Middle East Respiratory Syndrome. N Engl J Med
*376*, 584-594.

Banerjee, S., Chakrabarti, A., Jha, B.K., Weiss, S.R., and Silverman, R.H. (2014). Cell-type-specific
effects of RNase L on viral induction of beta interferon. mBio *5*, e00856-00814.

Basler, C.F., Mikulasova, A., Martinez-Sobrido, L., Paragas, J., Muhlberger, E., Bray, M., Klenk,
H.D., Palese, P., and Garcia-Sastre, A. (2003). The Ebola virus VP35 protein inhibits activation of
interferon regulatory factor 3. Journal of virology *77*, 7945-7956.

Belouzard, S., Chu, V.C., and Whittaker, G.R. (2009). Activation of the SARS coronavirus spike
protein via sequential proteolytic cleavage at two distinct sites. Proceedings of the National
Academy of Sciences of the United States of America *106*, 5871-5876.

Bender, S.J., Phillips, J.M., Scott, E.P., and Weiss, S.R. (2010). Murine coronavirus receptors are
differentially expressed in the central nervous system and play virus strain-dependent roles in
neuronal spread. Journal of virology *84*, 11030-11044.

Birdwell, L.D., Zalinger, Z.B., Li, Y., Wright, P.W., Elliott, R., Rose, K.M., Silverman, R.H., and
Weiss, S.R. (2016). Activation of RNase L by murine coronavirus in myeloid cells is dependent on
basal Oas gene expression and independent of virus-induced interferon. Journal of virology.

Blanco-Melo, D., Nilsson-Payant, B.E., Liu, W.C., Uhl, S., Hoagland, D., Moller, R., Jordan, T.X.,
Oishi, K., Panis, M., Sachs, D.*, et al.* (2020). Imbalanced Host Response to SARS-CoV-2 Drives
Development of COVID-19. Cell *181*, 1036-1045 e1039.

Boni, M.F., Lemey, P., Jiang, X., Lam, T.T., Perry, B.W., Castoe, T.A., Rambaut, A., and Robertson,
D.L. (2020). Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for the
COVID-19 pandemic. Nat Microbiol.

Canton, J., Fehr, A.R., Fernandez-Delgado, R., Gutierrez-Alvarez, F.J., Sanchez-Aparicio, M.T.,
Garcia-Sastre, A., Perlman, S., Enjuanes, L., and Sola, I. (2018). MERS-CoV 4b protein interferes
with the NF-kappaB-dependent innate immune response during infection. PLoS pathogens *14*,
e1006838.

Castelli, J.C., Hassel, B.A., Wood, K.A., Li, X.L., Amemiya, K., Dalakas, M.C., Torrence, P.F., and
Youle, R.J. (1997). A study of the interferon antiviral mechanism: apoptosis activation by the 2-
5A system. The Journal of experimental medicine *186*, 967-972.

Chakrabarti, A., Banerjee, S., Franchi, L., Loo, Y.M., Gale, M., Jr., Nunez, G., and Silverman, R.H.
(2015). RNase L activates the NLRP3 inflammasome during viral infections. Cell host & microbe
*17*, 466-477.

Channappanavar, R., Fehr, A.R., Vijay, R., Mack, M., Zhao, J., Meyerholz, D.K., and Perlman, S.
(2016). Dysregulated Type I Interferon and Inflammatory Monocyte-Macrophage Responses
Cause Lethal Pneumonia in SARS-CoV-Infected Mice. Cell host & microbe *19*, 181-193.

Channappanavar, R., Fehr, A.R., Zheng, J., Wohlford-Lenane, C., Abrahante, J.E., Mack, M.,
Sompallae, R., McCray, P.B., Jr., Meyerholz, D.K., and Perlman, S. (2019). IFN-I response timing
relative to virus replication determines MERS coronavirus infection outcomes. The Journal of
clinical investigation *129*, 3625-3639.

Channappanavar, R., and Perlman, S. (2017). Pathogenic human coronavirus infections: causes
and consequences of cytokine storm and immunopathology. Semin Immunopathol *39*, 529-539.
819
820    Comar, C.E., Goldstein, S.A., Li, Y., Yount, B., Baric, R.S., and Weiss, S.R. (2019). Antagonism of
821    dsRNA-Induced Innate Immune Pathways by NS4a and NS4b Accessory Proteins during MERS
822    Coronavirus Infection. mBio *10*.
823
824    Cruz, J.L., Sola, I., Becares, M., Alberca, B., Plana, J., Enjuanes, L., and Zuniga, S. (2011).
825    Coronavirus gene 7 counteracts host defenses and modulates virus virulence. PLoS pathogens
826    *7*, e1002090.
827
828    Cui, J., Li, F., and Shi, Z.L. (2019). Origin and evolution of pathogenic coronaviruses. Nature
829    reviews Microbiology *17*, 181-192.
830
831    de Haan, C.A., Stadler, K., Godeke, G.J., Bosch, B.J., and Rottier, P.J. (2004). Cleavage inhibition
832    of the murine coronavirus spike protein by a furin-like enzyme affects cell-cell but not virus-cell
833    fusion. Journal of virology *78*, 6048-6054.
834
835    Dedeurwaerder, A., Olyslaegers, D.A., Desmarets, L.M., Roukaerts, I.D., Theuns, S., and
836    Nauwynck, H.J. (2014). ORF7-encoded accessory protein 7a of feline infectious peritonitis virus
837    as a counteragent against IFN-alpha-induced antiviral response. The Journal of general virology
838    *95*, 393-402.
839
840    Deng, X., Hackbart, M., Mettelman, R.C., O'Brien, A., Mielech, A.M., Yi, G., Kao, C.C., and Baker,
841    S.C. (2017). Coronavirus nonstructural protein 15 mediates evasion of dsRNA sensors and limits
842    apoptosis in macrophages. Proceedings of the National Academy of Sciences of the United
843    States of America *114*, E4251-E4260.
844
845    Fehr, A.R., and Perlman, S. (2015). Coronaviruses: an overview of their replication and
846    pathogenesis. Methods Mol Biol *1282*, 1-23.
847
848    Forero, A., Ozarkar, S., Li, H., Lee, C.H., Hemann, E.A., Nadjsombati, M.S., Hendricks, M.R., So,
849    L., Green, R., Roy, C.N.*, et al.* (2019). Differential Activation of the Transcription Factor IRF1
850    Underlies the Distinct Immune Responses Elicited by Type I and Type III Interferons. Immunity
851    *51*, 451-464 e456.
852
853    Frieman, M., Yount, B., Heise, M., Kopecky-Bromberg, S.A., Palese, P., and Baric, R.S. (2007).
854    Severe acute respiratory syndrome coronavirus ORF6 antagonizes STAT1 function by
855    sequestering nuclear import factors on the rough endoplasmic reticulum/Golgi membrane.
856    Journal of virology *81*, 9812-9824.
857
Galani, I.E., Triantafyllia, V., Eleminiadou, E.E., Koltsida, O., Stavropoulos, A., Manioudaki, M.,
Thanos, D., Doyle, S.E., Kotenko, S.V., Thanopoulou, K.*, et al.* (2017). Interferon-lambda

Mediates Non-redundant Front-Line Antiviral Protection against Influenza Virus Infection
without Compromising Host Fitness. Immunity *46*, 875-890 e876.

Giamarellos-Bourboulis, E.J., Netea, M.G., Rovina, N., Akinosoglou, K., Antoniadou, A.,
Antonakos, N., Damoraki, G., Gkavogianni, T., Adami, M.E., Katsaounou, P.*, et al.* (2020).
Complex Immune Dysregulation in COVID-19 Patients with Severe Respiratory Failure. Cell host
& microbe *27*, 992-1000 e1003.

Gombold, J.L., Hingley, S.T., and Weiss, S.R. (1993). Fusion-defective mutants of mouse
hepatitis virus A59 contain a mutation in the spike protein cleavage signal. Journal of virology
*67*, 4504-4512.

Gordon, D.E., Jang, G.M., Bouhaddou, M., Xu, J., Obernier, K., White, K.M., O'Meara, M.J.,
Rezelj, V.V., Guo, J.Z., Swaney, D.L.*, et al.* (2020). A SARS-CoV-2 protein interaction map reveals
targets for drug repurposing. Nature *583*, 459-468.

Hadjadj, J., Yatim, N., Barnabei, L., Corneau, A., Boussier, J., Smith, N., Pere, H., Charbit, B.,
Bondet, V., Chenevier-Gobeaux, C.*, et al.* (2020). Impaired type I interferon activity and
inflammatory responses in severe COVID-19 patients. Science.

Hou, Y.J., Okuda, K., Edwards, C.E., Martinez, D.R., Asakura, T., Dinnon, K.H., 3rd, Kato, T., Lee,
R.E., Yount, B.L., Mascenik, T.M.*, et al.* (2020). SARS-CoV-2 Reverse Genetics Reveals a Variable
Infection Gradient in the Respiratory Tract. Cell *182*, 429-446 e414.

Hur, S. (2019). Double-Stranded RNA Sensors and Modulators in Innate Immunity. Annu Rev
Immunol *37*, 349-375.

Jacob, A., Vedaie, M., Roberts, D.A., Thomas, D.C., Villacorta-Martin, C., Alysandratos, K.D.,
Hawkins, F., and Kotton, D.N. (2019). Derivation of self-renewing lung alveolar epithelial type II
cells from human pluripotent stem cells. Nature protocols *14*, 3303-3332.

Kang, R., and Tang, D. (2012). PKR-dependent inflammatory signals. Science signaling *5*, pe47.

Kikkert, M. (2020). Innate Immune Evasion by Human Respiratory RNA Viruses. J Innate Immun
*12*, 4-20.

Kindler, E., Gil-Cruz, C., Spanier, J., Li, Y., Wilhelm, J., Rabouw, H.H., Zust, R., Hwang, M.,
V'Kovski, P., Stalder, H.*, et al.* (2017). Early endonuclease-mediated evasion of RNA sensing
ensures efficient coronavirus replication. PLoS pathogens *13*, e1006195.

Knoops, K., Kikkert, M., Worm, S.H., Zevenhoven-Dobbe, J.C., van der Meer, Y., Koster, A.J.,
Mommaas, A.M., and Snijder, E.J. (2008). SARS-coronavirus replication is supported by a
reticulovesicular network of modified endoplasmic reticulum. PLoS Biol *6*, e226.

Koetzner, C.A., Kuo, L., Goebel, S.J., Dean, A.B., Parker, M.M., and Masters, P.S. (2010).
Accessory protein 5a is a major antagonist of the antiviral action of interferon against murine
coronavirus. Journal of virology *84*, 8262-8274.
907
908 Konno, Y., Kimura, I., Uriu, K., Fukushi, M., Irie, T., Koyanagi, Y., Nakagawa, S., and Sato, K.
909 (2020). SARS-CoV-2 ORF3b is a potent interferon antagonist whose activity is further increased
910 by a naturally occurring elongation variant. bioRxiv, 2020.2005.2011.088179.
911
912 Kopecky-Bromberg, S.A., Martinez-Sobrido, L., Frieman, M., Baric, R.A., and Palese, P. (2007).
913 Severe acute respiratory syndrome coronavirus open reading frame (ORF) 3b, ORF 6, and
914 nucleocapsid proteins function as interferon antagonists. Journal of virology *81*, 548-557.
915
916 Laflamme, M.A., Chen, K.Y., Naumova, A.V., Muskheli, V., Fugate, J.A., Dupras, S.K., Reinecke,
917 H., Xu, C., Hassanipour, M., Police, S.*, et al.* (2007). Cardiomyocytes derived from human
918 embryonic stem cells in pro-survival factors enhance function of infarcted rat hearts. Nat
919 Biotechnol *25*, 1015-1024.
920
921 Lee, R.J., Hariri, B.M., McMahon, D.B., Chen, B., Doghramji, L., Adappa, N.D., Palmer, J.N.,
922 Kennedy, D.W., Jiang, P., Margolskee, R.F.*, et al.* (2017). Bacterial d-amino acids suppress
923 sinonasal innate immunity through sweet taste receptors in solitary chemosensory cells.
924 Science signaling *10*.
925
926 Lee, R.J., Kofonow, J.M., Rosen, P.L., Siebert, A.P., Chen, B., Doghramji, L., Xiong, G., Adappa,
927 N.D., Palmer, J.N., Kennedy, D.W.*, et al.* (2014). Bitter and sweet taste receptors regulate
928 human upper respiratory innate immunity. The Journal of clinical investigation *124*, 1393-1405.
929
930 Li, Y., Banerjee, S., Goldstein, S.A., Dong, B., Gaughan, C., Rath, S., Donovan, J., Korennykh, A.,
931 Silverman, R.H., and Weiss, S.R. (2017). Ribonuclease L mediates the cell-lethal phenotype of
932 double-stranded RNA editing enzyme ADAR1 deficiency in a human cell line. eLife *6*.
933
934 Li, Y., Banerjee, S., Wang, Y., Goldstein, S.A., Dong, B., Gaughan, C., Silverman, R.H., and Weiss,
935 S.R. (2016). Activation of RNase L is dependent on OAS3 expression during infection with
936 diverse human viruses. Proceedings of the National Academy of Sciences of the United States of
937 America *113*, 2241-2246.
938
939 Li, Y., Dong, B., Wei, Z., Silverman, R.H., and Weiss, S.R. (2019). Activation of RNase L in Egyptian
940 Rousette Bat-Derived RoNi/7 Cells Is Dependent Primarily on OAS3 and Independent of MAVS
941 Signaling. mBio *10*.
942
943 Li, Y., and Weiss, S.R. (2016). Antagonism of RNase L Is Required for Murine Coronavirus
944 Replication in Kupffer Cells and Liver Sinusoidal Endothelial Cells but Not in Hepatocytes.
945 Journal of virology *90*, 9826-9832.
946

947　Lindner, D., Fitzek, A., Brauninger, H., Aleshcheva, G., Edler, C., Meissner, K., Scherschel, K.,
948　Kirchhof, P., Escher, F., Schultheiss, H.P.*, et al.* (2020). Association of Cardiac Infection With
949　SARS-CoV-2 in Confirmed COVID-19 Autopsy Cases. JAMA Cardiol.
950

951　Llanes, A., Restrepo, C.M., Caballero, Z., Rajeev, S., Kennedy, M.A., and Lleonart, R. (2020).
952　Betacoronavirus Genomes: How Genomic Information has been Used to Deal with Past
953　Outbreaks and the COVID-19 Pandemic. Int J Mol Sci *21*.
954

955　Lokugamage, K.G., Hage, A., de Vries, M., Valero-Jimenez, A.M., Schindewolf, C., Dittmann, M.,
956　Rajsbaum, R., and Menachery, V.D. (2020). Type I interferon susceptibility distinguishes SARS-
957　CoV-2 from SARS-CoV. Journal of virology.
958

959　Lopusna, K., Rezuchova, I., Betakova, T., Skovranova, L., Tomaskova, J., Lukacikova, L., and
960　Kabat, P. (2013). Interferons lambda, new cytokines with antiviral activity. Acta Virol *57*, 171-
961　179.
962

963　Lu, R., Zhao, X., Li, J., Niu, P., Yang, B., Wu, H., Wang, W., Song, H., Huang, B., Zhu, N.*, et al.*
964　(2020). Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for
965　virus origins and receptor binding. Lancet *395*, 565-574.
966

967　Lundin, A., Dijkman, R., Bergstrom, T., Kann, N., Adamiak, B., Hannoun, C., Kindler, E.,
968　Jonsdottir, H.R., Muth, D., Kint, J.*, et al.* (2014). Targeting membrane-bound viral RNA synthesis
969　reveals potent inhibition of diverse coronaviruses including the middle East respiratory
970　syndrome virus. PLoS pathogens *10*, e1004166.
971

972　Malathi, K., Dong, B., Gale, M., Jr., and Silverman, R.H. (2007). Small self-RNA generated by
973　RNase L amplifies antiviral innate immunity. Nature *448*, 816-819.
974

975　Mukaida, N. (2003). Pathophysiological roles of interleukin-8/CXCL8 in pulmonary diseases.
976　American journal of physiology Lung cellular and molecular physiology *284*, L566-577.
977

978　Ng, D.L., Al Hosani, F., Keating, M.K., Gerber, S.I., Jones, T.L., Metcalfe, M.G., Tong, S., Tao, Y.,
979　Alami, N.N., Haynes, L.M.*, et al.* (2016). Clinicopathologic, Immunohistochemical, and
980　Ultrastructural Findings of a Fatal Case of Middle East Respiratory Syndrome Coronavirus
981　Infection in the United Arab Emirates, April 2014. Am J Pathol *186*, 652-658.
982

983　Oh, M.D., Park, W.B., Choe, P.G., Choi, S.J., Kim, J.I., Chae, J., Park, S.S., Kim, E.C., Oh, H.S., Kim,
984　E.J.*, et al.* (2016). Viral Load Kinetics of MERS Coronavirus Infection. N Engl J Med *375*, 1303-
985　1305.
986

987　Palpant, N.J., Pabon, L., Friedman, C.E., Roberts, M., Hadland, B., Zaunbrecher, R.J., Bernstein,
988　I., Zheng, Y., and Murry, C.E. (2017). Generating high-purity cardiac and endothelial derivatives
989　from patterned mesoderm using human pluripotent stem cells. Nature protocols *12*, 15-31.
990

Pashos, E.E., Park, Y., Wang, X., Raghavan, A., Yang, W.L., Abbey, D., Peters, D.T., Arbelaez, J.,
Hernandez, M., Kuperwasser, N*., et al.* (2017). Large, Diverse Population Cohorts of hiPSCs and
Derived Hepatocyte-like Cells Reveal Functional Genetic Variation at Blood Lipid-Associated
Loci. Cell Stem Cell *20*, 558-+.

Perlman, S., and Netland, J. (2009). Coronaviruses post-SARS: update on replication and
pathogenesis. Nature reviews Microbiology *7*, 439-450.

Platanias, L.C. (2005). Mechanisms of type-I- and type-II-interferon-mediated signalling. Nature
reviews Immunology *5*, 375-386.

Puelles, V.G., Lutgehetmann, M., Lindenmeyer, M.T., Sperhake, J.P., Wong, M.N., Allweiss, L.,
Chilla, S., Heinemann, A., Wanner, N., Liu, S*., et al.* (2020). Multiorgan and Renal Tropism of
SARS-CoV-2. N Engl J Med *383*, 590-592.

Qian, Z., Travanty, E.A., Oko, L., Edeen, K., Berglund, A., Wang, J., Ito, Y., Holmes, K.V., and
Mason, R.J. (2013). Innate immune response of human alveolar type II cells infected with severe
acute respiratory syndrome-coronavirus. American journal of respiratory cell and molecular
biology *48*, 742-748.

Qiu, Z., Hingley, S.T., Simmons, G., Yu, C., Das Sarma, J., Bates, P., and Weiss, S.R. (2006).
Endosomal proteolysis by cathepsins is necessary for murine coronavirus mouse hepatitis virus
type 2 spike-mediated entry. Journal of virology *80*, 5768-5776.

Rabouw, H.H., Langereis, M.A., Knaap, R.C., Dalebout, T.J., Canton, J., Sola, I., Enjuanes, L.,
Bredenbeek, P.J., Kikkert, M., de Groot, R.J*., et al.* (2016). Middle East Respiratory Coronavirus
Accessory Protein 4a Inhibits PKR-Mediated Antiviral Stress Responses. PLoS pathogens *12*,
e1005982.

Roth-Cross, J.K., Bender, S.J., and Weiss, S.R. (2008). Murine coronavirus mouse hepatitis virus
is recognized by MDA5 and induces type I interferon in brain macrophages/microglia. Journal of
virology *82*, 9829-9838.

Sadler, A.J., and Williams, B.R. (2008). Interferon-inducible antiviral effectors. Nature reviews
Immunology *8*, 559-568.

Sanche, S., Lin, Y.T., Xu, C., Romero-Severson, E., Hengartner, N., and Ke, R. (2020). High
Contagiousness and Rapid Spread of Severe Acute Respiratory Syndrome Coronavirus 2.
Emerging infectious diseases *26*, 1470-1477.

Scobey, T., Yount, B.L., Sims, A.C., Donaldson, E.F., Agnihothram, S.S., Menachery, V.D.,
Graham, R.L., Swanstrom, J., Bove, P.F., Kim, J.D*., et al.* (2013). Reverse genetics with a full-
length infectious cDNA of the Middle East respiratory syndrome coronavirus. Proceedings of
the National Academy of Sciences of the United States of America *110*, 16157-16162.

1035

1036 Sharma, A., Garcia, G., Jr., Wang, Y., Plummer, J.T., Morizono, K., Arumugaswami, V., and
1037 Svendsen, C.N. (2020). Human iPSC-Derived Cardiomyocytes Are Susceptible to SARS-CoV-2
1038 Infection. Cell Rep Med *1*, 100052.

1039

1040 Shi, S., Qin, M., Shen, B., Cai, Y., Liu, T., Yang, F., Gong, W., Liu, X., Liang, J., Zhao, Q*., et al.*
1041 (2020). Association of Cardiac Injury With Mortality in Hospitalized Patients With COVID-19 in
1042 Wuhan, China. JAMA Cardiol.

1043

1044 Sims, A.C., Tilton, S.C., Menachery, V.D., Gralinski, L.E., Schafer, A., Matzke, M.M., Webb-
1045 Robertson, B.J., Chang, J., Luna, M.L., Long, C.E*., et al.* (2013). Release of severe acute
1046 respiratory syndrome coronavirus nuclear import block enhances host transcription in human
1047 lung cells. Journal of virology *87*, 3885-3902.

1048

1049 Siu, K.L., Yeung, M.L., Kok, K.H., Yuen, K.S., Kew, C., Lui, P.Y., Chan, C.P., Tse, H., Woo, P.C.,
1050 Yuen, K.Y*., et al.* (2014). Middle east respiratory syndrome coronavirus 4a protein is a double-
1051 stranded RNA-binding protein that suppresses PACT-induced activation of RIG-I and MDA5 in
1052 the innate antiviral response. Journal of virology *88*, 4866-4876.

1053

1054 Sola, I., Almazan, F., Zuniga, S., and Enjuanes, L. (2015). Continuous and Discontinuous RNA
1055 Synthesis in Coronaviruses. Annu Rev Virol *2*, 265-288.

1056

1057 Stark, G.R., and Darnell, J.E., Jr. (2012). The JAK-STAT pathway at twenty. Immunity *36*, 503-514.

1058

1059 Suthar, M.S., Shabman, R., Madric, K., Lambeth, C., and Heise, M.T. (2005). Identification of
1060 adult mouse neurovirulence determinants of the Sindbis virus strain AR86. Journal of virology
1061 *79*, 4219-4228.

1062

1063 Thi Nhu Thao, T., Labroussaa, F., Ebert, N., V'Kovski, P., Stalder, H., Portmann, J., Kelly, J.,
1064 Steiner, S., Holwerda, M., Kratzel, A*., et al.* (2020). Rapid reconstruction of SARS-CoV-2 using a
1065 synthetic genomics platform. Nature *582*, 561-565.

1066

1067 Thornbrough, J.M., Jha, B.K., Yount, B., Goldstein, S.A., Li, Y., Elliott, R., Sims, A.C., Baric, R.S.,
1068 Silverman, R.H., and Weiss, S.R. (2016). Middle East Respiratory Syndrome Coronavirus NS4b
1069 Protein Inhibits Host RNase L Activation. mBio *7*.

1070

1071 Turner, R.B., Weingand, K.W., Yeh, C.H., and Leedy, D.W. (1998). Association between
1072 interleukin-8 concentration in nasal secretions and severity of symptoms of experimental
1073 rhinovirus colds. Clinical infectious diseases : an official publication of the Infectious Diseases
1074 Society of America *26*, 840-846.

1075

1076 Uhlén, M., Fagerberg, L., Hallström, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A., Sivertsson,
1077 Å., Kampf, C., Sjöstedt, E., Asplund, A*., et al.* (2015). Proteomics. Tissue-based map of the
1078 human proteome. Science *347*, 1260419.

1079

1080 Volk, A., Hackbart, M., Deng, X., Cruz-Pulido, Y., O'Brien, A., and Baker, S.C. (2020). Coronavirus
1081 Endoribonuclease and Deubiquitinating Interferon Antagonists Differentially Modulate the Host
1082 Response during Replication in Macrophages. Journal of virology *94*.

1083

1084 Wan, Y., Shang, J., Graham, R., Baric, R.S., and Li, F. (2020). Receptor recognition by novel
1085 coronavirus from Wuhan: An analysis based on decade-long structural studies of SARS. Journal
1086 of virology.

1087

1088 Weiss, S.R., and Navas-Martin, S. (2005). Coronavirus pathogenesis and the emerging pathogen
1089 severe acute respiratory syndrome coronavirus. Microbiology and molecular biology reviews :
1090 MMBR *69*, 635-664.

1091

1092 Whelan, J.N., Li, Y., Silverman, R.H., and Weiss, S.R. (2019). Zika Virus Production Is Resistant to
1093 RNase L Antiviral Activity. Journal of virology *93*.

1094

1095 Xia, H., Cao, Z., Xie, X., Zhang, X., Yun-Chung Chen, J., Wang, H., Menachery, V.D., Rajsbaum, R.,
1096 and Shi, P.-Y. (2020). Evasion of type-I interferon by SARS-CoV-2. Cell Reports *10*, 8234.

1097

1098 Xie, X., Muruato, A., Lokugamage, K.G., Narayanan, K., Zhang, X., Zou, J., Liu, J., Schindewolf, C.,
1099 Bopp, N.E., Aguilar, P.V.*, et al.* (2020). An Infectious cDNA Clone of SARS-CoV-2. Cell host &
1100 microbe *27*, 841-848 e843.

1101

1102 Yamada, Y., and Liu, D.X. (2009). Proteolytic activation of the spike protein at a novel RRRR/S
1103 motif is implicated in furin-dependent entry, syncytium formation, and infectivity of
1104 coronavirus infectious bronchitis virus in cultured cells. Journal of virology *83*, 8744-8758.

1105

1106 Yang, W., Liu, Y., Slovik, K.J., Wu, J.C., Duncan, S.A., Rader, D.J., and Morrisey, E.E. (2015).
1107 Generation of iPSCs as a Pooled Culture Using Magnetic Activated Cell Sorting of Newly
1108 Reprogrammed Cells. PloS one *10*, e0134995.

1109

1110 Ye, Y., Hauns, K., Langland, J.O., Jacobs, B.L., and Hogue, B.G. (2007). Mouse hepatitis
1111 coronavirus A59 nucleocapsid protein is a type I interferon antagonist. Journal of virology *81*,
1112 2554-2563.

1113

1114 Zhao, L., Birdwell, L.D., Wu, A., Elliott, R., Rose, K.M., Phillips, J.M., Li, Y., Grinspan, J., Silverman,
1115 R.H., and Weiss, S.R. (2013). Cell-type-specific activation of the oligoadenylate synthetase-
1116 RNase L pathway by a murine coronavirus. Journal of virology *87*, 8408-8418.

1117

1118 Zhao, L., Jha, B.K., Wu, A., Elliott, R., Ziebuhr, J., Gorbalenya, A.E., Silverman, R.H., and Weiss,
1119 S.R. (2012). Antagonism of the interferon-induced OAS-RNase L pathway by murine coronavirus
1120 ns2 protein is required for virus replication and liver pathology. Cell host & microbe *11*, 607-
1121 616.

1122

1123  Zhao, L., Rose, K.M., Elliott, R., Van Rooijen, N., and Weiss, S.R. (2011). Cell-type-specific type I
1124  interferon antagonism influences organ tropism of murine coronavirus. Journal of virology *85*,
1125  10058-10068.
1126
1127  Zhou, A., Paranjape, J., Brown, T.L., Nie, H., Naik, S., Dong, B., Chang, A., Trapp, B., Fairchild, R.,
1128  Colmenares, C.*, et al.* (1997). Interferon action and apoptosis are defective in mice devoid of
1129  2',5'-oligoadenylate-dependent RNase L. The EMBO journal *16*, 6355-6363.
1130
1131  Zhou, P., Yang, X.L., Wang, X.G., Hu, B., Zhang, L., Zhang, W., Si, H.R., Zhu, Y., Li, B., Huang, C.L.*,*
1132  *et al.* (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin.
1133  Nature.
1134

**Figure Legends**

**Figure 1. Double-stranded RNA induced innate immune responses during SARS-CoV-2**

**infection.** Coronavirus double-stranded RNA (dsRNA) is produced through replication and

transcription and recognized by cytosolic OAS, MDA5, or PKR host receptors to activate innate

immune pathways. MDA5 signals through MAVS, leading to type I and type III IFN production and

release from the cell where it binds to cell surface receptors, which induces phosphorylation and

heterodimerization of STAT1 and STAT2 that then prompt ISG transcription and cytokine

responses. OASs produce 2'-5'-oligoadenylates (2-5A) that bind RNase L, leading to

homodimerization and catalytic activation of RNase L, which cleaves host and viral ssRNA to

trigger apoptosis and inflammation. PKR autophosphorylates before phosphorylating eIF2$\alpha$,

which leads to translational arrest, cell death, and inflammatory responses.  Graphic was created

with Biorender.com

1147

1148

**Figure 2. Infection of nasal epithelia-derived cells by SARS-CoV-2 and MERS-CoV.** Nasal

cells were cultured in air-liquid trans-wells, and mock infected or infected with SARS-CoV-2

(MOI=5), MERS-CoV (MOI=5) , or Sendai Virus (SeV) apically at MOI=10. (A) At indicated times,

apically released virus was collected and quantified by plaque assay on Vero-E6 cells. Values are

means ± SD (error bars). Statistical significance (not displayed) was determined by two-way

ANOVA (*, P < 0.05). One experiment was performed using four separate donors. (B) At 48 hpi,

nasal cells were fixed with 4% PFA and permeabilized. Expression of nucleocapsid (N) protein

(red) of SARS-CoV-2 and MERS-CoV was detected with an anti-N antibody, and cilia (green) with

an anti-type IV β-tubulin antibody by immunofluorescence assay (IFA). One representative image

is shown from at least three independent experiments, with four donors for each virus infection

shown. Scale bar = 100μm. (C) At 120 hpi, cells were lysed, and proteins were analyzed by

immunoblotting with antibodies as indicated. One experiment using three separate donors was

performed. (D) At 120 hpi, total RNA was harvested, and the mRNA expression level of *IFNB,*

*IFNL1, OAS2, IFIT1, IFIH1, CXCL8* was quantified by RT-qPCR. Cycle threshold ($C_T$) values were

normalized to 18S rRNA to generate $\Delta C_T$ values ($\Delta C_T = C_T$ gene of interest - $C_T$ 18S rRNA). Fold

change over mock values were calculated by subtracting mock infected $\Delta C_T$ values from virus

infected $\Delta C_T$ values, displayed as $2^{-\Delta(\Delta Ct)}$. Technical replicates were averaged, the means for each

replicate displayed, ± SD (error bars). One experiment was performed using three separate donor

samples. (E) Total RNA was harvested from two donors at 120 hpi and rRNA integrity determined

by Bioanalyzer. The position of 28S and 18S rRNA and indicated. Data shown are from one

representative experiment of two independent experiments. (See also Figures S1A&S2).

1170

**Figure 3. Infection of iPSC-derived AT2 cells (iAT2) by SARS-CoV-2.** iAT2 cells were mock

infected or infected with SARS-CoV-2 at MOI=5 or SINV at MOI=1. (A) At indicated times,

supernatants were collected and infectious virus was quantified by plaque assay on Vero-E6 cells.

Values are means ± SD (error bars). Data shown are one representative experiment from at least

three independent experiments. (B) At 48 hpi, cells were fixed with 4% PFA and permeabilized.

Expression of nucleocapsid (N) protein (green) of SARS-CoV-2 and the expression of SFTPC

promoter control tdTomato fluorescent protein (AT2 marker in red) was examined by IFA.

Channels are merged with DAPI nuclear staining. Images shown are representative from at least

three independent experiments. Scale bar = 100μm. (C) At 48 hours post infection, cells were

lysed and proteins were analyzed by immunoblotting with antibodies as indicated. Data shown are

from one representative experiment of two independent experiments. (D) At 16 (SINV) or 48 (SARS-

CoV-2) hpi, total RNA was harvested, and the mRNA expression level of *IFNB, IFNL1, OAS2,*

*IFIT1, IFIH1, CXCL8* was quantified by RT-qPCR. $C_T$ values were normalized to 18S rRNA to

generate $\Delta C_T$ values ($\Delta C_T = C_T$ gene of interest - $C_T$ 18S rRNA). Fold change over mock values

were calculated by subtracting mock infected $\Delta C_T$ values from virus infected $\Delta C_T$ values,

displayed as $2^{-\Delta(\Delta Ct)}$. Technical replicates were averaged, the means for each replicate displayed,

± SD (error bars). Statistical significance was determined by Student *t* test (*, $P < 0.05$; **, $P <$

$0.01$; ***, $P < 0.001$). Data shown are from one representative experiment of two independent

experiments. (E) Total RNA was harvested at 16 (SINV) or 48 (SARS-CoV-2) hpi and rRNA integrity

determined by Bioanalyzer. The position of 28S and 18S rRNA and indicated. Data shown are from

one representative experiment of two independent experiments. (See also Figures S1B&S2).

1192

**Figure 4. Infection of iPSC-derived cardiomyocytes (iCM) by SARS-CoV-2.** iCM were mock

infected or infected at MOI=1 with SARS-CoV-2 or SINV. (A) At indicated times, supernatants

were collected and virus quantified by plaque assay on Vero-E6 cells. Values are means ± SD

(error bars). Data shown are one representative experiment from at least three independent

experiments. (B) At 48 hpi, iCM were fixed with 4% PFA and permeabilized, the expression of

SARS-CoV-2 N (green) of and of cTnT protein (cardiomyocyte marker, red) was examined by

IFA. Channels are merged with DAPI nuclear staining. Images shown are representative from

three independent experiments. Scale bar = 50μm. (C) At 16 (SINV) or 48 (SARS-CoV-2) hpi,

cells were lysed and proteins were analyzed by immunoblotting with antibodies as indicated.

Immunoblots were performed at least two times and one representative blot is shown. (D) At 16

(SINV) or 48 (SARS-CoV-2) hpi, total RNA was harvested, the mRNA expression level of *IFNB,*

*IFNL1, OAS2, IFIT1, IFIH1, CXCL8* was quantified by RT-qPCR. $C_T$ values were normalized to

18S rRNA to generate $\Delta C_T$ values ($\Delta C_T$ = $C_T$ gene of interest - $C_T$ 18S rRNA). Fold change over

mock values were calculated by subtracting mock infected $\Delta C_T$ values from virus infected $\Delta C_T$

values, displayed as $2^{-\Delta(\Delta Ct)}$. Technical replicates were averaged, the means for each replicate

displayed, ± SD (error bars). Statistical significance was determined by Student *t* test (*, P < 0.05;

****, P < 0.0001; ns = not significant). Data shown are from one representative experiment of two

independent experiments. (E) Total RNA was harvested at 16 (SINV) or 48 (SARS-CoV-2) hpi, and

rRNA integrity determined by Bioanalyzer. The position of 28S and 18S rRNA and indicated. Data

shown are from one representative experiment of two independent experiments. (See also Figures

S1C&S2).

1214

**Figure 5. Replication of SARS-CoV-2 in A549$^{ACE2}$ and Calu-3 cell lines.** (A) Vero-E6 or

A549$^{ACE2}$ cells were infected with SARS-CoV-2 at MOI=1. At the indicated times, supernatant was

collected and virus quantified by plaque assay on Vero-E6 cells. Values are means ± SD (error

bars). (B) Calu-3 cells were infected with SARS-CoV-2, MERS-CoV or MERS-CoV-ΔNS4ab at

MOI=1. Supernatant was collected at the indicated times and virus quantified by plaque assay on

Vero-E6 cells (SARS-CoV-2) or VeroCCL81 cells (MERS-CoV and MERS-CoV-Δ4ab). Values

represent means ± SEM (error bars). Statistical significance was determined by Student *t* test (**,

P < 0.01). Data shown are one representative experiment of three independent experiments. (C)

Vero-E6, A549$^{ACE2}$, and Calu-3 cells were grown on untreated (Vero-E6 and A549$^{ACE2}$) or

collagen-coated (Calu-3) glass coverslips before infection with SARS-CoV-2 at MOI = 1. At

indicated hpi, cells were fixed with 4% PFA and permeabilized for N (green) and dsRNA (red)

expression detection by IFA using anti-N and J2 antibodies, respectively. Channels are merged

with DAPI nuclear staining. Images shown are representative from two independent experiments.

Scale bar = 25μm. (See also Figure S3&S4A).

1229

**Figure 6. SARS-CoV-2 IFN responses in the lung epithelia-derived A549$^{ACE2}$ cell line.**

A549$^{ACE2}$ cells were mock infected or infected with SINV (MOI=1) or SARS-CoV-2 (MOI=5). (A)

Total RNA was harvested at 24 and 48 hpi. Expression of *IFNB, IFNL1, OAS2, IFIT1, IFIH1,* and

*CXCL8* mRNA was quantified by RT-qPCR. C$_T$ values were normalized to 18S rRNA to generate

$\Delta$C$_T$ values ($\Delta$C$_T$ = C$_T$ gene of interest - C$_T$ 18S rRNA). Fold change over mock values were

calculated by subtracting mock infected $\Delta$C$_T$ values from virus infected $\Delta$C$_T$ values, displayed as

2$^{-\Delta(\Delta Ct)}$. Technical replicates were averaged, the means for each replicate displayed, ± SD (error

bars). (B) Viral genome copies per ug of total RNA were calculated at 24 and 48hpi by RT-qPCR

standard curve generated using a digested plasmid encoding SARS-CoV-2 nsp12. Values are

means ± SD (error bars). Statistical significance was determined by one-way ANOVA (*, P < 0.05;

**, P < 0.01; ***, P < 0.001; ****, P < 0.0001; ns = not significant). (C) At 24 hpi, A549$^{ACE2}$ cells

were lysed and proteins harvested. Protein expression was analyzed by immunoblot using the

indicated antibodies. All data are one representative experiment of three independent

experiments. (See also Figures S2 and S4B&C).

1244

**Figure 7. SARS-CoV-2 and MERS-CoV IFN responses in the lung-derived Calu-3 cells.** Calu-

3 cells were mock treated or infected with SARS-CoV-2, MERS-CoV or MERS-CoV-$\Delta$NS4ab at

MOI=5. (A) At 24 or 48 hpi, total RNA was harvested. Expression of *IFNB, IFNL1, OAS2, IFIT1,*

*IFIH1,* and *CXCL8* mRNA was quantified by RT-qPCR. C$_T$ values were normalized to 18S rRNA

to generate $\Delta$C$_T$ values ($\Delta$C$_T$ = C$_T$ gene of interest - C$_T$ 18S rRNA). Fold change over mock values

were calculated by subtracting mock infected $\Delta$C$_T$ values from virus infected $\Delta$C$_T$ values,

displayed as 2$^{-\Delta(\Delta Ct)}$. Technical replicates were averaged, the means for each replicate displayed,

± SD (error bars). Statistical significance was determined by two-way ANOVA (*, P < 0.05; **, P

< 0.01; ***, P < 0.001; ****, P < 0.0001; ns = not significant). (B) Viral genome copies per ug of

total RNA were calculated by RT-qPCR standard curve generated using a digested plasmid

encoding SARS-CoV-2 nsp12 or plasmid encoding a region of MERS-CoV orf1ab. Values are

means ± SD (error bars). Statistical significance was determined by two-way ANOVA (*, $P < 0.05$;

**, $P < 0.01$; ns = not significant). (C) At 24 hpi, Calu-3 cells were lysed and proteins harvested.

Proteins were analyzed by immunoblotting using the indicated antibodies. All data are one

representative experiment of three independent experiments. (See Figure S2).

1260

**Figure 8. SARS-CoV-2 infection leads to activation of RNase L and PKR in A549$^{ACE2}$ and**

**Calu-3 cells.** A549$^{ACE2}$ and Calu-3 cells were mock infected or infected with SARS-CoV-2, MERS-

CoV, or MERS-CoV-ΔNS4ab at MOI=5. Total RNA was harvested from A549$^{ACE2}$ cells (A) or

Calu-3 cells (B) at 24 and 48 hpi. 28S and 18S rRNA integrity was assessed by Bioanalyzer. 28S

and 18s rRNA bands are indicated. At 24 hpi, A549$^{ACE2}$ cells (C) or Calu-3 cells (D) were lysed

and proteins harvested for analysis by immunoblotting using the indicated antibodies. All data are

one representative experiment of three independent experiments. (See also Figure S4D&E).

1268

**Figure 9. Replication of SARS-CoV-2 is restricted by RNase L independent of PKR or**

**MAVS.** Indicated genes were knocked out (KO) from one clone of A549$^{ACE2}$ cells using CRISPR-

Cas9 engineering. (A) KO cell lines were grown in culture with or without 1000U IFN-α treatment

for 24 hours. Cells were lysed and proteins harvested for analysis by immunoblotting using

indicated antibodies. (B) Indicated cell lines were infected with SARS-CoV-2 at MOI=1. At the

indicated time points, supernatant was collected and virus quantified by plaque assay on Vero-

E6 cells. Values represent mean ± SD (error bars). Statistical significance was determined by

two-way ANOVA (****, $P < 0.0001$; ns = not significant). Data are one representative experiment

from at least three independent experiments. (C) Indicated cell lines were mock treated or infected

with SARS-CoV-2 at MOI=1. At 48 hpi, cells were fixed with 4% PFA and stained with 1% crystal

violet as a marker for live cells. The image is one representative experiment from two independent

experiments. (D) The indicated cell lines were mock infected or infected with SARS-CoV-2 or

SINV at MOI=1. RNA was harvested 24 hpi (SINV) or 24 and 48 hpi (SARS-CoV-2). Integrity of

rRNA was assessed by Bioanalyzer. 28S and 18S rRNA bands are indicated. Data is one

representative experiment of two independent experiments.

1284

**Figure S1. Genome replication in nasal cells, iAT2, and iCM.** Nasal (A) and iAT2 cells (B)

were infected at MOI=5 with SARS-CoV-2, and (C) iCM at MOI=1 with SARS-CoV-2 or SINV.

Total RNA was harvested at 48 hpi(SARS-COV-2) or 16 hpi(SINV)  for iAT2 and iCM cells and

120 hpi for nasal cells. Viral genome copies per ug of harvested RNA were calculated by RT-

qPCR standard curve generated using a digested plasmid encoding SARS-CoV-2 nsp12. Values

are means ± SD (error bars). For SINV (C), cycle threshold ($C_T$) values of SINV nsP4 polymerase

sequences were normalized to 18S rRNA to generate $\Delta C_T$ values ($\Delta C_T = C_T$ gene of interest - $C_T$

18S rRNA). Technical triplicates were averaged and displayed using the equation $2^{-(\Delta CT)}$. Data are

from one representative experiment of two independent experiments.

1294

**Figure S2. Host basal mRNA expression of uninfected cells.** Total RNA was harvested from

mock treatment from all indicated cell types after 24 hours incubation. mRNA expression levels

of *IFNB, IFNL1, OAS2, IFIT1, IFHI1,* and *CXCL8* were quantified by RT-qPCR. $C_T$ values were

normalized to 18S rRNA to generate $\Delta C_T$ values ($\Delta C_T = C_T$ gene of interest - $C_T$ 18S rRNA). (A)

Basal level of gene expression is displayed for nasal cells, iAT2 and iCM, Calu-3 cells and two

clones of A549$^{ACE2}$ cells, displayed as $2^{-\Delta Ct}$. (B) Fold expression over A549$^{ACE2}$ C44 values were

calculated by subtracting $\Delta C_T$ values from the indicated cell line from A549$^{ACE2}$ C44 $\Delta C_T$ values,

displayed as $2^{-\Delta(\Delta CT)}$. Biological replicates were averaged and values are means ± SD (error bars).

Data were generated from at least two independent experiments.

1304

1305 **Figure S3. ACE2 protein expression in A549^ACE2 and Calu-3 cell lines.** Parental A549 cells,

1306 two A549^ACE2 clones, and Calu-3 cells were grown in culture before lysis and protein harvest.

1307 Protein expression was analyzed by immunoblotting using the indicated antibodies.

1308

**Figure S4. SARS-CoV-2 replication and host responses in a second lung epithelia-derived**

**A549^ACE2 cell line clone (C34).** (A) Vero-E6 or A549^ACE2 cells were infected with SARS-CoV-2 at

MOI=1 and supernatant harvested at indicated times post infection. Infectious virus was quantified

by plaque assay on Vero-E6 cells. Values are means ± SD (error bars). (B) A549^ACE2 cells (C34)

were mock infected or infected with SARS-CoV-2 or SINV at MOI=5 and total RNA total RNA

harvested at 24 (SINV) or 24 and 48 (SARS-CoV-2) hpi. Expression of *IFNB, IFNL1, OAS2, IFIT1,*

*IFIH1,* and *CXCL8* mRNA was quantified by RT-qPCR. $C_T$ values were normalized to 18S rRNA

to generate $\Delta C_T$ values ($\Delta C_T = C_T$ gene of interest - $C_T$ 18S rRNA). Fold change over mock values

were calculated by subtracting mock infected $\Delta C_T$ values from virus infected $\Delta C_T$ values,

displayed as $2^{-\Delta(\Delta Ct)}$. Statistical significance for each gene was determined by one-way ANOVA

(***, $P < 0.001$; ****, $P < 0.0001$; ns = not significant). Technical replicates were averaged, the

means for each replicate displayed, ± SD (error bars). (C&D) A549^ACE2 cells were infected at

MOI=5, lysed at 24 hpi, and proteins harvested for analysis by immunoblotting using the indicated

antibodies. (E) A549^ACE2 cells were infected at MOI=1 (SINV) or MOI=5 (SARS-CoV-2) and total

RNA harvested at 24 (SINV) or 24 and 48 (SARS-CoV-2) hpi. Integrity of rRNA was assessed by

Bioanalyzer. 28S and 18s rRNA bands are indicated. All data are representative of two or three

independent experiments.

1326

**A**

SARS-CoV-2 in iAT2

Titer (log10 PFU/ml) vs hours post-infection (6, 24, 48)

**B**

SFTPC:tdTomato    SARS-CoV-2 N    Merge DAPI

Mock

SARS-CoV-2

100µm

**C**

Mock    SARS-CoV-2

pSTAT1
STAT1
pPKR
PKR
peIF2α
eIF2α
RNase L
MDA5
MAVS
ACE2
SARS-CoV-2
GAPDH

**D**

Fold change over mock

SARS-CoV-2
SINV

IFNB    IFNL1    OAS2    IFIT1    IFIH1    CXCL8

**E**

Mock    SARS-CoV-2    SINV

28S
18S

**Figure 3**

**A** SARS-CoV-2 in iCardiomyocytes

**B** cTnT | SARS-CoV-2 N | Merge DAPI

Mock

SARS-CoV-2

**C** Mock | SINV | SARS-CoV-2

pSTAT1
STAT1
pPKR
PKR
peIF2α
eIF2α
RNase L
MDA5
MAVS
ACE2
SARS-CoV-2 N
GAPDH

**D**

**E** Mock | SARS-CoV-2 | SINV

28S
18S

**Figure 4**

Figure 5

# Figure 6

# Figure 7

**Figure 8**

**Figure 9**

**Figure S1**

**A**

**B**

**Figure S2**

**Figure S3**

**Figure S4**