# Disinhibitory circuitry gates associative synaptic plasticity in olfactory cortex

Martha Canto-Bustos[1,2], F. Kathryn Friason[1,2], Constanza Bassi[1]
and Anne-Marie M. Oswald[1,2,3]*

[1]Department of Neuroscience, [2]Center for the Neural Basis of Cognition,

University of Pittsburgh, Pittsburgh, PA, United States, 15213

**Corresponding Author**:
Anne-Marie M. Oswald
Associate Professor
Department of Neuroscience
University of Pittsburgh
A210 Langley Hall
Pittsburgh, PA, 15260

[3]Current Address
Department of Neurobiology
Neuroscience Institute
University of Chicago
Abbott Memorial Hall
947 East 58th St. MC0928
Chicago, IL, USA, 60637
amoswald@uchicago.edu

**Abstract**

Inhibitory microcircuits play an essential role in regulating cortical responses to sensory stimuli.

Interneurons that inhibit dendritic or somatic integration in pyramidal neurons act as

gatekeepers for neural activity, synaptic plasticity and the formation of sensory representations.

Conversely, interneurons that specifically inhibit other interneurons can open gates through

disinhibition. In the rodent piriform cortex, relief of dendritic inhibition permits long-term

potentiation (LTP) of the recurrent synapses between pyramidal neurons (PNs) thought to

underlie ensemble odor representations. We used an optogenetic approach to identify the

inhibitory interneurons and disinhibitory circuits that regulate LTP. We focused on three

prominent inhibitory neuron classes- somatostatin (SST), parvalbumin (PV), and vasoactive

intestinal polypeptide (VIP) interneurons. We find that LTP is gated by the inactivation SST or

PV interneurons or by activation of VIP interneurons. Further, activation of VIP interneurons

strongly inhibits putative SST-cells during LTP induction, but only weakly inhibit PV-

interneurons. Taken together, these findings suggest that VIP-interneurons mediate a

disinhibitory circuit that can regulate synaptic plasticity during olfactory processing.

Throughout the cortex, the response properties of individual neurons as well as

coordinated ensemble activity are refined by sensory experience. One underlying feature of

experience-dependent plasticity is long-term changes in synaptic strength within cortical circuits.

While the mechanisms underlying excitatory synaptic plasticity have been extensively studied

(Abbott and Nelson 2000, Malenka and Bear 2004), less is known about the role cortical

circuitry plays in gating changes in synaptic strength. Inhibitory interneurons regulate both

dendritic integration and neural activity, two major factors in synaptic plasticity. Thus, inhibitory

circuits can play a key role in the enhancement of synaptic connections (Artinian and Lacaille

2018, Lucas and Clem 2018). In this study, we elucidate the inhibitory and disinhibitory circuit

motifs that gate synaptic plasticity at recurrent excitatory synapses in the olfactory cortex.

The anterior piriform cortex (APC) processes olfactory information, and performs both

sensory and associative cortical functions. Located two synapses from the periphery; the APC is

a primary cortical area representing odor inputs. Olfactory receptor neurons in the nose project

to Mitral and Tufted (M/T) neurons in the olfactory bulb (OB)(Mombaerts, Wang et al. 1996).

M/T neurons then project directly to APC and synapse with pyramidal neurons (PNs)(Haberly

and Price 1977). Unlike other primary sensory cortices, APC lacks a topological representation

of odor identity. M/T axons project diffusely and randomly (Sosulski, Bloom et al. 2011, Igarashi,

Ieki et al. 2012) to activate distributed neural ensembles (Illig and Haberly 2003, Rennaker,

Chen et al. 2007, Stettler and Axel 2009). Distributed odor representations are further supported

by uniform intracortical excitatory connectivity across the APC (Franks, Russo et al. 2011). It is

postulated that odor-specific ensembles are constructed by strengthening excitatory synapses

between pyramidal neurons co-activated by odor components (Haberly 2001, Wilson and

Sullivan 2011). In support of this hypothesis, intracortical synapses between PNs are

strengthened following odor learning *in vivo* (Saar, Grossman et al. 2002) and through pairing of

afferent and intracortical stimulation *in vitro* (Kanter and Haberly 1993, Johenning, Beed et al.

2009). Hence, APC circuitry also supports early stage, associative odor processing.

The APC is an ideal structure for investigating the circuit and synaptic plasticity

mechanisms that underlie sensory representations. PNs receive compartmentalized excitatory

inputs on their apical dendrites from two distinct fiber tracts. M/T cell afferents synapse distally,

while PN axons form a proximal intracortical fiber tract (Haberly and Price 1977, Haberly and

Price 1978). Co-activation of afferent and intracortical fiber tracts strengthens intracortical

synapses onto PNs through NMDA receptor (R) dependent, associative LTP (Kanter and

Haberly 1993).  Further, NMDAR EPSPs and associative LTP induction are facilitated by

dendritic disinhibition by $GABA_A$ receptor antagonists (Kanter and Haberly 1993, Kanter, Kapur

et al. 1996). While these studies suggest interplay between dendritic inhibition and disinhibition

gates LTP induction at intracortical synapses, the inhibitory circuitry involved has not been

identified.

Olfactory stimuli recruit both feedforward and recurrent inhibition onto PNs in APC (Poo

and Isaacson 2009, Poo and Isaacson 2011). Feedforward inhibition is comparatively weak and

diminishes with high frequency stimulation of the afferent pathway, whereas recurrent inhibition

is strong, and increases with stimulation through synaptic facilitation and PN recruitment

(Stokes and Isaacson 2010, Suzuki and Bekkers 2010a, Large, Vogler et al. 2016). Recurrent

inhibition is mediated by inhibitory interneurons that express somatostatin (SST-INs) or

parvalbumin (PV-INs) (Stokes and Isaacson 2010, Suzuki and Bekkers 2010a, Suzuki and

Bekkers 2010b, Large, Kunz et al. 2016). SST-INs inhibit PN apical dendrites proximal to the

soma, and are optimally located to regulate plasticity of intracortical synapses (Suzuki and

Bekkers 2010b, Large, Kunz et al. 2016). PV-INs regulate spike activity and could also impact

LTP through backpropagation (Johenning, Beed et al. 2009). A third class of interneurons-

vasoactive intestinal polypeptide interneurons (VIP-INs); inhibit SST and PV-INs and could

disinhibit pyramidal neurons (Lee, Kruglikov et al. 2013, Pfeffer, Xue et al. 2013, Karnani,

Jackson et al. 2016). VIP-INs are a prominent in APC (Suzuki and Bekkers 2010b) but their

inhibitory connections and function are unknown. We investigated the connectivity and

functional roles of SST, PV and VIP-INs in APC. We find that activation of VIP-INs as well as

inactivation of SST-INs or PV-INs promote LTP of intracortical synapses. VIP-INs strongly inhibit

SST-INs during LTP induction, but only weakly inhibit PV-INs and PNs. Our finding suggests a

VIP->SST>-PN disinhibitory circuit gates associative LTP in APC.

**Methods:**

*Mice:* VIP-Cre *(*B6:Viptm1(cre)Zjh/J) , SST-Cre (B6:Sst<tm2.1(cre)Zjh>/J)  and PV-Cre mice

express cre-recombinase (Taniguchi, He et al. 2011). These mice were crossed with Ai32 mice

(B6;129S-Gt ROSA)26Sortm32 (CAG-COP4*H134R/EYFP)Hze/J) to express channelrhodopsin

(ChR2) or Ai35 mice (B6.129S-*Gt(ROSA)*26Sortm35.1(*CAG- aop3/GFP)Hze*/J) to express

archaerhodopsin (Arch) (Madisen, Mao et al. 2012).  All mice are from Jackson Laboratory. All

animals were bred, handled and treated in manner that was evaluated and approved by the

Animal Care and Use Committee at the University of Pittsburgh IACUC, protocol #17070877.

*Slice preparation:*  APC brain slices were prepared from mice aged P19-35.  The mice were

anesthetized with isoflurane and the brain was removed and immersed in ice cold oxygenated

(95% $O_2$-5% $CO_2$) ACSF (in mM: 125 NaCl, 2.5 KCl, 25 $NaHCO_3$, 1.25 $NaH_2PO_4$, 1.0 $MgCl_2$, 25

Glucose, 2.5 $CaCl_2$) (all chemicals from Sigma, USA unless otherwise stated). Parasagittal

slices (300 $\mu$m) were cut on a vibratome (Leica Biosystems) in ice-cold ACSF. The slices were

transferred to warm ACSF (36$°$C) for 30 min, then 20-22$°$C for 1 hour, and recorded at 25-28$°$C.

*Electrophysiology*:  Recordings were performed using a MultiClamp 700B amplifier (Molecular

Devices, Union City, CA). Data were low pass filtered (4 kHz) and digitized at 10 kHz using an

ITC-18 (Instrutech) controlled by custom software (Recording Artist,

https://bitbucket.org/rgerkin/recording-artist) written in IgorPro (Wavemetrics).  Recording

pipettes (4-10 M$\Omega$) were pulled from borosilicate glass (1.5 mm, outer diameter) on a

Flaming/Brown micropipette puller (Sutter Instruments). The series resistance (<20 M$\Omega$) was not

corrected. For PSPs the intracellular solution consisted of (in mM) 130 K-gluconate, 5 KCl, 2

$MgCl_2$, 4 ATP-Mg, 0.3 GTP, 10 HEPES, and 10 phosphocreatine, 0.05% biocytin. For IPSC

recordings, Qx-314 was added to the K-gluconate internal (holding potential 0 mV) or Cs-Glu-

Qx solution was used (in mM, 130 Cs-Gluconate, 5 KCl, 2 $MgCl_2$, 4 Mg-ATP, 0.3 GTP, 10

HEPES, 10 Phosphocreatine, 1 Qx-314, holding potential +30 mV). Neurons were visualized

using infrared-differential interference contrast microscopy (IR-DIC, Olympus).  For all recorded

neurons, subthreshold response properties were obtained using a series of hyperpolarizing and

depolarizing current steps (-50 pA to 50 pA, 1 s duration). We used specific criteria for

identification of putative PV and SST interneurons based on intrinsic properties of identified PV

and SST neurons in APC (see (Large, Kunz et al. 2016, for details). Neural identity was

confirmed post hoc using intrinsic properties and biocytin fills.

_LTP induction:_ Electrical stimulation was delivered using concentric bipolar electrodes (FHC).

The electrodes were placed in the LOT (L1a) and the L1b/L2 border. Stimuli (100 μs pulse

width) were delivered through a stimulus isolation unit. Theta burst stimulation (TBS) of L1a

consisted of 10 bursts of 4 pulses (100 Hz) delivered at 250 ms intervals (Kanter and Haberly

1993). TBS stimulation intensity was set near spike threshold for the recorded neuron. L1b was

stimulated with a single weak pulse delivered between the 3[rd] and 4[th] pulse of each burst.

Stimulation intensity was <30% of the maximum subthreshold EPSP (~1-6 mV).  Pre and post

induction test pulses were delivered to L1b every 30 s. Baseline was collected for ~5 min and

LTP was induced within 10 min of patching the neuron. Input resistance was monitored

throughout and neurons with deviations greater than 20% from baseline were excluded from

analysis (n=10 neurons). We also monitored membrane potential throughout recording. Small

fluctuations <5 mV were offset with constant current to maintain driving force. However, neurons

with drift greater than 5 mV or exhibiting unstable membrane potential fluctuations were

excluded from analysis (n=8).

*Pharmacology*:  The GABA$_A$ receptor antagonist, gabazine (GZ, 20 μM in ACSF) was loaded

into a regular patch pipette and focally applied to L1b using a gentle positive pressure (<1-5 s

duration) see. Slices were oriented such that bath flowed from the soma to dendrite of the

pyramidal cell to maintain GZ in the region of the dendrite. Every effort was made to ensure

stability of GZ application including supplemental applications. In a subset of slices (n=5) from

mice that expressed ChR2 in SST-INs, GZ application to L1b diminished optically evoked SST-

mediated inhibition onto pyramidal cells (n=5).  We verified that optically evoked IPSPs (Control:

6.3 ± 0.69 mV) in these slices were stably diminished by Gabazine application throughout the

LTP protocol and did not differ between baseline (IPSP GZ: 0.89 ± 0.27 mV) and post-induction

test pulses (IPSP GZ: 0.92 ± 0.28, p>0.05 WSR). In additional experiments, the NMDA receptor

antagonist, DL-2-Amino-5-phosphonopentanoic acid (DL-APV, 10 mM, Sigma-Aldrich) was

bath-applied as indicated in the main text.

*Optogenetic Stimulation:*  Shutter controlled full field stimulation with blue (473 nm) or green

(520 nm) light (Prior) was delivered through the epifluorescence pathway of the microscope

(Olympus) using a water-immersion objective (40x). Light intensity (5-10 mW) was adjusted to

induce spike responses (ChR2 activation) or spike suppression (Arch inactivation). Light pulse

duration varied by experiment as indicated in the main text.

EPSP Analysis:  EPSPs were analyzed using custom software written in IgorPro (Wavemetrics).

Because we do not block inhibition during L1b test-pulses, we set narrow criteria to ensure that

only the excitatory portion of mixed EPSP-IPSPs was analyzed. Recurrent inhibition is

disynaptic and IPSPs are delayed with respect to excitation. Only EPSP onsets within 5 ms of

the stimulus artifact were analyzed. Across all conditions the average EPSP onset was 3.7 ±

0.41 ms.  EPSP peak amplitude was taken as the max amplitude within 5 ms of EPSP onset.

We also measured the slope of the rising phase of the EPSP from onset to 80% of peak

amplitude to further isolate the EPSP. Our core findings did not differ between peak and slope

measurements.

<u>Statistics:</u> All data is presented as mean ± SE. Statistical tests were performed using two tailed,

one or two-sample, paired or unpaired Student's t-test as appropriate. In cases of small sample

sizes non-parametric tests were used, including the Mann-Whitney U-test (MWU) for unpaired

data and the Wilcoxon Signed Ranks test (WSR) for paired data. For multiple comparisons we

used ANOVA with post hoc Tukey Test (ANOVA).

**Results:**

We investigated the roles of SST, PV and VIP-INs, in gating associative LTP at intracortical

excitatory synapses onto PNs. Afferent input to PNs arrives via L1a on distal L2 PN dendrites,

while the intracortical fiber tract (L1b) is proximal (**Fig 1A**). L1a and L1b were easily identified

under IR-DIC and independently stimulated using bipolar electrodes. Associative LTP is induced

by pairing L1a and L1b stimulation using a theta burst stimulation (TBS) protocol (Kanter and

Haberly 1993) consistent with respiration coupled M/T spike frequencies (Kepecs, Uchida et al.

2007, Carey and Wachowiak 2011). Briefly, strong TBS of L1a was paired with weak, single

pulse stimulation of L1b (see methods, **Fig 1A**). This L1a+L1b pairing is hereafter denoted

*induction*. L1a-TBS evoked low PN firing rates ranging from 0-8 Hz and L1b EPSPs ranged from

1-6 mV. Pre and post induction, L1b stimulation was delivered every 30 s. To avoid drift in

recording integrity, potentiation was quantified as the average L1b EPSP amplitude 25-30 min

following induction versus average baseline EPSP amplitude (5 min prior to induction).

However, in neurons where stability could be verified, potentiation typically lasted for the

duration of recording (45-60 min). Both raw and normalized (to baseline) EPSP amplitude and

rising slope were analyzed. PNs were excluded if input resistance or membrane potential varied

from baseline (±20% or ±5 mV respectively).

***Disinhibition of PN dendrites promotes LTP***

With inhibition intact, the induction protocol did not induce LTP of L1b synapses.  Neither

EPSP amplitude (in mV, pre: 2.6 ± 0.60, post: 2.6 ± 0.51, p: 0.85, paired t-test, n=9) nor slope

(in mV/ms, pre: 0.64 ± 0.20, post: 0.61 ± 0.18, p: 0.27, paired t-test, n=9) significantly differed

from baseline (pre) at 30 min post induction (**Fig 1B**).  Likewise, normalized EPSP amplitude

and area did not differ significantly from 1 (Amplitude: 1.0 ± 0.071, p: 0.65; Slope: 0.96 ± 0.079,

p: 0.65, one sample t-test, **Fig 1C, D**). However, when we focally applied the $GABA_A$ receptor

antagonist, Gabazine (GZ, 20 µM), to L1b (schematic, **Fig 1A**), induction significantly enhanced

EPSP amplitude (mV, Pre: 1.3 ± 0.32, Post: 3.1 ± 0.95, p<0.050 WSR, n=6, **Fig 1B**) and slope

(mV/ms, Pre: 0.28 ± 0.070, Post: 0.52 ± 0.12, p<0.050 WSR, n=6). Normalized EPSP amplitude

and slope were significantly greater than 1 (Amplitude: 2.3 ± 0.35, p<0.05 WSR; Slope: 2.0 ±

0.27, p<0.05 WSR, **Fig 1C, D**). Consistent application of dendritic Gabazine for the duration of

recording is difficult to maintain. In a some PNs, EPSP amplitudes were potentiated but

unstable, these were excluded from analysis (n=5). This highlights the need for a more selective

and reliable optogenetic approach to investigating disinhibition.

***Inactivation of SST-INs promotes LTP***

We have previously shown that SST-interneurons strongly inhibit PN dendrites in L1B

(Large, Kunz et al. 2016).  Although SST-INs typically receive weak direct L1a input (Suzuki and

Bekkers 2010a), SST-INs are recruited by strong TBS of L1a through the recruitment of

recurrent excitation (**Fig 2A1,2).** To investigate the influence of SST-INs on LTP induction,

Archaerhodopsin (Arch) was selectively expressed in SST-INs for optical inactivation. During

TBS, optical inactivation reduced SST-IN firing rate (FR) from 6.8 ± 1.2 Hz to 1.8 ± 0.4 Hz

(p<0.05, WSR test, **Fig 2A2**) and enhanced depolarization in PNs (**Fig 2B**). These findings

suggest that SST-interneurons mediate inhibition during TBS stimulation and could prevent

associative synaptic plasticity at L1B synapses. To test this, we optically inactivated SST-INs

solely during LTP induction (2 sec). EPSP amplitude and slope was significantly enhanced 30

min post induction (Amplitude (mV), Pre: 2.6 ± 0.41, Post: 4.3 ± 0.76, p: 0.0071; Slope (mV/ms),

Pre: 0.55 ± 0.14, Post: 1.1 ± 0.23, p: 0.0048; n=9, paired t-test, **Fig 2C1,2, E**). Normalized

EPSP amplitude and slope were significantly greater than 1 (Amplitude: 1.7 ± 0.16, p: 0.00010;

Slope: 2.0 ± 0.24, p: 0.0033, one sample t-test, **Fig 2D**). EPSP traces from representative PNs

pre and post induction are shown in **Fig 2E**. The normalized amplitude and slope averaged

across neurons over 30 min are shown in **Fig 2F1,2**.  To confirm that associative synaptic

plasticity was NMDA-receptor dependent, we bath applied the NMDAR antagonist, DL-APV (10

mM) to a second set of slices. Antagonism of NMDAR prevented LTP induction despite

inactivation of SST-INs. EPSP amplitude and slope 30 min post induction did not significantly

differ from baseline (Amplitude (mV): Pre: 4.7 ± 0.81, Post: 4.0 ± 0.73, p: 0.21; Slope (mV/ms),

Pre: 0.83 ± 0.19, Post: 0.74 ± 0.18, p: 0.26; n=10, paired t-test, **Fig 2C, black circles)**;

(normalized Amplitude: 0.88 ± 0.13, p: 0.38; Slope: 0.85 ± 0.090, p: 0.28, one sample t-test. **Fig**

**2D,F black circles**). These findings suggest that inhibition by SST-INs regulates NMDA-

dependent associative LTP at L1b intracortical synapses.

***Inactivation of PV-INs promotes LTP***

PV-INs strongly inhibit cortical neurons. We investigated whether optogenetic

inactivation of PV-INs expressing Arch also promotes associative LTP (**Fig 3A1**). PV-INs are

robustly activated during TBS **(Fig 3A2)** and light inactivation significantly decreased FR (Light

OFF: 7.2 ± 2.5 Hz, ON: 2.8 ± 1.7 Hz, n=5, p<0.05, WSR, **Fig 3A2, right)**. In a number of PNs,

this enhanced EPSP summation **(Fig 3B1**) and/or FR consistent with somatic disinhibition. PV-

IN inactivation during induction promoted LTP of EPSP amplitude and slope **(Fig 3C1,C2,**

Amplitude (mV), Pre: 1.6 ± 0.34, Post: 2.4 ± 0.40, p: 0.00095; Slope (mV/ms): 0.38 ± 0.10, Post:

0.66 ± 0.18, p: 0.016; n=9, paired t-test). Normalized amplitude and slope were significantly >1

post induction (Amplitude: 1.6 ± 0.16, p: 0.0063; Slope: 1.8 ± 0.15, p: 0.00061, one sample t-test

**Fig 3C3, D**). Antagonism of NMDAR prevented LTP induction mediated by inactivation of PV-

INs. EPSP amplitude and slope post induction did not significantly differ from baseline

(Amplitude (mV): Pre: 3.3 ± 0.45, Post: 3.0 ± 0.37, p: 0.48; Slope (mV/ms), Pre: 0.44 ± 0.082,

Post: 0.52 ± 0.10, p: 0.74; n=8, paired t-test, **Fig 3C, black circles)**; (normalized Amplitude:

0.92 ± 0.076, p: 0.65; Slope: 1.0 ± 0.13, p: 0.56, one sample t-test. **Fig 3C,D black circles**).

Thus, PV-INs can also regulate the induction of NMDA-dependent associative LTP at L1b

intracortical synapses.

### *Inhibition of interneurons by VIP-INs*

If both SST-IN and PV-INs can inhibit LTP induction, the question remains as to which

circuits might provide a means for disinhibition and the promotion of LTP.  In other cortices, VIP

interneurons inhibit SST-INs and PV-INs (Pfeffer, Xue et al. 2013, Pi, Hangya et al. 2013).

Although there are numerous VIP-INs in piriform cortex (Suzuki and Bekkers 2010b), their

targets have not been identified.  We crossed VIP-cre mice with Ai32 mice to express

channelrhodopsin (ChR2) in VIP-INs. We then used light activation to drive single action

potentials in VIP-INs while recording potential postsynaptic targets. Typically, VIP-cre mice are

crossed with mice that express GFP in SST-INs (GIN-mice) or PV-INs (G42-mice). However,

these lines sparsely label SST and PV-INs in APC (Large, Kunz et al. 2016). Instead, we used

intrinsic properties to identify putative (p) SST-INs and pPV-INs according criteria determined

from recordings in genetically identified SST-INs and PV-INs in APC (Large, Kunz et al.

2016)(see methods). Interneurons that could not be confidently identified were excluded from

analysis.

VIP-INs were activated by brief blue light pulses (5 ms) and IPSCs were recorded in

voltage clamp with either Cs-Gluconate-Qx internal (IPSCs, +30 mV) or K-Gluconate-Qx internal

(IPSCs, 0 mV). IPSC recordings were most reliable with Cs-Glu-Qx and the data presented (**Fig**

**4)** are from this condition**.**  VIP-INs inhibited nearly all recorded pSST-INs (86%), most pPV-INs

(90%) and PNs (88%). Although high connectivity with SST and PV cells was expected, this

was unexpected for PNs (Pfeffer, Xue et al. 2013). VIP-INs strongly inhibited pSST-INs (IPSC

amplitude: 303 ± 57 pA, n=19).  However, despite a high probability of connection, VIP-INs very

weakly inhibited pPV-INs (35 ± 5.8 pA, n=9, p: 0.0008) and PNs (61 ± 14 pA, p: 0.001 n=14,

ANOVA, **Fig 4A1**). Qualitatively similar results were found with K-Glu-Qx, however IPSC

amplitude and connection probability were considerably lower (% connected, amplitude, #

connected: pSST: 73%, 47 ± 14 pA, n=19; pPV: 73%, 31 ± 7.6 pA, n=11; PN: 29%, 21 ± 7.0 pA,

n=4).

We find that VIP-INs typically show strongly adapting spike responses within the first 500

ms of suprathreshold depolarizations by current steps. For this reason, long light pulses (2 s)

during LTP induction would likely result in significant adaptation of VIP-INs spike responses.

Instead, we drove VIP-INs with shorter light pulses (100 ms) delivered at theta frequency (10

pulses, **Fig 4A2**). This pulsed-light stimulation evoked strong and relatively sustained IPSCs in

pSST-INs (**Fig 4B2 top**). Overall, IPSC strength decreased by only ~30% by the $5^{th}$ pulse in all

cell types then stabilized (SST: 31 ± 10%, p: 0.002; PV: 31± 5%, p: 0.06; PC: 28 ± 12%, p: 0.02,

paired t-test, **Fig 4B2**).

To investigate how VIP-IN mediated inhibition could influence action potential activity we

depolarized target neurons to near or suprathreshold membrane potentials in current clamp. In

pSST-INs activation of VIP-INs evoked strong IPSPs (2.02 ± 0.38 mV, n=18, **Fig 4C1**). These

IPSPs were sufficient to delete or delay spikes by an average of 77 ± 26 ms (p: 0.016, n=11,

paired t-test) compared to non-light trials (**Fig 4C1,3 left**). Finally, pSST-IN spike responses

evoked by TBS were significantly decreased by pulsed activation of VIP-INs compared to non-

light trials (OFF: 5.0 ± 0.6 Hz, ON: 3.5 ± 0.5, p: 0.002, n=11, paired t-test, **Fig 4C**). We could not

record IPSPs in PNs (n=7) or pPV-INs (n=4) likely due to a combination of low input resistance

and weak VIP-mediated IPSCs.  Taken together, these findings suggest that VIP-INs could

strongly influence pSST-INs during LTP induction but would have considerably less impact on

pPV-IN or PNs.

*Activation of VIP-INs promotes LTP*

VIP-INs are numerous in APC and we find that VIP-INs inhibit pSST-INs. VIP-to-SST

inhibition is a candidate circuit motif for dendritic disinhibition of PNs that could promote LTP.

So, why isn't LTP induced without antagonism of inhibition? We recorded VIP-INs during TBS

stimulation of L1a. We found that in contrast to SST-INs or PV-INs, VIP-INs were only weakly

driven to fire action potentials by TBS ($0.7 \pm 0.6$ Hz, n=6, Fig 5A2). This lack of VIP-IN

recruitment is consistent with the inability to induce LTP under control conditions (**Fig 1B**).

However, light activation of ChR2+ VIP-INs enhanced FR during TBS ($14 \pm 4.0$ Hz, $p<0.05$,

WSR test **Fig 5A2**). Activating VIP-INs also enhanced EPSP summation in PNs during TBS

**(Fig 5B)**. Activation of VIP-INs during induction lead to robust LTP of EPSP amplitude and slope

(**Fig 5C1,C2**, amplitude (mV), Pre: $2.1 \pm 0.50$, Post: $3.4 \pm 0.48$, $p<0.02$; slope (mV/ms) Pre: 0.47

$\pm 0.081$, Post: $0.76 \pm 0.072$, $p<0.02$, n=7, WSR). Normalized amplitude and slope were

significantly >1 post induction (Amplitude: $2.0 \pm 0.33$, $p<0.02$; Slope: $1.9 \pm 0.32$, $p<0.02$, WSR,

**Fig 5C3**). Antagonism of NMDARs prevented LTP induction mediated by activation of VIP-INs.

EPSP amplitude and slope post induction did not significantly differ from baseline (Amplitude

(mV): Pre: $6.0 \pm 1.6$, Post: $5.5 \pm 1.5$, $p>0.05$; Slope (mV/ms), Pre: $1.2 \pm 0.23$, Post: $1.1 \pm 0.28$,

p: $>0.05$; n=5, WSR, **Fig 5C,D black circles)**; (normalized Amplitude: $0.93 \pm 0.072$, $p>0.05$;

Slope: $0.99 \pm 0.17$, $p>0.05$, WSR. **Fig 5C,D black circles**). Altogether, these findings support a

VIP-SST-PN disinhibitory circuit that gates associative LTP at intracortical synapses.

**Discussion**

It has been hypothesized that odor ensembles are formed through associative plasticity of

recurrent excitatory synapses between co-activated pyramidal neurons (Haberly 2001, Wilson

and Sullivan 2011). These intracortical synapses are stronger in animals that have learned an

olfactory discrimination task (Saar, Grossman et al. 2002, Saar, Reuveni et al. 2012) and the

induction of associative LTP *in vitro* is occluded in animals that have learned tasks (Lebel,

Grossman et al. 2001). This capacity for enhancement as well as highly recurrent excitation

(Poo and Isaacson 2011) necessitates strong inhibition to regulate neural activity (Luna and

Schoppa 2008, Poo and Isaacson 2011, Bolding and Franks 2018) and synaptic plasticity

(Kanter and Haberly 1993, Kanter, Kapur et al. 1996). Our present findings are consistent with a

VIP->SST->PN circuit that transiently disinhibits PN dendrites to promote synaptic plasticity

during odor learning while overall inhibition remains intact.

Associative LTP at intracortical synapses within APC is well-characterized (Stripling,

Patneau et al. 1988, Kanter and Haberly 1993, Poo and Isaacson 2007). TBS provides strong

afferent excitation that depolarizes apical PN dendrites and promotes NMDAR dependent

potentiation of co-activated intracortical synapses. However, strong TBS also recruits strong

inhibition and LTP is rarely induced in the absence of $GABA_AR$ antagonists (Kanter and Haberly

1993). We have reproduced previous findings that LTP is gated by dendritic application of

Gabazine (Kanter, Kapur et al. 1996, Kumar, Schiff et al. 2018). Further, we have shown that

dendritic application of Gabazine blocks inhibition mediated by SST-INs (Large et al., 2016).  In

the present study, we demonstrate that inactivation of SST-INs at an opportune moment during

TBS, is sufficient to promote LTP. Thus, a circuit mechanism that inhibits SST-INs during LTP

induction could gate synaptic plasticity. Throughout the cortex, VIP-INs inhibit SST-INs and

potentially disinhibit PNs (Lee, Kruglikov et al. 2013, Pfeffer, Xue et al. 2013, Karnani, Jackson

et al. 2016). We show that VIP-INs strongly inhibit pSST-INs and weakly inhibit pPV-INs as well

as PNs. Moreover, optogenetic activation of VIP-INs decreases pSST-IN spike responses

during TBS.  Finally, activation of VIP-INs specifically during TBS is sufficient gate associative

LTP. Based on these findings, we propose that a VIP->SST->PN disinhibitory circuit gates

intracortical synaptic plasticity in APC.

However, it is relevant to discuss three additional circuits that are potentially recruited

with VIP-IN activation.  For example, VIP-INs could also gate plasticity through a VIP-PV-PN

disinhibitory circuit. Although optogenetic inactivation of PV-INs promotes LTP, we predict this

circuit motif contributes minimally to associative LTP induction.  VIP-INs very weakly inhibit

pPV-INs, which is likely insufficient to prevent PV-IN spiking during TBS.  Conversely, SST-INs

strongly inhibit PV-INs (Pfeffer, Xue et al. 2013, Xu, Jeong et al. 2013, Large, Kunz et al. 2016)

and inactivation of SST-INs in APC increases pPV-IN activity (Sturgill and Isaacson, 2015).

Instead of disinhibiting the PN, a VIP-SST-PV circuit could transiently increase somatic PV-IN

inhibition to compensate for a loss of SST mediated inhibition at the dendrite and/or soma.

Finally, VIP-INs also inhibit PNs directly, possibly through basket like synapses on the soma

(Suzuki and Bekkers 2010a). As with VIP-PV synapses, this VIP-PN inhibitory circuit is also

weak and its role is unknown. Importantly, recruitment of these additional circuits does not

impair LTP induction with VIP-IN activation. Rather, the interplay of these circuits suggests a

shift in inhibition from the dendrite to soma that may stabilize neural responses while opening a

window for plasticity.

Our study complements a recent study in somatosensory cortex that used

chemogenetics to inhibit VIP-INs or SST-INs and prevent or promote LTP induction respectively

(Williams and Holtmaat 2019). Together, these two studies support a common VIP->SST->PN

disinhibitory circuit motif regulates synaptic plasticity across cortical areas. A benefit of the

present study is that by using optogenetics, we were ability to isolate the influence of

disinhibition to a brief time window (~2 s) when afferent and intracortical pathways are co-

activated. This suggests that transient dendritic disinhibition is sufficient to gate the cascade of

intracellular mechanisms underlying synaptic enhancement. Thus, circuit mechanisms that

provide a means for VIP cells to be activated in conjunction with odor sampling could play a key

role in odor learning and ensemble formation.

Interestingly, VIP-INs are not recruited by pairing afferent and intracortical stimulation,

this may underlie the inability induce associative LTP with inhibition intact (**Fig 1**). At present, it

is not known how VIP-INs in APC are recruited. In other cortices, VIP-IN activity is enhanced by

arousal, locomotion, task engagement or reward (Lee, Kruglikov et al. 2013, Pi, Hangya et al.

2013, Fu, Tucciarone et al. 2014, Jackson, Ayzenshtat et al. 2016), either through additional

excitatory drive from other cortical areas (Lee, Kruglikov et al. 2013, Williams and Holtmaat

2019) or through neuromodulation (Porter, Cauli et al. 1999, Lee, Hjerling-Leffler et al. 2010,

Alitto and Dan 2012, Kuchibhotla, Gill et al. 2017, Pronneke, Witte et al. 2019).  Direct excitatory

inputs to VIP-INs would have to be well-timed with respect to sensory stimulation.  Conversely,

neuromodulation could raise the excitability of VIP-INs such that afferent and intracortical drive

are sufficient to drive VIP-INs with appropriate timing. The APC receives input from higher

cortices including orbitofrontal cortex (Illig 2005) as well as neuromodulatory centers

(Zaborszky, Carlsen et al. 1986, Linster, Wyble et al. 1999). Both of these pathways have been

implicated in olfactory learning and plasticity (Patil, Linster et al. 1998, Patil and Hasselmo 1999,

Linster, Maloney et al. 2003, Li, Luxenberg et al. 2006, Chapuis and Wilson 2013, Cohen,

Wilson et al. 2015, Strauch and Manahan-Vaughan 2018). Future studies are needed to

ascertain the potential links between descending excitation and/or neuromodulation, the

recruitment of VIP-INs, and the gating of synaptic plasticity during olfactory processing.

Though many previous studies have focused on the influence of disinhibitory circuits on

FR or behavior, our study highlights a role for disinhibition in the dynamics that gate long-term

plasticity. Specifically, our findings suggest that the VIP-SST-PN circuit motif plays a central role

in gating synaptic plasticity in PN dendrites. Additional circuit mechanisms mediated through

VIP-INs that regulate PV-INs or PNs directly could serve a stabilizing role. These findings

demonstrate the challenge of the delineating roles for the individual circuit motifs that are nested

in complex neural networks. It remains to be determined if these mechanisms work in concert

during afferent and recurrent olfactory processing.

**Figure 1: Associative LTP of intracortical synapses is gated by dendritic disinhibition.**

**A)** Schematic of APC circuit and stimulation paradigm. Strong (s) TBS of afferents (L1a) is paired with weak (w) single-pulses at the intracortical pathway (L1b). Gabazine (GZ) is focally applied to the dendrite region in L1b (tan oval). **B1)** Representative average traces pre (gray) and post induction in control (black) and with GZ(Tan). **B2)** Average EPSP amplitude (mV) pre (-5 to 0 min) and post induction (25-30 min) with inhibition intact (black) or with dendritic GZ (tan). **B3)** Average slope (mV/ms) of the 20-80% rising phase of the EPSP pre and post induction. **C1)** Time course of normalized EPSP amplitude pre and post induction (grey box, t=0) in control (black) and with dendritic Gabazine (tan). **C2)** Time course of normalized EPSP slope, colors as in C2. **C3)** Normalized EPSP amplitude, and slope 30 min post induction in control (black) and with GZ (tan).

**Figure 2: SST-interneuron inactivation promotes associative LTP. A1)** Schematic: SST-INs express Archaerhodopsin (Arch). **A2)** SST-IN spike responses during TBS in control (black) and inactivated (green) conditions (*p<0.05, WSR test). **B)** PN responses for a single TBS burst. **Left :** Inactivation of SST-INs enhanced PN depolarization (green vs. black trace). **Right:** Depolarization during SST-IN inactivation (green) is reduced by APV (gray trace). **C1)** Average EPSP amplitude (mV) pre (-5 to 0 min) and post induction (25-30 min) with SST-inactivation (Magenta) or with SST-inactivation plus bath application of the NMDAR antagonist, APV (black). **C2)** Average slope (mV/ms) of the 20-80% rising phase of the EPSP pre and post induction, colors as in B2. **C3)** Normalized EPSP amplitude and slope following pairing with SST-IN inactivation (magenta circles). LTP is blocked by bath application APV (black). **D1)** Representative average traces pre (gray) and post induction in with SST-IN inactivation (magenta) or inactivation plus APV (Black). **D2)** Time course of normalized EPSP amplitude pre and post induction with inactivation of SST-INs (green box, t=0) in control (magenta) and with bath application of APV (black). **D3)** Time course of normalized EPSP slope, colors as in D2.

**Figure 3: PV-interneuron inactivation during induction promotes associative LTP. A1)**

Schematic: PV-INs express Archaerhodopsin (Arch). **A2)** PV-IN spike responses during TBS in

control (black) and inactivated (green) conditions. **B)** Representative PN responses for a single

TBS for control (black) versus PV-IN inactivation (green). **C1)** Average EPSP amplitude (mV)

pre (-5 to 0 min) and post induction (25-30 min) with PV-inactivation (Blue) or with PV-

inactivation plus bath application of the NMDAR antagonist, APV (black). **C2)** Average slope

(mV/ms) of the 20-80% rising phase of the EPSP pre and post induction, colors as in B2. **C3)**

Normalized EPSP amplitude and slope following pairing with PV-IN inactivation (blue circles) or

with inactivation plus bath application APV (black). **D1)** Representative average traces pre

(gray) and post induction in with PV-IN inactivation (blue) or inactivation plus APV (Black). **D2)**

Time course of normalized EPSP amplitude pre and post induction with inactivation of PV-INs

(green box, t=0) in control (blue) and with bath application of APV (black). **D3)** Time course of

normalized EPSP slope, colors as in D2.

**Figure 4: Inhibition by VIP-interneurons in piriform cortex. A1)** VIP-INs express ChR2.

Optically evoked IPSCs were recorded in putative(p) pSST-INs (magenta), pPV-INs (blue) and

PNs (black). **A2)** IPSCs recorded in response to ten light pulses (100 ms duration, 4 Hz, same

neurons as in A1). **B1)** IPSC amplitudes were stronger in pSST-INs versus pPV-INs (*p:0.014)

or PNs (*p:0.042, ANOVA). **B2**) In pSST-INs, IPSC amplitude diminishes by the $5^{th}$ pulse of

theta stimulation (**p: 0.009, paired t-test). **C1)** IPSCs from VIP-INs delay pSST-IN spike

responses during suprathreshold depolarization (4 overlaid traces) **Left:** Control, **Right**:

activation of VIP-INs (100 ms pulse, blue). Magenta trace: VIP-IN mediated IPSC during

subthreshold depolarization. **C2)** Interspike interval (ISI) was significantly increased during

optical activation of VIP-INs (blue circles, p: 0.016, paired t-test, n=11) compared to light off

trials (black circles). **D1)** Spike responses in pSST-INs during TBS in control (magenta trace)

and during pulsed light (blue trace).  **D2)** pSST-IN firing rates decreased during TBS on light

trials (blue) versus control (black circles, p:0.002, paired t-test, n=11).

**Figure 5: VIP-interneuron activation promotes associative LTP. A1)** Circuit schematic: VIP-

INs express ChR2 and were activated using theta pulsed light during L1a+L1b pairing. **A2)** VIP-

IN responses during TBS without (black) and with light (blue). FRs increase during pairing with

light (blue circles, p<0.05, WSR, n=7). **B)** Top: Activation of VIP-INs enhanced PN

depolarization during TBS stimulation (blue vs. black trace). Bottom: PN depolarization during

VIP-IN activation (blue trace) is reduced by APV (gray trace).  **C1)**  Average EPSP amplitude

(mV) pre (-5 to 0 min) and post induction (25-30 min) with VIP-IN activation (green) or with VIP-

IN activation plus bath application of APV (black). **C2)**  Average slope (mV/ms) of the rising

phase of the EPSP pre and post induction, colors as in C1. **C3)** Normalized EPSP amplitude

and slope following pairing with VIP-IN activation (green circles) or with activation plus APV

(black). **D1)** Representative average traces pre (gray) and post induction with VIP-IN activation

(green) or activation plus APV (Black). **D2)** Time course of normalized EPSP amplitude pre and

post induction with activation of VIP-INs (blue box, t=0) in control (green) and with APV (black).

**D3)** Time course of normalized EPSP slope, colors as in D2.

**References**

**References**

Abbott, L. F. and S. B. Nelson (2000). "Synaptic plasticity: taming the beast." <u>Nat Neurosci</u> **3**

**Suppl**: 1178-1183.

Alitto, H. J. and Y. Dan (2012). "Cell-type-specific modulation of neocortical activity by basal

forebrain input." <u>Front Syst Neurosci</u> **6**: 79.

Artinian, J. and J. C. Lacaille (2018). "Disinhibition in learning and memory circuits: New

vistas for somatostatin interneurons and long-term synaptic plasticity." <u>Brain Res Bull</u> **141**:

20-26.

Bathellier, B., T. W. Margrie and M. E. Larkum (2009). "Properties of piriform cortex

pyramidal cell dendrites: implications for olfactory circuit design." <u>J Neurosci</u> **29**(40):

12641-12652.

Bolding, K. A. and K. M. Franks (2018). "Recurrent cortical circuits implement

concentration-invariant odor coding." <u>Science</u> **361**(6407).

Carey, R. M. and M. Wachowiak (2011). "Effect of sniffing on the temporal structure of

mitral/tufted cell output from the olfactory bulb." <u>J Neurosci</u> **31**(29): 10615-10626.

Chapuis, J. and D. A. Wilson (2013). "Cholinergic modulation of olfactory pattern

separation." <u>Neurosci Lett</u> **545**: 50-53.

Cohen, Y., D. A. Wilson and E. Barkai (2015). "Differential modifications of synaptic weights

during odor rule learning: dynamics of interaction between the piriform cortex with lower

and higher brain areas." <u>Cereb Cortex</u> **25**(1): 180-191.

Franks, K. M., M. J. Russo, D. L. Sosulski, A. A. Mulligan, S. A. Siegelbaum and R. Axel (2011).

"Recurrent circuitry dynamically shapes the activation of piriform cortex." <u>Neuron</u> **72**(1):

49-56.

Fu, Y., J. M. Tucciarone, J. S. Espinosa, N. Sheng, D. P. Darcy, R. A. Nicoll, Z. J. Huang and M. P.

Stryker (2014). "A cortical circuit for gain control by behavioral state." <u>Cell</u> **156**(6): 1139-

1152.

Haberly, L. B. (2001). "Parallel-distributed processing in olfactory cortex: new insights

from morphological and physiological analysis of neuronal circuitry." <u>Chem Senses</u> **26**(5):

551-576.

Haberly, L. B. and J. L. Price (1977). "The axonal projection patterns of the mitral and tufted
cells of the olfactory bulb in the rat." Brain Res **129**(1): 152-157.

Haberly, L. B. and J. L. Price (1978). "Association and commissural fiber systems of the
olfactory cortex of the rat. II. Systems originating in the olfactory peduncle." J Comp Neurol
**181**(4): 781-807.

Igarashi, K. M., N. Ieki, M. An, Y. Yamaguchi, S. Nagayama, K. Kobayakawa, R. Kobayakawa,
M. Tanifuji, H. Sakano, W. R. Chen and K. Mori (2012). "Parallel mitral and tufted cell
pathways route distinct odor information to different targets in the olfactory cortex." J
Neurosci **32**(23): 7970-7985.

Illig, K. R. (2005). "Projections from orbitofrontal cortex to anterior piriform cortex in the
rat suggest a role in olfactory information processing." J Comp Neurol **488**(2): 224-231.

Illig, K. R. and L. B. Haberly (2003). "Odor-evoked activity is spatially distributed in
piriform cortex." J Comp Neurol **457**(4): 361-373.

Jackson, J., I. Ayzenshtat, M. M. Karnani and R. Yuste (2016). "VIP+ interneurons control
neocortical activity across brain states." J Neurophysiol **115**(6): 3008-3017.

Johenning, F. W., P. S. Beed, T. Trimbuch, M. H. Bendels, J. Winterer and D. Schmitz (2009).
"Dendritic compartment and neuronal output mode determine pathway-specific long-term
potentiation in the piriform cortex." The Journal of Neuroscience **29**(43): 13649-13661.

Kanter, E. D. and L. B. Haberly (1993). "Associative long-term potentiation in piriform
cortex slices requires GABAA blockade." J Neurosci **13**(6): 2477-2482.

Kanter, E. D., A. Kapur and L. B. Haberly (1996). "A dendritic GABAA-mediated IPSP
regulates facilitation of NMDA-mediated responses to burst stimulation of afferent fibers in
piriform cortex." J Neurosci **16**(1): 307-312.

Karnani, M. M., J. Jackson, I. Ayzenshtat, A. Hamzehei Sichani, K. Manoocheri, S. Kim and R.
Yuste (2016). "Opening Holes in the Blanket of Inhibition: Localized Lateral Disinhibition
by VIP Interneurons." J Neurosci **36**(12): 3471-3480.

Kepecs, A., N. Uchida and Z. F. Mainen (2007). "Rapid and precise control of sniffing during
olfactory discrimination in rats." J Neurophysiol **98**(1): 205-213.

Kuchibhotla, K. V., J. V. Gill, G. W. Lindsay, E. S. Papadoyannis, R. E. Field, T. A. Sten, K. D.
Miller and R. C. Froemke (2017). "Parallel processing by cortical inhibition enables context-
dependent behavior." Nat Neurosci **20**(1): 62-71.

Kumar, A., O. Schiff, E. Barkai, B. W. Mel, A. Poleg-Polsky and J. Schiller (2018). "NMDA
spikes mediate amplification of inputs in the rat piriform cortex." Elife **7**.

Large, A. M., N. A. Kunz, S. L. Mielo and A. M. Oswald (2016). "Inhibition by Somatostatin
Interneurons in Olfactory Cortex." Front Neural Circuits **10**: 62.

Large, A. M., N. W. Vogler, S. Mielo and A. M. Oswald (2016). "Balanced feedforward
inhibition and dominant recurrent inhibition in olfactory cortex." Proc Natl Acad Sci U S A
**113**(8): 2276-2281.

Lebel, D., Y. Grossman and E. Barkai (2001). "Olfactory learning modifies predisposition for
long-term potentiation and long-term depression induction in the rat piriform (olfactory)
cortex." Cereb Cortex **11**(6): 485-489.

Lee, S., J. Hjerling-Leffler, E. Zagha, G. Fishell and B. Rudy (2010). "The largest group of
superficial neocortical GABAergic interneurons expresses ionotropic serotonin receptors."
The Journal of Neuroscience **30**(50): 16796-16808.

Lee, S., I. Kruglikov, Z. J. Huang, G. Fishell and B. Rudy (2013). "A disinhibitory circuit
mediates motor integration in the somatosensory cortex." Nat Neurosci **16**(11): 1662-
1670.

Li, W., E. Luxenberg, T. Parrish and J. A. Gottfried (2006). "Learning to smell the roses:
experience-dependent neural plasticity in human piriform and orbitofrontal cortices."
Neuron **52**(6): 1097-1108.

Linster, C., M. Maloney, M. Patil and M. E. Hasselmo (2003). "Enhanced cholinergic
suppression of previously strengthened synapses enables the formation of self-organized
representations in olfactory cortex." Neurobiol Learn Mem **80**(3): 302-314.

Linster, C., B. P. Wyble and M. E. Hasselmo (1999). "Electrical stimulation of the horizontal
limb of the diagonal band of broca modulates population EPSPs in piriform cortex." J
Neurophysiol **81**(6): 2737-2742.

Lucas, E. K. and R. L. Clem (2018). "GABAergic interneurons: The orchestra or the
conductor in fear learning and memory?" Brain Res Bull **141**: 13-19.

Luna, V. M. and N. E. Schoppa (2008). "GABAergic circuits control input-spike coupling in
the piriform cortex." J Neurosci **28**(35): 8851-8859.

Madisen, L., T. Mao, H. Koch, J. M. Zhuo, A. Berenyi, S. Fujisawa, Y. W. Hsu, A. J. Garcia, 3rd, X.
Gu, S. Zanella, J. Kidney, H. Gu, Y. Mao, B. M. Hooks, E. S. Boyden, G. Buzsaki, J. M. Ramirez, A.

R. Jones, K. Svoboda, X. Han, E. E. Turner and H. Zeng (2012). "A toolbox of Cre-dependent
optogenetic transgenic mice for light-induced activation and silencing." Nat Neurosci
**15**(5): 793-802.

Malenka, R. C. and M. F. Bear (2004). "LTP and LTD: an embarrassment of riches." Neuron
**44**(1): 5-21.

Mombaerts, P., F. Wang, C. Dulac, S. K. Chao, A. Nemes, M. Mendelsohn, J. Edmondson and R.
Axel (1996). "Visualizing an olfactory sensory map." Cell **87**(4): 675-686.

Patil, M. M. and M. E. Hasselmo (1999). "Modulation of inhibitory synaptic potentials in the
piriform cortex." J Neurophysiol **81**(5): 2103-2118.

Patil, M. M., C. Linster, E. Lubenov and M. E. Hasselmo (1998). "Cholinergic agonist
carbachol enables associative long-term potentiation in piriform cortex slices." J
Neurophysiol **80**(5): 2467-2474.

Pfeffer, C. K., M. Xue, M. He, Z. J. Huang and M. Scanziani (2013). "Inhibition of inhibition in
visual cortex: the logic of connections between molecularly distinct interneurons." Nat
Neurosci **16**(8): 1068-1076.

Pi, H. J., B. Hangya, D. Kvitsiani, J. I. Sanders, Z. J. Huang and A. Kepecs (2013). "Cortical
interneurons that specialize in disinhibitory control." Nature **503**(7477): 521-524.

Poo, C. and J. S. Isaacson (2007). "An early critical period for long-term plasticity and
structural modification of sensory synapses in olfactory cortex." J Neurosci **27**(28): 7553-
7558.

Poo, C. and J. S. Isaacson (2009). "Odor representations in olfactory cortex: "sparse" coding,
global inhibition, and oscillations." Neuron **62**(6): 850-861.

Poo, C. and J. S. Isaacson (2011). "A major role for intracortical circuits in the strength and
tuning of odor-evoked excitation in olfactory cortex." Neuron **72**(1): 41-48.

Porter, J. T., B. Cauli, K. Tsuzuki, B. Lambolez, J. Rossier and E. Audinat (1999). "Selective
excitation of subtypes of neocortical interneurons by nicotinic receptors." The Journal of
Neuroscience **19**(13): 5228-5235.

Pronneke, A., M. Witte, M. Mock and J. F. Staiger (2019). "Neuromodulation Leads to a
Burst-Tonic Switch in a Subset of VIP Neurons in Mouse Primary Somatosensory (Barrel)
Cortex." Cereb Cortex.

Rennaker, R. L., C. F. Chen, A. M. Ruyle, A. M. Sloan and D. A. Wilson (2007). "Spatial and
temporal distribution of odorant-evoked activity in the piriform cortex." J Neurosci **27**(7):
1534-1542.

Saar, D., Y. Grossman and E. Barkai (2002). "Learning-induced enhancement of
postsynaptic potentials in pyramidal neurons." J Neurophysiol **87**(5): 2358-2363.

Saar, D., I. Reuveni and E. Barkai (2012). "Mechanisms underlying rule learning-induced
enhancement of excitatory and inhibitory synaptic transmission." J Neurophysiol **107**(4):
1222-1229.

Sosulski, D. L., M. L. Bloom, T. Cutforth, R. Axel and S. R. Datta (2011). "Distinct
representations of olfactory information in different cortical centres." Nature **472**(7342):
213-216.

Stettler, D. D. and R. Axel (2009). "Representations of odor in the piriform cortex." Neuron
**63**(6): 854-864.

Stokes, C. C. and J. S. Isaacson (2010). "From dendrite to soma: dynamic routing of
inhibition by complementary interneuron microcircuits in olfactory cortex." Neuron **67**(3):
452-465.

Strauch, C. and D. Manahan-Vaughan (2018). "In the Piriform Cortex, the Primary Impetus
for Information Encoding through Synaptic Plasticity Is Provided by Descending Rather
than Ascending Olfactory Inputs." Cereb Cortex **28**(2): 764-776.

Stripling, J. S., D. K. Patneau and C. A. Gramlich (1988). "Selective long-term potentiation in
the pyriform cortex." Brain Res **441**(1-2): 281-291.

Sturgill, J. F. and J. S. Isaacson (2015). "Somatostatin cells regulate sensory response fidelity
via subtractive inhibition in olfactory cortex." Nat Neurosci **18**(4): 531-535.

Suzuki, N. and J. M. Bekkers (2010a). "Distinctive classes of GABAergic interneurons
provide layer-specific phasic inhibition in the anterior piriform cortex." Cereb Cortex
**20**(12): 2971-2984.

Suzuki, N. and J. M. Bekkers (2010b). "Inhibitory neurons in the anterior piriform cortex of
the mouse: classification using molecular markers." J Comp Neurol **518**(10): 1670-1687.

Taniguchi, H., M. He, P. Wu, S. Kim, R. Paik, K. Sugino, D. Kvitsiani, Y. Fu, J. Lu, Y. Lin, G.
Miyoshi, Y. Shima, G. Fishell, S. B. Nelson and Z. J. Huang (2011). "A resource of Cre driver

lines for genetic targeting of GABAergic neurons in cerebral cortex." <u>Neuron</u> **71**(6): 995-

1013.

Williams, L. E. and A. Holtmaat (2019). "Higher-Order Thalamocortical Inputs Gate Synaptic

Long-Term Potentiation via Disinhibition." <u>Neuron</u> **101**(1): 91-102 e104.

Wilson, D. A. and R. M. Sullivan (2011). "Cortical processing of odor objects." <u>Neuron</u> **72**(4):

506-519.

Xu, H., H. Y. Jeong, R. Tremblay and B. Rudy (2013). "Neocortical somatostatin-expressing

GABAergic interneurons disinhibit the thalamorecipient layer 4." <u>Neuron</u> **77**(1): 155-167.

Zaborszky, L., J. Carlsen, H. R. Brashear and L. Heimer (1986). "Cholinergic and GABAergic

afferents to the olfactory bulb in the rat with special emphasis on the projection neurons in

the nucleus of the horizontal limb of the diagonal band." <u>J Comp Neurol</u> **243**(4): 488-509.

**Figure 1: Associative LTP of intracortical synapses is gated by dendritic disinhibition.**
A) Schematic of APC circuit and stimulation paradigm. Strong (s) TBS of afferents (L1a) is
paired with weak (w) single-pulses at the intracortical pathway (L1b). Gabazine (GZ) is focally
applied to the dendrite region in L1b (tan oval). B1) Representative average traces pre (gray)
and post induction in control (black) and with GZ(Tan). B2) Average EPSP amplitude (mV) pre
(-5 to 0 min) and post induction (25-30 min) with inhibition intact (black) or with dendritic GZ
(tan). B3) Average slope (mV/ms) of the 20-80% rising phase of the EPSP pre and post
induction. C1) Time course of normalized EPSP amplitude pre and post induction (grey box,
t=0) in control (black) and with dendritic Gabazine (tan). C2) Time course of normalized EPSP
slope, colors as in C2. C3) Normalized EPSP amplitude, and slope 30 min post induction in
control (black) and with GZ (tan).

**Figure 2: SST-interneuron inactivation promotes associative LTP.** A1) Schematic: SST-INs express Archaerhodopsin (Arch). A2) SST-IN spike responses during TBS in control (black) and inactivated (green) conditions (*p<0.05, WSR test). B) PN responses for a single TBS burst. Left : Inactivation of SST-INs enhanced PN depolarization (green vs. black trace). Right: Depolarization during SST-IN inactivation (green) is reduced by APV (gray trace). C1) Average EPSP amplitude (mV) pre (-5 to 0 min) and post induction (25-30 min) with SST-inactivation (Magenta) or with SST-inactivation plus bath application of the NMDAR antagonist, APV (black). C2) Average slope (mV/ms) of the 20-80% rising phase of the EPSP pre and post induction, colors as in B2. C3) Normalized EPSP amplitude and slope following pairing with SST-IN inactivation (magenta circles). LTP is blocked by bath application APV (black). D1) Representative average traces pre (gray) and post induction in with SST-IN inactivation (magenta) or inactivation plus APV (Black). D2) Time course of normalized EPSP amplitude pre and post induction with inactivation of SST-INs (green box, t=0) in control (magenta) and with bath application of APV (black). D3) Time course of normalized EPSP slope, colors as in D2.

**Figure 3: PV-interneuron inactivation during induction promotes associative LTP.** A1) Schematic: PV-INs express Archaerhodopsin (Arch). A2) PV-IN spike responses during TBS in control (black) and inactivated (green) conditions. B) Representative PN responses for a single TBS for control (black) versus PV-IN inactivation (green). C1) Average EPSP amplitude (mV) pre (-5 to 0 min) and post induction (25-30 min) with PV-inactivation (Blue) or with PV-inactivation plus bath application of the NMDAR antagonist, APV (black). C2) Average slope (mV/ms) of the 20-80% rising phase of the EPSP pre and post induction, colors as in B2. C3) Normalized EPSP amplitude and slope following pairing with PV-IN inactivation (blue circles) or with inactivation plus bath application APV (black). D1) Representative average traces pre (gray) and post induction in with PV-IN inactivation (blue) or inactivation plus APV (Black). D2) Time course of normalized EPSP amplitude pre and post induction with inactivation of PV-INs (green box, t=0) in control (blue) and with bath application of APV (black). D3) Time course of normalized EPSP slope, colors as in D2.

**Figure 3: Inhibition by VIP-interneurons in piriform cortex.** A1) VIP-INs express ChR2. Optically evoked IPSCs were recorded in putative(p) pSST-INs (magenta), pPV-INs (blue) and PNs (black). A2) Responses of the same neurons (A1) to ten light pulses (100 ms duration, 4 Hz). B1) IPSC amplitude was stronger in pSST-INs versus pPV-INs (*p:0.014) or PNs (*p:0.042, ANOVA). B2) In pSST-INs, IPSC amplitude diminishes by the 5th pulse of theta stimulation (**p: 0.009, paired t-test). C1) IPSCs from VIP-INs delay pSST-IN spike responses during suprathreshold depolarization (4 overlaid traces) Left: Control, Right: activation of VIP-INs (100 ms pulse, blue). Magenta trace: VIP-IN mediated IPSC during subthreshold depolarization. C2) Interspike interval (ISI) was significantly increased during optical activation of VIP-INs (blue circles, p: 0.016, paired t-test, n=11) compared to light off trials (black circles). D1) Spike responses in pSST-INs during TBS in control (magenta trace) and during pulsed light (blue trace). D2) pSST-IN firing rates decreased during TBS on light trials (blue) versus control (black circles, p:0.002, paired t-test, n=11).

**Figure 5: VIP-interneuron activation promotes associative LTP**. A1) Circuit schematic: VIP-INs express ChR2 and were activated using theta pulsed light during L1a+L1b pairing. A2) VIP-IN responses during TBS without (black) and with light (blue). FRs increase during pairing with light (blue circles, p<0.05, WSR, n=7). B) Top: Activation of VIP-INs enhanced PN depolarization during TBS stimulation (blue vs. black trace). Bottom: PN depolarization during VIP-IN activation (blue trace) is reduced by APV (gray trace). C1) Average EPSP amplitude (mV) pre (-5 to 0 min) and post induction (25-30 min) with VIP-IN activation (green) or with VIP-IN activation plus bath application of APV (black). C2) Average slope (mV/ms) of the rising phase of the EPSP pre and post induction, colors as in C1. C3) Normalized EPSP amplitude and slope following pairing with VIP-IN activation (green circles) or with activation plus APV (black). D1) Representative average traces pre (gray) and post induction in with VIP-IN activation (green) or activation plus APV (Black). D2) Time course of normalized EPSP amplitude pre and post induction with activation of VIP-INs (blue box, t=0) in control (green) and with APV (black). D3) Time course of normalized EPSP slope, colors as in D2.