Telomere length and *TERT* expression are associated with age in almond (*Prunus dulcis*)

Katherine M. D'Amico-Willman[1,2,¶], Elizabeth Anderson[3,¶], Thomas M. Gradziel[4], and Jonathan

Fresnedo-Ramírez[1,2*]

[1] Department of Horticulture and Crop Science, Ohio Agricultural Research and Development

Center, The Ohio State University, Wooster, OH 44691

[2] Center for Applied Plant Sciences, The Ohio State University, Columbus, OH 43210

[3] College of Wooster, Wooster, OH 44691

[4] Department of Plant Sciences, University of California, Davis, CA 95616

[*]For correspondence (fresnedoramirez.1@osu.edu)

[¶]Authors contributed equally

Abstract

While it is well known that all organisms age, our understanding of how aging occurs varies
dramatically among species. The aging process in perennial plants is not well defined, yet can
have implications on production and yield of valuable fruit and nut crops. Almond, a relevant nut
crop, exhibits an age-related disorder known as non-infectious bud failure (BF) that affects
vegetative bud development, indirectly affecting kernel-yield. This species and disorder present
an opportunity to address aging in a commercially-relevant and vegetatively-propagated,
perennial crop threatened by an aging-related disorder. In this study, we tested the hypothesis
that telomere length and/or *TERT* expression can serve as biomarkers of aging in almond using
both whole-genome sequencing data and leaf samples collected from distinct age cohorts over a
two-year period. To measure telomere lengths, we employed both *in silico* and molecular
approaches. We also measured expression of *TERT*, a subunit of the enzyme telomerase, which
is responsible for maintaining telomere lengths. Results from this work show a marginal but
significant association between both telomere length measured by monochrome multiplex
quantitative PCR and *TERT* expression, and age of almond seedlings. These results suggest that
as almonds age, *TERT* expression decreases and telomeres shorten. This work provides valuable
information on potential biomarkers of perennial plant aging, contributing to our limited
knowledge of this process. In addition, translation of this information will provide opportunities
to address BF in almond breeding and nursery propagation.
32

33   **Keywords**

34   Perennial plant aging, biomarker, telomerase

Introduction

The current concept and study of aging is centered primarily around mammals with research focused on circumventing deleterious impacts on health (Kirkwood, 2005; Sanders and Newman, 2013). However, all eukaryotic organisms exhibit signals of aging, resulting in the deterioration of key biological processes and subsequent decrease in health, performance, and fitness of individuals. Perennial plants represent a unique model to address the aging process and its impact since these species undergo cycles of dormancy and growth, and maintain the ability to reproduce for multiple years. The aging process of perennial plants is relevant due to the longevity and economic importance of perennial crops such as fruit and nut trees (Munné-Bosch, 2007; Brutovská *et al.*, 2013; Thomas, 2013). Individual trees can remain productive in orchards for decades; however, aging in plants and its implications for growth and reproduction are neglected areas of research with potential consequences for production, management, conservation, and breeding.

The lack of understanding of aging in perennials is partly due to the complexity in measuring and conceptualizing age in perennial plant species since aging occurs chronologically and ontogenetically in opposite directions (Poethig, 2003). Chronologic age can be defined as the amount of time since tissue/organ formation (e.g. human skin cells replenish every few days, meaning each cell is typically a day or 2-days old), while ontogenetic age refers more to developmental time and allows for the accumulation of mutations or chromosomal alterations (e.g. 2-day old skin cells at age 6 compared to 2-day old skin cells at age 60). Agriculturally relevant perennials are often vegetatively propagated (i.e. cloned), blurring the distinction between ontogenetic and chronologic age, and tend to be grown under intensive management. The limitation in determining age in perennials creates a need to identify biomarkers in these species that enable ontogenetic age estimation.

Almond (*Prunus dulcis* [Mill.] D.A.Webb; Fig. 1) is an economically relevant, Rosaceous crop, subject to intense horticultural management to maintain maximum nut production. In California, the almond industry is estimated to contribute ~$11 billion to the state's GDP annually (Almond Board of California, 2019). Top-producing almond cultivars, some of which were first obtained more than 100 years ago, are produced for commercial orchards via vegetative propagation (Micke, 1996; Wickson, 1914). As orchards age (after 20-25

65 years), trees are replaced with "new" clones of typically the same cultivar to maintain

66 homogeneity in quality and high levels of production (Micke, 1996).

67   Almond exhibits an age-related disorder known as non-infectious bud-failure (BF)

affecting vegetative bud development in the spring (Micke, 1996). Genotypes exhibiting this

disorder show characteristic dieback at the top of the canopy, and severe levels of BF can result

in up to 50% yield loss (Gradziel et al. 2013). Empirical evidence shows BF is associated with

age (Kester *et al.*, 2004); however, as almonds are produced primarily through vegetative

propagation rather than by seed, their true ontogenetic age and thus susceptibility to BF can be

difficult to assess (Micke, 1996). Biomarkers indicative of age would be valuable to growers,

breeders, and producers to screen germplasm. Thus, almond represents a potential model species

for the study of aging in perennials due to its economic relevance, the abundance of available

germplasm and breeding programs, and the exhibition of an age-related disorder.

77   One biomarker of aging is telomere length measurement (Sanders and Newman, 2013;

Marioni *et al.*, 2016; Runov *et al.*, 2015), which has been primarily studied in animals.

Telomeres are nucleoproteins that cap the end of chromosomes, preventing premature instability

of genomic material and cellular senescence (Watson and Riha, 2011). Telomeres tend to shorten

over mitotic cellular divisions due to decreased levels of telomerase, an enzyme that supports

telomere replication during the S-phase of the cell cycle (Nelson *et al.*, 2014). Over mitotic cell

divisions, telomeres eventually reach a critical minimum length at which point the cell senesces

and dies due to genome instability resulting from single stranded DNA at the ends of

chromosomes (Hemann *et al.*, 2001). This progressive shortening is proposed as a marker of

aging in mammalian cells and is linked to physiological deterioration and some age-related

disorders (Sanders and Newman, 2013; Watson and Riha, 2011; Aviv and Shay, 2018). Plant

chromosomes also contain telomeres with similar functions. While the relationship between

telomeres and the aging process is not as clearly defined in plants as in animals, previous work

shows associations between telomere length and various stages of plant development (Zachová *et*

al., 2013; Watson and Riha, 2011; Procházková Schrumpfová *et al.*, 2019) suggesting telomere

length could be a suitable biomarker of age in plants.

93   Given that telomerase activity modulates telomere length, expression of genes involved

in the telomerase biosynthetic pathway could also serve as biomarkers for aging (Fitzgerald *et*

al., 1996; De la Torre-Espinosa *et al.*, 2020; Boccardi and Paolisso, 2014; Anchelin *et al.*, 2011;

Fossel, 1998). TERT is the catalytic subunit of the telomerase enzyme (Oguchi *et al.*, 1999) and

the RNA subunit (TR) functions as the template for reverse transcription (Procházková

Schrumpfová *et al.*, 2019). Expression of *TERT* is shown to affect telomerase activity (Sweetlove

and Gutierrez, 2019; Jurečková *et al.*, 2017). In Arabidopsis, increased *TERT* expression is

linked to proportional increases in telomerase activity and telomere length (Zangi *et al.*, 2019;

Fitzgerald *et al.*, 1999) which is in turn linked to age. Since *TERT* is tied to telomere length in

both plants and animals, its expression may also serve as an indicator of age in plants (Watson

and Riha, 2010).

This study tests the hypothesis that telomere length and *TERT* expression in almond are

associated with age and can thus serve as biomarkers of aging in this species. Both telomere

length and *TERT* expression show promise as diagnostic biomarkers since they can be measured

in a high-throughput manner via *in silico* and molecular approaches (Nersisyan and Arakelyan,

2015; Montpetit *et al.*, 2014; Cawthon, 2009). These approaches build on previous research

examining the relationship between telomere lengths and age in perennial plants (Flanary and

Kletetschka, 2005; Moriguchi *et al.*, 2007; Liang *et al.*, 2015; Liu *et al.*, 2007). The goal of this

work is to advance our understanding and provide a model for the study of aging and its

implications in perennial plant species.

Materials and Methods

Whole-genome Sequencing Data

To perform *in silico* mean telomere length estimation, fastq files produced from whole-genome

sequencing of nine almond accessions were downloaded to the Ohio Super Computer (Anon,

1987) from the National Center for Biotechnology Information Sequence Read Archive from

bioprojects PRJNA339570 and PRJNA339144. Accessions were selected for use in this study if

the age of the individual was included in the metadata for the biosample entry on NCBI SRA.

Table 1 includes the SRA biosample number for each almond accession used as well as

accession name, cultivar name (if available), and age.

In silico Mean Telomere Length Estimation

Mean telomere lengths were estimated for nine almond accessions using files in fastq format

containing whole-genome sequencing data for each individual. These fastq files were used as

input in the program Computel v. 1.2 (Nersisyan and Arakelyan, 2015). To estimate mean

telomere lengths, the computel.sh script was run with the following parameters: -nchr 8 -

lgenome 227411381 -pattern CCCTAAA -minseed 14 using the estimated length of the peach

genome based on the v2 assembly (Initiative *et al.*, 2013). All work was performed using the

Ohio Supercomputer Center computational resources (Anon, 1987).

Plant Material

Leaf samples for this study were collected in May 2018 and 2019 from almond breeding

selections located at the Wolfskill Experimental Orchards (Almond Breeding Program,

University of California – Davis, Winters, CA). Tissue was harvested from the canopy of a total

of 21 unique individuals representing distinct age cohorts (Table 2). Samples were immediately

frozen on ice and stored at -20 °C until shipment overnight on dry ice to the Ohio Agricultural

Research and Development Center (OARDC – Wooster, OH). Samples were stored at -20 °C

until processing, and all subsequent experimental procedures were conducted at the OARDC.

DNA and RNA Extraction

DNA was extracted from the age cohort samples using the Omega E-Z 96® Plant DNA Kit

(Omega Bio-tek, Norcross, GA) with slight modification. Briefly, 100 mg of leaf material was

weighed in 2.0 mL tubes containing two 1.6 mm steel beads and kept frozen in liquid nitrogen.

Samples were ground in a 2000 Geno/Grinder® (SPEX SamplePrep, Metuchen, NJ) in two 48-

well cryo-blocks frozen in liquid nitrogen. Following a 65 °C incubation, samples were

incubated on ice for 20 minutes, treated with 10 µL of RNase solution (2.5 ul RNase [Omega

Bio-tek, Norcross, GA] + 7.5 µl TE pH 8), equilibrated through addition of 150 µl Equilibration

Buffer (3 M NaOH), incubated at room temperature for four minutes, and centrifuged at 4,400

rpm for two minutes prior to the addition of SP3 buffer. Concentration and quality were analyzed

using a NanoDrop™ 1000 spectrophotometer and a Qubit 4 Fluorometer with a dsDNA HS

Assay Kit (ThermoFisher Scientific, Waltham, MA).

RNA was extracted following the protocol outline in Gambino *et al.* (2008) with slight

modifications. Briefly, leaf material was ground in liquid nitrogen using a mortar and pestle, and

150 mg of tissue was weighed into a 2.0 mL microfuge tube frozen in liquid nitrogen. To extract

RNA, 900 µL CTAB extraction buffer (2% CTAB, 2.5% PVP-40, 2 M NaCl, 100 mM Trish-HCl

pH 8.0, 25 mM EDTA pH 8.0, 2% Beta-mercaptoethanol added before use) was added to each

tube and samples were incubated at 65 °C for ten minutes. Following incubation, two phase

separations were performed using an equal volume of chloroform:isoamyl alcohol (24:1). RNA

was precipitated in 3 M lithium chloride and incubated on ice for 30 minutes, and samples were

pelleted by centrifugation at 21,000 x g for 15 minutes. Pellets were then resuspended in 500 μL

pre-warmed SSTE buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA pH 8.0, 1% SDS, 1 M NaCl)

followed by a phase separation with an equal volume of chloroform:isoamyl alcohol (24:1). A

final precipitation was performed using 0.7 volumes chilled 100% isopropanol. RNA was

pelleted and washed with 70% ethanol before being resuspended in 30 μL nuclease-free water. A

DNase treatment was performed using DNA-*free*™ DNA Removal Kit (ThermoFisher

Scientific) according to the manufacturer's instructions. All materials used for extraction were

nuclease- free and cleaned with RNaseZap™ RNase decontamination wipes (ThermoFisher

Scientific) prior to use. All centrifugation steps were performed at 4°C. RNA quality and

concentration were assessed using a NanoDrop™ 1000 spectrophotometer and a Qubit 4

Fluorometer with an RNA HS Assay Kit (ThermoFisher Scientific).

Monochrome Multiplex Quantitative PCR (MMQPCR) to Measure Relative Telomere Lengths

MMQPCR was conducted following the protocol outlined in Vaquero-Sedas and Vega-Palas

(2014) with minimal modifications. Primer sequences for genes used in this study are shown in

Table 3, including primers for the single copy gene, *PP2A*, and for the telomere sequence (Wang

et al., 2014; Vaquero-Sedas and Vega-Palas, 2014). Oligos were synthesized by MilliporeSigma

(Burlington, MA) and resuspended to a concentration of 100 μM upon arrival. Standard curves

were created for each primer pair by pooling six aliquots of DNA isolated from a single clone of

the almond cultivar 'Nonpareil', and performing successive dilutions to 20 ng/μL, 10 ng/μL, 1

ng/μL, 0.5 ng/μL, and 0.25 ng/μL. Reactions were carried out in triplicate for each primer by

concentration combination.

Isolated DNA from the age cohort samples was diluted to 20 ng/μL. Multiplex reactions

were carried out in sextuplicate for each replicate within the age cohorts in a 10 μL volume using

QuantaBio PerfeCTa SYBR® Green SuperMix (Quanta Biosciences, Beverly, MA) (2X),

forward and reverse primers (100 nM each), and 20 ng template DNA according to the

manufacturer's instructions. Reactions were performed in a Bio Rad C1000 Touch Thermal

Cycler (Bio Rad Laboratories, Hercules, CA) using the following program: initial denaturation at

95 °C for 3 minutes followed by 2 cycles of incubation at 94 °C for 15 seconds and annealing at

49 °C for 15 seconds; telomere and *PP2A* amplicons were generated following 35 cycles at 95

°C for 30 seconds, 59 °C for 1 minute, 72 °C for 30 seconds, 84 °C for 15 seconds and 85 °C for

15 seconds; final incubation at 72 °C for 1 minute. Melting curve analysis was performed at a

temperature range of 74-85 °C for both primer pairs to ensure no non-specific amplification.

cDNA Synthesis and Quantitative Reverse Transcriptase PCR (qRT-PCR) to Measure Relative

Expression of TERT

Reactions were carried out in a 20 μL volume using the Verso™ cDNA synthesis Kit

(ThermoFisher Scientific). One reaction was prepared for each age cohort sample according to

the manufacturer's instructions. Reactions were performed in a MJ Research PTC-200 thermal

cycler using the following program: 42 °C for 30 minutes followed by 95 °C for 2 minutes.

To quantify expression of *TERT* in age cohort individuals, qRT-PCR was performed in

triplicate for each sample. The gene *RPII* from peach was used as a reference (Bastias *et al.*,

2020; Tong *et al.*, 2009), and the sequence for the *TERT* gene was derived from the 'Texas'

genome (https://www.rosaceae.org/analysis/295) using the homologous peach gene sequence as

a reference (Alioto *et al.*, 2020). Primer sequences are shown in Table 2, and all oligos were

synthesized by MilliporeSigma (Burlington, MA) and resuspended to a concentration of 100 μM

upon arrival.

To generate cDNA from the age cohort samples, 100 ng of RNA was used as input in the

Verso cDNA Synthesis Kit (ThermoFisher Scientific) according to the manufacturer's

instructions. To test for relative expression of *TERT*, reactions were carried out in triplicate for

each biological replicate within the age cohorts in a 10 μL volume using QuantaBio PerfeCTa

SYBR® Green SuperMix (Quanta Biosciences) (1X), forward and reverse primers (100 nM),

and cDNA (1 μL) according to the manufacturer's instructions. Reactions were performed in Bio

Rad C1000 Touch Thermal Cycler (Bio Rad Laboratories) using the following program: initial

denaturation at 95 °C for 3 minutes followed by 40 cycles at 95 °C for 15 seconds and 55 °C for

45 seconds. Melt curves were generated at a temperature range of 74-85 °C for both primer pairs

to ensure no non-specific amplification.

Statistical Analysis

Mean telomere lengths generated *in silico* using Computel v. 1.2 for each almond accession were

square root transformed prior to analysis. Mean telomere length was regressed on chronological

age to test for a linear relationship based on *in silico* predictions. Normality was confirmed using

a Shapiro-Wilks test.

Using the standard curve generated with *PP2A* (S) and telomere (T) primers for a

reference almond sample, relative T/S ratios were calculated for each individual sample based on

Cq values for the telomere and *PP2A* products (Vaquero-Sedas and Vega-Palas, 2014). Z-scores

were calculated from the T/S ratios as recommend in Verhulst (2020) for each replicate within

the age cohorts. Normality and homogeneity of variance were confirmed using Shapiro-Wilks

and Bartlett tests. Analysis of variance (ANOVA) was performed for each age cohort followed

by *post hoc* Fisher's LSD and pairwise t-tests. Gene expression data were analyzed according to

guidelines in Bustin et. al (2009), first by normalizing *TERT* expression to that of the reference

gene, *RPII*. Following normalization, data were log-transformed, and normality and homogeneity

of variance were confirmed using Shapiro-Wilks and Bartlett tests. ANOVA was performed for

each age cohort followed by *post hoc* analysis with Tukey's HSD. All analyses were performed

using R v. 3.6.1 and plots were generated using ggplot2 v. 3.3.0.

Results

Associations of telomere length and age in almond

Mean telomere lengths were estimated *in silico* using whole-genome sequencing data for nine

select almond accessions and regressed against accession age following a square root

transformation. Normality of residuals was confirmed using a Shapiro-Wilks test (p-value =

0.318). Linear regression suggests a negative relationship between mean telomere length and age

as depicted in Fig. 2.

Relative telomere lengths were generated for the almond individuals within each of the

age cohorts collected in 2018 (1, 5, 9, and 14 years) and 2019 (2, 7, and 11 years old) using the

MMQPCR approach. Normality of residuals and homogeneity of variance of relative telomere

lengths were confirmed using Shapiro-Wilks (**2018:** p-value = 0.2578; **2019:** p-value = 0.4682)

and Bartlett (**2018:** p-value = 0.1408; **2019:** p-value = 0.4613) tests. ANOVA results for the

linear model, z-score ~ age, were marginally significant in both 2018 and 2019, and subsequent

post hoc Fisher's LSD and pairwise t-tests revealed significant differences between ages 1 and

14 years and 5 and 14 years (Fig. 3a) in the 2018 cohorts, and between ages 2 and 11 years old

(Fig. 3b) in the 2019 cohorts.

TERT gene expression patterns associated with age in almond

Normalized expression of *TERT* was measured for almond samples among the age cohorts

collected in 2018 and 2019 for this study using *RPII* as the reference gene. Normality of

residuals and homogeneity of variance were confirmed using Shapiro-Wilks (**2018:** p-value =

0.694; **2019:** p-value = 0.09456) and Bartlett (**2018:** p-value = 0.6976; **2019:** p-value = 0.3579)

tests. ANOVA results comparing the average log(expression) values for each age cohort revealed

significant differences between cohorts in both 2018 and 2019. *Post hoc* analysis with Tukey's

HSD revealed significant differences in *TERT* expression between ages 1 and 14 years old in the

2018 cohorts (Fig. 4a) and between ages 2 and 11 years old in the 2019 age cohorts (Fig. 4b).

Discussion

Almond, an economically-valuable nut crop, exhibits an aging-related disorder known as non-

infectious bud failure that negatively impacts vegetative development and ultimately, yield. As a

clonally propagated crop, tracking age and thus susceptibility to bud failure is difficult, making

biomarkers of age a valuable resource to circumvent the impacts of aging-related disorders in

almond germplasm. Telomere length is used as a biomarker of age and development of age-

related disorders in mammals, but the association between telomere length and age in plants is

not well-defined (Watson and Riha, 2011; Procházková Schrumpfová *et al.*, 2019). The present

study tests the hypothesis that telomere length and/or *TERT* expression are associated with age in

almond. To test this, both *in silico* and qPCR approaches were utilized to measure telomere

length and estimate *TERT* expression in sets of almond accessions of known chronological age.

In silico analysis was performed using whole-genome sequencing data from nine almond

accessions of known age. Samples were collected from three and four sets of age cohorts over

two years to test for an association between relative telomere length and individual age using the

MMQPCR method as well as between *TERT* expression and age using qRT-PCR.

In silico and quantitative PCR approaches suggest a marginal association between average or

relative telomere length and age in almond

Average telomere length estimated *in silico* using the program Computel (Nersisyan and

Arakelyan, 2015) revealed a non-significant, negative association with age in the almond

accessions tested. This same pattern was shown utilizing MMQPCR and almond leaf samples
collected from different almond age cohorts in 2018 and 2019 where telomere length decreases
with increasing age. The association demonstrated in this study adds to the growing body of
knowledge regarding the complex relationship between telomere length and plant aging.

Previous studies in both *Ginkgo biloba* and *Panax ginseng* showed a pattern of increased
telomere length with increased age, suggesting plants do not follow the same patterns of
telomere shortening as seen in mammals (Liang *et al.*, 2015; Liu *et al.*, 2007). Work in apple
(*Malus domestica*) and *Prunus yedoensis*, both members of Rosaceae like almond, show no
change in telomere lengths with increased plant age over a five year timespan (Moriguchi *et al.*,
2007). In bristlecone pine (*Pinus longaeva*), a long-lived perennial gymnosperm, telomere
lengths measured in needle and root tissues between 0 – 3,500 years old showed a cyclical
pattern of lengthening and shortening with age (Flanary and Kletetschka, 2005). Further, when
analyzing telomere length in relation to tissue differentiation, studies in both barley (*Hordeum*
vulgare) and Scots pine (*Pinus sylvestris*) showed telomere shortening through embryo
development to leaf or needle formation (Aronen and Ryynänen, 2012; Kilian *et al.*, 1995).
Similarly, in silver birch (*Betula pendula*), telomeres shorten when plant are grown in tissue
culture conditions compared to those grown outdoors, suggesting abiotic stressors may induce
telomere shortening (Aronen and Ryynänen, 2014).

The results in almond suggest a pattern closest to what was observed in bristlecone pine
where telomere lengths shorten and lengthen throughout an individual's lifetime. This pattern
could be unique to gymnosperms, however, and needs to be further characterized in angiosperms
including Rosaceous species. While the commercial lifespan of almond is typically less than 30
years, almond can live more than 150 years (Micke, 1996). In this study, the maximum age
tested via the *in silico* approach was 32 years-old and via qPCR was 14 years-old, suggesting
that a wider age-range of trees and a larger sample size could produce a more refined model of
telomere length patterns over time.

Current almond cultivars may also be ontogenetically old, such as 'Nonpareil', the most
relevant US cultivar representing ~40% of acreage, which was first described almost 140 years
ago and has been propagated by budding since (Wickson, 1914; Almond Board of California,
2019). The ontogenetic age of a cultivar may be a factor to consider in the onset of aging-related

disorders like BF in almond. Additionally, it would be interesting to track the change in telomere

length following clonal propagation (through budding) in which plants experience a rejuvenation

process, reverting to a juvenile state for a short period of time (Bonga, 1982). It was further

found that propagating almond from basal epicormic buds, potentially representing

ontogenetically young meristematic tissue, seemed to alleviate BF in resulting clones (Gradziel

et al., 2019). Testing telomere lengths in epicormic tissues could present another avenue to both

track aging in almond and develop biomarkers to predict BF potential in almond.

TERT expression measured by qRT-PCR is a putatively associated with age in almond

accessions

To test the hypothesis that *TERT* expression can serve as a biomarker of age in almond,

expression patterns were tested in cohorts representing either three or four distinct ages over two

years. Results from this work showed a consistent pattern of marginally significant, decreased

expression with increased ontogenetic age. Telomerase was shown to be a modulator of

longevity in humans and other mammals, but work describing telomerase patterns in plants is

limited (Boccardi and Paolisso, 2014; Fitzgerald *et al.*, 1996).

319   A comprehensive study examining telomerase protein activity in carrot (*Daucus carota*),

cauliflower (*Brassica oleracea*), soybean (*Glycine max*), *Arabidopsis thaliana*, and rice (*Oryza*

sativa) demonstrated that, like telomere lengths, protein activity tends to be highest in

undifferentiated tissues like meristematic tissues and is lower in differentiated tissues such as

leaves (Fitzgerald *et al.*, 1996). This result was supported by further work in barley and maize

showing little activity in differentiated tissues (Kilian *et al.*, 1998). These studies were all

performed in annuals or biennials, however, suggesting that telomerase activity does in fact

decrease with increased plant age in these crops. Work in perennials including bristlecone pine,

P. ginseng, and *G. biloba* showed an association between telomerase activity and age, suggesting

patterns unique to perennial plant species (Liang *et al.*, 2015; Flanary and Kletetschka, 2005;

Song *et al.*, 2011). A study in almond could be performed using a wider age-range and larger

sample size to elucidate the effect of age on telomerase activity, similar to what was referenced

above for telomere length measurements. Additionally, many of the studies performed in other

plants examining patterns of telomerase activity focused on protein activity rather than gene

expression. A future study will be necessary in almond to examine the telomerase protein

activity potentially by Western blot or other proteomics approaches to corroborate the

association between *TERT* expression and protein activity.

While a pattern was established in plants demonstrating a direct relationship between

telomerase activity and telomere length, regulation of telomerase is still not well understood in

the plant kingdom (Zachová *et al.*, 2013; Jurečková *et al.*, 2017; Fitzgerald *et al.*, 1996).

Interestingly, work in Arabidopsis has shown a link between DNA methylation and telomere

length, suggesting that this epigenetic mark likely has a role in regulating telomere lengths

potentially by modulating telomerase activity (Vega-Vaquero *et al.*, 2016; Lee and Cho, 2019;

Ogrocká *et al.*, 2014). A study is ongoing in almond to analyze DNA methylation patterns in a

set of almond accessions representing three distinct age cohorts to determine what, if any, impact

age has on methylation profiles.

Despite the limited age-range and small sample size used in this study, a consistent

pattern of both decreased telomere length and decreased *TERT* expression with increased age

was observed over two years of sampling. These results provide a basis for future study and

exploration into the utility of telomere length measurement and/or *TERT* expression or

telomerase activity as biomarkers of aging in almond. Developing a robust biomarker to track

aging in almond, a primarily clonally propagated crop, would allow growers, producers, and

breeders to screen germplasm to eliminate selections or clones with a high susceptibility to age-

related disorders due to advanced ontogenetic age.

References

Alioto, T., Alexiou, K.G., Bardil, A., et al. (2020) Transposons played a major role in the diversification between the closely related almond and peach genomes: results from the almond genome sequence. *The Plant Journal*, **101**, 455–472. 10.1111/tpj.14538.

Almond Board of California (2019) *Almond Almanac*, Modesto, CA: Almond Board of California.

Anchelin, M., Murcia, L., Alcaraz-Pérez, F., García-Navarro, E.M. and Cayuela, M.L. (2011) Behaviour of telomere and telomerase during aging and regeneration in zebrafish. *PLoS One*, **6**. 10.1371/journal.pone.0016955.

Ohio Supercomputer Center. 1987. Ohio Supercomputer Center. Columbus OH: Ohio Supercomputer Center. http://osc.edu/ark:/19495/f5s1ph73.

Aronen, T. and Ryynänen, L. (2014) Silver birch telomeres shorten in tissue culture. *Tree Genetics & Genomes*, **10**, 67–74. 10.1007/s11295-013-0662-4.

Aronen, T. and Ryynänen, L. (2012) Variation in telomeric repeats of Scots pine (*Pinus sylvestris L.*). *Tree Genetics & Genomes*, **8**, 267–275. 10.1186%2F1753-6561-5-S7-O42.

Aviv, A. and Shay, J.W. (2018) Reflections on telomere dynamics and ageing-related diseases in humans. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.*, **373**. 10.1098/rstb.2016.0436.

Bastias, A., Oviedo, K., Almada, R., Correa, F. and Sagredo, B. (2020) Identifying and validating housekeeping hybrid *Prunus* spp. genes for root gene-expression studies. *PLOS ONE*, **15**. 10.1371/journal.pone.0228403.

Boccardi, V. and Paolisso, G. (2014) Telomerase activation: A potential key modulator for human healthspan and longevity. *Ageing Research Reviews*, **15**, 1–5. 10.1016/j.arr.2013.12.006.

Bonga, J.M. (1982) Vegetative propagation in relation to juvenility, maturity, and rejuvenation. In J. M. Bonga and D. J. Durzan, eds. *Tissue Culture in Forestry*. Forestry Sciences. Dordrecht: Springer Netherlands, pp. 387–412. 10.1007/978-94-017-3538-4_13.

Brutovská, E., Sámelová, A., Dušička, J. and Mičieta, K. (2013) Ageing of trees: Application of general ageing theories. *Ageing Research Reviews*, **12**, 855–866. 10.1016/j.arr.2013.07.001.

Bustin, S.A., Benes, V., Garson, J.A., et al. (2009) The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin. Chem.*, **55**, 611–622. 10.1373/clinchem.2008.112797.

Cawthon, R.M. (2009) Telomere length measurement by a novel monochrome multiplex quantitative PCR method. *Nucleic Acids Res*, **37**, e21. 10.1093/nar/gkn1027.

De la Torre-Espinosa, Z.Y., Barredo-Pool, F., Castaño de la Serna, E. and Sánchez-Teyer, L.F. (2020) Active telomerase during leaf growth and increase of age in plants from *Agave tequilana* var. Azul. *Physiology and Molecular Biology of Plants*, **26**, 639-647. 10.1007/s12298-020-00781-7.

Fitzgerald, M.S., McKnight, T.D. and Shippen, D.E. (1996) Characterization and developmental patterns of telomerase expression in plants. *PNAS*, **93**, 14422–14427. 10.1073/pnas.93.25.14422.

Fitzgerald, M.S., Riha, K., Gao, F., Ren, S., McKnight, T.D. and Shippen, D.E. (1999) Disruption of the telomerase catalytic subunit gene from *Arabidopsis* inactivates telomerase and leads to a slow loss of telomeric DNA. *PNAS*, **96**, 14813–14818. 10.1073/pnas.96.26.14813.

Flanary, B.E. and Kletetschka, G. (2005) Analysis of telomere length and telomerase activity in tree species of various life-spans, and with age in the bristlecone pine *Pinus longaeva*. *Biogerontology*, **6**, 101–111. 10.1007/s10522-005-3484-4.

Fossel, M. (1998) Telomerase and the aging cell: Implications for human health. *JAMA*, **279**, 1732–1735. 10.1001/jama.279.21.1732.

Gambino, G., Perrone, I. and Gribaudo, I. (2008) A Rapid and effective method for RNA extraction from different tissues of grapevine and other woody plants. *Phytochemical Analysis*, **19**, 520–525. 10.1002/pca.1078.

Gradziel, T., Lampinen, B. and Preece, J.E. (2019) Propagation from basal epicormic meristems remediates an aging-related disorder in almond clones. *Horticulturae*, **5**, 28. 10.3390/horticulturae5020028.

Gradziel, T.M., Thorpe, M.A., Fresnedo-Ramírez, J., et al. (2013) *Molecular marker based diagnostics for almond non-infectious bud-failure*, Almond Board of California.

Hemann, M.T., Strong, M.A., Hao, L.-Y. and Greider, C.W. (2001) The shortest telomere, not average telomere length, is critical for cell viability and chromosome stability. *Cell*, **107**, 67–77. 10.1016/s0092-8674(01)00504-9.

Initiative, T.I.P.G., Verde, I., Abbott, A.G., et al. (2013) The high-quality draft genome of peach (*Prunus persica*) identifies unique patterns of genetic diversity, domestication and genome evolution. *Nature Genetics*, **45**, 487–494. 10.1038/ng.2586.

Jurečková, J.F., Sýkorová, E., Hafidh, S., Honys, D., Fajkus, J. and Fojtová, M. (2017) Tissue-specific expression of telomerase reverse transcriptase gene variants in Nicotiana tabacum. *Planta*, **245**, 549–561. 10.1007/s00425-016-2624-1.

443 **Kester, D.E. and Jones, R.W.** (1970) Non-infectious bud-failure from breeding programs of
444   almond (*Prunus amygdalus* Batsch). *Journal of the American Society of Horticultural*
445   *Science*, **95**, 492–96.
446

447 **Kester, D.E., Shackel, K.A., Micke, W.C., Viveros, M. and Gradziel, T.M.** (2004)
448   Noninfectious bud failure in `Carmel' almond: I. Pattern of development in vegetative
449   progeny trees. *J. Amer. Soc. Hort. Sci.*, **129**, 244–249. 10.21273/JASHS.129.2.0244.
450

451 **Kilian, A., Heller, K. and Kleinhofs, A.** (1998) Development patterns of telomerase activity in
452   barley and maize. *Plant Mol Biol*, **37**, 621–628. 10.1023/A:1005994629814.
453

454 **Kilian, A., Stiff, C. and Kleinhofs, A.** (1995) Barley telomeres shorten during differentiation
455   but grow in callus culture. *Proc Natl Acad Sci USA*, **92**, 9555. 10.1073/pnas.92.21.9555.
456

457 **Kirkwood, T.B.L.** (2005) Understanding the odd science of aging. *Cell*, **120**, 437–447.
458   10.1016/j.cell.2005.01.027.
459

460 **Lee, W.K. and Cho, M.H.** (2019) Epigenetic aspects of telomeric chromatin in *Arabidopsis*
461   *thaliana*. *BMB Rep*, **52**, 175–180. 10.5483/BMBRep.2019.52.3.047.
462

463 **Liang, J., Jiang, C., Peng, H., Shi, Q., Guo, X., Yuan, Y. and Huang, L.** (2015) Analysis of
464   the age of *Panax ginseng* based on telomere length and telomerase activity. *Sci Rep*, **5**.
465   10.1038/srep07985.
466

467 **Liu, D., Qiao, N., Song, H., Hua, X., Du, J., Lu, H. and Li, F.** (2007) Comparative analysis of
468   telomeric restriction fragment lengths in different tissues of *Ginkgo biloba* trees of
469   different age. *Journal Of Plant Research*, **120**, 523–528. 10.1007/s10265-007-0092-1.
470

471 **Marioni, R.E., Harris, S.E., Shah, S., McRae, A.F., Zglinicki, T. von, Martin-Ruiz, C.,**
472   **Wray, N.R., Visscher, P.M. and Deary, I.J.** (2016) The epigenetic clock and telomere
473   length are independently associated with chronological age and mortality. *Int J*
474   *Epidemiol*, **45**, 424–432. 10.1093/ije/dyw041.
475

476 **Micke, W.C.** (1996) *Almond Production Manual*, UCANR Publications.
477

478 **Montpetit, A.J., Alhareeri, A.A., Montpetit, M., et al.** (2014) Telomere length: A review of
479   methods for measurement. *Nurs Res*, **63**, 289–299. 10.1097/NNR.0000000000000037.
480

481 **Moriguchi, R., Kato, K., Kanahama, K., Kanayama, Y. and Kikuchi, H.** (2007) Analysis of
482   telomere lengths in apple and cherry trees. *Acta Horticulturae*, 389–395.
483   10.17660/ActaHortic.2007.738.47.
484

485 **Munné-Bosch, S.** (2007) Aging in perennials. *Critical Reviews in Plant Sciences*, **26**, 123–138.
486   10.1080/07352680701402487.
487

488   **Nelson, A.D.L., Beilstein, M.A. and Shippen, D.E.** (2014) Plant telomeres and telomerase. In
489   S. H. Howell, ed. *Molecular Biology*. New York, NY: Springer New York, pp. 25–49.
490   10.1007/978-1-4614-7570-5_4.

492   **Nersisyan, L. and Arakelyan, A.** (2015) Computel: Computation of mean telomere length from
493   whole-genome next-generation sequencing data. *PLOS ONE*, **10**.
494   10.1371/journal.pone.0125201.

496   **Ogrocká, A., Polanská, P., Majerová, E., Janeba, Z., Fajkus, J. and Fojtová, M.** (2014)
497   Compromised telomere maintenance in hypomethylated *Arabidopsis thaliana* plants.
498   *Nucleic Acids Res*, **42**, 2919–2931. 10.1093/nar/gkt1285.

500   **Oguchi, K., Liu, H., Tamura, K. and Takahashi, H.** (1999) Molecular cloning and
501   characterization of AtTERT, a telomerase reverse transcriptase homolog in *Arabidopsis*
502   *thaliana*. *FEBS Letters*, **457**, 465–469. 10.1016/s0014-5793(99)01083-2.

504   **Poethig, R.S.** (2003) Phase change and the regulation of developmental timing in plants.
505   *Science*, **301**, 334–336. 10.1126/science.1085328.

507   **Procházková Schrumpfová, P., Fojtová, M. and Fajkus, J.** (2019) Telomeres in plants and
508   humans: Not so different, not so similar. *Cells*, **8**, 58. 10.3390/cells8010058.

510   **Runov, A.L., Vonsky, M.S. and Mikhelson, V.M.** (2015) DNA methylation level and telomere
511   length as a basis for modeling of the biological aging clock. *Cell and Tissue Biology*, **9**,
512   261–264. 10.1134/S1990519X15040094.

514   **Sanders, J.L. and Newman, A.B.** (2013) Telomere length in epidemiology: A biomarker of
515   aging, age-related disease, both, or neither? *Epidemiol Rev*, **35**, 112–131.
516   10.1093/epirev/mxs008.

518   **Song, H., Liu, D., Li, F. and Lu, H.** (2011) Season- and age-associated telomerase activity in
519   *Ginkgo biloba* L. *Mol Biol Rep*, **38**, 1799–1805. 10.1007/s11033-010-0295-8.

521   **Sweetlove, L. and Gutierrez, C.** (2019) The journey to the end of the chromosome: delivering
522   active telomerase to telomeres in plants. *Plant J*, **98**, 193–194. 10.1111/tpj.14328.

524   **Thomas, H.** (2013) Senescence, ageing and death of the whole plant. *New Phytol.*, **197**, 696–
525   711. 10.1111/nph.12047.

527   **Tong, Z., Gao, Z., Wang, F., Zhou, J. and Zhang, Z.** (2009) Selection of reliable reference
528   genes for gene expression studies in peach using real-time PCR. *BMC Mol Biol*, **10**, 71.
529   10.1186/1471-2199-10-71.

531   **Vaquero-Sedas, M.I. and Vega-Palas, M.A.** (2014) Determination of *Arabidopsis thaliana*
532   telomere length by PCR. *Scientific Reports*, **4**. 10.1038/srep05540.

Vega-Vaquero, A., Bonora, G., Morselli, M., Vaquero-Sedas, M.I., Rubbi, L., Pellegrini, M.
and Vega-Palas, M.A. (2016) Novel features of telomere biology revealed by the
absence of telomeric DNA methylation. *Genome Res*, **26**, 1047–1056.
10.1101/gr.202465.115.

Verhulst, S. (2020) Improving comparability between qPCR-based telomere studies. *Molecular*
Ecology Resources, **20**, 11–13. 10.1111/1755-0998.13114.

Wang, T., Hao, R., Pan, H., Cheng, T. and Zhang, Q. (2014) Selection of suitable reference
genes for quantitative real-time polymerase chain reaction in *Prunus mume* during
flowering stages and under different abiotic stress conditions. *J. Amer. Soc. Hort. Sci.*,
139, 113–122. 10.21273/JASHS.139.2.113.

Watson, J.M. and Riha, K. (2011) Telomeres, aging, and plants: From weeds to Methuselah – a
mini-review. *GER*, **57**, 129–136. 10.1159/000310174.

Wickson, E.J. (1914) *The California fruits and how to grow them*, San Francisco, CA: Pacific
rural Press. https://lccn.loc.gov/14011514.

Zachová, D., Fojtová, M., Dvořáčková, M., Mozgová, I., Lermontova, I., Peška, V.,
Schubert, I., Fajkus, J. and Sýkorová, E. (2013) Structure-function relationships
during transgenic telomerase expression in *Arabidopsis*. *Physiol Plantarum*, **149**, 114–
126. 10.1111/ppl.12021.

Zangi, M., Bagherieh Najjar, M.B., Golalipour, M. and Aghdasi, M. (2019) met1 DNA
methyltransferase controls TERT gene expression: A new insight to the role of
telomerase in development. *CellJ*, **22**. 10.22074/cellj.2020.6290.

Tables

SRA Biosample	Accession Name	Cultivar	Age (years)
SRR4045222	DPRU 1207.2	N/A	29
SRR4045223	DPRU 210	'Languedoc'	32
SRR4045224	DPRU 2331.9	N/A	20
SRR4045225	DPRU 1791.3	N/A	23
SRR4045226	DPRU 2301	'Tuono'	21
SRR4045227	DPRU 2374.12	N/A	19
SRR4045228	DPRU 1456.4	'Badam'	28
SRR4045229	DPRU 1462.2	N/A	28
SRR4036105	DPRU 2578.2	N/A	15

Table 1. NCBI SRA biosample entry for almond accessions used for *in silico* mean telomere length estimation.

Sampling Year	Age (years)	Sample Size
2018	1	3
2018	5	3
2018	9	3
2018	14	3
2019	2	3
2019	7	3
2019	11	3

Table 2. Sampling scheme for 2018 and 2019 almond age cohort collections.

Oligo Name	Oligo Sequence (5' – 3')
PP2A Forward	**CGGCGGCGGGCGGCGCGGG**CAGGATAGACATTGGAGGGTTCGGCTCGCAA
PP2A Reverse	**CGGCGGCGGGCGGCGCGGG**ACCACTGCATGCAAAGGGACCCAAGCTTAT
Telomere Forward	CCCCGGTTTTGGGTTTTGGGTTTTGGGTTTTGGGT
Telomere Reverse	GGGGCCCTAATCCCTAATCCCTAATCCCTAATCCCT
TERT Forward	GCATCAGAGAAGGGTCAGATT
TERT Reverse	CTCTGGCTCCTTGAATCGTATAG
RPII Forward	TGAAGCATACACCTATGATGATGAAG
RPII Reverse	CTTTGACAGCACCAGTAGATTCC

591   **Table 3.** Oligos used for all MMQPCR and qRT-PCR studies.

Figure Legends

Figure 1. Image of almond cultivar 'Nonpareil' (photo taken by K. D'Amico-Willman in May 2018).

Figure 2. Linear regression showing the relationship between $\sqrt{mean\ telomere\ length}$ and accession age in nine almond accessions. Dashed line represents the best-fit linear model (p-value = 0.1458; R^2 = 0.2767).

Figure 3. Boxplots depicting the calculated z-score of the T/S ratio for almond samples within the age cohorts tested. **(a)** Age cohort collected in 2018. **(b)** Age cohort collected in 2019. Significant differences in z-scores between age cohorts based on ANOVA followed by *post hoc* Fisher's LSD (α = 0.1) are denoted by letter groupings (ANOVA 2018 p-value = 0.1077; ANOVA 2019 p-value = 0.06548).

Figure 4. Normalized expression of *TERT* for almond samples within the age cohorts test. **(a)** Age cohort collected in 2018. **(b)** Age cohort collected in 2019. Significant differences in relative expression between age cohorts based on ANOVA followed by *post hoc* Tukey's HSD (alpha = 0.1) are denoted by the letter groupings (ANOVA 2018 p-value = 0.09087; ANOVA 2019 p-value = 0.1414).

Acknowledgements

We would like to acknowledge Matthew Willman for his assistance with the statistical analyses and Cheri Nemes for her assistance with wet lab portions of this project. This work is supported by the Ohio State University CFAES-SEEDS program 2018113, the Translational Plant Sciences Graduate Fellowship, the AFRI-EWD Predoctoral Fellowship 2019-67011-29558 from the USDA National Institute of Food and Agriculture, and the Ohio Supercomputer Center.

Conflict of Interest Statement

The authors declare no conflicts of interest.

622   **Figures**

647   **Figure 1**

$y = 38 - 0.35\,x$

650   **Figure 2**

Figure 3

Figure 4