FisB relies on homo-oligomerization and lipid-binding to catalyze membrane fission in bacteria

Ane Landajuela[1,2†*], Martha Braun[2,3†], Christopher D. A. Rodrigues[4], Alejandro Martínez-Calvo[5], Thierry Doan[6], Florian Horenkamp[7#], Anna Andronicos[1‡], Vladimir Shteyn[1,2,] Nathan D. Williams[2,7], Chenxiang Lin[2,7], Ned S. Wingreen[8,9], David Z. Rudner[10], Erdem Karatekin[1,2,3,11*]

Cellular and Molecular Physiology, Yale University, New Haven, CT, USA

Nanobiology Institute, Yale University, West Haven, CT, USA

Molecular Biophysics and Biochemistry, Yale University, New Haven, CT, USA

iThree Institute, University of Technology Sydney (UTS), Australia

Grupo de Mecánica de Fluidos, Universidad Carlos III de Madrid, Spain

Laboratoire d'Ingénierie des Systèmes Macromoléculaires, Aix-Marseille Université - CNRS UMR7255, Marseilles, France

Cell Biology, Yale University

Department of Molecular Biology, Princeton University, Princeton, NJ 08544, USA.

Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ 08544, USA.

Department of Microbiology, Harvard Medical School, Boston MA

Université de Paris, SPPIN - Saints-Pères Paris Institute for the Neurosciences, Centre National de la Recherche Scientifique (CNRS), F-75006 Paris, France.

† These authors contributed equally

Present address: Pfizer, New London, CT

‡ Present address: Department of Biological Chemistry, School of Medicine, U. California Irvine

* To whom correspondence should be addressed

ane.landajuela@yale.edu

erdem.karatekin@yale.edu (lead contact)

ABSTRACT

Little is known about mechanisms of membrane fission in bacteria despite their
requirement for cytokinesis. The only known dedicated membrane fission machinery in
bacteria, FisB, is expressed during sporulation in *Bacillus subtilis* and is required to
release the developing spore into the mother cell cytoplasm. Here we characterized the
requirements for FisB-mediated membrane fission. FisB forms mobile clusters of ~12
molecules that give way to an immobile cluster at the engulfment pole containing ~40
proteins at the time of membrane fission. Analysis of FisB mutants revealed that binding
to acidic lipids and homo-oligomerization are both critical for targeting FisB to the
engulfment pole and membrane fission. Experiments using artificial membranes and
filamentous cells suggest FisB does not have an intrinsic ability to sense or induce
membrane curvature but can bridge membranes. Finally, modeling suggests homo-
oligomerization and trans interactions with membranes are sufficient to explain FisB
accumulation at the membrane neck that connects the engulfment membrane to the
rest of the mother cell membrane during late stages of engulfment. Together, our results
show that FisB is a robust and unusual membrane fission protein that relies on homo-
oligomerization, lipid-binding and the unique membrane topology generated during
engulfment for localization and membrane scission, but surprisingly, not on lipid
microdomains, negative-curvature lipids, or curvature-sensing.

INTRODUCTION

Membrane fission is a fundamental process required for endocytosis[1], membrane trafficking[2], enveloped virus budding[3], phagocytosis[4], cell division[5] and sporulation[6-8]. During membrane fission, an initially continuous membrane divides into two separate ones. This process requires dynamic localization of specialized proteins, which generate the work required to merge membranes[9-13]. Dynamin[14] and the endosomal sorting complex required for transport III (ESCRT-III) catalyze many eukaryotic membrane fission reactions[15]. Both fission machineries bind acidic lipids, assemble into oligomers, and use hydrolysis of a nucleoside triphosphate (GTP or ATP) to achieve membrane fission. However, membrane fission can also be achieved by friction[16], stress accumulated at a boundary between lipid domains[17], forces generated by the acto-myosin network[18-21] or protein crowding[22]. By contrast, very little is known about membrane fission in bacteria, even though they rely on membrane fission for every division cycle.

We previously found that fission protein B (FisB) is required for the final membrane fission event during sporulation in *B. subtilis*[23]. When nutrients are scarce, bacteria in the orders *Bacillales* and *Clostridiales* initiate a developmental program that results in the production of highly resistant endospores[24]. Sporulation starts with an asymmetric cell division that generates a larger mother cell and a smaller forespore (Figure 1A). The mother cell membranes then engulf the forespore in a process similar to phagocytosis. At the end of engulfment, the leading membrane edge forms a small pore. Fission of this membrane neck connecting the engulfment membrane to the rest of the mother cell membrane releases the forespore, now surrounded by two membranes, into the mother cell cytoplasm (Figure 1A,B). At this late stage the mother nurtures the forespore as it prepares for dormancy. Once mature, the mother cell releases the spore into the environment through lysis. Spores can withstand heat, radiation, drought, antibiotics, and other environmental assaults for many years[25-28]. Under favorable conditions, the spore will germinate and restart the vegetative life cycle.

Conserved among endospore-forming bacteria, FisB is a mother-cell transmembrane protein expressed under the control of the transcription factor, σ^E, after asymmetric division[29]. In sporulating cells lacking FisB, engulfment proceeds normally but the final membrane fission event, detected using a lipophilic dye, is impaired[23] (Figure 1C,F and S1 Appendix Fig. 1A). During engulfment, FisB fused to a fluorescent protein forms dim, mobile clusters in the engulfment membrane (Figure 1D,E, Movie 1). When the engulfing membranes reach the cell pole, approximately 3 hours (t = 3h) after the onset of sporulation, a cluster of FisB molecules accumulates at the pole forming a more intense, immobile focus, where and when fission occurs (Figure 1D,E, Movie 2).

We had previously reported[23] that FisB interacts with cardiolipin (CL), a lipid enriched at cell poles[30-32] whose levels increase during sporulation[33] and is implicated in membrane fusion[34-36] and fission reactions[37]. In addition, CL was reported to act as a landmark for the polar recruitment of the proline transporter ProP, and the mechanosensitive channel MscSm[38,39]. Thus, it seemed plausible that CL could act as a landmark to recruit FisB to the membrane fission site and facilitate membrane fission. Apart from this hypothesis, no information has been available about how FisB localizes to the membrane fission site and how it may drive membrane scission.

Here, we determined the requirements for FisB's sub-cellular localization and
membrane fission during sporulation. Using quantitative analysis, we find small clusters
of ~12 FisB molecules diffuse around the mother cell membrane and ~40 copies of FisB
accumulate at the fission site as an immobile cluster to mediate membrane fission.
When FisB expression was lowered, ~6 copies of FisB were sufficient to drive
membrane fission, but fission took longer. Unexpectedly, FisB dynamics and membrane
fission are independent of both CL and phosphatidylethanolamine (PE), another lipid
implicated in membrane fusion and fission. We found FisB binds phosphatidylglycerol
(PG) with comparable affinity as CL, after adjusting for charge density. Thus, we
suspect that, as a more abundant lipid in the cell, PG can substitute for CL to bind FisB.
We tested other factors that may be important for the sub-cellular localization of FisB
and membrane fission. We found FisB dynamics are independent of flotillins, which
organize bacterial membranes into functional membrane microdomains[40], cell wall
synthesis machinery, and proton or voltage gradients across the membrane. Using
mutagenesis, we show that both FisB oligomerization and binding to acidic lipids are
required for proper localization and membrane fission. *B. subtilis ΔfisB* cells were
partially complemented by *C. perfringens* FisB, despite only ~23% identity between the
two proteins, suggesting a common localization and membrane fission mechanism
based on a few conserved biophysical properties. The membrane neck that eventually
undergoes fission and where FisB accumulates is the most highly curved membrane
region in the late stages of engulfment. Thus, FisB could potentially localize at the
membrane neck due to a preference for highly curved membrane regions. We tested
this possibility in experiments with both artificial giant unilamellar vesicles (GUVs) and
live cells. Surprisingly, these experiments failed to reveal any intrinsic affinity of FisB for
highly curved membranes. However, we found that FisB bridges membranes and
accumulates at membrane adhesion sites. Using modeling, we found that self-
oligomerization of FisB, coupled with its ability to bridge negatively charged membranes
is sufficient to explain its localization to the membrane neck. Thus, proteins can localize
to highly curved membrane regions through mechanisms independent of intrinsic
curvature sensitivity. Together, these results suggest FisB-FisB and FisB-lipid
interactions, combined with the unique membrane topology generated at the engulfment
pole during sporulation, provide a simple mechanism to recruit FisB to mediate
membrane fission independent of other factors.

RESULTS

Membrane fission occurs in the presence of a cluster of FisB molecules

To correlate FisB dynamics with membrane fission, we devised a labeling strategy that
allowed us to monitor both simultaneously, using a modified version of a fission assay
developed previously[41]. In this assay, synchronous sporulation is induced by placing *B.*
subtilis cells in a nutrient-poor medium. At different time points after the nutrient
downshift, aliquots are harvested from the culture, stained with the lipophilic membrane
dye FM4-64, mounted on an agar pad, and imaged using fluorescence microscopy. The
dye is virtually non-fluorescent in the medium, and it cannot cross the cell membrane.
Thus, before fission, FM4-64 labels the outer leaflet of both the mother cell and the
forespore membranes. After fission, only the outer leaflet of the mother cell is labeled

(S1 Appendix Figure 1B). Because post-fission cells and cells that never entered
sporulation are labeled identically, in addition to FM4-64, a fluorescent protein is
expressed in the forespore under the control of the forespore-specific transcription
factor σ^F to distinguish between the two cell types[42] (S1 Appendix Figure 1B). This
makes it challenging to monitor FisB dynamics simultaneously, which requires a third
channel. As an alternative, we used another lipophilic dye, TMA-DPH, that has partial
access to internal membranes but can distinguish between pre- and post-fission stages
without need for a forespore reporter[23] (Figure 1C and S1 Appendix Fig. 1D-G). Using
TMA-DPH as the fission reporter, we quantified the percentage of cells that have
undergone fission as a function of time, for wild-type, *fisB* knock-out (*ΔfisB*, strain
BDR1083, see S1 Appendix Table 2 for strains used), and *ΔfisB* cells complemented
with FisB fused to monomeric EGFP (mGFP-FisB, strain BAM003) as shown in Figure
1D and 1F. These kinetic measurements reproduced results obtained using FM4-64 (S1
Appendix Figure 1C). Thus, TMA-DPH can be used as a faithful reporter of membrane
fission, leaving a second channel for monitoring dynamics of FisB fused to a fluorescent
reporter.

In the experiments of Figure 1D and 1F, we simultaneously monitored dynamics of
mGFP-FisB and membrane fission. We found that membrane fission is almost always
accompanied by an intense, immobile mGFP-FisB signal at the engulfment pole (Figure
1D, time= 3hr into sporulation). This intense spot at the engulfment pole (ISEP) is
distinct from the dimmer, mobile clusters (DMC) that appear at earlier times elsewhere
(Figure 1D). By 3 h into sporulation, around 70 % of the cells expressing mGFP-FisB at
native levels had an ISEP (Figure 1G), a number that was close to the percentage of
cells that had undergone fission by then (Figure 1F). Scoring individual cells, we found
>90% (212/235) of cells that had undergone membrane fission also had an ISEP.

We also monitored membrane fission and mGFP-FisB signals in a strain with lower FisB
expression. Here, lower FisB expression is achieved by reducing the spacing between
the ribosome binding site (RBS) and the ATG start codon[43]. In this strain (BAL003),
there was an initial delay in the fraction of cells that had undergone fission, but fission
accelerated after t=3 h to reach near wild-type levels at around t=4h (Figure 1E,F). The
fraction of cells with an ISEP followed a similar pattern (Figure 1G). The fraction of cells
that had undergone fission at a given time was strongly correlated with the fraction of
cells with an ISEP at that time (Figure 1H). Scoring individual cells, we found >93%
(258/277) of cells that had undergone membrane fission had an ISEP. We conclude that
membrane fission occurs in the presence of a large immobile cluster of FisB molecules
at the site of fission.

About 40 FisB molecules accumulate at the engulfment pole to mediate
membrane fission

We asked how many copies of FisB are recruited to the engulfment pole at the time of
membrane fission and how this number is affected by the expression level. For this
quantification, we used DNA-origami based fluorescence standards we recently
developed[44]. These standards consist of DNA rods (~410 nm long and 7 nm wide)
labeled with AF647 at both ends and a controlled number of mEGFP molecules along

the rod (Figure 2A).

DNA-origami standards carrying different mEGFP copies were imaged using widefield
fluorescence microscopy (Figure 2B). For each type of rod, the average total
fluorescence intensity of single-rods was computed and plotted against the number of
mEGFP molecules per rod, generating the calibration curve in Figure 2D. We generated
B. subtilis cells expressing mEGFP-FisB at native levels (BAL001) in a *ΔfisB*
background so that images of these cells obtained under identical imaging conditions as
for the calibration curve in Figure 2D could be used to compute mEGFP-FisB copy
numbers. We imaged mEGFP-FisB cells at t=3 h after sporulation was induced. From
these same images, we estimated the total fluorescence of dim, mobile clusters (DMC)
and ISEP in *B. subtilis* cells as a sum of background-corrected pixel values (Figure 2C).
Using the average values of these total intensities, we estimate ~40 copies at the ISEP,
and ~12 per DMC from the calibration in Figure 2D. From the total intensity of cells (S1
Appendix Fig. 2E), we also estimate there are ~1000 FisB molecules per cell. Two
independent estimates, based on *B. subtilis* calibration strains[45] and quantitative
immunoblotting, resulted in slightly larger and smaller estimates of these copy numbers,
respectively (S1 Appendix and S1 Appendix Figs. 2, 3).

We tracked the DMC to estimate how rapidly they move. From the tracks, we calculated
the mean squared displacement (MSD) as a function of time (Figure 2E). The short-time
diffusion coefficient estimated from the MSD is $D_{DMC} \approx 2.8 \times 10^3 \ \mathrm{nm}^2/\mathrm{s}$ (95%
confidence interval CI=$2.76 - 2.85 \times 10^3 \ \mathrm{nm}^2/\mathrm{s}$). This value is comparable to the
diffusivity of FloA and FloT clusters of ~100 nm with $D \approx 6.9 \times 10^3$ and $4.1 \times 10^3 \ \mu\mathrm{m}^2/\mathrm{s}$,
respectively[46]. By comparison, ISEP have $D_{ISEP} \approx 28 \ \mathrm{nm}^2/\mathrm{s}$ (CI=$22.9 - 33.1 \ \mathrm{nm}^2/\mathrm{s}$), two
orders of magnitude smaller.

We performed similar estimations of FisB copy numbers for the low expression strain
(BAL004) (S1 Appendix Fig. 4). We found ~160±66, 122±51, or 83±6 (±SD) copies per
cell using *B. subtilis* standards, DNA-origami, or the quantitative WB methods,
respectively. For the ISEP, we found 8±2, 6±2, or 5±3 (±SD) copies of mGFP-FisB
using the three approaches, respectively (S1 Appendix Table 1). About 6 % of the total
mGFP-FisB signal accumulated in ISEP, close to the ~4% in the native-expression
strain (S1 Appendix Fig. 4E). The DMC were too dim to quantify reliably. Assuming
DMCs to be ~3-fold dimmer than ISEP like in the native-expression strain, each DMC
would contain 2-3 mGFP-FisB, just below our detection limit. Interestingly, lowering the
total expression of FisB per cell ~8-fold resulted in a ~6-fold reduction in the average
number of FisB molecules found at the membrane fission site. Thus, only ~6 copies of
FisB are sufficient to mediate membrane fission, but only after some delay (Figure
1E,F).

In summary, ~40 FisB molecules accumulate at the fission site to mediate membrane
fission. Only 3-4 DMCs need to reach the fission site to provide the necessary numbers.
When FisB expression is lowered ~8-fold, ~6 FisB molecules accumulate at the
engulfment pole to mediate membrane fission, but fission takes longer.

FisB localization and membrane fission are independent of cardiolipin,
phosphatidylethanolamine and flotillins

To investigate how FisB is recruited to the membrane fission site, we began by testing a
potential role for the cell wall remodeling machinery, the protonmotive force, and the
membrane potential, and found none influenced FisB dynamics (S1 Appendix Results
and S1 Appendix Fig. 6).

We then tested whether lipid microdomains play a role in recruitment to the site of
fission. Previously, we reported that the recombinant, purified extracytoplasmic domain
(ECD, see Figure 4A) of FisB interacts with artificial lipid bilayers containing CL[23]. To
test if FisB-CL interactions could be important for the subcellular localization of FisB and
membrane fission, we generated a strain (BAM234) that carries deletions of the three
known CL synthase genes *ywnE (clsA)*, *ywjE (clsB)* and *ywiE (clsC)*[47] (Figure 3A). The
CL synthase-deficient strain did not contain detectable levels of CL at t=3 hours after
sporulation was initiated (Figure 3B). CL-deficient cells grew normally but had a
reduction in sporulation efficiency as assayed by heat-resistant (20 min at 80˚C) colony
forming units (S1 Appendix Table 2 and S1 Appendix Figure 5)[33]. A reduction in
sporulation efficiency measured in this manner can be due to a defect at one or several
steps during sporulation or germination. Importantly, the membrane fission time course
of *ΔclsABC* cells was indistinguishable from those of wild-type cells (Figure 3C,D),
indicating the defect in sporulation occurs at a stage after membrane fission. In addition,
mYFP-FisB localization and dynamics were similar in *ΔclsABC* (BAL037) and wild-type
(BAL002) cells (Figure 3F-H). The fraction of cells that had an ISEP, and the intensity of
the ISEP, reflecting the number of FisB molecules recruited to the membrane fission
site, were indistinguishable for wild-type and *ΔclsABC* cells (Figure 3G,H). We conclude
that CL is not required for the subcellular localization of FisB or membrane fission.

Next, we tested a potential role for phosphatidylethanolamine (PE), another lipid
implicated in membrane fusion and fission[48,49] and that forms microdomains[50]. We
deleted the *pssA* gene which encodes phosphatidylserine synthase that mediates the
first step in PE synthesis (Figure 3A) to generate cells lacking PE (strain BAL031,
Figure 3B). Kinetics of membrane fission during sporulation were identical in *ΔpssA* and
wild-type cells (Figure 3D), indicating PE does not play a significant role in membrane
fission.

PE and CL domains in *B. subtilis* membranes tend to occur in the same sub-cellular
regions[50], raising the possibility that CL and PE may compensate for each other. To test
whether removing both CL and PE affects fission, we generated a quadruple mutant
(BAL030) lacking both CL and PE (Figure 3B), leaving PG as the major phospholipid
component of the membrane. Surprisingly, the quadruple mutant underwent fission with
indistinguishable kinetics compared to wild-type (Figure 3C,D). Thus, two lipids with
negative spontaneous curvature and implicated in membrane fusion and fission
reactions in diverse contexts have no significant role in membrane fission mediated by
FisB during sporulation.

In addition to CL and PE microdomains, bacteria also organize many signal
transduction cascades and protein-protein interactions into functional membrane
microdomains (FMMs), loose analogs of lipid rafts found in eukaryotic cells[40]. The
FMMs of *B. subtilis* are enriched in polyisoprenoid lipids and contain flotillin-like
proteins, FloT and FloA, that form mobile foci in the plasma membrane[51,52]. FloT-
deficient cells have a sporulation defect, but which sporulation stage is impaired is not

known[46]. We observed that *ΔfloA* (BAL035)*,* but not *ΔfloT* (BAL036), cells are impaired
in sporulation as assayed by heat-resistant colony forming units (S1 Appendix Table 2,
and S1 Appendix Figure 5). However, when we monitored engulfment and membrane
fission, we found both proceeded normally in *ΔfloA* cells (Figure 3D). Thus, the
sporulation defect in *ΔfloA* cells lies downstream of engulfment and membrane fission.
This was confirmed by blocking formation of FMMs during sporulation by addition of 50
μM zaragozic acid[53] to the sporulation medium which had no effect on the localization of
mGPF-FisB (Figure 3E).

Together, these results imply that FisB-mediated membrane fission that marks the end
of engulfment during sporulation is insensitive to the negative-curvature lipids CL, PE,
and to FloA/T-dependent lipid domains.

FisB binds to acidic lipids

PG can substitute for CL as a binding partner for many proteins[54,55]. To see if this might
also be the case for the FisB ECD, we quantified the affinity of this domain for both
lipids.

Most, but not all, algorithms (S1 Appendix Fig. 7) predict FisB to possess a single
transmembrane domain (TMD) with a small N-terminal cytoplasmic domain and a larger
(23-kDa) ECD, as depicted in Figure 4A. We first confirmed this predicted topology
using a cysteine accessibility assay[56] (S1 Appendix Fig. 8, Materials and Methods, and
S1 Appendix Results). Our attempts to determine the structure of recombinant, purified
FisB ECD were unsuccessful, but a computational model of FisB for residues 44 to 225,
covering most of the ECD is available[57] and is shown in Figure 4B. The model predicts
a curved ECD structure, with ~3 nm and ~5 nm for the inner and outer radii of
curvatures. The overall topology of FisB, with the predicted ECD structure is depicted in
Figure 4B.

We probed interactions of FisB ECD with PG using a liposome co-flotation assay,
illustrated in Figure 4C. Purified recombinant, soluble FisB ECD (Fig. 4A, bottom) was
incubated with liposomes and subsequently layered at the bottom of a discontinuous
density gradient. Upon equilibrium ultracentrifugation, the lighter liposomes float up to
the interface between the two lowest density layers together with bound protein, while
unbound protein remains at the bottom of the gradient. We collected fractions and
determined the percentage of protein co-floated with liposomes using SDS-PAGE and
densitometry, as shown in Figure 4D. We first determined that binding of FisB ECD to
liposomes containing CL was not dependent on pH or the divalent ion Ca^{2+} (S1
Appendix Fig. 9F,G). By contrast, the fraction of liposome-bound protein decreased
rapidly as the ionic strength increased (S1 Appendix Fig. 9H). These results indicated
binding was mainly electrostatic in nature.

At neutral pH, CL carries two negative charges, whereas PG and phosphatidylserine
(PS), a lipid not normally found in *B. subtilis*[58], carry only a single negative charge. If
binding is mediated mainly by electrostatic interactions, then liposomes carrying PG or
PS at two times the mole fraction of CL should bind nearly the same amount of FisB
ECD, since the surface charge density would be the same. Indeed, similar amounts of

FisB ECD were bound to liposomes carrying 30% CL, 60% PG, or 60% PS (Figure 4E).
FisB ECD did not bind neutral phosphatidylcholine PC liposomes[23].

To quantify the affinity of recombinant soluble FisB ECD for CL vs PG, we then titrated
liposomes containing 45 mole % CL or PG and measured binding of 100 nM FisB ECD
(Figure 4F). In these experiments, we used iFluor555 labeled FisB ECD (iFluor555-FisB
ECD) and detected liposome-bound protein using fluorescence rather than densitometry
of SYPRO-stained gels, which extended sensitivity to much lower protein
concentrations. The titration data were fit to a model to estimate the apparent
dissociation constant, K_d (see Materials and Methods), which was 1.0 µM for CL (95%
confidence interval CI=0.7-2.1 µM) and 3.6 µM for PG, respectively (CI=2.8-5.0, Figure
4F,G).

Together, these results suggest that while FisB has higher affinity for CL than for PG,
the higher affinity results mainly from the higher charge carried by CL. FisB does not
bind CL with much specificity; at the same surface charge density, FisB ECD binds PG,
or even PS which is not a *B. subtilis* lipid, with similar affinity. Thus, *in vivo* FisB is likely
to bind CL as well as PG which is much more abundant.

Purified FisB ECD forms soluble oligomers

FisB forms clusters of various sizes in cells as described above (Figure 1, 2) and does
not appear to have other protein interaction partners[23]. Thus, homo-oligomerization of
FisB may be important for its function. We explored oligomerization of recombinant,
soluble FisB ECD (Figure 5). When FisB ECD bearing a hexa-histidine tag is expressed
in *E. coli* and purified to homogeneity by affinity chromatography, samples analyzed by
SDS-PAGE show multiple bands corresponding to different oligomeric states (Figure 5D
and S1 Appendix Figure 9B). Size-exclusion chromatography (SEC) analysis resolved
the purified protein into predominant high molecular weight oligomeric structures eluting
over a wide range of sizes, and low molecular weight peaks comprising minor
components (Figure 5E and S1 Appendix Figure 9C, top). The minor peak at ~23 kDa
(18 ml elution volume) corresponds to monomeric FisB ECD, whereas the peak at ~400
kDa (15 ml) is FisB ECD that co-elutes with another protein, likely the 60 kDa
chaperone GroEL, a common contaminant in recombinant proteins purified from *E. coli*
(S1 Appendix Figure 9D). To rule out potential artefacts caused by the hexa-histidine
affinity tag, we also purified FisB ECD using a GST-tag, which yielded similar results.
The SEC of high molecular weight peaks collected from the initial chromatogram did not
show a redistribution when re-analyzed (S1 Appendix Figure 9C, bottom), suggesting
that once formed, the oligomeric structures are stable for an hour or longer.

We analyzed the high molecular-weight SEC fractions (peaks 1 and 2) using electron
microscopy (EM) after negative staining. This analysis revealed rod-like structures quite
homogeneous in size, ~50 nm long and ~10 nm wide (Figure 5F and S1 Appendix
Figure 9E). These structures displayed conformational flexibility, which precluded
structural analysis using cryoEM (and likely hampered our attempts to crystallize FisB
ECD). We estimate every rod-like oligomer can accommodate ~40 copies of the
predicted structure of FisB[44-225] shown in Figure 4B, similar to the number of FisB
molecules recruited to the membrane fission site in cells (Figure 2).

A FisB mutant that is selectively impaired in homo-oligomerization

To determine whether self-oligomerization and lipid-binding interactions are important for FisB's function, we generated a series of mutants, characterized oligomerization and lipid-binding of the mutant proteins *in vitro*, and analyzed FisB localization dynamics and membrane fission during sporulation *in vivo*.

We suspected self-oligomerization of FisB was at least partially due to hydrophobic interactions. Accordingly, we first mutated conserved residues G175, I176, I195 and I196 in a highly hydrophobic region of FisB ECD (Figure 5A,B), producing a quadruple mutant, G175A,I176S, I195T, I196S (FisB[GIII]). These residues are on the surface of the predicted structure of FisB ECD (Figure 5C), so are not expected to interfere with folding. Purified FisB[GIII] ECD displayed reduced oligomerization when analyzed using SDS PAGE or size exclusion chromatography (Figure 5D,E). Though much reduced in amplitude, a broad, high molecular weight peak was still present in size exclusion chromatograms (Figure 5E). Negative-stain EM analysis of this fraction revealed oligomerization with less defined size and structure compared to wild type FisB ECD (Figure 5G).

To test whether lipid binding of the GIII mutant was affected, we used the co-flotation assay described above, except only two fractions were collected (Figure 5H,I). This analysis revealed that, despite being impaired in self-oligomerization, FisB[GIII] ECD has lipid binding properties similar to wild-type with a dissociation constant $K_d^{GIII} = 1.6$ µM (95% confidence interval CI=0.9-5.1 µM), indistinguishable from that of wild type FisB ECD[WT] ($K_d^{wt} = 1.0$ µM, CI = 0.7 − 2.1 µM, Figure 5J).

FisB[K168D,K170E] (FisB[KK]) is selectively impaired in binding acidic lipids

To engineer lipid-binding mutants, we took advantage of our observation that FisB binding to anionic lipids is principally mediated through electrostatic interactions (S1 Appendix Figure 9H). We generated a series of mutants in which we either neutralized or inverted up to four charges (S1 Appendix Fig. 11 and S1 Appendix Table 2). The ECD of a set of charge neutralization mutants were expressed in *E. coli*, purified and tested for lipid binding using the liposome co-floatation assay. The largest reductions in lipid binding were observed when lysines in a region comprising residues 168-172 were neutralized (S1 Appendix Fig. 11A). This region corresponds to a highly positively charged pocket in the predicted model of FisB 44-225 (Figure 5C).

A partially overlapping set of FisB mutants were expressed in a *ΔfisB* background and tested for sporulation efficiency by monitoring formation of heat-resistant colonies (S1 Appendix Fig. 11B-E). Again, the strongest reductions in sporulation efficiency were found when lysines 168, 170 or 172 were mutated (S1 Appendix Fig. 11D). We decided to characterize the K168D, K170E mutation in more detail, as it produced the strongest reduction in sporulation efficiency.

We purified the ECD of FisB[K168D,K170E] (FisB[KK]) from *E. coli* and tested its binding to liposomes containing 45 mole % CL using the co-floatation assay (Figure 5H-J). The

404   dissociation constant for FisB^{KK}-acidic lipid binding was $K_d^{KK} = 9.1$ μM (CI=6.5-15.3
405   μM), nearly 10-fold lower than that for wild-type FisB ECD ($K_d^{wt} = 1.0$ μM, CI $= 0.7 -$
406   2.1 μM, Figure 5I,J). Importantly, formation of oligomers was not affected (Figure 5D,E).
407   Thus, FisB^{KK} is specifically impaired in binding to acidic lipids.

409 FisB-lipid interactions and homo-oligomerization are important for targeting FisB
410 to the fission site

Using the FisB mutants selectively impaired in binding to lipids or homo-oligomerization,
we investigated whether these activities are important for FisB's function *in vivo*. To
analyze FisB clustering and targeting to the fission site, we fused wild-type FisB or the
two mutants to an N-terminal monomeric YFP (mYFP) and expressed the fusions at
lower levels, which facilitated observation of ISEPs (Figure 6A). We induced these
strains to sporulate and monitored FisB dynamics and membrane fission. Both the lipid-
binding (FisB^{KK}) and the oligomerization mutant (FisB^{GIII}) were targeted to the cell
membrane, unlike many other mutants we tested (S1 Appendix Figure 11E and S1
Appendix Table 2). At t=1.5 h after the nutrient downshift, mYFP-FisB signals were
visible in all strains without any distinguishing features. At t=2.5 h, a subset of cells
expressing the wild-type FisB fusion had undergone membrane fission and these cells
had an ISEP. By contrast, membrane fission was not evident in either of the mutants.
By t=3 h, 25% of WT FisB cells had undergone fission, nearly always with an
accompanying ISEP. In the lipid binding FisB^{KK} mutant, only 8% of the sporulating cells
had accomplished membrane fission (Figure 6B), but more than 90% of those that did
had an ISEP (53/58 cells). Membrane fission events and the accompanying bright
mYFP-FisB spots were very rare (0.6%) in the oligomerization-deficient FisB^{GIII} mutant.

The distribution of fluorescence intensities of the foci from low-expression WT and KK
cells were indistinguishable (Figure 6C). Using the DNA-origami fluorescence intensity
calibration (Figure 2), we estimate 6±2 copies of low-expression FisB WT or the KK
mutant to have accumulated at the fission site. For the GIII mutant, there were not
enough cells with an intense spot to perform a similar analysis.

From TMA-DPH labeling, we determined the fraction of cells that successfully
completed fission as a function of time (Figure 6D). Oligomerization-deficient FisB^{GIII}
was not able to induce fission, whereas the lipid-binding mutant FisB^{KK} had a partial, but
severe defect (~50% reduction compared to wild-type). Importantly, both mutants were
expressed at levels similar to the wild-type (S1 Appendix Fig. 10), so the defects to form
an ISEP and undergo membrane fission are not due to lower expression levels.

Together, these results suggest FisB-lipid and FisB-FisB interactions are both important
for targeting FisB to the fission site.

442 *C. perfringens* FisB can substitute for *B. subtilis* FisB

So far, our results suggest FisB-FisB and FisB-acidic lipid interactions are the main
drivers for targeting FisB to the membrane fission site. If no other partners are involved,
FisB should be largely an independent fission module, i.e. FisB homologs from different

sporulating bacteria should be able to substitute for one another at least partially, even if
sequence homology is low outside the consensus region. To test this idea, we
expressed *Clostridium perfringens* FisB (FisB^{Cperf}) in *B. subtilis* cells lacking FisB
(BAL005). The sequence identity is only 23% between FisB sequences from these two
species. In the heat-kill assay, FisB^{Cperf} fully rescued *B. subtilis ΔfisB* defects (Figure
7A). *C. perfringens* FisB fused to mEGFP (mEGFP-FisB^{Cperf}) had similar dynamics as
FisB^{Bsubti}, forming DMCs at early times that gave way to an ISEP where membrane
fission occurs (Figure 7B). Population kinetics of membrane fission were slower with
FisB^{Cperf} (Figure 7C), but nearly every cell that underwent fission had an ISEP as for the
wild type protein (220/239, or 92%). The intensity distribution of mEGFP-FisB^{Cperf} ISEP
was shifted to smaller values compared to mEGFP-FisB^{Bsubti} ISEP (Figure 7D). Since
the average ISEP intensity for FisB^{Bsubti} corresponds to ~40 copies (Figure 2), we
deduce ~9 copies of FisB^{Cperf} accumulate at ISEP at the time of membrane fission. At
t=3 h into sporulation, the percentage of cells with an ISEP was lower for cells
expressing mEGFP-FisB^{Cperf} (Figure 7E).

In all conditions tested so far, nearly all cells that had undergone membrane fission also
had an intense FisB spot at the engulfment pole (Figs. 2,3,6, and 7). When we plotted
the percentage of cells having an ISEP against the percentage of cells that have
undergone fission at t=3 h, we found a nearly perfect correlation (Figure 7F). FisB^{Cperf} fit
this pattern well, despite having a low sequence identity to FisB^{Bsubti}, suggesting a
common localization and membrane fission mechanism, likely based on a few
conserved biophysical properties.

FisB does not have a preference for highly curved membrane regions, but can bridge membranes

A number of proteins localize to sub-cellular sites due to their preference for curved
membrane regions[59-62]. During late stages of engulfment, the most highly curved region
in the cell is the membrane neck connecting the engulfment membrane with the rest of
the mother cell membrane and this is where FisB accumulates. We therefore asked
whether curvature-sensing could be a mechanism driving FisB's localization. To test this
possibility, we undertook three independent series of experiments.

First, we used the principle that any protein which preferentially binds curved
membranes at low membrane coverage can also induce membrane curvature when
present at sufficiently high coverage[60,63]. Thus, we tested whether the soluble ECD of
FisB could generate curved regions in highly malleable membranes of giant unilamellar
vesicles (GUVs) at high coverage. We incubated 2 µM purified soluble FisB ECD
labeled with iFluor555 with GUVs and monitored protein coverage and membrane
deformations using spinning-disc confocal microscopy. Even when the GUV
membranes were covered uniformly with iFluor555-FisB ECD we could not observe any
GUV membrane deformations (Figure 8A). As a positive control, we used purified
Endophilin A1 (EndoA1, labeled with Atto395), an N-BAR domain containing endocytic
protein[64-66]. We incubated 2 µM EndoA1 with GUVs composed of 45% DOPS, 24.5%
DOPC, 30% DOPE and 0.5% DiD, which resulted in extensive tubulation of GUV
membranes (Figure 8A), as reported previously[67]. Importantly, the difference in the

membrane sculpting ability of the two proteins is not due a weaker affinity of FisB ECD
for membranes ($K_d \approx 1$ µM for membranes with 45 mole % CL, Figure 4F) compared to
endophilin ($K_d = 1.15$ µM for membranes containing 45% DOPS, 30% DOPE, 24.5%
DOPC, 0.5% TR-DHPE[65]).

Second, we slowly deflated GUVs to facilitate any potential membrane curvature
generation by FisB ECD (which works against membrane tension) and/or to provide
curved regions to test if FisB ECD accumulated there. Deflated GUVs displayed curved
regions because their larger surface-to-volume ratios no longer allowed spherical
shapes. Even under these favorable conditions, FisB ECD was not able to generate
highly curved regions on these deflated GUVs (Figure 8B). In addition, if FisB ECD had
a preference for negatively (positively) curved regions, it should accumulate at such
regions while being depleted from positively (negatively) curved areas. Quantification of
FisB ECD coverage at negatively or positively curved membrane regions showed no
curvature preference (Figure 8B).

Third, we tested if FisB's localization in live *B. subtilis* cells depended on membrane
curvature. To avoid potentially confounding effects of other cues that may be present
during sporulation, we expressed GFP-FisB under an inducible promoter during
vegetative growth. In addition, we blocked cell division by inducing expression of MciZ[68].
MciZ normally blocks binary cell division during sporulation, but when expressed during
vegetative growth, cells grow into long flexible filaments that are bent to varying
degrees, providing regions with different membrane curvatures. We imaged GFP-FisB
spots along curved edges of these filaments and plotted the linear density of GFP-FisB
spots (spots/µm) as a function of filament curvature (Figure 8C). There was no clear
correlation between GFP-FisB spot density and filament curvature. Although this
method generates a limited amount of curvature, a similar approach was previously
used to show that DivIVA preferentially localizes to negatively curved regions[69].

In the GUV experiments, we noticed that FisB ECD caused GUVs to adhere to one
another when they came into contact, accumulating at the adhesion patch between the
membranes and at the rims (Figure 8D). Absorbance measurements using small
unilamellar vesicles (SUVs) confirmed that FisB ECD can bridge membranes and
aggregate liposomes (Figure 8E).

Overall, these experiments suggest that FisB does not have any intrinsic membrane
curvature sensing/sculpting ability, but it can bridge membranes.

Modeling suggests self-oligomerization and membrane bridging are sufficient to localize FisB to the membrane neck

To test the hypothesis that the homo-oligomerization, lipid-binding and the unique
membrane topology could be sufficient to recruit FisB to the membrane neck, we
considered a minimal model based on free energy minimization. As depicted in Figure
9, we consider the free energy F of an axisymmetric membrane neck of radius R and
length L connecting two membrane sheets, corresponding to the local geometry where
the engulfment membrane meets the rest of the mother cell membrane. We assume the
surface density ϕ of FisB proteins in the neck is uniform and reaches equilibrium with a

surface density φ_0 of FisB in the surrounding membranes, and ask whether the neck
geometry alone is enough to account for the observed FisB accumulation. The energy
functional consists of a term accounting for membrane bending and tension, F_m, and
another term accounting for FisB protein-protein interactions, F_p. We employ the
classical Helfrich-Canham theory[70-75] for the energy of the membrane

$F_m = \int_{S_n} dS_n \left[\frac{\kappa}{2} H^2 + \gamma \right] + \int_{S_s} dS_s \gamma,$ (1)

where S_n and S_s are the surfaces of the membrane neck and sheets, H is twice the
mean curvature, κ is the bending modulus, and γ is the surface tension. The two
membrane sheets are assumed to be planar, thus their only contribution to the energy
comes from membrane tension.

For FisB, we include translational entropy, the energy of homo-oligomerization in trans
between opposing membranes, and an energy that limits crowding. As shown above,
FisB proteins do not exhibit curvature sensing, so we do not include a term coupling
FisB density to membrane curvature in Eq. (1). This results in the following expression
for the protein free energy

$F_p = \int_{S_n} dS_n \left\{ k_B T \phi \ln \left(\frac{\phi}{\phi_0} \right) + a V_{LJ}(R) \phi^2 + U(\phi) \right\},$ (2)

where the first term accounts for translational entropy, the second term is an energy per
unit area accounting for trans interactions of FisB, and which for simplicity is assumed
to be proportional to the standard Lennard–Jones (LJ) potential accounting for a longer-
range attraction and shorter-range repulsion,

$V_{LJ}(r) = \left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^{6},$ (3)

and finally $U(\phi)$ is an energy penalty for crowding that increases rapidly above a certain
FisB concentration. To obtain $U(\phi)$ we assume a purely repulsive, truncated and shifted
LJ potential between cis-neighboring FisB molecules, which we take to occupy a
triangular lattice. Therefore, $U(\phi) = \epsilon \left[V_{LJ}(r(\phi)) - V_{LJ}(r_{max}) \right]$ when $r \le r_{max}$ and 0 when
$r > r_{max}$, where we have chosen $r_{max} = 2^{1/6} \sigma_{cis}$, namely the minimum of the LJ potential
with length scale σ_{cis}. The result is

$U(\phi) = \epsilon \frac{(\phi^3 - \phi_{r_{max}}^3)^2}{4 \phi_{r_{max}}^6}$ for $\phi \ge \phi_{r_{max}}$ or 0 for $\phi < \phi_{r_{max}}$, (4)

Where $\phi_{r_{max}} = 2^{\frac{2}{3}} / (3^{\frac{3}{2}} \sigma_{cis}^2) = 2 / (3^{\frac{3}{2}} r_{max}^2)$ is the FisB concentration corresponding to a
nearest neighbor distance r_{max}.

We minimize $F = F_m + F_p$ with respect to ϕ to obtain an equation for the equilibrium
density of FisB proteins in the neck

$k_B T \left[1 + \ln \left(\frac{\phi}{\phi_0} \right) \right] + 2 a \phi \left[\left(\frac{\sigma}{R} \right)^{12} - \left(\frac{\sigma}{R} \right)^{6} \right] + \partial_\phi U(\phi) = 0.$ (5)

Then, minimizing F with respect to R yields an equation that determines the equilibrium
radius of the neck

$\quad \gamma_{\text{eff}} - \frac{\kappa}{2R^2} + a\phi^2 \left[6 \left(\frac{\sigma}{R} \right)^6 - 12 \left(\frac{\sigma}{R} \right)^{12} \right] - \frac{2\gamma R}{L} = 0, \quad$ (6)

$\quad$ where $\gamma_{\text{eff}} = \gamma + k_B T \phi \ln \left(\frac{\phi}{\phi_0} \right) + U(\phi).$

Figure 9B shows R and ϕ as functions of the FisB trans homo-oligomerization strength
a, for different values of surface tension γ and neck length L. For realistic parameters
(the dot), we find that FisB trans interactions are strong enough to stabilize the neck at
$R \sim 3$ nm, with a close-packed concentration of FisB in the neck $\phi \approx \phi_{r_{\text{max}}}$. For these
same parameters, there is a critical lower limit of a below which the FisB interactions
are too weak to stabilize the neck, so the neck opens, i.e. $R \rightarrow \infty$ in our simple model.
Additionally, Figure 9B shows that the shorter the length of the neck L, the stronger the
trans interactions needed to stabilize the neck at a finite radius. This makes intuitive
sense: the longer the neck, the more FisB can be present to hold the neck together in
opposition to membrane tension. (Note that expanding the radius of the neck actually
decreases the total membrane area, which is the sum of the membrane in the neck and
in the parallel sheets, so that surface tension tends to make the neck expand – see S2
Appendix).

While the above results suggest that an accumulation of FisB at the neck can be
energetically stable, one question is how long it might take to reach that state? We
expect nucleation of a critical cluster of FisB to be rate limiting, since the time required
for diffusion and capture to reach ~40 FisB in the neck is quite short ~3.9 s (see S2
Appendix). To obtain a simple estimate of the nucleation time for both low-expression
and native-expression strains, we assume that FisB proteins diffuse independently on
the entire membrane and that nucleation of a stable cluster in the neck occurs when n
proteins happen to be in the neck at the same time. To this end, we need to estimate
the fraction of time there are n or more FisB in the neck, as well as the correlation time,
i.e. the time between uncorrelated samples. Since we assume FisB proteins are
independent, the number of proteins in the neck will be Poisson distributed, so we only
need to know the average in the neck to obtain the full distribution. The average number
of FisB in the neck is its area, $2\pi R L$, times the background concentration, ϕ_0.
Furthermore, the correlation time is simply the time for a FisB to diffuse the length of the
neck L^2/D. Using $\phi_0 \simeq 20$ FisB μm^{-2} (see "About 40 FisB molecules accumulate at the
engulfment pole to mediate membrane fission" above) for the low-expression strain
yields <FisB>$\simeq 0.03$ in the neck. Assuming that the one-hour delay in membrane fission
during sporulation of low-expression strain is due to the time for nucleation, we can infer
that the number of FisB proteins required for nucleation is $n \approx 3$ (see S2 Appendix). If
the native-expression strain also needs $n \approx 3$ to nucleate, we can estimate its
corresponding nucleation time using $\phi_0 \simeq 100$ FisB μm^{-2}, which yields <FisB>$\simeq$
0.15 and a nucleation time of ~30 s. We conclude that for native expression levels of
FisB, nucleation of a stable cluster of FisB at the neck is not likely to be rate limiting for
the process of membrane fission.

DISCUSSION

Previously, we showed that FisB is required for the membrane fission event that marks

the completion of engulfment of the forespore by the mother cell[23]. Here, we found that
a cluster of FisB molecules is nearly always present at the membrane fission site as
evidenced by an intense fluorescent spot at the engulfment pole (ISEP) using
fluorescently tagged FisB. The number of FisB molecules accumulated at the ISEP
correlates well with the fraction of cells having undergone membrane fission at a given
time point after induction of sporulation (Figs. 1,7). In addition, the number of wild-type
FisB molecules per ISEP correlates with the total number of FisB molecules per cell (S1
Appendix Figure 4). Thus, the kinetics of membrane fission is determined by the
accumulation of FisB molecules at the fission site. Lowering FisB expression could slow
membrane fission by slowing the accumulation of FisB at the pole, or by reducing the
number of FisB molecules driving fission after they are localized at the fission site. Our
modeling results are consistent with slower ISEP nucleation in the low-expression
strain, however, currently, we cannot experimentally distinguish between the two
possibilities, and both may be operating simultaneously.

How is FisB recruited to the fission site? Our results suggest FisB does not rely on
existing landmarks, lipid microdomains, cell-wall remodeling machinery, pH or voltage
gradients across the cell membrane, or membrane curvature cues for its dynamic
localization. In addition, we could not detect proteins interacting with FisB other than
itself using an anti-GFP resin pulling on YFP-FisB[23]. By contrast, we found self-
oligomerization and binding to acidic lipids to be critical for FisB's function, and purified
FisB ECD can bridge artificial membranes. Together, these results suggest FisB-FisB
and FisB-lipid interactions are key drivers for FisB clustering and accumulation at the
membrane fission site.

Can FisB oligomerization and lipid binding be sufficient to accumulate an immobile
cluster of FisB molecules at the engulfment pole? Modeling suggests this is indeed the
case. First, the narrow neck enables FisB's on opposing membranes to come close
enough to interact in trans. We infer this to be the preferred orientation for FisB-FisB
interaction, since otherwise large clusters would be expected to form elsewhere as well.
Second, the unique geometry of the neck connecting the engulfment membrane to the
rest of the mother cell membrane plays an important role, as this is the only region in
the cell where a cluster of FisB molecules can be "trapped", i.e., once a cluster is
formed inside the neck, it cannot diffuse away without breaking apart. This idea is
supported by the fact that we do not observe any FisB accumulation at the leading edge
of the engulfment membrane until a thin neck has formed at the end of engulfment.

The first FisB oligomers that appear during sporulation are dim, mobile clusters (DMCs),
each containing about a dozen FisB molecules. (One possibility is that the DMCs may
correspond to local membrane folds stabilized by FisBs interacting in trans.) Diffusion of
DMCs appears to be Brownian on the 10-20 s time scale (Figure 2), though a rigorous
analysis would require taking into account the geometry of the system. A DMC can
diffuse a typical distance of ~1 μm in ~ 5 min ($D_{DMC} \approx 3 \times 10^{-3} \mu m^2/s$, Figure 2E). By
comparison, engulfment in individual cells takes ~60 min on average[76]. Though the
engulfment time is much longer than the DMC diffusion time, the neck region, with an
inner diameter of several nanometers, only forms at the very end of the engulfment
process. Thus, ~40 FisB molecules could be recruited to the neck through diffusion-
limited capture of a few DMCs. However, we could not image such capture events

directly, and cannot rule out that FisB can also diffuse as monomers and could be
recruited to the neck in that form. Indeed, a simple model of the rate of nucleation of a
cluster of FisBs at the neck suggests that as few as three FisBs interacting in trans
could be sufficient to form a stable cluster there, with a nucleation time significantly
shorter than the engulfment time at native expression levels.

How many FisB molecules are needed for efficient membrane fission? In cells
completely lacking FisB, ~5% of the cells undergo membrane fission by t=3 h,
compared to ~80 % or ~30% for cells expressing FisB at native or ~8-fold reduced
levels, respectively (Figure 1F). The former achieve fission with ~40 copies, while the
latter with only ~6. Thus, FisB is not absolutely required for membrane fission, but it
makes it much more efficient, i.e. FisB catalyzes membrane fission. The variable
stoichiometry suggests that FisB does not oligomerize into a specific quaternary
structure with a definite stoichiometry. This variability appears to be a common property
among proteins catalyzing membrane fusion and fission, such as SNAREs[77-79] or
dynamin[14]. The smallest clusters associated with membrane fission had ~6 FisB copies
on average. This number is likely sufficient to form at least one ring inside the
membrane neck that eventually undergoes fission. Given that fission can occur in the
absence of FisB, it is likely that the FisB cluster cooperates with other cellular processes
to produce stress on this membrane neck.

We found FisB dynamics and membrane fission are not affected by removal of CL, PE,
or both. CL and PE are widely implicated in membrane fission and fusion reactions due
to their tendency to form non-bilayer structures[48,80-83]. The fact that CL or PE do not
affect membrane fission during sporulation is remarkable, because such lipids usually
affect the kinetics and/or the extent of fusion/fission reactions even if they are not
absolutely required[81]. We tested the role of CL in a strain that lacked all three known CL
synthases, with no detectable CL levels. A previous study reported that in *ΔclsABC B.*
subtilis cells, CL levels increase from undetectable during vegetative growth to readily
detectable during sporulation[31], suggesting a yet unidentified sporulation-specific CL
synthase may exist. Our results differ from those of Kawai et al. in that we were unable
to detect any CL in *ΔclsABC B. subtilis* cells during vegetative growth or sporulation. We
suggest the differences may be due to the different strain backgrounds used[84], PY79[85]
here vs. Bs168[86] in Kawai et al. or differences in detection sensitivities.

Overall, our results suggest FisB localizes to the membrane fission site using only lipid-
binding, homo-oligomerization, and the unique geometry encountered at the end of
engulfment. We propose that accumulation of a high enough density of FisB leads to
membrane fission, possibly by generating increased stress in the FisB network-
membrane composite, or in cooperation with another cellular process. A FisB
homologue with low sequence identity partially rescued fission defects in *ΔfisB B.*
subtilis cells, consistent with the idea that FisB acts as an independent module relying
mainly on homo-oligomerization, lipid-binding, and sporulation geometry.

MATERIALS AND METHODS

Materials

E. coli cardiolipin (CL), *E. coli* L-α-phosphatidylglycerol (PG), egg L-α-phosphatidylcholine (eggPC), *E.coli* L-α-phosphatidylethanolamine (PE), 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-(7-nitro-2-1,3-benzoxadiazol-4-yl) (NBD-PE), 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE), 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1,2-dioleoyl-sn-glycero-3-phospho-L-serine (DOPS) were purchased from Avanti Polar Lipids. 1-(4-Trimethylammoniumphenyl)-6-Phenyl-1,3,5-Hexatriene *p*-Toluenesulfonate (TMA-DPH) and *N*-(3-Triethylammoniumpropyl)-4-(6-(4-(Diethylamino) Phenyl) Hexatrienyl) Pyridinium Dibromide (FM4-64), and 1,1'-Dioctadecyl-3,3,3',3'-Tetramethylindodicarbocyanine (DiD) were from Thermo Fisher Scientific. Molybdenum Blue spray reagent was from Sigma-Aldrich. Carbonyl cyanide m-chlorophenyl hydrazone (CCCP) was purchased from Abcam and valinomycin was purchased from VWR. 3-(N-maleimidylpropionyl)biocytin (MBP) was obtained from Invitrogen and the HRP-conjugated antibody from eBioscience. Zaragozic acid was purchased from Sigma-Aldrich. 4-acetamido-4′-maleimidylstilbene-2,2′-disulfonic acid (AMS) and zaragozic acid were from obtained from Cayman Chemical Company.

General *B. subtilis* methods

B. subtilis strains were derived from the prototrophic strain PY79[85]. Sporulation was induced in liquid medium at 37°C by nutrient exhaustion in supplemented DS medium (DSM)[87] or by resuspension according to the method of Sterlini & Mandelstam[88]. Sporulation efficiency was determined in 24–30 h cultures as the total number of heat-resistant (80°C for 20 min) colony forming units (CFUs) compared to wild-type heat-resistant CFUs. Lipid synthesis mutants were from the *Bacillus* knock-out (BKE) collection[89] and all were back-crossed twice into *B. subtilis* PY79 before assaying and prior to antibiotic cassette removal. Antibiotic cassette removal was performed using the temperature-sensitive plasmid pDR244 that constitutively expresses Cre recombinase[89]. Cassette removal was further confirmed by PCR with primers flanking the deletion. *B. subtilis* strains were constructed using plasmidic or genomic DNA and a 1-step competence method. Site directed mutagenesis was performed using Agilent's Quick-change Lightning kit following manufacturer's instructions and mutations were confirmed by sequencing. The strains and plasmids used in this study are listed in S1 Appendix Tables 2 and 3, respectively.

Live-cell fluorescence microscopy of *B. subtilis*

Cells were mounted on a 2% agarose pad containing resuspension medium using a gene frame (Bio-Rad). Cells were concentrated by centrifugation (3300g for 30 s) prior to mounting and visualization. This step had no impact on the localization of the fusion proteins. Fluorescence microscopy was performed using a Leica DMi8 wide-field inverted microscope equipped with an HC PL APO 100×DIC objective (NA=1.40) and an iXon Ultra 888 EMCCD Camera from Andor Technology. Membranes were stained with TMA-DPH at a final concentration of 100 μM. Excitation light intensity was set to 50% and exposure times were 300 ms for TMA-DPH (λ_{ex}=395/25 nm; λ_{em}=460/50 nm); 500 ms for m(E)GFP (λ_{ex}=470/40; λ_{em}=500-550) and 1 s for mYFP (λ_{ex}=510/25; λ_{em}>530) respectively. Images were acquired with Leica Application Suite X (LAS X)

742 and analysis and processing were performed using the ImageJ software[90].

Determination of FisB's topology

We used the substituted cysteine accessibility method (SCAM[91]) to determine the
topology of FisB. We first generated stains expressing FisB versions with a single
cysteine substitution at position G6, L137, or A245, in a *ΔfisB* background. FisB does
not have any endogenous cysteines. These point mutations decreased the sporulation
efficiency slightly (S1 Appendix Table 2), we assume without affecting the topology. We
selectively biotinylated extra- or intracellular cysteines of *B. subtilis* protoplasts,
produced by addition of 0.5 mg/ml lysozyme and incubating cells at 37°C for 1h with
gentle rocking. Protoplasts were then incubated with the membrane-impermeant
reagent 3-(N-maleimidylpropionyl)biocytin (MPB). To selectively label extracellular
cysteines, protoplasts of sporulating cells at 2.5 h into sporulation were incubated with
100 µM MPB. The reaction was quenched with 50 mM DTT before cells were lysed with
hypotonic shock. To label intracellular cysteines selectively, extracellular cysteines of
protoplasts were first blocked AMS before cells were lysed and incubated with 100 µM
MPB. The reaction was quenched by addition of 100 µM MPB. FisB was pulled down
from the cell lysates as described in[91] using an anti-Myc antibody (mAb #2276) and
biotinylated proteins were detected by Western Blot using a HRP-conjugated-Avidin
antibody. Further details are provided in the S1 Appendix.

Expression, purification, and labeling of recombinant FisB protein

Recombinant soluble FisB ECD was purified as described in[23] but with slight
modifications. Briefly, His_6-FisB ECD was expressed in *E. coli* BL21 (DE3) from New
England Biolabs and purified using HisPur™ Ni-NTA Resin from Thermo Fisher
Scientific. Protein expression was induced with 1 mM IPTG at OD_{600} = 0.6 overnight at
16°C. Cells were harvested by centrifugation and the pellet was resuspended in Lysis
Buffer (20 mM HEPES, 500 mM NaCl, 0.5 mM TCEP, 20 mM Imidazole, 2% glycerol,
20 mM $MgCl_2$) and flash-frozen in liquid nitrogen. Pellets were thawed on ice and cells
were lysed by 5 passes through a high-pressure homogenizer (Avestin EmulsiFlex-C3).
The lysate was spun down at 100,000×g and the soluble fraction was incubated with
HisPur™ Ni-NTA Resin for 2.5 h at 4°C while rotating. The bound protein was washed
with Lysis Buffer, Lysis Buffer containing 50 mM and finally 100 mM Imidazole. The
protein was eluted in Elution Buffer (20 mM HEPES, 500 mM NaCl, 0.5 mM TCEP, 200
mM Imidazole, 2% glycerol, 20 mM MgCl2). The protein was concentrated using a
Vivaspin centrifugal concentrator with a 10 kDa molecular weight cutoff and the
concentration determined by Bradford protein assay. The protein was stored at -80ºC.

In experiments with labeled FisB ECD, we used a cysteine mutation, G123C (FisB ECD
does not have any endogenous cysteines). After expression and purification as above,
iFluor555-maleimide (AAT Bioquest) was reacted with FisB ECD[G123C] following the
manufacturer's instructions. G123 is in a loop that if removed does not interfere with
FisB's function (S1 Appendix Figure 10).

**Analytical size-exclusion chromatography (SEC) and negative-stain electron
microscopy (EM)**

For SEC analysis His_6-FisB ECD was loaded onto a Superose 6 Increase 10/300 GL

column (GE) previously equilibrated with 20 mM HEPES, pH 7.5, 500 mM NaCl, 0.5 mM
TCEP, 2% glycerol, 20 mM MgCl₂, running at a flow rate of 0.5 ml/min at 4ºC. The
column was calibrated with Bio-Rad's Gel Filtration Standards. For negative stain EM
analysis, 4 µL of the indicated elution fractions were applied to 200-mesh copper grids
coated with ~10 nm amorphous carbon film, negatively stained with 2% (wt/vol) uranyl
acetate, and air-dried. Images were collected on a FEI Tecnai T12 microscope, with a
LaB6 filament operating at 120 kV, and equipped with a Gatan CCD camera.

Inhibition of cell wall synthesis and analyses of FisB motions

Overnight cultures of GFP-Mbl (BDR2061) or IPTG-induced mGFP-FisB (BMB014)
were diluted in CH medium to OD600 = 0.05. Expression of GFP-FisB was induced with
1 mM IPTG for 2h at 37°C. Expression of GFP-Mbl was induced with 10 mM xylose for
30 min when BDR2061 reached OD600 = 0.5. For imaging untreated cells, 1 ml of cells
was washed twice with 1 ml PBS and finally resuspended in 10 µl PBS. 2 µl of cell
suspension was spread on a 2% PBS agar pad for imaging. To inhibit cell-wall
synthesis 50 µg/ml fosfomycin was added to the cultures 45 min before imaging. 1 ml of
cells was washed twice with PBS containing 50 µg/ml fosfomycin and mounted on a
PBS agar pad also containing fosfomycin. Cells were imaged using a Olympus IX81
microscope with a home-built polarized TIRF setup[92,93]. Exposure times were 50 ms for
BDR2061 and 100 ms for BMB014. Movies were acquired at 1 frame/s. Movies
collected for BMB014 were corrected for bleaching using the Bleaching Correction
function (exponential method) in ImageJ. Kymographs were created with imageJ along
the indicated axes. GFP fusion proteins were tracked using the ImageJ plugin
TrackMate[94]. A Laplacian of Gaussian (LoG) filter was used to detect particles with an
estimated blob diameter 400 µm. Particles were tracked using the Simple LAP tracker
with a 0.25 µm maximum linking distance and no frame gaps. MATLAB (Mathworks,
Natick, MA) was used for further processing of the tracks. Mean squared displacement
(MSD) was calculated using the MATLAB class @msdanalyzer[95].

The asymmetry of individual tracks (S1 Appendix Fig. 5F) was calculated as described
in[96] using:

$$814 \quad Asym = -log\left(1 - \frac{(R_1^2 - R_2^2)^2}{(R_1^2 + R_2^2)^2}\right)$$

815 where R_1 and R_2 are the principal components of the radius of gyration, equal to the
816 square roots of the eigenvalues of the radius of gyration tensor $\boldsymbol{R_g}$:

$$817 \quad \boldsymbol{R_g(i,j)} = \langle x_i x_j \rangle - \langle x_i \rangle \langle x_j \rangle.$$

Tracking fluorescently labeled FisB spots and estimation of diffusion coefficients

819 For estimating the mobility of DMC and ISEP, time-lapse movies were recorded with a
820 frame rate of 1 s using wide-field microscopy (50% LED intensity, 300 ms exposure
821 time, gain 300). Spot positions were tracked using SpeckleTrackerJ[97], a plugin for the
822 image analysis software ImageJ[90]. Mean-squared displacements (MSDs) were
823 calculated using the MATLAB class @msdanalyzer[95].

Dissipation of membrane potential

825 Cells were concentrated by centrifugation (3300xg for 30 s) and 100 µM CCCP or 30
826 µM valinomycin was added just prior to mounting cells onto a 2% PBS agar pad also
827 containing 100 µM CCCP or 30 µM valinomycin.

Lipid extraction and thin-layer chromatography (TLC)

829 Lipids were extracted from *B. subtilis* cells at 3 h into sporulation according to the
830 method of Lacombe and Lubochinsky[98]. Lipid extracts were analyzed by TLC on silica
831 gel plates in mixtures of chloroform:hexane:methanol: acetic acid (50:30:10:5).
832 Phospholipids were detected with Molybdenum Blue Reagent (Sigma-Aldrich).

Liposome preparation

Small unilamellar vesicles (SUVs) were prepared by mixing 1 µmol of total lipids at
desired ratios. A thin lipid film was created using a rotary evaporator (Buchi). Any
remaining organic solvent was removed by placing the lipid film under high vacuum for
2h. The lipid film was hydrated with 1 ml of RB-EDTA buffer [25 mM HEPES at pH 7.4,
140 mM KCl, 1 mM EDTA, 0.2 mM tris(2-carboxyethyl) phosphine] by shaking using an
Eppendorf Thermomix for >30 min. The lipid suspension was then frozen and thawed 7
times using liquid nitrogen and a 37°C water bath and subsequently extruded 21 times
through a 100 nm pore size polycarbonate filter using a mini-extruder (Avanti). All SUVs
contained 1% NBD-PE to determine the final lipid concentration.

Giant unilamellar vesicles (GUVs) were prepared by electroformation[99]. Briefly, lipids
dissolved in chloroform were mixed in a glass tube at desired ratios and spotted on two
indium tin oxide (ITO) coated glass slides. Organic solvent was removed by placing the
lipid films in a vacuum desiccator for at least 2 h. A short strip of copper conductive tape
was attached to each ITO slide which were then separated by a 3 mm thick
Polytetrafluoroethylene (PTFE) spacer and held together with binder clips. The chamber
was filled with 500 µl Swelling Buffer (1 mM HEPES, 0.25 M sucrose, 1 mM DTT) and
sealed with Critoseal (VWR International, Radnor, PA). GUVs were formed by applying
a 1.8 V sinusoidal voltage at 10 Hz for at least 2 h at room temperature.

For experiments involving FisB ECD the GUVs were composed of (all in mole
percentages): 25 *E. coli* PE, 5 *E. coli* CL, 50 *E. coli* PG, 19 eggPC and 1 DiD or 1 NBD-
PE. For experiments in which EndoA1 was used, GUV composition was (all in mole %):
45 DOPS, 24.5 DOPC, 30 DOPE and 0.5 DiD.

Liposome-protein co-floatation

For initial experiments, 40 nmol total lipid was incubated with 200 pmol FisB ECD for 1h
at room temperature in a total volume of 100 µl. 200 µl of 60% Optiprep (iodixanol,
Sigma-Aldrich) was added to the sample creating a 40% Optiprep solution. The sample
was then layered at the bottom of a 5 mm x 41 mm Beckman ultracentrifuge tube
(#344090) and overlaid with 200 µl of 20% Optiprep and finally 150 µl of buffer (Figure 4
C). Liposome-bound proteins co-float to a light density, while unbound proteins pellet
upon ultracentrifugation for 1.5 h at 48 krpm. Fractions were collected as shown in
Figure 4C and the amount of recovered protein was determined by SDS-PAGE (Nu-
PAPGE 12% Bis-tris gel, Thermo Fisher Scientific) stained with SYPRO™ Orange
(Invitrogen).

Liposome aggregation using absorbance

SUVs were prepared by extrusion as described above but using a 50 nm polycarbonate filter. SUVs were composed of 50 mole % *E. coli* PG, 25 mole % *E. coli* PE, 20 mole % eggPC, 5 mole % *E. coli* CL. The absorbance at 350 nm of 50 µM total lipid was measured for 5 min, before addition of 1 µM FisB ECD. Absorbance increases with increasing liposome aggregation due to increased scattering[100].

Filamentous *B. subtilis* cells to test for curvature-sensitive localization of FisB

An overnight culture of BMB014 was diluted into fresh CH medium[101] to OD_{600}=0.05. 1 mM IPTG and 20 mM xylose were added to induce the expression of GFP-FisB and MciZ, respectively. The latter inhibits cytokinesis[68]. The culture was grown at 37°C for 30 min before 3-5 µl of cells were transferred onto a 3% agar pad also containing 1 mM IPTG and 20 mM xylose. Cells were grown on the agar pad for 2h at 37°C prior to imaging. GFP-FisB foci were detected using the ImageJ plugin TrackMate as described above. Radii of the inner and outer edges were determined by manually fitting a circle to the cells using ImageJ.

Determination of binding constants

For determination of binding constants, the floatation protocol was slightly modified. Varying amounts of lipids were incubated with 100 nM iFluor555-FisB ECD for 1 h at room temperature in a total volume of 100 µl. Density gradients were created as before using Optiprep (iodixanol), however only 2 fractions were collected (Figure 5H). The protein concentration in fraction A was too small to be quantified by SDS-PAGE. Therefore, the sample was concentrated by trichloroacetic acid (TCA) precipitation. Briefly, 50 µl of TCA was added to fraction A and incubated for 30 min at 4°C. The sample was spun at 14 krpm in an Eppendorf microfuge for 5 min. The pellet was washed twice with ice-cold acetone and subsequently dried for 10 min in a 95°C heating block. 10 µl of 2X SDS sample buffer was added to the dried pellet and the sample was boiled for 10 min at 95°C and loaded on a 12% bis-tris gel. The amount of recovered protein was determined by fluorescence intensity of the labeled FisB ECD band on the gel using a Typhoon FLA 9500 (GE Healthcare). The dissociation constant K_d was determined following ref. 102. Titration curves were fitted to:

$$f_b = \frac{K[L]}{1+K[L]}, \qquad (1)$$

where f_b is the fraction of bound protein and K the apparent association constant ($K = 1/K_d$). Eq. (1) assumes that the total lipid concentration $[L]$ is much larger than the concentration of bound protein, a condition satisfied in our experiments for $[L] > 10^{-7}$ M.

Image analysis

To estimate the fraction of cells that have undergone membrane fission at a particular time after sporulation was initiated by nutrient downshift, cells were labeled with TMA-DPH (see "Live-cell fluorescence microscopy of *B. subtilis*" above). The dye labels the forespore contours intensely before membrane fission, as it has access to three membranes in close proximity that cannot be resolved (forespore, engulfment, and mother cell membranes). After membrane fission, the dye dimly labels forespore

contours (see S1 Appendix Fig. 1 for examples and quantification). Due to the clear
separation between the two labeling patterns (S1 Appendix Fig. 1) cells can be scored
visually, with 6-7% of cells having intermediate labeling that prevents categorization.
Thus, we underestimate the percentage of cells that have undergone membrane fission
by at most ~7%.

For the analyses shown in S1 Appendix Figs. 4A,B,C,E, and 9, we calculated the total
intensity (sum ox pixel values) inside the cell contour (indicated in yellow in Fig. S4A)
using MicrobeJ[103]. Mean integrated auto-fluorescence (~1300 a.u) was calculated by
analyzing in the same way an equivalent number of individual wild-type cells, imaged
under identical conditions.

For the analyses shown in Figure 2 and S1 Appendix Fig. 4D, FisB foci were semi-
automatically selected using SpeckleTrackerJ[97]. For each spot the sum of pixel values
in a 6 × 6 pixel (0.5 μm × 0.5 μm) box around the center of the spot were calculated. For
each corresponding cell the same operation was performed at a membrane area where
no clusters were present and subtracted from the FisB cluster intensity.

Preparation of DNA Origami-based mEGFP standards

These standards were prepared and characterized as described in[44]. Briefly, DNA
"rods" consisted of six-helix-bundle DNA origami nanotubes. Rods carried varying
numbers of single stranded "handle" sequences for DNA-conjugated fluorophore
hybridization. A long scaffold DNA (p7308[104]) was folded into the desired shape by self-
assembly with a six-fold molar excess of designed "staple strands" by heating and
cooling cycles over an 18-hour period in a thermocycler (Bio-Rad). Excess staples
were removed by PEG precipitation[105], and DNA-conjugated fluorophores were
hybridized to the DNA origami nanotubes by coincubation for 2 hours at 37ºC. Finally,
excess fluorophore-DNA conjugates were removed by a second PEG precipitation[105].
To estimate fluorophore labeling efficiency, standards designed to host 5 copies of
Alexa Fluor 488 were similarly prepared. These standards were imaged on a TIRF
microscope (Eclipse Ti, Nikon) until fully bleached. The photobleaching steps of the
fluorescence traces were fit to a binomial function to estimate the labeling efficiency to
be ~80% (95% CI = 76%-84%).

Quantitative Western Blot

mYFP was cloned into pVS001 (His$_6$-Sumo-mYFP) and purified using affinity
chromatography. For immunoblotting, cells in 100 ml sporulation medium were pelleted
and the supernatant removed. The pellets were suspended in ice-cold lysis buffer
(pH=7.5; 50 mM HEPES, 100 mM KCl, 3 mM MgCl$_2$, 1 mM EGTA, 0.1% Triton X-100, 1
mM DTT, 1 mM PMSF, with one complete protease inhibitor tablet (Roche) to a final
volume of 300 µl, and then we added 0.3 g acid-washed glass beads (425-600 µm,
Sigma). After adding 150 µl boiling sample buffer (250 mM Tris-HCl, pH 6.8, 50%
glycerol, 3.58 µM β-mercaptoethanol, 15% SDS, and 0.025% Bromophenol Blue),
samples were incubated at 100°C for 5 min. Samples were centrifuged at 14,000 rpm in
a desktop centrifuge at room temperature for 10 min and stored at -80°C. The blots
were probed with peroxidase-conjugated anti-GFP antibody (ab13970). Images were
scanned and quantified using ImageJ.

ACKNOWLEDGEMENTS

ACKNOWLEDGEMENTS

We thank members of the Karatekin and Rudner laboratories for stimulating
discussions. This work was supported by National Institute of General Medical Sciences
and National Institute of Neurological Disorders and Stroke of the National Institutes of
Health (NIH) under award numbers R01GM114513 and R01NS113236 (to EK),
DP2GM114830 and R01GM132114 (to CL). The content is solely the responsibility of
the authors and does not necessarily represent the official views of the National
Institutes of Health. We thank Vladimir Polejaev and Jeorg Nikolaus (directors of the
Yale West Campus Imaging Core), and Josh Lees (Yale Center for Cellular and
Molecular Imaging Electron Microscopy Facility) for their help with imaging, Karin
Reinisch in whose laboratory work by FH was carried out, Daniel R. Zeigler (Bacillus
Genetic Stock Center) for helpful advice, and Alexander J. Meeske for some of the
strains used in this study. NDW was supported by a NIH training grant (T32-EB09941).
We gratefully acknowledge a Yale University Predoctoral Fellowship to MB. This work
was supported in part by the National Science Foundation, through the Center for the
Physics of Biological Function (PHY-1734030). AM-C acknowledges support from the
Spanish MECD through the program Ayudas para la Formación de Profesorado
Universitario (FPU), Grant No. FPU16/0256. We are grateful to Aurelien Roux (U.
Genève) for the generous gift of Atto395-EndoA1.

AUTHOR CONTRIBUTIONS

AL, MB, VS, and EK conceived the study. AL and MB performed experiments whose
results are shown in the main figures. AA (Fig. S10), FH (Fig. 5, 7, S8), VS (Fig. S3)
performed additional experiments. NDW and CL developed the DNA-origami
fluorescence calibration method and contributed to the data in Fig. 2. CR, TD, and DR
provided resources, training, and technical and conceptual input. They introduced EK,
AL, MB and VS to *B. subtilis* and sporulation. EK and DR provided supervision and
acquired funding. AMC and NW developed the model and wrote the corresponding
sections. AL, MB, and EK and wrote the manuscript, with input from other co-authors.
We thank Aurelien Roux for the generous gift of Atto395-EndoA1.

CONFLICT OF INTEREST

None.

FIGURE LEGENDS

Figure 1. Membrane fission during sporulation is nearly always accompanied by
accumulation of a FisB cluster at the fission site. **A**. Vegetatively growing cells enter
sporulation when nutrients become scarce. Asymmetric division creates a forespore
(FS) and a mother cell (MC). The MC engulfs the FC in a phagocytosis-like event. At

the end of engulfment, a membrane neck connects the engulfment membrane to the
rest of the MC (i). Fission of the neck (ii) releases the FS, now surrounded by two
membranes, into the MC cytoplasm. Once the forespore becomes a mature spore, the
MC lyses to release it. **B**. The membrane fission step shown in more detail. **C**. Detection
of membrane fission. The lipophilic dye TMA-DPH does not fluoresce in the aqueous
solution and crosses membranes poorly. If membrane fission has not yet taken place,
the dye has access to the engulfment, FS and MC membranes, thus shows intense
labeling where these membranes are adjacent to one another (i). If fission has already
taken place, the dye labels internal membranes poorly (ii). **D.** Images show mGFP-FisB
(strain BAM003, native expression level) at indicated times during sporulation.
Membranes were visualized with TMA-DPH. Examples of sporulating cells with mGFP-
FisB enriched at the septum (1.5 h), forming dim mobile cluster (DMC; 2 h) and with a
discrete mGFP-FisB focus at the cell pole (intense spot at engulfment pole, ISEP, 3 h)
are highlighted with arrowheads. **E.** Similar to D, but using a strain (BAL003) that
expresses mGFP-FisB at lower levels in a *ΔfisB* background. **F.** Time course of
membrane fission for wild-type cells, *ΔfisB* cells, or *ΔfisB* cells complemented with
mGFP-FisB expressed at native (BAM003) or low levels (BAL003). Lower expression of
mGFP-FisB leads to a delay in membrane fission kinetics. **G.** The percentage of cells
with an intense spot at the engulfment pole (ISEP) for low and native level expression of
mGFP-FisB as a function of time into sporulation. **H**. Correlation between percentage of
cells that have undergone fission and percentage of cells having an ISEP for all time
points shown in F and G. The fitted dashed line passing through the origin has slope
1.06 ($R^2 = 0.9$). Scale bars represent 1 μm.

Figure 2. Estimation of mEGFP-FisB copies at the engulfment pole at t=3 h using
DNA-origami calibration standards and mobility of FisB clusters. **A.** Simplified
schematic of the DNA-origami-based mEGFP standards used in this study. Using DNA
origami, DNA rods bearing AF647 at both ends and the indicated numbers of mEGFP
molecules along the rod were designed. In the actual rods, the labeling efficiency was
found to be ~80%, so the actual copies of mEGFP per rod were 4, 20, 40, 56, and 80.
B. Representative wide field images of the DNA-origami-based mEGFP standards used
in this study. Bars are 1 μm. **C**. Distributions of total fluorescence intensities (sum of
pixel values) for the intense spot at the engulfment pole (ISEP) and the dim, mobile
clusters (DMC). Background was defined individually for every cell where an ISEP or
DMC intensity measurement was performed. Examples are shown on the left. **D.** Total
fluorescence intensity (sum of pixel values) for DNA-origami rods as a function of
mEGFP copy numbers. The best fit line passing through the origin has slope 29.56
au/mEGFP ($R^2 = 0.97$). The total intensity of the ISEP and DMCs correspond to ~40
and ~12 copies of mEGFP respectively. **E.** Mean-squared displacement (MSD) as a
function of delay time for DMCs (magenta) and ISEPs (blue). Cells expressing mGFP-
FisB (strain BAM003) were imaged using time-lapse microscopy. Forty-five cells from
10 different movies at t=2.5 hr and 30 cells from 10 different movies at t=3 hr after
nutrient downshift were analyzed. (See S1 Appendix Movie 1 for a representative single
bacterium at t=2.5 hrs showing several mobile DMCs and Movie 2 for a representative
single bacterium at t=3 hrs showing an immobile ISEP.) Fits to the initial 25 s (~10 % of

delays) yielded $D_{DMC} = 2.80 \pm 0.05 \times 10^3$ nm$^2/s$ (± 95% confidence interval, R² = 0.999,
24 tracks) and $D_{ISEP} = 2.80 \pm 0.51 \times 10$ nm^2/s (± 95% confidence interval, R² = 0.850,
25 tracks). **F.** Summary of FisB copy number and cluster mobility estimation.

Figure 3. Membrane fission is insensitive to membrane lipid composition. **A.**
Pathways for membrane lipid synthesis in *B. subtilis*. Lipid synthetases responsible for
each step are highlighted in blue. **B.** Thin-layer chromatography (TLC) of the total lipid
extracts of wild-type and indicated lipid synthesis-deficient cells. Cells were collected 3
hrs after induction of sporulation by nutrient downshift. Phospholipid spots (PLs) were
visualized by staining with Molybdenum Blue spray reagent. Purified CL, PG, and PE
were used as standards to identify the PLs of *B. subtilis*. Arrows indicate locations to
which individual standards migrate. **C.** Membranes from cells of the indicated genetic
backgrounds were visualized with TMA-DPH at t=3h. The images are from cells
mounted on agarose pads containing sporulation medium. Bar,1 µm. **D.** Percentage of
cells from indicated strains that have undergone membrane fission as a function of time
after initiation of sporulation. For every strain, 150-220 cells from 3 independent
experiments were analyzed at the indicated times during sporulation. **E.** mGFP-FisB
(strain BAM003) treated with the squalene-synthase inhibitor zaragozic acid, imaged at
t=3 h. **F.** Cells expressing mYFP-FisB (low expression levels) in either wild type
(BAL002) or in a CL deficient strain (BAL037) at t=3h. Membranes were visualized with
the fluorescent dye TMA-DPH. Examples of sporulating cells with a discrete mYFP-FisB
focus at the cell pole (ISEP) are highlighted (white arrows). Foci were semi-
automatically selected with SpeckletrackerJ[97]. **G.** The percentage of cells with an
intense spot at engulfment pole for wild-type (BAL002) or cardiolipin-deficient (BAL037)
mYFP-FisB expressing cells at t=3h (low expression). For each strain, 150-220 cells
from 3 independent experiments were analyzed. **H.** Distributions of total fluorescence
intensities (sum of pixel values) at ISEP for wild-type (BAL002) or cardiolipin-deficient
(BAL037) mYFP-FisB cells at 3hr into sporulation. For every strain, 150 ISEPs were
analyzed. Scale bars are 1 µm.

Figure 4. **Binding of FisB ECD to acidic lipids. A**. Domain structure of FisB and its
His$_6$-tagged extracytoplasmic domain (ECD) used in floatation experiments. **B.**
Predicted model of FisB$^{44-225}$ comprising most of the ECD[57], schematically attached to
the membrane. **C**. Schematic of the floatation assay. Liposomes (40 nmol total lipid)
and FisB ECD (200 pmol) were incubated for 1 hour (total volume of 100 µl) at room
temperature and layered at the bottom of an iodixanol density gradient. Upon
ultracentrifugation, liposomes float to the top interface, whereas unbound protein
remains at the bottom. Four fractions were collected as indicated and analyzed by SDS-
PAGE. **D**. SYPRO orange stained gel of FisB ECD incubated with liposomes containing
45 mole % CL. The percentage of recovered protein is determined by comparing the
intensity of the band in fraction B to the input band intensity. **E**. Indistinguishable
amounts of FisB ECD are recovered when FisB ECD is incubated with liposomes
containing different acidic lipid species as long as the charge density is similar. CL30,
PG60, PS60 indicate liposomes containing 30 mole % CL, 60 mole % PG and 60 mole
% PS, respectively. CL carries 2 negative charges, whereas PG and PS carry one each.
The rest of the liposome composition is PC. **F**. Fraction of liposome-bound iFluor555-
labeled FisB ECD (iFluor555-FisB ECD, 100 nM) recovered after floatation as a function

of lipid concentration. Titration curves were fit to $f_b = K[L]/(1 + K[L])$, where f_b is the
bound fraction of protein, $[L]$ is the total lipid concentration (assumed to be $\gg$ [protein
bound]), and $K = 1/K_d$ the apparent association constant, and K_d is the apparent
dissociation constant. **G**. Best fit values for K_d were 1.0 µM for CL (95% confidence
interval, CI=0.7-2.1 µM) and 3.6 µM for PG (CI=2.8-5.0 µM), respectively. iFluor555-
FisB ECD (100 nM) was incubated with 10^{-8} to 10^{-4} M lipids for 1 h at room temperature
before flotation. Liposomes contained 45 mole % of CL or PG and 55% PC.

Figure 5. FisB mutants selectively impaired in oligomerization and membrane
binding. A. Mutated residues shown on the FisB domain structure. **B**. Kyle-Doolittle
hydrophobicity profile of the FisB sequence for wild-type (WT), FisB K168D,K170E
(FisB^{KK}), and FisB G175A,I176S, I195T, I196S (FisB^{GIII}) mutants. **C**. Mutations shown
on the predicted model[57] of FisB$^{44-225}$. Residue conservation (top) and electrostatic
potential (bottom) are mapped onto the structure. **D**. Western blot of cell lysates from *E.*
coli cells expressing FisB-ECD^{WT}, FisB-ECD^{GIII}, or FisB-ECD^{KK}, probed with an anti-
histidine antibody. High molecular weight bands in the WT and KK lanes are largely
absent in the GIII lane, indicating FisB^{GIII} is less prone to forming oligomers. **E**. Size-
exclusion chromatography of FisB WT and the GIII and KK mutants. Intensities of high
and low molecular weight peaks are reversed for FisB WT and the GIII mutant, whereas
the KK mutant has a profile similar to WT. **F**. A fraction corresponding to the high-
molecular peak in E (indicated by *) for FisB WT was collected and imaged using
negative-stain electron microscopy (EM), which revealed flexible, elongated structures
~50 nm × 10 nm. **G**. A similar analysis for FisB^{GIII} revealed more heterogeneous and
less stable structures. Scale bars in F, G are 50 nm. **H**. Schematic of the floatation
experiments to determine the apparent affinity of FisB mutants for liposomes containing
acidic lipids. Experiments and analyses were carried out as in Figure 4, except only two
fractions were collected. iFluor555-FisB ECD (100 nM) was incubated with 10^{-8} to 10^{-4} M
lipids for 1 h at room temperature before floatation. Liposomes contained 45 mole % of
CL and 55% PC. **I**. Fraction of protein bound to liposomes as a function of total lipid
concentration. Data was fitted to a model as in Figure 4F. The data and fit for FisB WT
is copied from Figure 4F for comparison. **J.** Best fit values for K_d were 1.0 µM for WT
(95% confidence interval, CI=0.7-2.1 µM), 9.1 µM for KK (CI=6.5-15.3 µM), and 1.6 for
GIII (CI=0.9-5.1 µM), respectively**.**

Figure 6. FisB clustering and binding to acidic lipids are both required for ISEP
formation and membrane fission. A. Snapshots of sporulating *ΔfisB* cells expressing
mYPF-FisB^{WT} (BAL002), mYPF-FisB^{KK} (BAL006), or mYPF-FisB^{GIII} (BAL007), at low
levels. For each time point after downshifting to the sporulation medium, cell
membranes were labeled with TMA-DPH and images were taken both in the membrane
(left) and the YFP (right) channels. By t=2.5 h, some foci at the engulfment pole (ISEP)
are visible for WT cells that have undergone membrane fission (red boxes), but not for
the KK or GIII mutants (white boxes). A small fraction of KK mutants (7.3%)
accumulated FisB at the engulfment pole and underwent membrane fission at t=3h.
Scale bars represent 1 µm. **B.** Percentage of cells with an intense spot at the
engulfment membrane (ISEP) at t=3 h into sporulation, for WT FisB, FisB^{KK}, or FisB^{GIII}.
For every strain, 200-300 cells from three independent experiments were analyzed at
the indicated times during sporulation. **C**. Distribution of background-corrected

integrated intensities (sum of pixel values) of ISEP fluorescence for *ΔfisB* cells
expressing mYFP-FisB^{WT} or mYPF-FisB^{KK}. The distributions are indistinguishable.
Since low-expression cells accumulate, on average, 6±2 FisB^{WT} molecules at the ISEP
(Fig. S4D), so do FisB^{KK} cells. 175 and 68 ISEPs were analyzed for WT and KK mutant
strains. **D**. Percentage of cells that have undergone membrane fission at the indicated
time points. (For every strain, 200-300 cells from 3 independent experiments were
analyzed at the indicated times during sporulation.)

Figure 7. *C. perfringens* FisB can substitute for *B. subtilis* FisB despite poor
sequence identity. A. Heat-resistant colony forming units for *ΔfisB* cells expressing *B.*
subtilis (BAL001) or *C.perfringens* FisB (BAL005) at native levels, presented as a
percentage of the WT sporulation efficiency. Results are shown as means ± SD for
three replicates per condition. **B.** Snapshot of *ΔfisB* cells expressing mEGFP-
FisB^{Cperfringens}. Aliquots were removed at the indicated times, membranes labeled with
TMA-DPH, and both the TMA-DPH and the EGFP channels imaged after mounting into
agar pads. White boxed areas are shown on an expanded scale in yellow boxes.
Arrows indicate cells with ISEP that have undergone membrane fission. Bar, 1 µm. **C**.
Percentage of cells that have undergone membrane fission as a function of sporulation
time for wild-type cells, *ΔfisB* cells, *ΔfisB* cells expressing *B. subtilis* mEGFP-FisB at
native levels, or *ΔfisB* cells expressing mEGFP-FisB^{Cperfringens}. The plots for the first
three conditions are reproduced from Figure 1F for comparison. **D.** Distribution of
background-corrected total fluorescence intensity of ISEP for *ΔfisB* cells expressing
mEGFP-FisB^{Cperfringens} or mEGFP-FisB^{Bsubtilis} at native levels. From the calibration in
Figure 2D, we estimate 9±7 FisB^{Cperfringens} per ISEP. The distribution for mEGFP-
FisB^{Bsubtilis} is reproduced from Figure 2C for comparison. (150 and 93 ISEPs were
analyzed for mEGFP-FisB^{Bsubtilis} and mEGFP-FisB^{Cperfringens}, respectively.) **E**.
Percentage of cells with ISEP, for *ΔfisB* cells expressing mEGFP-FisB^{Cperfringens} or
mEGFP-FisB^{Bsubtilis}. (For each strain, 200-300 cells from 3 independent experiments
were analyzed.) **F**. Percentage of cells that have undergone membrane fission at t=3 h
vs. the percentage of cells with ISEP at the same time point, for the conditions
indicated. There is a nearly perfect correlation between these two quantities (the
dashed line is a best-fit, $y = 1.03x, R^2 = 0.96$).

Figure 8. FisB does not sense or induce membrane curvature. A. FisB ECD does
not induce deformation of GUV membranes. Left: GUVs incubated with 2 µM iFluor555-
labeled FisB ECD did not show any tubulation or invagination of the GUV membrane.
GUVs were composed of (in mole %: 25 *E. coli* PE, 5 *E. coli* CL, 50 *E. coli* PG, 19
eggPC, 1 NBD-PE). Right: incubation of 2 µM endophilin A1 (EndoA1, labeled with
Atto395) with GUVs (45% DOPS, 24.5% DOPC, 30% DOPE and 0.5% DiD) resulted in
extensive tubulation of membranes, as reported previously[22]. The two proteins have
similar affinities for GUV membranes under these conditions (Figure 4F and ref. 106).
B. FisB ECD cannot deform deflated GUVs and its membrane localization is
independent of curvature. To avoid potential issues with high membrane tension
preventing membrane deformation, GUVs were deflated using osmotic stress, which
resulted in deformed GUVs with both negatively and positively curved regions. FisB
ECD bound to these GUVs was unable to induce any high-curvature deformations. The
intensity of iFluor555-FisB ECD along a membrane contour (proportional to coverage)
was plotted against membrane curvature in the corresponding region. There was no

correlation between membrane curvature and FisB ECD coverage. **C**. FisB localization
does not depend on curvature in filamentous *B. subtilis* cells. GFP-FisB was expressed
under an inducible promoter during vegetative growth and cell division was blocked by
inducing expression of MciZ[68]. Cells grew into long flexible filaments that were bent to
varying degrees. The linear density of GFP-FisB spots (spots/µm) was independent of
filament curvature. **D**. FisB ECD bridges GUV membranes. iFLuor555-FisB ECD (100
nM) was incubated with GUVs (same composition as in A and B). Many GUVs were
found adhering to one another. iFluor555-FisB ECD signals were enhanced in the
adhesion patches, in particular at the rims. Intensity profiles along the highlighted
contours are shown below the examples. **E**. FisB ECD aggregates small liposomes.
Liposomes (in mole %: 25 *E. coli* PE, 5 *E. coli* CL, 50 *E. coli* PG, 19 eggPC, 50 µM total
lipid) were incubated in the absence and presence of FisB ECD (unlabeled) and their
aggregation monitored by absorbance at 350 nm. FisB was added at 5 min (1 µM final),
which caused the absorbance to increase, indicating increased liposome aggregation.

Figure 9. Modeling supports recruitment of FisB to membrane neck via
oligomerization without curvature sensing. A. Left: Schematic of the late stages of
engulfment, when a small membrane neck connects the engulfment membrane to the
rest of the mother cell membrane. Right: Schematic of FisB accumulation at the fission
site. FisB freely moves around the engulfment membrane and other regions of the
mother cell membrane, forming clusters of up to ~12 molecules. Cluster motions are
independent of lipid microdomains, flotillins, the cell-wall synthesis machinery, and
voltage or pH gradients. About 40 copies of FisB accumulate at the membrane neck in
an immobile cluster. Bottom: Modeled axisymmetric membrane neck of radius R and
length L connecting two membrane sheets. The uniform areal concentration of FisB in
the neck is ϕ. **B.** Top: Equilibrium radius of the neck as a function of FisB trans homo-
oligomerization strength, a, for several values of neck length, L. Below a minimum
interaction strength, FisB cannot stabilize the neck and the neck opens. The horizontal
line is the radius corresponding to the minimum of the potential describing the trans
interaction, $R = 2^{1/6}\sigma$ (Eq. 3). Bottom: Equilibrium FisB concentration in the neck as a
function of a. The horizontal line is $\phi_{r_{max}} = 2/(3^{3/2}r_{max}^2)$, the concentration of FisB at the
onset of in-plane crowding. Model parameters (see Eqs. 5 and 6): $\kappa = 20\ k_B T$ (ref. 107),
$\phi_0 = 100$ FisB μm^{-2}, $\gamma = 10^{-4}$ N m^{-2} (ref. [108]), $\sigma_{cis} \simeq 2.47$nm, $\phi_{r_{max}} = 5 \times 10^4$ FisB μm^{-2},
$L = 40$ nm, and for the dot $a \simeq 10^4\ k_B T$ nm^2 FisB^{-1}. For details, see S2 Appendix.

1206
1207
1208

REFERENCES

1. Haucke V, Kozlov MM. Membrane remodeling in clathrin-mediated endocytosis.
J Cell Sci **131**, (2018).
2. Campelo F, Malhotra V. Membrane fission: the biogenesis of transport carriers.
Annu Rev Biochem **81**, 407-427 (2012).
3. Ahmed I, Akram Z, Iqbal HMN, Munn AL. The regulation of Endosomal Sorting
Complex Required for Transport and accessory proteins in multivesicular body

sorting and enveloped viral budding - An overview. *Int J Biol Macromol* **127**, 1-11
(2019).

4. Jaumouille V, Waterman CM. Physical Constraints and Forces Involved in
Phagocytosis. *Front Immunol* **11**, 1097 (2020).

5. Carlton JG, Jones H, Eggert US. Membrane and organelle dynamics during cell
division. *Nat Rev Mol Cell Biol* **21**, 151-166 (2020).

6. Errington J. Regulation of endospore formation in Bacillus subtilis. *Nat Rev*
Microbiol **1**, 117-126 (2003).

7. Higgins D, Dworkin J. Recent progress in Bacillus subtilis sporulation. *FEMS*
Microbiol Rev **36**, 131-148 (2012).

8. Tan IS, Ramamurthi KS. Spore formation in Bacillus subtilis. *Environ Microbiol*
Rep **6**, 212-225 (2014).

9. Rand RP, Parsegian VA. Mimicry and mechanism in phospholipid models of
membrane fusion. *Annu Rev Physiol* **48**, 201-212 (1986).

10. Wong JY, Park CK, Seitz M, Israelachvili J. Polymer-cushioned bilayers. II. An
investigation of interaction forces and fusion using the surface forces apparatus.
Biophys J **77**, 1458-1468 (1999).

11. Kozlovsky Y, Kozlov MM. Membrane fission: model for intermediate structures.
Biophys J **85**, 85-96 (2003).

12. Bashkirov PV, Akimov SA, Evseev AI, Schmid SL, Zimmerberg J, Frolov VA.
GTPase cycle of dynamin is coupled to membrane squeeze and release, leading
to spontaneous fission. *Cell* **135**, 1276-1286 (2008).

13. Kozlov MM, McMahon HT, Chernomordik LV. Protein-driven membrane stresses
in fusion and fission. *Trends Biochem Sci* **35**, 699-706 (2010).

14. Ferguson SM, De Camilli P. Dynamin, a membrane-remodelling GTPase. *Nat*
Rev Mol Cell Biol **13**, 75-88 (2012).

15. Schoneberg J, Lee IH, Iwasa JH, Hurley JH. Reverse-topology membrane
scission by the ESCRT proteins. *Nat Rev Mol Cell Biol* **18**, 5-17 (2017).

16. Simunovic M*, et al.* Friction Mediates Scission of Tubular Membranes Scaffolded
by BAR Proteins. *Cell* **170**, 172-184 e111 (2017).

17. Roux A, Cuvelier D, Nassoy P, Prost J, Bassereau P, Goud B. Role of curvature
and phase transition in lipid sorting and fission of membrane tubules. *EMBO J*
24, 1537-1545 (2005).

18. Hatch AL, Gurel PS, Higgs HN. Novel roles for actin in mitochondrial fission. *J*
Cell Sci **127**, 4549-4560 (2014).

19. Yang C, Svitkina TM. Ultrastructure and dynamics of the actin-myosin II
cytoskeleton during mitochondrial fission. *Nat Cell Biol* **21**, 603-613 (2019).

20. Lacy MM, Ma R, Ravindra NG, Berro J. Molecular mechanisms of force
production in clathrin-mediated endocytosis. *FEBS Lett* **592**, 3586-3605 (2018).

21. Nickaeen M, Berro J, Pollard TD, Slepchenko BM. Actin assembly produces
sufficient forces for endocytosis in yeast. *Mol Biol Cell* **30**, 2014-2024 (2019).

22. Snead WT*, et al.* BAR scaffolds drive membrane fission by crowding disordered
domains. *J Cell Biol* **218**, 664-682 (2019).

23. Doan T*, et al.* FisB mediates membrane fission during sporulation in Bacillus
subtilis. *Genes Dev* **27**, 322-334 (2013).

24. Stragier P, Losick R. Molecular genetics of sporulation in Bacillus subtilis. *Annu*
Rev Genet **30**, 297-241 (1996).
25. Gest H, Mandelstam J. Longevity of microorganisms in natural environments.
Microbiol Sci **4**, 69-71 (1987).
26. Potts M. Desiccation tolerance of prokaryotes. *Microbiol Rev* **58**, 755-805 (1994).
27. Ulrich N, Nagler K, Laue M, Cockell CS, Setlow P, Moeller R. Experimental
studies addressing the longevity of Bacillus subtilis spores - The first data from a
500-year experiment. *PLoS One* **13**, e0208425 (2018).
28. Brown JK, Hovmøller MS. Aerial dispersal of pathogens on the global and
continental scales and its impact on plant disease. *Science* **297**, 537-541 (2002).
29. Eichenberger P*, et al.* The sigmaE regulon and the identification of additional
sporulation genes in Bacillus subtilis. *J Mol Biol* **327**, 945-972 (2003).
30. Mileykovskaya E, Dowhan W. Visualization of phospholipid domains in
Escherichia coli by using the cardiolipin-specific fluorescent dye 10-N-nonyl
acridine orange. *J Bacteriol* **182**, 1172-1175 (2000).
31. Kawai F, Shoda M, Harashima R, Sadaie Y, Hara H, Matsumoto K. Cardiolipin
domains in Bacillus subtilis marburg membranes. *J Bacteriol* **186**, 1475-1483
(2004).
32. Koppelman CM, Den Blaauwen T, Duursma MC, Heeren RM, Nanninga N.
Escherichia coli minicell membranes are enriched in cardiolipin. *J Bacteriol* **183**,
6144-6147 (2001).
33. Kawai F, Hara H, Takamatsu H, Watabe K, Matsumoto K. Cardiolipin enrichment
in spore membranes and its involvement in germination of Bacillus subtilis
Marburg. *Genes Genet Syst* **81**, 69-76 (2006).
34. Lewis RN, McElhaney RN. The physicochemical properties of cardiolipin bilayers
and cardiolipin-containing lipid membranes. *Biochim Biophys Acta* **1788**, 2069-
2079 (2009).
35. Haines TH. A new look at Cardiolipin. *Biochim Biophys Acta* **1788**, 1997-2002
(2009).
36. Ortiz A, Killian JA, Verkleij AJ, Wilschut J. Membrane fusion and the lamellar-to-
inverted-hexagonal phase transition in cardiolipin vesicle systems induced by
divalent cations. *Biophys J* **77**, 2003-2014 (1999).
37. Khalifat N, Puff N, Bonneau S, Fournier JB, Angelova MI. Membrane deformation
under local pH gradient: mimicking mitochondrial cristae dynamics. *Biophys J* **95**,
4924-4933 (2008).
38. Romantsov T, Helbig S, Culham DE, Gill C, Stalker L, Wood JM. Cardiolipin
promotes polar localization of osmosensory transporter ProP in Escherichia coli.
Mol Microbiol **64**, 1455-1465 (2007).
39. Romantsov T, Stalker L, Culham DE, Wood JM. Cardiolipin controls the osmotic
stress response and the subcellular location of transporter ProP in Escherichia
coli. *J Biol Chem* **283**, 12314-12323 (2008).
40. Lopez D, Koch G. Exploring functional membrane microdomains in bacteria: an
overview. *Curr Opin Microbiol* **36**, 76-84 (2017).
41. Sharp MD, Pogliano K. An in vivo membrane fusion assay implicates SpoIIIE in
the final stages of engulfment during Bacillus subtilis sporulation. *Proc Natl Acad*
Sci U S A **96**, 14553-14558 (1999).

42. Meyer P, Gutierrez J, Pogliano K, Dworkin J. Cell wall synthesis is necessary for
membrane dynamics during sporulation of Bacillus subtilis. *Mol Microbiol* **76**,
956-970 (2010).
43. Doan T*, et al.* Novel secretion apparatus maintains spore integrity and
developmental gene expression in Bacillus subtilis. *PLoS Genet* **5**, e1000566
(2009).
44. Williams ND*, et al.* DNA-Origami-Based Fluorescence Brightness Standards for
Convenient and Fast Protein Counting in Live Cells. *Nano Lett* **20**, 8890-8896
(2020).
45. Guiziou S*, et al.* A part toolbox to tune genetic expression in Bacillus subtilis.
Nucleic Acids Res **44**, 7495-7508 (2016).
46. Donovan C, Bramkamp M. Characterization and subcellular localization of a
bacterial flotillin homologue. *Microbiology* **155**, 1786-1799 (2009).
47. Kunst F*, et al.* The complete genome sequence of the Gram-positive bacterium
Bacillus subtilis. *Nature* **390**, 249-256 (1997).
48. Chernomordik LV, Kozlov MM. Mechanics of membrane fusion. *Nat Struct Mol*
Biol **15**, 675-683 (2008).
49. Schmid SL, Frolov VA. Dynamin: functional design of a membrane fission
catalyst. *Annu Rev Cell Dev Biol* **27**, 79-105 (2011).
50. Nishibori A, Kusaka J, Hara H, Umeda M, Matsumoto K.
Phosphatidylethanolamine Domains and Localization of Phospholipid Synthases
in *Bacillus subtilis* Membranes. *Journal of Bacteriology* **187**, 2163-2174 (2005).
51. Good MC, Zalatan JG, Lim WA. Scaffold proteins: hubs for controlling the flow of
cellular information. *Science* **332**, 680-686 (2011).
52. Langhorst MF, Reuter A, Stuermer CA. Scaffolding microdomains and beyond:
the function of reggie/flotillin proteins. *Cell Mol Life Sci* **62**, 2228-2240 (2005).
53. López D, Kolter R. Functional microdomains in bacterial membranes. *Genes Dev*
24, 1893-1902 (2010).
54. Sohlenkamp C, Geiger O. Bacterial membrane lipids: diversity in structures and
pathways. *FEMS Microbiology Reviews* **40**, 133-159 (2015).
55. Oliver PM, Crooks JA, Leidl M, Yoon EJ, Saghatelian A, Weibel DB. Localization
of anionic phospholipids in Escherichia coli cells. *J Bacteriol* **196**, 3386-3398
(2014).
56. Bogdanov M, Heacock PN, Dowhan W. Study of polytopic membrane protein
topological organization as a function of membrane lipid composition. *Methods*
Mol Biol **619**, 79-101 (2010).
57. Ovchinnikov S*, et al.* Protein structure determination using metagenome
sequence data. *Science* **355**, 294-298 (2017).
58. den Kamp JA, Redai I, van Deenen LL. Phospholipid composition of Bacillus
subtilis. *J Bacteriol* **99**, 298-303 (1969).
59. Updegrove TB, Ramamurthi KS. Geometric protein localization cues in bacterial
cells. *Curr Opin Microbiol* **36**, 7-13 (2017).
60. Simunovic M, Voth GA, Callan-Jones A, Bassereau P. When Physics Takes
Over: BAR Proteins and Membrane Curvature. *Trends in Cell Biology* **25**, 780-
792 (2015).

61. Antonny B. Mechanisms of membrane curvature sensing. *Annu Rev Biochem* **80**,
101-123 (2011).
62. Baumgart T, Capraro BR, Zhu C, Das SL. Thermodynamics and mechanics of
membrane curvature generation and sensing by proteins and lipids. *Annu Rev*
Phys Chem **62**, 483-506 (2011).
63. Sorre B*, et al.* Nature of curvature coupling of amphiphysin with membranes
depends on its bound density. *Proc Natl Acad Sci U S A* **109**, 173-178 (2012).
64. Kjaerulff O, Brodin L, Jung A. The structure and function of endophilin proteins.
Cell Biochem Biophys **60**, 137-154 (2011).
65. Chen Z, Zhu C, Kuo CJ, Robustelli J, Baumgart T. The N-Terminal Amphipathic
Helix of Endophilin Does Not Contribute to Its Molecular Curvature Generation
Capacity. *J Am Chem Soc* **138**, 14616-14622 (2016).
66. Hohendahl A*, et al.* Structural inhibition of dynamin-mediated membrane fission
by endophilin. *Elife* **6**, (2017).
67. Shi Z, Baumgart T. Membrane tension and peripheral protein density mediate
membrane shape transitions. *Nat Commun* **6**, 5974 (2015).
68. Handler AA, Lim JE, Losick R. Peptide inhibitor of cytokinesis during sporulation
in Bacillus subtilis. *Mol Microbiol* **68**, 588-599 (2008).
69. Renner LD, Eswaramoorthy P, Ramamurthi KS, Weibel DB. Studying
biomolecule localization by engineering bacterial cell wall curvature. *PLoS One* **8**,
e84143 (2013).
70. Canham PB. Minimum Energy of Bending as a Possible Explanation of
Biconcave Shape of Human Red Blood Cell. *J Theor Biol* **26**, 61-& (1970).
71. Helfrich W. Elastic Properties of Lipid Bilayers - Theory and Possible
Experiments. *Z Naturforsch C* **C 28**, 693-703 (1973).
72. Zhongcan OY, Helfrich W. Instability and Deformation of a Spherical Vesicle by
Pressure. *Phys Rev Lett* **59**, 2486-2488 (1987).
73. Zhongcan OY, Helfrich W. Bending Energy of Vesicle Membranes - General
Expressions for the 1st, 2nd, and 3rd Variation of the Shape Energy and
Applications to Spheres and Cylinders. *Phys Rev A* **39**, 5280-5288 (1989).
74. Seifert U, Langer SA. Viscous Modes of Fluid Bilayer-Membranes. *Europhys Lett*
23, 71-76 (1993).
75. Seifert U. Configurations of fluid membranes and vesicles. *Adv Phys* **46**, 13-137
(1997).
76. Ojkic N, López-Garrido J, Pogliano K, Endres RG. Cell-wall remodeling drives
engulfment during Bacillus subtilis sporulation. *Elife* **5**, (2016).
77. Hernandez JM, Kreutzberger AJ, Kiessling V, Tamm LK, Jahn R. Variable
cooperativity in SNARE-mediated membrane fusion. *Proc Natl Acad Sci U S A*
111, 12037-12042 (2014).
78. Mostafavi H*, et al.* Entropic forces drive self-organization and membrane fusion
by SNARE proteins. *Proc Natl Acad Sci U S A* **114**, 5455-5460 (2017).
79. Wu Z*, et al.* Dilation of fusion pores by crowding of SNARE proteins. *Elife* **6**,
(2017).
80. Stepanyants N, Macdonald PJ, Francy CA, Mears JA, Qi X, Ramachandran R.
Cardiolipin's propensity for phase transition and its reorganization by dynamin-

related protein 1 form a basis for mitochondrial membrane fission. *Mol Biol Cell*
26, 3104-3116 (2015).

81. Chernomordik LV, Kozlov MM, Melikyan GB, Abidor IG, Markin VS,
Chizmadzhev YA. The shape of lipid molecules and monolayer membrane
fusion. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **812**, 643-655
(1985).

82. Cullis PR, de Kruijff B, Verkleij AJ, Hope MJ. Lipid polymorphism and membrane
fusion. *Biochem Soc Trans* **14**, 242-245 (1986).

83. Landajuela A*, et al.* Lipid Geometry and Bilayer Curvature Modulate
LC3/GABARAP-Mediated Model Autophagosomal Elongation. *Biophysical*
journal **110**, 411-422 (2016).

84. Zeigler DR*, et al.* The origins of 168, W23, and other Bacillus subtilis legacy
strains. *J Bacteriol* **190**, 6983-6995 (2008).

85. Youngman PJ, Perkins JB, Losick R. Genetic transposition and insertional
mutagenesis in Bacillus subtilis with Streptococcus faecalis transposon Tn917.
Proc Natl Acad Sci U S A **80**, 2305-2309 (1983).

86. Spizizen J. TRANSFORMATION OF BIOCHEMICALLY DEFICIENT STRAINS
OF BACILLUS SUBTILIS BY DEOXYRIBONUCLEATE. *Proc Natl Acad Sci U S*
A **44**, 1072-1078 (1958).

87. Schaeffer P, Millet J, Aubert JP. Catabolic repression of bacterial sporulation.
Proc Natl Acad Sci U S A **54**, 704-711 (1965).

88. Sterlini JM, Mandelstam J. Commitment to sporulation in Bacillus subtilis and its
relationship to development of actinomycin resistance. *Biochem J* **113**, 29-37
(1969).

89. Koo BM*, et al.* Construction and Analysis of Two Genome-Scale Deletion
Libraries for Bacillus subtilis. *Cell Syst* **4**, 291-305.e297 (2017).

90. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* **9**, 671-675 (2012).

91. Bogdanov M, Zhang W, Xie J, Dowhan W. Transmembrane protein topology
mapping by the substituted cysteine accessibility method (SCAM(TM)):
application to lipid-specific membrane protein topogenesis. *Methods* **36**, 148-171
(2005).

92. Stratton BS*, et al.* Cholesterol Increases the Openness of SNARE-Mediated
Flickering Fusion Pores. *Biophys J* **110**, 1538-1550 (2016).

93. Nikolaus J, Karatekin E. SNARE-mediated Fusion of Single Proteoliposomes
with Tethered Supported Bilayers in a Microfluidic Flow Cell Monitored by
Polarized TIRF Microscopy. *J Vis Exp*, (2016).

94. Tinevez JY*, et al.* TrackMate: An open and extensible platform for single-particle
tracking. *Methods* **115**, 80-90 (2017).

95. Tarantino N*, et al.* TNF and IL-1 exhibit distinct ubiquitin requirements for
inducing NEMO-IKK supramolecular structures. *J Cell Biol* **204**, 231-245 (2014).

96. Huet S, Karatekin E, Tran VS, Fanget I, Cribier S, Henry JP. Analysis of transient
behavior in complex trajectories: application to secretory vesicle dynamics.
Biophys J **91**, 3542-3559 (2006).

97. Smith MB, Karatekin E, Gohlke A, Mizuno H, Watanabe N, Vavylonis D.
Interactive, computer-assisted tracking of speckle trajectories in fluorescence

1443   microscopy: application to actin polymerization and membrane fusion. *Biophys J*
1444   **101**, 1794-1804 (2011).

1445   98.   Lacombe C, Lubochinsky B. Specific extraction of bacterial cardiolipin from
1446   sporulating Bacillus subtilis. *Biochim Biophys Acta* **961**, 183-187 (1988).

1447   99.   Angelova MI, Dimitrov DS. Liposome Electroformation. *Faraday Discuss* **81**, 303-
1448   + (1986).

1449   100.   Connell E, Scott P, Davletov B. Real-time assay for monitoring membrane
1450   association of lipid-binding domains. *Anal Biochem* **377**, 83-88 (2008).

1451   101.   Sterlini JM, Mandelstam J. Commitment to Sporulation in Bacillus Subtilis and Its
1452   Relationship to Development of Actinomycin Resistance. *Biochem J* **113**, 29-+
1453   (1969).

1454   102.   Buser CA, Sigal CT, Resh MD, McLaughlin S. Membrane binding of myristylated
1455   peptides corresponding to the NH2 terminus of Src. *Biochemistry* **33**, 13093-
1456   13101 (1994).

1457   103.   Ducret A, Quardokus EM, Brun YV. MicrobeJ, a tool for high throughput bacterial
1458   cell detection and quantitative analysis. *Nat Microbiol* **1**, 16077 (2016).

1459   104.   Douglas SM, Dietz H, Liedl T, Högberg B, Graf F, Shih WM. Self-assembly of
1460   DNA into nanoscale three-dimensional shapes. *Nature* **459**, 414-418 (2009).

1461   105.   Stahl E, Martin TG, Praetorius F, Dietz H. Facile and scalable preparation of pure
1462   and dense DNA origami solutions. *Angew Chem Int Ed Engl* **53**, 12735-12740
1463   (2014).

1464   106.   Chen ZM, Atefi E, Baumgart T. Membrane Shape Instability Induced by Protein
1465   Crowding. *Biophysical Journal* **111**, 1823-1826 (2016).

1466   107.   Powers TR. Dynamics of filaments and membranes in a viscous fluid. *Rev Mod*
1467   *Phys* **82**, (2010).

1468   108.   Ojkic N, Lopez-Garrido J, Pogliano K, Endres RG. Bistable Forespore
1469   Engulfment in Bacillus subtilis by a Zipper Mechanism in Absence of the Cell
1470   Wall. *Plos Comput Biol* **10**, (2014).

1471

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

FIGURE 7

FIGURE 8

FIGURE 9