Processing of novel food reveal payoff and rank-biased social learning in a wild primate

C. Canteloup[1,2]*, M.B. Cera[1], B.J. Barrett[3,4,5]† & E. van de Waal[1,2]†

[1] Inkawu Vervet Project, Mawana Game Reserve, KwaZulu Natal, 3115, South Africa

[2] Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland

[3] Max Planck Institute for Animal Behavior, Department for the Ecology of Animal Societies,

Konstanz, Germany

[4] University of Konstanz, Department of Biology, Konstanz, Germany

[5] Max Planck Institute of Evolutionary Anthropology, Department of Human Behavior,

Ecology, and Culture, Leipzig, Germany

† These authors contributed equally to this work

*corresponding author: Charlotte Canteloup (charlotte.canteloup@gmail.com)

Abstract

Cultural complexity is strongly shaped by the efficiency by which new knowledge is

propagated. Different social learning strategies (SLS), where individuals biasedly learn

particular behaviors or from specific demonstrators, can contribute to an individual's success.

While SLS have been mostly studied in isolation, their interaction and the interplay between

individual and social learning is less understood. We performed a field-based open diffusion

experiment in a wild primate. We provided two groups of vervet monkeys with a novel food,

unshelled peanuts, and documented how three different peanut opening techniques spread

within the groups. We analyzed data using hierarchical Bayesian dynamic learning models that

explore the integration of multiple SLS with individual learning. We show that vervets

preferentially use the technique yielding the highest observed payoff, and also bias attention

toward individuals of higher rank. This shows that traditions may arise when individuals

integrate information about the efficiency of a behavior alongside cues related to the rank of a

demonstrator.

Introduction

Organisms may learn about their environment and acquire new skills individually, by

trial and error, or socially, by directly observing and copying others' behavior (1). Social

learning has been considered adaptive since organisms may gain fitness benefits when learning

from others, especially in a changing environment (2). Socially transmitted behaviors resulting

in animal cultural traditions have been described in diverse taxa and domains such as songbird

vocal dialects (3), tool use in crows (4), hunting techniques in whales (5), social conventions,

extractive foraging techniques and manipulation behaviors in numerous primates (6-8).

Social learning biases direct an individual's attention towards a behavior based on its

content, relative frequency in a population, or its association with a particular type of

demonstrator. These biases are typically, but not always, assumed to be shortcuts to acquiring

adaptive behavior and are theoretically well explored (10-12). Content-biases focus on

characteristics of the observed behavior such as a bias towards better payoff. Much theoretical

work has explored the evolution of payoff-biased learning (13, 14), which is useful when

payoffs are stochastic or adaptive behavior is rare. Empirical work in humans show payoff-bias

for its concurrent use with other social learning strategies (15). In some cases, payoff-biased

learning might be under-utilized by humans (16). Researchers have recently expanded their

investigation of payoff-biased learning to wild animals. Wild capuchins adopted the most

successful and efficient technique when processing novel fruits (17). Great tits do not use

payoff-biased learning but acquire a high payoff foraging technique through a combination of

positive frequency-dependent learning and individual learning that is highly sensitive to

personal experiences in payoffs (18). Male vervet monkeys, but not females, used the technique

displayed by male demonstrators more when a difference in payoff was introduced with males

receiving more food than females (19). The prevalence of payoff-biased learning is likely

underestimated in wild populations particularly if adaptive behavior is common, easy to

innovate, or if there is a single optimal solution (17).

Context biases focus on other cues, such as their frequency in a population (e.g.

preferentially use the most common behavior or the one exhibited by the majority of

individuals: conformity bias) or particular traits of models (e.g. preferentially use the behavior

displayed by familiar, prestigious, related, or similarly aged individuals: (20)). Such biases have

been reported in various animal species. "Moss sponging" by chimpanzees (21) and "sponging"

by dolphins (22) were both transmitted within matrilines; "Shelling", another foraging

technique, spread non-vertically among associated dolphins (23). Younger female guppies

chose the mate choice of older females (24). Capuchins and chimpanzees used the foraging

method displayed by older individuals (17, 25). In female-philopatric vervet monkeys, females

were preferentially chosen over males as demonstrators (26), but demonstrators were all of

high-rank. The techniques used by higher-ranked individuals were preferentially chosen by

chimpanzees (27) and vervet monkeys (28) when researchers simultaneously tested for several

biases. Animals also conform to the most frequently observed behavior: in vervet monkeys,

dispersing males conformed to the local foraging norm, abandoning their native preference

(29); great tits and chimpanzees disproportionately adopted the most frequent local foraging

technique (30, 31 but see 32); nine-spined sticklebacks preferentially chose the feeder favored

by the majority of individuals, even when this feeder was less rewarding than an alternative

(33). Fruit flies chose their mating partner according to the preference of the majority of

copulating conspecifics (34). All these findings suggest that different contextual characteristics

strongly influence social learning patterns. However, relatively little research has investigated

how organisms jointly integrate multiple social learning strategies with individual experience.

Open diffusion experiments with experimentally trained demonstrators in the wild have

been well utilized (18, 29, 30), and show what types of social learning animals are capable of

in a well-defined experimental context. However, open diffusion experiments without

experimentally trained demonstrators may better show how traditions endogenously arise in the

wild in more realistic social settings; yet, they are rare (17, 28).

Recently developed dynamic statistical models including Network Based Diffusion

Analysis (hereafter NBDA: (35, 36) and Experience-Weighted Attraction models (hereafter

EWA: (15, 17, 37)) are powerful statistical modeling approaches to identify evidence for social

learning and the pathways of social transmission. NBDA is useful to identify an evidence for

social learning and the typical pathways of social transmission but it only analyzes the initial

spread of a new behavior, i.e. the origin of a cultural tradition. It cannot thus evaluate what

social learning strategies individuals use to inform behavioral choice once they have learned

multiple options (36). Nor can NBDA examine how individuals uniquely combine individual

and social information – the combination of which makes social learning adaptive. EWA

models analyze the entire unique behavioral sequence chosen by each individual and investigate

how this is influenced over time by both personal experience and observation of others, i.e. the

origin and maintenance of a cultural tradition (15, 17). EWA models permit extant theoretical

models of social learning strategies to be turned into statistical models to examine which

strategy, or combination of strategies, best predict the behavior of individuals. These

predictions are conditional upon an individual's unique personal experiences and opportunities

for observing social information, thus linking individual variation in behavior to population-level cultural dynamics.

This study aims to determine which social learning strategies – content-biased: frequency-dependent; payoff-biased and/or context biased: based on demonstrator specificities (sex, kin and rank) - are responsible for the transmission of novel food processing techniques - peanut shelling - in wild vervet monkeys (*Chlorocebus pygerythrus*). Here, we offered a novel food – unshelled peanuts – to two groups of wild vervet monkeys: Noha (hereafter 'NH') and Kubu (hereafter 'KB') during four months of field experiments. We recorded the exact time of each peanut consumption event and the identity of observers at each time to construct a dynamic observation network. We described the three different peanut extraction techniques used by monkeys: crack with the hand (hereafter 'CH'; Movie S1: https://youtu.be/BkiTJyxp3gQ); crack with the mouth from the side of the peanut (hereafter 'CMS'; Movie S2: https://youtu.be/3JOAH4khnjk) and crack with the mouth from the top of the peanut (hereafter 'CMT'; Movie S3: : https://youtu.be/ikcFeAjVZqI). We then analyzed data from this experiment with EWA models. We predict, in accordance with another study focusing on context biases (28), that higher-ranked vervet monkeys are more influential demonstrators than others. We hypothesize that other content biases, such as frequency or payoff based (i.e. the most successful technique), would also be possible. The major interest of this study remains in assessing the weight of each social learning strategy in a single analysis combining individual and social information through the use of a field open diffusion experiment with no trained model.

Results

Effects of group, rank, sex and age on success, manipulation and observation

In NH, 25 individuals (9 adults; 12 juveniles; 4 infants) succeeded in opening 2104 peanuts and attempted to open a further 1401 peanuts. In KB, 9 individuals (5 juveniles; 4 infants) succeeded in opening 45 peanuts and 10 individuals (6 juveniles; 4 infants) attempted to open 102 peanuts (Table S1). We found an effect on rank and sex both on the rate of peanut opening successes and on the rate of manipulation. Higher rankers succeeded significantly more to extract peanuts (GLMs: estimate=-2.67; standard error=0.67; t value=-4.00; P=0.0004) and manipulated more peanuts than lower rankers (estimate=-2.22; standard error=0.65; t value=-3.56; P=0.002). Males succeeded significantly more to extract peanuts (18/24 males; 16/29 females; estimate=1.12; standard error=0.32; t value=3.56; P=0.001) and manipulated significantly more peanuts than females (19/24 males; 16/29 females; estimate=0.93; standard error=0.32; t value=2.95; P=0.007).

The latency of first success ranged from 2.03 minutes (122s) to 448.1 minutes (26,886s) in NH (mean=218.35 min (13,101.52s); standard error of the mean=25.29 min (1517.54s)) and from 78.95 min (4737s) to 393.6 min (23,616s) in KB (mean=223.33 min (13,400.89s); standard error of the mean=38.85 min (2330.88s); Table S1). The innovator in NH was a low-ranked, newly immigrated adult male at the first exposure to peanuts (Avo) while, in KB, it was an infant male (Aar) at the third exposure. We found a significant effect of age on latency of first peanut opening success. Juveniles had a lower latency compared to infants (GLMs: estimate=-0.57; standard error=0.25; t value=-2.34; P=0.05), but we found no significant difference between infants and adults (estimate=0.13; standard error=0.31; t value=0.41; P=0.91) nor between juveniles and adults (estimate=-0.45; standard error=0.25; t value=-1.82; P=0.16). We found no significant difference between groups (estimate=-0.16; standard error=0.24; t value=-0.67; P=0.51), between males and females (estimate=0.12; standard

error=0.21; t value=0.57; P=0.58) and no significant effect of rank (estimate=-0.11; standard

error=0.42; t value=-0.26; P=0.80) on latency of first success.

We found no clear difference in the rate of observation of successes between groups

(GLMs: estimate=2.79; standard error=1.66; t value=1.68; P=0.10), between high rankers and

low rankers (estimate=-1.64; standard error=0.86; t value=-1.91; P=0.07), between adults and

infants (estimate=0.64; standard error=0.59; t value=1.08; P=0.29), between adults and

juveniles (estimate=-0.007; standard error=0.42; t value=-0.017; P=0.99) nor between males

and females (estimate=0.81; standard error=0.44; t value=1.85; P=0.08) . We found no evidence

for any difference on the rate of observation of manipulations between groups (estimate=2.42;

standard error=1.36; t value=1.78; P=0.09), between high and low rankers (estimate=-1.56;

standard error=0.83; t value=-1.89; P=0.07), between adults and infants (estimate=0.57;

standard error=0.58; t value=0.98; P=0.33), between adults and juveniles (estimate=-0.01;

standard error=0.41; t value=-0.03; P=0.98) and between males and females (estimate=0.79;

standard error=0.42; t value=1.88; P=0.07).

We found effects of rank, age and sex on the rate of being observed when manipulating

and successfully opening peanuts. High-rankers were observed significantly more than low-

rankers when succeeding (GLMs: estimate=-3.61; standard error=0.65; t value=-5.57; P<0.001)

and manipulating (estimate=-3.31; standard error=0.65; t value=-5.08; P<0.001). Adults were

observed significantly more than juveniles when succeeding (estimate=-0.91; standard

error=0.29; t value=-3.12; P=0.004) and manipulating (estimate=-0.73; standard error=0.29; t

value=-2.56; P=0.02). Males were observed significantly more than females when successfully

opening (estimate=1.15; standard error=0.27; t value=4.20; P<0.001) and manipulating peanuts

(estimate=1.02; standard error=0.28; t value=3.63; P=0.001).

EWA model interpretations

For most individuals, and at the population level, we found that CMS was the most successful, and became the most common technique in the population. CMS was successful on 46.8% occasions in NH and 24.8% in KB over all manipulations. This was followed by CMT (8.1% in NH, 5.4% success in KB) and CH (5.1% in NH, 0.7% in KB). Overall, we found the best support for the global model which receives the most weight and has the lowest WAIC value (Support 100%; Table 1) and predicts well at the population-level (Fig. 1) and across all individuals (Fig. 2; Fig. S1). Model predictions from the global model show that the highest-payoff behavior became the most common for nearly all individuals in the population (Fig. 2; Fig. 3 and Fig. S1). The parameters' magnitudes and certainties in the global model, as well as WAIC score and parameter estimates in individual models, suggest that payoff-bias, followed by rank-bias, were the two singular most important learning strategies (Table 1; Table 2). Figure 3 shows predictions for five evaluated models for a single individual ('Gran' in NH). In this case, all models allow the highest payoff behavior (CMS; displayed in purple) to become the most common in the population. The global model tends to show more uncertainty around any particular prediction, whereas models evaluating pure individual learning strategies tend to be overconfident in their estimates, less accurate, or do not perform well for all individuals in the population. Payoff-biased and rank-biased learning models perform well, but do not adequately predict the frequency of all three behaviors across the entire time series, whereas frequency-dependent learning over-predicts the frequency of the most common, high-payoff behavior. The individual learning model's predictions are overconfident, tend to drift around, and underestimate the frequency of CMS, particularly at the beginning of the time series. We found little support that individuals preferentially used the technique displayed by females or individuals of the same sex (Table 1). Kin-biased learning might be important for a few, but not most individuals. We encourage readers to examine Figure S1, as we believe it is the most informative about model fit compared to real data and is informative about individual variation.

Variation between individuals and groups in parameters

We see considerable variation between the two groups and among individuals for many varying effects parameters in these models, this occurred for several reasons. In KB, fewer individuals participated in the experiment and they had fewer foraging bouts generating more uncertain estimates as reflected by the population posterior mean estimates in Table 2. They were also less likely to observe social information, based on the 20-minute time windows we selected in this analysis. For these reasons, many of the individual-level parameter estimates of varying effects have tighter 89% High Posterior Density Intervals than the population mean estimates. Posterior estimates of group-level varying effects also differ greatly for some parameters. These estimates are more precise and parameter values related to social learning are higher for NH due to the greater number of individuals who participated to the experiment. This can be visualized when looking at the shaded HPDIs around model predictions in Figure S1 for individuals in both groups of different sample sizes. It is also visualized in Figures S2-S4. Estimates of σ extracted from the variance-covariance matrix of the global model are in Figure S5.

Sensitivity to attraction scores (λ)

Females (Posterior Mean (λ_f=8.38; [89% HPDI: 4.01-14.86]) are more sensitive than males (λ_m=6.27 [3.35-10.35]) to differences in attraction scores while juveniles are more sensitive than adults (λ_j=10.04 [5.45-17.07]; λ_a=5.22 [2.64-9.03]). A higher attraction score typically means that individuals are more sensitive to personally-experienced differences of behavioral payoffs. The direction of these effects is consistent across nearly all evaluated models (Table 2). We see considerable variation in varying effects of λ at the group and individual level (Fig. S2A), but much of this is due to sampling differences.

Weight given to recent experiences (ϕ)

Overall, we see small values for ϕ, suggesting that individuals tend to weight past experience more heavily than recent experience; memory strongly influences behavior. Adults tend to weight new experiences more heavily than juveniles (ϕ_a=0.08 [0.02-0.17]; ϕ_j=0.04 [0.01-0.08]). We see little support for sex differences (ϕ_f =0.06 [0.02-0.12]; ϕ_m=0.05 [0.01-0.11]). The direction and magnitude of these effects are consistent across all evaluated models (Table 2). Groups and individuals (particularly in KB) tend to have larger and more uncertain estimates of ϕ as a wider range of parameter values can predict the observed data (Fig. S2B).

Weight given to social information (γ)

In the global model, social information influences behavior slightly more heavily for adults than juveniles (γ_a=0.23 [0.05-0.48]; γ_j=0.20 [0.05-0.38]), and for females than males (γ_f =0.23 [0.05-0.43]; γ_m=0.20 [0.04-0.43]). However, these patterns are not consistent across all models. This is because for each individual model, one learning bias might be particularly salient for a particular age or sex class. We can gain insight into this from comparing predictions from the global model to models representing individual learning strategies (Fig. S2C). For example, the payoff-bias learning model has higher values of γ for juveniles and females (γ_j=0.31 [0.13-0.50] ; γ_a=0.28 [0.11-0.49]; γ_f =0.30 [0.13-0.51]; γ_m=0.28 [0.11-0.49]), while the rank-bias learning model has higher values for of γ for adults and males (γ_a=0.29 [0.12-0.50]; γ_j=0.18 [0.07-0.33]; γ_m=0.27 [0.11-0.47]; γ_f =0.20 [0.08-0.37]).

Payoff-bias

We find reliable support for payoff-biased learning, for nearly all individuals in the population (Fig. S3A). Of the social learning biases, we also see the most salient differences

between age and sex classes. Juveniles rely more on payoff-biased social learning than adults (βpay_{j}=1.24 [-0.36-2.76]; βpay_{j}=0.45 [-1.06-1.97]), while females rely more heavily on it than males (βpay_{f}=1.07 [-0.49-2.63]; βpay_{m}=0.62 [-0.88-2.08]). Of the EWA models representing individual learning strategies, payoff-biased learning had the lowest WAIC value (Table 1) and its predictions were most similar to the global model for most individuals in the population. Similar differences and stronger effect sizes were estimated in the payoff-biased learning model when compared to the global model (Table 2).

Rank-bias

We also see good support for rank-biased social learning for most individuals in the population, although there is more uncertainty around these predictions (Fig. S3B). The global model suggests that adults are more likely to learn from high ranking individuals than juveniles ($\beta rank_{a}$=1.26 [-0.25,2.72] ; $\beta rank_{j}$=0.80 [-0.85,2.53]), and females are more likely to learn from high-ranking individuals than males ($\beta rank_{f}$=1.18 [-0.40,2.75]; $\beta rank_{m}$=0.89 [-0.60,2.48]). These differences between age and sex parameters are much smaller in the rank-biased learning model (Table 2). This is likely because higher ranking individuals are more successful than low ranking individuals. When including both rank and payoff in the global models, differences in reliance on rank cues, instead of payoff cues, emerge.

Frequency-dependence

On average, the population tends to be somewhere between linear and slightly positive-frequency-dependent. We see posterior means slightly larger than 1 for most individuals (Fig. S4). However, boundary conditions (f^{c}>0) often make posterior point estimates misleading; nearly half of the posterior mass for many individuals and all age and sex classes lies in a parameter space consistent with anti-conformity. The frequency-dependent learning model

estimated a stronger degree of conformist transmission, but it often under-predicts the

probability of an individual choosing a less common behavior.

Matrilineal kin-, Female- and Same-sex biased learning

We see some evidence for matrilineal kin-biased learning (Table 2), but this might only

be for some individuals in the population (Fig. S3C). We see little evidence for sex-biased and

female-biased transmission (Table 2; Fig. S3D; Fig. S3E).

Discussion

In the present study we used a time-series model predicting each technique's probability

of being chosen in order to infer which social learning strategies were utilized in two groups of

wild vervet monkeys when learning to process unshelled peanuts, a novel food. The best

supported statistical model, and evidence from a suite of models, suggests that vervet monkeys

used the most successful, highest payoff technique, and displayed by the higher ranked

demonstrators. We found weak support for a positive frequency-dependent conformity bias

defined as 'use the most observed technique'. We also found no evidence that they show a bias

toward the technique displayed by females, individuals of the same sex, or related individuals.

They do not ignore social information and also use individual learning. We reported high

variation between groups and among individuals for many parameters in the models. This is

due to the difference in data sample size between the two groups; one group (NH) with many

more participants eating peanuts than the other (KB). Finally, we reported effects of age, sex

and rank on individual and social learning which are discussed in the remainder of the

discussion.

Note that we avoided using the term "copying" across the manuscript which is strongly

associated to "imitation" by some scholars (38, 39), although not by all (9, 40). Here, we do not

make any assumption and conclusion regarding the social learning mechanisms (action form

copying versus non-form copying/socially mediated individual learning: 39) implied in the

acquisition of the peanuts processing techniques. We acknowledge that the three peanuts

processing techniques we describe are within the Zone of Latent Solution of vervet monkeys

(39) and we do not claim for form copying, i.e. that vervet monkeys faithfully copied the exact

movements – the form of the behavior - displayed by their conspecifics.

Wild vervet monkeys acquire novel food processing technique via payoff-biased learning

We found the strongest support for the global model suggesting that vervet monkeys

use multiple social learning strategies simultaneously. Here, payoff- and rank-biased learning

guided diffusion of peanut processing techniques in both groups of wild vervet monkeys. These

results do not fully align with those of a previous study on vervet monkeys at our field site (19).

In Bono and colleagues' study, payoff consisted of a different quantity of food provided by a

two-action design – one item versus five. In our study, socially observed payoffs were an

individual's mean observed probability of a technique being successful in the previous 20

minutes. Bono et al. found that only males chose the technique used by the male model more

when it received five food items compared to a female model receiving only one item (19). We

find that females and juveniles relied more on payoff-biased social learning than males and

adults conversely. We believe this discrepancy in results can be explained by methodological

differences between both studies. First, in Bono and colleagues' study (19), only two models,

a male and a female, that were both adults and of high rank, were available in each group, while

in our study, numerous individuals of both sexes and various ages and ranks could

opportunistically serve as demonstrators. Second, we tested here behavioral responses of

individuals towards a novel and unknown food while Bono and colleagues (19) tested monkeys

in a two-action box experiment filled with known food. It is possible that monkeys got

accustomed to boxes' affordance across the numerous experiments of this kind they had

experienced in previous years at the Inkawu Vervet Project. Different strategies could be at play

in different environmental contexts, i.e. in a usual foraging situation *versus* an unprecedented

one. Our results suggest thus that females - the philopatric sex – are maybe less conservative in

their social learning than previously thought and that their conservatism decreases in a novel

context after an environmental change. When it has been suggested that females would stick to

their habits compared to males who are expected to show more behavioral flexibility (19), our

findings challenge this hypothesis and filled a gap. Females possibly possess more detailed

knowledge than males about the distribution of existing food resources or objects such as boxes

they encounter in their territory, which could make them a valuable source of directed social

learning in these common situations. However, we suggest that when facing an unknown

foraging situation, females can show flexibility, decrease their conservatism and rely more than

males on the social information available.

Other recent studies used a similar multilevel dynamic learning model in great tits (18)

and in capuchins (17). Great tits have been found to switch to an alternative higher payoff

variant when given a choice between a low and a high payoff option (18). As we found in vervet

monkeys, young birds were more likely to use social information than older birds (18). Barrett

and colleagues (17) offered *Sterculia apetala* fruits to a naïve group of wild capuchins and

followed the spread of the processing techniques. They found that, of multiple biases, payoff-

biased social learning was the one responsible for the spread of the most successful techniques

(17), and that age was a strong predictor in the heterogeneity of learning among individuals.

We can hypothesize that this maintained preference for CMS technique could be at the origin

of cultural transmission of peanut processing techniques if this food was made available on the

long term in vervet monkeys' environment.

Rank also matters: vervet monkeys preferentially used the technique displayed by higher

rankers

Here, we also report a rank bias in agreement with the 'use preferentially the behavior

displayed by higher-ranked individuals' strategy found in chimpanzees (27) and vervet

monkeys (28). In this latter study, we used another modeling approach – NBDA - to identify

the typical pathways of transmission of boxes opening techniques in an open diffusion

experiment. We tested whether the diffusion followed specific social networks representing

different pathways of learning such as learning from females, older individuals, related

individuals or higher rankers. We discovered evidence of a transmission bias favoring learning

from higher-ranked individuals, with no evidence for age, sex or kin bias (28). Present results

are thus in accordance with these previous findings. We could not in this previous study test for

frequency-dependent learning and payoff-biased learning nor for the interplay between the use

of social and individual information, but this is precisely what we assessed here. This means

that higher rankers are not only considered as models for the first learning event but that this

bias is maintained across time.

While higher rankers were manipulating more boxes and were more successful, they

were not observed more than low rankers in our previous study (28). Instead, observations of

higher-ranked individuals had a greater effect on observers' behavior than observations of

lower-ranked individuals. Here, we found that higher rankers manipulated and successfully

processed more peanuts, probably because they can monopolize access to food. We did not find

any difference in observation rate between individuals but, we reported that higher rankers,

adults and males were observed more than lower rankers, juveniles and females. This difference

might be due to experimental novelty arousing a different attentional pattern. Indeed, in a box experiment, vervet monkeys showed a selective attention to female models (26) when all models were of high rank. This is in accordance with literature on chimpanzees and capuchins in which older and more proficient nut crackers are more observed than others (41, 42).

No clear evidence of conformity, kin or sex biased learning

We tested for frequency-dependent learning with a model estimating whether individuals used i) positively frequency dependence as conformity bias; ii) unbiased transmission/copy proportional to frequency and iii) negative frequency dependence/anti-conformity- bias toward copying rarer traits. We found little evidence for any frequency-dependent learning, our population being slightly positive frequency-dependent. A previous study on vervet monkeys reported that migrant males conformed to their new group norm by abandoning their natal feeding preference, thus following a functional definition of conformity (29). The authors' claim of conformity (29) has been considered a premature conclusion regarding the mechanism of conformity, as the authors did not have the observational data of who the males attended to before switching and could not prove that the males observed the foraging choice of a majority before switching (43). Here, we did not test for such a conformity bias based on the number of individuals using the same processing technique, but we rather tested for a conformity bias based on the frequency of technique used. We thus tested whether individuals were i) more likely to use a technique frequently observed; ii) as likely to use a technique as it is observed or iii) less likely to use it. Such biases have been tested in great tits where the establishment of a foraging tradition has been found to rely on positive frequency-dependence; individuals adopting the most frequent local variant (30). Barrett and colleagues (18) found no evidence for a conformity bias but they found evidence of weak anti-conformity meaning that rare techniques attracted more attention in capuchins.

We found no evidence for kin- or sex-biased social learning. Previous studies reported

that infant vervet monkeys ate the same food as their mother (29) and that females were

preferred as models over males (26). However, these studies tested for a single bias at a time

while we tested here for several biases simultaneously, weighting each strategy against the other

which makes the analysis more powerful. Our results are furthermore consistent with a previous

study testing for several model biases in a single analysis that did not identify neither any such

biases (28). We believe that testing for several biases in a single analysis allow to disentangle

between potential confounding factors such as age, sex and rank and can highlight instances of

equifinality where multiple social learning strategies produce the same signature in a population

(44).

Sociodemographic effects on individual variation in social and individual learning

In the present study, the two innovators were an infant and a low-ranking adult male. In

another open diffusion experiment (28), the alpha male and female were the two innovators in

the same two groups when tested one year before. We believe that this difference can be linked

to the aforementioned experiments' methodological considerations. In Canteloup and

colleagues' study (28), groups were tested in a two-action box experiment for which individuals

may get accustomed to boxes' affordances despite the use of new design. Some studies

suggested that juveniles and low rankers are often innovators (45, 46) although this depends on

the type of behavior (47), while other studies suggest neophilia and boldness have been found

to better predict innovation (48). We hypothesize that innovators' identity also depends on task

novelty. Monkeys being neophobic, an unknown food can be considered as a source of stress,

which suggests that low rankers were the initiators and that juveniles were quicker to first

succeed to extract peanuts because of their usual limited access to resources. When the task is

not completely new, dominants are more expected to be the innovators.

We reported that, in the global model, social information influenced behavior more

heavily in adults than juveniles and in females than males, but these patterns were not consistent

across all models. In the payoff model, social information influenced behavior more heavily for

juveniles and females whereas in the rank model, adults and males relied more on social

information. Despite being in disagreement with what has been previously found in vervet

monkeys (19), this suggests that juveniles and females would be more sensitive to behaviors'

payoffs while adults and males would be more sensitive to rank cues associated with a

demonstrator. Overall, we found that adults weighed past experience more heavily than recent

experience compared to juveniles, suggesting that memory strongly influences their behavior

as previously found in birds and capuchins (17, 18).

Conclusions

We show that payoff bias could be at the origin of the spread of novel food processing

techniques in vervet monkeys, and that this content bias likely acts concurrently with rank-

biased learning. EWA models consider exactly what and whom individuals had uniquely

observed both prior to their observing a successful solution and after they learn how to solve

the task. This detailed time-varying modeling approach makes the analysis particularly

powerful in uncovering the interplay between personal and social information use, which makes

social learning adaptive and efficient. Because it is very rare that first innovation event and its

spread are recorded through observational studies, we believe that controlled experiments

coupled with dynamic, theory-informed modeling are a good approach to track the diffusion of

new behaviors and to disentangle between different social learning strategies.

Materials and Methods

Experimental model and subject details

We studied two groups of wild vervet monkeys (*Chlorocebus pygerythrus*): 'Noha'

(NH) and 'Kubu' (KB). NH was composed of 34 individuals (5 adult males; 9 adult females;

11 juveniles males; 9 juveniles females) and KB was composed of 19 individuals (1 adult male;

6 adult females; 6 juveniles males; 6 juveniles females; Table S1). Males were considered as

adults once they dispersed, and females were considered as adults after they gave their first

birth. Individuals that did not fulfill these criteria were considered as juveniles (28). Each group

had been habituated to the presence of human observers: since 2010 for NH and since 2013 for

KB. All individuals were identifiable thanks to portrait photographs and specific individual

body and face features (scars, colors, shape etc.).

Ethics guidelines: Our study adhered to the "Guidelines for the use of animals in

research" of Association for Study of Animal Behavior and was approved by the relevant local

authority, Ezemvelo KZN Wildlife, South Africa.

Study site

The study was conducted at the Inkawu Vervet Project (IVP) in a 12,000-hectare private

game reserve: Mawana (28°00.327S, 031°12.348E) in KwaZulu Natal province, South Africa.

The vegetation of the study site consisted in a savannah characterized by a mosaic of grasslands

and clusters of trees of the typical savannah thornveld, bushveld and thicket patches. Mawana

houses various species of animals including elephants, hippopotamus, giraffes, zebras and

numerous species of antelopes. The common predators of vervet monkeys consist of leopards,

hyenas, jackals, caracals, servals and several species of snakes and raptors.

Hierarchy establishment

Agonistic interactions (aggressor behavior: stare, chase, attack, hit, bite, take place; victim behavior: retreat, flee, leave, avoid, jump aside) were collected from May 2018 to October 2018, aside from experiment days, on all the adults and juveniles of both groups via *ad libitum* sampling (49) and food competition tests (i.e. corn provided to the whole group from a plastic box). Data were collected by CC, MBC and different observers from the IVP team. Before beginning data collection, observers had to pass an inter-observer reliability test with 80% reliability for each data category between two observers. Data were collected on tablets (Vodacom Smart Tab 2) equipped with the Pendragon version 8.

Individual hierarchical ranks were determined by the outcome of dyadic agonistic interactions recorded *ad libitum* and through food competition tests using Socprog software version 2.7 (50). Hierarchies in both groups were significantly linear (NH: h' = 0.27; P < 0.0001; KB: h' = 0.42; P < 0.0001) and ranks were assessed by I&SI method (51).

Open diffusion experiment

The experimental apparatus consisted of provisioning the group with two transparent rectangular plastic boxes (dimensions: 34x14x12 cm) containing unshelled peanuts in quantities (about 2 kg) to avoid a single individual monopolizing the boxes. The monkeys had never been provided with peanuts before the experiment and peanuts were not available in their environment. Thus, unshelled peanuts were a novel, nutritious food that required processing to be extracted from their shells before consumption.

Experiments took place at sunrise at monkeys' sleeping sites during the dry, food-scarce winter to maximize their motivation for novel food. The two boxes of peanuts were offered to the monkeys, spaced by about 1 to 10 meters from each other. CC led the experiment with the help of two to four field assistants. All monkeys were free to come to the boxes within the constraints of the social group dynamics. Experiments were video recorded using JVC cameras

(EverioR Quad Proof GZ-R430BE) to which the experimenter said aloud the identities of the

actor and of the attending neighbors for each manipulation event. A manipulation event was

defined either as an attempt to extract a peanut from its shell (i.e. the individual acted on the

peanut failing to fully open it and to get access to the food) or as a success (i.e. the individual

succeeded to fully open a shell and to extract the peanut from the shell). A conspecific was

considered as attending when it had its head or body oriented in an unobstructed line towards

the subject manipulating the peanut and was located within 0-30m from the actor. Several

individuals could thus be registered as attending to one or several actors simultaneously.

The open diffusion experiments ran from May 2018 to August 2018 to have a maximum

of individuals participating in the experiment. A total of 11 sessions of open diffusion

experiment were run in NH and 10 in KB. The average duration of an experimental session was

46:46 minutes for NH and 42:47 minutes for KB.

Video analysis

All video recordings were later analyzed by MBC with Media Player Classic Home

Cinema software version 1.7.11. Twenty percent of the video were analyzed by CC and inter-

observer reliability was substantial ($\kappa = 0.78$). During video analysis in slow motion or frame

by frame, the following variables were encoded: date, exact time of each manipulative event,

actor identity, the technique used (crack with hand: 'CH'; crack with mouth from the top of the

peanut: 'CMT'; crack with mouth from the side of the peanut: 'CMS'; see Movies S1-S3) and

the identity of attending individuals.

Quantification and statistical analysis

Following Barrett and colleagues (17), we used a suite of hierarchical experience-

weighted attraction (EWA) models to analyze data collected in the open-diffusion experiment.

EWA models are time-series models that evaluate the joint influence of personal experience

and social information on the probability of individual displaying a behavior (37). While

originally a tool in behavioral economics, they are being further developed and utilized in cross-

taxa studies of cultural transmission (16-18, 27, 52).

This analytical approach has several strengths: It permits evaluation of multiple

hypothesized learning strategies, utilizes a dynamic social learning network unique to each

individual, and links individual variation in behavior and cognition to population level-cultural

dynamics. Working with time-series of behavior unique to each individual is important as

population-level signatures can often be misleading (53) or exhibit equifinality, particularly if

individuals vary in experience, observation opportunities or the social learning strategies they

employ (44). The mathematical specification of our analytical approach also minimizes

ambiguity of what types of social learning we are evaluating. This is important as verbal

definitions are imprecise, and terminologies are differently interpreted in studies of social

learning. Most importantly, this approach links theory to data. Instead of using a theoretically

uninformed analytical approach to find results consistent with theory, we bypass quantitative

proxies and translate theoretical models to statistical models. We discuss the models in

mathematical detail in the supplemental material.

We fit a series of EWA models evaluating the following learning strategies:

1.   Individual learning

2.   Frequency-dependent learning (using preferentially behavior using information about

its frequency in population)

3.   Female-biased learning (bias toward using preferentially the technique displayed by

females in group i.e. matrilineal sex in vervets)

4.   Matrilineal kin-biased learning (bias towards using preferentially the technique

displayed by closely related individuals)

5.   Compare means payoff-biased learning (using preferentially most successful or efficient

behavior)

6.   Rank-biased learning (using preferentially the technique displayed by high-ranking

individual)

7.   Sex-biased learning (using preferentially behaviors of individuals that are the same sex

as you)

8.   Global model that includes 1-7.

555   For each behavioral choice, social information used by an actor was the average value

556   of each cue observed in a time window of 20 minutes prior to the observation (Table 3). As

557   individuals access social information at different timescales, and this window choice was

558   somewhat arbitrary, we also evaluated social info at 30, 10, and 5-minute timescales. These

559   analyses yielded similar results and were robust to time windows.

560   We ran the models using regularizing priors and a Cholesky decomposition for

561   estimating varying effects. They were fit using RStan version 2.19.3 (54). Models were

562   compared using widely applicable information criteria (WAIC). Reproducible model code and

563   data can he found here: https://github.com/bjbarrett/vervet_peanut_EWA

565   EWA Model Specification

566   EWA models have two parts: a set of expressions that specify how individuals

567   accumulate experience and a second set of expressions that specify the probability of each

568   option being chosen. Accumulated experience is represented by *attraction scores*, $A_{ij,t}$, unique

569   to each behavior i, individual j, and time t. We update $A_{ij,t}$ with an observed payoff $\pi_{ij,t}$:

570             $$A_{ij,t+1} \quad (1 - \phi_j)A_{ij,t} + \phi_j\pi_{ij,t}$$

The parameter ϕ_j controls the importance of recent payoffs in influencing attraction scores.

When ϕ_j is high, more weight is given to recent experience over past experiences – memory

has less of an influence on behavioral choice. This parameter is unique to an individual j, and

we also estimate how it varies by age-class and sex.

Attraction scores are converted into probabilities of behavioral choice with a standard

multinomial logistic choice rule:

577
$$Pr\left(i \vee A_{ijt}, \lambda_j\right) \quad \frac{exp\left(\lambda_j A_{ij,t}\right)}{\sum_k exp\left(\lambda_j A_{kj,t}\right)} = I_{ij}$$

λ_j controls sensitivity to differences in attraction scores on behavioral choice and is unique to

an individual j. A very large λ_j ,means the option with the largest attraction score is nearly

always selected. Choice is random with respect to attraction score when $\lambda_j = 0$. Individuals

were assigned a payoff of zero, $\pi_{ij,t} = 0$, if they failed to open a peanut. If they were successful

$\pi_{ij,t} = 1$.

Social learning may directly influence choice distinctly from individual learning. $S_{ij} =$

$S\left(i \vee \Theta_j\right)$ is the probability an individual j chooses behaviour i on the basis of a set of social

cues and parameters Θ_j. Behavioral choice is a convex combination specified by:

586
$$Pr\left(i \vee A_{ij,t}, \Theta_j\right) = \left(1 - \gamma_j\right)I_{ij,t} + \gamma_j S_{ij,t}$$

587   where γ_j is the weight assigned to social cues bounded by 0 and 1.

588   Social cues are incorporated into $S_{ij,t}$ by use of a multinomial probability expression with a

589   log-linear component $B_{ij,t}$ that is an additive combination of cue frequencies. The probability

590   of displaying each behavior i, solely as a function of social cues, is:

$$591 \qquad S_{ij,t} \quad \frac{N_{ij,t}^{f_c} expB_{ij,t}}{\sum_m N_{mj,t}^{f_c} expB_{mj,t}}$$

$N_{ij,t}$ are the observed frequencies of each technique i at time t by individual j. The

exponentiated parameter f_c controls the amount and type of frequency dependence. When $f_c =$

1, social learning is unbiased by frequency and techniques influence choice in proportion to

their occurrence (sometimes referred to as unbiased transmission). When $f > 1$, social learning

is positive frequency-dependent or conformist. When $f < 1$, social learning is negative

frequency-dependent, and a bias is shown towards rare behaviors.

Other social cues associate with individuals (i.e. rank, age, or relatedness) or behaviors (i.e

payoffs), are incorporated via:

$$600 \qquad B_{ijt} \quad \sum_k \beta_k$$

B_{ijt} is the sum of the products of the influence parameters β_k and the cue values $\kappa_{k,ijt}$. Cues

evaluated in these models are explained in detail in Table 3.

Other statistics

We used generalized linear models (GLMs) with quasi-Poisson error and log link

function to test for the effect of group, normalized rank, sex and age on i) the latency of first

success of shelling a peanut; ii) the number of peanut shelling successes; iii) the number of

peanuts manipulation (attempts + successes); iv) the number of successes observed; v) the

number of manipulations observed; vi) the number of times being observed when succeeding

and vii) the number of times being observed when manipulating. We added the log of the time

each individual had to process peanuts once they first succeeded as an offset (a standard

statistical technique for converting a Poisson GLM for analysis of counts into a model for analyzing counts per unit of time). All tests were performed with package lme4.

Acknowledgments: We thank Arend van Blerk and the whole IVP team for their help and support in the field. We are also particularly thankful to Loïc Brun, Adwait Deshpande, Manon Kerréveur-Lavaud, Maria Teresa Martinez Navarrete, Anaïs Rilly and Claudia Seminara for their assistance in data collection. We are grateful to the van der Walt family for their permission to conduct the study on their land. We warmly thank Matthias Wubs for writing us some R scripts to extract social data as matrices. We greatly thank Rachel Harrison for her comments on the manuscript. **Funding:** This study has been funded by two postdoctoral fellowships from the Fyssen Foundation and the Fondation des Treilles granted to C.C. IVP was funded by the Swiss National Science Foundation (31003A_159587 and PP03P3_170624) and the Branco Weiss Fellowship – Society in Science granted to E.v.d.W. The authors do not have any conflict of interest. **Author contributions:** Data collection and video analysis: C.C. and M.B.C; statistical analysis: B.J.B (EWA) and C.C (GLMs); experimental design: C.C. and E.v.d.W; writing first draft: C.C. and B.J.B; reviews and editing: C.C., B.J.B. and E.v.d.W; funding: C.C. and E.v.d.W. **Competing interests**: The authors declare that they have no competing interests. **Data and materials availability**: All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

References

1. C.M. Heyes, Social learning in animals: categories and mechanisms. *Biol. Rev. Camb.*

Philos. Soc. **69**, 207e231 (1994).

2. J. Heinrich, R. McElreath, The evolution of cultural evolution. *Evol. Anthropol*. **12**,

123–135 (2003).

3. P. Marler, M. Tamura, Song 'dialects' in three populations of white-crowned sparrows.

Science. **146**, 1483-1486 (1964).

4. J.C. Holzhaider, G.R. Hunt, R.D. Gray, Social learning in new Caledonian crows.

Learn. & Behav. **38(3)**, 206-219 (2010).

5. L. Rendell, H. Whitehead, Culture in whales and dolphins. *Behav. & Brain Sci*. **24**, 309-

382 (2001).

6. J.B. Leca, N. Gunst, M. Huffman, Japanese macaque cultures: Inter- and intra- troop

behavioural variability of stone handling patterns across 10 troops. *Behaviour*, **144**, 251-

281 (2007).

7. S. Perry, M. Baker, L. Fedigan, J. Gros-Louis, K. Jack, K.C. MacKinnon, J.H. Manson,

M. Panger, K. Pyle, L. Rose, Social conventions in wild white-faced capuchin monkeys:

evidence for traditions in a neotropical primate. *Curr. Anthropol*. **44***, 241–268 (2003).

8. C.P. van Schaik, M. Ancrenaz, G. Borgen, B. Galdikas, C.D. Knott, I. Singleton, A.

Suzuki, S.S. Utami, M. Merrill, Orangutan cultures and the evolution of material

culture. *Science*, **299**, 102– 105. (2003).

9. A. Whiten, J. Goodall, W.C. McGrew, T. Nishida, V. Reynolds, Y. Sugiyama, C.E.G.

Tutin, R.W. Wrangham, C. Boesch, Cultures in chimpanzees. *Nature*, **399**, 682–685.

(1999).

10. L.L. Cavalli-Sforza, M.W. Feldman, *Cultural transmission and evolution: a*

quantitative approach. Princeton University Press. (1981).

11. R. Boyd, P.J. Richerson, *Culture and the Evolutionary Process*. Chicago: Univ. Chicago

Press. (1985).

12. A. Rogers, Does biology constrain culture? *Am. Anthropol*. **90**, 819-830. (1988).

13. R.L. Kendal, J.R. Kendal, W. Hoppitt, K.N. Laland, Identifying social learning in

animal populations: A new 'option-bias' method. ***PLoS ONE***, 4(8)*, e6541 (2009).

14. R. Baldini. Two Success-Biased Social Learning Strategies. *Theo. Pop. Biol.* **86**, 43–49

(2013).

15. R. McElreath, A.V. Bell, C. Efferson, M. Lubell, P.J. Richerson, T. Waring, Beyond

Existence and Aiming Outside the Laboratory: Estimating Frequency-Dependent and

Pay-Off-Biased Social Learning Strategies. *Phil. Trans. R. Soc. B*. **363 (1509)**, 3515–

28 (2008).

16. A. Mesoudi, An Experimental Comparison of Human Social Learning Strategies:

Payoff-Biased Social Learning Is Adaptive but Underused. *Evol. & Hum. Behav*. **32 (5)**,

334–42 (2011).

17. B.J. Barrett, R.L. McElreath, S.E. Perry, Pay-off-biased social learning underlies the

diffusion of novel extractive foraging traditions in a wild primate. *Proc. Roy. Soc. B*.

284, 20170358. (2017).

18. L.M. Aplin, B.C. Sheldon, R. McElreath, Conformity Does Not Perpetuate Suboptimal

Traditions in a Wild Population of Songbirds. *Proc. Nat. Aca. Sci.* **114 (30)**, 7830–7837

(2017).

19. A.E.J. Bono, A. Whiten, C. van Schaik, M. Krützen, F. Eichenberger, A. Schnider, E.

van de Waal, Payoff- and Sex-Biased Social Learning Interact in a Wild Primate

Population. *Curr. Biol.* **28 (17)**, 2800–2805.e4 (2018).

20. L. Rendell, L. Fogarty, W.J.E. Hoppitt, T.J.H. Morgan, M.M. Webster, M. M., K.N.
Laland, Cognitive culture: Theoretical and empirical insights into social learning
strategies. *Tr. Cogn. Sci*, **15***, 68–76 (2011).

21. N. Lamon, C. Neumann, T. Gruber, K. Zuberbühler, Kin-based cultural transmission
of tool use in wild chimpanzees. *Sci. Adv*. **3**, e1602750 (2017).

22. S. Wild, S.J. Allen, M. Krützen, S.L. King, Gerber, L., W.J.E. Hoppitt, Multi-network-
based diffusion analysis reveals vertical cultural transmission of sponge tool use within
dolphin matrilines. *Biol. Lett*. **15**, 20190227 (2019).

23. S. Wild, W.J.E. Hoppitt, S.J. Allen, M. Krützen, Integrating genetic, environmental and
social networks to reveal transmission pathways of a dolphin foraging innovation. *Curr.*
Biol. **30**, 1-7 (2020).

24. L.E. Dugatkin, J.G.J. Godin, Female mate copying in the guppy (*Poecilia reticulata*):
age- dependent effects. *Behav. Ecol*. **4(4)**, 289-292 (1993).

25. V. Horner, D. Proctor, K.E. Bonnie, A. Whiten, F.B.M. de Waal, Prestige affects
cultural learning in chimpanzees. *PLoS ONE*, **5**, e10625 (2010).

26. E. van de Waal, N. Renevey, C.M. Favre, R. Bshary, R, Selective attention to philopatric
models causes directed social learning in wild vervet monkeys. *Proc. Roy. Soc. B*, **277**,
2105-2111 (2010).

27. R.L. Kendal, L.M. Hopper, A. Whiten, S.F. Brosnan, S.P. Lambeth, S.J. Schapiro, S. J.
W. Hoppitt, Chimpanzees copy dominant and knowledgeable individuals: implications
for cultural diversity. *Evol. & Hum. Behav*. **36(1)***, 65-72 (2015).

28. C. Canteloup, W. Hoppitt, E. van de Waal, Wild primates copy higher-ranked
individuals in a social diffusion experiment. *Nat. Comm*, **11**, 459 (2020).

29. E. van de Waal, C. Borgeaud, A. Whiten, Potent social learning and conformity shape
a wild primate's foraging decisions. *Science*, **340**, 483-485 (2013).

30. L.M. Aplin, D.R. Farine, J. Morand-Ferron, A. Cockburn, A. Thornton, B.C. Sheldon,
Experimentally induced innovations lead to persistent culture via conformity in wild
birds. *Nature*, **518 (7540)**, 538–541 (2015).

31. D.B.M. Haun, Y. Rekers, M. Tomasello, Majority biased transmission in chimpanzees
and human children, but not orangutans. *Curr. Biol*. **22**, 727-731 (2012).

32. G.L. Vale, S.J. Davis, E. van de Waal, S.J. Schapiro, S.P. Lambeth, A. Whiten, Lack of
conformity to new local dietary preferences in migrating captive chimpanzees. *Anim*
Behav. **124**, 135-144 (2017).

33. T.W. Pike, K.N. Laland, Conformist learning in nine-spined sticklebacks' foraging
decisions. *Biol. Lett*. **6**, 466–468 (2010).

34. E. Danchin, S. Nöbel, A. Pocheville, A.C. Dagaeff, L. Demay, M. Alphand, S. Ranty-
Roby, L. van Renssen, M. Monier, L. Gazagne, M. Allain, G. Isabel, Cultural flies:
Conformist social learning in fruitflies predicts long-lasting mate-choice traditions.
Science, **362**, 1025-1030 (2018).

35. M. Franz, C.L. Nunn, Network-based diffusion analysis: a new method for detecting
social learning. *Proc. Roy. Soc. B*, **276**, 1829–1836 (2009).

36. W. Hoppitt, The conceptual foundations of network-based diffusion analysis: choosing
networks and interpreting results. *Phil. Trans. Roy. Soc. B*. **372**, 20160418 (2017).

37. C. Camerer, T-H. Ho, Experience-weighted attraction learning in normal form games.
Econometrica. **67(4)**, 827-874 (1999).

38. E. Bandini, C. Tennie. Exploring the role of individual learning in animal tool-use.
PeerJ. 8:e9877 (2020).

39. C. Tennie, A. Bandini, C.P. van Schaik, L.M. Hopper. The zone of latent solutions and
its relevance to understanding ape cultures. *Biol. & Phil*. 35-55 (2020).

40. R.L. Kendal, N.J. Boogert, L. Rendell, K.N. Laland, M. Webster, P.L. Jones. Social

learning strategies: bridge-building between fields. *Trends Cogn. Sci*. **22**, 651-665

(2018).

41. D. Biro, N. Inokue-Nakamura, R. Tonooka, G. Yamakoshi, C. Sousa, T. Matsuzawa,

Cultural innovation and transmission of tool use in wild chimpanzees: Evidence from

field experiments. *Anim. Cogn*. **6**, 213–223 (2003).

42. C.G. Coelho, T. Falotico, P. Izar, M. Mannu, B.D. Resende, J.O. Siqueira, E.B. Ottoni,

Social learning strategies for nut-cracking by tufted capuchin monkeys (*Sapajus* spp.).

Anim. Cogn. **18**, 911-919 (2015).

43. E.J.C. van Leeuwen, Kendal, R.L., C. Tennie, D.B.M. Haun, Conformity and its look-

a-likes. *Anim. Behav*. e1-e5, (2015)

44. B.J. Barrett. Equifinality in Empirical Studies of Cultural Transmission. *Behav. Proc*.

161, 129–38 (2019).

45. H. Kummer, J. Goodall, Conditions of innovative behaviour in primates. *Phil. Trans*.

Roy. Soc. B. **308**, 203–214 (1985).

46. S.M. Reader, K.N. Laland, Animal innovation: An introduction in *Animal innovation*

S. M. Reader, K. N. Laland Eds. (Oxford University Press, 2003), pp. 3–35.

47. S.E. Perry, B.J. Barrett, I. Godoy, Older, sociable capuchins (*Cebus capucinus*) invent

more social behaviors, but younger monkeys innovate more in other contexts. *Proc*.

Nat. Aca. Sci. **114(30)**, 7806-7813 (2017).

48. F. Amici, A. Widdig, J. Lehmann, B. Majolo, A meta-analysis of interindividual

differences in innovation. *Anim. Behav*. **155**, 257–268 (2019).

49. J. Altmann. Observational study of behavior: sampling methods. *Behav*. **49**, 227–267

(1974).

50. H. Whitehead. SOCPROG programs: analyzing animal social structure. *Behav. Ecol.*
Sociobiol. **63**, 765–778 (2009).

51. H. de Vries. Finding a dominance order most consistent with a linear hierarchy: a new
procedure and review. *Anim. Behav.* **55**, 827–843 (1998).

52. W. Toyokawa, A. Whalen, K.N. Laland. Social Learning Strategies Regulate the
Wisdom and Madness of Interactive Crowds. *Nat. Hum. Behav.* **3(2)**, 183–93 (2019).

53. A. Kandler, B. Wilder, L. Fortunato. Inferring Individual-Level Processes from
Population-Level Patterns in Cultural Evolution. *Roy. Soc. Op. Sci.* **4(9)**, 170949 (2017).

54. Stan Development Team. RStan: The R Interface to Stan, http://mc-stan.org/. (2020).

Figures and Tables

Figure 1

Figure 1. Daily mean group probability of processing techniques over experimental days for

NH. Filled lines and points are global model predictions. Dashed lines and circles are

probabilities of raw data. Diameter of points of raw data scales with daily sample size.

Figure 2

Figure 2. Visualization of posterior estimates from the global model of daily mean probability

of processing techniques across all foraging individuals in NH. Point diameter scales with

probability, with many individuals approaching fixity of CMS on the final day.

Figure 3

785
786

Figure 3. Model predictions of the probability of using a technique across foraging bout from

five models for the behavioral time series of individual 'Gran' in NH group. Colors correspond

to the three processing techniques (purple: 'CMS'; orange: 'CMT'; green: 'CH'). Line shows

posterior mean at each timestep, shaded interval is a 89% High Posterior Density Intervals

(HPDI). Points along top row are raw data showing observed behavioral choice at each foraging

bout. Filled circles are successes, empty circles are failures.

Table 1

Model	WAIC	SE	dWAIC	weight
Global Model	4707.71	91.09	0.00	1.00
Payoff-bias	4756.90	91.12	49.18	0.00
Rank-bias	4780.21	90.55	72.50	0.00
Frequency-dependence	4796.59	90.92	88.88	0.00
Matrilineal Kin Bias	4856.82	92.46	149.11	0.00
Same-sex-bias	4874.49	92.68	166.78	0.00
Female-bias	4875.25	92.73	167.54	0.00
Individual learning	4920.83	92.00	213.12	0.00

796   Table 1. Widely applicable information criteria estimates for all evaluated models with a 20

797   minute window of social information. Standard errors, difference in WAIC and model weight

798   are included.

800   Table 2
801

Parameter	Global	Payoff-bias	Rank-bias	Freq-dep.	Kin-bias	Female-bias	Sex-bias	Ind. learn.
λ_f	8.38	20.89	19.17	8.79	15.76	13.73	13.55	7.31
λ_m	6.27	16.93	14.86	6.24	14.69	13.66	13.41	6.36
λ_j	10.04	20.37	19.94	10.73	16.72	15.42	15.13	9.31
λ_a	5.22	17.22	14.22	5.09	13.81	12.14	12.02	4.99
ϕ_f	0.06	0.04	0.04	0.06	0.04	0.05	0.05	0.06
ϕ_m	0.05	0.04	0.04	0.05	0.04	0.04	0.04	0.05
ϕ_j	0.04	0.03	0.03	0.04	0.03	0.03	0.03	0.05
ϕ_a	0.08	0.05	0.05	0.07	0.05	0.06	0.06	0.07
γ_f	0.23	0.30	0.20	0.21	0.14	0.13	0.13	
γ_m	0.20	0.28	0.26	0.21	0.16	0.13	0.13	
γ_j	0.20	0.31	0.18	0.16	0.12	0.09	0.09	
γ_a	0.23	0.28	0.28	0.26	0.19	0.18	0.18	
f_f^c	1.26			1.66				
f_m^c	1.04			1.29				
f_j^c	1.08			1.60				
f_a^c	1.20			1.33				
$\beta_f em_f$	-0.03					-0.01		
$\beta_f em_m$	0.09					-0.16		
$\beta_f em_j$	-0.05					-0.20		
$\beta_f em_a$	0.11					0.03		
$\beta_k in_f$	0.53				1.07			
$\beta_k in_m$	0.75				1.00			
$\beta_k in_j$	0.63				0.98			

$\beta_k in_a$	0.65			1.09	
$\beta_p ay_f$	1.07	1.46			
$\beta_p ay_m$	0.62	0.87			
$\beta_p ay_j$	1.24	1.46			
$\beta_p ay_a$	0.45	0.87			
$\beta_r ank_f$	1.18		2.37		
$\beta_r ank_m$	0.89		2.25		
$\beta_r ank_j$	0.80		2.22		
$\beta_r ank_a$	1.26		2.40		
$\beta_s ex_f$	-0.40				0.17
$\beta_s ex_m$	-0.80				0.00
$\beta_s ex_j$	-0.86				-0.07
$\beta_s ex_a$	-0.34				0.24

802
803
804 Table 2. Posterior mean estimates of learning parameters for all evaluated EWA models.

805 Subscripts correspond to male (m), female (f), adult (a), and juvenile (j).

807   Table 3

Model	Social and Individual Cues
1) Individual learning	$\pi_{ij,t} = 1$ if successful, 0 if not
2) Frequency-dependence	$N_{ij,t}$= count of times j observed demonstrators perform behavior i
3) Female-bias	$\kappa_{k,ijt}$= 1 if adult female, 0 otherwise
4) Matrilineal kin-bias	$\kappa_{k,ijt}$= coefficient of relatedness, r, between observer j and demonstrator
5) Payoff-biased learning	$\kappa_{k,ijt}$= mean observed payoff, $\overline{\pi_{ij,t}}$, of behavior i observed by j
6) Rank-bias	$\kappa_{k,ijt}$= rank of demonstrator determined by I&SI method
7) Sex-bias	$\kappa_{k,ijt}$= 1 if same sex as observe, 0 if different
8) Global model	all cues in 1-7

808
809   Table 3. Individual information and social cues used in the EWA models. Models 2-8 also

810   incorporate individual experience from model 1. Information observed at timestep t for models

811   2-8 was the mean of each cue, κ evaluated in the 20 minutes prior to timestep t.