A high affinity human monoclonal antibody against Pfs230 binds multiple parasite stages and blocks oocyst formation in mosquitoes

Camila H. Coelho[1,11], Wai Kwan Tang[2,11], Martin Burkhardt[3], Jacob D. Galson[4,5], Olga Muratova[3], Nichole D. Salinas[2], Thiago Luiz Alves e Silva[6], Karine Reiter[3], Nicholas J. MacDonald[3], Vu Nguyen[3], Raul Herrera[3], Richard Shimp[3], David L. Narum[3], Miranda Byrne-Steele[7], Wenjing Pan[7], Xiaohong Hou[7], Brittany Brown[7], Mary Eisenhower[7], Jian Han[7], Bethany J. Jenkins[1], Justin Yai Alamou Doritchamou[1] , Margery G. Smelkinson[8], Joel Vega-Rodriguez[6], Johannes Trück[4], Justin J. Taylor[9], Issaka Sagara[10], Jonathan P. Renn[3], Niraj H. Tolia[2,12*], Patrick E. Duffy[1,12*]

1 - Pathogenesis and Immunity Section, Laboratory of Malaria Immunology and Vaccinology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA
2 - Host-Pathogen Interactions and Structural Vaccinology Section, Laboratory of Malaria Immunology and Vaccinology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA
3- Vaccine Development Unit, Laboratory of Malaria Immunology and Vaccinology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA
4 - University Children's Hospital, and the Children's Research Center, University of Zurich, Zurich, Switzerland
5- Alchemab Therapeutics Ltd, 55-56 Russell Square, London, WC1B 4HP, UK
6 - Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD, United States
7 - iRepertoire Inc., Huntsville, AL, USA.
8- Biological Imaging Section, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA
9 - Fred Hutchinson Cancer Research Center, Seattle, WA, USA
10 - Malaria Research and Training Center, University of Sciences, Techniques, and Technology, Bamako, Mali
11-These authors contributed equally: Camila H. Coelho, Wai Kwan Tang
12- These authors jointly directed this work Niraj H. Tolia1, Patrick E. Duffy

* Corresponding authors:
patrick.duffy@nih.gov and niraj.tolia@nih.gov

Laboratory of Malaria Immunology and Vaccinology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, 9000 Rockville Pike, Bethesda MD 20892, USA.

ABSTRACT

Malaria elimination requires tools that interrupt parasite transmission. Here, we characterized B cell receptor responses among Malian adults vaccinated against the first domain of the cysteine-rich 230kDa gamete surface protein Pfs230[1-3] to neutralize sexual stage *P. falciparum* parasites and halt their further spread. We generated nine Pfs230 human monoclonal antibodies (mAbs). One mAb potently blocked transmission to mosquitoes in a complement-dependent manner and reacted strongly to gamete surface while eight mAbs showed only low or no blocking activity. This study provides a rational basis to improve malaria vaccines and develop therapeutic antibodies for malaria elimination.

MAIN TEXT

Malaria eradication is a global priority and will require innovative strategies that, in addition to preventing or controlling human infection, might block parasite transmission through mosquitoes. Sequences of matched heavy and light chain variable

regions from single human B cells have been used to identify antibodies generated in response to infection or vaccination and inform vaccinology[4-7]. In this study, we apply this approach to examine human antibodies elicited in response to a transmission blocking vaccine (TBV), that used a Pfs230 fragment as antigen. Pfs230 is present on the surface of *P. falciparum* gametocytes and gametes and mediates binding of exflagellating microgametes to red blood cells, thus parasites lacking this protein cannot bind to red blood cells or further develop into oocysts.[1] We collected Pfs230 domain 1 (D1)-specific single memory B cells **(Extended Data Fig. 1, Extended Data Fig. 2a)** from eight Malian adults immunized with four doses of Pfs230D1-EPA/Alhydrogel® vaccine (Clinicaltrials.gov NCT02334462) to identify functional monoclonal antibodies elicited in response to a TBV. This vaccine aims to neutralize sexual stage *P. falciparum* parasites by targeting Pfs230, a 230kDa gamete surface protein comprised of fourteen 6-cysteine (6-Cys) domains[1-3]. All samples were chosen from subjects presenting high serum Transmission-Reducing Activity (TRA), measured by the capacity of serum antibodies from immunized subjects to reduce the number of oocysts that develop in mosquitoes fed on in vitro cultured *P. falciparum* gametocytes **(Extended Data Table 1)**.

We obtained 272 VH and 351 VL sequences of B cell receptor (BCR) from Pfs230D1-specific single memory B cells from the vaccinees via amplification and sequencing of the V(D)J region **(Extended Data Fig. 3).** When analysing V gene usage of the BCR sequences, 87.5% of the subjects presented Pfs230D1-specific memory B cells using kappa chains derived from IGKV4-1 **(Extended Data Fig. 2e).** This light chain gene has also been identified in sequences of functional human mAbs obtained in

response to other *Plasmodium* antigens[4-6,8]. For the heavy chain, IGHV1-69 was the

most commonly expressed gene and detected in 100% (8/8) of vaccinees **(Extended**

Data Fig. 2f). IGHV1-69 is one of the most polymorphic loci of the IGHV gene cluster[9]

and is frequently found in broadly neutralizing antibodies generated in response to

influenza haemagluttinin[10,11].

Nine pairs of BCR sequences were chosen for expression of fully human

Pfs230D1 IgG1 antibodies by assessing whether the CDR3 sequences were shared

between sorted B cells. This approach identifies identical sequences in multiple B cells

from the same subject, indicating that they have been activated in response to

vaccination. These nine pairs **(Fig. 1a)** represented distinct heavy and light chain

germline genes with an overabundance of IGHV1-18 (N=6), IGHV1-69 (N=3), and

IGKV4-1 (N=7). The resulting recombinant antibodies bound to Pfs230D1 antigen

(Figure 1d,e, Extended Data Figure 4). Competitive epitope binning of the nine mAbs

suggested they bind three non-overlapping epitopes in Pfs230D1 **(Fig. 1b)**. LMIV230-01

forms a distinct group (Group 1) and has potent neutralizing activity **(Fig. 1b, c)**. The

remaining mAbs do not compete with LMIV230-01 and may form two additional epitope

groups. Group 2 and 3 mAbs possess low or no neutralizing activity **(Fig. 1c)**. We

therefore focused most of our subsequent analyses on LMIV230-01 and to a lesser

extent on LMIV230-02, which demonstrated low functional activity.

LMIV230-01 and 02 bound to Pfs230D1 recombinant protein **(Fig. 1d)** with

strong and similar binding affinities **(Fig. 1e, Extended Data Fig. 4, Extended Data**

Table 2). We confirmed the two mAbs bind distinct epitopes using competition ELISA

(Extended data Figure 5d) consistent with the epitope binning results (Fig. 1b). Despite their shared use of IGHV1-69, LMIV230-01 and LMIV230-02 displayed numerous differences in their heavy chain CDRs, consistent with their recognition of distinct epitopes (Extended Data Figure 12).

Although presenting similar affinity to Pfs230D1, the mAbs differed in their functional activity as measured by SMFA. LMIV230-01 ablated *P. falciparum* oocyst burden in mosquitoes in a dose-dependent manner with 91.7% neutralization (TRA) at 1000 µg/mL (Fig. 1f). Importantly, 80.3% neutralization was retained at 60 µg/mL. On the other hand, LMIV230-02 reduced oocyst burden by only 58.7% at the maximum concentration of 1000 µg/mL and activity was lost at 250 µg/mL. As previously reported, TRA values higher than 80% are highly reproducible across independent experiments[12,13].Combining the two antibodies did not increase their overall activity: TRA values were not statistically different when 500µg of LMIV230-02 was combined with 10µg of LMIV230-01 (TRA= 58.7%) versus 10µg of LMIV230-01 alone (TRA= 52.5%) in mosquito feeding assays (Extended Data Figure 5e).

To understand the differences in functional activity of the two mAbs, we assessed binding to the native protein. Both mAbs reacted to the protein extract of parasites and were sensitive to reduction of the two disulfide-bonds, suggesting the presence of conformational epitopes (Fig. 1g, Extended Data Fig. 5c). Interestingly, LMIV230-01 strongly labelled the surface of live *P. falciparum* gametes purified 2 hours post-exflagellation, while LMIV230-02 did not (Fig. 1h). This suggests that the LMIV230-02 epitope is not completely accessible on the surface-displayed native

134   protein, possibly due to structural limitations imposed by the multi-domain protein

135   Pfs230, as has been seen for other proteins[14,15] including another 6-Cys TBV

136   candidate[16].

a

mAb	Heavy chain	Light chain
LMIV230-01	IGHV1-69	IGKV1-5
LMIV230-02	IGHV1-69	IGKV4-1
GKV16	IGHV1-69	IGKV4-1
GKV08	IGHV1-18	IGKV4-1
GKV01	IGHV1-18	IGKV4-1
GKV07	IGHV1-18	IGKV4-1
GKV05	IGHV1-18	IGKV4-1
GKV06	IGHV1-18	IGKV4-1
GKV22	IGHV1-18	IGKV2-28

b

	Group 1	Group 2							Group 3	
	LMIV230-01	LMIV230-02	GKV01	GKV05	GKV06	GKV07	GKV22	GKV08	GKV16
LMIV230-01		95	100	100	99	100	100	100	100
LMIV230-02	85		10	13	14	18	4	51	60
GKV01	100	0		1	1	1	3	41	50
GKV05	100	0	0		0	0	0	40	49
GKV06	100	0	0	0		1	0	40	51
GKV07	100	0	0	0	1		0	41	53
GKV22	89	0	0	4	3	7		38	41
GKV08	71	0	0	0	0	0	0		18
GKV16	24	0	0	0	0	0	0	0	

c

		LMIV230-01	LMIV230-02	GKV01	GKV05	GKV06	GKV07	GKV22	GKV08	GKV16
TRA%	375 µg/mL	- - -	- - -	20%	0%	10%	10%	0%	0%	0%
	250 µg/mL	85%	51%	- - -	- - -	- - -	- - -	- - -	- - -	- - -

Fig. 1| Human recombinant mAbs were generated from Pfs230D1-specific single memory B cells of Malian adults vaccinated with the Pfs230D1-EPA/Alhydrogel® TBV. a, VH and VL genes corresponding to each mAb. LMIV230-01 and LMIV230-02 sequences originate from the IGHV1-69 heavy chain gene but utilize different kappa chain genes. Complete V gene usage determined in Pfs230-specific memory B cells is described in **Extended Data Figures 2e,f**. **b,** Epitope binning of human anti-Pfs230D1scFvs. The primary binding scFv is listed on the left and the competing scFv are listed on the top. Reported scores are a percentage of total binding of that antibody in the absence of a competitor scFv. Values greater that 50% display low amounts of competition, while values lower than 50% exhibit greater competition. Any experiment with >100% binding was given a score of 100, while negative values were given a score of 0. Potential epitope bins are grouped and labelled above the table. **c,** Functional activity of each mAb, assessed by Standard Membrane Feeding Assay (SMFA) and measured as the % reduction (versus control mAb) in the number of *P. falciparum* NF54 oocysts in midguts of *Anopheles* mosquitoes ("TRA"). **d,** LMIV230-01 and LMIV230-02 mAbs bound similarly to Pfs230D1 and **e,** show high affinity to recombinant Pfs230D1 **(Extended Data Fig. 4, Extended Data Table 2)** **f,** LMIV230-01 reduces *P. falciparum* NF54 oocyst numbers by 91.7% at 1000 µg/mL, 86.7% at 500 µg/mL, 84.4% at 250 µg/mL and 80.3% at 60 µg/mL, while LMIV230-02 displays only modest activity with 58.7% reduction at the maximum concentration of 1000 µg/mL, in SMFA. Data from eleven independent SMFA and each concentration was evaluated in at least three biological replicates for each mAb. N ≥ 20 mosquitos per assay. Average oocyst numbers in the control mosquitoes (fed with mouse IgG1 mAb targeting *P. yoelii* P140 protein) for each experiment were: exp. 1 = 29.73; exp.2= 7.18; exp. 3= 57.86; exp. 4= 36.41; exp. 5= 51.71, exp. 6= 4.55; exp. 7= 62.35; exp. 8= 20.50, exp.9= 8.71, exp 10= 18.05, exp. 11= 5.86. Negative oocyst reduction values were set to zero. Human isotype IgG1 and US human serum pool were used as additional negative controls **(Extended Data Fig. 5b)**. Values are shown as mean ± s.e.m. **g,** LMIV230-01 and LMIV230-02 bind to non-reduced (NR) protein extract of *P. falciparum* NF54 gametes purified on Nycodenz after 2 hours in exflagellation medium. **h,** LMIV230-01 strongly binds to gametes at 7.5µg/mL while LMIV230-02 does not bind at 7.5µg/mL. or 30µg/mL. Both mAbs were labelled with Alexa Fluor 488. Scale bars: 5µM.

LMIV230-01 bound strongly to fixed parasites in numerous developmental stages including gametocytes, exflagellation centers, microgametes, macrogametes and round forms (zygotes) collected 4 hours after mosquito feeding. As expected, the mAb did not bind to the post-fertilization stage ookinetes, obtained 24 hours after the mosquito bloodmeal **(Fig. 2a)**.

Pfs230 antibody activity depends on complement fixation to lyse *P. falciparum*[17]. To test whether the activity of LMIV230-01 was dependent on activation of the complement system, we incubated parasites with LMIV230-01 in the presence of intact or heat-inactivated sera from US donors then assessed lysis of gametes **(Fig. 2b)** as well as transmission of parasites fed to mosquitoes after treatment using the same

conditions **(Fig. 2c)**. Functional activity of LMIV230-01 to lyse gametes and block oocyst

formation in mosquitoes was substantially reduced in the heat-inactivated sera **(Figs.**

2b and 2c), demonstrating complement-dependency. Activation of complement leads to

the formation of the membrane attack complex (MAC), an assembly of the complement

molecules C5b, C6, C7, C8, and C9[18,19] on the parasite surface. Using an antibody that

recognizes assembled MAC, we demonstrated complement fixation on the surface of

live *P. falciparum* gametes that were incubated with LMIV230-01 in the presence of

intact but not heat-inactivated serum **(Fig. 2d and Extended Data Fig. 7).**

a

| | Gametocyte | Exflagellation center | Microgamete | Macrogamete | Zygote | Ookinete |

LMIV230-01

5 μm

b

Parasite lysis assay

Parasite lysis (%)

p< 0.0001

ns

LMIV230-01   IgG1 control

Intact serum
HI serum

c

SMFA

Number of oocysts per *A. stephensi* midgut

p< 0.0001

ns

LMIV230-01   Anti- *P. yoelii* P140

d

LMIV230-01

IgG1 control

C5b-9+C5b-8
DNA

Intact serum   HI serum

C5b-9+C5b-8
DNA

Intact serum   HI serum

Fig. 2| LMIV230-01 binds to multiple parasite stages and its activity is complement-dependent. a, LMIV230-01 strongly binds to permeabilized gametocytes, gametes and zygotes and does not bind to ookinetes. Parasites were fixed and permeabilized, and 7.5 µg/mL of antibody was used to stain the different parasite stages. Scale bars: 5µM. **b,** *In vitro* parasite lysis by LMIV230-01 is complement-dependent. Samples were tested in two independent assays, using two different parasite cultures **c,** Functional activity of LMIV230-01 is also complement-dependent *in vivo* (SMFA with mosquitoes). Data from three independent SMFA assays. N ≥ 20 mosquitos per assay. Oocyst averages in the control mosquitoes (fed with IgG1 targeting *P. yoelli* P140) for each of the experiments were: exp. 1= 4.55; exp. 2= 20.50, exp. 3=5.86. Data obtained from mosquitoes fed with LMIV230-01 at 1000 µg/mL with intact sera were also used to generate figure 1f. Values are shown as mean ± s.e.m. One-Way ANOVA and Turkey's multiple comparisons test were used to compare the different groups **d,** Live imaging of *P. falciparum* NF54 female gametes incubated with LMIV230-01 in the presence of intact serum from a healthy donor revealed surface-deposited MAC (membrane attack complex) using anti-C5b-9+C5b-8 antibody (magenta color). MAC deposition was not detected in the presence of heat-inactivated (HI) serum. Scale bars: 5µM.

To assess whether LMIV230-01 would also bind to other *P. falciparum* strains, we prepared gametocytes from a culture-adapted Malian isolate [20] and from St. Lucia strain (originally from El Salvador) [21]. LMIV230-01 labelled in vitro-induced gametes from both strains **(Fig. 3a and b)**. Induction of gamete stage from the newly characterized Malian isolate was confirmed using a murine anti-Pfs48/45 mAb **(Fig. 3c)**. LMIV230-01 fixed complement on the gamete surface of both strains, confirming that the antibody is functional against heterologous parasites **(Figs. 3d and e).**

To assess the abundance of antibodies that share paratope specificity with LMIV230-01, we developed an ELISA assay to demonstrate the competition between post-vaccination sera (tested at a 1:2500 dilution) and LMIV230-01 for binding the vaccine antigen **(Figure 3f)**. Among subjects who received the vaccine, levels of competing antibody ranged from ~10-70% displacement of Pfs230D1 binding to LMIV230-01, with a normal distribution confirmed by Shapiro-Wilk test (p= 0.52) **(Figure 3g)**. Levels of competition strongly correlated with total Pfs230D1 IgG titers in sera (Spearman's rho= 0.604, p<0.0001) **(Figure 3h)**. Increasing levels of competing

antibody also corresponded to serum functional activity measured by SMFA. Because serum TRA levels of vaccines were high with minimal variability ranging from 95-100% **(Extended Data Figure 14)**, our primary correlation analysis used TBA (Transmission Blocking Activity) which indicates the % reduction in the proportion of infected mosquitoes, a high bar for TBV activity generally seen only when TRA is very high. Correlation analyses showed that % serum competition was related to TBA (Spearman's rho= 0.324, p= 0.05) **(Figure 3i)**, suggesting that antibodies that compete for the LMIV230-01 epitope play an important role in serum functional activity. This result, however, does not exclude the possible role of antibodies that do not compete with LMIV230-01 in mediating vaccine activity, and notably some sera with high TBA demonstrated low levels of LMIV230-01 competing antibodies.

a

St. Lucia strain

LMIV230-01

b

Malian isolate 20-2217-0

LMIV230-01

c

Malian isolate 20-2217-0

| Bright field | LMIV230-01 |
| DNA | LMIV230-01 |

| Anti-Pfs48/45 | Merged |
| Anti-Pfs48/45 | |

d

St. Lucia strain

| LMIV230-01 + intact serum | LMIV230-01 + HI serum |
| C5b-9+C5b-8 | C5b-9+C5b-8 |

e

Malian isolate 20-2217-0

| LMIV230-01 + intact serum | LMIV230-01 + HI serum |
| C5b-9+C5b-8 | C5b-9+C5b-8 |

f

| Plates coated with LMIV230-01 | Serum and Pfs230D1-biotin preincubated then added | Plates washed, streptavidin-HRP and substrate added |

Vaccinee serum

Non-immune serum

LMIV230-01-competing Abs   Other Pfs230D1 Abs   Nonspecific Abs   Pfs230D1 protein   Biotin   Streptavidin-HRP   Substrate

g

Serum competition for binding of LMIV230-01 mAb

% of subjects

% Competition

h

Spearman's rho= 0.604, p< 0.0001

% Competition

Pfs230D1 total IgG

i

Spearman's rho= 0.324, p= 0.05

% Competition

TBA%

Figure 3| LMIV230-01 binds to heterologous *P. falciparum* strains and antisera from Pfs230D1
vaccinees vary widely in levels of antibody that compete with LMIV230-01 for binding. a, LMIV230-
01 bound to gametes of *St. Lucia* parasite strain and **b,** of an isolate obtained from a Malian adult and
adapted to culture. **c,** Murine anti-48/45 mAb confirms formation of gametes by Malian isolate and its
signal colocalizes with LMIV230-01. "Merged" refers to combination of green and red channels. **d,**
Membrane attack complex forms on gametes of St. Lucia strain and **e,** of a Malian isolate incubated with
LMIV230-01 in the presence of intact but not heat-inactivated serum**.** Scale bars for all images in this
panel: 5μM. **f,** Cartoon schematizing LMIV230-01 competition ELISA assay. **g,** Distribution of serum
antibody levels that compete with LMIV230-01 for binding to Pfs230D1 in 36 subjects who received
Pfs230D1-EPA vaccine. Values displayed represent mean from three independent experiments. **h,**
Relationship of LMIV230-01-competing antibody levels to total Pfs230D1 antibody titers, or **i**, to serum
functional activity (TBA, transmission blocking activity) measured by SMFA.

Altogether, our data confirm that vaccination with TBV can elicit strong

neutralizing antibodies, capable of binding to gametocytes, gametes, and zygotes, and

of impairing fertilization in the mosquito. Due to its complex domains and repeating

motifs with numerous disulfide bonds, expression of full length Pfs230 has been

difficult[22,23]. Preclinical studies of Pfs230 fragments have shown that immunization with

recombinant domain 1 of Pfs230 (Pfs230D1), but not other domains, induces strong

functional TRA in SMFA[3,22,24].

Our data support further development of TBV strategies to induce potent

antibody responses against mosquito sexual stage parasites.

ACKNOWLEDGMENTS

This work was funded by the Intramural Research Program of the National Institute of

Allergy and Infectious Diseases, National Institutes of Health. Calvin Eigsti provided

support for single cell sorting; Sundar Ganesan for live cell imaging; Ashley McCormack

and Emily Higbee for Standard Membrane Feeding Assays; and J. Patrick Gorres for

proofreading and editing this manuscript. We are thankful to Jillian Neal and Robert

Morrison for determining the Pfs230D1 sequence for the Malian *P. falciparum* isolate.

JT and JDG were supported by the Swiss National Science Foundation (Ambizione-

SCORE fellowship to JT: PZ00P3_161147; PZ00P3_183777)**.** We thank the staff

members of GM/CA beamline at the Advanced Photon Source, Argonne National

Laboratory for beamline support.

AUTHOR CONTRIBUTIONS

C.H.C. and P.E.D. conceived the single B cell sorting of Pfs230D1-specific B cells, V

gene repertoire analyses, antibody generation, conventional and competitive ELISAs,

Western blot, microscopy-based binding assays and in vitro and in vivo functional

characterization of mAbs. W.K.T. and N.H.T. conceived the epitope binning and

biophysical studies. C.H.C, W.K.T., N.H.T and P.E.D conceived the analysis of

polymorphisms. C.H.C., W.K.T., M.B., J.R., A.S., T.A.S., W.P., X.H., B.B., O.M., B. J,

M.S. and N.D.S. performed the experiments. M.E., C.H.C., and J.D.G. performed

bioinformatic analyses. N.J.M., K.R., V.N., R.H., R.S. and D.N. generated recombinant

Pfs230D1. I.S., J.J.T., J.V.R., J.T., J.H., M.B.S, J.R., N.H.T. and P.E.D. supervised the

experiments and interpreted the data. C.H.C., W.K.T., N.H.T. and P.E.D. wrote the

manuscript, with input from all authors.

COMPETING INTERESTS

M.B.S, W.P., X.H., B.B., and M.E. declare competing financial interests as all are

employees of iRepertoire Inc., and J.H. is co-founder and CEO. J.D.G. is an employee

of Alchemab Therapeutics Limited.

CODE AVAILABILITY

Code is available on request from the corresponding author.

EXTENDED DATA - FIGURES

Extended Data Fig. 1| Experimental pipeline. Pfs230D1-specific single B cells were sorted from PBMCs of eight Malian adults who had been immunized with four doses of 40µg of Pfs230D1-EPA/Alhydrogel®. After extraction of single B cells, a 500 bp fragment of the BCR variable regions of VH/VL were amplified and sequenced. Matched VH/VL pairs that were identified in more than one B cell were preferentially selected for cloning in an IgG1 vector for expression in 293 HEK cells and subsequent analyses.

Extended Data Fig. 2| Pfs230D1-specific mAbs belong to the same heavy chain germline subgroup but differ for kappa chain. a, Sorted memory B cells were gated as live, single cells, excluded for CD3, CD14 and CD56, and gated on CD19+, CD20+, CD27+ cells. Then, a tetramer approach was used to select antigen-specific cells and reduce nonspecific binding. Cells binding to the decoy tetramer (BSA) were excluded and only those binding to Pfs230D1 were selected for sorting. **b,** Serum from each subject was used to measured antibody titers against Pfs230D1 and functional activity to reduce oocyst burden in Standard Membrane Feeding Assays (SMFA). TRA= Transmission Reducing Activity measured as the reduction in average oocyst count; TBA= Transmission Blocking Activity measured as the reduction in the proportion of infected mosquitoes. **c,** Proportion of memory B cells for each subject that are Pfs230D1-specific. **d,** Complementarity-determining regions (CDRs) of each sequence selected for mAb expression. **e,** IGKV4-1 germline (gene sequence in LMIV230-02) was the most frequent for the kappa chain genes. IGKV1-5 germline (gene sequence in LMIV230-01) was found in only three subjects **f,** Sequences related to germline 1-69 of the IGHV gene were the most frequently elicited in response to the vaccination.

327   **Extended Data Fig. 3| Amplification of V(D)J region.** 500 bp fragment amplified from cDNA of sorted
328   Pfs230D1-specific single B cell. This fragment was obtained using primers targeting the V(D)J region
329   (iRepertoire Inc.).

Extended Data Fig.5| Additional binding and functional characterization of LMIV230-01 and -02. a, Both mAbs failed to bind to the ookinete protein Pfs25. **b,** Additional controls for the Standard Membrane Feeding Assay (SMFA). Human IgG1 isotype control was expressed using the same conditions as LMIV230-01 and -02 and was used in this assay at 1000μg/mL. Sixty microliters of undiluted human pooled serum obtained from US healthy donors were used as additional control. Values are shown as mean ± s.e.m. **c,** Full depiction of the Western blot gel displayed in Fig. 1g .**d,** The two mAbs do not compete for the same epitope in the recombinant Pfs230D1 protein, since unlabelled LMIV230-01 blocks binding of LMIV-230-01-HRP to immobilized Pfs230D1 but LMIV230-02 does not. **e,** Combination of LMIV230-01 and LMIV230-02 did not increase functional activity over LMIV230-01 alone. Control mosquitoes were fed with mouse IgG1 mAb targeting *P. yoelii* P140 protein, or with non-immune human serum.

[EXT. DATA FIGURE 6 WILL BE AVAILABLE IN THE PUBLISHED VERSION]

Extended Data Fig.7| Pfs230 mAbs activity is complement-dependent and LMIV230-01 competing
antibodies are acquired at varying levels by vaccinees. a, Activity of LMIV230-01 and LMIV230-02 is
complement-dependent in the vitro lysis assay and **b,** in the vivo mosquito feeding assay. **c,** Membrane
attack complexes (MAC) on parasites were detected using an Alexa 488-labeled antibody that recognizes
the assembled MAC complex (anti C5b-9+ C5b-8). Gametes incubated with LMIV230-01 and intact
serum produced MAC-positive parasites. Heat-inactivating serum to degrade the heat-labile components
of the complement pathway eliminated deposition of MAC on gametes. MAC-positive *P. falciparum* strain
NF54 gametes were enumerated in a large, tiled confocal image and normalized to the number of
Hoechst-stained nuclei.

[EXT. DATA FIGURES 8- 13 WILL BE AVAILABLE IN THE PUBLISHED VERSION

Extended Data Fig. 14| Correlation between levels of LMIV230-01 competing antibodies and Transmission-Reducing Activity (TRA) measured in SMFA.

 EXTENDED DATA – TABLES

Subject ID	Antibody titers	TRA (%)	TBA (%)
1	944	99.9	96
2	1581	99.3	73.9
3	2115	100	100
4	1382	99.6	79.2
5	2100	100	100
6	5277	100	100
7	800	99.9	95.8
8	774	95.1	20.8

398 **Extended Data Table 1-** Antibody titers and functional activity of sera from the eight subjects whose
399 sequences were analyzed in this study. TRA= Transmission-reducing activity. TBA=Transmission
400 blocking activity.

	K_D (x 10^{-10} ± SEM M)	k_a (x 10^5 ± SEM 1/Ms)	k_{dis} (x 10^{-4} ± SEM 1/s)	N
LMIV230-01				
Biological Replicate 1	1.58 ± 0.77	1.71 ± 0.06	0.28 ± 0.15	3
Biological Replicate 2	2.06 ± 0.99	1.80 ± 0.04	0.37 ± 0.18	3
LMIV230-02				
Biological Replicate 1	6.36 ± 0.24	7.67 ± 0.21	4.87 ± 0.06	3
Biological Replicate 2	4.27 ± 0.22	6.37 ± 0.13	2.71 ± 0.10	3

404 **Extended Data Table 2-** Binding of mAbs LMIV230-01 and LMIV230-02 to Pfs230D1 using Biolayer
405 Interferometry. Binding data for each mAb was fitted using a 1:1 binding model. The averages for two
406 biological replicates, composed of three technical replicates each, are shown for both mAbs.

409 *[EXT. DATA TABLES 3-7 WILL BE AVAILABLE IN THE PUBLISHED VERSION]*

REFERENCES

1. Eksi, S.*, et al.* Malaria transmission-blocking antigen, Pfs230, mediates human red blood cell binding to exflagellating male parasites and oocyst production. *Mol Microbiol* **61**, 991-998 (2006).
2. Williamson, K.C., Keister, D.B., Muratova, O. & Kaslow, D.C. Recombinant Pfs230, a Plasmodium falciparum gametocyte protein, induces antisera that reduce the infectivity of Plasmodium falciparum to mosquitoes. *Mol Biochem Parasitol* **75**, 33-42 (1995).
3. Tachibana, M.*, et al.* Identification of domains within Pfs230 that elicit transmission blocking antibody responses. *Vaccine* **37**, 1799-1806 (2019).
4. Triller, G.*, et al.* Natural Parasite Exposure Induces Protective Human Anti-Malarial Antibodies. *Immunity* **47**, 1197-1209 e1110 (2017).
5. Kisalu, N.K.*, et al.* A human monoclonal antibody prevents malaria infection by targeting a new site of vulnerability on the parasite. *Nat Med* **24**, 408-416 (2018).
6. Alanine, D.G.W.*, et al.* Human Antibodies that Slow Erythrocyte Invasion Potentiate Malaria-Neutralizing Antibodies. *Cell* **178**, 216-228 e221 (2019).
7. Rappuoli, R., Bottomley, M.J., D'Oro, U., Finco, O. & De Gregorio, E. Reverse vaccinology 2.0: Human immunology instructs vaccine antigen design. *J Exp Med* **213**, 469-481 (2016).
8. Tan, J.*, et al.* A LAIR1 insertion generates broadly reactive antibodies against malaria variant antigens. *Nature* **529**, 105-109 (2016).
9. Watson, C.T.*, et al.* Complete haplotype sequence of the human immunoglobulin heavy-chain variable, diversity, and joining genes and characterization of allelic and copy-number variation. *Am J Hum Genet* **92**, 530-546 (2013).
10. Avnir, Y.*, et al.* Molecular signatures of hemagglutinin stem-directed heterosubtypic human neutralizing antibodies against influenza A viruses. *PLoS Pathog* **10**, e1004103 (2014).
11. Avnir, Y.*, et al.* IGHV1-69 polymorphism modulates anti-influenza antibody repertoires, correlates with IGHV utilization shifts and varies by ethnicity. *Sci Rep* **6**, 20842 (2016).
12. Lensen, A.*, et al.* Measurement by membrane feeding of reduction in Plasmodium falciparum transmission induced by endemic sera. *Trans R Soc Trop Med Hyg* **90**, 20-22 (1996).
13. Miura, K.*, et al.* Transmission-blocking activity is determined by transmission-reducing activity and number of control oocysts in Plasmodium falciparum standard membrane-feeding assay. *Vaccine* **34**, 4145-4151 (2016).
14. Doolan, K.M. & Colby, D.W. Conformation-dependent epitopes recognized by prion protein antibodies probed using mutational scanning and deep sequencing. *J Mol Biol* **427**, 328-340 (2015).
15. Verrier, F.C.*, et al.* Antibodies to several conformation-dependent epitopes of gp120/gp41 inhibit CCR-5-dependent cell-to-cell fusion mediated by the native envelope glycoprotein of a primary macrophage-tropic HIV-1 isolate. *Proc Natl Acad Sci U S A* **94**, 9326-9331 (1997).
16. Outchkourov, N.*, et al.* Epitope analysis of the malaria surface antigen pfs48/45 identifies a subdomain that elicits transmission blocking antibodies. *J Biol Chem* **282**, 17148-17156 (2007).
17. Healer, J.*, et al.* Complement-mediated lysis of Plasmodium falciparum gametes by malaria-immune human sera is associated with antibodies to the gamete surface antigen Pfs230. *Infect Immun* **65**, 3017-3023 (1997).
18. Tegla, C.A.*, et al.* Membrane attack by complement: the assembly and biology of terminal complement complexes. *Immunol Res* **51**, 45-60 (2011).
19. Bayly-Jones, C., Bubeck, D. & Dunstone, M.A. The mystery behind membrane insertion: a review of the complement membrane attack complex. *Philos Trans R Soc Lond B Biol Sci* **372**(2017).
20. Doritchamou, J.Y.A.*, et al.* Placental malaria vaccine candidate antigen VAR2CSA displays atypical domain architecture in some Plasmodium falciparum strains. *Commun Biol* **2**, 457 (2019).
21. Collins, W.E., Warren, M., Skinner, J.C., Richardson, B.B. & Kearse, T.S. Infectivity of the Santa Lucia (El Salvador) strain of Plasmodium falciparum to different anophelines. *J Parasitol* **63**, 57-61 (1977).
22. MacDonald, N.J.*, et al.* Structural and Immunological Characterization of Recombinant 6-Cysteine Domains of the Plasmodium falciparum Sexual Stage Protein Pfs230. *J Biol Chem* **291**, 19913-19922 (2016).
23. Farrance, C.E.*, et al.* A plant-produced Pfs230 vaccine candidate blocks transmission of Plasmodium falciparum. *Clin Vaccine Immunol* **18**, 1351-1357 (2011).
24. Singh, S.K.*, et al.* Pfs230 and Pfs48/45 Fusion Proteins Elicit Strong Transmission-Blocking Antibody Responses Against Plasmodium falciparum. *Front Immunol* **10**, 1256 (2019).

25.   Tonkin, M.L.*, et al.* Structural and biochemical characterization of Plasmodium falciparum 12 (Pf12) reveals
a unique interdomain organization and the potential for an antiparallel arrangement with Pf41. *J Biol Chem*
288, 12805-12817 (2013).
26.   Parker, M.L., Peng, F. & Boulanger, M.J. The Structure of Plasmodium falciparum Blood-Stage 6-Cys
Protein Pf41 Reveals an Unexpected Intra-Domain Insertion Required for Pf12 Coordination. *PLoS One* **10**,
e0139407 (2015).
27.   Kundu, P.*, et al.* Structural delineation of potent transmission-blocking epitope I on malaria antigen Pfs48/45.
Nat Commun **9**, 4458 (2018).
28.   Holm, L. & Rosenstrom, P. Dali server: conservation mapping in 3D. *Nucleic Acids Res* **38**, W545-549
(2010).