Degeneracy and stability in neural circuits of dopamine and serotonin neuromodulators: A theoretical consideration

Chandan K. Behera[1,^], Alok Joshi[1], Da-Hui Wang[2,3], Trevor Sharp[4] and KongFatt Wong-Lin[1,*]

[1]Intelligent Systems Research Centre, School of Computing, Engineering and Intelligent Systems, Ulster University, Derry~Londonderry, Northern Ireland, UK

[2]State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal University, Beijing, China

[3]School of Systems Science, Beijing Normal University, Beijing, China

[4]Department of Pharmacology, University of Oxford, Oxford, UK

*Corresponding author: KongFatt Wong-Lin (k.wong-lin@ulster.ac.uk)

^Current address: Yale School of Medicine, New Haven, Connecticut, USA

Author contributions: C.K.B. and K.W.-L. conceptualized and designed the study. C.K.B., A.J., D.-H.W. and K.W.-L. acquired the data and conducted analyses. K.W.-L. supervised the study. T.S. provided guidance on the study. C.K.B., A.J. and K.W.-L. wrote the first draft of the manuscript. All authors interpreted the data and revised the manuscript.

Competing Interest Statement: The authors declare that they have no competing interests.

Keywords: Degeneracy, computational modelling, serotonin, dopamine, reward and punishment

Abstract

Degenerate neural circuits perform the same function despite being structurally different. However, it is unclear whether neural circuits with interacting neuromodulator sources can themselves be degenerate while maintaining the same neuromodulatory function. Here, we address this by computationally modelling the neural circuits of neuromodulators serotonin and dopamine, local glutamatergic and GABAergic interneurons, and their possible interactions, under reward/punishment-based conditioning tasks. We show that a single parsimonious, sparsely connected neural circuit model can recapitulate several separate experimental findings, but not all, suggesting potentially multiple circuits acting in parallel. Using computational simulations and dynamical systems analysis, we demonstrate that several different stable circuit architectures can produce the same observed network activity profile. Further, simulating dopamine (D2) receptor agonist can distinguish among sub-groups of these degenerate networks; a testable model prediction. Overall, this theoretical work suggests the plausibility of degeneracy within neuromodulator circuitry and has important implication for the stable and robust maintenance of neuromodulatory functions.

Introduction

The nervous system can be modulated by endogenous chemical messengers, called neuromodulators [1]. Neurons or synapses, and hence neural circuits, that succumb to neuromodulation can change circuit configuration and function [2, 3]. It is also known that neural circuits can be degenerate, that is, circuits can comprise different elements and/or structure while performing the same function or yielding the same output [4]. This can allow robust maintenance of functions and behaviour in the face of changes in the underlying structure [5, 6]. Although it has been shown that neuromodulators can selectively regulate degenerate neural circuits [3, 4], it is unclear whether neural circuits with neuromodulator-containing neurons can themselves be degenerate, which could in turn provide stable widespread neuromodulator influences on targeted neural circuits (Fig. 1A).

In this theoretical study, we investigate the plausibility of degenerate and stable neuromodulator circuits by focusing on the neural circuits in the midbrain which are the source of ascending pathways of two highly studied monoaminergic neuromodulators, serotonin (5-hydroxytrptamine; 5-HT) and dopamine (DA). These neuromodulators have major roles in modulating cognition, emotion and behaviour, and are linked to the pathogenesis and pharmacological treatment of many common neuropsychiatric and neurological disorders [7]. The majority of 5-HT-producing neurons are found in the dorsal and median raphe nuclei (DRN and MRN), while most DA-producing neurons reside in the ventral tegmental area (VTA) and substantia nigra compacta (SNc) [7].

At the functional level, DA and 5-HT are known to play important roles in reward- and punishment- based learning and decision-making [8]. For example, there is strong evidence that DA neuronal activity signals reward prediction error (difference between predicted and actual reward outcome) to guide reinforcement learning [9]. Specifically, DA neurons are phasically excited upon unexpected reward outcome or reward-predictive cues, and inhibited upon unexpected reward omission or punishment [10-12], although there is heterogeneity amongst DA neurons in this regard [11, 13].

Fig. 1. Degenerate neuromodulator circuits constrained by observed neuronal signalling. A) Multiple neuromodulators that influence neural circuits, cognition and behaviour, may be embedded within degenerate neural circuits. Neuromod: Specific neuromodulator type. **B)** Schematic of DRN and VTA activity profiles in reward and punishment tasks. Activities (firing rates) aligned to timing of unexpected punishment outcome (left, vertical red dashed lines) and learned reward-predictive cue (right, vertical green dashed lines) and reward outcome (right, vertical red dashed lines). Top-to-bottom: VTA DA neural activity exhibits phasic excitation (inhibition) at reward-predictive cue (punishment) outcome (e.g. [11, 21]). VTA GABAergic neural activity shows phasic excitation upon punishment (e.g. [21, 22]), while exhibiting post-cue tonic activity which is not modulated by the presence/absence of actual outcome (e.g. [11]). DRN Type-I 5-HT neurons shows phasic activation by reward-predicting cue (right) but not punishment (left). DRN Type-II 5-HT neurons signal punishment outcome (left) and sustained activity towards expected reward outcome (right) (e.g. [13]). DRN GABAergic neurons have phasic activation upon punishment but have tonic inhibition during waiting and reward delivery (e.g. [16]). DRN glutamatergic neurons deduced to be excited by reward-predicting cue, in line with VTA DA neural activation [23], and assumed not to respond to punishment outcome. Baseline activity for Type-I 5-HT DRN neurons is higher in reward than punishment tasks (e.g. [13]).

In comparison to DA neurons, DRN 5-HT neurons exhibit greater complexity in function, with studies reporting that 5-HT neuronal activity encodes both reward and punishment. For instance, it has been shown that certain 5-HT neurons (labelled "Type I") were phasically activated only by reward predicting cues, but not punishment in a classical conditioning paradigm [13]. Yet, in the same study, a different population of 5-HT neurons ("Type II") signalled both expected reward and punishment with sustained elevated activity towards reward outcome [13]. The latter study also found that, unlike DA neurons, baseline firing of Type-I 5-HT neurons was generally higher in rewarding than punishment trials, and this effect lasted across many trials, suggesting information processing over a long timescale, possibly

used to perform meta-learning via learning rate modulation [14]. Similar 5-HT neuronal responses to reward and punishment were reported in other rodent [15-18] and non-human primate [19] studies. This differential responses of DA and 5-HT neurons to reward and punishment cannot be trivially reconciled with a simple model e.g. using two opposing neuromodulatory systems as previously proposed [20].

Other studies reveal further complexity in reward/punishment processing, specifically in the form of altered activity of non-5-HT/DA midbrain neurons. For example, DRN neurons utilising gamma-aminobutyric (GABA) were tonically inhibited during reward-waiting with further inhibition during reward acquisition, but phasically activated by aversive stimuli [16]. In contrast, GABAergic neuronal activity in the VTA exhibited sustained activity upon rewarding cue onset but no response to the presence or absence of actual reward outcome [11]. Further, other studies found that VTA GABAergic neuronal activity was potently and phasically activated by punishment outcome, which in turn inhibited VTA DA neuronal activity [21, 22]. Another study showed that glutamatergic (Glu) neurons in the DRN reinforced instrumental behaviour through VTA DA neurons [23].

This complexity of signalling within the DRN-VTA system in response to reward and punishment may reflect the DRN and VTA having shared afferent inputs [12, 24-29]. Another possibility is that the DRN and VTA interact with each other. Indeed, a growing number of studies have suggested that there are direct and indirect interactions among distinctive neuronal types between and within the DRN and VTA [20, 23-25, 27, 30-35].

Taken together, information of reward and punishment signalled by neuronal activities within the DRN and VTA seems to be diverse, heterogeneous, distributed and mixed. Some of these signalling responses are illustrated in Fig. 1B. This led to the following questions (Fig. 1A). First, can these experimental findings from separate studies be reconciled and understood in terms of a single, parsimonious neural circuit model encompassing both the DRN and VTA? Second, can there be several degenerate DRN-VTA circuits, which are stable, that can signal reward and punishment information as observed in experiments? Third, if there are degenerate and stable DRN-VTA circuits, can some of them be distinguishable from others, for example, through perturbative means?

To address these questions, we developed a biologically plausible DRN-VTA computational neural circuit model based on our previous multiscale modelling framework for neuromodulators [36, 37], constrained by some of the observed neuronal signalling profiles under reward and punishment tasks. The modelling considers known direct and indirect

pathways between DRN 5-HT and VTA DA neurons. Upon simulating the model under reward and punishment conditions, we found that many, but not all of the experimental findings, could be captured in a single, parsimonious DRN-VTA model. Further, several distinct model architectures could replicate the same neural circuit activity response profile, hence reflecting the possibility of degeneracy. Applying dynamical systems theory, we found that all these circuits were dynamically stable. To distinguish among these degenerated models, we simulated drug effects of DA D2-receptor based agonist and were able to distinguish sub-groups of these degenerate model architectures. Overall, our study provides theoretical support for degeneracy and stability of neural circuits of neuromodulators.

Results

DRN-VTA network modelling

To develop the DRN-VTA neural circuit models, we made use of our dynamic mean-field (neuronal population-based) modelling framework [36] for neuromodulator interactions. The modelling approach was constrained by data from known electrophysiological, neuropharmacological and voltammetry parameters (see [36] and below). Each neural circuit model architecture investigated consisted of DRN 5-HT, VTA DA, VTA GABA, DRN GABA, and DRN Glu neuronal populations. Direct and indirect interactions among these five neuronal populations were then explored. The main aim of this work was to evaluate the plausibility of neuromodulator circuit degeneracy and stability rather than replicate every neuronal populations in these brain regions.

Directed interaction between two neuronal populations, in which the source was a neuromodulator, was mathematically described by the neuromodulator neuronal population firing (rate) activity, followed by the release-and-uptake dynamics of the neuromodulator, which in turn induced certain population-averaged currents on the targeted neuronal population [36]. An induced current, I_x, which could be effectively excitatory or inhibitory depending on experimental findings, can be described by

$$\tau_x \frac{dI_x}{dt} = -I_x + \frac{k_x}{1+e^{-g_x([x]-[x]_0)}}$$

(1)

where x was some neuromodulator (5-HT or DA), τ_x the associated time constant, k_x some constant that determined the current amplitude, and constants g_x and $[x]_0$ that controlled the

slope and offset of the function on the right-hand-side of Eq. (1). The release-and-uptake dynamics for a neuromodulator x was described by

$$\frac{d[x]}{dt} = [x]_p R_x - \frac{V_{max,x}[x]}{K_{m,x}+[x]}$$

(2)

where $[x]$ was the concentration of x, $[x]_p$ the release per neural firing frequency [36], and constants $V_{max,x}$ and $K_{m,x}$ were constants determined from voltammetry measurements.

If the source of the interaction came from GABAergic (or Glu) neurons, then for simplicity, we assumed an instantaneous inhibitory (or excitatory) current-based synaptic influence on the targeted neuronal populations (see Materials and methods). This reflected the faster ionotropic receptor-based synaptic dynamics compared to metabotropic receptor-based neuromodulatory effects, while also reducing the number of free model parameters. Similarly, we also ignored the relatively faster neuronal membrane dynamics. Threshold-linear input-output function for each neuronal population was used [36], and described by

$$F = g[I - I_0]_+$$

(3)

where F was the neural population firing rate (output), I the total input current into a neural population, g was the slope, and I_0 some threshold current, and with $[z]_+ = z$ if $z \geq 0$, and 0 otherwise. Here, I is a function of the summed currents, including the neuromodulator-induced currents I_x's and ionotropic receptor-based currents (proportional to presynaptic neural firing rates; see Materials and methods). For simplicity, fast co-transmission of neurotransmitters was only considered in one modelling instance (co-release of 5-HT and glutamate via fast 5-HT$_3$ and ionotropic receptors) based on findings by [34]. From a computational modelling perspective, the DRN (Type I) 5-HT and Glu neuronal populations, which have rather similar activity profiles (Fig. 1B, 3rd and 6th rows) could also be effectively grouped and considered as a single 5-HT neuronal population that "co-transmit" both 5-HT and Glu to DA neurons [34].

For each model circuit architecture that we investigated, we considered separately the inclusion of either Type I or II 5-HT neurons in the circuit [13], and the possibility of excitatory and inhibitory projections from 5-HT to DRN Glu/GABA and DA neurons, and from DA to GABA neurons in the VTA. To allow tractability in the search for the many possible connectivity structures, the models' connections were largely based on known experimental evidence. For example, the connections between VTA GABAergic and DRN Glu neurons were not considered as, to date, there is little experimental support. We also focused only on learned

reward (with reward-predicting cue followed by reward outcome) and unexpected punishment conditions, simulated using a combination of tonic and/or phasic afferent inputs (Materials and methods). It should be noted that we did not consider other conditions and network learning effects (e.g. [8]) as the main aim was to demonstrate the plausibility of DRN-VTA circuit degeneracy and stability, constrained by observed neuronal signalling in reward and punishment tasks. In particular, we investigated various DRN-VTA circuit architectures with network activity profiles that closely resembled those in Figure 1B. Note also that all activity profiles in Fig. 1B were based on experimental studies except VTA Glu neural activity, which we deduced to be similar to that of DA neural activity in the reward task [23] and assumed to be non-responsive in the punishment task. See Materials and methods for further details regarding the modelling procedures, parameters, simulations, and analyses.

A DRN-VTA model can reconcile many signalling patterns

We began with a parsimonious, sparsely connected DRN-VTA model architecture and adjusted the strength of the afferent inputs and the internal connection weights of the network such that the network activity profiles attained qualitatively like some of the stereotypical profiles as illustrated in Fig. 1B (Materials and methods). Variations in some of the model parameters could still recapitulate these profiles, exhibiting model robustness (Supplementary Information, Table 1). The resultant network configuration (Fig. 2) was our most sparsely connected DRN-VTA circuit model. All other subsequent architectures considered would henceforth be derived from this basic architecture.

This minimal network architecture readily recapitulated many of the neuronal signalling changes in the DRN and VTA in separate experimental studies, both for reward (Fig. 3, black dashed) and punishment (Fig. 3, orange bold) tasks. Moreover, only modifications to the afferent inputs to DRN 5-HT and VTA GABA neurons (Materials and methods) were needed to replicate the signalling of Type I or II 5-HT neurons (Figs. 3A and B, respectively), while maintaining the same internal connectivity structure. For example, a lack of sustained reward-based activity of Type I 5-HT activity (Fig. 3B, 2nd row, black dashed) required additional external input to sustain VTA GABAergic neural activity (Fig. 2; Fig. 3A, bottom row, black dashed). It should be noted that this assumed that the activity profiles for these non-5-HT neurons were qualitatively similar, regardless of the 5-HT neuronal types (Fig. 2).

To understand across-trial reward versus punishment effects in the model, we implemented a higher constant excitatory input into both DRN 5-HT and VTA DA neurons under reward compared to punishment conditions (Fig. 2, long black arrows). This particular model required

differential inputs to 5-HT and DA neurons (Materials and methods) such that the overall tonic 5-HT neural activity was higher for reward than punishment trials, while DA neural activity remained unchanged (Fig. 3, black dashed vs orange bold lines in top two rows), again consistent with experimental observation (e.g. [13]). In the model, although both 5-HT and DA neurons directly received constant across-trial reward-based excitatory inputs, the indirect inhibitory pathway from 5-HT neurons through VTA GABA neurons onto VTA DA neurons nullified the overall effects on DA neurons (Figs. 2 and 3). In other words, increased firing of 5-HT neurons could be activating VTA GABAergic neurons to a level sufficient to inhibit VTA DA neurons and thereby cancelling out the net long-term reward signals (Fig. 2). Note that this is just one out of several possible models; e.g. a more complex case (with more model parameters) which we are not considering could be that the VTA DA, VTA, GABA and DRN 5-HT neurons individually receive very different inputs.

Fig. 2. A parsimonious, sparsely connected DRN-VTA circuit model. Grey: brain region. Coloured circle: neuronal population. Legend: network's afferent inputs. Model architecture implicitly encompasses either Type I or II 5-HT neurons with two different inputs for reward/punishment task (bright red arrows if Type I; blue arrows if Type II; black arrows denote common inputs for reward/punishment task for both Types). Circuit connections: triangular-end arrows (excitatory); circle-end arrows (inhibitory). Thicker arrows: stronger connection weights. Constant long-term reward inputs simultaneously to 5-HT and DA neurons to alter baseline activities. Sustained activity for expectation of reward outcome implemented with tonic input between cue and reward outcome. All other inputs are brief, at cue or reward/punishment outcome, producing phasic excitations/inhibitions. Note: Self-

inhibitory (self-excitatory) connections within GABAergic (Glu) neurons, and autoreceptor inhibitions within 5-HT or DA neurons were implemented but not shown here (see Materials and methods, and Supplementary Information, Fig. 1). This is the most basic, sparsely connected model architecture considered.

Fig. 3. DRN-VTA model replicates signalling patterns and suggests multiple parallel circuits. A-B) Model with reward (black dashed lines) and punishment (orange bold lines) tasks with 5-HT neurons that are of Type I **(A)** or Type II **(B)** Time label from cue onset. Green (red) vertical dashed-dotted lines: cue (outcome) onset time (as in Fig. 1B). Top-to-bottom: VTA DA, DRN 5-HT, DRN GABAergic, DRN Glu, and VTA GABAergic neural populations. **C)** Hypothesis for multiple different DRN-VTA circuits operating in parallel, which may consist of different clusters of neuronal sub-populations and different set of afferent inputs, leading to different outputs. Vertical dots denote the potential of having more than two distinctive circuits.

However, under these conditions, with both Type I and II 5-HT neurons, the baseline DRN and VTA GABAergic activities in the reward task were slightly different than those in the punishment task (Fig. 3, 3rd and 5th rows, black dashed vs orange bold), which have yet to be observed in experiments. The model's inability to recapitulate this specific phenomenon might perhaps suggest that there are more complex features in the system, such as further division of neuronal subgroups. This also holds for other model architectures (see below, Fig. 4). For example, it could be possible that a sub-population of DRN GABAergic neurons be directly connected to 5-HT neurons (as in Fig. 1B), but not another DRN GABAergic neuronal sub-

population, such that across-trial reward signal inputs are distributed differently than that of Fig. 1B. Moreover, high chemical and functional diversity amongst DRN 5-HT neurons is now well recognised [38]. Hence, it might be possible that there could exist multiple neural circuits with different circuit architectures operating in parallel, as illustrated in Fig. 3C. Another possibility could be due to mere different inputs from e.g. different sources.

In summary, we have shown that, under reward and punishment conditions, many of the observed signalling patterns in different DRN and VTA neuronal types could readily be reconciled within a single sparsely connected DRN-VTA circuit model. However, not all the signalling patterns could be captured, suggesting that multiple different neuronal sub-populations and circuits may be operating in parallel within the DRN-VTA system. Next, we shall investigate whether various different DRN-VTA neural circuits could produce the same output, i.e. be degenerate.

Multiple degenerate DRN-VTA circuits

To search for degenerate DRN-VTA circuit models, we evaluated various combinations of connections within and between the DRN and VTA, and where necessary, adjusted any afferent inputs (Materials and methods). We used the network activity profile template as illustrated in Figs. 3A and B as output "target" to check whether other different circuits could replicate similar activity profiles.

Given the variability of neuronal firing rates reported in the literature (Supplementary Information, Table 2), to be considered a degenerate neural circuit, we set an inclusion criterion that allowed the time-averaged percentage changes of the neural population firing activities to be within certain acceptable ranges as compared to the activities in Figs. 3A-B. Specifically, the time-averaged percentage change in activities of the DA, 5-HT, DRN GABA, VTA GABA, and Glu neural populations for any qualified (degenerate) circuits had to be less than 10%, 10%, 16%, 16% and 10% from those of the template activity profiles, respectively. These mean percentage changes were calculated within a 3 s time duration, from 1 s before cue onset to 1 s after outcome onset, encompassing both baseline and stimulus-evoked activities. Any neural circuit architecture which did not satisfy the inclusion criterion for any neural population activity was discarded. Note that here, we aimed to proof a concept and the

specific values of the inclusion criterion were theoretical and not critical - the general results, i.e. overall trends, remained even if the values were altered (not shown).

Various neural circuits were created by systematic addition and modification of connections of our parsimonious, sparsely connected DRN-VTA circuit model (now presented as model architecture 'k' in Fig. 4). For each model architecture, model parameters were searched, in both reward and punishment tasks using both Type I and II 5-HT neurons, until the activity outputs lie within the inclusion criterion. Both excitatory and inhibitory connections from DRN 5-HT to DRN Glu/GABA and VTA DA neurons were also explored. Once the inclusion criterion was met, the model parameter values were varied to check for robustness (Supplementary Information, Table 1). This was repeated for different model architecture until we reached a highly connectivity model structure (architecture 'a' in Fig. 4).

Based on this extensive search process, we obtained a total of 84 different neural circuit model architectures that recapitulated the activity profiles in Figs. 3A and B. Their high-level model architectures were illustrated in Fig. 4 (see also Fig. 5, and Supplementary Information, Tables 1 and 3). For instance, model architecture 'a' in Fig. 4 (see Supplementary Information, Fig. 1, for detailed architecture) actually consisted of 32 distinctive models with different 5-HT neuron types and connectivity signs (Supplementary Information, Table 3). Interestingly, this architecture's model parameters remained the same with either excitatory or inhibitory connections (Fig. 2, diamond connections; Supplementary Information, Table 1).

We observed that all models with connection from 5-HT to DA neurons to be relatively weak or not required, consistent with previous work [23] (but see below, [30, 34]). Further, a substantial number of other connections in the DRN-VTA model were identified to be redundant or relatively weak, at least in the context of the conditions investigated (Fig. 4). In particular, the relatively weak VTA DA-to-GABA connection was consistent with studies that showed either a weak or non-existent direct effect of VTA DA on VTA GABA neurons [39].

With these degenerate models, we could also investigate the effects of specific DRN-VTA connectivity in which there are mixed findings or lack of evidence in the literature. For example, the influence of 5-HT on DRN GABA neurons could be excitatory or inhibitory [40]. In the case when this connection was inhibitory, and not excitatory, we found it to involve several degenerate circuits (model architectures 'b' to 'l'). In another example, with model architecture 'b', when the connection from DRN 5-HT to DRN Glu neurons was inhibitory, it led to more restricted values in its connectivity strength (0-0.1 a.u.) and hence less robust than when it was excitatory (0-1 a.u.) (Supplementary Information, Table 1). It has been reported that there

are mixed findings of 5-HT on VTA DA neurons [31]. In our simulations, we found that an inhibitory connection from 5-HT to VTA DA neurons could lead to several degenerate models (architectures 'a'-'d', 'i' and 'l'). Note that in model 'l' (labelled with an extra asterisk in Fig. 4), we had also successfully simulated its fast connectivity version, mimicking either 5-HT₃ receptor mediated transmission or 5-HT-glutamate co-transmission [34].

In terms of compensatory network effects, we found that upon the removal of the 5-HT to DRN GABA connection (from architecture 'b' to 'c'), be it excitatory or inhibitory, the excitatory connection from DRN Glu to 5-HT neurons had to be weakened by ~16% to satisfy the inclusion criterion. With additional removal of the connection from VTA DA to VTA GABA neurons, we found that the strength of the inhibitory connection from 5-HT to DRN GABA neurons had to be increased by ~150% (architecture 'd'). Interestingly, unlike other models, only the models with architectures 'a' and 'l' had to include inhibitory connection from VTA GABA to DRN GABA neurons, in which such connection was observed in e.g. [35].

Fig. 4. Neural circuit model architectures with similar network activity profiles. The activity profiles were similar to that in Figs. 3A and B, satisfying the inclusion criterion. Model architectures 'a'-'k' denote architectures of decreasing connectivity, with Fig. 3 as architecture 'k'. Model architecture 'l' has an asterisk to denote that it was the only model with fast 5-HT to VTA DA connection, simulating fast 5-HT₃ or Glu receptor mediated connection or their combination (co-transmission). Architectures 'a' and 'l' have additional inhibitory input to DRN GABA neurons in reward task. All labels, connections and nomenclature have the same meaning as that in Fig. 2, except that, for simplicity, the relative connection

weights (thickness) are not illustrated, and the diamond-end arrows denote connections which are
either excitatory or inhibitory, with both explored. Self-connectivity not illustrated (see the general
example in Supplementary Information, Fig. 1 for a detailed version of model architecture 'a'). Note:
Each architecture consists of several distinctive model types (with a total of 84 types) with different 5-
HT neuronal or excitatory/inhibitory connectivity types (see Supplementary Information, Tables 1 and
3, and Fig. 5).

Degenerate DRN-VTA circuit models are dynamically stable

After identifying the theoretical existence of degenerate models, we used dynamical systems
theory to determine whether they were dynamically stable, i.e. whether (local) perturbation
from their steady states would eventually cause a return to their initial steady states (see
Materials and methods). Specifically, the stability of each neural circuit could be determined
by first finding the possible steady state(s) (i.e. fixed point(s)). This was achieved by setting
all the dynamical (differential) equations to zero and finding the algebraic solutions for the
dynamical variables (Materials and methods). Then the eigenvalues of the system's Jacobian
matrix at the steady states were computed (see Materials and methods for mathematical
derivation of the steady states and the Jacobian matrix). For a neural circuit to be dynamically
stable, the real part of all the eigenvalues associated with the steady state has to be negative.
This was exactly what we found for all the identified degenerate neural circuits in Fig. 4 (Fig.
5; Materials and methods), with no imaginary part in the eigenvalues.

Fig. 5A displayed the complete set of the real part of the eigenvalues for model #1 with
architecture 'a' in Fig. 4. This model had inhibitory connections from VTA DA to VTA GABA
neurons and from 5-HT to VTA DA/Glu neurons, using Type I 5-HT neurons and under
punishment conditions (Supplementary Information, Table 3). It was observed that the
eigenvalues with phasic input (blue) were generally larger, magnitude wise, than those with
tonic input (red). This was more pronounced for the eigenvalues with the largest magnitude
(maximal eigenvalues) (e.g. see Fig. 5A). Importantly, all the eigenvalues were negative,
indicating a dynamically stable network model even in the presence of additional phasic
stimulus input.

We repeated the analysis for all 84 models, under both phasic and tonic input conditions. This
analysis was presented in Fig. 5B only for the maximal eigenvalues (red circles and blue
crosses). The non-maximal eigenvalues had similar trends across the other models. In
general, with phasic activities (blue crosses), the models were more stable than with tonic

activities (red circles). However, during phasic activations, there were 18 models with rather

small (close to zero) eigenvalues (magnitude wise), albeit still negative. Hence, close to

instability. This was not observed for tonic activations, where the (most negative) eigenvalues

were found to hover within a small range of values (-0.017 to -0.016), except models with

architecture 'l' (~-0.03). In fact, the latter models, which were the only ones with a fast 5-HT-

to-DA connection (Fig. 5B, models #77-84), were the most stable under both phasic and tonic

conditions. Further, there was no difference identified between the excitatory (models #77-80)

and inhibitory (models #81-84) connections. Moreover, most of their eigenvalues in phasic

condition were substantially more negative than their tonic counterparts.

Fig. 5. Negative real eigenvalues at steady states of degenerate models. A) Complete set of the

real part of the eigenvalues for model #1 (Supplementary Information, Tables 1 and 3) with architecture

'a' in Fig. 4. Horizontal axis: Eigenvalues ranked from the largest to the smallest (magnitude wise). Blue

(red): More negative eigenvalues with phasic (blue) than tonic (red) input. Asterisk: Maximal eigenvalue

(largest magnitude) for each condition. **B)** For each of the 84 models, only the real part of the eigenvalue

with the largest magnitude is plotted under phasic (blue cross) and tonic (red circle) input condition.

Model architectures 'a' to 'l' refer to the different architectures as in Fig. 4, in which each has their own

distinctive model types (e.g. different 5-HT neuronal or excitatory/inhibitory connectivity types).

Eigenvalues for all model types have negative real parts, indicating dynamically stable. For most

models, the eigenvalues are generally more negative during phasic than tonic activities.

D$_2$ mediated drugs can distinguish some degenerate DRN-VTA circuits

Given the large number of degenerate and stable DRN-VTA circuits identified, how could one

distinguish among at least some of them? To address this, we investigated the neural circuit

responses to simulated dopaminergic D$_2$ receptor agonist due to the extensive D$_2$ receptor

mediated connectivity within the degenerate DRN-VTA circuits (Fig. 4; Supplementary

Information, Fig. 1). In particular, in the degenerate models, D$_2$ receptor mediated connections

involved those from VTA DA neurons to DRN 5-HT, DRN GABA and VTA GABA neurons, and
also the self-inhibitory connection (D_2 autoreceptor-mediated) of DA neurons (see Materials
and methods for references). To mimic the effects in the model of D_2 receptor agonist, we
gradually increased the strengths equally on the connections mediated by D_2 receptors
(Supplementary Information, Fig. 1, connections emanating from DA neurons) by some factor
(X) and observed how the network activity changed.

As we gradually increased the strengths of these specific sets of connection, subsets of the
degenerate models gradually behaved differently from the activity profile template in Figs. 3A
and B (Fig. 6A, model architectures embedded in non-black regions), i.e. not satisfying the
inclusion criterion, thus, possibly allowing us to distinguish them. (See Supplementary
Information, Tables 4-8 for details.) When at low dose with ten-fold (X=10) increase in D_2
mediated connections, all models showed substantial changes to their DA activities, beyond
the inclusion criterion, regardless of reward/punishment condition and 5-HT neuronal type
composition. Hence, they could not be distinguished at this dosage.

Fig. 6. D$_2$ receptor agonist can distinguish subsets of DRN-VTA neural circuits. Simulated drug
administered during punishment and reward tasks with efficacy factor X increments of 1, 10, 40, 70 and
100 times. Baseline condition: X=1. Colours other than black in the pie chart denote that at least one
neural population activity in specific model architecture(s) (labelled in yellow, and as in Fig. 4) has
deviated (increased or decreased) beyond the inclusion criterion. Specific colours denote changes in
specific neural population activities (See Supplementary Information, Tables 4-8 for details.)

For moderately higher D$_2$ agonist doses (X=40 and 70), models with architecture 'a' can be
distinguished from others, with substantial alteration to the DRN GABA and 5-HT activities
(Fig. 6A, light blue). Models with architectures 'b' and 'c' were also beginning to reveal
substantial activity changes in reward condition with Type II 5-HT neurons. Further, with an
increased factor of 70 under punishment condition, model with architectures 'd', 'e', 'h' and 'l'
could be distinguished from others with additional changes to their 5-HT activities. With a
factor of 100, a maximum of 5 subsets of model architectures could be distinguished based
on activity enhancement of different combination of neuronal types. An interesting observation
we found was that for the same composition of 5-HT neuronal type (I or II), the subsets of
distinguishable neural circuits were different between reward and punishment tasks.

Overall, our drug simulation predicted that a gradual increment of the level of D$_2$ receptor
activation could lead to differential enhancements of firing rate activities that could be used to
distinguish subsets of the degenerate DRN-VTA circuits.

116
117

Discussion

In this work, we have shown that neural circuits which are sources of neuromodulators can be
degenerate. In particular, the work focuses on computationally modelling and analysing DRN-
VTA circuits, as the DRN and VTA share structural and functional relationships among their
constituent neuronal types. In particular, these circuits are involved in the regulation of several
cognitive, emotional and behavioural processes, and implicated in many common and
disabling neuropsychiatric conditions [7]. Moreover, their neuronal signalling in reward and
punishment tasks are well studied [8].

In this work, we developed biologically based mean-field computational models of the DRN-
VTA circuit (Fig. 2) with several neuronal types, and tested them under classic conditions of
(learned) reward and (unexpected) punishment. The modelling was partially constrained by

known connectivity within and between the DRN and VTA regions, and their inputs from
multiple other brain regions, including mixed combinations of inputs [12, 24-29].

We found that a parsimonious, sparsely connected version of the DRN-VTA model could
reconcile many of the diverse phasic and tonic neural signalling events reported in the DRN
and VTA in punishment and reward tasks observed across separate experimental studies
(Figs. 1 and 3A-B). This model was evaluated using Type I and Type II 5-HT neurons in the
DRN as defined electrophysiologically in a previous study [13]. In the case of Type II 5-HT
neurons in reward condition, the model suggested that sustained 5-HT neuron activity
between cue and reward outcome [13] would lead to the gradual inhibition of DRN GABA
neuron activity and enhancement of VTA GABA neuron activity [11, 16]. The sparsely
connected model could also reproduce experimental observations [13] of an increase in
baseline firing of Type I 5-HT neurons across several trials in the rewarding task, without
similar effects on VTA DA neurons, or in the punishment task (Fig. 3). Thus, this specific model
suggested that slow, across-trial reward-based excitatory inputs could potentially be directly
targeted to both DRN 5-HT and VTA DA neurons, and that inhibitory 5-HT to GABA to DA
connectivity could cancel out the effects of the direct input to DA neurons, rendering only long
timescale changes to the baseline activity of 5-HT neurons but not DA neurons (Fig. 2). It
should however be noted that this is just one possible theoretical speculation offered out of
many. For instance, another trivial possibility would be simply to have different separate inputs
to DRN 5-HT and VTA DA and GABA neurons.

Despite the success of readily recapitulating the main observed phenomena, this relatively
simple model was unable to capture some specific activity profiles with Type I 5-HT neurons,
namely, the differential baseline activities of VTA GABA neurons and DRN GABA neurons
between reward and punishment conditions. We later found that this was the case even for
the more complex neural circuit models. Thus, perhaps additional neuronal populations and
DRN-VTA circuits could be operating in parallel (Fig. 3C) or it could be just be these neurons
receive different inputs from e.g. different sources. Such parallel circuits could be validated
experimentally in the future, for example, using gene-targeting of specific DRN and VTA
neuron subtypes and projections. Indeed, there is increasing evidence that DRN 5-HT neurons
are more chemically diverse than previously expected, and that there is a high level of
functional diversity in output pathways of the DRN and VTA [12, 24-29, 38-39, 41-45].

To demonstrate degeneracy in the DRN-VTA system, we showed that several variants of the
DRN-VTA circuit model could readily recapitulate the same neuronal signalling profiles, with
only occasional slight changes made to their afferent inputs (Fig. 4). Our results also suggest

experimental studies on the DRN and/or VTA system, e.g. using tracing methods, to search

for more than one neural circuit configuration.

Consistent with previous work [23], the degenerate models showed relatively weaker direct

connection from DRN 5-HT to VTA DA neurons as compared to the connection from DRN Glu

to VTA DA neurons. However, previous studies had also demonstrated a direct influence of

DRN 5-HT on VTA DA neuron activity (e.g. [31]). More recent work has shown that DRN 5-

HT terminals in the VTA co-release glutamate and 5-HT, eliciting fast excitation (via ionotropic

receptors) onto VTA DA neurons and increased DA release in the nucleus accumbens to

facilitate reward [34]. Hence, we also incorporated a model with fast 5-HT-to-DA connection

(model architecture 'I*' in Fig. 4) and found such circuit to be plausible in terms of capturing

the stereotypical reward and punishment signalling.

Next, we applied dynamical systems theory and showed that all the found degenerate circuits

were dynamically stable (Fig. 5). Interestingly, we found that model architecture 'I*' with fast

5-HT-to-DA connection was dynamically more stable than all other architectures investigated

(Fig. 5B). Future computational modelling work could explore the effects of co-transmission of

neurotransmitters on neural circuit degeneracy and functioning, and using more biologically

realistic spiking neuronal network models across multiple scales, e.g. [37, 45].

Finally, we simulated the gradual increase in dopaminergic D_2 receptor activation, due to

extensive D_2 mediated connections in the investigated circuits. This was done by increasing

the connection strengths emanating from the VTA DA neurons. This allowed us to distinguish

subsets of the degenerate DRN-VTA circuits by identifying substantial and differential

deviations in specific neural populations' activities (Fig. 6). Interestingly, we found that for the

same composition of 5-HT neuronal type (I or II), the subsets of distinguishable DRN-VTA

circuits were different between reward and punishment tasks (Fig. 6A). Future experimental

work could verify this model prediction.

From a more general perspective, our computational modelling and analytical framework

could be applied to the study of degeneracy and stability of neural circuits involving the

interactions of other neuromodulators such as norepinephrine/noradrenaline (e.g. [36]). It

should be also noted that in the development of the models, we had resorted to a minimalist

approach by focusing only on sufficiently simple neural circuit architectures that could replicate

experimental observations. Future modelling work may investigate the relative relevance of

these connections with respect to larger circuits involving cortical and subcortical brain regions

across multiple scales, especially during adaptive learning (e.g. [37, 46]).

Taken together, our computational modelling and analytical work suggests the existence of degeneracy and stability in DRN-VTA circuits. Some of these degenerate circuits can be dynamically more stable than others, and subsets of the degenerate circuits can be distinguished through pharmacological means. Importantly, our work opens up a new avenue of investigation on the existence, robustness and stability of neuromodulatory circuits, and has important implication on the stable and robust maintenance of neuromodulatory functions.

Materials and methods

Input-output functions of neural population firing rates

The computational models developed were based on our previous mean-field, neural population based modelling framework for neuromodulator circuits [36], in which the averaged concentration releases of neuromodulators (e.g. [5-HT]) were monotonic functions of the averaged firing rate of (e.g. 5-HT) neuronal populations via some neuromodulator induced slow currents. All 5 neural populations' firing rates were described by threshold-linear functions (general form in Eq. (3)) [36, 47]:

$$F_{5-HT} = g_{5-HT} \left[I_{5-HT} - I_{0,5-HT} \right]_+ \tag{4}$$

$$F_{DA} = g_{DA} \left[I_{DA} - I_{0,DA} \right]_+ \tag{5}$$

$$F_{Glu} = g_{Glu} \left[I_{Glu} - I_{0,Glu} \right]_+ \tag{6}$$

$$F_{GABA-DRN} = g_{GABA-DRN} \left[I_{GABA-DRN} - I_{0,GABA-DRN} \right]_+ \tag{7}$$

$$F_{GABA-VTA} = g_{GABA-VTA} \left[I_{GABA-VTA} - I_{0,GABA-VTA} \right]_+ \tag{8}$$

where $[z]_+ = z$ if $z \geq 0$, and 0 otherwise. The threshold input values for $I_{0,DA}$ was -10 (a.u.) for DA neurons, and $I_{0,5-HT}$ was 0.13 (a.u.) for 5-HT neurons, to allow spontaneous activities mimicking *in vivo* conditions (Fig. 3). 5-HT neurons had a threshold-linear function with gain value g_{5-HT} of about 1.7 times higher than that for DA neurons, and so we set their for DA and 5-HT neurons to be 0.019 and 0.033 (Hz), respectively (e.g. [45, 48-51]). For simplicity, we assumed the same current-frequency or input-output function in either tonic or phasic activity mode [36, 47].

Afferent currents and connectivity

The afferent current, I, for a neural population consisted of summed contributions from external excitatory inputs I_{ext} including those induced by reward or aversive stimuli, and

recurrent interactions with other neural populations (see below) e.g. $I_{5-HT,DA}$ for effective DA-

induced currents in 5-HT neurons. Additionally, for a neuromodulator population,

autoreceptor-induced current, I_{auto}, was included.

Due to limited experimental evidence, and to reduce the parameter search space, we did not

consider the following connections from: (i) DRN GABA to VTA DA neurons; (ii) DRN GABA

to VTA GABA neurons; (iii) VTA GABA to DRN Glu neurons; (iv, v) DRN Glu to DRN GABA

neurons, and vice versa; and (vi) VTA DA to DRN Glu neurons. Then the total (population-

averaged) afferent input currents to DA and 5-HT neurons were, respectively, described by

246
$$I_{DA} = -I_{auto,DA} \pm I_{DA,5-HT} + I_{DA,Glu} - I_{DA,GABA-VTA} + I_{DA,ext} \tag{9}$$

247
$$I_{5-HT} = -I_{auto,5-HT} + I_{5-HT,DA} + I_{5-HT,Glu} - I_{5-HT,GABA-DRN} - I_{5-HT,GABA-VTA} + I_{5-HT,ext}$$

248
$$\tag{10}$$

where the first terms on the right-hand sides of Eqs. (6) and (7) were autoreceptor-induced

self-inhibitory currents, the second terms were the 5-HT-to-DA (labelled with subscript $DA, 5-$

HT) and DA-to-5-HT (with subscript $5-HT,DA$) interactions, the third terms were excitatory

interactions from DRN Glu neurons, the fourth/fifth terms were inhibitory interactions from local

DRN/VTA GABAergic neurons, and the last terms were additional external constant biased

inputs from the rest of the brain and the influence of behaviourally relevant stimuli (due to

rewards or punishments; see below). 5-HT neurons have a possible additional negative

interaction from VTA GABA neurons (second last term on right-hand-side) [35]. Negative or

positive sign in front of each term indicated whether the interaction was effectively inhibitory

or excitatory. The $\pm$ sign indicated effectively excitatory $(+)$ or inhibitory $(-)$ interactions which

we investigated (Figs. 2, 4 and 5), given their mixed findings in the literature (see also Eqs.

(11-13)). This form of summed currents was consistent with some experimental evidence that

showed different afferents modulating the tonic and phasic activation (e.g. [28, 52]).

Similarly, the total (population-averaged) afferent current to the glutamatergic (Glu), and VTA

and DRN GABAergic neurons, can respectively be described by

264
$$I_{Glu} = I_{self,Glu} \pm I_{Glu,5-HT} + I_{Glu,ext} \tag{11}$$

265
$$I_{GABA-DRN} = -I_{self,GABA-DRN} \pm I_{GABA-DRN,5-HT}$$

266
$$+ I_{GABA-DRN,DA} - I_{GABA-DRN,GABA-VTA} + I_{GABA-DRN,ext}$$

267
$$\tag{12}$$

268
$$I_{GABA-VTA} = -I_{self,GABA-VTA} + I_{GABA-VTA,5-HT} \pm I_{GABA-VTA,DA} + I_{GABA-VTA,ext} \tag{13}$$

where, the subscript Glu denoted DRN Glu neural population, and subscripts $GABA - VTA$

and $GABA - DRN$ denoted GABAergic neural populations in the VTA and DRN, respectively.

The subscript $self$ denoted self-connection.

The averaged synaptic currents of non-5-HT/DA ionotropic glutamatergic and GABAergic

neurons, namely, $I_{DA,Glu}$, $I_{5-HT,Glu}$, $I_{DA,GABA-VTA}$, $I_{5-HT,GABA-VTA}$, $I_{5-HT,GABA-DRN}$, $I_{self,Glu}$,

$I_{self,GABA-DRN}$, $I_{self,GABA-VTA}$ and $I_{GABA-DRN,GABA-VTA}$ were typically faster than currents

induced by (metabotropic) 5-HT or DA currents. Thus, we assumed the former currents to

reach quasi-steady states and described [47] and represented by $I_{e/i} = \pm J_{e/i} F_{e/i}$, where the

subscript e/i denoted excitatory/inhibitory synaptic current, J the connectivity coupling

strength, F the presynaptic firing rate for neural population e/i, and the sign $\pm$ for excitatory

or inhibitory currents. Further, dimensionless coefficients or relative connectivity weights, $W's$

(with values ≥ 0), were later multiplied to the above neuromodulator induced current terms

(right-hand-side of terms in Eqs. (9-13); see Eqs. (20-24)). Both the $J's$ and $W's$ were allowed

to vary to fit the network activity profiles of Fig. 1B within certain tolerance ranges (see below)

while exploring different neural circuit architectures (Fig. 4) (see Supplementary Information,

Table 1 for specific values). The self-connection weights $J's$ within the DRN Glu, DRN GABA

and VTA GABA neurons were set at 0.5, 0.5, and 10 respectively, for all network activity's

response profiles.

Autoreceptor-induced currents were known to trigger relatively slow G protein-coupled

inwardly-rectifying potassium (GIRK) currents [53]. For 5-HT$_{1A}$ autoreceptors, the inhibitory

current $I_{5-HT,auto}$ was described by [36, 53, 54]

$$\tau_{auto,5-HT} \frac{dI_{auto,5-HT}}{dt} = -I_{auto,5-HT} + \frac{k_{auto,5-HT}}{1+e^{-g_{auto,5-HT}([5-HT]-[5-HT]_0)}} \tag{14}$$

291 and similarly, for DA autoreceptor induced inhibitory current $I_{auto,DA}$:

$$\tau_{auto,DA} \frac{dI_{auto,DA}}{dt} = -I_{auto,DA} + \frac{k_{auto,DA}}{1+e^{-g_{auto,DA}([DA]-[DA]_0)}} \tag{15}$$

where $\tau_{auto,5-HT}$ was set at 500 ms [36] and $\tau_{auto,DA}$ at 150 ms [55-57]. The threshold values

$[5-HT]_0$ and $[DA]_0$ were set at 0.1 μM. These parameters can be varied to mimic the effects

of autoreceptor antagonists/agonist [36]. The gains $g_{auto,5-HT}$ and $g_{auto,DA}$ were set at 10 μM^{-}

^{1} each, and $k_{auto,5-HT} = k_{auto,DA} = 80$ a.u.. These values were selected to allow reasonable

spontaneous neural firing activities and baseline neuromodulator concentration levels (see

below), and similar to those observed in experiments.

Similarly, we assumed sigmoid-like influence of $[5-HT]$ ($[DA]$) on DA (5-HT) neural firing

activities between the DRN and VTA populations such that the induced current dynamics

could be described by [36, 58]:

302
$$\tau_{DA,5-HT} \frac{dI_{DA,5-HT}}{dt} = -I_{DA,5-HT} + \frac{k_{DA,5-HT}}{1+e^{-g_{DA}([5-HT]-[5-HT]_1)}}$$
(16)

303
$$\tau_{5-HT,DA} \frac{dI_{5-HT,DA}}{dt} = -I_{5-HT,DA} + \frac{k_{5-HT,DA}}{1+e^{-g_{5-HT}([DA]-[DA]_1)}}$$
(17)

with the time constants $\tau_{5-HT,DA} = 1$ s, and $\tau_{DA,5-HT} = 1.2$ s [59-60]. We set $k_{5-HT,DA} = k_{DA,5-HT} = 0.03$ a.u. and $g_{5-HT} = g_{DA} = 20\ \mu M^{-1}$, $[5-HT]_1 = 0.3$nM, $[DA]_1 = 0.1$ nM such that the neural firing activities and baseline neuromodulator concentration levels were at reasonable values (see below) and similar to those in experiments (e.g. [61, 62]). For simplicity, we assumed Eqs. (16) and (17) to be applied equally to all targeted neural populations, but with their currents multiplied by their appropriate weights $w's$ (see above).

Release-and-reuptake dynamics of neuromodulators

The release-and-reuptake dynamics of 5-HT followed the form of a Michaelis-Menten equation [36, 61-63]:

$$\frac{d[5-HT]}{dt} = [5-HT]_p\ R_{5-HT} - \frac{V_{max,5-HT}\ [5-HT]}{K_{m,5-HT} + [5-HT]}$$
(18)

where $[5-HT]_p = 0.08$ nM was defined as the release per firing frequency [36, 63-64] (value selected to fit to reasonable baseline activities [62]; see below), and the Michaelis-Menten constants $V_{max,5-HT} = 1.3\ \mu M/s$ (maximum uptake rate) and $K_{m,5-HT} = 0.17\ \mu M$ (substrate concentration where uptake proceeds at half of maximum rate) were adopted from voltammetry measurements [62].

Similarly, the release-and-reuptake dynamics for DA was described by

$$\frac{d[DA]}{dt} = [DA]_p\ R_{DA} - \frac{V_{max,DA}\ [DA]}{K_{m,DA} + [DA]}$$
(19)

where $V_{max,DA} = -0.004\ \mu M/s$ and $K_{m,DA} = 0.15\ \mu M$ [65]. We set $[DA]_p = 0.1$ nM to constrain the ratio $[DA]_p/[5-HT]_p = 1.25$ [61, 62, 65]. For simplicity, we assumed Eqs. (18) and (19) to be applied equally to all targeted neural populations.

Reward and punishment conditions with Type I and Type II 5-HT neurons

To limit the size of the parameter search space, we focused on only the classical, fully learned reward conditioning task, and unexpected punishment task. For each simulated trial or realisation within a set of conditions (reward/punishment, excitatory/inhibitory connection), we set the cue onset time at 4.5 s to allow the network time to stabilise and reach steady state. The within-trial protocol for the external input current, I_{ext}, was implemented as a function of

time t as followed, depending on the simulated conditions. Note that, for simplicity, all external

input currents were assumed to be excitatory, regardless of reward or punishment task, unless

stated.

For reward task with Type-I 5-HT neurons: (i) constant input currents $I_{5-HT,ext}$ and $I_{DA,ext}$ to 5-

HT and VTA DA neurons, respectively, of amplitude $50\ a.u.$ to simulate long-term reward; (ii)

brief 0.2 s pulse $I_{Glu,ext}$ to DRN Glu neurons of amplitude $1000\ a.u.$ at 4.5 s; (iii) constant step

input current $I_{GABA-VTA,ext}$ to VTA GABAergic neurons with amplitude $200\ a.u.$ from 4.5 to 5.7

s to simulate sustained activity; and (iv) no external input $I_{GABA-DRN,ext}$ to DRN GABAergic

neurons. For punishment task with Type-I 5-HT neurons: (i) no external input to VTA DA, 5-

HT neurons and DRN Glu neurons; (ii) brief 0.2 s pulse to VTA (I_{app}) and DRN GABAergic

(I_{app5i}) neurons with amplitude $1000\ a.u.$ at 5.7 s.

For reward task with Type II 5-HT neurons: (i) constant input current to 5-HT and VTA DA

neurons with amplitude 50 a.u. to simulate long-term reward; (ii) additional step input current

to 5-HT neurons with amplitude 100 a.u. from 4.5 to 5.7 s to simulate sustained activity; (iii)

brief 0.2 s pulse to DRN Glu neurons of 1000 a.u. at 4.5 s; (iv) no external input to VTA and

DRN GABAergic neurons. For punishment task with Type II 5-HT neurons: (i) no external input

to VTA DA and GABAergic neurons, and DRN Glu neurons; (ii) brief 0.2 s pulse to DRN

GABAergic and 5-HT neurons with amplitude of 1000 a.u. at 5.7 s.

In addition, for transient inputs, we have used multiplicative exponential factors $\exp(-t/\tau)$ with

τ of 50 ms to smooth out activity timecourses (e.g. afferent synaptic filtering), but they do not

affect the overall results. To simulate long-term, across-trial reward/punishment signalling, we

assumed a higher constant excitatory input to both 5-HT and DA neurons in reward than

punishment trials. When searching for the neural circuit architecture using either Type I or II

5-HT neurons, we limited ourselves as much as possible to the same internal DRN-VTA circuit

structure. This also reduced the complexity of the parameter search space.

In reward task, to replicate a sustained Type II 5-HT neuronal reward signalling between cue

onset and reward outcome (Fig. 3B, 2nd row, black dashed) [13, 15, 16], tonic excitatory input

to DRN Type II 5-HT was implemented. The resulting sustained 5-HT activity led to gradual

suppression of DRN GABAergic activity (Fig. 3B, 3rd row, black dashed) [15-16] but also

gradual rise in VTA GABAergic activity regardless of reward outcome (Fig. 3B, last row, black

dashed) [11, 22]. The effects on these two GABAergic populations were respectively due to

5-HT's inhibitory connection to DRN GABAergic neurons and excitatory connection to VTA

GABAergic neurons. With regards to the latter, evidence of such excitatory influence via 5-

HT$_{2C}$ receptors was reported [33].

Excitatory connection from DRN Glu to VTA DA neurons was particularly strong in the models

(Fig. 2). This, together with no (or weak) 5-HT-to-DA connection, was consistent with previous

work [23]. In reward task, brief excitatory input to the DRN Glu neuronal population led to its

reward-sensitive phasic activation (Figs. 3A-B, 4th row, black dashed). Alternatively, we could

have directly implemented phasic excitatory input to the VTA DA neurons. In any case, the

phasic activation of DA activity was consistent with the reported DA neuronal response to a

fully learned reward-predicting cue [10-12].

In punishment task with Type I 5-HT neurons, brief excitatory inputs to GABAergic neurons in

the DRN and VTA led to their punishment-based phasic activation (Fig. 2; Fig. 3A, 3rd and 5th

rows, orange) [16, 21]. To replicate the activity profiles for a circuit with Type II 5-HT neurons,

brief excitatory inputs to the DRN GABAergic and 5-HT neurons were implemented instead,

leading to punishment-based phasic activation for both [11, 16, 21] (Fig. 2; Fig. 3B, 2nd and 3rd

rows, orange). With both Type I and II 5-HT neurons, there was also phasic inhibition of VTA

DA activity via VTA GABAergic neurons (Fig. 3, 1st row, orange), in line with findings from

previous studies [10-12, 21]. It should be noted that with Type II 5-HT neurons, the phasic

activation of VTA GABAergic neurons was not as potent due to the filtering by the slow

excitatory connection from 5-HT to VTA GABA neurons. Alternatively, a phasic input might

instead be sent to VTA GABAergic neurons, leading to a more potent activation of the latter.

Baseline neural activities and inclusion criterion

We define the neural circuit activities under baseline condition (right before cue onset) to follow

that in Figs. 3A-B. Namely, the baseline firing rates for 5-HT, DRN GABA, DRN Glu, DA and

VTA GABA neurons in punishment task were 3.0, 21.5, 4.1, 4.8 and 13.5 Hz, and those in

reward task were 4.5, 19.4, 4.1, 4.8 and 16.3 Hz, respectively (compared with Supplementary

Information, Table 2). Baseline $[5 - HT]$ and $[DA]$ levels were constrained to be at 10 and 1.5

nM, respectively. However, it is known that these activities can vary widely across subjects,

species and studies (Supplementary Information, Table 2).

While searching for degenerate neural circuit architecture or investigating substantial changes

due to simulated D_2 agonist, we had to define acceptable ranges of neural activities to evaluate

whether the variant neural circuit still behaved similarly to that of the model in Figs. 3A-B.

Specifically, an inclusion criterion was set that evaluated the time-averaged percentage

change in the neural population activities of any new model with respect to that of the template

activity profiles (Figs. 3A-B); the time-averaged percentage change in activities of the DA, 5-

HT, DRN GABA, VTA GABA, and Glu neural populations for any qualified (degenerate)

circuits had to be less than 10%, 10%, 16%, 16% and 10% from those of the template activity

profiles, respectively. These mean percentage changes were calculated within a 3 s time
duration, from 1 s before cue onset to 1 s after outcome onset, encompassing both baseline
and stimulus-evoked activities. Any neural circuit architecture which did not satisfy the
inclusion criterion for any neural population activity was discarded when in search of
degenerate neural circuits, or was highlighted (embedded in non-black region in Fig. 6A) when
investigating simulated D_2 agonist (see below). As this part of the study was to proof a concept,
specific values of the inclusion criterion were theoretical and not critical - the general results,
i.e. overall trends, remained even if these values were altered.

Simulating the effects of D_2 agonist

DA and 5-HT induced currents can lead to overall excitatory or inhibitory effects, depending
on receptor subtype(s) and the targeted neurons. In particular, DA enhances VTA GABAergic
neuronal activity via D_2 receptors and depolarizes the membrane of 5-HT neurons (e.g. [56,
59, 60, 66, 67]). DA also regulates the activity of other DA neurons via D_2 auto-inhibitory
receptors [68]. To study how D_2 receptor mediated drugs can affect DRN-VTA architecture
differently, we simulated different D_2 agonist dosage levels simultaneously by multiplying the
connection weights of D_2 receptor mediated currents (see above; Fig. 4; orange connections
in Supplementary Information, Fig. 1) by a factor 'X' (Fig. 6) of 10, 40, 70 and 100 times (default
of X=1). Then for each dosage, we separately observed the deviation in activity profiles for
each neural population with respect to the allowed range. Again, we considered the time-
averaged percentage changes due to the drug to be substantial if the mean percentage
change in at least one neural population activity had violated the inclusion criterion (see above,
and Supplementary Information, Tables 2 and 4-8).

Numerical simulations and codes

Simulations were performed using MATLAB with forward Euler numerical integration method
on the dynamical (ordinary differential) equations (see above). A simulation time step of 1 ms
was used and smaller time steps did not affect the results. MATLAB were used for analyses
of network stability and sensitivity. Source codes and generated data are available from the
corresponding author on reasonable request.

Network stability analysis

The 5 neural population firing rates (Eqs. (4-8)), when combined with their associated afferent
currents (Eqs. (9-13)) with explicit parameter values and relative connectivity weights, W's

and J's (Supplementary Information, Fig. 1 and Table 1), can be rewritten, respectively, as:

$F_{5-HT} = g_{5-HT}[-W_{auto,5-HT} I_{auto,5-HT} + W_{5d} I_{5-HT,DA} + J_{55e} I_{5-HT,Glu} - J_{55i} I_{5-HT,GABA-DRN} -$

$\qquad J_{5ii} I_{5-HT,GABA-VTA} + 99.87 + I_{5-HT,ext}]_+$ (20)

$F_{DA} = g_{DA}[-W_{auto,DA} I_{auto,DA} \pm W_{d5} I_{DA,5-HT} + J_{d5e} I_{DA,Glu} - J_{di} I_{DA,VTA-GABA} + 210 +$

$I_{DA,ext}]_+$ (21)

$F_{Glu} = g_{Glu}[J_{self,Glu} F_{Glu} \pm W_{5e5} I_{Glu,5-HT} + 100 + I_{Glu,ext}]_+$ (22)

$F_{GABA-DRN} =$

$\qquad g_{GABA-DRN}[-J_{self,GABA-DRN} F_{GABA-DRN} \pm$

$\qquad W_{5i5} I_{GABA-DRN,5-HT} + W_{5id} I_{GABA-DRN,DA} - W_{5ii} I_{GABA-DRN,GABA-VTA} + 450 +$

$\qquad I_{GABA-DRN,ext}]_+$ (23)

$F_{GABA-VTA} =$

$\qquad g_{GABA-VTA}[-J_{self,GABA-VTA} F_{GABA-VTA} + W_{i5} I_{GABA-VTA,5-HT} \pm W_{id} I_{GABA-VTA,DA} +$

$\qquad 200 + I_{GABA-VTA,ext}]_+$ (24)

We also inserted the explicit parameter values to the dynamical equations (Eqs. (14-19)) to

obtain:

$$500\frac{dI_{auto,5-HT}}{dt} = -I_{auto,5-HT} + \frac{80}{1+e^{-10([5-HT]-0.1)}}$$ (25)

$$150\frac{dI_{auto,DA}}{dt} = -I_{auto,DA} + \frac{80}{1+e^{-10([DA]-0.1)}}$$ (26)

$$1200\frac{dI_{DA,5-HT}}{dt} = -I_{DA,5-HT} + \frac{0.03}{1+e^{-20([5-HT]-0.1)}}$$ (27)

$$1000\frac{dI_{5-HT,DA}}{dt} = -I_{5-HT,DA} + \frac{0.03}{1+e^{-20([DA]-0.3)}}$$ (28)

$$\frac{d[5-HT]}{dt} = \frac{0.08F_{5-HT}}{1000} - \frac{0.0013[5-HT]}{0.17+[5-HT]}$$ (29)

$$\frac{d[DA]}{dt} = \frac{0.1F_{DA}}{1000} - \frac{0.004[DA]}{0.15+[DA]}$$ (30)

To check for network stability for each of the considered degenerate neural circuits, we first

find each network's steady state (or fixed point) by setting the rate of change for all the

dynamical equations to zero, i.e. $\frac{dI_{auto,5-HT}}{dt} = \frac{dI_{auto,DA}}{dt} = \frac{dI_{DA,5-HT}}{dt} = \frac{dI_{5-HT,DA}}{dt} = \frac{d[5-HT]}{dt} =$

$\frac{d[DA]}{dt} = 0$, and then solving them algebraically. The solution of these equations will give the

steady-state value (equilibrium point) of the system. Specifically, the currents (dynamical

variables) from Eqs. (25-28) (e.g. $I_{auto,5-HT} = \frac{80}{1+e^{-10([5-HT]-0.1)}}$) were substituted into Eqs. (20-

24). See Supplementary Information, Supplementary Method, for more detailed mathematical

derivation.

Using only the linear parts of the above threshold-linear functions (which were validated post-

hoc), and after some algebraic manipulations, we obtained the following system of equations,

in matrix form:

$$\begin{pmatrix} F_{Glu} \\ F_{GABA-VTA} \\ F_{GABA-DRN} \end{pmatrix}$$

$$= \begin{pmatrix} 1 - g_{Glu} J_{self-Glu} & 0 & 0 \\ 0 & 1 + g_{GABA-VTA} J_{self,GABA-VTA} & 0 \\ 0 & g_{GABA-DRN} J_{5ii} & 1 + g_{GABA-DRN} J_{self,GABA-DRN} \end{pmatrix}^{-1}$$

$$\begin{pmatrix} \frac{0.03\ W_{5e5} g_{Glu}}{1+e^{-20([5-HT]-0.1)}} + \frac{0}{1+e^{-16([DA]-0.1)}} + g_{Glu}(100 + I_{Glu,ext}) \\ \frac{0.03\ W_{i5} g_{GABA-VTA}}{1+e^{-20([5-HT]-0.1)}} + \frac{0.03\ W_{id} g_{GABA-VTA}}{1+ae^{-20([DA]-0.1)}} + g_{GABA-VTA}(200 + I_{GABA-VTA}) \\ \frac{0.03 W_{5i5} g_{GABA-DRN}}{1+e^{-20([5-HT]-0.1)}} + \frac{0.03\ W_{5id} g_{GABA-DRN}}{1+ae^{-20([DA]-0.1)}} + g_{GABA-DRN}(450 + I_{GABA-DRN}) \end{pmatrix} \quad (31)$$

$$\begin{pmatrix} F_{5-HT} \\ F_{DA} \end{pmatrix} = \begin{pmatrix} g_{5-HT}\left(\frac{-80}{1+e^{-10([5-HT]-0.1)}} + \frac{0.03\ W_{5d}}{1+ae^{-20([DA]-0.1)}} + 99.87 + I_{5-HT,ext} \right) \\ -g_{DA}\left(\frac{0.03\ W_{d5}}{1+e^{-20([5-HT]-0.1)}} + \frac{80}{1+e^{-10([DA]-0.1)}} - 210 - I_{DA,ext} \right) \end{pmatrix} +$$

$$\begin{pmatrix} g_{5-HT} & 0 \\ 0 & g_{DA} \end{pmatrix} \begin{pmatrix} J_{55e} & -J_{5i} & -J_{55i} \\ J_{d5e} & -J_{di} & 0 \end{pmatrix} \begin{pmatrix} F_{Glu} \\ F_{GABA-VTA} \\ F_{GABA-DRN} \end{pmatrix} \quad (32)$$

But by setting Eqs. (29) and (30) to be zero, we obtained, $F_{DA} = \frac{40[DA]}{0.15+[DA]}$ and $F_{5-HT} =$

$\frac{16.25[5-HT]}{0.17+[5-HT]}$ at steady state. Substituting these two into Eqns. (31) and (32), we can solve for

the values of [DA] and [5-HT] at steady state.

Next, to check whether the system is stable at a steady state, we compute the 5-by-5 Jacobian

matrix $M_{Jacobian}$ [69] for the 5 dynamical equations (Eqs. (25-30)), which can be computed

using partial derivatives on the right-hand-side of these equations:

$M_{Jacobian} =$

$$\begin{pmatrix} -1/500 & 0 & 0 & 0 & \frac{80}{500}\frac{10e^{-10([5-HT]-0.1)}}{(1+e^{-10([5-HT]-0.1)})^2} & 0 \\ 0 & -1/150 & 0 & 0 & 0 & \frac{80}{150}\frac{10e^{-10([DA]-0.1)}}{(1+e^{-10([DA]-0.1)})^2} \\ 0 & 0 & -1/1200 & 0 & \frac{0.03}{1200}\frac{20e^{-20([5-HT]-0.1)}}{(1+e^{-20([5-HT]-0.1)})^2} & 0 \\ 0 & 0 & 0 & -1/1000 & 0 & \frac{0.03}{1000}\frac{20e^{-20([DA]-0.3)}}{(1+e^{-20([DA]-0.3)})^2} \\ \frac{0.008}{1000}g_{5-HT} & 0 & 0 & \frac{0.08}{1000}g_{5-HT} & -0.0013 + \frac{0.0013[5-HT]}{(0.17+[5-HT])^2} & 0 \\ 0 & \frac{0.1}{1000}g_{DA} & -\frac{0.1}{1000}g_{DA} & 0 & & -\frac{0.004}{0.15+[DA]} + \frac{0.004[DA]}{(0.15+[DA])^2} \\ & & & & 0 & \end{pmatrix}$$

483   (33)

The eigenvalues of this Jacobian matrix were computed for each steady state for each model under each simulated condition (e.g. reward/punishment task). If the real parts of all the eigenvalues of the Jacobian matrix were negative at a given steady state, then the model was considered to be dynamically stable at that steady state [69].

Acknowledgments

CKB was supported by Ulster University Research Challenge Fund. AJ, TS and KFW-L were supported by BBSRC (BB/P003427/1). KFW-L and D-HW were jointly supported by The Royal Society – NSFC International Exchanges. KFW-L was further supported by COST Action Open Multiscale Systems Medicine (OpenMultiMed) supported by COST (European Cooperation in Science and Technology), and Northern Ireland Functional Brain Mapping Facility (1303/101154803) funded by Invest NI and the University of Ulster. D-HW received further support by NSFC under grants 31671077 and 31511130066.

References

1. Kaczmarek LK, Levitan IB. Neuromodulation : The biochemical control of neuronal excitability. New York: Oxford University Press; 1987.
2. Marder E. Neuromodulation of neuronal circuits: back to the future. Neuron. 2012;76(1): 1–11.
3. Marder E, O'Leary T, Shruti S. Neuromodulation of circuits with variable parameters: single neurons and small circuits reveal principles of state-dependent and robust neuromodulation. Annu Rev Neurosci. 2014;37: 329–346.
4. Cropper EC, Dacks AM, Weiss KR. Consequences of degeneracy in network function. Curr Opin Neurobiol. 2016;41: 62–67.
5. Edelman GM, Gally JA. Degeneracy and complexity in biological systems. Proc Nat Acad Sci USA. 2001;98(24): 13763–13768.
6. Whitacre JM. Degeneracy: a link between evolvability, robustness and complexity in biological systems. Theor Biol Med Model. 2010:7:6. doi:10.1186/1742-4682-7-6.
7. Müller CP, Cunningham KA. Handbook of the Behavioral Neurobiology Serotonin. Academic Press; 2020.
8. Hu H. Reward and aversion. Annu Rev Neurosci. 2016;39: 297-324.
9. Doya K. Metalearning and neuromodulation. Neural Netw. 2002;15(4–6): 495–506.

10. Schultz W, Dayan P, Montague PR. A neural substrate of prediction and reward. Science.
1997;275(5306): 1593–1599.

11. Cohen J, Haesler S, Vong L, Lowell BB, Uchida N. Neuron-type specific signals for reward
and punishment in the ventral tegmental area. Nature. 2012;482(7383): 85-88.

12. Watabe-Uchida M, Eschel N, Uchida N. Neural circuitry of reward prediction error. Annu
Rev Neurosci. 2017;40: 373-394.

13. Cohen JY, Amoroso MW, Uchida N. Serotonergic neurons signal reward and punishment
on multiple timescales. eLife. 2015;4:e06346. doi:10.7554/eLife.06346

14. Grossman CD, Bari BA, Cohen JY. Serotonin neurons modulate learning rate through
uncertainty. Curr Biol. 2022;32(3): 586-599.

15. Liu Z, Zhou J, Li Y, Hu F, Lu Y, Feng Q, Zhang J-E, Wang D, Zeng J, Bao J, Kim JY, Chen
Z-F, Mestikaway SE, Luo M. Dorsal raphe neurons signal reward through 5-HT and
glutamate, Neuron. 2014;81(6): 1360–1374.

16. Li Y, Zhong W, Wang D, Feng Q, Liu Z, Zhou J, Jia C, Hu F, Zeng J, Guo Q, Fu L, Luo M.
Serotonin neurons in the dorsal raphe nucleus encode reward signals. Nat Commun.
2016;7:10503. doi: 10.1038/ncomms10503.

17. Zhong W, Li Y, Feng Q, Luo M. Learning and stress shape the reward response patterns
of serotonin neurons. J Neurosci. 2017;37(37): 8863-8875.

18. Matias S, Lottem E, Dugué GP, Mainen ZF. Activity patterns of serotonin neurons
underyling cognitive flexibility. eLife. 2017;6:e20552. doi: 10.7554/eLife.20552.

19. Hayashi K, Nakao K, Nakamura K. Appetitive and aversive information coding in the
primate dorsal raphé nucleus, J Neurosci. 2015;31(15): 6195-6208.

20. Boureau YL, Dayan P. Opponency revisited: Competition and cooperation between
dopamine and serotonin. Neuropsychopharmacology. 2011;36(1): 74–97.

21. Tan KR, Yvon C, Turiault M, Mirzabekov JJ, Doehner J, Labouèbe G, Deisseroth K, Tye
KM, Lüscher C. GABA neurons of the VTA drive conditioned place aversion. Neuron.
2012;73(6): 1173–1183.

22. Eshel N, Bukwich M, Rao V, Hemmelder V, Tian J, Uchida N. Arithmetic and local circuitry
underlying dopamine prediction errors. Nature. 2015;525: 243–246. doi:
10.1038/nature14855.

23. McDevitt RA, Tiran-Cappello A, Shen H, Balderas I, Britt JP, Marino RAM, Chung SL,
Richie CT, Harvey BK, Bonci A. Serotonergic versus non-serotonergic dorsal raphe
projection neurons: differential participation in reward circuitry. Cell Rep. 2014;8(6): 1857-
1869.

24. Watabe-Uchida M, Zhu L, Ogawa SK, Vamanrao A, Uchida N. Whole-brain mapping of
direct inputs to midbrain dopamine neurons. Neuron. 2012;74: 858–873.

25. Ogawa SK, Cohen JY, Hwang D, Uchida N, Watabe-Uchida M. Organization of
monosynaptic inputs to the serotonin and dopamine neuromodulatory systems. Cell Rep.
2014;8: 1105–1118.
26. Dorocic IP, Furth D, Xuan Y, Johansson Y, Pozzi L, Silberberg G, Carlen M, Meletis K. A
whole-brain atlas of inputs to serotonergic neurons of the dorsal and median raphe nuclei.
Neuron. 2014;83(3): 663-678.
27. Beier KT, Steinberg EE, DeLoach KE, Xie S, Miyamichi K, Schwarz L, Gao XJ, Kremer
EJ, Malenka RC, Luo L. Circuit architecture of VTA dopamine neurons revealed by
systematic input-output mapping. Cell. 2015;162(3): 622-634.
28. Tian J, Huang R, Cohen JY, Osakada F, Kobak D, Machens CK, Callaway EM, Uchida N,
Watabe-Uchida M. Distributed and mixed information in monosynaptic inputs to dopamine
neurons. Neuron. 2016;91: 1374-1389.
29. Ogawa SK, Watabe-Uchida M. Organization of dopamine and serotonin system:
anatomical and functional mapping of monosynaptic inputs using rabies virus. Pharmacol
Biochem Behav. 2018;174: 9-22.
30. Di Giovanni G, Di Matteo V, Pierucci M, Esposito E. Serotonin-dopamine interaction:
electrophysiological evidence. Serotonin-Dopamine Interaction: Experimental Evidence
and Therapeutic Relevance. Prog Brain Res. 2008;172.
31. De Deurwaerdère P, Di Giovanni G. Serotonergic modulation of the activity of
mesencephalic dopaminergic systems: Therapeutic implications. Prog Neurobiol.
2017;151: 175–236.
32. Xu P, He Y, Cao X, Valencia-Torres L, Yan X, Saito K, Wang C, Yang Y, Hinton Jr A, Zhu
L, Shu G, Myers Jr MG, Wu Q, Tong Q, Heisler LK, Xu Y. Activation of serotonin 2C
receptors in dopamine neurons inhibits binge-like eating in mice. Biol Psychiatry.
2017;81(9): 737-747.
33. Valencia-Torres L, Olarte-Sanchez CM, Lyons DJ, Georgescu T, Greenwald-Yarnell M,
Myers Jr MG, Bradshaw CM, Heisler LK. Activation of ventral tegmental area 5-HT2C
receptors reduces incentive motivation. Neuropsychopharmacology. 2017;42(7): 1511-
1521.
34. Wang HL, Zhang S, Qi J, Wang H, Cachope R, Mejias-Aponte CA, Gomez JA, Mateo-
Semidey GE, Beaudoin GMJ, Paladini CA, Cheer JF, Morales M. Dorsal raphe dual
serotonin-glutamate neurons drive reward by establishing excitatory synapses on VTA
mesoaccumbens dopamine neurons. Cell Rep. 2019;26(5): 1128-1142.
35. Li Y, Li CY, Xi W, Jin S, Wu ZH, Jiang P, Dong P, He XB, Xu FQ, Duan S, Zhou YD, Li
XM. Rostral and caudal ventral area GABAergic inputs to different dorsal raphe neurons
participate in opioid dependence. Neuron. 2019;101(4): 748-761.

36. Joshi A, Youssofzadeh V, Vemana V, McGinnity TM, Prasad G, Wong-Lin K. An integrated
modelling framework for neural circuits with multiple neuromodulators. J R Soc Interface.
2017;14(126): 20160902. doi: 10.1098/rsif.2016.0902.
37. Wong-Lin K, Wang D-H, Moustafa AA, Cohen JY, Nakamura K. Toward a multiscale
modeling framework for understanding serotonergic function. J Psychopharmacol.
2017;31(9): 1121-1136.
38. Okaty BW, Commons KG, Dymecki SM. Embracing diversity in the 5-HT neuronal
system. Nat Rev Neurosci. 2019;20(7): 397-424.
39. Morales M, Margolis EB. Ventral tegmental area: cellular heterogeneity, connectivity and
behaviour. Nat Rev Neurosci. 2017;18(2): 73-85.
40. Hernández-Vázquez F, Garduño J, Hernández-López S. GABAergic modulation of
serotonergic neurons in the dorsal raphe nucleus. Rev Neurosci. 2019;30(3): 289-303,
2019. doi: 10.1515/revneuro-2018-0014.
41. Weissbourd B, Ren J, DeLoach KE, Guenthner CJ, Miyamichi K, Luo L. Presynaptic
partners of dorsal raphe serotonergic and GABAergic neurons. Neuron. 2014;83: 645-662.
42. Fernandez SP, Cauli B, Cabezas C, Muzerelle A, Poncer J-C, Gaspar P. Multiscale single-
cell analysis reveals unique phenotypes of raphe 5-HT neurons projecting to the forebrain.
Brain Struct Funct. 2016;221(8): 4007-4025.
43. Zhou L, Liu M-Z, Li Q, Deng J, Mu D, Sun Y-G. Organization of functional long-range
circuits controlling the activity of serotongergic neurons in the dorsal raphe nucleus. Cell
Rep. 2017;18: 3018-3032.
44. Ren J, Friedman D, Xiong J, Liu CD, Ferguson BR, Weerakkody T, DeLoach KE, Ran C,
Pun A, Sun Y, Weissbourd B, Neve RL, Huguenard J, Horowitz MA, Luo L. Anatomically
defined and functionally distinct dorsal raphe serotonin sub-systems. Cell. 2018;175(2):
472-487.
45. Wong-Lin K, Joshi A, Prasad G, McGinnity TM. Network properties of a computational
model of the dorsal raphe nucleus. Neural Netw. 2012;32: 15–25.
46. Zhou H, Wong-Lin K, Wang D-H. Parallel excitatory and inhibitory neural circuit pathways
underlie reward-based phasic neural responses. Complexity. 2018;2018:4356767. doi:
10.1155/2018/4356767.
47. Jalewa J, Joshi A, McGinnity TM, Prasad G, Wong-Lin K, Hölscher C. Neural circuit
interactions between the dorsal raphe nucleus and the lateral hypothalamus: an
experimental and computational study. PLoS One. 2014;9(2): 1–16. doi:
10.1371/journal.pone.0088003.
48. Shepard PD, Bunney BS. Repetitive firing properties of putative dopamine-containing
neurons in vitro: regulation by an apamin-sensitive Ca(2+)-activated K+ conductance. Exp
Brain Res. 1991;86(1): 141-50.

49. Richards CD, Shiroyama T, Kitai ST. Electrophysiological and immunocytochemical
characterization of GABA and dopamine neurons in the substantia nigra of the rat.
Neuroscience. 1997;80(2): 545-557.

50. Crawford LK, Craige CP, Beck SG. Increased intrinsic excitability of lateral wing serotonin
neurons of the dorsal raphe: a mechanism for selective activation in stress circuits. J
Neurophysiol. 2010;103(5): 2652-2663.

51. Challis C, Boulden J, Veerakumar A, Challis C, Boulden J, Veerakumar A, Espallergues
J, Vassoler FM, Pierce RC, Beck SG, Berton O. Raphe GABAergic neurons mediate the
acquisition of avoidance after social defeat. J Neurosci. 2013;33(35): 13978–13988.

52. Floresco SB, West AR, Ash B, Moore H, Grace AA. Afferent modulation of dopamine
neuron firing differentially regulates tonic and phasic dopamine transmission. Nat
Neurosci. 2003;6(9): 968-973.

53. Tuckwell HC, Penington NJ. Computational modeling of spike generation in serotonergic
neurons of the dorsal raphe nucleus. Prog Neurobiol. 2014;118: 59–101.

54. Ritter J, Lewis L, Mant T, Ferro A. A textbook of clinical pharmacology and therapeutics.
CRC Press; 2008.

55. Benoit-Marand M, Borrelli E, Gonon F. Inhibition of dopamine release via presynaptic D2
receptors: time course and functional characteristics in vivo. J Neurosci. 2001;21(23):
9134–9141.

56. Courtney NA, Mamaligas AA, Ford CP. Species differences in somatodendritic dopamine
transmission determine D2-autoreceptor-mediated inhibition of ventral tegmental area
neuron firing. J Neurosci. 2012;32(39): 13520-13528.

57. Cullen M, Wong-Lin K. Integrated dopaminergic neuronal model with reduced intracellular
processes and inhibitory autoreceptors. IET Syst Biol. 2015;9(6): 245–258.

58. Wang D-H, Wong-Lin K. Comodulation of dopamine and serotonin on prefrontal cortical
rhythms: a theoretical study. Front Integr Neurosci. 2013;7:54. doi:
10.3389/fnint.2013.00054.

59. Haj-Dahmane S. D2-like dopamine receptor activation excites rat dorsal raphe 5-HT
neurons in vitro. Eur J Neurosci. 2001;14(1): 125-134.

60. Aman TK, Shen R-Y, Haj-Dahmane S. D2-like dopamine receptors depolarize dorsal
raphe serotonin neurons through the activation of nonselective cationic conductance. J
Pharmacol Exp Ther. 2007;320(1): 376-385.

61. Bunin MA, Prioleau C, Mailman RB, Wightman RM. Release and uptake rates of 5-
hydroxytryptamine in the dorsal raphe and substantia nigra reticulata of the rat brain. J
Neurochem. 1998;70(3): 1077–1087.

62. Hashemi P, Dankoski EC, Wood KM, Ambrose RE, Wightman RM. 5-HT and histamine

release in the rat substantia nigra pars reticulata following medial forebrain bundle

stimulation. J Neurochem. 2011;118(5): 749-759.

63. Joshi A, Wong-Lin K, McGinnity TM, Prasad G. A mathematical model to explore the

interdependence between the serotonin and orexin/hypocretin systems. In: Conf Proc

IEEE Eng Med Biol Soc. 2011;2011: 7270–7273.

64. Flower G, Wong-Lin K. Reduced computational models of serotonin synthesis, release,

and reuptake. IEEE Trans Biomed Eng. 2014;61(4): 1054-1061.

65. May LJ, Kuhr WG, Wightman RM. Differentiation of dopamine overflow and uptake

processes in the extracellular fluid of the rat caudate nucleus with fast-scan in vivo

voltammetry. J Neurochem. 1988;51(4): 1060-1069.

66. Ludlow KH, Bradley KD, Allison DW, Taylor SR, Yorgason JT, Hansen DM, Walton CH,

Sudweeks SN, Steffensen SC. Acute and chronic ethanol modulate dopamine D2-subtype

receptor responses in ventral tegmental area GABA neurons. Alcohol Clin Exp Res.

2009;33(5): 804-811.

67. Ford CP. The role of D2-autoreceptors in regulating dopamine neuron activity and

transmission. Neuroscience. 2014;282: 13-22.

68. Adell A, Artigas F. The somatodendritic release of dopamine in the ventral tegmental area

and its regulation by afferent transmitter systems. Neurosci Biobehav Rev. 2004;28(4):

415-31.

69. Strogatz SH. Nonlinear dynamics and chaos: with applications to physics, biology,

chemistry, and engineering. CRC Press; 2018.