Whole genome comparisons reveal panmixia among fall armyworm (*Spodoptera*

***frugiperda*) from diverse locations**

Katrina A. Schlum[1], Kurt Lamour[2], Caroline Placidi de Bortoli[2], Rahul Banerjee[2], Scott J.

Emrich[1,3], Robert Meagher[4], Eliseu Pereira[5], Maria Gabriela Murua[6], Gregory A. Sword[7], Ashley

E. Tessnow[7], Diego Viteri Dillon[8], Angela M. Linares Ramirez[9], Komivi S. Akutse[10], Rebecca

Schmidt-Jeffris[11], Fangneng Huang[12], Dominic Reisig[13], and Juan Luis Jurat-Fuentes[2]

[1]Genome Science and Technology Graduate Program, University of Tennessee, Knoxville, TN

37996, USA

[2]Department of Entomology and Plant Pathology, University of Tennessee, Knoxville, TN

37996, USA

[3]Department of Electrical Engineering and Computer Science, University of Tennessee,

Knoxville, TN, USA

[4]USDA-ARS Center for Medical, Agricultural and Veterinary Entomology (CMAVE), Insect

Behavior and Biocontrol Research Unit, Gainesville, FL 32608, USA

[5]Departamento de Entomologia, Universidade Federal de Viçosa, Viçosa, MG 36570, Brazil

[6]Estación Experimental Agroindustrial Obispo Colombres, Las Talitas, Tucumán T4101XAC,

Argentina

[7]Department of Entomology, Texas A&M University, College Station, TX 77843, USA

[8]Isabel Research Substation, Department of Agro-Environmental Sciences, University of Puerto

Rico, Isabela, PR 00662, USA

[9]Lajas Research Substation, Department of Agro-Environmental Sciences, University of Puerto

Rico, Lajas, PR 00667, USA

[10]International Centre of Insect Physiology and Ecology, Nairobi, Kenya

[11]USDA-ARS Temperate Fruit & Vegetable Research Unit, Wapato, WA 98951, USA

[12]Department of Entomology, Louisiana State University Agricultural Center, Baton Rouge, LA

70803, USA

[13]Department of Entomology and Plant Pathology, North Carolina State University, Raleigh, NC

27695, USA

Abstract

The fall armyworm (*Spodoptera frugiperda* (J.E. Smith)) is a highly polyphagous

agricultural pest with long-distance migratory behavior threatening food security worldwide.

This pest has a host range of >80 plant species, but two host strains are recognized based on their

association with corn (C-strain) or rice and smaller grasses (R-strain). In this study, the

population structure and genetic diversity in 55 *S. frugiperda* samples from Argentina, Brazil,

Kenya, Puerto Rico and the United States (USA) were surveyed to further our understanding of

whole genome nuclear diversity. Comparisons at the genomic level suggest panmixia in this

population, other than a minor reduction in gene flow between the two overwintering populations

in the continental USA that also corresponded to genetically distinct host strains. Two maternal

lines were detected from analysis of mitochondrial genomes. We found members from the

Eastern Hemisphere interspersed within both continental USA overwintering subpopulations,

suggesting multiple individuals were likely introduced to Africa. Comparisons between

laboratory-reared and field collected *S. frugiperda* support similar genomic diversity, validating

the experimental use of laboratory strains. Our research is the largest diverse collection of United

States *S. frugiperda* whole genome sequences characterized to date, covering eight continental

states and a USA territory (Puerto Rico). The genomic resources presented provide foundational

information to understand gene flow at the whole genome level among *S. frugiperda*

populations.

Introduction

Larvae of the fall armyworm, *Spodoptera frugiperda* (J.E. Smith, 1797) (Lepidoptera:

Noctuidae), are highly polyphagous agricultural pest affecting key food and fiber staples such as

corn (*Zea mays* L.), cotton (*Gossypium spp.* L.), sorghum (*Sorghum bicolor* L.), rice (*Oryza*

sativa L.), and vegetable crops [1]. Damage by *S. frugiperda* in its native subtropical range in the

Americas resulted in 15-100% yield loss, depending on the level of infestation [2]. In most of the

continental USA this pest does not diapause or survive cold winters. Overwintering populations

in southern Texas and Florida migrate yearly northward over several generations to populate

northern regions in USA and Canada [3].

For more than a decade, effective control of *S. frugiperda* in the Western hemisphere has

been provided mainly by genetically modified corn and cotton producing insecticidal Cry and

Vip3A proteins from the bacterium *Bacillus thuringiensis* (*Bt*). However, resistance to Cry1F,

Cry1Ab and Cry1A.105 insecticidal proteins quickly developed in Puerto Rico and the

continental USA (Florida and North Carolina) [4, 5], as well as Brazil [6] and Argentina [7].

More recently, the economic importance of this pest has further increased with its introduction in

sub-Saharan Africa [8], subsequent spread to India and Southeastern Asia [9], and more recently

Australia [10]. An estimate using data from twelve African countries indicates that yield losses

resulting from *S. frugiperda* injury could be 21-53% of their annual corn production, which

equals to a loss of US$2.5-$6.2 billion in losses [11]. While yet to be detected in molecular

screens [12], resistance alleles to *Bt* toxins could have been carried by invasive *S. frugiperda* to

the Eastern hemisphere.

Populations of *S. frugiperda* are composed of sympatric mixtures of two genetically

differentiated strains based on host preference, a "rice" (R) strain feeding on rice, millet and

smaller grasses, and a "corn" (C) strain feeding preferentially on corn and sorghum [13, 14].

There is evidence supporting that this differentiation involves reproductive incompatibility [15]

and differential susceptibility to xenobiotics [16]. Signatures of positive selection for genes

involved in chemoreception, detoxification, and digestion were also detected in whole-genome

comparisons of nine C- and R-strain individuals [17]. Being morphologically indistinguishable,

C and R individuals are discriminated using genetic markers located on the mitochondrial *COI*

(cytochrome oxidase subunit I) and sex-linked (Z chromosome) *Tpi* (*triosephosphate isomerase*)

genes [18-21]. However, host preference is not absolute and discrepancies among haplotype

markers have been reported. For example, the *Tpi* marker agrees with assortative mating and host

assessments [22] in describing *S. frugiperda* samples from corn in the Eastern Hemisphere as C-

strain, yet a predominant *COI*-R strain marker is obtained in these collections [9]. These

conflicting results may reflect interstrain hybridization [22] or be driven by maternally inherited

symbionts that skew the distribution and diversity of certain haplotypes [23]. Additional factors

influencing these conflicting haplotyping results may include incomplete lineage sorting and

plant host behavioral plasticity.

Nucleotides found at *COI* host strain marker sites have been commonly grouped into

"haplotypes" with relative proportions associated to geographic origin [24], enabling

differentiation of "Texas" and "Florida" overwintering *S. frugiperda* populations in the USA.

Migratory studies with these haplotypes determined that the "Texas" population is distributed

throughout South, Central and North America [25, 26], while the "Florida" population locates to

Florida and the Caribbean but migrates through the eastern USA seaboard to reach Canada [27,

28]. Haplotype profiling of *S. frugiperda* from Africa, India and Southeastern Asia supports

introductions in Africa of individuals from the "Florida" population and subsequent spread to

Asia [9, 12].

In this work, we collected and sequenced 55 genomic DNA (gDNA) samples of *S.*

frugiperda from three continents, with an emphasis on C-strain individuals from North and South

America. Genome-wide comparisons allowed testing for gene flow between geographically

distant *S. frugiperda* populations and comparing genetic diversity between field and laboratory-

reared *S. frugiperda.* We find that based on F_{st} values there is low genetic differentiation between

Texas vs. Florida and that samples from these locations display similar genetic nucleotide

diversity. Host strain differentiation was supported by nuclear and mitochondrial genome

differences. Comparison of mitochondrial genomes detected two clusters where the clusters

correlate by and large with host strain and resistance to the Cry1F toxin. The exceptions to this

correlation suggest that C-strain individuals based on nuclear genome have likely mated with R-

strain individuals. We also observed no detectable reduction in genetic diversity in well-

established *S. frugiperda* laboratory-reared colonies, which supports using laboratory strains of

this pest as a model for field populations.

Methods

Samples and strain typing

Details of the 55 laboratory and field-collected *S. frugiperda* samples analyzed are found

in Supplementary Table 1. Adults (moths) were captured using sex pheromone baited traps [29],

with most collection sites near corn plantings to optimize trap capture efficiency of C-strain

males. Laboratory-reared samples were obtained from rearing facilities. Susceptibility to Cry1F

toxin from *B. thuringiensis* for these laboratory samples was tested in bioassays presented

elsewhere [5, 30-32]. Larval samples were collected at field and laboratory rearing locations and

used when available. The collected specimens were identified as *S. frugiperda* by morphology

features [33] and stored at -20°C until required for analysis.

We followed a sample naming protocol that included the first three letters representing

the country of origin (Bra for Brazil, Arg for Argentina, USA for United States of America, Ken

for Kenya, and Pue for Puerto Rico), the next two letters representing the first two letters of the

state/province of origin (SP for Sao Paulo, TX for Texas, etc. XX when unknown), and the third

letter representing the Cry1Fa susceptibility phenotype when known ("r" for resistant, "s" for

susceptible and "u" for unknown). A number was used when necessary to differentiate samples

with the same geographic origin and Cry1F susceptibility phenotype.

Host strain (C versus R) was determined based on sequence identity at specific marker

positions in reference mt*CoI*1164 and *Tpi*183 sequences, as described elsewhere [12]. The

mitochondrial *COI1287* marker identified all samples as C-strain, so the nuclear *Tpi*183 marker

was used as a more reliable marker of host strain [34]. The mitochondrial *COI*1164 marker was

not used because it did not fall under any of the pre-defined *COI* haplotypes [34] for any of the

samples.

DNA extraction, library preparation, and sequencing

Genomic DNA was isolated from individual carcasses of fifth instar *S. frugiperda* larvae

after dissecting the gut tissue or from legs or heads of adults using the Pure Link Genomic DNA

mini kit (Invitrogen), following manufacturer's protocols, and then quantified using a Nanodrop

spectrophotometer (Thermo Scientific). Extracted DNA was then sheared randomly to between

250-500 bp using a Covaris M220 focused ultrasonicator (Woburn, MA), according to

manufacturer's instructions. The fragmented DNA was then ligated with dual-indices using a

KAPA Hyper prep PCR-free library kit (Roche) according to the manufacturer's directions. The

ligated fragments were quantified using quantitative PCR and a KAPA Library quantification kit

(Roche), and then submitted for sequencing on an Illumina HiSeqX device running a 2 x 150 bp

paired-end configuration (Admera Health, NJ). Raw paired-end sequence reads for each sample

(55) are available for download in the NCBI Sequence Read Archive (SRA) under

SRR12044614-SRR12044668 with associated metadata available under NCBI BioProject id

PRJNA640063. The raw data was processed to remove low quality reads using the CLC

Genomics Workbench v9.5.2 (Qiagen) trim function with default parameters.

Filtering and mapping

Raw reads were quality-trimmed at both ends, filtered for adapter sequences and error

corrected for known Illumina artifacts and PhiX sequencing control using BBDuk

(https://jgi.doe.gov/data-and-tools/bbtools/) with options trimq = 15 and filter = 23. Further

quality control and confirmation that adapter and non-relevant (contaminants, primer artifacts)

sequences were removed from filtered reads were performed using FastQC [35]. Given the

prevalence of C-strain among our samples and similar number of variants called when using both

S. frugiperda corn and rice genomes [36], we mapped the remaining reads to the *S. frugiperda*

corn reference genome (v3.1) downloaded from

https://bipaa.genouest.org/sp/spodoptera_frugiperda_pub/ (assembled length of 312 Mb across

29,949 scaffolds [17]) using a Burrows-Wheeler aligner algorithm (bwa-mem) [37]. Variants

were then called using the SAMtools mpileup utility [38], resulting in detection of 126,977,977

single nucleotide polymorphisms (SNPs) and indels. The unfiltered dataset contained

120,398,863 SNPs. Three samples (USATXu1, BraSPr2 and BraSPr1) were removed from SNP

analysis because they had missing variant call rate greater than 50%, resulting in 52 samples

being used for SNP PCA-based analyses. For nucleotide diversity analyses, SNPs were filtered

using VCFtools [39] based on both minor allele frequency being less than or equal to 0.05 and

presence in at least 50% of the 52 samples surveyed. This filtering further reduced SNP number

to 204,784.

Phylogenomics

The Mash distance method was used to measure pairwise dissimilarity of genomic

sequences [40]. One key advantage of sketching-based approaches is that they neither require *de*

novo genome assembly nor a pre-existing reference genome to identify related individuals. The

Mash method estimates sequence similarity via the Jaccard similarity coefficient (the ratio of

shared k-mers to total k-mers) for a "random" subset of k-mer pairs for each genome surveyed.

The Jaccard similarity (J) is then used to calculate dissimilarity between two genomes (*D*) as D =

1-J. All *S. frugiperda* individuals were sketched with Mash [40] using a 10,000 sketch size, k-

mer size of 21 and minimum copy of each k-mer equal to 2, per previous recommendations of

representing genomes with minimal computational costs [40]. All sample sketches were screened

against the Mash Refseq database and other *Spodoptera* species (*Spodoptera exigua*,

GCA_011316535.1; and *Spodoptera litura*, GCA_002706865.2) as outgroups.

SNP diversity

Weighted F_{st} values as indicators of total genetic variance in a subpopulation were

calculated using VCFtools [39] by the Weir and Cockerham estimator on 1 kbp, 3 kbp and 5 kbp

windows. The variation in window size was used to confirm that there was little to no change in

F_{st} when the window was varied. For F_{st} calculations using the mitochondrial genome, each base

pair position was compared. Nuclear genome (pi) values were calculated using 1 kbp windows

when comparing corn vs. rice host strain, and 3 kbp when comparing F_{st} values for overwintering

FL/TX vs. USA laboratory samples. We also used 3 kbp windows for F_{st} comparisons across the

nuclear genome with populations defined by mitochondria haplotype network groupings, Mash

genome defined "haplotypes", and host strain haplotypes. For all nucleotide diversity analyses, a

total of 3,558,854 SNPs were used to assess diversity using VCFtools [39].

Mitochondrial genome assembly

Mitochondrial sequences were extracted from the whole genome sequence reads using

NOVOplasty [41] with the mitochondrial partial *COI* sequence MH932092.1 as the seed. A

continuous complete mitochondrial chromosome of around 15 kbp was generated for all 55

samples, except for BraBAr5, which was excluded from further analysis. The whole

mitochondrial FASTA sequences were assembled for each sample and then aligned via bwa-

mem (https://github.com/lh3/bwa) with default options. Mitochondrial variants were called using

SAMtools mpileup based on the *S. frugiperda* corn reference genome at LepidoDB

(https://bipaa.genouest.org/sp/spodoptera_frugiperda_pub/). The vcf files were compressed using

SAMtools bgzip, indexed using SAMtools tabix and merged using SAMtools BCFtools merge

[38]. Missing genotypes were replaced with sequence at the mitochondrial reference genome

from the *S. frugiperda* corn genome at LepidoDB. The vcf files were then filtered by removing

16 multi-allelic sites and 92 uninformative sites due to being found in 2 samples or less via the

informloci function in the adegenet R package [42], resulting in 298 sites remaining. These

remaining mitochondrial sites were used to build a haplotype network using the R package pegas

haploNet function [43] with Euclidean distance and an infinite site model used for building the

network.

Structure

PLINK –make-bed [44] was used to generate the input file to FastStructure [45], a

variation of Structure made for larger SNP datasets. FastStructure is a generative model-based

approach based on Hardy-Weinberg equilibrium assumptions between alleles and linkage

disequilibrium between genotyped loci. The FastStructure script structure.py was used to

determine k, the number of assumed populations or genetic groups that share a subset of allele

frequencies from k = 2 to 10. Then k =2 was chosen by the FastStructure's choosek.py script for

model complexity that maximizes marginal likelihood, and k = 1 for model components to

explain structure. Additionally, PCA eigenvalues and eigenvectors were generated using PLINK

2.0 [44, 46] and visualized in R using ggplot2 [47].

Results

Host strain typing

The majority (26, 47.3%) of the 55 *S. frugiperda* samples used in this study originated

from field collections representing eight states in the USA. The average mapping rate for all

samples was 88.2% with a range of between 60.62% to 96.82% range per SAMtools flagstat

[38]. Identification of the C- and R- host strains for each individual was performed using nuclear

Tpi and mitochondrial *COI* genetic markers described elsewhere [26]. Given that sample

collections were from or around cornfields, we expected to observe a majority of C-strain

samples. The *COI*1164 marker identified all samples as C-strain. In contrast, typing based on the

nucleotides at the *TPi*183 site resulted in 42 samples identified as C-strain individuals (C/C), 8 as

R-strain individuals (T/T), 3 as interstrain hybrids (C/T) and 2 that were undetermined due to

low sequencing coverage at this locus. All eight identified R-strain samples were from the USA

(3 from Texas, 2 from Maryland and Tennessee each, and 1 from Puerto Rico).

Whole genome diversity, nucleotide diversity and differentiation

The FastStructure tool was used first to determine possible population structure among

the samples. The number of groups (k) was analyzed for k = 1 to k = 10, but all samples

clustered as one group even when the number of groups was increased. This lack of clear

population structure was further confirmed using principal component analysis (PCA) on a

filtered set of 204,784 nuclear SNPs (see Methods). PC1 through PC10 was plotted and no

structure was found based on country of origin, host strain or susceptibility to the Cry1F

insecticidal protein from *Bt* (data not shown for PC2 to PC10). We noted that PCA using the first

two principal components uncovered weak substructure within the USA and Brazil samples,

where the majority of samples clustered in a larger cluster and four samples (USAFLu9 from

Florida, and BraMGr2, BraBAr2, and BraMGr5 from Minas Gerais in Brazil) grouped together in a

smaller cluster with 1 Florida sample (USAFLr2) the most differentiable from the two clusters

(Fig. 1). All these small cluster samples except USAFLu9 are laboratory samples, and therefore

this grouping may be a result of the side effect of genetic drift associated with the colony

formation process.

Given the overall lack of clear population structure observed based on origin, we next

applied genome sketching [40] and traditional distance-based hierarchical clustering as a third

strategy to analyze these samples. The hierarchical cluster dendrogram defined by the genome-

wide distance estimates based on the fraction of shared k-mers between each pair of genomes

was cut at height 0.05 as it resulted in at least two primary clusters (Texas vs. non-Texas) among

the samples analyzed (Fig. 2). This cut height generated 8 clusters, 6 containing less than 3

samples and were merged to the major cluster, while the remaining 10 samples were left in the

remaining 2 minor cluster. The first cluster was comprised of samples from the USA

(USATXu2-5 from Texas, USATNu1-2 from Tennessee, USAMDu1-2 from Maryland,

USAFlu2 from Florida and PueSlu1 from Puerto Rico), while the second cluster included

individuals from all countries sampled with the majority of samples from USA representing

Florida/eastern seaboard. The three samples from Puerto Rico were interspersed across both

clusters and always clustered with Florida samples, yet genomic distances between clusters were

small (< 0.05 Mash distance). Similar to the PCA clusters, these two clusters did not correlate

with host strain designation or geographic origin, with the first cluster including both R- and C-

strain samples and the second cluster containing both R- and C-strain samples as well as three

interstrain hybrids (Supplemental Fig. S2).

Mash distance is a proxy for genome pairwise distance, with 0 representing identical

genomes (all k-mers that comprise the sketches are present in both sequences). The Mash

distances across all countries surveyed were averaged to be <0.05 (Table S2, range of average

distance was 0.043 to 0.045), probably reflecting recent population expansion. Moreover, these

Mash distances support that R and C-strains are members of the same species, as the estimated

genetic variation is comparable to at least 95% average nucleotide identity (ANI), which is the

ANI cutoff for eukaryotic species [40].

Gene flow between overwintering populations in the USA

Previous work based on the ratios of four pre-determined *COI* haplotypes suggested that

the overwintering Florida *S. frugiperda* population is mostly reproductively isolated from the

overwintering population in Texas, with genetic exchange mostly occurring at the north and

south ends of the Appalachian mountain chains [27]. As mentioned above, we observed that C-

strain samples from Texas and Florida/eastern seaboard did not separate based on hierarchical

clustering of genome-wide distances, and Florida samples were found across both groups (Fig.

2). Further, when comparing the two R-strain samples from Texas and the fourteen C-strain

samples from Florida/eastern seaboard, we detected a low pairwise weighted F_{st} value (0.0111)

based on 3,558,854 nuclear sites. This very low F_{st} value suggests no population separation at the

nuclear level, supporting recent population separation or ongoing gene flow between Texas and

Florida overwintering populations, independently of host strain.

Based on the same Texas R-strain and Florida C-strain groupings, mean diversity was

determined. Mean diversity for the field samples from Texas R-strain and Florida C-strain

samples was estimated at 0.0631 and 0.0556, respectively (Fig. 3). Next, we determined the F_{st}

value based on 406 mitochondrial SNPs found across all samples in these groups. The

mitochondrial F_{st} for Texas R-strain and Florida C-strain samples was estimated at 0.7961. This

F_{st} estimate being close to 1 suggests genetic differentiation between the R and C strains is the

strongest at the mitochondrial level, possibly due to two main ancestral lineages. Although we

see a stronger difference between geographic regions than between R and C strains in this study,

the estimated mitochondrial to nuclear F_{st} ratio of 46.8 is consistent with previously reported sex-

based differences in dispersal [48] and/or mating preference differences between strains [49].

Even so, our data suggest hybridization occurs and some C-strain nuclear genomes have a clearly

R-strain associated mitochondrial genome (see Fig. 5).

Laboratory vs. field diversity

Laboratory-reared strains of *S. frugiperda* are commonly used for diverse physiological,

toxicological and genetic studies, yet their genetic resemblance to field populations has not been

investigated. To investigate the differentiation level between two well established laboratory

strains (>20 years) originating from the same locations in the USA (Mississippi, MS) and all

field USA populations surveyed, we estimated a weighted F_{st} (fixation index). As shown in Table

1, there was moderate differentiation between the MS laboratory strains and field USA field

populations, with an F_{st} range of 0.0056 to 0.1109. Moderate differentiation was also observed

when comparing the MS laboratory strains with other more recently established (<10 years)

laboratory populations from North Carolina (NC) and Florida (FL), with an F_{st} range of 0.0024 to

0.0874. The samples least differentiated ($F_{st} = 0.0631$) from the MS laboratory populations were

collected from TX/MN locations, while the most distinct ($F_{st} = 0.1263$) were the samples from

Maryland (MD). Overall, these moderate F_{st} values [50] suggest small differences at the allelic

level between laboratory and field-collected groups. Genetic diversity in the two laboratory MS

colonies was slightly different but comparable to diversity in field USA populations (F_{st} = 0.0981

on average) and to either of the two overwintering populations (F_{st} = 0.0594).

Additionally, we surveyed overall genomic diversity using 3 kbp windows and estimated

mean diversity for the same two MS laboratory strains compared to all field and laboratory *S.*

frugiperda samples surveyed. Mean nuclear diversity was 0.0011 for Benzon (USAMSs1) and

SIMRU (USAMSs2) laboratory strains (n=2), compared to 0.0015 for field Texas samples (n=7),

and 0.0015 for field Florida/eastern USA samples (n=14).

Diversity across the mitochondrial genome

We investigated the maternal lineage among all but one of the *S. frugiperda* samples (due

to lack of a complete mitochondrial genome in sample BraBAr4) by comparing genetic diversity

of whole mitochondrial (mt) genomes. Using a total of 390 SNPs, we generated a haplotype

network as described in Materials and Methods (Fig. 4). Consistent with a large amount of

standing variation in the species, even the mitochondrial sequences appeared unique with all but

eight samples forming a unique haplotype. These exceptions included three haplogroups of only

Brazil samples, a second group of Kenya and USA samples, and a third haplogroup of samples

from Kenya and two haplogroups of Florida, USA and Puerto Rico samples (Fig. 4). There were

two main "haplotype" clusters that separated in the network, with average genetic difference

between the two clusters of 29.53, supporting evidence for two main maternal lineages within the

samples surveyed. Using these two haplotype clusters as labels, we mapped the mt haplotype

clusters to the two whole genome sequence (WGS) "haplotype" clusters found in Mash distance-

based tree against a mitochondrial variant UPGMA distance-based tree (Fig. 5). In accordance

with the previously estimated F_{st}, the mt and WGS clusters did not correlate completely.

However, all samples in one of the mt haplotypes clustered in one of the WGS haplotypes ("one"

in purple in Fig. 5), while the other mt haplotype contained a mix of WGS haplotypes.

Interestingly, and while not all samples were phenotyped, all samples classified as resistant to the

Cry1F toxin clustered in one of the mt haplotype groups.

Discussion

We report on the genome-level comparison of 55 *S. frugiperda* samples from diverse

locations, mostly from the native range in the Americas but also including samples from an

African location (Kenya) where the insect is a devastating exotic pest. While using the C-strain

S. frugiperda genome for alignments could have underestimated diversity in R-strain individuals,

this approach is supported by previous surveys using both corn and rice genomes with minimal

differences found based on the number of variants called and F_{st} values [36].

The population structure of the 55 genomic *S. frugiperda* samples was surveyed using

parametric and non-parametric methods. Based on FastStructure (parametric), no population

structure was found either within samples from the Eastern Hemisphere or between the Eastern

and Western Hemisphere samples. Since methods based on parametric estimation of allele

frequencies methods are sensitive to sample size, one possibility is that the lack of structure

detected is a reflection of the relatively small number of R-strain samples (only 8 individuals and

potentially 2 hybrids) analyzed. A more even distribution would address potential limited

sampling of some locations surveyed, especially samples representing the Eastern Hemisphere

(e.g. Africa).

Although PCA based on the nuclear genome agreed with the little to no population

structure as found by FastStructure, the Mash distance-based analysis on the nuclear genome and

SNP Hamming-based distance method based on the mitochondrial genome detected two

subpopulations within the samples surveyed, which associate with the previously speculated two

subpopulations in the continental U.S. based on four mitochondrial *COI* haplotypes [51]. The

two clusters clearly indicate that the corn and rice strain individuals are more similar locally than

between geographic areas, suggesting some level of hybridization between host strains in the

USA. Further, we have analyzed individuals that appear to have a C-strain nuclear genome with

an R-strain associated mitochondria (maternal lineage). This is consistent with the conflicting

host strain assignments using specific genetic markers between R-strain and interstrain hybrids

that was previously reported [22]. We still observe host differentiation at the nuclear genome

level, though, and may suggest current single locus markers are not sufficient for surveying all *S.*

frugiperda.

Our analysis included samples collected from locations (Maryland & Tennessee) close to

predicted hybridization zones for TX and FL overwintering *S. frugiperda* populations [27], and

as expected these associated with both geographic regions. There were also three exceptions to

the geographic separation: two Texas samples were found in the predominately Florida

represented group and one Florida sample was found in the predominately Texas population.

This observation suggests that the reproductive isolation between Texas and Florida/Eastern

seaboard is not absolute and mating may be occurring in overlapping locations, as previously

suggested [27]. Future work should include increased sampling from potential hybrid zones

(Georgia/New York) and Texas. The Mash distance-based clusters suggested that Puerto Rico

samples are closer to Florida, since they were found in sub-clusters with Florida-based samples.

However, due to low representation and small genomic distances between each cluster, our data

is not able to show conclusive support for migration models suggesting significant exchanges

between *S. frugiperda* from Puerto Rico and Florida [28].

The low amount of genetic variation across the whole genome between the C- and R-

strains is in line with previous reports, [17] and suggest that the genome differentiated after the

ecological differentiation of host strains or that barriers are isolating C- and R-strains (i.e.

prezygotic reproductive isolation or mating preference). In testing this hypothesis, future

research should involve analyses of a larger sample size of C- and R-strain individuals from

Texas and Florida.

Mitochondrial F_{st} estimates of 0.7961 obtained between C- and R- host strains were

highly similar to the F_{st} value (0.938) from a prior analysis of 9 corn and 9 rice sympatric

samples from Mississippi (USA) [17], and the nuclear DNA F_{st} found in our study (0.0173) is

also similar to the previous study's estimate (0.019). The two mitochondrial haplotypes found in

our analysis could be explained by selective mating or some form of reproductive

incompatibility. Although susceptibility to Cry1F was not known for all samples, we observed

that all Cry1F-resistant strains clustered to one of the mitochondrial haplotypes, despite their

geographic origin. While speculative given the comparatively low number of confirmed Cry1F-

resistant samples, this grouping may reflect partitioning of resistance with the haplotype group

only containing C-strain samples. Further work including Cry1F-resistant R-strain samples

would be necessary to test for associations between mitochondrial haplotype and resistance to

Cry1F.

One possibility to explain the mitochondrial haplotypes detected could be that they are

based on reproductive incompatibility from infection by parasites such as *Wolbachia* [52].

However, we only detected short fragments (<100 bp) matching to the RefSeq *Wolbachia*

genomes on NCBI as of May 9, 2020 using BLAST (data not shown). This observation is in

agreement with leg and head tissues being mostly used for isolating genetic material. In testing

for *Wolbachia* infection, we extracted genomic DNA from a limited number of available

abdominal samples and performed sequencing of PCR amplicons using *Wolbachia*-specific

primers (Fig. S3). *Wolbachia* amplicons suggestive of infection were detected in all samples

tested, independently of mitochondrial haplotype, supporting no influence of *Wolbachia*

infection on reproductive isolation of clustered populations. Further, lack of clear

correspondence between genome-based and mitochondria-based haplotypes suggest on-going

hybridization between C- and R-strains, consistent with detection of two potential hybrids among

our 54 samples. Consistent lack of concordance between the nuclear and mitochondrial F_{st} values

suggests that biological mitochondria-nuclear interactions are not maintained, further supporting

that the samples analyzed are part of a panmictic population with little to no population structure.

Previous reports comparing genetic diversity in laboratory-reared insects to wild

populations mostly reported loss of allele number and heterozygosity during adaptation to mass

rearing [53, 54], although in some examples high genetic variability remained [55], even in long-

established lines [56]. We compared diversity in two long-established (>20 years) *S. frugiperda*

laboratory lines. Both laboratory lines, Benzon (USAMSs1) and SIMRU (USAMSs2), originated

from field collections in Mississippi and they were the least differentiated from the TX/MN field

collections. Interestingly, more recently laboratory populations from NC and FL [5] presented

lower F_{st} values compared to field samples. This moderate differentiation between laboratory and

field samples could be indicative of good culture management involving maintenance of large

populations and/or regular introgression of field individuals. One possibility could also be that

some of the field samples compared display low diversity, yet this is unlikely given samples

from Central and South America (for Texas population) and the Caribbean (for Florida

populations) based on current *S. frugiperda* migratory models [25]. However, these observations

of moderate differentiation in laboratory and field populations are in contrast to previous

research on locusts where genetic differentiation was decreased in laboratory colonies compared

to field samples [57]. It is important to consider that despite the lack of overall differentiation

between laboratory and field *S. frugiperda*, differentiation probably occurs during adaptation to

mass rearing, resulting in selection of specific alleles that may not be representative of field

populations. Thus, while our observations suggest laboratory strains contain genetic diversity

similar to field populations, laboratory strains may have specific allele frequencies that may not

be representative of field populations.

Overall, this study provides a diverse characterization of *S. frugiperda* whole and

mitochondrial genome diversity. Findings from the study support lack of clear *S. frugiperda*

population structure among the locations sampled, with only mitochondrial genomes indicating

two different maternal lines mostly separating host strains. The genomic resources generated

allow further exploration of gene flow and how it may affect management of *S. frugiperda* as an

expanding global superpest using laboratory-reared and field-collected individuals.

Acknowledgements

This project was funded by Agriculture and Food Research Initiative Foundational

Program competitive grant 2018-67013-27820 and Hatch Multistate NC-246 from the US

Department of Agriculture National Institute of Food and Agriculture. We would like to thank

Rodney Nagoshi for review and helpful comments.

References

1. Montezano DG, Specht A, Sosa-Gómez D, Roque-Specht V, Sousa-Silva J, Paula-Moraes SV, Peterson J, Hunt T: **Host plants of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the Americas**. *African Entomology* 2018, **26**(2):286-300.
2. Hruska AJ, Gould F: **Fall armyworm (Lepidoptera: Noctuidae) and *Diatraea lineolata* (Lepidoptera: Pyralidae): Impact of larval population level and temporal occurrence on maize yield in Nicaragua**. *Journal of Economic Entomology* 1997, **90**(2):611-622.
3. Westbrook J, Fleischer S, Jairam S, Meagher R, Nagoshi R: **Multigenerational migration of fall armyworm, a pest insect**. *Ecosphere (Washington, DC)* 2019, **10**(11):n/a.
4. Storer NP, Babcock JM, Schlenz M, Meade T, Thompson GD, Bing JW, Huckaba RM: **Discovery and characterization of field resistance to Bt maize: *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in Puerto Rico**. *Journal of Economic Entomology* 2010, **103**(4):1031-1038.
5. Huang F, Qureshi JA, Meagher RL, Jr., Reisig DD, Head GP, Andow DA, Ni X, Kerns D, Buntin GD, Niu Y *et al*: **Cry1F resistance in fall armyworm *Spodoptera frugiperda*: single gene versus pyramided Bt maize**. *PLoS One* 2014, **9**(11):e112958.
6. Farias JR, Andow DA, Horikoshi RJ, Sorgatto RJ, Fresia P, dos Santos AC, Omoto C: **Field-evolved resistance to Cry1F maize by *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in Brazil**. *Crop Protection* 2014, **64**:150-158.
7. Chandrasena DI, Signorini AM, Abratti G, Storer NP, Olaciregui ML, Alves AP, Pilcher CD: **Characterization of field-evolved resistance to *Bacillus thuringiensis*-derived Cry1F delta-endotoxin in *Spodoptera frugiperda* populations from Argentina**. *Pest Management Science* 2018, **74**(3):746-754.
8. Goergen G, Kumar PL, Sankung SB, Togola A, Tamò M: **First report of outbreaks of the fall armyworm *Spodoptera frugiperda* (J E Smith) (Lepidoptera, Noctuidae), a new alien invasive pest in West and Central Africa**. *PLoS ONE* 2016, **11**(10).
9. Nagoshi RN, Htain NN, Boughton D, Zhang L, Xiao Y, Nagoshi BY, Mota-Sanchez D: **Southeastern Asia fall armyworms are closely related to populations in Africa and India, consistent with common origin and recent migration**. *Scientific Reports* 2020, **10**(1).
10. Department of Argiculture and Fisheries. **First mainland detection of fall armyworm.** [https://www.daf.qld.gov.au/news-media/media-centre/biosecurity/news/first-mainland-detection-of-fall-armyworm]
11. Day R, Abrahams P, Bateman M, Beale T, Clottey V, Cock M, Colmenarez Y, Corniani N, Early R, Godwin J *et al*: **Fall armyworm: Impacts and implications for Africa**. *Outlooks on Pest Management* 2017, **28**(5):196-201.
12. Nagoshi RN, Koffi D, Agboka K, Tounou KA, Banerjee R, Jurat-Fuentes JL, Meagher RL: **Comparative molecular analyses of invasive fall armyworm in Togo reveal**

strong similarities to populations from the eastern United States and the Greater
Antilles. *PLoS One* 2017, **12**(7):e0181982.
13. Pashley DP: **Host-associated genetic differentiation in fall armyworm (Lepidoptera:**
Noctuidae): A sibling species complex? *Annals of the Entomological Society of*
America 1986, **79**:898-904.
14. Pashley DP: **Quantitative genetics, development, and physiological adaptation in**
host strains of fall armyworm. *Evolution* 1988, **42**(1):93-102.
15. Pashley DP, Martin JA: **Reproductive incompatibility between host strains of the fall**
armyworm (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America*
1987, **80**:731-733.
16. Hay-Roe MM, Meagher RL, Nagoshi RN: **Effects of cyanogenic plants on fitness in**
two host strains of the fall armyworm (*Spodoptera frugiperda*). *Journal of Chemical*
Ecology 2011, **37**(12):1314-1322.
17. Gouin A, Bretaudeau A, Nam K, Gimenez S, Aury J-M, Duvic B, Hilliou F, Durand N,
Montagné N, Darboux I *et al*: **Two genomes of highly polyphagous lepidopteran pests**
(*Spodoptera frugiperda*, Noctuidae) with different host-plant ranges. *Sci Rep* 2017,
7(1):11816-11816.
18. Levy H, Garcia-Maruniak A, Maruniak J: **Strain identification of *Spodoptera***
***frugiperda* (Lepidoptera: Noctuidae) insects and cell line; PCR-RFLP of cytochrome**
oxidase C subunit 1 gene. *Florida Entomologist* 2002, **85**(1):186.
19. Nagoshi RN, Meagher RL, Adamczyk JJ, Jr., Braman SK, Brandenburg RL, Nuessly G:
New restriction fragment length polymorphisms in the cytochrome oxidase I gene
facilitate host strain identification of fall armyworm (Lepidoptera: Noctuidae)
populations in the southeastern United States. *Journal of Economic Entomology* 2006,
99(3):671-677.
20. Nagoshi RN, Meagher RL: **Using intron sequence comparisons in the triose-**
phosphate isomerase gene to study the divergence of the fall armyworm host strains.
Insect Mol Biol 2016, **25**(3):324-337.
21. Nagoshi RN: **The fall armyworm triose phosphate isomerase (Tpi) gene as a marker**
of strain identity and interstrain mating. *Annals of the Entomological Society of*
America 2010, **103**(2):283-292.
22. Nagoshi RN: **Evidence that a major subpopulation of fall armyworm found in the**
Western Hemisphere is rare or absent in Africa, which may limit the range of crops
at risk of infestation. *PLoS ONE* 2019, **14**(4):e0208966.
23. Cook James M, Delgado Ana M: **Effects of a sex-ratio distorting endosymbiont on**
mtDNA variation in a global insect pest. *BMC evolutionary biology* 2009, **9**(1):49.
24. Nagoshi RN, Silvie P, Meagher RL: **Comparison of haplotype frequencies**
differentiate fall armyworm (Lepidoptera: Noctuidae) corn-strain populations from
Florida and Brazil. *Journal of Economic Entomology* 2007, **100**(3):954-961.
25. Nagoshi RN, Fleischer S, Meagher RL, Hay-Roe M, Khan A, Murua MG, Silvie P,
Vergara C, Westbrook J: **Fall armyworm migration across the Lesser Antilles and the**
potential for genetic exchanges between North and South American populations.
PLoS One 2017, **12**(2):e0171743.
26. Nagoshi RN, Nagoshi BY, Canarte E, Navarrete B, Solorzano R, Garces-Carrera S:
Genetic characterization of fall armyworm (*Spodoptera frugiperda*) in Ecuador and

comparisons with regional populations identify likely migratory relationships. *PLoS*
ONE 2019, **14**(9):e0222332.

27. Nagoshi RN, Meagher RL, Hay-Roe M: **Inferring the annual migration patterns of fall armyworm (Lepidoptera: Noctuidae) in the United States from mitochondrial haplotypes**. *Ecol Evol* 2012, **2**(7):1458-1467.

28. Nagoshi RN, Rosas-Garcia NM, Meagher RL, Fleischer SJ, Westbrook JK, Sappington TW, Hay-Roe M, Thomas JM, Murua GM: **Haplotype profile comparisons between** *Spodoptera frugiperda* **(Lepidoptera: Noctuidae) populations from Mexico with those from Puerto Rico, South America, and the United States and their implications to migratory behavior**. *J Econ Entomol* 2015, **108**(1):135-144.

29. Meagher RL: **Trapping fall armyworm (Lepidoptera: Noctuidae) adults in traps baited with pheromone and a synthetic floral volatile compound**. *Florida Entomologist* 2001, **84**(2):288-292.

30. Banerjee R, Hasler J, Meagher R, Nagoshi R, Hietala L, Huang F, Narva K, Jurat-Fuentes JL: **Mechanism and DNA-based detection of field-evolved resistance to transgenic Bt corn in fall armyworm (*Spodoptera frugiperda*)**. *Scientific Reports* 2017, **7**(1):10877.

31. Oscar FS-A, João VCR, Thadeu CS, Clébson ST, Silverio OC, Raul NCG, Eliseu JGP: **Resistance to dual-gene Bt maize in *Spodoptera frugiperda*: selection, inheritance, and cross-resistance to other transgenic events**. *Scientific Reports* 2015, **5**(1).

32. Santos-Amaya OF, Tavares CS, Rodrigues JVC, Campos SO, Guedes RNC, Alves AP, Pereira EJG: **Fitness costs and stability of Cry1Fa resistance in Brazilian populations of *Spodoptera frugiperda***. *Pest Management Science* 2017, **73**(1):35-43.

33. **Fall armyworm. University of Florida Publication Number EENY-98** [http://entnemdept.ufl.edu/creatures/field/fall_armyworm.htm#host]

34. Nagoshi RN, Silvie P, Meagher RL, Lopez J, Machado V: **Identification and comparison of fall armyworm (Lepidoptera: Noctuidae) host strains in Brazil, Texas, and Florida**. *Annals of the Entomological Society of America* 2007, **100**(3):394-402.

35. **FastQC: A quality control tool for high throughput sequence data** [https://qubeshub.org/resources/fastqc]

36. Nam K, Nhim S, Robin S, Bretaudeau A, Nègre N, d'Alençon E: **Divergent selection causes whole genome differentiation without physical linkage among the targets in *Spodoptera frugiperda* (Noctuidae)**. *bioRxiv* 2018:452870.

37. Li H: **Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM**. 2013.

38. Li H: **A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data**. *Bioinformatics* 2011, **27**(21):2987-2993.

39. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST *et al*: **The variant call format and VCFtools**. *Bioinformatics* 2011, **27**(15):2156-2158.

40. Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM: **Mash: fast genome and metagenome distance estimation using MinHash.(Report)**. *Genome Biology (Online Edition)* 2016, **17**(1).

41. Dierckxsens N, Mardulyn P, Smits G: **NOVOPlasty: de novo assembly of organelle genomes from whole genome data**. *Nucleic acids research* 2017, **45**(4):e18.

42.   Jombart T, Ahmed I: **Adegenet 1.3-1: new tools for the analysis of genome-wide SNP**
data. *Bioinformatics (Oxford, England)* 2011, **27**(21):3070.
43.   Paradis E: **pegas: an R package for population genetics with an integrated–modular**
approach. *Bioinformatics* 2010, **26**(3):419-420.
44.   Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ: **Second-**
generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*
2015, **4**(1).
45.   Pritchard JK, Stephens M, Raj A: **fastSTRUCTURE: Variational inference of**
population structure in large SNP data sets. *Genetics* 2014, **197**(2):573-589.
46.   **PLINK 2.0.** [www.cog-genomics.org/plink/2.0/]
47.   Wickham H: **ggplot2 Elegant Graphics for Data Analysis**. 2016.
48.   Rašić G, Schama R, Powell R, Maciel-de Freitas R, Endersby-Harshman NM, Filipović I,
Sylvestre G, Máspero RC, Hoffmann AA: **Contrasting genetic structure between**
mitochondrial and nuclear markers in the dengue fever mosquito from Rio de
Janeiro: implications for vector control. *Evolutionary Applications* 2015, **8**(9):901-
915.
49.   Finley LWS, Haigis MC: **The coordination of nuclear and mitochondrial**
communication during aging and calorie restriction. *Ageing Research Reviews* 2009,
8(3):173-188.
50.   Stephan W: **Principles of Population Genetics. Fourth Edition.** *The Quarterly review*
of biology 2007, **82**(4):416-416.
51.   Nagoshi RN, Meagher RL, Flanders K, Gore J, Jackson R, Lopez J, Armstrong JS,
Buntin GD, Sansone C, Leonard BR: **Using haplotypes to monitor the migration of fall**
armyworm (Lepidoptera: Noctuidae) corn-strain populations from Texas and
Florida. *Journal of Economic Entomology* 2008, **101**(3):742-749.
52.   Correa CC, Ballard JWO: **Wolbachia associations with insects: winning or losing**
against a master manipulator. *Frontiers in Ecology and Evolution* 2016, **3**.
53.   Zygouridis NE, Argov Y, Nemny-Lavy E, Augustinos AA, Nestel D, Mathiopoulos KD:
Genetic changes during laboratory domestication of an olive fly SIT strain. *Journal*
of applied entomology (1986) 2014, **138**(6):423-432.
54.   Gilchrist AS, Cameron EC, Sved JA, Meats AW: **Genetic consequences of**
domestication and mass rearing of pest fruit fly *Bactrocera tryoni* **(Diptera:**
Tephritidae). *Journal of Economic Entomology* 2012, **105**(3):1051-1056.
55.   Simões P, Pascual M, Coelho MM, Matos M: **Divergent evolution of molecular**
markers during laboratory adaptation in *Drosophila subobscura*. *Genetica* 2010,
138(9-10):999-1009.
56.   Parreño MA, Scannapieco AC, Remis MI, Juri M, Vera MT, Segura DF, Cladera JL,
Lanzavecchia SB: **Dynamics of genetic variability in** *Anastrepha fraterculus* **(Diptera:**
Tephritidae) during adaptation to laboratory rearing conditions. *BMC genetics* 2014,
15 Suppl 2(Suppl 2):S14.
57.   Berthier K, Chapuis MP, Simpson SJ, Ferenz HJ, Habib Kane CM, Kang L, Lange A, Ott
SR, Babah Ebbe MA, Rodenburg KW *et al*: **Laboratory populations as a resource for**
understanding the relationship between genotypes and phenotypes: A global case
study in locusts. *Advances in insect physiology* 2010, **39**:1-37.
633

Figure legends

Figure 1: Principal component analysis (PCA) plot of 52 *Spodoptera frugiperda* samples (3 samples removed due to missing rate ≥ 50%) based on 3,558,854 SNPs. Country of origin and host strain as determined by the nuclear *Tpi* marker are visualized by color and shape, respectively, as shown in the legend.

Figure 2: Dissimilarity heatmap of Mash distances across all 55 *Spodoptera frugiperda* samples, colored by country of origin as indicated. Two outgroups (*Spodoptera litura* and *Spodoptera exigua*) were added. Samples were named as described in Materials and Methods.

Figure 3: Nucleotide diversity box plots between C-strain Florida (FL) vs. R-strain Texas (TX) samples.

Figure 4: Mitochondrial haplotype network of all 54 *S. frugiperda* mitochondrial chromosome level assembled individuals, colored by country of origin.

Figure 5: UPGMA distance-based tree on biallelic variant SNPs in 54 mitochondrial chromosome level assembled individuals rooted with *Spodoptera litura* as an outgroup.

Table 1: Pairwise F_{st} values across all populations surveyed. NA was used when the samples overlap in the defined populations.

2   2.828   4   5.292   29.53

DISTANCE

Cy1F resistance profile
- ■ resistant
- ■ susceptible
- ■ unknonwn/NA

Field vs. Laboratory
- ■ field
- ■ laboratory

WGS haplotype group
- ■ one
- ■ two
- ■ NA

Host strain
- ■ corn
- ■ rice
- ■ hybrid
- ■ unknonwn/NA

Genetic distance
— 0.01

USAFLr1
USAFLr2
ArgBAu1
BraMGr2
BraMGr1
BraBAr3
BraBAr2
BraBAr4
BraBAr1
USASCu2
USAFLu8
USAFLu10
ArgXXr2
USAMNu2
ArgXXr1
BraSPr2
BraSPr1
BraMGr5
BraMGr4
USAFLu6
BraSPr3
PueGUr1
USAFLu3
USAFLu7
USAFLu11
USAFLu4
PueLAu1
USANCr1
USAFLu1
USASCu1
USAFLu12
USATXu6
USAFLu5
USAFLu9
BraMGr3
USAMSs1
USAMSs2
USATXu2
ArgJs1
USAMNu1
USAMDu1
USATXu1
PueSlu1
USATXu3
KenXXu2
KenXXu1
USAMDu2
ArgSFu1
KenXXu3
USATXu4
USATXu5
USATNu1
USATNu2
USAFLu2
Spodoptera litura