Evolution of the canonical sex chromosomes

of the guppy and its relatives

M. Kirkpatrick[*], J.M. Sardell[*], B. J. Pinto[*,†], G. Dixon[*], C. L. Peichel[‡], M. Schartl [§,1]

Addresses:

[*] Dept. of Integrative Biology, University of Texas, Austin TX 78712 USA

[†] Milwaukee Public Museum, Milwaukee, WI 53233 USA

[‡] Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, 3012 Bern

Switzerland

[§] Developmental Biochemistry, University of Würzburg, Biozentrum, Am Hubland, 97074
Würzburg, Germany and The Xiphophorus Genetic Stock Center, Department of
Chemistry and Biochemistry, Texas State University, San Marcos, Texas, TX 78666, USA

Data repositories:

NCBI SRA Bioprojects PRJNA528814 and PRJNA353986

[1] Authors listed in order of their relative contributions.

<u>Running title</u>: Evolution of *Poecilia* sex chromosomes

<u>Key words</u>: Recombination, sex determination, heterochiasmy, X chromosome, Y chromosome

<u>Corresponding author</u>:

Mark Kirkpatrick

Dept. of Integrative Biology, University of Texas, Austin TX 78712

(512) 499-0925

kirkp@mail.utexas.edu

Abstract

The sex chromosomes of the guppy, *Poecilia reticulata*, and its close relatives are of particular

interest: they are much younger than the highly degenerate sex chromosomes of model

systems such as mammals and *Drosophila melanogaster*, and they carry many of the genes

responsible for the males' dramatic coloration. Over the last decade, several studies have

analyzed these sex chromosomes using a variety of approaches including sequencing genomes

and transcriptomes, cytology, and linkage mapping. Conflicting conclusions have emerged, in

particular concerning the history of the sex chromosomes and the evolution of suppressed

recombination between the X and Y. Here we address these controversies by reviewing the

evidence and reanalyzing data. We find no support for a nonrecombining sex determining

region (SDR) or evolutionary strata in *P. reticulata.* Further, we find that the evidence most

strongly support the hypothesis that the SDRs of *P. picta* and *P. wingei* evolved independently

after those lineages diverged. We identify possible causes of conflicting results in previous

studies and suggest best practices going forward.

46   1: INTRODUCTION

The origin and evolution of young sex chromosomes are of particular interest to evolutionary

genomics. They are the most rapidly evolving part of the genome in many animals and plants,

and they have features that give unique insights into the evolution of recombination, sexually

antagonistic selection, and other important processes (BACHTROG *et al.* 2011). The guppy,

Poecilia reticulata, holds a special place in the history of this subject. Because they carry many

of the genes responsible for the males' famed coloration, the *P. reticulata* sex chromosomes

have been studied since the 1920s (WINGE 1922; HASKINS *et al.* 1961; LINDHOLM AND BREDEN 2002;

CHARLESWORTH 2018). The last decade has seen a burst of research on the sex chromosomes of

the guppy and its relatives. Several recent studies have arrived at conflicting conclusions,

notably regarding the evolution of recombination. These controversies are the focus of this

paper.

Studies from the pre-genomic era also provided conflicting conclusions regarding

recombination between the sex chromosomes. Using linkage maps, TRIPATHI *et al.* (2009)

reported that recombination between the X and Y of *P. reticulata* is confined to a relatively

small region bounded on either side by large nonrecombining regions. NANDA *et al*. (2014) used

cytology and linkage maps to study the sex chromosomes of *P. reticulata*, its sister species *P.*

wingei, and the closely related *P. obscura*. They concluded that the Y chromosomes of these

three species descended from a common ancestral Y, and that the X and Y chromosomes of *P.*

reticulata recombine down their lengths (save perhaps for a small heterochromatic region

specific to the Y). Further, they reported that an extended region of heterochromatin at the end

of the Y distal to the centromere has evolved in *P. wingei* that would regionally block

recombination with the X. Conversely, by inferring meiotic crossovers visualized through the

localization of MHL1, LISACHOV *et al.* (2015) found that recombination between the X and Y in *P.*

reticulata is concentrated towards the end of the chromosome distal to the centromere.

The pace of discovery accelerated with the arrival of genome sequences for *P. reticulata*

(FRASER *et al.* 2015; KÜNSTNER *et al.* 2016). By analyzing molecular variation among re-sequenced

genomes from several natural populations of *P. reticulata*, WRIGHT *et al.* (2017) drew

conclusions at odds with the previous reports. They reported that crossing over between the X

and Y is completely blocked over about 40% (10 Mb) of their lengths, with recombining regions

on either side. They claimed that the nonrecombining sex determining region, or SDR, is divided

into two "strata", which are regions in which crossing over between the X and Y was completely

suppressed at different times in the past (LAHN AND PAGE 1999; CHARLESWORTH 2017). WRIGHT *et*

al. (2017) further concluded that the nonrecombining SDR has expanded independently in

three populations that inhabit the headwaters of rainforest streams. MORRIS *et al.* (2018)

followed up this study by identifying 40 loci that are unique to the putative nonrecombining

region of the Y chromosome, and proposed two of them as candidates for the sex determining

gene.

In the next study from the same research team, DAROLTI *et al.* (2019) enlarged the

phylogenetic picture by analyzing genomes and transcriptomes from five species of poeciliid

fish, including *P. reticulata*, its sister species *P. wingei*, and their congener *P. picta*. These

authors found that Chromosome 12 is responsible for sex determination in all three species.

They reported that the two strata that they found on the *P. reticulata* Y are shared with *P.*

wingei, which implies that they evolved in the common ancestor of those species. In the more

distantly related *P. picta*, they found the Y chromosome to be highly degenerate. They also

concluded that the SDRs in all three species descend from a common ancestor, which implies

that the rates at which the Y degenerates varies greatly between lineages. DAROLTI *et al.* (2020)

found from linkage mapping that completely suppressed recombination is confined only to the

first of the two strata in *P. reticulata*, while there is very rare crossing over between the X and Y

in the second "stratum". In the most recent paper from that research group, ALMEIDA *et al.*

(2021) carried out long-read sequencing on much larger samples of individuals from six natural

populations. They again concluded that the l data show that there is no recombination in

Stratum 1.

An independent research team studied the *P. reticulata* sex chromosomes using linkage

mapping (BERGERO *et al.* 2019). They found no evidence for completely suppressed

recombination between the X and Y anywhere on those chromosomes. Recombination

between the X and Y in natural populations is very rare, however, as there is elevated F_{ST}

between males and females down the entire length of Chromosome 12, in some regions

attaining values greater than 0.1. BERGERO *et al.* (2019) showed that the very low levels of

recombination between the X and Y are not unique to the sex chromosomes: males also have

extremely low recombination rates on autosomes except near the telomeres. They questioned

the conclusion of WRIGHT *et al.* (2017) that recombination between the X and Y has been

suppressed. BERGERO *et al.* also identified a region of high recombination on the end of the Y

chromosome distal to the centromere, consistent with the patterns of male recombination on

autosomes and the findings of LISACHOV *et al.* (2015). The localization of crossovers in males to

the tips of all chromosomes is consistent with the high GC content found there (CHARLESWORTH

et al. 2020b). CHARLESWORTH *et al.* (2020a) reported a crossover between the X and Y of *P.*

reticulata very close to the boundaries between Strata 1 and 2 as defined ALMEIDA *et al.* (2021).

Most recently, FRASER *et al.* (2020) assembled a new high quality reference genome for *P.*

reticulata that includes unphased scaffolds from both the X and Y, and they re-sequenced fish

from six natural populations. They concurred with BERGERO *et al.* (2019) that there is no

evidence for a nonrecombining stratum several megabases in size. Further, they found two

candidate regions for the sex determining gene contained in Y-specific regions that

(surprisingly) are located at opposite ends of the Y chromosome. Figure 1 summarizes some of

these results.

These conflicting conclusions have led to controversy and confusion in the scientific

community (BERGERO AND CHARLESWORTH 2019; WRIGHT *et al.* 2019). In an effort to remedy this

situation, in this paper we review, reanalyze, and reinterpret published data from the guppy

and its close relatives. Our goal is not to perform a forensic analysis to determine why different

research teams have come to different conclusions. Rather, we analyze the existing data in

order to independently evaluate conclusions regarding the evolutionary histories of their sex

chromosomes.

We first focus on the evolution of recombination and the origin of the Y chromosomes.

We find no evidence for strata in *P. reticulata* or for any region where crossing over between

the X and Y is completely suppressed. Our results confirm that a non-recombining SDR spans

much of the X and Y in *P. wingei* and *P. picta*, and that the Y in *P. picta* is highly degenerate.

We then consider hypotheses regarding homology of the SDRs in all three species. By

"homology", we mean that the SDRs in these species descended from a common ancestral SDR.

While the shared identity of the Y chromosomes of *P. reticulata* and *P. wingei* has been

established by molecular cytogenetics (NANDA *et al.* 2014), that approach cannot evaluate the

homology of their SDRs. To date, no genetic evidence has been adduced regarding homology of

the Y chromosome or the SDR in *P. picta*. Our analysis of gene trees does not provide support

for the hypothesis that the SDRs of *P. wingei* and *P. picta* descend from a common ancestral

SDR. A second hypothesis is that the Y chromosomes of *P. wingei* and *P. reticulata* originated in

their common ancestor when a new Y chromosome evolved from an X chromosome after

divergence from *P. picta* (BERGERO AND CHARLESWORTH 2019; MEISEL 2020). This would make their

Y chromosomes younger, and so explain why they are so much less degenerate than the Y of *P.*

picta. But again, we find that gene trees give little support to this hypothesis. The most strongly

supported hypothesis is that the SDRs of *P. wingei* and *P. picta* originated independently. We

are unable to determine whether the entire Y chromosome originated once in the common

ancestor of all three species and degenerated at different rates, as suggested by parsimony, or

if the variation in Y degeneration across species reflects different ages of Y chromosomes that

originated independently.

2: MATERIALS AND METHODS

2.1: The data

We downloaded all of the whole genome re-sequencing data available as of June 2020 from the

NCBI SRA under Bioprojects PRJNA528814 and PRJNA353986. These data were acquired by

WRIGHT *et al.* (2017) and DAROLTI *et al.* (2019). (We did not use the more recent data from

FRASER *et al.* (2020)) or ALMEIDA *et al.* (2021) for most of our analyses because they were

published after our paper was first submitted.) The data consist of paired-end Illumina

sequences from 52 samples of whole genome sequences of 5 species: *Gambusia holbrooki*,

Poecilia latipinna, *P. picta*, *P. reticulata*, and *P. wingei*. For a given data type, multiple records

from the same individual were concatenated. Read quality was assessed using FastQC [v0.11.5]

(ANDREWS 2010).

For most analyses, we mapped WGS reads to the *Xiphophorus maculatus* reference

genome (v5.0; GCA_002775205.2; SCHARTL *et al.* (2013)) because it is the highest-quality

poeciliid fish genome and is an equal phylogenetic distance from the *Poecilia* species. For the

analyses of gene trees (see below) we redid our original analyses using the new *P. reticulata*

reference genome (GCA_904066995.1) which was published after our paper was first

submitted. We used this second genome with the goal of increasing read mapping rates across

Poecilia species. Indeed, we found that the mapping efficiency with sequencing reads from *P.*

reticulata is more than twice as high when using the new *P. reticulata* reference (~93%) as

when using the *X. maculatus* reference (~41%), and it was similar across all three focal taxa.

However, there are no differences in read mapping efficiency between males and females using

either of the two reference genomes, which makes *X. maculatus* an appropriate reference for

the other analyses. There are small differences in the analysis pipelines using the *X. maculatus*

and *P. reticulata* reference genomes that are specified below.

There is some uncertainty about the physical coordinates on the *P. reticulata* sex

chromosomes. Different studies have used different reference genomes that vary in important

features (e.g. WRIGHT *et al.* (2017) *vs.* BERGERO *et al.* (2019) *vs.* FRASER *et al.* (2020) *vs.* this study).

Further, there are several reports of inversions and assembly errors on the *P. reticulata* sex

chromosomes (NANDA *et al.* 2014; BERGERO *et al.* 2019; CHARLESWORTH *et al.* 2020a; DAROLTI *et al.*

2020; FRASER *et al.* 2020). Accordingly, we interpret the chromosome coordinates with caution.

<u>Analyses based on the *X. maculatus* genome</u>

We trimmed adapters and low-quality bases from raw SRA sequence reads using cutadapt

[v1.14] (MARTIN 2011), mapped DNA and RNA reads to the *X. maculatus* reference genome

using Bowtie2 [v2.3.4] with default parameters and the –local argument (LANGMEAD AND SALZBERG

2012). We removed PCR duplicates using Picard [v2.21] (BROAD INSTITUTE 2019). We sorted and

subsetted alignment files by chromosome using SAMtools [v1.6] (LI *et al.* 2009) and calculated

fold coverage in 10 kb windows using BEDtools [v2.26.0] (QUINLAN AND HALL 2010). We called

variants with a quality score ≥20 using BCFtools mpileup [v1.10] (LI 2011). We removed indels

and singletons, selected for biallelic SNPs using VCFtools [v0.1.16] (DANECEK *et al.* 2011), and we

required a minimum read depth of 3.

To determine which SNPs may be in an SDR, we considered patterns of heterozygosity. An

allele was considered to be putatively Y-linked if it always appeared in males in heterozygotes

and was absent from females. We designate these alleles as "Y-like", the alternate alleles at

these SNPs as "X-like", and the sites at which these occur as "SDR-like SNPs". We imposed the

additional criterion that these sites have data from at least two individuals of each sex. Because

the sample sizes are small, SDR-like SNPs can occur by chance at autosomal loci. Nevertheless,

this set of sites will be highly enriched for SNPs that truly are sex-linked. As an internal control,

we also applied this analysis to the autosomes to assess how frequently SNPs with putative sex

linkage occur throughout the genome.

To study divergence between the X and Y chromosomes, we computed five statistics in

sliding windows: (1) the ratio of read depth in males and females ("read depth ratio"),

computed using BEDtools [v2.26.0] (QUINLAN AND HALL 2010); (2) F_{ST} between males and females

("F_{ST}"), based on WEIR and COCKERHAM'S (1984) estimator computed using VCFtools [v0.1.16]

(DANECEK *et al.* 2011); (3) the ratio of the density of all SNPs in males and in females ("SNP

density ratio"), computed using VCFtools and a custom R script; (4) the density of SNPs with

patterns of heterozygosity consistent with the SDR ("SDR-like SNP density"), computed using

VCFtools and a custom R script; and (5) the density of SNPs with alleles that are restricted to

females ("female-specific SNP density"), computed using VCFtools and a custom R script. (The

latter are unexpected in species with XY sex determination, but we will see below that they do

appear in one of the species.) Because all downstream analyses are dependent on successfully

mapping reads, we first calculated read depth differences between males and females in highly-

sensitive 10 kb windows. All other sliding-window statistics we calculated in 100 kb windows.

<u>Analyses based on the *P. reticulata* genome</u>

We quality and adapter trimmed the sequencing reads using Trim Galore! [v0.6.6] (ANDREWS

2010; MARTIN 2011), filtered PCR duplicates using bbmap [v38.90] (BUSHNELL 2014), and mapped

reads to the *P. reticulata* reference genome using minimap2 [2.17] (LI 2018). We called variants

one chromosome at a time at sites with quality score ≥20 using BCFtools mpileup [v1.11] (LI

2011). We removed indels and singletons, selected for biallelic SNPs using VCFtools [v0.1.16]

(DANECEK *et al.* 2011), and we required a minimum read depth of 10.

We implemented this strategy using a custom R script. We first split the genotypes by

species using VCFtools. We then filtered SNPs to include only those with no missing individuals

(--max-missing 1.0) and a minimum allele frequency of 0.2 (--maf 0.2). We then split the

genotypes further by sex and calculated allele frequencies (--freq) and observed heterozygosity

(--hardy) in VCFtools.

We used gene trees to test hypotheses regarding the evolution of suppressed

recombination. We applied several filters with a custom R script to identify SNPs used to

construct these trees. We refer to the SNPs that meet all of the following criteria as

"topologically informative." First, we required sequence from at least two *P. wingei* males, two

P. wingei females, two *P. picta* males, two *P. picta* females, one *P. latpinna*, and one *G.*

holbrooki. Second, we only considered SNPs where alleles are fixed within X and Y

chromosomes. That is, we required that the females within a species are all homozygous for

the same allele, and that the males in that species either are all heterozygous or all

homozygous for the same allele found in females. Third, we required that all individuals from

the outgroup species (*P. latpinna* and *G. holbrooki*) possess the same allele, which we identified

as the "ancestral" allele. Finally, we required that the alternate "derived" allele is fixed in either

two or three of the following four chromosomes: the *P. wingei* X, *P. wingei* Y, *P. picta* X, or *P.*

picta Y. We used the same R script to calculate the number of topologically informative SNPs

that exhibit each of the 10 gene trees shown in Figure 5.

3: RESULTS

3.1: Divergence and recombination between the X and Y

To quantify divergence between the X and Y in the three species, we examined several statistics in sliding windows using the syntenic linkage group 8 of *X. maculatus* as the reference. Figure 2 shows the read depth ratio and male:female F_{ST}, while Figure 3 shows the male:female SNP density ratio, SDR-like SNP density, and female-specific SNP density. Figure 4 shows the densities of SDR-like SNPs on the sex chromosomes and autosomes. As anticipated, SDR-like SNPs do occur on autosomes, but are much less common (see Methods).

In *P. reticulata*, there is no sign that the X and Y have diverged very much anywhere along their lengths. We do not see a decreased read depth ratio, which is a telltale signature of Y chromosome degeneration (Vicoso and Bachtrog 2015). The maximum value of the smoothed regression for F_{ST} between males and females is less than 0.003 down the entire length of the sex chromosomes (Figure 2). The maximum value in any 100 kb window is F_{ST} = 0.15. (We obtained similar results for F_{ST} using both the *P. reticulata* and the *X. maculatus* reference genomes.) The other four statistics fall within the ranges typical for autosomes. These results do not provide any evidence of a nonrecombining sex determining region (SDR).

Our results are consistent with the results but not the interpretation of previous analyses based on gene trees (Darolti *et al.* 2020; Almeida *et al.* 2021). Following the origin of a non-recombining stratum on the Y by an inversion (or any other mechanism involving a *cis* recombination modifier), the SDRs on all Y chromosomes that inherit that stratum will form a monophyletic group with respect to the X chromosomes (Lahn and Page 1999; Zhou *et al.* 2014).

The monophyly persists through speciation events: the homologous strata on the Ys from all

the descendant species continue to form a clade with respect to the X chromosomes, and the

monophyly extends across the entire length of the stratum. In recombining regions of the sex

chromosomes, however, this monophyly breaks down. Recombination causes gene copies from

the X and Y chromosomes of each species to be intermingled on a gene tree. Recombination

events more recent than a speciation event cause gene copies from the X and Y of a species to

cluster together rather than with their gametologs in other species. Gene trees thus offer a

sensitive way to distinguish the SDR from the recombining pseudoautosomal region (PAR)

because they integrate signals of recombination that have accumulated in natural populations

over many generations (DIXON *et al.* 2018; SARDELL *et al.* 2021).

DAROLTI *et al.* (2020) estimated the gene trees at 42 loci spread along the length of the sex

chromosomes. Figure 1 shows their locations. Only one of these loci, *alad*, shows a gene tree

topology consistent with a nonrecombining stratum shared between *P. reticulata* and *P. wingei*,

but the statistical support for this tree is weak (bootstrap value: 51%). (Three other gene trees

are also stated to be consistent with that hypothesis, but in fact their topologies are not: see

Figures 2c-e in DAROLTI *et al.* (2020).) At the locus *npr2*, which is less than 3 kb away from *alad*,

there is strong statistical support for the node joining the X and Y in *P. reticulata* together

(bootstrap value: 97%) and for a node joining the X and Y in *P. wingei* (bootstrap value: 96%).

This is evidence in both species that there has been recombination between the X and Y

chromosomes in the region between the *npr2* locus and the sex determining region since the

two species diverged. The remaining 40 gene trees are also inconsistent with a stratum that

predates the divergence of *P. reticulata* and *P. wingei*. Most recently, ALMEIDA *et al.* (2021)

analyzed gene trees with much larger samples. In five of the six populations studied, the

majority of trees in the proposed Stratum 1 and elsewhere on Chromosome 12 have topologies

that are consistent with ongoing recombination between the X and Y in *P. reticulata*.

In the species that is sister to guppies, *P. wingei*, we find quite a different picture. Four of

the five statistics (all but read depth ratio) fall far outside the autosomal range of values over

much of the center part of the sex chromosomes (Figures 2 and 3). These patterns are

consistent with very little or no recombination. The read depth ratio remains near 1, however,

suggesting that there has not been extensive degeneration on the Y chromosome, possibly

because recombination was suppressed recently. These conclusions concur with Darolti *et al.*

(2019).

Within the region of reduced recombination in *P. wingei*, there is a smaller segment

between about 10 Mb and 20 Mb where the female-specific allele density is greatly elevated,

with very sharp boundaries at each end (Figure 3). As noted earlier, this pattern in unexpected

in species with XY sex determination. The other three statistics just mentioned show no unusual

patterns specific to this 10 Mb region. We speculate that these patterns might result from a

polymorphic inversion on the X chromosome of *P. wingei* that is evident in the cytological data

of Nanda *et al.* (2014).

The third species, *P. picta*, shows yet another distinctive set of patterns. Most strikingly,

the read depth ratio is about one half over most of the end of the sex chromosome proximal to

the centromere (Figure 2). This is indicative of large-scale deletions and/or divergence of the Y

sequence to the point that reads from it no longer map to the reference (Vɪᴄᴏsᴏ 2019). The

SDR-like SNP density, female-specific allele density, and F_{ST} are much lower than in *P. wingei*,

presumably for the same reasons. These conclusions agree with DAROLTI *et al.* (2019).

3.2: Evolutionary relations between the SDRs

We used gene trees to investigate the evolutionary relations between the SDRs on the Y

chromosomes of *P. wingei* and *P. picta*. One hypothesis is that the oldest parts of their SDRs

are homologous, meaning that they descend from an SDR in their common ancestor. This

possibility is favored by DAROLTI *et al.* (2019). A second hypothesis is that the Y chromosome of

the shared ancestor of *P. wingei* and *P. reticulata* was derived from its X, an idea favored by

BERGERO AND CHARLESWORTH (2019) and MEISEL (2020). A third hypothesis is that the SDRs of *P.*

wingei and *P. picta* originated independently, either due to the independent evolution of

suppressed recombination of a largely recombining ancestral Y, or independent recruitment of

the same autosome as a sex chromosome.

These hypotheses make contrasting predictions regarding the gene trees and

corresponding patterns of allele sharing for SNPs in the SDRs. If the SDRs of *P. wingei* and *P.*

picta descend from a common ancestral SDR, then the Y chromosomes from both species will

cluster together to the exclusion of the X chromosomes (Trees 1 and 2 in Figure 5). If the SDRs

evolved independently in the two species, then the X and Y of each species will cluster together

(Trees 3 and 4 in Figure 5). If the Y of *P. wingei* was derived from its X chromosome after that

lineage diverged from *P. picta*, then the X and Y chromosomes of *P. wingei* will cluster together

and form a clade that is sister to the *P. picta* X chromosomes (Tree 5 in Figure 5). Other

evolutionary histories involving turnover events in which a Y was derived from an X or *vice*

versa result in three additional topologies (Trees 6 - 8 in Figure 5).

None of the 339,397 SNPs have either of the topologies expected if the SDRs of *P. picta*

and *P. wingei* are homologous (Trees 1 and 2). Instead, *P. picta* and *P. wingei* are fixed for

different alleles at 99.8% of the SNPs (Trees 3 and 4), suggesting that their SDRs evolved

independently. Unfortunately, bioinformatic artifacts resulting from Y degeneration can cause

these gene trees to be erroneously inferred. At hemizygous sites (where there is a deletion on

the Y), BCFtools and other widely-used SNP calling software impute a homozygote, which

implies the presence of a Y-linked allele that is the same as the X-linked allele. This artifact

results in gene trees in which the X and Y chromosomes from *P. picta* appear to cluster together

to the exclusion of the X and Y chromosomes of *P. wingei*, which can falsely suggest that their

SDRs evolved independently. Thus, the ubiquity of Trees 3 and 4 could result from the extreme

degeneration of the Y in *P. picta* rather than from the true evolutionary history. It is possible to

use criteria such as read depth to identify and remove hemizygous sites, but these filters are

not always accurate, especially with the very small sample size used in this study.

Further analysis suggests, however, that hemizygosity is unlikely to be responsible for all

the SNPs associated Trees 3 and 4. Tree 3, in which the derived allele is unique to the X and Y

chromosomes of *P. picta*, is 2.7 times more common than Tree 4, in which the derived allele is

unique to the X and Y chromosomes of *P. wingei*. The predominance of Tree 3 is likely to be an

artifact of degeneration in the Y of *P. picta*, as that tree will be erroneously inferred if a

mutation that is unique to the X chromosome in *P. picta* occurs in a region deleted from the Y

chromosome. Likewise, Tree 4 can be erroneously inferred because of the same bioinformatic

error if the true evolutionary history is Tree 7. That evolutionary scenario is unlikely, however,

since the origin of a Y from an X to our knowledge has never been reported. Tree 4 can also

result from a mutation that is unique to the X chromosome of *P. wingei* in a region that is

deleted from its Y chromosome. We found little evidence, however, of extensive Y

degeneration in *P. wingei* based on the read depth ratio (Figure 2). These findings suggest that

the large number of SNPs exhibiting Tree 4 (about 10% of all SNPs in our dataset, including non-

fixed sites) likely reflect the true evolutionary history of the *Poecilia* sex chromosomes. This

conclusion implies that the SDRs of *P. wingei* and *P. picta* originated independently.

The trees associated with sex chromosome turnover hypotheses (Trees 5 - 8 in Figure 5)

are all very rare, and likely result from homoplasy and/or degeneration of the Y in *P. picta*. We

first consider hypotheses based on homoplasy. At 42 SNPs, the gene tree is consistent the

origin of a new Y chromosome from the X in the ancestor of *P. wingei* (Tree 5). But at more than

twice as many SNPs, the X and Y of *P. wingei* both share a derived allele with the Y of *P. picta*

(Tree 7). As explained above, this tree is consistent with an evolutionary hypothesis that is

implausible. A more likely explanation is that Trees 5 and 7 arose when the same mutation

independently occurred on one of the *P. picta* sex chromosomes and on a recombining region

in *P. wingei* that later became part of the SDR. The greater frequency of Tree 7 relative to Tree

5 could be due to higher mutation rates on the Y than on the X in *P. picta*, as observed in

species with male-biased mutation (WILSON SAYRES AND MAKOVA 2011). Male-biased mutation

would also explain the greater frequency of Tree 8 relative to Tree 6.

Now consider how degeneration of the Y in *P. picta* might lead to the patterns seen in

Trees 5 – 8. Tree 5 can only be inferred at those rare sites in *picta* that are not hemizygous,

while Tree 6 can be erroneously inferred at hemizygous sites. This contrast could explain why

Tree 6 is much more common. The same logic could explain why Tree 8 is much more common

than Tree 7. Finally, Trees 5 and 6 could also result if the SDRs of the two species are

homologous and a SNP that evolved in the ancestral X occurs in a region that was lost from one

of the Y chromosomes. We believe this hypothesis is unlikely as that scenario is incompatible

with Trees 7 and 8, which are more common. Additionally, the SNPs showing Trees 5 and 6

occur throughout the chromosome rather than being colocalized in a region, as we expect for

an ancestral SDR.

In sum, there is no evidence that supports the homology of the SDRs in *P. wingei* and *P.*

picta. The gene trees instead suggest that these SDRs originated independently. Several details

of the data can be understood as resulting from homoplasy and bioinformatic artifacts caused

by the extreme degeneration of the Y in *P. picta*.

4: DISCUSSION

4.1: Current state of recombination on the X and Y chromosomes in the three

species

The results presented above and those from previous studies provide several insights into the

state of the sex chromosomes in these three species. In the guppy *P. reticulata*, five lines of

evidence argue that the sex chromosomes have not evolved suppressed recombination or

evolutionary strata and in fact continue to recombine in the region near to the sex determining

gene (see Figure 1).

First, we see no signs of increased divergence between males and females (a proxy for

divergence between the X and Y) in the candidate regions for sex determination or anywhere

else along the chromosome (Figures 2 and 3).

Second, gene trees on the sex chromosomes presented by DAROLTI *et al.* (2020) suggest

that crossovers have occurred in the region identified as nonrecombining by Wright *et al.*

(2017) and later studies by that research team (MORRIS *et al.* 2018; DAROLTI *et al.* 2019; DAROLTI

et al. 2020; ALMEIDA *et al.* 2021). We expect that the most recent recombinant Y chromosome

was established within the last few thousand generations, based on the effective population

size of Y chromosomes relative to autosomes and the small census population sizes of guppies

(FRASER *et al.* 2015).

Third, a male crossover has been directly observed in a region that had previously been

proposed as a nonrecombining stratum (BERGERO *et al.* 2019; CHARLESWORTH *et al.* 2020a) (see

Figure 1). It occurred near the 21 Mb position, which is also the approximate boundary

between Strata 1 and 2 as defined by ALMEIDA *et al.* (2021). While that paper states there is no

X-Y crossing over in Stratum 1, it does allow for the possibility of rare events in Stratum 2.

Fourth, SNPs with heterozygosity patterns consistent with sex linkage are scattered

across the length of the sex chromosomes (BERGERO *et al.* 2019; DAROLTI *et al.* 2020; ALMEIDA *et*

al. 2021), rather than being restricted to the proposed nonrecombining region.

Fifth and finally, Y haplotypes have high molecular diversity in guppies (ALMEIDA *et al.*

2021). This is consistent with the ongoing introduction of genetic variation by recombination

from the X to the Y. It does not seem, however, compatible with a hypothesis proposed by

Almeida *et al.*, which is that recombination in Stratum 1 was suppressed gradually. No matter

what the mechanism (an inversion, change in chromatin structure, etc.), the fixation of any *cis*

modifier that expands the SDR by so much as a single base pair eliminates all variation

throughout the entire SDR of the chromosome on which that change occurred. If all the

suppression results from inversion(s) on the X, then diversity would be reduced on that

chromosome, but that is not what the data show (Fig. 3c in ALMEIDA *et al.* (2021)). Variation

could be maintained if the SDR expanded -- either gradually or rapidly -- by one more more

unlinked (*trans*) recombination modifiers. We do not know, however, of any species in which

that has been observed, nor of any genetic mechanism that has that effect.

BERGERO *et al.* (2019) and CHARLESWORTH *et al.* (2020a) found that all chromosomes in

males have extremely low recombination rates over most of their length, consistent with earlier

reports in guppies (TRIPATHI *et al.* 2009; LISACHOV *et al.* 2015). They suggested that the very low

rate of crossing over between the X and Y is a simple consequence of this situation. We concur.

Apparently, recombination in males is sufficiently rare in at least some populations that the X

and Y chromosomes have diverged at the molecular level, *e.g.* F_{ST} between males and females

in the range 0.05 to 0.1 over large regions of the sex chromosomes (BERGERO *et al.* 2019;

ALMEIDA *et al.* 2021). (For unknown reasons, our analyses show lower levels of differentiation

between males and females, with no clear peaks that exceed values found on the autosomes

(Figure 2).)

While the degree of sex bias in recombination (heterochiasmy) is extreme in guppies,

qualitatively similar differences between males and females are seen across the eukaryotes

(SARDELL AND KIRKPATRICK 2020). In fact, there are examples of heterochiasmy even more extreme

than guppies. Recombination along almost the entire lengths of all chromosomes is extremely

rare in males of ranid frogs (BRELSFORD *et al.* 2016; JEFFRIES *et al.* 2018). As a result, F_{ST} between

males and females approaches its maximum possible value on the sex chromosomes in some

populations (TOUPS *et al.* 2019).

Variation between populations in recombination rates on the sex chromosomes is a

recurring theme in the guppy literature. Using laboratory crosses, HASKINS *et al.* (1961) showed a

significant difference between two populations in the recombination rate between the sex-

linked locus *Sb* and the sex determining gene. FRASER *et al.* (2015) reported differences between

populations in the linkage of certain color patterns to the X and Y chromosome, but their

findings regard differences in the frequencies of color pattern alleles on the X and Y

chromosomes rather than differences in recombination rates. In contrast, WRIGHT *et al.* (2017)

and ALMEIDA *et al.* (2021) argue that differences between populations in patterns of molecular

variation reflect differences in the actual rates of recombination between the X and Y.

We suggest that the very subtle differences they reported between populations in the

degree of molecular differentiation between the guppy X and Y could occur even in the absence

of differences in the recombination landscape. Consider the consequences of a rare crossover

between the X and Y, for example near to the male determining gene. If the new Y haplotype

spreads to high frequency by selection or drift, differentiation between the X and Y will be

erased from the crossover breakpoint down the rest of the chromosome distal to the male

determining gene. In the headwaters of the streams where they live, population sizes of *P.*

reticulata are only a couple thousand individuals (FRASER *et al.* 2015). Consequently, crossovers

between the X and Y may occur very infrequently, giving the X and Y time to diverge slightly

before the next recombinant Y chromosome is established. Downstream populations are many

times larger (FRASER *et al.* 2015). The result will be that recombinant Y chromosomes appear

much more frequently in downstream populations and so divergence between the X and Y has

less opportunity to build up.

A second factor that could contribute to patterns of molecular variation is sexually

antagonistic selection, or SAS. More than 50 traits that are under sexual selection in male *P.*

reticulata have been mapped to the sex chromosomes (LINDHOLM AND BREDEN 2002). These are

expected to generate peaks in the divergence between the X and Y in neutral genetic variation

and can generate patterns that give the appearance of suppressed recombination

(CHARLESWORTH 2018; BERGERO *et al.* 2019). Recombination in *P. reticulata* males is so rare that

the map length of almost the entire Y chromosome is much less than 1 cM (BERGERO *et al.* 2019).

The small census population sizes suggest that many targets of SAS may lie within less than one

ρ (= $N_e r/2$) of the sex determining factor, a condition favorable to inflated values of F_{ST}

between the X and Y (KIRKPATRICK AND GUERRERO 2014). Indeed, peaks in F_{ST} between males and

females (a proxy for divergence between the X and Y) consistent with SAS have been reported

by BERGERO *et al.* (2019) and are visible in Figure 1 of ALMEIDA *et al.* (2021). Thus differences

between populations in the intensity of sexual selection and the frequencies of alleles at loci

that experience SAS could also contribute to the differences between populations in the degree

of X-Y divergence even in the absence of variation in recombination rates.

The recent publication of a greatly improved reference genome for *P. reticulata* that

includes the Y (FRASER *et al.* 2020) adds additional insight. Those authors concluded that if an

SDR exists, it is very small (less than 1 Mb). Of the two candidate regions for sex determination

that they found, sequence from the most likely one is absent from the X chromosome. This

would explain why it was not detected in studies (including ours) that are based on earlier

reference genomes that lack Y-specific sequences. Analysis of gene trees might determine

which of the two candidate regions identified by FRASER *et al.* (2020) harbors the sex

determination gene.

In the guppy's sister species, *P. wingei*, molecular divergence between males and

females suggests that the X and Y have extremely low or no recombination over most of their

lengths. Divergence is clearly evident in four of the five statistics shown in Figures 2 and 3.

Within that large region, there is a smaller segment of about 10 Mb that shows greatly elevated

female-specific allele density. This region could reflect a polymorphic inversion on the X

chromosome of *P. wingei* (NANDA *et al.* 2014). If so, the patterns seen in the figures could result

if the three X chromosomes in our sample of *P. wingei* males are monomorphic for one

arrangement while the six X chromosomes in the females include both arrangements. We are

not able to determine if there is more than one stratum blocking recombination between the X

and Y in *P. wingei*. A segment of the sex chromosomes distal to the centromere shows no sign

of molecular divergence between the X and Y, suggesting high rates of recombination there.

The most dramatic patterns are seen in *P. picta*. Consistent with the findings of Darolti

et al. (2019), our analyses suggest that most of its Y chromosome no longer recombines, and

much of it has degenerated by sequence divergence or deletion (Figures 2 and 3).

4.2: History of the sex chromosomes and their SDRs

Our analysis of gene trees lead us to reject the hypotheses that the SDRs of *P. picta* and

P. wingei are largely homologous. Although this conclusion is complicated by degeneration of

the Y in *P. picta*, a study of *Gasterosteus* sticklebacks showed a clear signal of SDR homology

using gene trees, even though the Y chromosome of one of the species shows about the same

degree of degeneration as the Y of *P. picta* (SARDELL *et al.* 2021). We likewise reject an

alternative hypothesis is that a new Y was derived from its X in *P. wingei* after that lineage

diverged from *P. picta*.

The most parsimonious explanation for the data is one originally proposed by DAROLTI *et*

al. (2019): the guppy and its two congeners share an ancestral Y chromosome, one with an

extremely small SDR (as in *P. reticulata*). This would explain the absence of signal of homology

in our gene trees, as they can only detect homology of non-recombining regions of the Y

chromosome. But in contrast to the conclusions of Darolti *et al.*, we find that suppressed

recombination between the X and Y most likely evolved independently in the ancestors of *P.*

picta and *P. wingei*. Under this hypothesis, the Y may be much more degenerate in *P. picta*

either because recombination between its X and Y was suppressed much earlier or because

degeneration proceeded much more rapidly than in *P. wingei*. Both situations are seen in

Gasterosteus sticklebacks (SARDELL *et al.* 2021). The homology of the SDRs in *P. picta* and *P.*

wingei might best be investigated by long-read sequencing technology to determine if these

species share Y-specific sequences identified in the guppy by Fraser *et al.* (2020).

4.3: Reconciliation of past studies and best practices going forward

How can the many discrepancies between the conclusions from previous studies be reconciled,

and how best can the field move forward? Some of the problems can be traced to semantic

differences. We defer to the terminology for sex chromosomes defined by BACHTROG *et al.*

(2011). The sex determining region, or SDR, is a segment of the sex chromosomes that includes

the sex determining factor and in which crossing over between the X and Y is completely

suppressed. Within the SDR, there can be one or more strata, which are regions in which

crossing over was suppressed at different times in the past (LAHN AND PAGE 1999; CHARLESWORTH

2018). The SDR may be as small as a nucleotide, as in the case of the fugu, or as large as most of

the Y chromosome, as in mammals (BACHTROG *et al.* 2014). By this definition, it is a *non sequitur*

to say there is a stratum on the guppy sex chromosomes in which there is rare crossover

recombination (DAROLTI *et al.* 2020; ALMEIDA *et al.* 2021). Again following BACHTROG *et al.* (2011),

all of the sex chromosomes that fall outside of the SDR make up the pseudoautosomal region,

or PAR, regardless of whether the local recombination rate (measured as cM/Mb) is very high,

as in mammals, or very low, as in some frogs (BACHTROG *et al.* 2014). By this definition, it is also

a *non sequitur* to refer to only part of the recombining region as the PAR (BERGERO *et al.* 2019).

Authors are of course free to adopt noncanonical definitions, but in that case, we urge them to

give explicit definitions and use terms consistently.

There are, however, three substantive reasons why different studies have arrived at

differing conclusions. First, some statistical approaches used to define the SDRs and PARs of the

guppy and its relatives are problematic. Two nonrecombining strata were identified in *P.*

reticulata and *P. wingei* using the criterion that molecular differences between males and

females over a region of the sex chromosome fall outside the 95% confidence interval seen on

autosomes (WRIGHT *et al.* 2017; DAROLTI *et al.* 2019). By that standard, 5% of the entire genome

is expected to be identified as nonrecombining sex chromosome strata, and indeed several

regions of autosomes do meet that criterion (Supplementary Figure 1 in WRIGHT *et al.* 2017).

Further, the differences between males and females used to define nonrecombining strata in

those studies are extremely small and likely not biologically meaningful (*e.g.* a male:female SNP

density ratio differing much less than 1% from the autosomal average (WRIGHT *et al.* 2017)).

Additionally, three studies have given unambiguous evidence that the Y chromosome of *P.*

reticulata recombines in or very near the region that had been proposed as a nonrecombining

stratum (Figure 1) (BERGERO AND CHARLESWORTH 2019; CHARLESWORTH *et al.* 2020a; DAROLTI *et al.*

2020). Thus the bioinformatic strategy used by WRIGHT *et al.* (2017), DAROLTI *et al.* (2019), and

ALMEIDA *et al.* (2021) appears to be inadequate to identify SDRs and strata. Finally, although the

authors state otherwise, the gene trees presented by DAROLTI *et. al*. (2020) provide strong

evidence of ongoing recombination, as only one poorly-supported gene tree shows the

topology expected for a non-recombining region.

Second, the results from linkage mapping experiments have been interpreted in

different ways. It is difficult to draw conclusions from a failure to observe crossovers, especially

in crosses with few offspring and when recombination is rare. Mapping experiments are

underpowered to distinguish between partial and full linkage of loci to the male determining

gene (MUYLE *et al.* 2016; WRIGHT *et al.* 2019). This limitation is particularly acute in species with

extremely low recombination rates in males, such as the guppy. Conclusions about regions

where the X and Y do not recombine based on linkage mapping should be treated with caution.

Methods that define nonrecombining sex determining regions using gene trees are much more

sensitive because they integrate genetic signatures of recombination over long periods of

evolutionary time (*e.g.* (LAHN AND PAGE 1999; ZHOU *et al.* 2014; DIXON *et al.* 2018; SARDELL *et al.*

2021).

Third, sequencing studies have used different reference genomes and different

strategies to map reads. WRIGHT *et al.* (2017) used a *de novo* assembly of *P. reticulata* (N50 =

0.017 Mb) whose scaffolds were then oriented according to a guppy reference. BERGERO *et al.*

(2019) and CHARLESWORTH *et al.* (2020a) mapped their DNA reads to the old *P. reticulata*

reference (scaffold N50 = 31.4 Mb; KÜNSTNER *et al.* (2016)). While this reference has the

advantage of being from the species of interest, it is based entirely on scaffolded short-read

sequences. DAROLTI *et al.* (2019; 2020) used one of the publicly-available *Xiphophorus helleri*

genomes (scaffold N50 = 29.4 Mb; SHEN *et al.* (2016)) to order scaffolds from their own *de novo*

genome assemblies of *P. reticulata*. Last, for most of the analyses in this paper we used the

Xiphophorus maculatus reference because it has the highest quality of any species in this family

(SCHARTL *et al.* 2013). The current assembly (version 5) is based on long-read as well as short-

read sequencing and optical mapping to produce the chromosome-level assembly (scaffold N50

= 31.5 Mb). To verify our findings in *X. maculatus*, we re-analyzed the gene trees on Chr 12

using the new *P. reticulata* reference genome (scaffold N50 = 32.8 Mb) that was recently

released by FRASER *et al.* (2020).

Many of the obstacles that have made studies of the guppy sex chromosomes so

difficult will hopefully soon be in the past as the use of experimental phasing of sex

chromosomes (SARDELL *et al.* 2021) and advances in long-read sequencing eliminate many of the

bioinformatic problems.

ACKNOWLEDGEMENTS

We are grateful to members of the Kirkpatrick lab and two anonymous reviewers for discussion

and comments. This study was supported by NIH grant R01-GM116853 to M.K..

DATA ACCESSIBILITY

All the data used in the study were acquired from the publicly-accessible sources cited in the

text. The custom scripts used for data processing, analysis, and figure generation are available

on GitHub at https://github.com/grovesdixon/guppy_sex_chroms.

AUTHOR CONTRIBUTIONS

M.K. conceived of the study. J.M.S., B.J.P., and G.D.D. conducted the analyses. All authors

contributed to the writing.

REFERENCES

Almeida, P., B. A. Sandkam, J. Morris, I. Darolti, F. Breden *et al.*, 2021 Divergence and remarkable diversity of the Y chromosome in guppies. Molecular Biology and Evolution 38**:** 619-633.

Andrews, S., 2010 FastQC: a quality control tool for high throughput sequence data, pp.

Bachtrog, D., M. Kirkpatrick, J. E. Mank, S. F. McDaniel, J. C. Pires *et al.*, 2011 Are all sex chromosomes created equal? Trends in Genetics 27**:** 350-357.

Bachtrog, D., J. Mank, C. Peichel, S. Otto, M. Kirkpatrick *et al.*, 2014 Sex determination: Why so many ways of doing it? PLoS Biology 12**:** e1001899.

Bergero, R., and D. Charlesworth, 2019 Reply to Wright *et al.*: How to explain the absence of extensive Y-specific regions in the guppy sex chromosomes. Proceedings of the National Academy of Sciences of the United States of America 116**:** 12609-12610.

Bergero, R., J. Gardner, B. Bader, L. Yong and D. Charlesworth, 2019 Exaggerated heterochiasmy in a fish with sex-linked male coloration polymorphisms. Proceedings of the National Academy of Sciences of the United States of America 116**:** 6924-6931.

Brelsford, A., N. Rodrigues and N. Perrin, 2016 High-density linkage maps fail to detect any genetic component to sex determination in a *Rana temporaria* family. Journal of Evolutionary Biology 29**:** 220-225.

Broad Institute, 2019 *Picard Toolkit*.

Bushnell, B., 2014 BBMap: a fast, accurate, splice-aware aligner pp. Lawrence Berkeley National Laboratory, Berkeley, CA.

Charlesworth, D., 2017 Evolution of recombination rates between sex chromosomes. Philosophical Transactions of the Royal Society B-Biological Sciences 372**:** 20160456.

Charlesworth, D., 2018 The guppy sex chromosome system and the sexually antagonistic polymorphism hypothesis for Y chromosome recombination suppression. Genes 9**:** 264.

Charlesworth, D., R. Bergero, C. Graham, J. Gardner and L. Yong, 2020a Locating the sex determining region of linkage group 12 of guppy (*Poecilia reticulata*). G3-Genes Genomes Genetics 10**:** 3639-3649.

Charlesworth, D., Y. X. Zhang, R. Bergero, C. Graham, J. Gardner *et al.*, 2020b Using GC content to compare recombination patterns on the sex chromosomes and autosomes of the guppy, *Poecilia reticulata*, and its close outgroup species. Molecular Biology and Evolution 37**:** 3550-3562.

Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks *et al.*, 2011 The variant call format and VCFtools. Bioinformatics 27**:** 2156-2158.

Darolti, I., A. E. Wright and J. E. Mank, 2020 Guppy Y chromosome integrity maintained by incomplete recombination suppression. Genome Biology and Evolution 12**:** 965-977.

Darolti, I., A. E. Wright, B. A. Sandkam, J. Morris, N. I. Bloch *et al.*, 2019 Extreme heterogeneity in sex chromosome differentiation and dosage compensation in livebearers. Proceedings of the National Academy of Sciences of the United States of America 116**:** 19031-19036.

Dixon, G., J. Kitano and M. Kirkpatrick, 2018 The origin of a new sex chromosome by
introgression between two stickleback fishes. Molecular Biology and Evolution 36: 28-
38.
Fraser, B. A., A. Kunstner, D. N. Reznick, C. Dreyer and D. Weigel, 2015 Population genomics of
natural and experimental populations of guppies (*Poecilia reticulata*). Molecular Ecology
24: 389-408.
Fraser, B. A., J. R. Whiting, J. R. Paris, C. J. Weadick, P. J. Parsons *et al.*, 2020 Improved reference
genome uncovers novel sex-linked regions in the guppy (*Poecilia reticulata*). Genome
Biology and Evolution 12: 1789-1805.
Haskins, C. P., E. F. Haskins, J. J. A. McLaughlin and R. E. Hewitt, 1961 Polymorphism and
population structure in *Lebistes reticulatus*, a population study, pp. 320-395 in
Vertebrate Speciation, edited by W. F. Blair. University of Texas Press, Austin.
Jeffries, D. L., G. Lavanchy, R. Sermier, M. J. Sredl, I. Miura *et al.*, 2018 A rapid rate of sex-
chromosome turnover and non-random transitions in true frogs. Nature
Communications 9: 4088.
Kirkpatrick, M., and R. F. Guerrero, 2014 Signatures of sex-antagonistic selection on
recombining sex chromosomes. Genetics 197: 531-541.
Künstner, A., M. Hoffmann, B. A. Fraser, V. A. Kottler, E. Sharma *et al.*, 2016 The genome of the
Trinidadian guppy, *Poecilia reticulata*, and variation in the Guanapo population. Plos
One 11: e0169087.
Lahn, B. T., and D. C. Page, 1999 Four evolutionary strata on the human X chromosome (vol
286, pg 964, 1999). Science 286: 2273-2273.
Langmead, B., and S. L. Salzberg, 2012 Fast gapped-read alignment with Bowtie 2. Nature
Methods 9: 357-359.
Li, H., 2011 A statistical framework for SNP calling, mutation discovery, association mapping
and population genetical parameter estimation from sequencing data. Bioinformatics
27: 2987-2993.
Li, H., 2018 Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34: 3094-
3100.
Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The sequence alignment/map
format and SAMtools. Bioinformatics 25: 2078-2079.
Lindholm, A., and F. Breden, 2002 Sex chromosomes and sexual selection in poeciliid fishes.
American Naturalist 160: S214-S224.
Lisachov, A. P., K. S. Zadesenets, N. B. Rubtsov and P. M. Borodin, 2015 Sex chromosome
synapsis and recombination in male guppies. Zebrafish 12: 174-180.
Martin, M., 2011 Cutadapt removes adapter sequences. from high-throughput sequencing
reads. EMBnet Journal 17: 10-12.
Meisel, R. P., 2020 Evolution of sex determination and sex chromosomes: a novel alternative
paradigm. BioEssays 2020: 1900212.
Morris, J., I. Darolti, N. I. Bloch, A. E. Wright and J. E. Mank, 2018 Shared and species-specific
patterns of nascent Y chromosome evolution in two guppy species. Genes 9: 238.
Muyle, A., J. Kafer, N. Zemp, S. Mousset, F. Picard *et al.*, 2016 SEX-DETector: a probabilistic
approach to study sex chromosomes in non-model organisms. Genome Biology and
Evolution 8: 2530-2543.

Nanda, I., S. Schories, N. Tripathi, C. Dreyer, T. Haaf *et al.*, 2014 Sex chromosome polymorphism
in guppies. Chromosoma 123**:** 373-383.
Quinlan, A. R., and I. M. Hall, 2010 BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics 26**:** 841-842.
Sardell, J. M., M. P. Josephson, A. C. Dalziel, C. L. Peichel and M. Kirkpatrick, 2021
Heterogeneous histories of recombination suppression on stickleback sex
chromosomes. Molecular Biology and Evolution msab179.
Sardell, J. M., and M. Kirkpatrick, 2020 Sex differences in the recombination landscape.
American Naturalist 195**:** 361-379.
Schartl, M., R. B. Walter, Y. J. Shen, T. Garcia, J. Catchen *et al.*, 2013 The genome of the
platyfish, *Xiphophorus maculatus*, provides insights into evolutionary adaptation and
several complex traits. Nature Genetics 45**:** 567-572.
Shen, Y. J., D. Chalopin, T. Garcia, M. Boswell, W. Boswell *et al.*, 2016 *X. couchianus* and *X.*
hellerii genome models provide genomic variation insight among *Xiphophorus* species.
BMC Genomics 17**:** 67.
Toups, M. A., N. Rodrigues, N. Perrin and M. Kirkpatrick, 2019 A reciprocal translocation
radically reshapes sex-linked inheritance in the common frog. Molecular Ecology 28**:**
1877-1889.
Tripathi, N., M. Hoffmann, E.-M. Willing, C. Lanz, D. Weigel *et al.*, 2009 Genetic linkage map of
the guppy, *Poecilia reticulata*, and quantitative trait loci analysis of male size and colour
variation. Proceedings of the Royal Society of London. Series B: Biological Sciences 276**:**
2195-2208.
Vicoso, B., 2019 Molecular and evolutionary dynamics of animal sex-chromosome turnover.
Nature Ecology & Evolution 3**:** 1632-1641.
Vicoso, B., and D. Bachtrog, 2015 Numerous transitions of sex chromosomes in Diptera. Plos
Biology 13**:** e1002078.
Weir, B. S., and C. C. Cockerham, 1984 Estimating F-statistics for the analysis of population
structure. Evolution 38**:** 1358-1370.
Wilson Sayres, M. A., and K. D. Makova, 2011 Genome analyses substantiate male mutation
bias in many species. BioEssays 33**:** 938-945.
Winge, O., 1922 One-sided masculine and sex-linked inheritance in *Lebistes reticulatus*. Journal
of Genetics 12**:** 145-162.
Wright, A. E., I. Darolti, N. I. Bloch, V. Oostra, B. Sandkam *et al.*, 2017 Convergent
recombination suppression suggests role of sexual selection in guppy sex chromosome
formation. Nature Communications 8**:** 14251.
Wright, A. E., I. Darolti, N. I. Bloch, V. Oostra, B. A. Sandkam *et al.*, 2019 On the power to detect
rare recombination events. Proceedings of the National Academy of Sciences of the
United States of America 116**:** 12607-12608.
Zhou, Q., J. L. Zhang, D. Bachtrog, N. An, Q. F. Huang *et al.*, 2014 Complex evolutionary
trajectories of sex chromosomes across bird taxa. Science 346**:** 1246338.
728

Figure 1: The sex chromosome of the guppy, *P. reticulata*. The horizontal green bar

shows the interval in which a crossover was observed by BERGERO *et al.* (2019; D.

CHARLESWORTH pers. comm.). Stratum 1 and Stratum 2 are regions where WRIGHT *et al.*

(2017) reported that the X and Y do not recombine. Blue boxes labeled "Sex" are

candidate regions for the male-determining factor (FRASER *et al.* 2020). At far right is a

region with a high local recombination rate (LISACHOV *et al.* 2015, BERGERO *et al.* 2019,

DAROLTI *et al.* 2019). Vertical blue lines show locations of the 42 loci at which gene trees

from *P. reticulata* (Xr, Yr) and *P. wingei* (Xw, Yw) were estimated by DAROLTI *et al.* (2020).

Only the gene tree highlighted in the red box has a topology consistent with an ancestral

SDR shared by the two species; examples of four other representative gene trees are

also shown. Numbers at their nodes give the bootstrap support. The centromere is

shown as the green circle at left.

Figure 2. Divergence between the X and Y chromosomes in the three species as

reflected by two statistics that measure differences between the sexes: the male:female

read depth, and the male:female F_{ST}. The gray horizontal dashed lines show the bottom

2.5% and top 97.5% intervals based on windows from all autosomes. The blue curves

are smoothed regressions. Green circles at the left of the Y axes represent the

centromere. The phylogeny is shown at top.

Figure 3. Divergence between the X and Y chromosomes as measured by three additional statistics: the male:female SNP density ratio, the SDR-like SNP density, and the female-specific SNP density. The gray horizontal dashed lines show the bottom 2.5% and top 97.5% intervals based on windows from all autosomes. The blue curves are smoothed regressions. Green circles at the left of the Y axes represent the centromere. The phylogeny is shown at top.

767   **Figure 4.** Densities per base pair of SDR-like SNPs (see Methods) that map to the sex

768   chromosomes and to the autosomes.

Figure 5. Ten trees for the SDRs of *P. picta* and *P. wingei* inferred from allele sharing patterns at topologically informative SNPs. X_p and Y_p represent the sex chromosomes of *P. picta*; X_w and Y_w represent the sex chromosomes of *P. wingei*. Derived mutations are shown by stars. Indicated are the numbers of SNPs at which each tree is seen and a biological interpretation of that tree. All trees can also result from introgression, homoplasy, and sequencing error. Note that the subtrees descending from the mutations shown in Trees 5 – 8 are consistent with the proposed biological scenario, but the allele sharing patterns observed at those SNPs give no information about the evolutionary relations of the three types of sex chromosomes that carry the mutant allele.