Apoptosis in plants: from semantic appeal to empirical rejection

Elena A. Minina[1,2], Adrian N. Dauphinee[1], Florentine Ballhaus[1], Vladimir Gogvadze[3,4], Andrei P. Smertenko[5] and Peter V. Bozhkov[1]

[1]Department of Molecular Sciences, Uppsala BioCenter, Swedish University of Agricultural Sciences and Linnean Center for Plant Biology, P.O. Box 7015, Uppsala, SE-750 07, Sweden

[2]COS, Heidelberg University. Im Neuenheimer Feld 230. 69120 Heidelberg, Germany

[3]Institute of Environmental Medicine, Division of Toxicology, Karolinska Institutet, Box 210, Stockholm, SE-171 77, Sweden.

[4]Faculty of Medicine, MV Lomonosov Moscow State, University, 119991 Moscow, Russia

[5]Institute of Biological Chemistry, College of Human, Agricultural, and Natural Resource Sciences, Washington State University, Pullman, WA 99164, USA.

Correspondence to Elena A. Minina (alena.minina@slu.se) or Peter V. Bozhkov (peter.bozhkov@slu.se)

* E. A. Minina and A. N. Dauphinee contributed equally to this paper.

Abstract

Animals and plants diverged over one billion years ago and evolved unique mechanisms for many cellular processes, including cell death. Plants lack key instruments of apoptotic cell death, including caspases, Bcl-2 family proteins, and phagocytosis. This casts serious doubts on the occurrence of apoptosis in these organisms. Nevertheless, the concept of plant apoptosis or "apoptotic-like programmed cell death" ("AL-PCD") is wide-spread, primarily due to superficial resemblance between protoplast shrinkage in plant cells dying in response to stress and apoptotic volume decrease preceding animal cell fragmentation into apoptotic bodies. Here, we provide a comprehensive spatio-temporal analysis of cytological and biochemical events occurring during plant cell death previously classified as "AL-PCD". We show that under "AL-PCD" inducing conditions, protoplast shrinkage does not proceed to membrane blebbing and formation of apoptotic bodies. Instead, it coincides with instant ATP depletion and irreversible loss of plasma membrane integrity. Furthermore, neither uncoupling of mitochondria nor Ca^{2+} chelation can prevent protoplast shrinkage and cell death, pointing to passive, energy-independent cell collapse perfectly matching definition of necrosis. Although one cannot exclude the possibility that the very early steps of this cell death are genetically regulated, classifying it as apoptosis or "AL-PCD" is misconception. Our study invalidates the notion that apoptosis is conserved across animal and plant kingdoms and demonstrates the importance of focusing on plant-specific aspects of cell death pathway.

Keywords: apoptosis, apoptotic-like programmed cell death, ferroptosis, heat shock, mitochondrial dysfunction, necrosis, plant cells, plasma membrane integrity, protoplast shrinkage.

Introduction

Programmed cell death (PCD) is a process of cellular suicide aimed at maintaining proper development by counteracting cell proliferation and removing aged cells (Jacobson et al., 1997). PCD also plays an essential role in morphogenesis by eliminating surplus cells and shaping new structures (Suzanne and Steller, 2013; Daneva et al., 2016; Huysmans et al., 2017). Furthermore, stress-triggered PCD restricts the spread of pathogens through tissues and confines damage caused by abiotic factors (Coll et al., 2011; Huysmans et al., 2017; Jorgensen et al., 2017).

In their seminal work published almost half a century ago, Kerr and colleagues (Kerr et al., 1972) named the newly described type of PCD in animals after the process of "falling off" of flower petals or leaves, apoptosis (ancient Greek ἀπόπτωσις, apóptōsis, "falling off"). Since then the apoptotic cell death in animals became one of the most studied pathways (for review see Letai, 2017), while our understanding of plant PCD remains fragmented.

The utmost importance of PCD for all multicellular organisms gave weight to the hypothesis that this process should be evolutionary conserved. Indeed, PCD in animals and plants share quite a number of general similarities attributed to the common features of all eukaryotic cells, e.g. dependency on ATP and protein synthesis, regulated proteolysis, and control over toxicity of the dying cell to its neighbours (Kroemer et al., 2005; van Doorn et al., 2011). On the other hand, differences in plant and animal cell morphology, motility, genome maintenance, trophic strategies and body plan plasticity (Murat et al., 2012; Drost et al., 2017) create very different sets of challenges for development, homeostasis and adaptation to the environment, powering diversification of cell death mechanisms (Bozhkov and Lam, 2011).

After the discovery of apoptosis, numerous non-apoptotic types of PCD were described in animals (Galluzzi et al., 2015, 2018), such as paraptosis (Sperandio et al., 2000), necroptosis (Molnár et al., 2019), ferroptosis (Dixon and Stockwell, 2019), pyroptosis (Miao et al., 2010), and others. Since animals possess multiple PCD pathways, each optimized for a specific physiological requirement, there is no reason to assume that plant evolution did not shape plant-specific PCD mechanisms.

A strong evidence for the existence of plant-specific PCD mechanisms is the lack of plant

orthologues for the core apoptotic regulators: caspases and Bcl-2 family proteins (Ameisen, 2002).

The closest plant orthologues of caspases, metacaspases, have a different substrate specificity and

regulatory features (Minina et al., 2020). It is not yet fully explored what subset of proteases is

responsible for caspase-like activity detected during plant PCD (for review see Salvesen et al.,

2016). Bcl-2 family proteins play either pro- or anti-apoptotic functions in animals by regulating

release of cytochrome c from mitochondria (Edlich, 2018). Interestingly, ectopic expression of

genes encoding pro- or anti-apoptotic Bcl-2 family members in plants can trigger cell death or

reduce its frequency, respectively (Lacomme and Santa Cruz, 1999; Mitsuhara et al., 1999),

suggesting potential conservation of the cytochrome c release function in animal and plant cell

deaths. However, the lack of Bcl-2 genes in plant genomes clearly indicates that although

cytochrome c release might contribute to plant PCD under certain conditions, it is regulated

differently from the apoptotic pathway (Yu et al., 2002; Martínez-Fábregas et al., 2013).

Nevertheless, original research and reviews on apoptotic features in plants PCD are still being

published (for recent reviews, see Dickman et al., 2017; Kabbage et al., 2017; Valandro et al.,

2020) propagating controversial conclusions that require careful consideration.

Apoptotic cell death has typical hallmarks: (i) it is an active, ATP- and caspase-dependent process

(Tsujimoto, 1997; Hengartner, 2000); (ii) the plasma membrane (PM) integrity is retained

throughout the cell death process and phosphatidylserine is exposed on the outer membrane surface

as an "eat-me" signal for phagocytes (Segawa and Nagata, 2015; Zhang et al., 2018); (iii) cell

shrinkage (apoptotic volume decrease, AVD) is followed by chromatin condensation and nuclear

segmentation (Galluzzi et al., 2012, 2015, 2018); (iv) PM blebbing results in cell fragmentation

into apoptotic bodies (Kroemer et al., 2005; Atkin-Smith and Poon, 2017) and their subsequent

phagocytosis (Arandjelovic and Ravichandran, 2015).

The term "apoptotic-like PCD" ("AL-PCD") was coined due to morphological and biochemical

resemblance of stress-associated plant cell death to apoptosis, in particular AVD-like protoplast

shrinkage, but also Ca^{2+}- and ATP-dependency, maintenance of PM integrity and involvement of

caspase-like activity (Balk et al., 1999; Vacca et al., 2004; Hogg et al., 2011; Kacprzyk et al.,

2017). Notably, "AL-PCD" is induced by an acute stress, e.g. a pulse heat shock (HS) at 55°C, a

temperature approximately 30°C above the physiologically relevant conditions that might in fact
cause fatal damage to the cell leading to necrosis. Unlike apoptosis, necrosis is associated with a
drop of cellular ATP content, does not depend on energy or caspases, and displays early PM
permeabilization leading to the leakage of cellular contents into the extracellular environment
(Ankarcrona et al., 1995; Majno and Joris, 1995).

To reconcile proposed features of "AL-PCD" with the apparent lack of the key components of the
apoptotic machinery in plants, we systematically analysed reported "AL-PCD" hallmarks using
appropriate controls for potential technical pitfalls. This revealed that while protoplast shrinkage
superficially resembles AVD, it is neither dependent on ATP nor regulated by Ca^{2+}, but instead
coincides with instant and irreversible loss of PM integrity. We also demonstrate the absence of
key apoptotic events such as PM blebbing, nuclear segmentation, and formation of apoptotic
bodies. In summary, we establish plant "AL-PCD" as an energy-independent process bearing only
superficial resemblance to apoptosis, but perfectly matching definition of necrosis.

Results and discussion

HS-induced plant cell death is morphologically distinct from apoptosis

One of the earliest hallmarks of apoptosis is AVD (Bortner and Cidlowski, 2002, 2007), followed by blebbing of the PM, nuclear segmentation and finally fragmentation of the cell into apoptotic bodies that facilitates their uptake by phagocytes (Arandjelovic and Ravichandran, 2015). Death of plant cells caused by abiotic or biotic stress can be accompanied by the protoplast shrinkage, a phenomenon repeatedly exploited to support existence of plant apoptosis or "AL-PCD" (Kabbage et al., 2017; Dickman et al., 2017). However, as plants lack phagocytes, apoptotic body formation during plant cell death would unlikely serve a purpose (Bozhkov and Lam, 2011; Green and Fitzgerald, 2016). Nevertheless, there are studies claiming the occurrence of the PM blebbing and the formation of apoptotic bodies during plant PCD (for review see Dickman et al., 2017).

To examine AVD, PM blebbing, nuclear segmentation and cellular fragmentation in plants, we analysed morphology of Bright Yellow-2 tobacco cell cultures (BY-2) under HS conditions that induce "AL-PCD" (Balk et al., 1999; Kacprzyk et al., 2017). After a pulse HS at 55°C, the cell culture was stained with Sytox Orange (SO) nucleic acid dye to visualize cells with compromised PM integrity and with the styryl dye FM4-64 to visualize cell membranes (**Fig. 1, Supplementary Video S1**).

SO-negative cells lacked any apparent AVD or PM blebbing (**Fig. 1A, B**). On the contrary, the SO-positive cells exhibited protoplast shrinkage and atypical vesicle-like structures at the inner side of the PM (**Fig. 1A, B**). The cells became SO-positive already within 10 min of the HS indicating rapid PM permeabilization. The PM and shrunken protoplast were imaged at 15 min, 1, 2, 4, 6, 24, 48, and 72 h after the pulse HS (data not shown). Nevertheless, we failed to detect PM blebbing or protoplast fragmentation into discrete bodies at any time point. Furthermore, although we did observe moderate nuclear condensation upon HS, it was not followed by segmentation of the organelle (**Fig. 1C, D**). In summary, the gross morphological changes of cells undergoing "AL-PCD" do not resemble apoptosis.

PM integrity is irreversibly compromised during or shortly after pulse HS

Intact PM is a pivotal hallmark of apoptosis (Kroemer et al., 2005). However, SO staining

suggested permeabilization of PM in most BY-2 cells already within 10 min of the HS. To

investigate dynamics of the PM permeabilization, we analysed cellular content leakage after two

types of HS: 55°C or 85°C, which were reported to induce "AL-PCD" or necrosis, respectively

(McCabe et al., 1997; Reape and McCabe, 2013). The leakage of cellular content was assessed

using a fluorescein diacetate (FDA)-based fluorochromatic assay (Rotman and Papermaster,

1966). In brief, the non-fluorescent FDA molecules can passively diffuse into the living cells

where their acetate groups are cleaved off by esterases. The resulting fluorescein molecules have

poor membrane permeability and are retained in cells with an intact PM, but are released into

extracellular space upon PM permeabilization.

We imaged BY-2 cells loaded with FDA prior to the pulse HS at 55°C or 85°C (**Fig. 2A**). Both

types of HS caused rapid (within 10 min) leakage of the dye into the extracellular space (**Fig. 2A**).

We measured the amount of fluorescein accumulated in the extracellular space immediately after

the HS and found that the rate of cellular content leakage in HS-treated cells is comparable to that

occurring after freeze-thaw of cells in liquid nitrogen (**Fig. 2B**).

To determine whether PM permeabilization was transient or permanent, we added SO to the cell

cultures either before or 1 h after HS. If PM permeabilization upon HS was transient, cultures

stained after HS would show a significantly lower frequency of SO staining. However, SO staining

before and after 55°C or 85°C HS showed no differences in the proportion of SO-positive cells

(**Fig. 2C, D**), indicating that both treatments caused irreversible rapid permeabilization of PM

typical for necrosis (Kroemer et al., 2005; Zong and Thompson, 2006; van Doorn et al., 2011).

The protoplast shrinkage during HS-induced cell death is ATP- and Ca^{2+}-independent

Dismantling of the apoptotic cells is ATP-dependent (Tsujimoto, 1997; Zamaraeva et al., 2005).

Yet, loss of the PM integrity would lead to rapid depletion of intracellular ATP, rendering all

energy-dependent processes defunct. To examine whether the HS-induced cell death requires ATP,

we first imaged mitochondria after 55°C and 85°C HS. Within 4 min after HS, under both

temperatures, mitochondrial dye MitoTracker localized to aberrant structures similar to those

observed after treatment with mitochondrial uncoupler and ATP synthesis inhibitor, carbonyl

cyanide m-chlorophenylhydrazone (CCCP) (Perry et al., 2011), indicating disruption of

mitochondrial membrane potential (MMP; **Fig. 3A**). Furthermore, intracellular ATP content

dropped dramatically after both HS treatments (**Fig. 3B**), most probably due to dissipation of the

MMP and leakage of cytoplasmic content through the permeabilized PM.

In addition to ATP depletion, PM permeabilization would also cause entry of Ca^{2+} into the cells,

potentially followed by its accumulation in mitochondria. MMP-driven accumulation of Ca^{2+} can

trigger mitochondrial permeability transition (MPT) due to the opening of a nonspecific pore

(MPTP) in the inner mitochondrial membrane (Hunter and Haworth, 1979). Opening of MPTP

leads to a collapse of the MMP, thereby causing arrest in ATP synthesis, and production of reactive

oxygen species leading to necrotic cell death (Kristián and Siesjö, 1998). Although HS-induced

plant cell death was previously suggested to be a Ca^{2+}-dependent process (Kacprzyk et al., 2017),

those experiments lacked controls for mitochondrial phenotype, respiration, or ATP production.

To test whether MPT plays a role in the observed mitochondrial phenotype, experiments were

performed in the presence of cyclosporin A (CsA), an inhibitor of MPTP opening, or the Ca^{2+}

chelator EGTA. Neither CsA nor EGTA could alleviate the mitochondrial phenotype in cells

subjected to the HS (**Fig. 3C, D**), indicating that mitochondrial malfunction was caused by the

direct loss of the mitochondrial membrane integrity during the HS, independently on PM

permeabilization.

Next, we examined the importance of intracellular ATP for the AVD-like protoplast shrinkage.

We found that pre-treatment of the cell cultures with CCCP prior to HS had no effect on the

protoplast shrinkage (**Figs. S1A, B**), demonstrating that the protoplast shrinkage does not require

ATP. Time-resolved quantitative analysis of cell death (frequency of SO-positive cells) and

protoplast shrinkage upon HS of cells with normal or uncoupled mitochondria revealed that both

parameters were independent of the mitochondrial bioenergetic function (**Fig. 3E, F**). Pre-

treatment with CsA or EGTA prior to HS did not alleviate the cell death rate either (**Fig. 3G, H,**

Fig. S1C, D).

The discrepancies between our work and the previous studies could be caused by technical issues,
i.e. precision of the temperature measurement during HS. To examine this possibility, we
performed HS at 40, 45 and 50°C. Although cell viability was inversely proportional to the
temperature, the morphology of dead cells in all cases was identical to that observed at 55°C, and
the rate of cell death did not depend on mitochondrial activity (**Fig. S1E-H**).

A recent study by Distéfano et al., (2017) proposed that HS at 55°C causes ferroptosis in
Arabidopsis thaliana root hair cells. Although mitochondrial dysfunction is known to suppress
ferroptosis in animal cells (Gao et al., 2019) and, as shown above, HS-induced plant cell death is
mitochondria-independent, we still examined whether "AL-PCD" could be classified as a
ferroptosis. For this, BY-2 cells were treated with a ferroptosis inhibitor, ferrostatin-1 (Fer-1), prior
to HS at 55°C and cell death rate was measured during 24 h after the stress. Treatment with Fer-1
did not alleviate the cell death (**Fig. 4A, B;** data is shown for the first 12 h after HS). Furthermore,
two independent experiments replicating conditions that were reported to induce ferroptosis in
Arabidopsis root hair cells did not confirm such type of cell death (**Fig. 4C**). Taken together, our
results reject the notion that HS-induced "AL-PCD" is a programmed process. On the contrary,
they demonstrate that HS triggers rapid destruction of cellular components and passive decay of
plant cells, i.e. necrotic cell death.

Necrotic deaths caused by 55°C or 85°C display different cell morphologies due to a fixating effect
of higher temperature
The lack of protoplast shrinkage during cell death induced by 85°C was used to classify it as
necrosis (McCabe et al., 1997; Reape and McCabe, 2013). However, both 55°C and 85°C HS
trigger instant and irreversible permeabilization of the PM, MMP dissipation, and drop in
intracellular ATP content, i.e. hallmark features of necrosis. Plausibly, morphological differences
between necrotic cell deaths triggered by 55°C and 85°C could be explained by rapid protein
denaturation occurring at 85°C that would crosslink cellular components. Such "fixation" would
prevent protoplast shrinkage. Consistent with this suggestion, high-temperature cell fixation
protocols have been used as an alternative to chemical fixation (Login and Dvorak, 1988).

To test whether 85°C HS acts as a fixative, we induced protoplast retraction from the cell wall by
exposing stressed cell cultures to hypertonic conditions at 185 mM D-mannitol (**Fig. 5**). The high
osmotic pressure would induce dehydration and protoplast shrinkage in the non-fixed cells. As
expected, the living cells treated with D-mannitol underwent typical plasmolysis manifested by
reversible protoplast detachment from the cell wall. Cells exposed to 55°C displayed irreversible
protoplast shrinkage phenotype both with and without D-mannitol treatment. However, although
cells treated at 85°C HS exhibited visible signs of dehydration in the hypertonic solution,
protoplasts of virtually all cells remained attached to the cell wall. These data provide compelling
evidence that the fixing effect of 85°C HS prevents protoplast shrinkage. Thus, morphological
differences between 55°C and 85°C HS-induced cell deaths do not reflect differences in the cell-
death execution mechanism.
252
Conclusions
Research on plant cell death suffers from superficial extrapolation of the rich knowledge about
various types of animal cell death, and in particular *bona fide* apoptosis that is especially appealing
due to immense track record of ground-breaking discoveries. Results of our current work
schematically summarized in **Fig. 6** indicate that morphological and molecular characteristics of
plant HS-induced cell death termed in the literature "AL-PCD" *de facto* fit the definition of
necrosis. The set of assays described here can be easily implemented to various plant model
systems used for the investigation of cell death with "apoptotic-like" features (e.g., Ryerson and
Heath, 1996; Navarre and Wolpert, 1999; Houot et al., 2001; Behboodi and Samadi, 2004; Tada
et al., 2005; Lytvyn et al., 2010; Rybaczek et al., 2015; Nyalugwe et al., 2016; Ambastha et al.,
2017; Żabka et al., 2017; Araniti et al., 2018; Malerba and Cerana, 2018; Ghasemi et al., 2020).
This will hopefully bring the question about existence of plant apoptosis to rest before long. Our
study highlights the need for critical classification of plant cell death modalities (van Doorn et al.,
2011) which is crucial for understanding the evolution of eukaryotic cell death machinery.
267
Methods
Plant material and growth conditions
BY-2 cell cultures (Nagata et al., 1992) were grown at 25°C (unless stated otherwise), 120 rpm, in
the darkness. Cells were subcultured every 7 days to the standard Murashige and Skoog (MS)

medium supplemented with vitamins (M0222, Duchefa), 1.875 mM KH_2PO_4, 0.2 mg/L 2,4-D, and

3% (w/v) sucrose; pH 5.6.

For growing *Arabidopsis thaliana* seedlings, Col-0 seeds were sterilized for 30 min using a 2.7

g/L sodium hypochlorite solution (Klorin, Colgate Palmolive) with 0.05% (v/v) Tween 20. The

seeds were then rinsed three times with filter-sterilized Milli-Q water prior to being plated on

solidified medium. The MS medium was brought to pH 5.8 with 1M KOH prior to autoclaving

and contained half-strength MS salts and vitamins (M0222, Duchefa), 1% (w/v) sucrose, 10 mM

MES and 0.8% (w/v) plant agar (P1001, Duchefa). The plates were incubated vertically with

cycles of 16 h of 150 $\mu E\, m^{-2}\, s^{-1}$ light at 22°C/8 h of dark at 20°C.

Fluorescein leakage assay (fluorochromatic assay)

Cells were stained with 4 µg/mL FDA (Merck, F7378) in the culture medium for 10 min at room

temperature. Half a mL aliquots of the stained cell culture were then transferred into 2 mL

Eppendorf tubes using a cut 1-mL tip to avoid mechanical damage of the cells. Each tube was

treated as a biological replicate. At least three replicates were used for each condition in each

experiment.

The tubes were exposed to four conditions: (i) left on the bench at room temperature; incubated

for 10 min at (ii) 55°C or (iii) 85°C in a preheated thermoblock (wells were filled with water for

better thermoconductivity); (iv) snap frozen in liquid nitrogen. After treatments, cultures were let

to cool down or thaw for 5 min at room temperature and then either mounted on a sample glass to

be imaged using CLSM or processed further for quantitative assay.

For quantitative fluorescein leakage assay cells were pelleted on nylon meshes with 50 µm pores.

Supernatants were collected entirely with a pipette, 150 µL of each supernatant were transferred

to the empty wells of a 96-well flat bottom plate (Sarstedt). Cells on the mesh were then washed

off the meshes into wells of the same plate using 150 µL of fresh BY-2 media. Fluorescence was

measured in the FLUOstar Omega Microplate Reader (BMG LABTECH) using 485 nm

excitation/520 nm emission filters (gain 800, 20 flashes/well). Importantly, the measurement had

to be performed within 20 min after treatment to avoid passive diffusion of fluorescein into

extracellular space in the control samples. Leaked fluorescein intensity was calculated for each

sample as a percentage of total fluorescence intensity detected in the cells and supernatant. A one-

way ANOVA with a Dunnet's test was performed using JMP 10, and the graph was built in

Origin2019.

Sytox Orange and FM4-64 staining

One-mL aliquots of 4-5-day old BY-2 cell culture were transferred to 1.5 mL Eppendorf tubes

using a cut 1 mL tip. The final concentrations of Sytox Orange (SO; ThermoFisher, S11368) and

FM4-64 (ThermoFisher, T3166) in cell culture were 1 µM and 0.5 µM, respectively. Five different

treatments were applied to the cell cultures: (i) As a control, stained cell culture was incubated on

the lab bench for 30 min. (ii) Stained cell culture was incubated in a preheated thermoblock at

55°C for a 10-min HS. (iii) Cell culture was subjected to a 55°C HS for 10 min, and stains were

added after additional 30 min at room temperature. (iv) Stained cell culture was incubated in a

preheated thermoblock at 85°C for a 10-min HS. (v) Cell culture was subjected to a 85°C HS for

10 min, and stains were added after additional 30 min at room temperature. Cells were mounted

on a microscope slide and imaged by CLSM 30 min after staining. More than one hundred cells

were counted for each treatment to ensure robustness of the results. Note that there is some degree

of crosstalk in the fluorescence emission profiles of the SO and FM4-64 stains, which was most

prominent following 85°C HS; however this crosstalk had no influence on the results of our

experiments.

Importantly, following a 10-min pulse HS at 55°C, we observed that there are two distinct SO

staining patterns: (i) cells with collapsed protoplasts that have intense staining (arrow,

Supplementary Fig. S2A) and (ii) cells that have not collapsed with low intensity nuclear staining

(arrowhead, **Supplementary Fig. S2A**). We found that the majority of cells were SO-positive

within 10 min after the HS and that the proportion of cells with low intensity staining (i.e. those

with only SO-positive nuclei) decreased over time as more cells collapsed and displayed intense

SO staining (**Supplementary Fig. S2B**).

Nuclear phenotyping

A transgenic BY-2 line expressing GFP fused to nuclear localization signal (NLS) under a double

35S promoter (2x35::Rluc-sGFP-NLS) was used to evaluate nuclear phenotype following HS. The

transgenic culture was generated according to (Dauphinee et al., 2019). Treatments included: (i)

No HS, (ii) 10-min pulse HS at 55°C (iii) 10-min pulse HS at 55°C with SO staining as stated

above, which was used to validate results following HS (data not shown). Six hours following HS,

z-stack acquisitions were captured with Zeiss LSM 800 as described in the *Confocal microscopy.*

Quantification was performed using maximum intensity projections. The areas of nuclei were

approximated using threshold segmentation of images with Fiji software (ImageJ version 1.52r).

Three independent experiments were carried out.

MitoTracker Red Staining

BY-2 cells were stained with 100 µM MitoTracker Red (ThermoFisher, M7512) for 10 min at

room temperature. Cells were mounted on a microscope slide and imaged using Zeiss LSM 780

as described in the *Confocal microscopy.*

EGTA and cyclosporin A treatments

Half a mL of a 4-5-day old BY-2 cell culture were transferred to 1.5 mL Eppendorf tubes. The

cells were stained with Mitotracker Red (100 µM) and FDA (2 µg/mL) 10 min prior to

treatment. Four different treatments (all at room temperature) were applied prior to 10-min HS at

55°C: (i) 10 mM EGTA (Merck, E3889), pH 8.0 for 10 min, (ii) 0.1% DMSO for 2 h, (iii) 15 µM

cyclosporin A (CsA) for 2 h, and (iv) 10 mM EGTA (applied 10 min prior to HS) and 15 µM CsA

(applied 2 h prior to HS). Treatment times were staggered and samples were scanned within 10

min following HS. Cells were mounted on a microscope slide and imaged using Zeiss LSM 800.

In order to assess the cell death rate following EGTA treatment, an additional experiment was

carried out using BY-2 cells that were treated and then stained with SO and FM4-64 as described

above. There were three biological replicates for each treatment group: (i) No HS - control

(water), (ii) No HS – EGTA, (iii) 10-min 55°C HS – control, and (iv) 10-min 55°C HS - EGTA.

After HS, 300 µL of each sample was transferred to a 24-well plate designed for fluorescent

microscopy (µ-Plate 24, Ibidi 82406) and imaged using Zeiss LSM 780 as described in the

Confocal microscopy section.

ATP content measurement

The assay was performed based on the previously published protocol (Minina et al., 2013). Half a mL aliquots of a 4-day-old BY-2 cell culture were transferred with a cut tip into 2 mL Eppendorf tubes. Six replicates were measured for each treatment in each experiment. The aliquoted cells were subjected to the treatments described in the *Fluorescein leakage* assay. Additionally, six aliquots were incubated with 48 µM carbonyl CCCP on the bench for 1 h. After treatment, cells were pelleted on a nylon mesh with 50 µm pores, rinsed with 3 mL of medium and washed off into new Eppendorf tubes with 600 µL of boiling buffer (100 mM Tris-HCl, 4 mM EDTA, pH 7.5, autoclaved and supplemented with phosphatase inhibitor cocktail 2 (Sigma-Aldrich, P5726)). During sample collection, the tubes were kept on ice. Samples were then boiled at 100°C for 10 min, and debris was spun down at 4°C, $10,000g$ for 20 min. The supernatants were transferred into new tubes and stored on ice.

For each sample, 83.5 µL of supernatant was transferred into wells of white Nunc plates in three technical replicates. The background signal was detected using FLUOstar Omega Microplate Reader (BMG LABTECH), with a lens filter taking ten measurements per well. 41.5 µL of freshly prepared reaction buffer (1.5 mM DTT, 100 µM Luciferin (Sigma L 6152), 5 µg/mL Luciferase from *Phontius pyralis* (Sigma L9506)) were pipetted into each well by the automated pump in a microplate reader. Luminescence was detected for each well after adding the reaction buffer and shaking the plate to ensure good mixing in the reaction volume. A serial dilution of ATP ranging from 1.8×10^{-2} M to 10^{-8} M was used to build a standard curve and calculate the amount of ATP in each sample. A one-way ANOVA with Dunnet's test was performed using JMP 10, and the graph was built in Origin 2019.

CCCP treatment

Half a mL of 4-5-day old BY-2 cell culture were transferred to 1.5 mL Eppendorf tubes using a cut 1 mL tip. The four treatment groups included: (i) 10-min treatment with 48 µM CCCP at room temperature followed by 55°C HS for 10 min, (ii) 0.1% DMSO (vehicle control) at room temperature for 10 min followed by 55° HS for 10 min, (iii) 10-min treatment with 48 µM CCCP and no HS, and (iv) 0.1% DMSO with no HS. The cultures were stained with 1 µM SO and then

395 transferred to a 24-well plate designed for fluorescent microscopy (μ-Plate 24, Ibidi 82406).
396 Scanning of the wells was performed automatically using CLSM, 10 minutes after treatment and
397 then every 2 h for 24 h. Three replicates were carried out per treatment group.
398
399 A HS gradient experiment was also carried out, comparing control (DMSO-) and CCCP-treated
400 BY-2 cells. The cultures were stained and treated as per the CCCP experiment described above
401 and then subjected to 10-min exposure to one of the following conditions: (i) no HS, (ii) 40°C HS,
402 (iii) 45°C, (iv) 50°C, and (v) 55°C. The cultures were scanned after 6 h and 24 h using CLSM.
403 Three technical replicates were carried out per group.
404
405 *Osmotic stress assay*
406 Two hundred μL of BY-2 cell culture were transferred into Eppendorf tubes. Cells were stained
407 with FDA and exposed to HS as described in the *Fluorescein leakage assay*. 60 μL of 0.8 M D-
408 Mannitol were added to each tube (final concentration 185 mM), cells were incubated at room
409 temperature for 5 min and mounted on sample glass for CLSM imaging.
410
Ferroptosis assays
Three mL of 4-5-day old BY-2 cell culture were transferred to a 6-well plate using a cut 1 mL tip.
Five different treatments were applied to the cell cultures for 16 h during which time they were
returned to the incubator and grown as described above: (i) 0.1% DMSO (vehicle control), (ii) 1
μM Ferostatin-1 (Fer-1, Sigma SML0583), (iii) 10 μM Fer-1, (iv) 50 μM Fer-1 and (v) 100 μM
Fer-1. After the 16-h treatments, 0.5 mL of culture from each well were transferred to 1.5 mL
Eppendorf tubes using a cut 1 mL tip and separated into three treatments: (i) no HS, (ii) 55° HS
for 10 min, or (iii) 85° HS for 10 min as described above. Following treatment, 10 μL of culture
were transferred to 490 μL of fresh BY-2 media and stained with 1 μM SO and 0.5 μM FM4-64
in 24-well plates designed for fluorescent microscopy (μ-Plate 24, Ibidi 82406). Three technical
replicates were carried out for each group and the wells were scanned automatically using CLSM
at the following intervals: 10 min, 3, 6, 9, 12, 15, 18, 21 and 24 h. Three independent experiments
were carried out.
424

The effect of Fer-1 was also tested on *Arabidopsis thaliana* seedlings which were grown as
described above. Six-day old seedlings were gently transferred to 6-well plates containing 3 mL
of liquid half-strength MS medium and given one of the following three treatments for 16 h: (i)
0.1% DMSO (vehicle control), (ii) 1 µM Fer-1, and (iii) 10 µM Fer-1. The treated medium was
pipetted gently over the seedlings to ensure that roots were submerged in the liquid prior to being
placed back into the growth cabinet overnight. The following morning, the seedlings were
carefully transferred to 1.5 mL Eppendorf tubes containing the same treatment media from the 6-
well plates before subjection to one of the following three treatments: (i) no HS, (ii) 55° HS for 10
min, or (iii) 85° HS for 10 min as described above. The seedlings were then stained with SO and
FM4-64 as described above and placed back into the growth cabinet until scanning with CLSM at
3 h and 6 h post-HS. The early differentiation zone of the root was scanned for three seedlings per
time point, and three independent experiments were carried out.

Confocal microscopy
Micrographs were acquired using either a LSM 800 or LSM 780 confocal laser scanning
microscope (Carl Zeiss) with GaAsP detectors. Micrographs were taken with four different
objectives: x10 (NA0.45), x20 (NA0.8), x40 (NA1.2, water immersion), and x63 (NA1.2, water
immersion). For Fig. 1A-D and supplementary video S1, z-stack acquisition with sequential
scanning was performed with a x63 objective. Nuclear phenotyping using the BY-2 transgenic line
(Fig. 1E) was done with z-stack acquisitions using a x20 objective. Fluorescein was excited at 488
nm and emission was detected from 499 – 560 nm. MitoTracker red was excited at 561 nm and
emission was detected from 582 – 754 nm. SO was excited at 561 nm and emission was detected
from 410 – 605 nm. FM4-64 excitation was 506 nm, the emission was detected from 650 – 700
nm. Images were acquired using ZEN blue software (version 2.5, Carl Zeiss) or Zen black (version
2.3).

Acknowledgements
This project was supported by grants from Carl Tryggers Foundation (to EAM), MSCA IF (to
EAM), the Swedish Research Councils VR (to PVB) and Formas (to AND and PVB), the Knut
and Alice Wallenberg Foundation (to PVB), the Swedish Foundation for Strategic Research (to
PVB), the Natural Sciences and Engineering Research Council (NSERC) of Canada (to AND) and

by the research programme "Crops for the Future" at the Swedish University of Agricultural

Sciences. APS is grateful to August T. Larsson Guest Researcher Programme for supporting his

visits to the Swedish University of Agricultural Sciences. VG was supported by the Russian

Science Foundation (grant 19-14-00122).

Competing Interests statement

The authors declare no conflicts of interest.

References

Ambastha, V., Sopory, S.K., Tiwari, B.S., and Tripathy, B.C. (2017). Photo-modulation of

programmed cell death in rice leaves triggered by salinity. Apoptosis **22**: 41–56.

Ameisen, J.C. (2002). On the origin, evolution, and nature of programmed cell death: a timeline

of four billion years. Cell Death Differ. **9**: 367–393.

**Ankarcrona, M., Dypbukt, J.M., Bonfoco, E., Zhivotovsky, B., Orrenius, S., Lipton, S.A.,

and Nicotera, P.** (1995). Glutamate-induced neuronal death: a succession of necrosis or

apoptosis depending on mitochondrial function. Neuron **15**: 961–973.

Arandjelovic, S. and Ravichandran, K.S. (2015). Phagocytosis of apoptotic cells in

homeostasis. Nat. Immunol. **16**: 907–917.

**Araniti, F., Costas-Gil, A., Cabeiras-Freijanes, L., Lupini, A., Sunseri, F., Reigosa, M.J.,

Abenavoli, M.R., and Sánchez-Moreiras, A.M.** (2018). Rosmarinic acid induces

programmed cell death in Arabidopsis seedlings through reactive oxygen species and

mitochondrial dysfunction. PLoS One **13**: e0208802.

Atkin-Smith, G.K. and Poon, I.K.H. (2017). Disassembly of the Dying: Mechanisms and

Functions. Trends Cell Biol. **27**: 151–162.

Balk, J., Leaver, C.J., and McCabe, P.F. (1999). Translocation of cytochrome c from the

mitochondria to the cytosol occurs during heat-induced programmed cell death in cucumber

plants. FEBS Lett. **463**: 151–154.

Behboodi, B.S. and Samadi, L. (2004). Detection of apoptotic bodies and oligonucleosomal

DNA fragments in cadmium-treated root apical cells of Allium cepa Linnaeus. Plant Sci.
167: 411–416.

Bortner, C.D. and Cidlowski, J.A. (2002). Apoptotic volume decrease and the incredible
shrinking cell. Cell Death Differ. **9**: 1307–1310.

Bortner, C.D. and Cidlowski, J.A. (2007). Cell shrinkage and monovalent cation fluxes: Role
in apoptosis. Arch. Biochem. Biophys. **462**: 176–188.

Bozhkov, P. V and Lam, E. (2011). Green death: revealing programmed cell death in plants.
Cell Death Differ. **18**: 1239–1240.

Coll, N.S., Epple, P., and Dangl, J.L. (2011). Programmed cell death in the plant immune
system. Cell Death Differ. **18**: 1247–1256.

Daneva, A., Gao, Z., Van Durme, M., and Nowack, M.K. (2016). Functions and Regulation of
Programmed Cell Death in Plant Development. Annu. Rev. Cell Dev. Biol. **32**: 441—468.

Dauphinee, A.N., Cardoso, C., Dalman, K., Ohlsson, J.A., Fick, S.B., Robert, S., Hicks,
G.R., Bozhkov, P. V., and Minina, E.A. (2019). Chemical screening pipeline for
identification of specific plant autophagy modulators. Plant Physiol. **181**: 855–866.

Dickman, M., Williams, B., Li, Y., Figueiredo, P., and Wolpert, T. (2017). Reassessing
apoptosis in plants. Nat. plants **3**: 773–779.

Distéfano, A.M. et al. (2017). Heat stress induces ferroptosis-like cell death in plants. J. Cell
Biol. **216**: 463–476.

Dixon, S.J. and Stockwell, B.R. (2019). The Hallmarks of Ferroptosis. Annu. Rev. Cancer Biol.
3: 35–54.

van Doorn, W.G. et al. (2011). Morphological classification of plant cell deaths. Cell Death
Differ. **18**: 1241–1246.

Drost, H.-G., Janitza, P., Grosse, I., and Quint, M. (2017). Cross-kingdom comparison of the
developmental hourglass. Curr. Opin. Genet. Dev. **45**: 69–75.

Edlich, F. (2018). BCL-2 proteins and apoptosis: Recent insights and unknowns. Biochem.
Biophys. Res. Commun. **500**: 26–34.

Galluzzi, L. et al. (2015). Essential versus accessory aspects of cell death: recommendations of
the NCCD 2015. Cell Death Differ. **22**: 58–73.

Galluzzi, L. et al. (2012). Molecular definitions of cell death subroutines: recommendations of
the Nomenclature Committee on Cell Death 2012. Cell Death Differ. **19**: 107–120.

Galluzzi, L. et al. (2018). Molecular mechanisms of cell death: recommendations of the
Nomenclature Committee on Cell Death 2018. Cell Death Differ. **25**: 486–541.

Gao, M., Yi, J., Zhu, J., Minikes, A.M., Monian, P., Thompson, C.B., and Jiang, X. (2019).
Role of Mitochondria in Ferroptosis. Mol. Cell **73**: 354-363.e3.

Ghasemi, R., Sharifi, R., and Ghaderian, S.M. (2020). Studying the roles of calcium and
magnesium in cell death in the serpentine native plant Alyssum inflatum NYÁRÁDY
through cell suspension culture technique. Plant Physiol. Biochem. PPB **151**: 362–368.

Green, D.R. and Fitzgerald, P. (2016). Just So Stories about the Evolution of Apoptosis. Curr.
Biol. **26**: R620–R627.

Hengartner, M.O. (2000). The biochemistry of apoptosis. Nature **407**: 770–776.

Hogg, B. V, Kacprzyk, J., Molony, E.M., O'Reilly, C., Gallagher, T.F., Gallois, P., and
McCabe, P.F. (2011). An in vivo root hair assay for determining rates of apoptotic-like
programmed cell death in plants. Plant Methods **7**: 45.

Houot, V., Etienne, P., Petitot, A.S., Barbier, S., Blein, J.P., and Suty, L. (2001). Hydrogen
peroxide induces programmed cell death features in cultured tobacco BY-2 cells, in a dose-
dependent manner. J. Exp. Bot. **52**: 1721–1730.

Hunter, D.R. and Haworth, R.A. (1979). The Ca2+-induced membrane transition in
mitochondria: I. The protective mechanisms. Arch. Biochem. Biophys. **195**: 453–459.

Huysmans, M., Lema A, S., Coll, N.S., and Nowack, M.K. (2017). Dying two deaths -
programmed cell death regulation in development and disease. Curr. Opin. Plant Biol. **35**:
37—44.

Jacobson, M.D., Weil, M., and Raff, M.C. (1997). Programmed Cell Death in Animal
Development. Cell **88**: 347–354.

Jorgensen, I., Rayamajhi, M., and Miao, E.A. (2017). Programmed cell death as a defence
against infection. Nat. Rev. Immunol. **17**: 151–164.

Kabbage, M., Kessens, R., Bartholomay, L.C., and Williams, B. (2017). The Life and Death
of a Plant Cell. Annu. Rev. Plant Biol. **68**: 375–404.

Kacprzyk, J., Brogan, N.P., Daly, C.T., Doyle, S.M., Diamond, M., Molony, E.M., and
McCabe, P.F. (2017). The retraction of the protoplast during PCD is an active, and
interruptible, calcium-flux driven process. Plant Sci. **260**: 50–59.

Kerr, J.F., Wyllie, A.H., and Currie, A.R. (1972). Apoptosis: a basic biological phenomenon
with wide-ranging implications in tissue kinetics. Br. J. Cancer **26**: 239–257.

Kristián, T. and Siesjö, B.K. (1998). Calcium in Ischemic Cell Death. Stroke **29**: 705–718.

Kroemer, G. et al. (2005). Classification of cell death: recommendations of the Nomenclature
Committee on Cell Death. Cell Death Differ. **12 Suppl 2**: 1463–1467.

Lacomme, C. and Santa Cruz, S. (1999). Bax-induced cell death in tobacco is similar to the
hypersensitive response. Proc. Natl. Acad. Sci. U. S. A. **96**: 7956–7961.

Letai, A. (2017). Apoptosis and Cancer. Annu. Rev. Cancer Biol. **1**: 275–294.

Login, G.R. and Dvorak, A.M. (1988). Microwave fixation provides excellent preservation of
tissue, cells and antigens for light and electron microscopy. Histochem. J. **20**: 373–387.

Lytvyn, D.I., Yemets, A.I., and Blume, Y.B. (2010). UV-B overexposure induces programmed
cell death in a BY-2 tobacco cell line. Environ. Exp. Bot. **68**: 51–57.

Majno, G. and Joris, I. (1995). Apoptosis, oncosis, and necrosis. An overview of cell death.
Am. J. Pathol. **146**: 3–15.

Malerba, M. and Cerana, R. (2018). Role of peroxynitrite in the responses induced by heat
stress in tobacco BY-2 cultured cells. Protoplasma **255**: 1079–1087.

Martínez-Fábregas, J., Díaz-Moreno, I., González-Arzola, K., Janocha, S., Navarro, J.A.,
Hervás, M., Bernhardt, R., Díaz-Quintana, A., and La Rosa, M.Á. (2013). New
Arabidopsis thaliana cytochrome c partners: a look into the elusive role of cytochrome c in
programmed cell death in plants. Mol. Cell. Proteomics **12**: 3666–3676.

McCabe, P.F., Levine, A., Meijer, P.-J., Tapon, N.A., and Pennell, R.I. (1997). A
programmed cell death pathway activated in carrot cells cultured at low cell density. Plant J.
12: 267–280.

Miao, E.A., Leaf, I.A., Treuting, P.M., Mao, D.P., Dors, M., Sarkar, A., Warren, S.E.,
Wewers, M.D., and Aderem, A. (2010). Caspase-1-induced pyroptosis is an innate
immune effector mechanism against intracellular bacteria. Nat. Immunol. **11**: 1136–1142.

Minina, E.A. et al. (2020). Classification and Nomenclature of Metacaspases and Paracaspases:
No More Confusion with Caspases. Mol. Cell **77**: 927–929.

Minina, E.A., Filonova, L.H., Sanchez-Vera, V., Suarez, M.F., Daniel, G., and Bozhkov, P.
V. (2013). Detection and Measurement of Necrosis in Plants. In K. McCall and C. Klein,
eds (Humana Press: Totowa, NJ), pp. 229–248.

Mitsuhara, I., Malik, K.A., Miura, M., and Ohashi, Y. (1999). Animal cell-death suppressors
Bcl-xL and Ced-9 inhibit cell death in tobacco plants. Curr. Biol. **9**: 775-S1.

Molnár, T., Mázló, A., Tslaf, V., Szöllősi, A.G., Emri, G., and Koncz, G. (2019). Current
translational potential and underlying molecular mechanisms of necroptosis. Cell Death Dis.
10: 860.

Murat, F., van de Peer, Y., and Salse, J. (2012). Decoding plant and animal genome plasticity
from differential paleo-evolutionary patterns and processes. Genome Biol. Evol. **4**: 917–
928.

Nagata, T., Nemoto, Y., and Hasezawa, S. (1992). Tobacco BY-2 Cell Line as the "HeLa" Cell
in the Cell Biology of Higher Plants. In International review of cytology. A survey of cell
biology : Volume 132, K.W. Jeon and M. Friedlander, eds, International Review of
Cytology TS - CrossRef. (Elsevier: London), pp. 1–30.

Navarre, D.A. and Wolpert, T.J. (1999). Victorin induction of an apoptotic/senescence-like
response in oats. Plant Cell **11**: 237–249.

Nyalugwe, E.P., Barbetti, M.J., Clode, P.L., and Jones, R.A.C. (2016). Programmed cell
death pathways induced by early plant-virus infection are determined by isolate virulence
and stage of infection. Plant Pathol. **65**: 1518–1528.

Perry, S.W., Norman, J.P., Barbieri, J., Brown, E.B., and Gelbard, H.A. (2011).
Mitochondrial membrane potential probes and the proton gradient: a practical usage guide.
Biotechniques **50**: 98–115.

Reape, T.J. and McCabe, P.F. (2013). Commentary: the cellular condensation of dying plant
cells: programmed retraction or necrotic collapse? Plant Sci. **207**: 135–139.

Rotman, B. and Papermaster, B.W. (1966). Membrane properties of living mammalian cells as
studied by enzymatic hydrolysis of fluorogenic esters. Proc. Natl. Acad. Sci. **55**: 134.

Rybaczek, D., Musiałek, M.W., and Balcerczyk, A. (2015). Caffeine-Induced Premature
Chromosome Condensation Results in the Apoptosis-Like Programmed Cell Death in Root
Meristems of Vicia faba. PLoS One **10**: e0142307.

Ryerson, D.E. and Heath, M.C. (1996). Cleavage of Nuclear DNA into Oligonucleosomal
Fragments during Cell Death Induced by Fungal Infection or by Abiotic Treatments. Plant
Cell **8**: 393–402.

Salvesen, G.S., Hempel, A., and Coll, N.S. (2016). Protease signaling in animal and plant-
regulated cell death. FEBS J. **283**: 2577–2598.

Segawa, K. and Nagata, S. (2015). An Apoptotic "Eat Me" Signal: Phosphatidylserine
Exposure. Trends Cell Biol. **25**: 639–650.

Sperandio, S., Belle, I., and Bredesen, D.E. (2000). An alternative, nonapoptotic form of
programmed cell death. Proc. Natl. Acad. Sci. U. S. A. **97**: 14376–14381.

Suzanne, M. and Steller, H. (2013). Shaping organisms with apoptosis. Cell Death Differ. **20**:
669–675.

Tada, Y., Kusaka, K., Betsuyaku, S., Shinogi, T., Sakamoto, M., Ohura, Y., Hata, S., Mori,
T., Tosa, Y., and Mayama, S. (2005). Victorin triggers programmed cell death and the
defense response via interaction with a cell surface mediator. Plant Cell Physiol. **46**: 1787–
1798.

Tsujimoto, Y. (1997). Apoptosis and necrosis: intracellular ATP level as a determinant for cell
death modes. Cell Death Differ. **4**: 429–434.

Vacca, R.A., Pinto, M.C., Valenti, D., Passarella, S., Marra, E., and Gara, L. (2004).
Production of reactive oxygen species, alteration of cytosolic ascorbate peroxidase, and
impairment of mitochondrial metabolism are early events in heat shock-induced
programmed cell death in tobacco Bright-Yellow 2 cells. Plant Physiol. **134**: 1100–1112.

Valandro, F., Menguer, P.K., Cabreira-Cagliari, C., Margis-Pinheiro, M., and Cagliari, A.
(2020). Programmed cell death (PCD) control in plants: New insights from the Arabidopsis
thaliana deathosome. Plant Sci. **299**: 110603.

Yu, X.-H., Perdue, T.D., Heimer, Y.M., and Jones, A.M. (2002). Mitochondrial involvement
in tracheary element programmed cell death. Cell Death Differ. **9**: 189–198.

Żabka, A., Winnicki, K., Polit, J.T., and Maszewski, J. (2017). Sanguinarine-induced
oxidative stress and apoptosis-like programmed cell death (AL-PCD) in root meristem cells
of Allium cepa. Plant Physiol. Biochem. PPB **112**: 193—206.

Zamaraeva, M. V, Sabirov, R.Z., Maeno, E., Ando-Akatsuka, Y., Bessonova, S. V, and
Okada, Y. (2005). Cells die with increased cytosolic ATP during apoptosis: a
bioluminescence study with intracellular luciferase. Cell Death Differ. **12**: 1390–1397.

Zhang, Y., Chen, X., Gueydan, C., and Han, J. (2018). Plasma membrane changes during
programmed cell deaths. Cell Res. **28**: 9–21.

Zong, W.-X. and Thompson, C.B. (2006). Necrotic death as a cell fate. Genes Dev. **20**: 1–15.

Figure Legends

Figure 1. Gross morphological changes in HS-treated plant cells do not match hallmarks of apoptosis.

A FM4-64 and Sytox Orange (SO) dyes were used to visualize all cells and cells with permeabilized plasma membrane (PM), respectively. Stained BY-2 cells were imaged under control conditions (No HS) or after a 10-min HS at 55°C. Scanning was performed within 1 h post-HS. The arrows indicate PM; note the PM (red dotted line) being tightly pressed to the cell wall (white dotted line) under control conditions and detached under stress conditions. **B** Higher magnification of the areas indicated with arrows in **A**. **C** BY-2 cells expressing GFP fused to nuclear localization signal (NLS) were subjected to the same treatments as in **A**. Images represent maximum intensity projections of z-stack scans acquired 6 h post-HS. **D** Quantification of nuclear area in samples shown in **C**. Data from three independent experiments, with $\geq$ 142 cells counted per treatment. Student's t-test, *p < 0.005. DIC, differential interference contrast microscopy. n, nucleus. IQR, interquartile range. Scale bars, 20 µm (**A, C**) or 5 µm (**B**).

Figure 2. Both 55°C and 85°C HS cause instant and irreversible PM permeabilization.

A Green fluorescence in the BY-2 cells loaded with FDA. Fluorescein leaks into extracellular space within 10 min after HS at either 55°C or 85°C. Arrows indicate shrunken protoplasts in 55°C HS-treated cells. **B** Similar fraction of fluorescein leaks out of cells after 10 min of 55°C, 85°C, or liquid nitrogen (N_2) treatment. **C** BY-2 cells stained with SO and FM4-64. To assess whether disruption of the PM integrity after pulse HS is transient or irreversible, SO staining was performed before HS or 1 h after HS. **D** Frequency of the SO-positive cells in the cultures shown in **C**. DIC, differential interference contrast microscopy. IQR, interquartile range. Scale bars in **A** and **C**, 50 µm. **B**, representative data from one out of three independent experiments. **D**, data from three independent experiments, with $\geq$ 115 cells counted per treatment. **B** and **D**, one-way ANOVA with Dunnet's test; *, p<0.005.

Figure 3. Protoplast shrinkage at 55°C HS is an ATP- and Ca^{2+}-independent process.

A Mitochondria in BY-2 cells stained with MitoTracker Red and imaged 10 min after 55°C or 85°C HS, or after treatment with 48 µM CCCP. Severely damaged mitochondria were observed

upon all three treatments. **B** Loss of intracellular ATP content upon HS. Snap freeze-thaw
treatment in liquid nitrogen (N_2) and CCCP treatment were used as positive controls for completely
disrupted and uncoupled mitochondria, respectively. The experiment was repeated twice, each
time using four biological replicates per treatment. **C** MitoTracker Red staining of BY-2 cells
exposed to 55°C in the presence or absence of 15 µM Cyclosporin A (CsA) reveals that inhibition
of MPTP opening does not rescue mitochondria from severe damage and loss of membrane
potential caused by HS. **D** MitoTracker Red localization in the cells pre-treated with 10 mM EGTA
prior to the HS reveals that chelation of extracellular Ca^{2+} does not rescue mitochondrial
phenotype. **E, F** Dynamics of cell death (% SO-positive cells; **E**) and protoplast shrinkage **(F)** in
cells with normal and uncoupled (48 µM CCCP treatment) mitochondria. **G, H** Pre-treatment with
10 mM EGTA before HS does not affect dynamics of cell death (% SO-positive cells; **G**) and
protoplast shrinkage **(H)**. Experiments shown in **E-H** were repeated three times, with ≥ 184 cells
per treatment and time point. Each microscopy experiment was performed at least twice. Scale
bars, 20 µm (**A**) or 50 µm (**C, D**). IQR, interquartile range. **B**, **E**-**H**, one-way ANOVA with
Dunnet's test; *, p<0.05.

Figure 4. HS-induced cell death response is not ferroptosis.
A Sytox Orange (SO) and FM4-64 staining of BY-2 cultures demonstrates that pre-treatment with
1 µM Ferrostatin-1 (Fer-1) does not affect 55°C HS-induced cell death. **B** Quantification of cell
death (% SO-positive cells) in the samples illustrated in **A**. The chart shows representative results
of three independent experiments, each including ≥ 280 cells per treatment and time point. **C** Pre-
treatment with 1 µM Fer-1 does not alleviate the HS-induced death of *Arabidopsis thaliana* root
hair cells. Arrows indicate SO-positive nuclei. Three independent experiments demonstrated the
same results. No quantification was performed in these experiments, since all cells were SO-
positive. DIC, differential interference contrast microscopy. Scale bars, 40 µm

Figure 5. 85°C HS prevents protoplast shrinkage in necrotic cells by fixing cellular content
Cells stained with FDA were exposed to 55°C or 85°C and mounted in the normal growth medium
or hypertonic medium supplemented with 185 mM D-mannitol. High osmotic pressure induced
plasmolysis in the non-stressed cells (no HS, arrows). Protoplasts of the cells treated at 55°C or
85°C only showed signs of dehydration in the hypertonic medium and no additional changes. The

frequency of protoplast detachment in the samples treated with 85°C remained close to zero even

under high osmotic pressure, indicating that cells were fixed by the high-temperature treatment.

White dotted line, cell wall; red dotted line, plasma membrane. Scale bars, 50 µm.

Figure 6. HS-induced plant cell death is necrotic rather than apoptotic process.

Plasma membrane (PM) integrity is maintained during apoptosis to prevent spillage of toxic dying

cell contents. However, plant cell undergoing the so called "AL-PCD" exhibit early irreversible

permeabilization of PM, a feature typical for uncontrolled cell decay, necrosis. Apoptosis is

associated with step-wise massive re-organization of the cellular morphology (nuclear

segmentation, PM blebbing, fragmentation into apoptotic bodies) and thus is an energy consuming

process that relies on controlled release of cytochrome c, functional mitochondria and intracellular

ATP. On the contrary, necrotic death is a passive lysis of the cellular contents associated with low

intracellular ATP and decoupled mitochondria. Similarly to necrotic animal cells, plant cells

undergoing "AL-PCD" have an extremely low intracellular ATP content and uncoupled

mitochondria. While apoptotic cells condense and disassemble into apoptotic bodies that are

engulfed by phagocytes, a necrotic cell typically undergoes swelling that is followed by a decrease

of the cell volume and leakage of the dead cell contents. Presence of the rigid cell wall most

probably does not permit necrosis-like swelling of plant cells during "AL-PCD", while rupture of

the vacuole coinciding with permeabilization of the PM would lead to rapid leakage of the cellular

contents and thus decrease in volume. Such shrinkage has very little in common with AVD, as it

is a passive process not associated with the formation of apoptotic bodies.

Supplementary Figure and Video Legends

Figure S1. HS-induced cell death is ATP- and Ca^{2+} -independent process.

A-D Morphology of FDA-stained cells under normal conditions (no HS) and after a 55°C HS.

Protoplast shrinkage is denoted by arrows. Pre-treatment with 48 µM CCCP for 10 min (**B**), 15

µM CsA for 2 h (**C**), or 10 mM EGTA for 10 min (**D**) did not alleviate protoplast shrinkage upon

HS. Each treatment was repeated at least twice. **E, G** Sytox Orange (SO) staining of BY-2 cells

heat-shocked for 10 min at 40, 45, or 50°C and imaged after 6 h (**E**) and 24 h (**G**). Pre-treatment

with CCCP provided no protection against cell death and protoplast shrinkage at any of the tested

HS temperatures. **F, H** Quantification of cell death (% SO-positive cells) in the samples shown in

E and **G,** respectively. DIC, differential interference contrast microscopy. IQR, interquartile range.

Experiments shown in **F** and **H** were repeated three times, with ≥ 170 cells per treatment and time

point. The data was subjected to one-way ANOVA with Bonferroni correction. Scale bars, 20 µm

(A-D) or 100 µm **(E, G).**

Figure S2. Different SO staining intensities following 55°C HS.

A Different intensity of SO staining in BY-2 cells exposed to 55°C HS for 10 min. Cells with

shrunken protoplast (arrow) are brightly stained with SO whereas cells with normal-looking

protoplasts have low-intensity nuclear staining (arrowhead). DIC, differential interference contrast

microscopy. Scale bars, 50 µm. **B** Quantification of SO-positive cells over time. Most cells are

SO-positive within 10 min of HS (0 h) and the proportion of cells with low SO intensity

(arrowhead, **A**) decreases over time as more cells collapse and show intense staining (arrow, **A**).

The experiment comprises three technical replicates, each including ≥ 180 cells per treatment and

time point.

Supplementary Video S1. Shrinkage of plant protoplast caused by 55°C HS is not followed by

fragmentation into apoptotic bodies. 3D reconstructions (maximum intensity projections) of BY-

2 cells stained with FM4-64 to visualize the PM. Scanning was performed within 1 h post-HS. The

arrows indicate membrane structures, which were not observed in the non-stressed (no HS) cells.

These structures are localized on the inner side of the PM and do not separate from the cell at the

later stages of cell death. Scale bars, 20 µm.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

Figure S2