Road to ruin? Urbanization increases herbivory in a tropical wildflower but does not

lead to the evolution of chemical defence traits.

Abstract

Urbanization is associated with numerous changes to the biotic and abiotic environment,

many of which degrade the environment and lead to a loss of biodiversity. Cities often have

elevated pollution levels that harm wildlife; however, the increased concentration of some

pollutants can fertilize urban plants, leading to corresponding positive effects on herbivore

populations. Increases in herbivory rates may lead to natural selection for greater defence

phenotypes in plants. However, evidence supporting increased herbivory leading to the

evolution of plant defence in urban environments is contradictory, and entirely absent from

tropical regions of the world. To address these research gaps, we evaluated herbivory on

Turnera subulata, a common urban wildflower, along an urbanization gradient in Joao

Pessoa, Brazil. We predicted that higher rates of herbivory in urban areas would lead these

populations to evolve cyanogenesis, a chemical defence found in a closely related *Turnera*

species. We assessed herbivory and screened for cyanogenesis in 32 populations along the

urbanization gradient, quantified by the Human Footprint Index. Our results show that

urbanization is significantly associated with increased herbivory rates in *T. subulata*

populations. Despite elevated herbivory, we found no evidence for the evolution of

cyanongenesis in any of the populations, suggesting that the fitness effects of leaf herbivory

are not extreme enough to select for the evolution of plant defence in these populations.

Habitat loss, predator release, and nutrient enrichment likely act together to increase the

abundance of herbivorous arthropods, influencing the herbivory patterns observed in our

study.

Key words: plant defence, urban ecology, cyanogenesis, insect herbivory.

Introduction

Urbanization is associated with numerous changes to the biotic and abiotic environment, which can have large impacts on urban biodiversity (Grimm et al., 2008; Fenoglio et al., 2020). Cities often exhibit increased pollution levels that may be harmful to urban wildlife (i.e., increased cancer risk (Sepp et al. 2019)). However, studies have indicated that increased levels of certain pollutants in the urban environment can positively benefit some species. For instance, elevated nitrogen oxides (NOx) from vehicle emissions can increase the nitrogen content and the nutritional quality of the foliage of certain species of plants (Bell et al. 2011; Honor et al, 2009; Spencer et al. 1988). This may in turn favour increased abundance of some herbivorous arthropods in urban environments (Dale & Frank, 2018; Raupp et al. 2010). Arthropod abundance in cities can also be affected through changes to their community structure (reviewed in Miles et al. 2019). Urbanization can lead to a decline in arthropod predators through habitat fragmentation and degradation, which offers a release from predation from their insect prey (Faeth et al 2005). As a result, the abundance of herbivorous arthropod species may be greater in cities relative to natural habitats.

Increased herbivorous species in cities means greater herbivory pressure on urban plants, which can compromise the growth, reproduction, and survival of plants (Mauricio & Rausher, 1997; Crawley, 1989). If this pressure sufficiently reduces plant fitness, there may have altered patterns of evolution of the defence mechanisms in urban plants (Hanley et al. 2007; Mauricio & Rausher, 1997; Miles et al., 2019; Johnson et al. 2018; Mauricio & Rausher, 1997; Züst et al. 2012). For instance, invasive wild parsnip rapidly reacquired its chemical defences when its specialist herbivore was introduced into its range (Zangerl and Berenbaum 2005; Zangerl et al. 2008). Sufficient herbivory pressure along urbanization gradients may trigger the evolution of chemical defence mechanisms in urban plant

populations (Miles et al., 2019; Raupp et al. 2010), however little work has paired herbivory

rates with the evolution of novel defence traits along urban gradients.

Studies that have estimated herbivory rates along urbanization gradients have shown

that there is considerable spatial variation in herbivory across the urbanization gradient.

However, studies differ in the results on herbivory rates between urban and rural areas; some

suggest an increase in herbivory with increased urbanization (Cuevas-Reyes et al. 2013, Dale

& Frank, 2017; Turrini et al. 2016) while others indicate a reduction in herbivory (Kozlov et

al. 2017; Moreira et al. 2019). In addition, most studies evaluating gradients in herbivory are

limited to cities in temperate regions (Miles et al. 2019). To our knowledge, the only study

carried out in a tropical region found increased herbivory rates of *Solanum lycorcapurm* in

habitats with greater levels urbanization (Cuevas-Reyes et al. 2013). Recent reviews on the

topic (Miles et al. 2019; Rivkin et al. 2019) emphasize the urgent need for studies in tropical

areas.

We assessed the effects of urbanization on herbivory and the evolution of

antiherbivore defence in *Turnera subulata* in Joao Pessoa, Brazil. This species is a good

system to test the effects of urbanization on herbivory and defence for several reasons.

Turnera subulata is native to Central and South America (Schlindwein and Medeiros, 2006),

and it thrives in urban habitats despite elevated pollution and habitat fragmentation.

Additionally, the close relative, *T. ulmifolia*, produces the antiherbivore defence chemical

hydrogen cyanide (HCN; Schappert & Shore, 1995). In cyanogenic plants, HCN is released

following tissue damage, and is toxic to insect herbivores (Hughes, 1991). Cyanogenesis

most likely evolved in *T. ulmifolia* if HCN as a mechanism to reduce their susceptibility to

herbivores (Schappert & Shore, 1999). Although *T. subulata* has not yet been documented to

express cyanogenesis (Shore & Obrist 1992), it is possible that this trait may evolve under

elevated herbivory pressure. Here, we sought to answer the two questions: (1) Is there an

effect of urbanization on herbivory rates on *T. subulata*? And (2) has HCN expression

evolved in urban populations in response to increased herbivory pressure? Addressing these

questions will help to fill the gaps in the field surrounding the effects of urbanization on

species interactions in the tropics.

Methods

Study system

Turnera subulata (Passifloraceae, common names: white buttercup, sulphur alder) is a

perennial herb with distylic flowers, native to Brazil and common in cities and agricultural

habitats (Fig. 1). In urban areas, *T. subulata* is typically found in dense populations alongside

pavement, lawns, and in dry fields. Individuals reproduce via outcrossing and is an important

nectar resource for many species of bees (Schlindwein & Medeiros, 2006). *Turnera subulata*

is depredated by arthropod herbivores from different feeding guilds (Cruz et al. 2019;

Schappert & Shore, 1999). Most leaf herbivory is caused 23 herbivore species, although the

most prevalent herbivores are lace bugs (*Gargaphia sp.*, Tingidae), *Euptoieta hegesia*

caterpillars, Coleoptora in the genii *Disonycha* and *Parchicola*, aphids, and leaf miners

(Schappert & Shore, 1999).

Sampling design

A total of 32 sites were selected in December 2018, along an east-west transect spanning a

gradient of urbanization in the city of Joao Pessoa (>800,000 people) capital of Paraiba,

north-eastern Brazil (Fig. 1). We considered a site a viable population if they consist of at

least 16 *T. subulata* plants within a 100 m radius of the centre of the population. Each site in

the city was at least 400 m away from the next and in the rural habitats they were at least one

kilometre apart (see Fig. 1). We collected 20 plants per population, except in sites which had

fewer than 20 plants ($N = 8$ sites). To avoid sampling close relatives, we sampled individual

plants spaced at least 2 m apart, except in sites with fewer than 20 individuals in which case

we sampled all of the plants.

We measured herbivory rate by calculating the percent area of each leaf that was

damaged by herbivory. We selected the largest four leaves on up to three ramets per plant

(N= 12 leaves per plant) and measured the amount of damage on each using the software

application BioLeaf (Machado et al. 2016). This method is an effective and provide reliable

measurements of leaf damage (see details in Machado et al. 2016). We calculated the per

plant herbivory rate by averaging the percent leaf damage across all leaves. We also recorded

whether the leaves had experienced herbivory from sap-suckling lace bugs (*Gargaphia sp.*),

which discolour the leaves (Fig. S1; Guidoti et al. 2015).

We quantified the degree of urbanization at each site using the Human Footprint

Index (HFI) from the GIS package generated by Venter et al. (2016a). The HFI combines

human land-use variables such as increasing impervious surface and built-up areas with

human demographic data to provide standardized measure of the cumulative human footprint

on the terrestrial landscape (Venter et al 2016a). We used the freely available, open-source

GIS software QGIS (v.3.16) to extract HFI values at 1 km^2 resolution from 2009 HFI

GeoTIFF dataset provided in Venter et al (2016b). We provide detailed methods on how to

extract HFI values using QGIS in the supplemental materials.

Cyanogenesis assays

We screened each plant for the presence of HCN using the Feigl-Anger assay, which use a

color change reaction to indicate the presence (blue) or absence (white) of HCN (Shore &

Obrist, 1992). We collected two to three young leaves from each plant, froze them at -20 ^{o}C

for at least 48 hours to lyse the cell wall and induce the formation of HCN. We then followed

the same protocol used by Thompson et al. (2016) to determine the frequency of cyanogenic

plants in each population.

Fig. 1. Locations of the sampled *Turnera subulata* population in Joao Pessoa in NE Brazil,

demarcated by yellow (urban) and black (rural) circles. The bottom left image shows a

typical urban population of *T. subulata*.

Statistical analysis

We tested for fine-scale patterns in herbivory along the urbanization gradient by running two

mixed effects regression models using R version 4.0.2 (R Core Team 2018). The first model

tested for differences in the percent herbivory rate experienced by each plant averaged across

all sampled leaves on a plant using a linear mixed effect model (*lmer* from the lme4 v. 1.1-23

package (Bates et al. 2015)). The second model tested for differences in the presence or

absence of herbivory experienced by a plant using a generalized linear mixed effect model

run on a binomial distribution (g*lmer* from lme4).

Each response variable was run against the same predictor variable: the HFI. We

incorporated population ID as a random effect in each model to account for variation between

plants in the population that was unrelated to urbanization. We assessed the assumptions of

the models by examining the distribution of residuals, and cube-root transformed percent

herbivory to better meet these assumptions. We assessed the significance of HFI on herbivory

using Analysis of Variance (ANOVA) implemented with the *Anova* function in the *car* v.

3.0-6 package (Fox and Weisburg 2011) to calculate Wald chi-square test statistics with Type

II sums of squares.

Results

Herbivory rates increased with urbanization. We found that there was a significant positive

effect of HFI on percent herbivory ($\chi^2_1 = 4.49$, p = 0.034), where populations with the highest

HFI experienced greater herbivory that populations with lower HFI (Fig. 2a). Similarly, we

found there was a significant positive effect of HFI on the likelihood of herbivory ($\chi^2_1 =$

10.17, p = 0.001), where urban populations were more likely to experience herbivory than

nonurban populations (Fig. 2b). Two populations experience greater herbivory than others

(Fig. 2), however removal of these outliers returned a similar result thus we retained them in

the analysis (percent herbivory: p = 0.063, presence: p = 0.001).

We confirmed that this pattern of herbivory was due to differences between urban and

rural habitats by running the models with habitat type (urban vs rural; Fig. 1) as the predictor.

Both percent herbivory ($\chi^2_1 = 7.09$, p = 0.008) and presence of herbivory ($\chi^2_1 = 15.89$, p <

0.001) were significantly greater in urban habitats (Fig. 2). Lastly, a two-sided t-test

demonstrated that 33% of plants in urban populations exhibited evidence of lace bug

herbivory whereas only 4.6% of plants in rural populations exhibited lace bug damage (t =

3.98; df = 30; p < 0.001, Figs. S2; S3).

We assayed five rural populations (97 plants) and seven urban populations (138

plants) for HCN. We detected no evidence of HCN production in any of the populations we

assayed, suggesting that this chemical defence has not evolved in these populations.

Fig. 2. A) Percent leaf herbivory averaged across all populations increases with Human

Footprint Index (p = 0.034), corresponding to greater herbivory in urban habitats (p = 0.008;

inset). B) Presence of leaf herbivory averaged across all populations also increases with

Human Footprint Index (p = 0.001), corresponding to increased likelihood of herbivory in

urban habitats (p < 0.001; inset). Overlapping points have been jittered for clarity.

Discussion

We found that urbanization led to increased levels of herbivory on *T. subulata*, however we did not find any evidence for the evolution of HCN in any of the plants we tested. These results suggest that elevated herbivore pressure due to urbanization is insufficient to lead to the evolution of chemical defence in the populations we studied. Our results reinforce earlier findings reporting that *T. subulata* has not evolved cyanogenesis (Shore & Obrist, 1992). Arthropod herbivory appears to be the main selective driver for cyanogenesis in the closely related *T. ulmifolia* (Schappert & Shore, 1999), and this herbivore-plant interaction can drive fast evolutionary changes in plant chemical defences (Agrawal et al. 2012; Zanger et al., 2008). Nevertheless, the absence of HCN in *T. subulata* populations might be linked to more efficient mechanical protection (e.g., pilosity in leaves) or mutualistic interactions in this species that could fend off herbivory without need of HCN. Moreover, a large proportion of herbivory in urban plants were caused by lace bugs (*Gargaphia sp.*) and their feeding methods are unlikely to cause cell rupture and release of HCN, minimizing selection for this type of chemical defence.

Lace bugs feed on sap of living plants using specialized buccal apparatus that pierce the cellular tissue to extract the sap and cause serious damage in the foliage (Guidoti et al. 2015), which lead to some species being used to control population of weed plants (Baars & Heystek 2003). For instance, lace bugs are considered as one of the three most successful biological control agents against the invasive plant *Lantana camara* (Guidoti et al., 2015; Baars & Heystek 2003). Lace bugs feed gregariously on plants and cause heavy damage to the leaves (Guidoti et al. 2015) and are likely to have a strong impact on plant populations where they are abundant. In our study, we found urban plant population were more likely to exhibit lace bug damage than rural populations, suggesting that urbanization drives changes in lace bug abundance. At a community level, lace bug herbivory on *T. subulata* may be

regulate urban community structure by inflicting reduction on competition within and among

plant species; although speculative, this possibility deserves further study.

City life: a road to ruin?

We found convincing evidence that urbanization is associated with elevated herbivory rates

in *T. subulata* populations. This trend may be the results of increased vehicular pollution, a

ubiquitous feature of urbanization. Despite its negative impacts on many organisms (Durrani

et al. 2004; Rai, 2016), pollutants like NOx can increase the nitrogen content in leaves and

the nutritional quality of plant foliage (Bell et al. 2011; Moreira et al. 2019; Spencer et al.

1988; Truscott et al. 2005). Increased foliage quality has been linked to high arthropod

herbivore abundance in urban environment (reviewed in Miles et al 2019) because many

arthropods are nitrogen limited (Dale & Frank, 2017; Jones & Leather, 2012; Raupp et al.

2010), and may be capitalizing on the increased nitrogen in roadside plants. Although we

collected rural plants along a highway with high traffic load, rural plants were surrounded by

fewer anthropogenic sources such as impervious surfaces, possibly reducing their exposure to

pollutants.

In addition to pollution, urban habitats often exhibit reduced plant species richness

due to increased habitat fragmentation (Aronson et al., 2014). Consequently, herbivory may

be elevated in urban *T. subulata* sites because there are fewer species for generalist

herbivores to feed upon. Lastly, rural populations may also harbour a greater diversity of

herbivore predators than urban areas (Raupp et al. 2010; Turrini et al., 2016). A more diverse

predatory guild in rural areas could be exerting greater top-down control compared to urban

sites, reducing herbivory on plants in rural habitats. A number of studies suggest urbanization

is associate with a reduction of predator and parasitoid species, leading to a release form

predation in lower-guild species (Burkman & Gardner, 2014; Denys & Schmidt 1998;

McIntyre 2000; Rocha & Fellowes, 2020). The decline in richness at high trophic levels in

urban environment appears to be a pervasive trend and has been associated with increased

herbivory in other systems (Rivkin et al 2020; Turrini et al. 2016; Raupp et al. 2010; Cuevas-

Reyes et al. 2013; McIntyre 2000).

A complex relationship between abiotic environmental factors and biotic interactions,

exists in urban system, and how these factors jointly influence the strength of the trophic

effects on food webs is yet not well understood (El-Sabaawi 2018; Lagucki et al. 2017;

Turrini et al. 2016). For example, habitat fragmentation and increased barriers to dispersal

(e.g., roads) in cities reduce connectivity, and may lead to lowered abundance and diversity

natural enemies to herbivores (Peralta et al. 2011; Rocha & Fellowes, 2020). Spatial

distribution and patches size of plants could affect the abundance of arthropod species,

leading to altered patterns of herbivory in cities (Fenoglio et al., 2020; Peralta et al. 2011;

Raupp et al., 2010). We suggest that the extensive human footprint found in cities causes

significant shifts in arthropod communities, releasing populations of herbivorous from top-

down control, and facilitating their increased abundance in cities. However, we also found

that this herbivory pressure was insufficient for *T. subulate* to evolve chemical defence to

resist herbivory, thus the role of herbivory as an ultimate driver of plant defence in urban

environments remains unclear.

Literature cited

Agrawal, A.A., Hastings, A.P., Johnson, M.T.J., Maron, J.L. & Salminen, J.-P. (2012). Insect

herbivores drive real-time ecological and evolutionary change in plant populations.

Science 338, 113–116. https://doi.org/10.1126/science.1225977

Aronson, M. F., La Sorte, F. A., Nilon, C. H., Katti, M., Goddard, M. A., Lepczyk, C. A., ...

& Winter, M. (2014). A global analysis of the impacts of urbanization on bird and plant

diversity reveals key anthropogenic drivers. Proceedings of the Royal Society B, 281,

20133330. https://doi.org/10.1098/rspb.2013.3330

Baars, J. R., & Heystek, F. (2003). Geographical range and impact of biocontrol agents

established on *Lantana camara* in South Africa. Biol Control 48: 743-759.

https://doi.org/10.1023/A:1026389207517

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models

using {lme4}. Journal of Statistical Software, *arXiv preprint arXiv:1406.5823*.

Bell, J.N.B., Honour, S.L. & Power, S.A. (2011). Effects of vehicle exhaust emissions on

urban wild plant species. Environmental Pollution, 159: 1984–1990.

https://doi.org/10.1016/j.envpol.2011.03.006

Burkman CE, Gardiner MM (2014) Urban greenspace composition and landscape context

influence natural enemy community composition and function. Biol Control 75:58–67.

https://doi.org/10.1016/j.biocontrol.2014.02.015

Crawley, M.J. (1989). Insect herbivores and plant population dynamics. Annual Review of

Entomology, 34: 531–564. https://doi.org/10.1146/annurev.en.34.010189.002531

Cruz, N.G., Almeida, C.S., Bacci, L., Cristaldo, P.F., Santana, A.S., Oliveira, A.P., Ribeiro,

E.J.M. & Araújo, A.P.A. (2019), Ant associations in the Neotropical shrub Turnera

subulata (Turneraceae): Costs or benefits to the host plant? Austral Ecology, 44: 60-69.

https://doi.org/10.1111/aec.12652

Cuevas-Reyes, P., Gilberti, L., Gonzalez-Rodrıguez, A. & Fernandes, G, (2013), Patterns of

herbivory and fluctuating asymmetry in Solanum lycocarpum St. Hill (Solanaceae) along

an urban gradient in Brazil. Ecological Indicators, 24:557–561.

https://doi.org/10.1016/j.ecolind.2012.08.011

Dale, A.G. & Frank, S. D. (2017). Warming and drought combine to increase pest insect

fitness on urban trees. PLoS ONE 12:e0173844.

https://doi.org/10.1371/journal.pone.0173844

Dale, A. G. & Frank, S. D. (2018) Urban plants and climate drive unique arthropod

interactions with unpredictable consequences. Curr. Opin. Insect Sci. 29, 27–33.

https://doi.org/10.1016/j.cois.2018.06.001

Denys C, Schmidt H (1998) Insect communities on experimental mugwort (Artemisia

vulgaris L.) plots along an urban gradient. Oecologia 113:269–277.

https://doi.org/10.1007/s004420050378

Durrani G.F.; Hassan, M.; Baloch, M.K.; Hameed, G., (2004). Effect of traffic pollution on

plant photosynthesis. Journal of the Chemical Society Pak., 26: 176-179.

El-Sabaawi R (2018) Trophic structure in a rapidly urbanizing planet. Functional Ecology 32:

1718–172. https://doi.org/10.1111/1365-2435.13114

Faeth, S. H., Marussich, W. A., Shochat, E., and Warren, P. S. (2005). Trophic dynamics in

urban communities. Bioscience 55, 399–407. https://doi.org/10.1641/0006-

3568(2005)055[0399:TDIUC]2.0.CO;2

Fenoglio, M. S., Rossetti, M. R., and Videla, M. (2020). Negative effects of urbanization on

terrestrial arthropod communities: A meta‐analysis. Global Ecology and Biogeography,

29, 1412-1429. https://doi.org/10.1111/geb.13107

Fox and Weisburg 2011

Grimm, N.B, Faeth, S.H., Golubiewski, N.E., Redman, C.L., Wu J, Bai X. & Briggs JM

(2008). Global change and the ecology of cities. Science 319: 756–760.

https://doi.org/10.1126/science.1150195

Guidoti M, Montemayor SI, Guilbert É, 2015. Lace bugs (Tingidae). In: Panizzi AR, Grazia

J. (eds). True Bugs (Heteroptera) of the Neotropics. Springer Science, Dordrecht. pp. 395–

419.

Hanley, M. E., Lamont, B. B., Fairbanks, M. M., & Rafferty, C. M. (2007). Plant structural

traits and their role in anti□herbivore defence. Perspectives in Plant Ecology Evolution

and Systematics, 8, 157– 178. https://doi.org/10.1016/j.ppees.2007.01.001

Honour, S.L, Bell, J.N.B., Ashenden, T.W., Cape, J.N. & Power, S.A. (2009). Responses of

herbaceous plants to urban air pollution: effects on growth, phenology and leaf surface

characteristics. Environental Pollution, 157: 1279–1286.

https://doi.org/10.1016/j.envpol.2008.11.049

Hughes MA. 1991 The cyanogenic polymorphism in Trifolium repens L. (white clover).

Heredity 66, 105–115.

Johnson, M. T. J., & Munshi-South, J. (2017). Evolution of life in urban environments.

Science 358:eaam8327. https://doi.org/10.1126/science.aam8327

Jones, E.L., & Leather, S.R. (2012). Invertebrates in urban areas: a review. European Journal

of Entomolology, 109, 463–478. https://doi.org/10.14411/eje.2012.060

Kozlov, M. V., Lanta, V., Zverev, V., Rainio, K., Kunavin, M. A., & Zvereva, E. L. (2017).

Decreased losses of woody plant foliage to insects in large urban areas are explained by

bird predation. Global Change Biology, 23, 4354–4364. https://doi.org/10.1111/gcb.13692

Lagucki E, Burdine JD, McCluney KE (2017) Urbanization alters communities of flying

arthropods in parks and gardens of a medium-sized city. PeerJ 5:e3620

Machado, B. B., Orue, J. P. M., Arruda, M. S., Santos, C. V., Sarath, D. S., Goncalves, W. N.

et al. (2016). BioLeaf: A professional mobile application to measure foliar damage caused

by insect herbivory. Computers and Electronics in Agriculture, 129, 44–55.

https://doi.org/10.1016/j.compag.2016.09.007

Mauricio, R., & Rausher, M. D. (1997). Experimental manipulation of putative selective

agents provides evidence for the role of natural enemies in the evolution of plant defense.

Evolution, 1435–1444. https://doi.org/10.1111/j.1558-5646.1997.tb01467.x

McIntyre, N.E., Rango, J., Fagan, W.F. & Faeth, S.H. (2001). Ground arthropod community

structure in a heterogeneous urban environment. Landscape and Urban Planning, 52, 257–

274. https://doi.org/10.1016/S0169-2046(00)00122-5

Miles, L.S., Breitbart, S.T., Wagner, H.H. &. Johnson, M.T.J. (2019). Urbanization shapes

the ecology and evolution of plant-arthropod herbivore interactions. Frontiers in Ecology

and Evolution, 7:310. | https://doi.org/10.3389/fevo.2019.00310

Moreira, X., Abdala-Roberts, L., Teran J.C.B., Covel, F., Mata, R., Francisco M. (2019)

Impacts of urbanization on insect herbivory and plant defences in oak trees. Oikos 128,

113–12. https://doi.org/10.1111/oik.05497

Peralta G, Fenoglio MS, Salvo A (2011) Physical barriers and corridors in urban habitats

affect colonisation and parasitism rates of a specialist leaf miner. Ecol Entomol 36:673–

679. https://doi.org/10.1111/j.1365-2311.2011.01316.x

Rai PK (2016) Impacts of particulate matter pollution on plants: implications for

environmental biomonitoring. Ecotoxicol Environ Saf 129: 120–136.

https://doi.org/10.1016/j.ecoenv.2016.03.012

Raupp, M. J., Shrewsbury, P. M., and Herms, D. A. (2010). Ecology of Herbivorous

Arthropods in Urban Landscapes. Annuall Review of Entomogy, 55, 19–38.

ttps://doi.org/10.1146/annurev-ento-112408-085351

Rivkin, L. R., Santangelo, J. S., Alberti, M., Aronson, M. F. J., de Keyzer, C. W., Diamond,

S. E., et al. (2019). A roadmap for urban evolutionary ecology. Evolutionary Applications

18, 1–15. https://doi.org/10.1111/eva.12734

Rivkin, L. R., Johnson, R. A., Chaves, J. A., Johnson, M. T. (2020). Urbanization alters

ecological and evolutionary interactions between Darwin's finches and *Tribulus cistoides*

on the Galápagos Islands. BioRxiv. https://doi.org/10.1101/2020.09.26.301507

Rocha EA, Fellowes MD (2020) Urbanisation alters ecological interactions: ant mutualists

increase and specialist insect predators decrease on an urban gradient. Sci Rep 10:6406.

https://doi.org/10.1038/s41598-020-62422-z

Schappert PJ, Shore JS (1995) Cyanogenesis in *Turnera ulmifolia* L. (Turneraceae). I.

Phenotypic distribution and genetic variation for cyanogenesis on Jamaica. Heredity

74:392–404. https://doi.org/10.1038/hdy.1995.57

Schappert, P.J. & Shore, J.S. (1999) Cyanogenesis, herbivory and plant defense in *Turnera*

ulmifolia on Jamaica. Écoscience, 6: 511-520.

https://doi.org/10.1080/11956860.1999.11682560

Schlindwein, C. and P. C. R. Medeiros. 2006. Pollination in Turnera subulata (Turneraceae):

Unilateral reproductive dependence of the narrowly oligolectic bee Protomeliturga

turnerae (Hymenoptera, Andrenidae). Flora Morphol. Distrib. Funct. Ecol. Plants.

201:178-188. https://doi.org/10.1016/j.flora.2005.07.002

Sepp, T., Ujvari, B., Ewald, P. W., Thomas, F., & Giraudeau, M. 2019. Urban environment

and cancer in wildlife: available evidence and future research avenues. Proceedings of the

Royal Society B, *286*: 20182434. https://doi.org/10.1098/rspb.2018.2434

Shore JS, Obrist CM. 1992. Variation in cyanogenesis within and among populations and

species of Turnera series Canaligerae (Turneraceae). Biochemical Systematics and

Ecology, 20: 9-15.

Spencer H.J., Scott N.E., Port G.R. & Davison A.W. (1988). Effects of roadside conditions

on plants and insects. I. Atmospheric conditions. Journal of Applied Ecology 25: 699–707.

https://doi.org/10.2307/2403855

Thompson K.A., Renaudin, M. & Johnson M.T.J. (2016). Urbanization drives the evolution

of parallel clines in plant populations. Proceedings of the Royal Society B 283: 20162180.

http://dx.doi.org/10.1098/rspb.2016.2180

Truscott, A.M., S.C.F. Palmer, G.M. McGowan, J.N. Cape, and S. Smart. 2005. Vegetation

composition of roadside verges in Scotland: The effects of nitrogen deposition,

disturbance and management. Environmental Pollution 136: 109–118.

https://doi.org/10.1016/j.envpol.2004.12.009.

Turrini, T., Sanders, D., and Knop, E. (2016). Effects of urbanization on direct and indirect

interactions. Ecological Application 26, 664–675. https://doi.org/10.1890/14-1787

Venter, O., Sanderson, E.W., Magrach, A., Allan, J.R., Beher, J., Jones, K.R., Possingham,

H.P., Laurance, W.F., Wood, P., Fekete, B.M. and Levy, M.A. (2016a). Sixteen years of

change in the global terrestrial human footprint and implications for biodiversity

conservation. Nature communications, 7:12558. https://doi.org/10.1038/ncomms12558

Venter, O., Sanderson, E.W., Magrach, A., Allan, J.R., Beher, J., Jones, K.R., Possingham,

H.P., Laurance, W.F., Wood, P., Fekete, B.M. and Levy, M.A. (2016b). Data from: Global

terrestrial Human Footprint maps for 1993 and 2009, Dryad Dataset.

https://doi.org/10.5061/dryad.052q5

Zangerl, A. R., & Berenbaum, M. R. (2005). Increase in toxicity of an invasive weed after

reassociation with its co□evolved herbivore. Proceedings of the National Academy of

Sciences of the United States of America, 102, 15529– 15532.

https://doi.org/10.1073/pnas.0507805102

Zangerl, A.R, Stanley, M.C. & Berenbaum, M.R. (2008). Selection for chemical trait

remixing in an invasive weed after reassociation with a coevolved specialist. Proc. Natl.

Acad. Sci. USA 105: 4547–52. https://doi.org/10.1073/pnas.0710280105

Züst, T., Heichinger, C., Grossniklaus, U., Harrington, R., Kliebenstein, D. J. & Turnbull, L.

A. (2012). Natural enemies drive geographic variation in plant defenses. Science 338,

116–119. https://doi.org/10.1126/science.1226397