Mechanosensitive Ion Channel Piezo1 Regulates Myocyte Fusion during Skeletal Myogenesis

Huascar Pedro Ortuste Quiroga[1], Shingo Yokoyama[2], Massimo Ganassi[3], Kodai Nakamura[1], Tomohiro Yamashita[1], Daniel Raimbach[4], Arisa Hagiwara[1], Atsushi Asakura[5], Yoshiro Suzuki[6], Makoto Tominaga[6], Peter S. Zammit[3] & Katsumasa Goto[1,2,]*

[1]Department of Physiology, Graduate School of Health Sciences, Toyohashi SOZO University

[2]Laboratory of Physiology, School of Health Sciences, Toyohashi SOZO University

[3]Randall Centre for Cell and Molecular Biophysics, King's College London, London, SE1 1UL, UK

[4]Centre of Human and Aerospace Physiological Sciences, King's College London, London, SE1 1UL, UK

[5]Stem Cell Institute, Paul & Sheila Wellstone Muscular Dystrophy Center, Department of Neurology, University of Minnesota Medical School

[6]Division of Cell Signalling, National Institute for Physiological Sciences

* Address for correspondence;

Katsumasa Goto, Ph.D.

Laboratory of Physiology, Toyohashi SOZO University

23   20-1 Matsushita, Ushikawa, Toyohashi City, Aichi 440-8511, Japan

24   TEL +81-50-2017-2272

25   FAX +81-532-55-0803

26   E-mail: gotok@sepia.ocn.ne.jp

Abstract

Mechanical stimuli such as stretch and resistance training are essential to regulate growth and function of skeletal muscle. However, the molecular mechanisms involved in sensing mechanical stress remain unclear. Here, the purpose of this study was to investigate the role of the mechanosensitive ion channel Piezo1 during myogenic progression. Muscle satellite cell-derived myoblasts and myotubes were modified with stretch, siRNA knockdown and agonist-induced activation of Piezo1. Direct manipulation of Piezo1 modulates terminal myogenic progression. *Piezo1* knockdown suppressed myoblast fusion during myotube formation and maturation. This was accompanied by downregulation of the fusogenic protein *Myomaker*. *Piezo1* knockdown also lowered Ca^{2+} influx in response to stretch. Conversely Piezo1 activation stimulated fusion and increased Ca^{2+} influx in response to stretch. These evidences indicate that Piezo1 is essential for myotube formation and maturation, which may have implications for msucular dystrophy prevention through its role as a mechanosensitive Ca^{2+} channel.

Key words: skeletal muscle, satellite cells, mechanosensation, Piezo1, Ca^{2+} channel, myocyte fusion

Introduction

Skeletal muscle is a highly specialised tissue composed of multi-nucleated, post-mitotic muscle fibres. Since myonuclei within a muscle fibre do not divide after development, the production of new myonuclei is entrusted to satellite cells (SCs), the skeletal muscle's resident stem cell. SCs are found on the surface/periphery of postnatal skeletal muscle fibres [1-3] and in response to muscle damage, rapidly activate to generate a myoblast progeny that proliferate, undergo myogenic differentiation, and fuse to repair damaged myofibres, resulting in regeneration of a functional muscle [4]. Although much is known about the myogenic progression program, some of the underlying mechanisms remain to be determined. Among these are the molecular mechanisms involved in sensing mechanical stress (mechanosensation) and its effect on myocyte fusion at the cellular and molecular levels.

Mechanosensitive (MS) ion channels are pore-forming membrane proteins which gate in response to mechanical stimuli applied on the cell membrane [5-7]. MS ion channels have been linked to many physiological processes associated with mechanosensory transduction; including osmoregulation, proprioception, hearing, touch, blood flow regulation to name but a few examples [8-10]. Piezo1 and Piezo2 were first identified by Coste et al, (2010) as the long-sought principal types of molecular force sensors (mechanosensors) in mammalian cells [5]. Piezo1 (and Piezo2) is a very large protein containing ~2500 amino acids with each subunit (a total of three subunits per channel) containing an estimated 24-40 transmembrane (TM) segments [6,8,11,12]. Characterisation of ionic selectivity revealed that Piezo1 was nonselective, permeating Na^+, K^+, Ca^{2+} and Mg^{2+} with a preference for Ca^{2+} [5,13]. With such a crucial role Ca^{2+} regulation plays in skeletal muscle maintenance and repair, understanding Piezo1's function may prove vital when looking at strategies for therapeutic interventions of muscular dystrophies [14,15].

Piezo1 is widely expressed in a range of tissues, including skeletal muscle, the bladder,

colon, lung, skin and stomach [5,16]. Piezo2 shows a similar pattern with the exception of

being highly expressed in the dorsal root ganglion (DRG) while being less represented in

skeletal muscle [5,16]. Another key difference between Piezo1 and Piezo2 is their activation

and inactivation kinetics. Overexpression analyses revealed that the kinetics of inactivation of

Piezo2-dependent mechano-activated currents was faster than Piezo1 both for inward and

outward currents, thus conferring distinct channel properties [5,13,11]. Although Tsuchiya et

al published the first report on Piezo1 in skeletal muscle cells [17], looking at the role of

transmembrane localisation of phosphatidylserine and Piezo1 activation, the physiological

function(s) of Piezo1 throughout the myogenic program remain unelucidated.

The current study aimed to analyse the role of Piezo1 in skeletal muscle proliferation,

differentiation and its role in stretch-induced Ca^{2+} influx of primary derived myotubes. Our

findings revealed that Piezo1 is dispensable for myoblast proliferation and onset of

differentiation, but its knockdown suppressed myotube formation and maturation in primary

myotubes derived from slow soleus and fast extensor digitorum longus (EDL) muscles in

mice. In line, *Piezo1* reduction was accompanied by downregulation of the fusogenic gene

Myomaker, decreased accumulation of f-actin and lowered Ca^{2+} influx of myotubes in

response to mechanical stretch. In contrast, administration of the Piezo1-specific agonist

Yoda1 increased fusion of myoblasts. Piezo1 activation also showed increased Ca^{2+} influx in

response to stretch. Using publicly available datasets, we showed a dysregulation in the

expression of Piezo1 in the skeletal muscle disease facioscapulohumeral muscular dystrophy

(FSHD). In summary, we propose that Piezo1 plays a crucial role at the terminal stage of

myotube fusion and maturation. Piezo1 may employ direct or indirect mechanisms that

regulate fusion proteins as well as proteins involved in cytoskeletal organisation. These

mechanisms may be regulated by intracellular Ca^{2+}-signals mediated by mechanosensitive

Piezo1 channels.

Materials and methods

Primary myoblast cell culture

All experimental procedures were carried out in accordance with the Guide for the

Care and Use of Laboratory Animals as adopted and promulgated by the National Institutes

of Health (Bethesda, MD, USA) and were approved by the Animal Use Committee of

Toyohashi SOZO University A2018006, A2019006). Male C57BL/6J mice (8-12 weeks of

age) were used. All mice were housed in a vivarium room with 12-h-12-h light-dark cycle;

with temperature and humidity maintained at ~23°C and ~50%, respectively. Solid food and

water were provided *ad libitum*.

After cervical dislocation, the EDL (*extensor digitorum longus*) and soleus muscles

were carefully dissected, and manipulated only by their tendons. Muscles were digested in

0.2% Collagenase Type 1 (Sigma, UK. Ref: SCR103) in Dulbecco's Modified Eagle Medium

(DMEM, Gibco, Thermo Fisher Scientific, Ref: 11885084) with 1% penicillin/streptomycin

(Pen Strep, Gibco, Thermo Fisher Scientific, Ref: 15140-122) for 2 hours. Individual

myofibres were then dissociated by trituration using heat polished glass Pasteur pipettes

(Marienfeld, Germany. Ref: 3233049) with variously sized apertures (coated with 5% BSA,

Sigma-Aldrich, Ref: A7906-100G) and washed as described by Collins and Zammit

(2009)[18]. Isolated myofibres were plated on Matrigel (Corning. Ref: 354234) and the

satellite cell-derived myoblasts were then expanded in proliferation medium, consisting of;

DMEM, with 30% heat-inactivated foetal bovine serum (FBS) (Gibco, Thermo Fisher

Scientific. Ref: 26140-079), 10% horse serum (Gibco, Thermo Fisher Scientific, Ref: 16050-

122), 1% chick embryo extract (Sera Laboratories. Ref: CE-650-TL), 10 ng/ml basic FGF

(bFGF, Gibco, Thermo Fisher Scientific. Ref: PHG0264) and 1% penicillin. Cells were kept

in a 37°C incubator (Panasonic, MCO-230AICUVH) under a humidified atmosphere with

95% air and 5% CO_2. Cells designated for proliferation and differentiation conditions were

seeded at different densities depending on the size of wells they were cultured in. For 96-well

plate proliferation conditions required 5,000 cells per well, and for differentiation, cells were

seeded at 10,000 cells per well. For 6-well plates, proliferating and differentiation cohorts

consisted of 50,000 cells and 70-80,000 per well, respectively. Differentiation medium was

made up of DMEM, 2% heat-inactivated horse serum, and 1% penicillin.

siRNA transfection

Small interfering RNAs (siRNAs) were purchased from (Qiagen, Hiden, Germany)

(Table 1) and diluted to 20 or 10 µM in double-distilled water (ddH2O) and stored at -20°C.

To investigate the effects *Piezo1* knockdown on proliferation, early entry into differentiation

and myotube formation, *Piezo1*-targeting or control scrambled siRNA (siScrambled; Qiagen,

Hiden, Germany) was transfected in proliferation medium. Cells were plated on 6-well plates

at 50,000 cells per well in proliferation medium. Following a 24 h incubation period, the

medium was replaced with 1.75 ml fresh proliferation medium and the transfection mixture

was prepared: A solution 150 µl Optimem (Gibco, Thermo Fisher Scientific. Ref: 31985-070)

medium with 9 µl of lipofectamine (lipofectamine RNAiMAX Thermo Fisher Scientific. Ref:

13778030) was made for each well. Separately, siRNA was diluted in 150 µl Optimem. The

two solutions were then mixed and incubated for 5 min. 250 µl of the siRNA/lipofectamine

mixture was added to corresponding wells dropwise. The final siRNA concentration was set

at 10 nM. Following overnight incubation in the transfection medium, cells were trypsinised

for RT-qPCR analysis and seeded in 96-well plates for proliferation and differentiation

cohorts (day 1 and day 3 differentiation) in proliferation and differentiation medium,

respectively. After a 24-hour incubation, proliferating cells were subjected to a 2-hour 5-

ethynyl-2'-deoxyuridine (EdU) pulse and fixed as below. Day 1 and day 3 differentiating

cohorts were also fixed.

To determine *Piezo1*'s role in myotube formation, siRNA transfection was performed

in early differentiated myotubes. Cells were seeded at confluency in differentiation medium.

Following 24-hour incubation in differentiation medium, siRNA transfection was performed.

Cohorts were designated for RT-qPCR analyses and immunolabelling.

Table 1. List of siRNAs used

Gene	Species	siRNA ID
scrambled non-targeting siRNA (All Stars Negative Control siRNA)	Mouse	Qiagen, 1027281
Piezo1	**Mouse**	**Qiagen, S104420409**
Piezo1	Mouse	Qiagen, S104420402
Piezo1	Mouse	Qiagen, S100814807
Piezo1	Mouse	Qiagen, S100814821

N.B. The highlighted *Piezo1* siRNA (S1044120409) was used for most of the

experiments. The other three were used as validators of our obtained results.

RNA Extraction and Reverse Transcription

RNA was extracted from cells using the RNeasy mini kit as per manufacturer's

requirements (Qiagen. Ref: 217004). Reverse transcription was carried out using PrimeScript

RT Master Mix (Takara Bio, Otsu, Japan. Ref: RR036A). Optical density analysis using a

Nanodrop ND-1000 spectrophotometer (Labtech, UK) quantified RNA concentration.

Samples were then loaded to a PCR thermal cycler (Takara, Dicemini). The resulting cDNA

was then diluted 1:9 to obtain a working dilution for RT-qPCR analysis.

Real-time Quantitative PCR (RT-qPCR)

Primers were designed using the Takara Bio Perfect Real Time Support System

(Takara Bio, Table 2). Primers were diluted to 50 μM in ddH_20 and stored at -20°C. Real-

time RT-qPCR was performed on the cDNA (Thermal Cycler Dice Real Time System

IIMRQ, Takara Bio) using Takara SYBR Premix Ex Taq II (Takara Bio. Ref: RR802A). 12.5

μl of SYBR Premix Ex were added to each RT-qPCR well. 8.5 μl of ddH20 and 2 μl of the

corresponding primers were then added (a final concentration of 2 μM per primer). 2 μl of

the respective cDNA was then added to the appropriate wells, bringing the total volume to

25μl per well. The RT-qPCR cycle consisted of 95°C for 30 s (for enzyme activation),

followed by 40 cycles at 95°C for 5 s and a qPCR amplification period of 30 s at 60°C. The

relative fold change of expression was calculated by the comparative threshold cycle (CT)

method using Takara Thermal Cycler Dice Real Time System Software Ver. 4.00 (Takara

Bio). To normalise for the amount of total RNA present in each reaction, *Gapdh* was used as

an internal standard.

185 **Table 2. Real Time quantitative PCR (RT-qPCR) primers.**

Gene	Species	Forward primer (5'-3')	Reverse primer (5'-3')	Reference
Gapdh	Mouse	TGTGTCCGTGGATCTGA	TTGCTGTTGAAGTCGCAGGAG	Takara Bio, MA050371
Piezo1	Mouse	CTTTATCATGAAGTGCAGCCGAG	CCAGATGATGGCGATGAGGA	Takara Bio, MA125411
Myomaker	Mouse	CATGCGCCGTGACATTCTG	AAGCATTGTGAAGGTCGATCTCTG	Takara Bio, MA131293
Myomixer	Mouse	GAATCCACCGCAGGCAAA	ACCATCGGGAGCAATGGAAC	Takara Bio, MA101853

Piezo1 activation

In order to induce Piezo1 activation, early forming myotubes were subjected to the

Piezo1 specific agonist Yoda1 (Cayman Chemical Company. Ref: 21904) diluted in dimethyl

sulfoxide (DMSO, Sigma. Ref: D2650-5x5ML). Yoda1 binds the agonist transduction motif

(ATM), located at the pore domain of the Piezo1 channel [19]. With each subunit displaying

such motif, Yoda1 has potentially three binding sites. This phase of the investigation

consisted of two main variables 1) drug concentration and 2) duration of time cells were

incubated with the drug. Five concentrations were chosen in order to cover an increasing

spectrum of Yoda1 final concentration, these were: 5, 10, 30 and 100 μM diluted in

differentiation medium. Preliminary findings from the group found that a 24-hour incubation

with any of the concentrations chosen, led to complete abolishment of myotube maturation

(data not show). Thus, time-points thought to have potential to maximise myotube

fusion/maturation were tested. The incubation time-points were set for 1 and 30 min, 1 and 4

hours. Control cohorts containing only DMSO were incubated at the allocated times to allow

comparisons to be made within each condition. Following the incubation of Yoda1 or DMSO,

cells were cultured in the differentiation medium for a further 2 days (i.e. myotubes were

analysed 3 days post initial induction of differentiation).

Immunolabelling

Throughout the protocol, all washes were performed with Dulbecco's phosphate-

buffered saline (DPBS, Gibco, Dulbecco's Phosphate Buffered Saline, Thermo Fisher

Scientific, Ref: 14190-144). Cells were fixed with 4% paraformaldehyde for 15 min. Samples

were then washed three times with PBS (5 minutes each wash) and permeabilised for 15 min

using 0.5% triton-X100/PBS (Sigma-Aldrich. Ref: T9284-500ml). Cells were blocked for 1

hour in 5% bovine serum albumin (BSA, Sigma-Aldrich Ref: A7906-100G). Primary

antibodies (diluted to the working concentration in PBS) (Table 3) were added to the samples

and incubated overnight at 4ºC. Primary antibodies were decanted, the samples were washed

three times and appropriate secondary antibodies diluted to the working concentration in PBS

added (Table 3) were added to the samples. The Samples were covered with aluminium foil

to avoid light exposure and left to stand at room temperature for 1 hour. Cells were washed

again (three times). To visualise nuclei, the cells were incubated for 10 min at room

temperature with 1 µg/ml 4', 6-diamidino-2-phenylindole (DAPI) (Sigma. Ref: D9542-

10MG) diluted 1:1000 in PBS. After a final wash with PBS (5 min), cells were replenished

with PBS and stored at 4 ºC until image analysis.

Table 3. Primary and Secondary antibodies used

Primary antibody	Dilution	Reference
Monoclonal mouse – myogenin	1:10	Development Studies Hybridoma Bank (DSHB), F5D-s
Monoclonal mouse – MF20 (Myosin Heavy Chain)	1:10	Development Studies Hybridoma Bank (DSHB), MF20-s
Secondary antibody	**Dilution**	**Reference**
Donkey anti-mouse IgG (H+L), Alexa Fluor® 555	1:500	Life Technologies, A21203

EdU incorporation

For the evaluation of cell proliferation, cells were incubated with 5-ehtynyl-2'-

deoxyuridine (EdU: Invitrogen, Thermo Fisher Scientific) at 10 μM, added in fresh

proliferation medium for 2 hours at 37 °C. EdU, the alkyne-containing thymidine analog, is

incorporated into DNA during active DNA synthesis. The click-iT EdU Alexa Flour kit

(Invitrogen, Thermo Fisher Scientific, Click-iT, EdU Alexa Fluor. Ref: 594 C10339) was

used as per manufacturer's instructions with either the 488 (green) or 594 (red) azide to

detect incorporated EdU.

Phalloidin labelling

To evaluate the cytoskeleton, cells were treated with phalloidin (Invitrogen, Thermo

Fisher Scientific, Alexa-Fluor. Ref: 488 A12379) diluted 1:40 in PBS. Phalloidin binds to f-

actin, a major cytoskeleton protein in skeletal muscle fibres. Cells were incubated with

phalloidin solution for 30 mins at room temperature. Cells were then washed with PBS twice

for 5 min.

Image quantification

Images were taken using a fluorescence microscope (BZ-X710, KEYENCE, Osaka,

Japan). Four to five images per each well (3 wells per repeat) consistent of a total of 12-15

images per repeat were analysed. One repeat refers to one mouse. For EdU incorporation, the

total number of DAPI-counterstained nuclei and total number of EdU-incorporated cells were

quantified. The proportion of EdU-incorporated cells relative the total number of nuclei was

subsequently expressed as percentages. The relative proportion of cells expressing myogenin

was also quantified in this manner.

The fusion index was calculated by quantifying the total number of nuclei within

MyHC positive myotubes and expressing this value as a proportion of the total number of

nuclei in each field of view. As a criterion, more than two nuclei must be within a MyHC

positive myotube to be quantified for the fusion index.

To measure myotube width, the "measure" tool on ImageJ imaging software was used.

This allows measurements of a chosen distance to be made. Before measurement, a scale was

applied to all images. On the "set scale" option pixels are converted into μm. Taking the

fluorescence microscope and magnification into account the program determines 100 μm to

be 133.00 pixels or 0.75 μm/pixel. Three independent images were chosen per condition. The

criterion for this analysis was to choose the widest possible distance between myotube edges

without the presence of any branching points. This was carried out in three independent

points within the field of view. The values were then averaged.

Stretch experiments and intracellular Ca^{2+} level imaging

Stretch experiments were performed at the National Institute for Physiological Sciences

(NIPS). Primary derived myoblasts were seeded (30,000 cells/chamber) on modified elastic

silicone chambers (Strexcell, Ooyodonaka, Ref: STB-CH-0.02). After 24-hour incubation,

cells were transfected with either control (siScrambled) or *Piezo1*-specific siRNA in

proliferation medium. After overnight incubation, cells were switched to differentiation

medium and cultured for a further 3 days. Stretch experiments were conducted on the third

day. For Piezo1 activation by Yoda1 administration, cells were seeded as above. Following

the initial overnight incubation, cells were switched to differentiate and the resulting

myotubes were analysed 3 days post differentiation induction.

For Ca^{2+} imaging, Fura 2-AM (Invitrogen) with 10% Pluronic® F-127 (Molecular

Probes, USA) diluted in double distilled water (ddH$_2$O), was administered to EDL and

soleus-derived myotubes followed by a 30-min incubation time. Chambers were attached to

an extension device (modified version of STB-150, Strex) on the microscope stage. Stretch

stimulation was applied using a pre-set stretch speed and distance. After an initial 1 min rest

period (0% stretch), stretch was applied at 3% (0.3 mm), 6% (0.6 mm) and 9% (0.9 mm) for

1 min followed by a 1-min resting period in between. During the initial 0% stretch timepoint,

Yoda1 cohorts were administered with 30 μM of the agonist before being subjected to stretch.

Ionomycin (Sigma-Aldrich) at 5 μM was applied at the final step in each experiment for

normalisation and to check cell viability.

Changes in intracellular calcium [Ca^{2+}]i were measured by ratiometric imaging with

Fura 2-AM at 340 and 380 nm, and the emitted light signal was read at 510 nm. Images were

then analysed on ImageJ imaging software. Three independent myotubes from each condition

were selected and analysed. The changes in ratios were calculated by subtracting basal values

from peak values. The values were then normalised to ionomycin data.

Statistical analysis

Data is presented as mean ± SEM from at least three experiments (at least three mice).

Significance was assessed by either paired Student's t-test or one-way ANOVA followed by

followed by the Tukey-Kramer post-hoc; wherein p-values of < 0.05 were considered to be

statistically significant. A paired t-test was adopted when comparing effects within the same

group e.g. analysing the effects of siRNA mediated down-regulation of *Piezo1* versus siRNA

controls in murine derived myoblasts. A one-way ANOVA was implemented when two or

more independent groups were analysed, for example; comparing the effects of varying

agonist concentrations across different timepoints.

300   **Results**

301   *Expression of Piezo1 in primary myoblasts and myotubes*

302   We investigated the expression level of *Piezo1* during myogenic progression in murine

303   fast EDL and slow soleus muscles-derived primary myoblasts (Figure 1a). The expression of

304   EDL-derived cells (Figure 1b) showed a significant increase in mRNA expression of *Piezo1*

305   in myotubes cultured at 3 days in differentiation medium, compared to the expression level in

306   proliferating myoblasts ($p<0.05$). Soleus-derived cells (Figure 1c) showed a significant

307   increase in *Piezo1* mRNA expression after 24 and 72 hours in differentiation medium,

308   compared to proliferation cohorts ($p<0.05$). The observed increase in *Piezo1* in soleus-

309   derived cells after 1 day of differentiation may be explained by the fact that soleus-derived

310   myoblasts tend to differentiate slightly quicker than EDL, i.e. myotubes could be seen sooner

311   in soleus-derived cells compared to EDL. Moreover, we found that *Piezo1* is expressed at a

312   higher level in day 1 and day 3 differentiated soleus cells compared to EDL counterparts

313   (Figure S1).

314

315   *siRNA-mediated downregulation of Piezo1*

316   We next set out to evaluate the effects of manipulating *Piezo1* on proliferation and

317   myogenic differentiation. This was achieved by siRNA-mediated knockdown (Figure 1d and

318   e). Cells were transfected with either control (siScrambled) siRNA (control-siRNA) or

319   targeting siRNA specific for *Piezo1* (*Piezo1*-siRNA). Following a 24-hours incubation period,

320   the medium was changed and cells were incubated for a further 24 hours. In EDL-derived

321   cells (Figure 1d), the treatment of *Piezo1*-siRNA showed a significant reduction (~48%) in

322   mRNA expression level of *Piezo1*, compared to control-siRNA ($p<0.05$). Analysis of soleus-

323   derived cells (Figure 1e) also revealed a significant reduction (~34%) in expression of *Piezo1*

324   post siRNA-mediated transfection, compared to control conditions (p < 0.05).

Figure 1. *Piezo1* **is expressed in satellite cell-derived myoblasts throughout myogenic differentiation. Piezo1**

downregulation does not alter the expression of *Piezo2***.**

a) Schematic representation of muscle isolation and satellite cell expansion procedure. b) and c) Relative fold changes in

expression of *Piezo1* in murine EDL and soleus muscle-derived myoblasts , during proliferation and through differentiation;

24 hours (Day 1) and 72 hours (Day 3) in differentiation medium. d) and e) EDL and soleus-derived myoblasts were

transfected with 10 nM of either control-siRNA (siScrambled) or targeting siRNA against *Piezo1* (*Piezo1*-siRNA. After

overnight incubation, cells were incubated for a further 24 hours and expression of *Piezo1* was measured. f-i) Relative fold

changes in expression of *Piezo1* and *Piezo2* in EDL and soleus muscle-derived myotubes. Following an initial differentiation

period (24 hours at high confluency), cells were transfected with 10 nM of either control-siRNA (siScrambled) or *Piezo1*-

siRNA. After overnight incubation, cells were incubated for a further 24 hours. Values were normalised to *Gapdh* and then

expressed as fold change compared to either level of proliferation (b and c) or to siScrambled conditions (f-i). Data is

presented as mean ± SEM from three experiments (n = 3 mice). Asterisks (*) denote significance at $p < 0.05$ compared to

control conditions (either proliferating myoblasts or siScrambled cohorts), using one-way ANOVA followed by the Tukey-

Kramer post-hoc (a and b) or a 2-tailed paired student t-test (f-i).

Effects of Piezo1 downregulation on Piezo2

Using RT-qPCR analysis, the expression of *Piezo2* in *Piezo1*-downregulated conditions

was measured in both EDL and soleus primary cells (Figures 1f-i). Furthermore as *Piezo1*

seemed to exert most of its expression during differentiation, we evaluated *Piezo2* expression

in early differentiated myotubes. This encompassed a 24-hour incubation in differentiation

medium at high confluence, followed by siRNA transfection. After overnight incubation with

siRNA, the medium was exchanged with fresh differentiation medium and incubated for a

further 24 hours before sampling. We again confirmed the reliability in our method of

siRNA-mediated knockdown by showing significant reduction of *Piezo1* expression in both

EDL- and soleus-derived cells (Figures 1h and i). Data from the EDL (Figure 1f) showed that

in *Piezo1*-downregulated conditions, the expression of *Piezo2* is relatively stable. Soleus-

derived cells subjected to *Piezo1*-siRNA (Figure 1g) showed a slight trending increase in

Piezo2 expression, compared to control-siRNA; however this did not reach statistical

significance, indicating that *Piezo2* did not provide a compensatory role in the down-

regulation of *Piezo1*. Though, we must acknowledge that perhaps if the degree of *Piezo1*

downregulation was more "potent", *Piezo2* may alter its expression in response. Furthermore

a prolonged downregulation of *Piezo1* may also affect the expression level of *Piezo*2.

Piezo1 is dispensable for muscle cell proliferation

We next turned our attention to the effects of *Piezo1* downregulation during

proliferation. Although we did not see a significant difference in the expression dynamics of

Piezo1 early in the myogenic program that is not to say that the knockdown of this channel

would not affect proliferating cells. Thus in order to address this question we used EdU

incorporation assays (Figures 2a-d).–Analyses of the proliferation rate in myoblasts derived

from EDL (Figures 2a and b) showed that the downregulation of *Piezo1* had no significant

effect on EdU incorporation, compared to siRNA control conditions. Similarly, the

downregulation of *Piezo1* in soleus-derived myoblasts (Figures 2c and d) showed no

statistical change in the proliferation rate between *Piezo1*-siRNA-treated cells and control

conditions. From these results, we can summarise that on plated cell conditions,

downregulation of *Piezo1* does not seem to affect the proliferation capacity of EDL- and

soleus-derived myoblasts.

Suppression of Piezo1 does not inhibit the onset of the myogenic program

Our next aim was to investigate whether knockdown of *Piezo1* would alter the capacity

of cells to enter into the differentiation phase of the myogenic program. siRNA-transfected

cohorts were cultured in differentiation medium for 24 hours. This early differentiation phase

was analysed by immunolabelling of the myogenic regulatory factor (MRF) Myogenin.

Quantification was performed by calculating the proportion of Myogenin-positive myoblasts

relative to the number of DAPI counterstained nuclei (Figures 2e-h). Analysis of EDL-

derived myoblasts (Figures 2e and 2f) showed no significant difference in the relative

proportion of myogenin-positive cells between control-siRNA and *Piezo1*-siRNA treated

conditions–Similarly, soleus-derived cells (Figures 2g and 2h) did not show a significant

difference in the relative proportion of myogenin-positive cells in samples treated with

Piezo1-siRNA, compared to control-siRNA. From these results, we may suggest that the

downregulation of *Piezo1* does not seem to have any effects on the ability for myoblasts to

enter the differentiation phase of the myogenic program. However, the question remained

whether the continuous knockdown of this receptor would affect the fusion and maturation of

myotubes. To answer this, we turn to myotube analyses.

Figure 2. *Piezo1* **knockdown does not affect proliferation rate nor early entry into differentiation.**

a) and c) Representative images of EDL and soleus primary-derived myoblasts , transfected with 10nM of siRNAcontrol

(siScrambled) or siRNA-*Piezo1*. Following overnight incubation, the medium was changed with fresh proliferation

medium and cells were incubated for a further 24 hours, and then subjected to a 2-hour pulse with EdU (red panels).

DAPI counterstained nuclei shown in black and white panels. Scale bar is 100 μm. b) and d) Proportion of EdU-

incorporated cells relative to total (DAPI) cell count. e) and g) Representative images of EDL and soleus muscle-derived

myoblasts transfected with 10nM of control-siRNA (siScrambled) or *Piezo1*-siRNA. Following overnight incubation,

cells were incubated for a further 24 hours, immunolabelled for Myogenin (red panels) and counterstained with DAPI

(black and white panels). Scale bar is 100 μm. f) and h) Percentage proportion of Myogenin positive cells relative to total

nuclei. Data is represented as percentages. Data is mean ± SEM from three experiments (n = 3 mice). *: significant at p

< 0.05 compared to siScrambled conditions using a 2-tailed paired student t-test.

Piezo1 regulates myocytes fusion

Piezo1-siRNA-treated cells were cultured in differentiation medium for 3 days post

transfection. By this timepoint myoblasts fuse to neighbouring differentiating myoblasts and

begin to upregulate the sarcomeric motor protein MyHC, which is expressed in the forming

myotubes. We can visualise MyHC containing myotubes through immunolabelling. Not only

does this allow a visual representation of myotube formation, but it can also be used to

quantify the level of fusion that has occurred - the fusion index.

The data derived from the EDL-derived cells (Figure 3a), showed a clear difference

between control-siRNA and *Piezo1*-siRNA conditions, with the latter displaying much fewer

myotubes. Indeed, we observed a significant ($p < 0.05$) reduction of fusion index in *Piezo1*-

siRNA conditions compared to control conditions (Figure 3b). These findings were also

translated to soleus-derived cells. From the myotube images (Figure 3c), we observed smaller

myotubes when *Piezo1* was downregulated. This was confirmed by the significant reduction

in myocytes fusion (Figure 3d) in *Piezo1*-siRNA conditions compared to control conditions

($p < 0.05$).

A potential concern of the data in Figures 3a-d, is the fact that cells were first

transfected in proliferating conditions (i.e. in proliferation medium) before being switched to

differentiate; and although the initial analysis of EdU- and myogenin-positive cells did not

reveal a change post siRNA-mediated knockdown of *Piezo1*, that is not to say that the initial

knockdown in proliferating conditions did not alter the behaviour of the channel and the

subsequent formation of myotubes. Thus, in order to address this, we carried out the

transfection of *Piezo1*-siRNA in differentiating conditions. This encompassed a 24-hour

incubation period in differentiation medium. By this point primary derived myoblasts seeded

at high confluency, already begin to show myotube formation in the relatively small 96-well

plates. These early formed myotubes were then subjected to *Piezo1*-siRNA transfection in

differentiation medium. Following overnight incubation, the transfection mixture was

exchanged with fresh differentiation medium. Myotubes were subsequently analysed 3 days

post the initial *Piezo1*-siRNA transfection (Figures 3e-h).

Downregulation of *Piezo1* in EDL-derived myotubes (Figures 3e and 3f), showed a

significant reduction ($p<0.05$) in the fusion index, compared to control-siRNA conditions.

The soleus muscle (Figures 3g and 3h) also showed a reduction in the ability of myoblasts to

fuse in *Piezo1*-siRNA-treated conditions, compared to control-siRNA cohorts ($p<0.05$). In

summary, the downregulation of the Piezo1 channel both in forming and maturing myotubes

significantly reduces the ability of cells to fuse into new or existing myotubes. This response

seems to be unique to this phase of the myogenic program as downregulation of *Piezo1* does

not affect proliferation or early entry into differentiation.

These thought-provoking results demand further exploration of; 1) the mechanisms of

Piezo1-mediated fusion, 2) other fusion regulators, and 3) the downstream signalling

pathways of the Piezo1 channel which enables correct myogenic fusion.

Figure 3. *Piezo1* knockdown reduced fusion indexes during myotube formation and maturation.

a) and c) Representative images of primary derived myotubes from EDL and soleus muscle, transfected with 10nM of

control-siRNA (siScrambled) or *Piezo1*-siRNA. Following overnight incubation, cells were incubated for a further 72

hours. Cells were immunolabelled for Myosin heavy chain (MyHC) (red panels) and counterstained with DAPI (black

and white panels). b) and d) The fusion index was calculated by counting the total number of nuclei within each

myofibre and representing this as a percentage relative to the total number nuclei in the image taken. e) and g)

Representative images of EDL and soleus muscle-derived. Early forming myotubes were transfected with 10nM of

control-siRNA (siScrambled) or *Piezo1*-siRNA. Following overnight incubation, cells were incubated for a further 72

hours. immunolabelled for Myosin heavy chain (MyHC) (red panels) and counterstained with DAPI (black and white

panels). Scale bar is 100 μm. f) and h) Bar graphs display the fusion index. Data is mean ± SEM from three

experiments for myotube formation data (n = 3 mice) and four experiments for myotube maturation data (n = 4 mice).

*: Significant at $p < 0.05$ compared to siScrambled conditions using a 2-tailed paired student t-test.

Piezo1 and actin remodelling

Previous research places Piezo1 regulation as a key modulator of this cytoskeletal protein [20,21]. In skeletal muscle actin plays a pivotal role not only in myoblast fusion but myotube alignment to attachment sites [22]. Therefore, a putative mechanism(s) of Piezo1 is the regulation of cytoskeletal structures, including filamentous actin (f-actin). As part of our analyses, we examined f-actin intensity using fluorescently labelled phalloidin as a proxy to evaluate its accumulation, and thus the extent of cytoskeletal reorganisation during myogenic differentiation in EDL or Soleus derived myoblasts. *Piezo1*-siRNA-treated cells showed a significant decrease in the accumulation of f-actin compared to control-siRNA (Figure S2). However, activation of Piezo1 using Yoda1 did not reveal a statistically significant increase in f-actin intensity during any of the incubation timepoints or concentrations (Figure S3). Further, high concentration and relatively longer incubation time of Yoda1, suggestive of an over-activation of Piezo1, showed a significant decrease in f-actin in EDL (100 µM; 4-hour incubation) and soleus (100 µM; 30-min, 1- and 4-hour incubation) derived myotubes. These findings suggest that cytoskeletal organisation may be affected by Piezo1 activity; however our methods of measuring such changes lacked resolution. For example, f-actin stress fibre formation as well as filopodia extension in response to Piezo1 manipulation should be quantified under higher magnification.

Lack of Piezo1 affects the expression of fusogenic genes

An ever-increasing topic of interest within the skeletal muscle field is the role fusion proteins play. The recently discovered fusion protein Myomaker and its associated micropeptide Myomixer/Myomerger/Minion (hereafter referred to as Myomixer) are at the

centre of myogenic fusion research [23]. Thus, we were curious to examine the effects *Piezo1*

may have on these fusion proteins. *Piezo1* was first downregulated in EDL- and soleus-

derived myoblasts. Cells were then collected and RT-qPCR against *Myomaker* and *Myomixer*

was carried out and compared to control-siRNA conditions (Figure 4a-d). EDL-derived cells

(Figure 4a) showed that the expression of *Myomaker* is significantly reduced in *Piezo1*-

siRNA treated cells (~43%), compared to control-siRNA conditions (p<0.05). In *Piezo1*-

downregulated conditions, *Myomixer* (Figure 4b) showed a trend to decrease, compared to

control-siRNA, (p=0.0505). In soleus-derived cells (Figure 4c), downregulation of *Piezo1*

significantly lowered the expression of *Myomaker* (~30%), compared to control-siRNA

(p<0.05). Similar to the EDL, we observed a trend for *Myomixer* to decrease post *Piezo1*

knockdown (Figure 4d, p>0.05). Taken together, these results show a potential interplay

between the stretch induced Piezo1 channel and the fusion protein gene *Myomaker* (and

perhaps *Myomixer*). The question remains as to whether the downregulation of *Piezo1*

actively employs mechanisms which lower *Myomaker* expression which in turn reduce

myogenic fusion or whether this is an indirect effect of another *Piezo1* governed event.

Answering these queries may have implication in skeletal muscle regeneration and muscular

dystrophies.

Relative fold changes in expression of EDL-derived myoblasts (a and b) and soleus-derived myoblasts (c and d). Cells

were transfected with 10 nM of either control-siRNA (siScrambled) or *Piezo1*-siRNA. After overnight incubation, cells

incubated for a further 24 hours. The expression of the fusogenic protein genes *Myomaker* and *Myomixer* were then

analysed. Values were normalised to *Gapdh* and then expressed as fold change compared to siScrambled conditions.

Data is mean ± SEM from three experiments (n = 3 mice). *: Significant at $p < 0.05$ compared to siScrambled conditions

using a 2-tailed paired student t-test.

Figure 4. *Myomaker* expression is reduced in *Piezo1*-downregulated conditions

Ca^{2+} influx in Piezo1 downregulated myotubes

The Piezo1 channel permeates Ca^{2+} influx at a greater preference than other cations

(Na$^+$, K$^+$ and Mg^{2+}) [5]. Ca^{2+} is itself a crucial regulatory and signalling molecule of the

myogenic program as well as skeletal muscle contraction and various intracellular events.

Thus, a logical query to pursue was to see whether selective downregulation of *Piezo1*

hinders or blunts the influx of Ca^{2+} in cultured myotubes exposed to stretch. To achieve this

we used customised stretch silicon bio-chambers [16]. Myoblasts from both EDL and soleus

muscles were transfected with either *Piezo1*-siRNA or control-siRNA (siScrambled). The

transfection mixture was then removed, and samples were cultured in differentiation medium

for 3 days. The resulting myotubes were then subjected to incremental bouts of stretch.

Throughout the experiment, intracellular Ca^{2+} level [Ca^{2+}]i was measured using Fura-2AM, a

membrane-permeant fluorescent Ca^{2+} indicator (Figure 5).

Focusing first on control-siRNA, we found that [Ca^{2+}]i was increased at certain stretch

conditions. In EDL-derived myotubes for example (Figures 5a and 5c), a 6% stretch

significantly increased [Ca^{2+}]i influx compared to 0% stretch controls (p<0.05). An increase

in [Ca^{2+}]i was also observed in EDL-derived myotubes at 9% stretch conditions (p<0.05). A

9% stretch bout also showed significantly higher [Ca^{2+}]i, compared to 3% stretch conditions

(p<0.05). Soleus-derived myotubes (Figures 5b and 5d) at stretch conditions of 6% and 9%,

showed a significant increase in [Ca^{2+}]i, compared to 0% stretch conditions (p<0.05). Soleus-

derived myotubes in 9% stretch conditions also demonstrated higher [Ca^{2+}]i, compared to 3%

stretch bouts (p<0.05). Neither EDL- nor soleus-derived myotubes at the relatively lower

stretch bout of 3%, showed a significant increase in [Ca^{2+}]i, compared to 0% stretch controls.

This implies that a certain magnitude of physical stretch is necessary to elicit measurable

changes in [Ca^{2+}]i in cultured myotubes.

Turning our attention to the effects of *Piezo1* downregulation, we witnessed a striking

suppression in $[Ca^{2+}]i$ in response to stretch compared to control-siRNA (Figures 5a and 5b).

First of all, the progressive increase in $[Ca^{2+}]i$ was completely abolished in EDL-derived

myotubes (Figure 5c). Indeed neither a 6%, nor a 9% stretch could elicit a significant increase

in $[Ca^{2+}]i$. In soleus-derived cells (Figure 5d), only *Piezo1*-downregulated myotubes at the

9% stretch condition, were able to show increased $[Ca^{2+}]i$, compared to 3% stretch

counterparts ($p < 0.05$).

Comparison of $[Ca^{2+}]i$ between control-siRNA and *Piezo1*-siRNA cohorts at each

stretch condition further cemented the remarkable reduction of $[Ca^{2+}]i$ in *Piezo1*

downregulated conditions in response to stretch (Figure 5c). EDL-derived myotubes at 3%

stretch in *Piezo1*-siRNA conditions, show a suppression in $[Ca^{2+}]i$ compared to control-

siRNA counterparts ($p<0.05$). This is also true for 6% cohorts, with *Piezo1*-siRNA

transfected EDL myotubes displaying a lower $[Ca^{2+}]i$, compared to control-siRNA ($p<0.05$).

At 9% stretch we continued to see a significant suppression of increase in $[Ca^{2+}]i$ in *Piezo1*-

siRNA treated conditions, compared to control-siRNA at the same stretch bout ($p<0.05$). In

the soleus (Figures 5b and 5d), the suppression of $[Ca^{2+}]i$ following *Piezo1* downregulation

persisted. Indeed, a 3% stretch showed a lower $[Ca^{2+}]i$ in *Piezo1*-downregulated conditions,

compared to 3% stretch in control-siRNA. At 6% stretch, increase in $[Ca^{2+}]i$ is again blunted

in *Piezo1*-downregulated cohorts, compared to siRNA controls ($p<0.05$). This is further

demonstrated at 9% stretch bouts, with *Piezo1*-downregulated myotubes showing a reduced

increase in $[Ca^{2+}]i$, compared to control-siRNA- ($p< 0.05$). The suppression of increase in

$[Ca^{2+}]i$ following *Piezo1* downregulation may be a key factor in the reduction of fusion

observed in Figure 3.

EDL

a)

622 **c)** Relative intracellular Ca²⁺ level in EDL myotubes

d) Relative intracellular Ca²⁺ level in Soleus myotubes

Figure 5. *Piezo1* downregulation inhibits stretch-mediated increase of intracellular Ca^{2+}.

a) and b) Representative images of intracellular Ca^{2+} imaging in EDL and soleus-derived myotubes. Myoblasts were

transfected with 10 nM of either control-siRNA (siScrambled) (top panels) or *Piezo1*-siRNA (bottom panels). After

overnight incubation, cells were incubated for a further 72 hours. Fura 2-AM was administered to myotubes followed

by a 30-min incubation time. Stretch was then applied at 3% (0.3 mm), 6% (0.6 mm) and 9% (0.9 mm) for 1 min

followed by a 1-min resting period in between. Ionomycin at 5 μM was then applied. Side vertical bar shows Fura 2-

AM ratio emittance from low to high. c) and d) Average changes in the intracellular Ca^{2+} level ([Ca^{2+}]i) calculated by

difference between base and peak pixel value) normalised to ionomycin and control-siRNA at 0% stretch. Values are

mean ± SEM from three experiments (n = 3 mice). *: Significant difference at $p < 0.05$ compared to control-siRNA at

each condition. ¶: Significant difference at $p < 0.05$ compared to 0% stretch counterparts. †: Significant difference at p

< 0.05 compared to 3% stretch counterparts. Ŧ: Significant difference at $p < 0.05$ compared to 9% stretch in *Piezo1*-

siRNA conditions using one-way ANOVA followed by the Tukey-Kramer post-hoc.

Piezo1 activation and myotube fusion

The next phase of the investigation demanded we look at the effects of Piezo1 activation in myotube fusion and Ca^{2+} influx. This was achieved using the Piezo1 specific agonist Yoda1. Our preliminary findings showed that a 24-hour incubation of myotubes with all the chosen Yoda1 concentrations significantly inhibited myotube maturation (data not shown). Thus the subsequent experiments set out to test a series timepoints with our chosen Yoda1 concentrations, in order to uncover a potential "sweet-spot" between agonist concentration, incubation time and our hypothesised increase in myogenic fusion. Early forming myotubes (cells incubated in differentiation medium for 24 hours at high cell confluency in 96-well plates) were subjected to varying timepoints and concentrations of Yoda1. After each allocated timepoint, the agonist containing medium was removed and replenished with fresh differentiation medium and incubated for a further 2 days (Figure 6).

EDL-derived myotubes (Figures 6a and c) at the 1-minute timepoint showed that a 30 μM and 100 μM incubation with Yoda1 significantly increased the fusion index compared to DMSO controls. Similarly, soleus-derived myotubes (Figures 6b and d) had a significant increase in fusion at 30 μM and 100 μM, compared to DMSO controls ($p < 0.05$). At the 30 min timepoint, we found that EDL-derived myotubes at Yoda1 concentrations of 30 μM had a significantly higher fusion index, compared to DMSO counterparts ($p < 0.05$). Soleus samples incubated for 30 min with Yoda1 at 10 μM and 30 μM had a significantly higher fusion index than DMSO controls ($p < 0.05$). Unlike the 1 min timepoint, a 100 μM Yoda1 incubation revealed a reduced fusion index when incubated for 30 minutes. Indeed, both soleus and EDL samples incubated at 100 μM beyond the 1 min mark, showed a significant decrease in fusion, compared to DMSO controls.

At the 1-hour incubation timepoint, EDL-derived myotubes exhibited increased fusion

index at 5 μM, 10μM and 30 μM, compared to DMSO controls (p<0.05). Similarly soleus-

derived samples incubated with Yoda1 for 1 hour showed an increase in fusion at 5 μM, 10

μM–and 30 μM, compared to DMSO controls. Four hours of consistent Piezo1 activation

showed a significant decrease in fusion of EDL-derived myotubes at a Yoda1 concentration

of 30 μM compared to DMSO control (p<0.05). Taken together, we showed that Piezo1

activation results in enhanced fusion of myotubes. This is dependent on the duration and

potency in which Piezo1 is activated, as too much of either inhibits myotube fusion

Figure 6. Piezo1 activation increases myogenic fusion.

Early forming myotubes were administered with either DMSO (control, white bars) or the Piezo1-agonist Yoda1 at the

following concentration: 5 µM (red bars), 10 µM (blue bars), 30 µM (green bars) and 100 µM (grey bars). Myotubes

were incubated for 1 min, 10 min, 30 min, 1 hour and 4 hours. Following the incubation period, the medium was

exchanged with fresh reduced medium (without agonist) and myotubes were incubated for a further 2 days. a) and b)

Representative images of cohorts at relevant timepoints and concentrations, immunostained for Myosin heavy (MyHC)

(red panels) and counterstained with DAPI (black and white panels). Micrographs taken at x20 magnification. Scale

bar is 100 µm. Bar graphs display fusion index in c) EDL and d) soleus-derived myotubes from each time point and

concentration variables. Values are mean ± SEM. * indicates significance at $p < 0.05$ using one-way ANOVA tests

followed by the Tukey-Kramer post-hoc. n = 3 mice.

Myotube width and Piezo1 activation

We noticed that Yoda1 treated samples that exhibited increased fusion, also appeared

to have thinner myotubes. In order to address this, we compared the myotube width of these

samples to DMSO controls (Figure 7). Interestingly, both EDL (Figure 7a and b) and soleus

derived myotubes (Figure 7c and d) showed reduced myotube width compared to controls.

We further subdivided myotube width into bins of 5 μm increments and compared the

distribution of Yoda1 treated myotubes at the 1 min incubation with 100 μM concentration to

DMSO controls. Both EDL (Figure 7e) and soleus derived myotubes (Figure 7f), showed that

Yoda1 treated cells, on average, have a greater proportion of smaller myotubes. In contrast

DMSO controls have a higher distribution of lager myotubes. A similar pattern was observed

in the rest of the Yoda1 treated cohorts that displayed increased fusion (Figure S5). Taken

together it appears that increased fusion post Yoda1 administration comes at the cost of

myotube syncytial maturation.

Figure 7. Yoda1-mediated activation of Piezo1 decreases myotube width.

Representative images of a) EDL and c) soleus derived myotubes immunolabelled for MyHC. We measured the width

of the myotubes to quantify potential differences between Piezo1-activated cohorts (lower panels) and DMSO controls

(upper panels). This was achieved by taking three independent points within a field of view (yellow bars) and

measuring the distance from one side of the myotube to the other. Example pictures from DMSO controls and 100 μM

of Yoda1 incubated for 1 min are displayed. Scale bar is 100 μm. b) and d) Average myotube width in DMSO and

Piezo1 activated samples. e) and f) Proportion of myotube width in EDL and soleus-derived myotubes. Myotube width

was divided into incremental bins of 5 μm and represented as percentages relative to the total number of myotubes

counted. Data is mean ± SEM from three experiments data (n = 3 mice). *: Significant at $p < 0.05$ compared to DMSO

controls conditions using one-way ANOVA followed by the Tukey-Kramer post-hoc. Please note that only cohorts

which showed increased myogenic fusion in both EDL and soleus myotubes from Figure 6 are displayed.

Ca^{2+} influx and Piezo1 activation

The next aim of was to determine whether Piezo1 activation leads to increased influx

of Ca^{2+} in cultured myotubes. Using the customised stretch silicon bio-chambers [16], we

cultured myotubes derived from both EDL and soleus muscle. We then divided the samples

into two groups; those given the Piezo1 agonist Yoda1 (at 30 μM) and those without. The

chambers were then subjected to incremental bouts of stretch, with a minute rest in between

each stretch. Throughout the experiment we measured [Ca^{2+}]i (Figure 8).

First of all our results confirmed that increasing stretch amplitude results in higher

[Ca^{2+}]i, which means a greater influx of Ca^{2+}, in both EDL- and soleus-derived myotubes

(Figures 8c and 8d). Indeed under control conditions (without Yoda1), EDL-derived

myotubes subjected to 6% stretch and 9% stretch show a higher [Ca^{2+}]i, compared to 0%

stretch controls (p<0.05). Similarly, soleus-derived myotubes at a stretch of 6% and 9%

exhibited a higher [Ca^{2+}]i, compared to 0% stretch counterparts (p< 0.05). Notably EDL- and

soleus-derived myotubes under 3% of stretch showed no significant difference in [Ca^{2+}]i,

compared to 0% stretch counterparts. In contrast when Yoda1 was administered prior to

stretch, we find that the apparent "threshold" for Ca^{2+} influx is decreased. Indeed at 3%

stretch bouts (Figures 8c and 8d), we found that both EDL- and soleus-derived myotubes

showed a significantly higher [Ca^{2+}]i compared to their respective 0% stretch control

conditions (p<0.05). Soleus-derived myotubes (Figure 8d) also exhibited a greater increase in

[Ca^{2+}]i post Yoda1 administration in both 6% and 9% stretch conditions, compared to 0%

stretch controls (p<0.05). In a similar pattern (Figure 8c), EDL-derived myotubes at 6% and

9% stretch bouts in Yoda1 administered groups, showed a significant increase in [Ca^{2+}]i

compared to 0% controls (p<0.05).

Comparing the differences between agonist treated groups and those without, at each

stretch condition, we are able to further demonstrate the increased $[Ca^{2+}]i$ in response to

Piezo1 activation. For example at bouts of 3% stretch, EDL-derived myotubes treated with

Yoda1 (Figure 8c) showed a significantly higher $[Ca^{2+}]i$, compared to untreated 3% stretch

counterparts. At bouts of 6% stretch, EDL-derived myotubes given Yoda1 showed a

significantly higher $[Ca^{2+}]i$, compared to untreated myotubes ($p<0.05$). At 9% stretch, EDL-

derived myotubes treated with Yoda1, once again showed a significantly higher $[Ca^{2+}]i$,

compared to cohorts without the agonist ($p<0.05$). Soleus-derived myotubes at 3% stretch

(Figure 8d), showed that cohorts given Yoda1 prior, exhibited a higher $[Ca^{2+}]i$, compared to

untreated counterparts ($p<0.05$). At 6% stretch the significant difference in $[Ca^{2+}]i$ is also

evident, with Yoda1 treated myotubes showing a higher $[Ca^{2+}]i$, compared to untreated

samples ($p<0.05$). In line with these findings we showed that soleus-derived myotubes at

bouts of 9% stretch had a higher $[Ca^{2+}]i$ when given Yoda1, compared to myotubes without

the Piezo1 agonist ($p<0.05$). Taken together, this series of experiments showed that Piezo1 is

an essential player in the regulation and function of myotubes. This process seems to involve

active modulation of Ca^{2+} influx. How this leads to increased myotube fusion is an area of

great interest and one which demands further research.

Figure 8. Piezo1 activation increases the intracellular Ca²⁺ level in primary derived myotubes

a) and b) Representative images of Ca²⁺ imaging in EDL and soleus-derived myotubes. Fura 2-AM was administered

to myotubes followed by a 30-min incubation time. Stretch was then applied at 3% (0.3 mm), 6% (0.6 mm) and 9%

(0.9 mm) for 1 min followed by a 1-min resting period in between. During the initial 0% stretch timepoint, Yoda1

cohorts were administered with 30 μM of the agonist (red *) before being subjected to stretch. Ionomycin at 5 μM was

then applied. Side vertical bar shows Fura 2-AM ratio emittance from low to high. c) and d) Average changes in the

intracellular Ca^{2+} level ($[Ca^{2+}]i$,) difference between base and peak pixel value) normalised to ionomycin. Data is mean

± SEM from three experiments (n = 3 mice). *: Significant difference at $p < 0.05$ compared to control at each condition.

¶: Significant difference at $p < 0.05$ compared to 0% stretch counterparts. †: Significant difference at $p < 0.05$

compared to 3% stretch counterparts using one-way ANOVA followed by the Tukey-Kramer post-hoc.

Discussion

The current study showed that the mechanosensitive ion channel Piezo1 is required for the fusion and formation of myotubes. Indeed reduced expression of this channel hindered the fusion of myocytes to forming or maturating myotubes – a phenotype that may be partly associated with interacting fusogenic proteins like Myomaker or Myomixer. We further showcased how agonist mediated activation of Piezo1 can enhance myogenic fusion at the expense of myotube width. Additionally, our findings revealed that over-activation of Piezo1 results in the loss of myotube integrity. Using live imaging assays, we discovered that Piezo1 activation enhances Ca^{2+} permeation in cultured myotubes. This led to an increased influx of Ca^{2+} in response to stretch. Conversely, we showed how the downregulation of *Piezo1* hinders Ca^{2+} entry into myotubes of both EDL and soleus muscle.

Piezo1 in myogenesis

The current study investigated the effects of Piezo1 regulation throughout the myogenic program. *Piezo1* is expressed at a higher proportion in terminally differentiated myotubes, compared to proliferating myoblasts. Moreover, we found that, differentiating primary muscle cells derived from the mainly slow-type muscle soleus displayed higher expression of *Piezo1* compared to the fast EDL muscle (Figure S1). Understanding the potential differences in muscle/fibre types and *Piezo1* regulation is an intriguing area for future research and could reflect differences in the dynamics of myogenic progression. We also confirmed that the expression of *Piezo2* is not altered by the down-regulation of *Piezo1*. This is perhaps not surprising given the fact that *Piezo2* is not as abundant in skeletal muscle compared to *Piezo1* [5,16]. Nevertheless it was important to see any potential compensatory effects *Piezo2* may impose. Specific downregulation of *Piezo1* by siRNA-mediated transfection showed no significant change in the proliferation rate of either EDL- or soleus-

derived primary myoblasts. However, our data do not exclude the possibility that *Piezo1* is

not involved in earlier myogenic events, perhaps in balancing quiescence and activation of

satellite cells. In proliferating myoblasts, lack of Piezo1 function does not alter onset of

myogenic differentiation, evaluated by the proportion of myogenin-positive cells. In contrast,

where a significant phenotype was observed was in terminally differentiated myotubes.

Indeed, our data found that knockdown of Piezo1 significantly reduced fusion of myocytes

and prevented myotube formation and maturation. In contrast, activation of this Ca^{2+}

permeable channel resulted in enhanced myogenic fusion. The main findings are summarised

in Figure 9a-c.

Piezo1 and muscular dystrophies

Piezo1 activation showed a significant increase in the fusion index of both EDL- and

soleus-derived myotubes. Although this phenotype could be viewed as beneficial in terms of

muscular dystrophy prevention, we must be aware of the potential dangers of an overactive

Piezo1 channel. In fact, we showed that even a 30 min incubation of myotubes with a high

agonist concentration (100 µM of Yoda1 treatment) led to decreased fusion in both EDL- and

soleus-derived myotubes (Figure 6). This adverse phenotype is most likely the result of a

dangerously high Ca^{2+} influx. The publicly available data obtained from MRI-guided human

biopsies from facioscapulohumeral muscular dystrophy (FSHD) affected individuals (Figure

9f and g) further highlights the necessity to examine mechanosensor channels like Piezo1 in

muscular dystrophies. FSHD is associated with mutations in the distal end of chromosome 4

(4q35), resulting in epigenetic deregulation of the D4Z4 macrosatellite repeat array, causing

uncontrolled expression of the transcription factor *Double homeobox 4* (*DUX4*) [24-29]. In

turn, *DUX4* accumulation hampers myogenic progression promoting cytotoxicity and is

considered the most possible root cause of FSHD pathogenesis [30]. We took advantage of a

transcriptomic study on murine *DUX4*-inducible myoblasts model elucidating *DUX4* role in

FSHD, to assess the level of *Piezo1* in such context and found that it is significantly

downregulated upon accumulation of *DUX4* [31] (Figure 9d), thus suggesting that *Piezo1*

dysregulation may contribute to FSHD pathology.

In line with the muscular atrophy usually found in patients, human primary myoblasts

derived from FSHD biopsies display reduced differentiation *in vitro* generating hypotrophic

myotubes [32,33], thus resembling the effect of *Piezo1* knockdown in our experimental setup.

Indeed, analysis of transcriptomic dataset on differentiating isogenic FSHD affected or

unaffected myoblasts previously performed by Banerji and colleagues [33], revealed that

while during unaffected (healthy) myogenic differentiation, *Piezo1* is upregulated from

myoblasts to myotubes, its expression fails to increase in FSHD myogenic progression

(Figure 9e). Moreover, our analyses of transcriptomic dataset from Wang and colleagues [34]

found that *PIEZO1* mRNA is significantly upregulated in MRI-guided muscle biopsy from

FSHD patients that show the worst pathological phenotype (Group 4) (Figure 9f).

Interestingly, it has recently been demonstrated that FSHD muscles display traces of ongoing

regeneration, as indicated by elevated levels of embryonic Myosin, a canonical marker of

ongoing muscle fibre regeneration [24]. Congruently with the accumulation of *PIEZO1*

mRNA, additional transcriptomic analysis also revealed a concomitant upregulation of

MYOMAKER and *MYOMIXER* in the most affected group of muscle biopsies (Figure 9g)

indicating an ongoing regeneration process. Such regeneration is further suggested by

significant accumulation of embryonic *MYH3* (myosin 3) mRNA (Figure 9g). In contrast, the

reduced level of *MYBPC1* (myosin binding protein C1), a myosin-associated protein involved

in the formation and maintenance of sarcomere structure confirmed the dystrophic status of

FSHD muscle biopsies (Figure 9g). Therefore, from these pieces of evidence we conclude

that *Piezo1* dysregulation may contribute to the pathomechanism underlying FSHD. However,

the precise definition of *PIEZO1* dynamics in FSHD progression would require specific

investigation.

Figure 9. Piezo1 which regulates myogenic fusion is dysregulated in FSHD muscle pathology.

a) Under normal conditions, membrane mechanosensors, such as Piezo1 (magenta arrowhead), regulate myocyte fusion thereby ensuring efficient muscle fibre formation and maturation. **b)** Suppression of Piezo1 expression blunts myocyte fusion and Ca^{2+} homeostasis, thus hampering the formation of proper myotubes. **c)** In contrast, selective chemical activation of Piezo1 function enhances myogenic fusion thus significantly increasing myotube formation and maturation. d) Statistical analysis derived from Choi et al, [31], shows that DUX4 expression induced by Doxycycline, which mimics human FSHD pathogenesis, results in significant suppression of *Piezo1* level compared to control non-induced cells (Unpaired two-tailed t-test). e) Time-course transcriptomics analysis, derived from Banerji et al, [33], on differentiating isogenic (bar D4Z4 repeats) myoblasts derived from a male mosaic FSHD1 patient show that while unaffected (blue) myoblasts upregulated significantly *PIEZO1* expression when differentiating into myotubes, in FSHD *PIEZO1* level remains unchanged from myoblasts to myotubes (red bar) (Unpaired two-tailed t-test). f) Statistical analysis of transcriptomic data on human FSHD biopsies derived from Wang et al [34]. Patients have been grouped according to FSHD severity, which increases from Group1 to Group 4. Control (ctrl) represents non-affected individual. Count of gene reads (cpm, count per million) *PIEZO1* is significantly upregulated in FSHD Group 4. g) The same dataset as (f) shows concomitant upregulation of *MYOMAKER*, *MYOMIXER* and *MYH3* (myosin 3) while reduced *MYBPC1* (myosin binding protein C1) level in line with regeneration process ongoing and a severe dystrophic phenotype respectively. One-way ANOVA with Tukey's post-hoc test.

The skeletal muscle has typically a slow turnover, thus putative fusion proteins like

Myomaker and Myomixer regulated by Piezo1 may be present at a relatively lower level (as

shown by control patients, Figure 9d) and called upon in response to muscle damage (injury,

resistive exercise etc.). However in FSHD, the disorganised and possibly inefficient

activation of the regeneration program may dysregulate Piezo1 expression. From a

pathophysiological angle and taking into consideration our over-activation data using Yoda1

(Figure 6), we suggest that an over-abundance of Piezo1 expression in FSHD may result in

uncontrolled increase in Ca^{2+} influx, thus hindering myotube integrity. On the other hand,

improper Piezo1 expression may also lead to defective myogenic progression and muscle

formation, as indicated by our knockdown analysis. In line with this, transcriptomic analysis

of human myoblasts induced to express a high level of DUX4, which is considered to mimic

some aspects of FSHD pathogenesis, revealed significant suppression of *PIEZO1* expression

(Figure 9d), thus suggesting that reduced PIEZO1 function may contribute to FSHD

pathogenesis.

Piezo1 and the myocyte fusion machinery

Piezo1 downregulation significantly reduced myocyte fusion during myotube

formation and myotube maturation. To the best of our knowledge there is only one other

paper published that examined *Piezo1* in skeletal muscle by Tsuchiya et al. [17]. Interestingly,

the findings from this group showed that *Piezo1* inhibition resulted in a sheet-like syncytium

of MyHC coupled with increased fusion. Although these findings show contrasting results to

the ones presented in this study, we must take into consideration potential factors which may

explain why this may be the case. One such factor is the method of *Piezo1* inhibition used by

Tsuchiya et al, [17]. They carried out many of their experiments using knockout lines of

Piezo1 through the gene editing tool CRISPR/Cas9. The fact that these cell lines did not

express *Piezo1* to begin with (unlike the cells we used) may yield completely different

phenotypes, compared to the transient inhibition achieved by siRNA mediated transfection.

Therefore, complete lack of *Piezo1* expression may favour the activation of a secondary, yet

unknown, alternative *Piezo1*-affected pathway(s) to fusion. Regarding the *Piezo1* siRNA

transfection experiments, although more than 60% reduction in gene expression is ideal; we

nevertheless found that our level of *Piezo1* knockdown produced very interesting effects on

myogenic regulation. Moreover, Miyamoto et al, [16] also obtained slightly below or just

about 60% *Piezo1* reduction, yet reported intriguing *Piezo1*-associated events in urothelial

cells. Perhaps Tsuchiya et al, obtained even greater knockdown of *Piezo1* in their siRNA-

mediated analyses [17], further suggesting that the timing and level of *Piezo1* expression may

yield varying phenotypes. There was also the likelihood that siRNA used in this study may

potentially have off-target effects on other genes which could influence myotube formation

by employing other mechanistic pathways. However this is excluded by the fact that we have

now tested four different *Piezo1*-specific siRNAs and all show a marked reduction in the

fusion of myoblasts, compared to siRNA controls (Figure S4). Therefore, we conclude that

impaired Piezo1 signalling inhibits myotube fusion. Further support comes from our Piezo1

activation experiments. Congruently, we found that selective activation of Piezo1 by the

agonist Yoda1 significantly increased the fusion index in a concentration- and time-

dependent fashion. These results do not disprove the investigation by Tsuchiya et al, but

rather highlight the complexity underlying Piezo1 dynamics. It would be interesting to see the

effects of constitutive overexpression of *Piezo1* by retroviral transduction or conditional

Piezo1 transgenic mice. Furthermore, obtaining the cells used by Tsuchiya et al [17] and

conducting myocyte fusion analysis using our setup would undoubtedly prove vital at

discerning potential reasons underlying some of the seemingly opposing results.

Additional support for the involvement of Piezo1 in myogenic fusion comes from our

results which showed that its downregulation significantly reduced the expression of

Myomaker - a muscle specific protein that localises to the plasma membrane and is crucial

for vertebrate myocyte fusion [23,35,36]. Whether this is a direct response for the

downregulation of Piezo1 or an indirect event remains to be determined. In addition, the

expression level of *Myomixer*, another essential molecule for cell fusion, showed a tendency

to reduce upon *Piezo1* knockdown further suggesting that *Piezo1* contributes in the regulation

of the cell fusion machinery. Removal of either Myomaker or Myomixer results in decreased

myoblast fusion, whereas overexpression results in enhanced fusion [36,37].

A feature that became apparent in our study was the seemingly "thinner" myotubes in

agonist treated samples which showed increased fusion. Indeed, compared to DMSO controls,

the average myotube width was smaller in Yoda1 treated cohorts (Figure 7 and Figure S5).

Thus, we wonder whether the enhanced fusion, measured by the number of myonuclei within

MyHC positive myotubes, comes at the expense of myotube syncytial maturation. In other

words, Piezo1 activation may have hastened myonuclear fusion leading to premature

myotube formation. Perhaps increased PIEZO1 expression (Figure 9f) contributes to the

inadequate regenerative state of FSHD stricken muscle.

DMSO controls at the 4 hours mark showed a trending decrease in fusion in soleus-

derived myotubes, compared to DMSO counterparts from other timepoints. EDL-derived

DMSO-treated myotubes at the 4-hour incubation timepoint, showed a statistically significant

decline in fusion compared to DMSO counterparts. Although these differences may be

negligible at lower timepoints, we should warrant concern over the potential for DMSO-

mediated inhibition of primary derived myotubes during longer timeframes. Indeed, the

observed myotube inhibition may be slightly overestimated in Yoda1-treated samples at the

4-hour timepoint.

Piezo1 activation and Ca^{2+} influx

The present study showed that selective downregulation of *Piezo1* dramatically

suppressed [Ca^{2+}]i, which most likely translates in the depression of the influx of Ca^{2+} into

cultured myotubes exposed to stretch. In contrast we showed that activation of Piezo1

significantly increased [Ca^{2+}]i, which means the enhancement of Ca^{2+} influx. Our results

propose that Piezo1 is a novel intracellular Ca^{2+} regulatory protein in skeletal muscle function.

Ca^{2+} plays a crucial role in skeletal muscle function, maintenance and plasticity. All muscle

fibres use Ca^{2+} as their main regulatory and signalling molecule [38-40]. Therefore, the

contractile functionalities of muscle fibres are dependent on the highly regulated expression

of proteins involved in Ca^{2+} handling and signalling. Our study showed that Piezo1 mediated

regulation of Ca^{2+} influx is a key driving factor in the respective decrease and increase in

myotube fusion in response to Piezo1 inhibition and activation. To the best of our knowledge

this is the first time this has been demonstrated.

The silicon bio-chamber experiments revealed that at relatively low stretch conditions

of 3% (0.3 mm) neither EDL- nor soleus-derived myotubes elicited a significant increase in

Ca^{2+} influx. However at higher stretch distances (6% and 9% stretch) this mechanical barrier

was crossed as demonstrated by the net increase in [Ca^{2+}]i, compared to 0% stretch

counterparts. In a similar set of experiments (albeit using urothelial cells), Miyamoto et al.

also showed a distance dependent increase of Ca^{2+} influx. Interestingly this response was

blunted in *Piezo1*-siRNA-treated conditions [16]. The researchers also showed that a high

enough [Ca^{2+}]i must be attained in order to elicit a response, in their case ATP efflux. The

data presented in this study supports the presence of a stretch-dependent increase in Ca^{2+}

influx. Remarkably we found that activation of Piezo1 resulted in increased [Ca^{2+}]i at 3%

stretch conditions, suggesting that the activation threshold of Piezo1 was lowered.

Furthermore the data showed that reduction of *Piezo1* expression significantly blunted any

significant increase of $[Ca^{2+}]i$ in response to stretch. These results, for the first time show the

need for Piezo1 to respond to stretch and permeate Ca^{2+} into myotubes. The findings also

propose the presence of a physical threshold that must be attained before Piezo1 mediated

Ca^{2+} influx is significantly increased. Like Miyamoto et al [14], we find that a stretch-

dependent increase in Ca^{2+} influx is suppressed when *Piezo1* expression is decreased.

Conversely, we see an increase of $[Ca^{2+}]i$ when Piezo1 is activated. Whether this leads to

altered cellular/myotube viability in the form of ATP release remains a subject for future

research.

1061

Conclusion

The data presented in this study showed that the Piezo1 channel is present in primary

derived myoblasts and myotubes but expressed at a higher proportion in the latter.

Downregulation of *Piezo1* significantly lowered the fusion capacity during myotube

formation and maturation. In contrast, Piezo1 activation increased fusion. Future research

examining changes in myotube function (integrity, Ca^{2+} influx, cytoskeletal organisation and

fusion) that are directly the result of mechanical stress should consider analysis of Piezo1. In

the context of therapeutic strategies against muscular dystrophies such as FSHD, not only

must we unravel the spatiotemporal regulation of *Piezo1* expression, but we must be aware of

this channel's ability to alter it Ca^{2+} influx threshold by adapting or inactivating its gating

capacity in response to repetitive stimuli. Pharmaceutically, small activating molecules such

as Yoda1 (and others like it) may prove beneficial. However careful attention must be given

to the half-life and pharmacokinetics of these agonists *in vivo* before even considering them

as viable drugs for human consumption. Piezo1's importance in skeletal muscle maintenance

and function will undoubtedly grow as new research aims to explore the mechanisms and

signalling pathways this remarkable mechanosensor employs.

1078

1079

Funding

This work was supported by KAKENHI (JP17K01762, K.G.; JP18H03160, K.G.;

JP19K22825, K.G.; 19KK0254. K.G.) from the Japan Society for the Promotion of Science,

the Science Research Promotion Fund from the Promotion and Mutual Aid Corporation for

Private Schools of Japan, and Graduate School of Health Sciences, Toyohashi SOZO

University (K.G.).

1086

1087   **Acknowledgments**

1088   The authors thank Ms. Yumiko Asakura and from Department of Physiology, Graduate

1089   School of Health Sciences, Toyohashi SOZO University for her assistance.

1090

1091   **Conflicts of Interest**

1092   The authors declare that there are no conflicts of interest.

1093

References

1. Mauro A (1961) Satellite cell of skeletal muscle fibers. The Journal of biophysical and biochemical cytology 9:493-495

2. Zammit PS, Relaix F, Nagata Y, Ruiz AP, Collins CA, Partridge TA, Beauchamp JR (2006) Pax7 and myogenic progression in skeletal muscle satellite cells. Journal of cell science 119 (Pt 9):1824-1832

3. Zammit PS, Partridge TA, Yablonka-Reuveni Z (2006) The skeletal muscle satellite cell: the stem cell that came in from the cold. J Histochem Cytochem 54:1177-1191

4. Relaix F, Zammit PS (2012) Satellite cells are essential for skeletal muscle regeneration: the cell on the edge returns centre stage. Development (Cambridge, England) 139 (16):2845-2856

5. Coste B, Mathur J, Schmidt M, Earley TJ, Ranade S, Petrus MJ, Dubin AE, Patapoutian A (2010) Piezo1 and Piezo2 are essential components of distinct mechanically activated cation channels. Science (New York, NY) 330 (6000):55-60

6. Wang Y, Chi S, Guo H, Li G, Wang L, Zhao Q, Rao Y, Zu L, He W, Xiao B (2018) A lever-like transduction pathway for long-distance chemical- and mechano-gating of the mechanosensitive Piezo1 channel. Nature communications 9 (1):1300-1300

7. Cinar E, Zhou S, DeCourcey J, Wang Y, Waugh RE, Wan J (2015) Piezo1 regulates mechanotransductive release of ATP from human RBCs. Proc Natl Acad Sci U S A 112 (38):11783-11788

8. Ge J, Li W, Zhao Q, Li N, Chen M, Zhi P, Li R, Gao N, Xiao B, Yang M (2015) Architecture of the mammalian mechanosensitive Piezo1 channel. Nature 527:64

9. Sachs F (2010) Stretch-activated ion channels: what are they? Physiology (Bethesda) 25 (1):50-56

10. Saotome K, Murthy SE, Kefauver JM, Whitwam T, Patapoutian A, Ward AB (2017)

Structure of the mechanically activated ion channel Piezo1. Nature 554:481

11. Zhao Q, Zhou H, Li X, Xiao B (2018) The mechanosensitive Piezo1 channel: a three-

bladed propeller-like structure and a lever-like mechanogating mechanism. The FEBS journal

12. Zhao Q, Zhou H, Chi S, Wang Y, Wang J, Geng J, Wu K, Liu W, Zhang T, Dong M-Q,

Wang J, Li X, Xiao B (2018) Structure and mechanogating mechanism of the Piezo1 channel.

Nature 554:487

13. Coste B, Murthy SE, Mathur J, Schmidt M, Mechioukhi Y, Delmas P, Patapoutian A

(2015) Piezo1 ion channel pore properties are dictated by C-terminal region. Nature

Communications 6 (1):7223

14. Franco-Obregón A, Lansman J (1990) Calcium entry through stretch-in-activated ion

channels in MDX myotubes. Nature 344:670-673

15. Mercuri E, Bönnemann CG, Muntoni F (2019) Muscular dystrophies. Lancet 394

(10213):2025-2038

16. Miyamoto T, Mochizuki T, Nakagomi H, Kira S, Watanabe M, Takayama Y, Suzuki Y,

Koizumi S, Takeda M, Tominaga M (2014) Functional role for Piezo1 in stretch-evoked

$Ca(2)(+)$ influx and ATP release in urothelial cell cultures. J Biol Chem 289 (23):16565-

16575

17. Tsuchiya M, Hara Y, Okuda M, Itoh K, Nishioka R, Shiomi A, Nagao K, Mori M, Mori

Y, Ikenouchi J, Suzuki R, Tanaka M, Ohwada T, Aoki J, Kanagawa M, Toda T, Nagata Y,

Matsuda R, Takayama Y, Tominaga M, Umeda M (2018) Cell surface flip-flop of

phosphatidylserine is critical for PIEZO1-mediated myotube formation. Nature

Communications 9 (1):2049

18. Collins CA, Zammit PS (2009) Isolation and grafting of single muscle fibres. Methods in

molecular biology (Clifton, NJ) 482:319-330

19. Lacroix JJ, Botello-Smith WM, Luo Y (2018) Probing the gating mechanism of the

mechanosensitive channel Piezo1 with the small molecule Yoda1. Nature Communications 9

(1):2029

20. Nourse JL, Pathak MM (2017) How cells channel their stress: Interplay between Piezo1

and the cytoskeleton. Seminars in cell & developmental biology 71:3-12

21. McHugh BJ, Murdoch A, Haslett C, Sethi T (2012) Loss of the integrin-activating

transmembrane protein Fam38A (Piezo1) promotes a switch to a reduced integrin-dependent

mode of cell migration. PLoS One 7 (7):e40346-e40346

22. Schnorrer F, Dickson BJ (2004) Muscle building; mechanisms of myotube guidance and

attachment site selection. Dev Cell 7 (1):9-20

23. Millay DP, O'Rourke JR, Sutherland LB, Bezprozvannaya S, Shelton JM, Bassel-Duby R,

Olson EN (2013) Myomaker is a membrane activator of myoblast fusion and muscle

formation. Nature 499 (7458):301-305

24. Banerji CRS, Henderson D, Tawil RN, Zammit PS (2020) Skeletal muscle regeneration

in facioscapulohumeral muscular dystrophy is correlated with pathological severity. Human

Molecular Genetics

25. Bosnakovski D, Xu Z, Gang EJ, Galindo CL, Liu M, Simsek T, Garner HR, Agha-

Mohammadi S, Tassin A, Coppee F, Belayew A, Perlingeiro RR, Kyba M (2008) An

isogenetic myoblast expression screen identifies DUX4-mediated FSHD-associated

molecular pathologies. EMBO J 27

26. Deidda G, Cacurri S, Piazzo N, Felicetti L (1996) Direct detection of 4q35

rearrangements implicated in facioscapulohumeral muscular dystrophy (FSHD). J Med Genet

33

27. Ferreboeuf M, Mariot V, Bessieres B, Vasiljevic A, Attie-Bitach T, Collardeau S, Morere

J, Roche S, Magdinier F, Robin-Ducellier J, Rameau P, Whalen S, Desnuelle C, Sacconi S,

Mouly V, Butler-Browne G, Dumonceaux J (2014) DUX4 and DUX4 downstream target

genes are expressed in fetal FSHD muscles. Hum Mol Genet 23

28. Tassin A, Laoudj-Chenivesse D, Vanderplanck C, Barro M, Charron S, Ansseau E, Chen

YW, Mercier J, Coppee F, Belayew A (2013) DUX4 expression in FSHD muscle cells: how

could such a rare protein cause a myopathy? J Cell Mol Med 17

29. Tawil R, van der Maarel SM, Tapscott SJ (2014) Facioscapulohumeral dystrophy: the

path to consensus on pathophysiology. Skeletal Muscle 4 (1):12

30. Himeda CL, Jones PL (2019) The Genetics and Epigenetics of Facioscapulohumeral

Muscular Dystrophy. Annual Review of Genomics and Human Genetics 20 (1):265-291

31. Choi SH, Bosnakovski D, Strasser JM, Toso EA, Walters MA, Kyba M (2016)

Transcriptional Inhibitors Identified in a 160,000-Compound Small-Molecule DUX4

Viability Screen. J Biomol Screen 21 (7):680-688

32. Barro M, Carnac G, Flavier S, Mercier J, Vassetzky Y, Laoudj-Chenivesse D (2010)

Myoblasts from affected and non-affected FSHD muscles exhibit morphological

differentiation defects. Journal of cellular and molecular medicine 14 (1-2):275-289

33. Banerji CRS, Panamarova M, Pruller J, Figeac N, Hebaishi H, Fidanis E, Saxena A,

Contet J, Sacconi S, Severini S, Zammit PS (2019) Dynamic transcriptomic analysis reveals

suppression of PGC1α/ERRα drives perturbed myogenesis in facioscapulohumeral muscular

dystrophy. Hum Mol Genet 28 (8):1244-1259

34. Wang LH, Friedman SD, Shaw D, Snider L, Wong C-J, Budech CB, Poliachik SL, Gove

NE, Lewis LM, Campbell AE, Lemmers RJFL, Maarel SM, Tapscott SJ, Tawil RN (2018)

MRI-informed muscle biopsies correlate MRI with pathology and DUX4 target gene

expression in FSHD. Human Molecular Genetics 28 (3):476-486

35. Ganassi M, Badodi S (2018) Myogenin promotes myocyte fusion to balance fibre number

and size. 9 (1):4232

36. Shi J, Cai M, Si Y, Zhang J, Du S (2018) Knockout of myomaker results in defective

myoblast fusion, reduced muscle growth and increased adipocyte infiltration in zebrafish

skeletal muscle. Human Molecular Genetics 27 (20):3542-3554

37. Goh Q, Millay DP (2017) Requirement of myomaker-mediated stem cell fusion for

skeletal muscle hypertrophy. 6

38. Berchtold MW, Brinkmeier H, Muntener M (2000) Calcium ion in skeletal muscle: its

crucial role for muscle function, plasticity, and disease. Physiological reviews 80 (3):1215-

1265

39. Gehlert S, Bloch W, Suhr F (2015) Ca2+-dependent regulations and signaling in skeletal

muscle: from electro-mechanical coupling to adaptation. Int J Mol Sci 16 (1):1066-1095

40. Allen DG, Lamb GD, Westerblad H (2008) Skeletal muscle fatigue: cellular mechanisms.

Physiological reviews 88 (1):287-332

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

Supplementary figure 1. ***Piezo1* is expressed at a higher level in soleus compared to EDL muscle-derived myoblasts.**

Relative fold changes in expression of *Piezo1* in myoblasts from EDL (green bars) and Soleus muscle (blue bars), during proliferation and through differentiation; 24 hours (Day 1) and 72 hours (Day 3) in differentiation medium. Values were normalised to *Gapdh* and then expressed as fold change compared to levels of proliferation in EDL samples. Data is presented as mean ± SEM from three experiments (n = 3 mice). Asterisks (*) denote significance at $p < 0.05$ compared to control proliferation conditions. ¶: Significant difference at $p < 0.05$ significant compared to either day 1 or day 3 differentiation cohorts using one-way ANOVA followed by the Tukey-Kramer post-hoc.

Supplementary Figure 2. *Piezo1* knockdown reduced f-actin intensity in EDL and soleus-derived myotubes

a) and b) Representative images of EDL and soleus muscle--derived myotubes. Early forming myotubes were transfected with 10nM of control-siRNA (siScrambled) or *Piezo1*-siRNA. Following overnight incubation, cells incubated for a further 72 hours. The cytoskeleton protein f-actin was visualised using fluorescently labelled phalloidin (green panels). Nuclei were counterstained with DAPI (black and white panels). Scale bar is 100 μm. c) and d) Overall fluorescence intensity measured by pixel/area in each field of view (six images per conditions). e) and f) is data from c) and d) expressed as percentages relative to the maximum pixel value. Data is mean ± SEM from three experiments (n = 3 mice). *: Significant at $p < 0.05$ compared to siScrambled conditions using a 2-tailed paired student t-test.

Supplementary figure 3. Yoda1-mediated activation of Piezo1 does not increase f-actin intensity

Early forming myotubes were administered with either DMSO (control, white bars) or Yoda1 at the following

concentration: 5 μM (red bars), 10 μM (blue bars), 30 μM (green bars) and 100 μM (grey bars). Myotubes were

incubated for 1 min, 10 min, 30 min, 1 hour and 4 hours. Following the incubation period, the medium was exchanged

with fresh reduced medium (without agonist) and myotubes were incubated for a further 2 days. b) and d) Representative

images at relevant timepoints and concentrations treated with fluorescently labelled phalloidin (green panels and

counterstained with DAPI (black and white panels). Images taken at x20 magnification. Scale bar is 100 μm. Bar graphs

display proportion of f-actin intensity in a) EDL and b) soleus derived myotubes from each time point and concentration

variables. Values are mean ± SEM. * indicates significance at $p < 0.05$ using one-way ANOVA tests followed by the

Tukey-Kramer post-hoc. n = 3 mice.

Supplementary figure 4. Different *Piezo1* siRNA targets continue to show decrease in myotube fusion

a) Location of siRNAs shown in Table 1 on *Piezo1* mRNA. Diagram shows *Piezo1* mRNA transcript (orange bar),

coding sequence (blue bar) and location of each siRNA (red line). bp: base pair. Diagram is adapted from Qiagen

(https://geneglobe.qiagen.com/product-groups/flexitube-sirna). b and c) Representative images of EDL and soleus

muscle-derived myotubes, transfected with 10nM of control-siRNA (siScrambled) or different siRNAs specific for

Piezo1 (*Piezo1*-siRNA 1 and *Piezo1*-siRNA 2). A mixture of four different *Piezo1*-siRNAs (*Piezo1*-siRNA Mix)

including the one in the main text (Table 2) was also used at 10nM (2.5 nm each). Myoblasts were transfected and

incubated overnight; cells were incubated for a further 72 hours. Cells were immunolabelled for Myosin heavy chain

(MyHC) (red panels) and counterstained with DAPI (black and white panels). d) and e) The fusion index was

calculated by counting the total number of nuclei within each myofibre and representing this as a percentage relative to

the total number nuclei in the image taken. Data is mean ± SEM from three experiments (n = 3 mice). *: Significant at

$p < 0.05$ compared to siScrambled conditions using one-way ANOVA followed by the Tukey-Kramer post-hoc. From

table 2, Piezo1 siRNA 1 corresponds to Qiagen, S104420402. Piezo1 siRNA 2 corresponds to Qiagen, S100814807.

1334

Supplementary Figure 5. Yoda1-mediated activation of Piezo1 decreases myotube width.

Myotube width was divided into incremental bins of 5 μm and represented as percentages relative to the total number

of myotubes counted. The overall proportion of myotube width distribution in DMSO controls and Yoda1-treated

samples is summarised as line graphs in a) EDL and b) soleus-derived myotubes. c-l) Bar graphs comparing DMSO

controls and samples which showed increased fusion post Yoda1 treatment in EDL (left) and soleus (right)-derived

myotubes Data is mean ± SEM from three experiments (n = 3 mice). *: Significant at $p < 0.05$ compared to DMSO

controls conditions using one-way ANOVA followed by the Tukey-Kramer post-hoc. Please refer to Figure 6 a) and b)

for representative images.

1369