A *cis*-regulatory atlas in maize at single-cell resolution

Alexandre P. Marand[1]*, Zongliang Chen[2], Andrea Gallavotti[2,3], and Robert J. Schmitz[1]*

Affiliations:
[1]Department of Genetics, University of Georgia, Athens, GA 30602, USA
[2]Waksman Institute, Rutgers University, Piscataway, NJ, 08854, USA
[3]Department of Plant Biology, Rutgers University, New Brunswick, NJ, 08901, USA

Correspondence: Alexandre P. Marand (marand@uga.edu), Robert J. Schmitz (schmitz@uga.edu)

ABSTRACT

***Cis*-regulatory elements (CREs) encode the genomic blueprints for coordinating spatiotemporal gene expression programs underlying highly specialized cell functions. To identify CREs underlying cell-type specification and developmental transitions, we implemented single-cell sequencing of Assay for Transposase Accessible Chromatin in an atlas of *Zea mays* organs. We describe 92 distinct states of chromatin accessibility across more than 165,913 putative CREs, 56,575 cells, and 52 known cell-types in maize using a novel implementation of regularized quasibinomial logistic regression. Cell states were largely determined by combinatorial accessibility of transcription factors (TFs) and their associated binding sites. Co-accessible chromatin recapitulated higher-order chromatin interactions, with distinct sets of TFs coordinating cell type-specific regulatory dynamics. Pseudotime reconstruction and alignment with *Arabidopsis thaliana* trajectories identified conserved TFs, associated motifs, and *cis*-regulatory regions specifying sequential developmental progressions. Cell-type specific accessible chromatin regions were enriched with phenotype-associated genetic variants and signatures of selection, revealing the major cell-types and putative CREs targeted by modern maize breeding. Collectively, our analysis affords a comprehensive framework for understanding cellular heterogeneity, evolution, and *cis*-regulatory grammar of cell-type specification in a major crop species.**

MAIN TEXT

Multicellular eukaryote development relies on precise spatiotemporal gene expression mediated

by *cis*-regulatory elements (CREs)[1-7]. Despite the economic and nutritional importance of crops

to society, our understanding of cell-type functions in plants has been precluded by technical

limitations imposed by the cell wall and an inability to culture homogenous cell lines. To

circumvent these limitations and comprehensively assess *cis*-regulatory variation among cell-

types in a major crop species, we isolated nuclei using fluorescence-activated nuclei sorting

(FANS) and generated single-cell accessibility profiles using Assay for Transposase Accessible

Chromatin (scATAC-seq) from six major *Zea mays* L. cultivar B73 organs, including axillary buds,

staminate and pistillate inflorescence, whole seedling (composed of stem, leaf, and coleoptile

tissues), embryonic root tips, and post-embryonic crown roots (**Fig. 1a**, **Extended Data Fig. 1**,

Supplementary Table 1). Rigorous evaluation of quality metrics, including doublet analysis via

genotype mixing (B73/Mo17), biological replicates, per nucleus read-depth, transcription start site

(TSS) enrichment, and fragment length distribution were reflective of high-quality scATAC-seq

data (Methods, **Extended Data Fig. 1, 2**). In total, we identified 56,575 high-quality nuclei with an

average of 31,660 unique Tn5 integration sites per nucleus (**Supplementary Data 1**). To group

nuclei with similar global patterns of accessibility, we developed a regularized quasibinomial

logistic regression framework that removes technical variation stemming from differences in nuclei

read depth by pooling information across accessible chromatin regions (ACRs) and extracting

model residuals as normalized accessibility scores for each nucleus and ACR (Methods).

Denoising the residual matrix with singular value decomposition (SVD) and projecting the

components into a two-dimensional space using Uniform Manifold Approximation Projection

(UMAP) revealed 10 clusters generally composed of nuclei from the same organ, highlighting

functional diversification of spatially distinct cells (**Fig. 1b, c; Extended Data Fig. 3a-e**). A second

round of clustering for each major grouping uncovered a total of 92 sub-clusters (hereafter

Figure 1: Atlas-scale profiling of single nuclei chromatin accessibility in *Zea mays*

a, Overview of experimental protocol. Homogenized samples release nuclei that are extracted by flow cytometry. Accessible chromatin is tagged by Tn5 and individual nuclei are distinctly barcoded in disparate droplets following microfluidic separation, allowing droplet pooling and library amplification prior to sequencing. **b,** Nuclei similarity clustering as a UMAP embedding derived from the denoised quasibinomial residuals across all ACRs for each nucleus. UMAP embedding of nuclei colored by **b,** organ identity and **c,** major cluster identity. **d,** UMAP embedding of sub-cluster assignments following a second round of clustering within each major cluster. **e,** Cell-type specific enrichment of gene accessibility for a subset of marker genes associated with six different cell-types. **f,** Sub-cluster specific chromatin accessibility profiles surrounding known marker genes for floral primordia, xylem precursors, and L1 epidermal cells. Bold, circled numbers indicate the cognate major cluster shown in panel **c**. Sub-cluster numeric identifications are present on the sides of the coverage plots. **g,** Top, gene accessibility for *ZmGRFTF36*, an inflorescence and spikelet meristem enriched TF with no previously known cell-type specificity. Bottom, RNA in situ hybridization of *ZmGRFTF36* in maize B73 staminate (tassel) primordia. FP, floral primordia; GP, glume primordia; IM, inflorescence meristem; LFM, lower floral meristem; SM, spikelet meristem; SPM, spikelet pair meristem; TF, transcription factor; UKN, unknown.

referred to as clusters) with an average of 551 nuclei and 165,913 reproducible ACRs covering

~4% of the maize genome (**Fig. 1d**).

To annotate clusters with corresponding cell-types, we integrated gene and promoter (1-

kb upstream TSSs) chromatin accessibility information with long-range distal ACRs on the basis

of co-accessibility (Methods) into a single accessibility score for each gene and nucleus as a

proxy for gene expression (bulk RNA-seq versus aggregate scATAC-seq Spearman's correlation

coefficient = 0.54-0.58, **Extended Data Fig. 3f**). We then (*i*) evaluated differential accessibility

among clusters for a manually curated list of 221 known marker genes (**Supplementary Table**

2), (*ii*) classified cell-types of individual nuclei with a multinomial logistic classifier trained on nuclei

with discriminative cell-type specific signatures, and visually assessed (*iii*) accessibility scores of

a priori marker genes over the UMAP embeddings and (*iv*) cluster-aggregated ATAC-seq

coverages (Methods, **Fig. 1e**,**f, Extended Data Fig. 4a**, **Supplementary Table 3**). Patterns of

gene accessibility were consistent with *a priori* information of cell-type/domain specific

expression, such as overlapping accessibility of bundle sheath-specific genes *DICARBOXYLIC*

ACID TRANSPORTER1 (*DCT2*) and *RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL*

SUBUNIT2 (*SSU2*) in bundle sheath cells, and mesophyll-specific genes *MALATE*

DEHYDROGENASE6 (*MDH6*) and *PYRUVATE DEHYDOGENASE KINASE1* (*PDK1*) in

mesophyll cells (**Fig. 1e, f, Extended Data Fig. 4a**)[8]. To further corroborate predicted cell-type

annotations, we performed RNA in situ hybridization for a subset of differentially accessible genes

with no prior evidence of cell-type specificity. In all cases, expression patterns were in line with

the predicted localization based on gene accessibility (**Fig. 1g**, **Extended Data Fig. 4b**).

Estimates of cell-type proportions within and across organs were also concordant with prior

observations, such as the occurrence of vascular bundle sheath and parenchymal mesophyll cells

within multiple organs, including those derived from stem and leaf tissues of seedlings and within

pistilate and staminate inflorescence (**Extended Data Fig. 4c**, **Supplementary Table 3**)[9].

Analysis of chromatin accessibility variation at genic loci revealed 74% (28,625/38,752) of

genes with significant differential accessibility (FDR < 0.05, log fold-change > 2), with an average

of 2,768 differentially accessible genes per cluster (**Extended Data Fig. 5b, Supplementary**

Data 2). Marker-agnostic gene set enrichment analysis (GSEA) of Gene Ontology (GO) terms

exemplified prior information regarding specific cell-type functions, with enriched terms such as

"root hair cell development" in root epidermal initials, "regulation of stomatal closure" in subsidiary

cells, and "malate transmembrane transport" for mesophyll cells (**Extended Data Fig. 4d**).

Distinct cell-types were generalized by highly specific GO annotations, as most (>51%) GO terms

were identified in only a handful of cell-types (five or fewer), implicating accessibility dynamics as

underling the signature hallmarks of cell-type identity and function (**Extended Data Fig. 4d**). In

summary, this combinatorial approach enabled the identification of 52 cell-types for 83% (76/92)

of clusters, capturing nearly all major expected cell-types in the profiled organs and suggesting

the existence of novel uncharacterized cell-types present in these data (**Supplementary Table**

3).

Characterization of chromatin accessibility variation

Deconvolution of nuclei into discrete cell-types provides an opportunity to identify CREs encoding

cell identity. To this end, we implemented a generalized linear model to catalog ACRs with

discrete patterns of chromatin accessibility across cell-types (Methods). In total, 52,520 ACRs

(31%) were differentially accessible (FDR < 0.05, fold-change > 2) and restricted to one or a

handful of clusters, with an average of 2,826 per-cluster (**Extended Data Fig. 5a**, **6a**, **b**). Similar

to the reported functions of transcriptional enhancers in metazoan genomes, we found cluster-

specific ACRs were associated with significantly greater enhancer activity based on STARR-seq

of maize leaf protoplasts[10] relative to controls (Methods, n=165,913) and non-specific (n=113,393)

ACRs (**Fig. 2a**). Deconvolution of chromatin accessibility by cell-type revealed accessible sites

primarily located distal to genic regions (>2-kb from any gene) compared to previously published

Figure 2: Characterization of accessible chromatin regions

a, Distribution of enhancer activity (maximum log2[RNA/input]) for control regions (n=165,913), non-specific (n=113,393) and cluster-specific ACRs (n=52,520). The dash red line indicates the overall mean. Orange lines reflect differences between the group mean and overall mean. **b,** Distribution of ACR distances to the nearest gene. Inset, distribution of ACR genomic context. **c,** Relative DNA methylation levels 2-kb flanking ACRs. **d,** Relative enrichment of polymorphisms after normalizing by mappability for 5-kb regions flanking cell-type specific and non-specific ACR summits. Smoothed splines are shown as dark lines. **e,** Relative enrichment of significant GWAS polymorphisms relative to all polymorphisms for 5-kb regions flanking cell-type specific and non-specific ACR summits. **f,** Enrichment of signatures of selection (XP-CLR) in the top 2,000 ACRs for all cell-type clusters. The 20 most enriched cell-types are highlighted on the left. AM, axillary meristem; CC, companion cell; FM, floral meristem; FP, floral primordia; IM, inflorescence meristem; QC, quiescent center; SM, spikelet meristem; SPM, spikelet pair meristem. **g,** Average motif coverage for all ACRs (n=165,913) and control regions (n=165,913). Shaded polygon, 95% confidence intervals. **h,** Enrichment of TF motifs in the top 2,000 ACRs ranked by Z-score for each cell-type compared to the top 2,000 most constitutive ACRs via binomial tests. FDR was estimated by the Benjamini-Hochberg method. **i,** Comparative heatmaps for matched TF gene accessibility (bottom) and motif deviation (top) Z-scores across clusters. **j,** Gene accessibility scores for five maize transcription factors (top) and their associated motif deviations (bottom).

bulk-level experiments (**Fig. 2b**)[10]. Notably, 30% (22,456/73,791) of distal ACRs overlapped with

LTR transposons, including the major maize domestication locus *tb1*-enhancer, and were

generally devoid of DNA methylation[11,12]. These findings are consistent with transposable

elements playing a prominent role in CRE evolution of the maize genome (**Fig. 2b, c, Extended**

Data Fig. 1a)[13-15].

Sequence variation underlying CREs contribute to disease emergence and phenotypic

innovation over evolutionary timescales[16,17]. In contrast to broadly accessible regions, analysis of

extant genetic variation in maize revealed lower polymorphism rates within cell-type specific

ACRs (**Fig. 2d**). However, of the genetic variants embedded within ACRs, those within cell-type

specific ACRs were more frequently associated with phenotypic variation[18] (**Fig. 2e**). To

investigate the contribution of domestication and signatures of selection in distinct cell-type

contexts, we assessed the relative enrichment of multi-locus allele-frequency differentiation signal

(XP-CLR scans) from chronologically sampled elite inbred maize lines within cell-type specific

ACRs[19]. Of the 21 cell-types with significant (FDR < 0.01) selection signature enrichment, 57%

(12) correspond to staminate and pistillate cell-types, such as spikelet meristems, spikelet pair

meristems, inflorescence meristems, floral meristems, and floral primordia (**Fig. 2f**). Thus, modern

maize breeding resulted in the selection of alleles containing floral-specific ACRs associated with

agronomically favorable inflorescence architecture[20].

To characterize the determinants of chromatin accessibility variation among cell-types, we

mapped the genome-wide occurrences of 570 transcription factor (TF) motifs. Expectedly, ACRs

were highly enriched with putative TF binding sites relative to control (n=165,913) and flanking

regions (**Fig. 2g**). Using the top 2,000 differential ACRs for each cell-type, we found 84% of TF

motifs (475/568) were enriched (binomial test: FDR < 0.05, Methods) in at least one type, with a

median of 43 enriched TF motifs per cell-type (**Fig. 2h**). To characterize TF motif accessibility

variation at the scale of individual nuclei, we estimated relative TF motif enrichment controlling for

background and sequence bias across ACRs for each nucleus (**Extended Data Fig. 6c-f**).

Diverse combinatorial TF motif landscapes were apparent across cell-types and closely mirrored

gene accessibility profiles from matched TFs (median PCC across cell-types = 0.45), establishing

synchronized chromatin accessibility of *cis* and *trans* cell autonomous factors as major

determinants of cell identity (**Fig. 2i, j**, **Extended Data Fig. 6g**). Classification of enriched TFs

and their cognate motifs identified several known cell-type specific regulators – including *WRKY*

family TFs in root epidermal progenitors and trichoblasts[21], *G2-like1* in parenchymal mesophyll[8],

and *AGAMOUS-like* and *SEPALLATA*[22] TFs in floral primordia – as well as previously

uncharacterized TFs with new potential roles as cell-type regulators (**Fig. 2i, j**, **Supplementary**

Table 4, 5). Taken together, these results are indicative of robust inference of CRE and TF activity

at the level of single cells and reveal TF dynamics central to *cis*-regulatory specification of diverse

cell states.

Coordinated dynamic chromatin accessibility recapitulates *in vivo* chromatin interactions

Correlated changes in chromatin accessibility of nearby loci represent putative physical chromatin

interactions with regulatory potential[1,23-26]. We identified 3.8 million (M) ACR-ACR linkages

(hereafter referred to as co-accessible ACRs) with significantly correlated (empirical FDR < 0.05,

Methods) patterns of chromatin accessibility across cell-types, capturing known physical

interactions such as between *TB1* and the genetically mapped domestication *tb1*-enhancer that

confers inflorescence/axillary meristem-specific expression (**Fig. 3a**, **Extended Data Fig.**

7a)[10,14,27,28]. To assess the broad interactive potential of co-accessible ACRs *in vivo*, we compared

co-accessible ACRs from leaf cell-types with maize whole-leaf chromatin conformation capture

data, recovering more than 78% (3,313/4,265), 57% (37,712/65,691) and 44% (17,108/38,567)

of chromatin loops from Hi-C, H3K4me3-HiChIP and H3K27me3-HiChIP experiments,

respectively (**Extended Data Fig. 7b**). Hi-C/HiChIP is a direct reflection of the proportion of cells

exhibiting a particular interaction, as ubiquitously interacting loci dominate rarer cell-type specific

contexts that are frequently missed by loop-calling algorithms. To determine the relative

Figure 3: Co-accessible ACRs reflect *in vivo* chromatin interactions and are established by cell-type specific TFs.
a, Comparison of ear and seedling co-accessible ACRs across nine cell-types at the genetically mapped *tb1-enhancer* domestication locus (highlighted region). Link height reflects co-accessibility scores between ACRs across cells in a cluster. **b**, Average normalized Hi-C signal across 4-kb windows centered on ACRs equally distributed into three groups (each group: n=54,904) based on the number and strength of participating co-accessible links. **c**, Co-accessible ACR interaction frequency across clusters. Inset: Single cell-type co-accessible links (n=1,018,417) for different genomic contexts. **d**, Standardized (row-wise) number of connections for each ACR (rows) by cell-type (columns). Column color map reflect cell-types from the legend in Extended Data Fig. 4. Gene-proximal ACRs for a subset of marker genes are indicated on the right. **e**, Right, chromatin accessibility and co-accessible ACR links surrounding the *UB2* locus associated with ear row number and tassel branch number quantitative traits. Black arrow indicates a distal ACR upstream of *UB2* present only in spikelet meristems. Co-accessible links with an edge within 2-kb of *UB2* are colored pink while remaining links are grey. Link height represents co-accessibility strength. Left, close-up of accessibility profiles of *UB2*. **f**, Distributions of average ACR-ACR links across cell-types for ACRs that overlap (purple) and do not overlap (grey) phenotype-associated genetic variants from maize GWAS. The median of each distribution is shown as a white horizontal line. Violin plots present the entire range of average number of connections on a log scale. Hypothesis testing was conducted within the *R* statistical framework via Wilcoxon rank sum test. **g,** Distributions of average ACR-ACR links celltypes for ACRs with (blue) and without (grey) enhancer activity (log2 RNA/input greater than 0). Hypothesis testing and distribution illustration was performed similarly as panel **f**.

predictability of *in vivo* interactions using co-accessible ACRs, we estimated the interaction

strength of each ACR by integrating correlative scores across cell-types (Methods). ACRs

classified by interaction strength recapitulated expected *in vivo* chromatin interaction frequencies,

where even the weakest class of co-accessible ACRs were associated with elevated interaction

frequencies relative to flanking and randomized control regions (**Fig. 3b**).

Cataloguing the usage of co-accessible ACRs across cell-types identified more than 27%

(~1M in total) that were unique to a single type, 49% of which were classified as distal-

genic/proximal, with an average of 11,069 distinct links per cell-type (**Fig. 3c**). Consistent with

regulatory models where a single gene can interact with multiple distant loci, proximal ACRs,

rather than distal or genic ACRs, were associated with the greatest number of links on average

(Wilcoxon rank sum test: *P* < 2.2e-308, **Extended Data Fig. 7c**). Highlighting long-range "hub"

interactions as a key contributors towards cell identity, cell-type specific co-accessible ACRs were

associated with greater number of links per site (Wilcoxon rank sum test: *P* < 2.2e-308) and a

greater proportion of links involving distal ACRs (Chi-squared test: *P* < 2.2e-308, **Extended Data**

Fig. 7c, d). Furthermore, the interactive capacity of any given ACR strongly depended on the cell-

type context (**Fig. 3d**). For example, *UNBRANCHED2* (*UB2*) – a major ear row number and tassel

branch number quantitative trait locus[29] – demonstrated preferential accessibility in spikelet

meristems that coincided with the greatest number of *UB2* proximal to distal ACR interactions,

including a cell-type specific ACR located upstream approximately 150-kb (**Fig. 3e**). We posited

that ACRs with expanded interactive capacity resemble enhancers with the potential to influence

organismal phenotypes. Indeed, ACRs with enhancer activity and co-localization with phenotype-

associated genetic variants from GWAS were associated with a significantly greater number of

ACR-ACR connections (Wilcoxon rank sum; *P* < 2.2e-308, **Fig. 3f**, **g**). These results highlight the

occurrence of diverse cell-type specific regulatory configurations among distal enhancer ACRs

and their target genes and implicate genetic variants perturbing highly interactive distal enhancers

as major contributors towards phenotypic variation.

The structural protein, CCCTC-binding factor (CTCF) plays an important role in metazoan

genome organization and is notably absent in plant lineages[30]. In search of an orthogonal factor

in maize, we hypothesized that higher-order chromatin structure captured by co-accessible ACRs

may be driven by TFs recognizing similar sequence motifs embedded within interacting

accessible regions. Comparison of motif occurrences between co-accessible link edges indicated

that ACRs in co-accessible links are more similar to one another than randomly linked ACRs

(empirical; *P* < 1e-4, **Extended Data Fig. 7e**). Furthermore, we identified several cases where

co-accessible ACR edges were reciprocally enriched (FDR < 0.05, Methods) for the same TF

motif in both cell-type specific and non-specific co-accessible links (**Extended Data Fig. 7f**). We

identified *TEOSINTE BRANCHED1/CYCLOIDEA/PROLIFERATING CELL NUCLEAR ANTIGEN*

FACTOR1 (*TCP*), *APETALA2/ETHYLENE-RESPONSIVE ELEMENT BINDING PROTEINS*

(*AP2-EREBP*) and *LATERAL ORGAN BOUNDARIES DOMAIN* (*LBD*) motifs that were not only

broadly associated with co-accessible edges across cell-types (FDR < 0.05), but also exhibited

strikingly similar GC-rich palindromic binding sites (**Extended Data Fig. 7f, g**). A role in chromatin

organization is supported by previous research demonstrating *TCP* motif overrepresentation in

topologically associated domain-like (TAD) boundaries in *Oryza sativa* and *Marchantia*

polymorpha, and the distal edges of chromatin loops in *Zea mays*[27,28,31,32].

Dynamic chromatin accessibility specifies cell developmental trajectories

The apical domains of maize enclose undifferentiated meristematic stem cells that continuously

give rise to all differentiated cell-types[33,34]. To define a *cis*-regulatory catalog of temporal cell-type

progressions, we estimated pseudotime scores for ordered nuclei across 18 developmental

trajectories reflecting meristematic to differentiated cell-states; identifying ACRs, TF loci, and TF

motifs with significant accessibility variation across pseudotime in a given trajectory (**Fig. 4a**,

Extended Data Fig. 8, **Supplementary Data 3-6**, Methods). To showcase the power of trajectory

construction to characterize a relatively understudied process, we focused our analysis on root

Figure 4: Chromatin accessibility is dynamic across pseudotime and conserved between _Arabidopsis thaliana_ and _Zea mays_

a, Overview of pseudotime trajectory analysis. Inflorescence development was performed for both pistillate and staminate inflorescence. **b,** UMAP embedding of companion cell developmental trajectory depicting cell-types (left) and pseudotime progression (right). **c,** Relative motif deviations for 440 TF motifs (left, rows), 402 TF gene accessibility scores (middle, rows), and relative accessibility of 8,094 ACRs (right, rows) associated with pseudotime (columns). Four motifs enriched along the trajectory gradient are shown on the left. ACRs, accessible chromatin regions; TF, transcription factor. **d,** Genome browser screenshots of cell-type specific chromatin accessibility profiles along the developmental trajectory for quiescent center, procambial/phloem sieve element precursor, and companion cell at associated marker gene loci. **e,** UMAP embedding based on whole-genome chromatin accessibility profiles of 4,655 _Arabidopsis thaliana_ root nuclei. **f,** UMAP embedding of compnaion cell developmental trajectories in _Arabidopsis thaliana_ depecting cell-types (left) and pseudotime progression (right). Motifs, TFs, and ACRs with significant association with pseudotime are shown as heatmaps below. **g,** Per-gene pseudotime shift scores from alignments between _Zea mays_ and _Arabidopsis thaliana_ companion cell development progressions, clustered by k-means into three groups. Distribution of gene-gene distances from the alignments are shown on the right, split by k-mean groups. **h,** Exemplary one-to-one homologs between _Arabidopsis thaliana_ and _Zea mays_ for the three groups split by pseudotime shifts. Acronyms: CC, compansion cell; LRC, lateral root cap; LRP, lateral root primordia. QC, quiescent center; SE, sieve elements.

phloem companion cell (PCC) development (**Fig. 4b**). Ordering nuclei along the PCC trajectory,

we identified 8,004 ACRs, 440 TF motifs, 7,955 genes, and 402 TF loci that were differentially

accessible across pseudotime (**Fig. 4c**, Methods). Several known meristem and phloem

developmental genes including *AT-RICH INTERACTIVE DOMAIN-CONTAINING 8* (*ARID8*)[35],

SUPPRESSOR OF MAX2 1-LIKE3 (*ZmSMXL3*)[36], *and SUCROSE TRANSPORTER 1*

(*ZmSUT1*)[37], were identified among the top differentially accessible genes throughout PCC

development (**Fig. 4d**). To realize conserved regulatory mechanisms in plant root development,

we profiled chromatin accessibility in 4,655 nuclei from 7-day old embryonic root tissues in the

eudicot model species *Arabidopsis thaliana* and constructed eight pseudotime trajectories

encompassing vascular, dermal and ground development (**Fig. 4e, f, Extended Data Fig. 9,**

Supplementary Data 7-11). Maintaining focus on PCC development, we aligned *Z. mays* and *A.*

thaliana trajectories using a time-warping algorithm. Consistent with recent comparative analysis

of vascular development in *O. sativa*, *A. thaliana,* and *Solanum lycopersicum*[38], only 206 out of

10,976 putative orthologs were significantly associated (FDR < 0.01) with PCC pseudotime in

both species, indicating that the majority PCC trajectories-associated genes are unique to each

lineage (97% *Z. mays*, 83% *A. thaliana*). However, of the 206 PCC-associated orthologs, ~50%

(102/206) exhibited similar patterns of gene accessibility across pseudotime (**Fig. 4g**, **Extended**

Data Fig. 10a, b). Several putative orthologs with matching gene accessibility patterns have been

previously associated with PCC development, such as *SCARECROW-LIKE8* (*SCL8*), which

displays an increasing expression gradient in nascent to mature *A. thaliana* PCCs in a previously

generated *A. thaliana* root cell-type gene expression atlas (**Fig. 4h**)[39]. Despite evolutionary

conserved sequence homology, the remaining homologs (n=104) exhibited differential

accessibility patterns clustered into two groups that reflect changes in the timing of gene

accessibility across PCC development, underscoring functional novelty in PCC development

between *Z. mays* and *A. thaliana*. To understand the extent of innovation in *cis* regulation along

the pseudotime continuum, we aligned *Z. mays* and *A. thaliana* TF motif accessibility profiles

associated with PCC progression (**Extended Data Fig. 10c**). Of the 440 motifs, 142 demonstrated

highly conserved *cis*-regulatory dynamics between species (**Extended Data Fig. 10d, e**). Indeed,

TFs recognizing the top four motifs ranked by normalized distances (Methods) included

HOMEOBOX25 (*HB25*), *HOMEOBOX18* (*HB18*), *NAC DOMAIN CONTAINING PROTEIN 55*

(*NAC055*)*,* and *NAC DOMAIN CONTAINING PROTEIN 83* (*NAC083*) that have been previously

implicated in regulation of hormonal responses and vascular development[40-42]. Gene expression

profiles of these TFs from published root cell-type resolved data in *A. thaliana* was largely

restricted to maturing procambial and companion cells, consistent with motif accessibility

dynamics in both *Z. mays* and *A. thaliana* ontogenies (**Extended Data Fig. 10f-h**)[39]*.*

In summary, we describe 92 distinct cell-states from a suite of *Z. mays* organs, many of

which previously lacked prior genome-wide characterization. We provide evidence supported by

RNA *in situ* hybridization for the use of chromatin accessibility as a robust proxy of gene

expression. We defined the TFs, CREs, and other genomic regions that discretize cell-type

identities and the sequential trajectories for an array of developmental ontogenies. We

demonstrate that floral cell-type specific ACRs have been the historical targets of modern

agronomic selection and while subject to considerable sequence constraint within maize, cell-

type specific ACRs harbor polymorphisms associated with significant phenotypic variation. By

considering patterns of gene and motif accessibility across two highly diverged species, our

results indicate that a majority of genes involved in cell-type development are unique to each

lineage. This finding contrasts with greater conservation of *cis*-regulatory dynamics, as a greater

proportion of TF motifs exhibited consistent spatiotemporal progressions. Lastly, we anticipate

that the dissemination of ACRs by cell-type presents an ideal framework for future crop

improvement efforts via genome editing and synthetic biology approaches.

ACKNOWLEDGEMENTS

We would like to acknowledge Dr. Zefu Lu for training in nuclei extraction, Julie Nelson from the

University of Georgia flow cytometry core for helping sort nuclei, Noah Workman and Julia

Portocarrero from the Georgia Genomics and Bioinformatics Core for library preparation, the

Georgia Advanced Computing Resource Center for providing access to computing resources,

and Tyler Earp for assistance with data management. This study was funded by support from the

National Science Foundation (IOS-1856627 and IOS-1802848), the Pew Charitable Trusts and

the UGA Office of Research to RJS, the National Science Foundation (IOS-1916804) to AG, and

an NSF Postdoctoral Fellowship in Biology (DBI-1905869) to APM. AG and RJS are also

supported by an NSF Collaborative award to support this research (IOS-2026554 and IOS-

2026561). RJS is a Pew Scholar in the Biomedical Sciences, supported by The Pew Charitable

Trusts.

AUTHOR CONTRIBUTIONS

APM and RJS designed the research. APM and ZC performed the experiments. APM, ZC, AG

and RSJ analyzed the data. APM and RJS wrote the manuscript.

COMPETING INTERESTS

RJS is a co-founder of REqest Genomics, LLC, a company that provides epigenomic services.

APM, ZC and AG declare no competing interests.

SUPPLEMENTARY INFORMATION

Supplementary information is available for this paper.

CORRESPONDENCE

Correspondence and requests for materials should be addressed to Robert J. Schmitz

(schmitz@uga.edu) and Alexandre P. Marand (marand@uga.edu).

METHODS

Plant material and growth conditions

For libraries derived from seedlings, kernels from genotypes B73 and Mo17 were obtained from USDA National Plant Germplasm System (https://npgsweb.ars-grin.gov) and sown in Sungro Horticulture professional growing mix (Sungro Horticulture Canada Ltd.). Soil was saturated with tap water and placed under a 50/50 mixture of 4100K (Sylvania Supersaver Cool White Delux F34CWX/SS, 34W) and 3000K (GE Ecolux w/ starcoat, F40CX30ECO, 40W) lighting. Seedlings were grown under a photoperiod of 16 hours of light, eight hours of dark. The temperature was approximately 25°C during light hours with a relative humidity of approximately 54%. Above ground seedling tissues were harvested between 8 and 9 AM six days (V1-stage) after sowing. We used both fresh (B73/Mo17 pooled) and flash frozen (B73 only) seedling tissue to construct scATAC-seq libraries (**Supplementary Table 1**).

Maize root samples were obtained as follows: B73 kernels were sterilized with 70% EtOH treatment for 5 minutes. After removing the ethanol solution, kernels were suspended with 50% bleach for 30 minutes, followed by five washes with autoclaved Milli-Q water. Sterilized kernels were then sown onto mesh plates with half strength MS (Phytotech laboratories, catalog: M519) media and wrapped in Millipore tape. Plates were incubated in a Percival growth chamber with a photoperiod of 16 hours of light, eight hours of dark. The growth chamber temperature was set to 25°C with a relative humidity of approximately 60%. Apical root tips (bottom 2 cm) of seminal and primary root samples were harvested six days (V1-stage) after sowing between 8 and 9 am. Crown root samples (21 days after sowing) were derived from the three developmental zones of greenhouse grown B73 plants between 8 and 9 am and rinsed with sterile water 3 times.

Data generated from young inflorescence (ear and tassel primordia) were derived from B73 maize grown in the greenhouse. Inflorescence primordia were extracted from shoots harvested approximately one month (V7-stage, 2-4 mm) after sowing, between 8 and 9 AM. Inflorescence primordia between three and eight millimeters from the base to the apical tip were

placed in sterile water and used for nuclei isolation. Axillary buds (~30 samples per library) were

taken from B73 maize plants grown in the greenhouse at approximately the same developmental

stage (V7) as tassel and ear primordia.

Single cell ATAC-seq library preparation

Each library was prepared by mixing at least three independent biological samples (3-4 seedlings,

3 tassel or ear primordia, 12-14 root tips, 12-14 crown root samples and ~30 axillary buds). One

scATAC-seq library (B73 seedling) was derived from flash frozen tissue (liquid nitrogen, followed

by 7-day -80°C storage), while the remaining nine libraries were constructed with freshly

harvested tissue (**Supplementary Table 1**).

To isolate individual plant nuclei, fresh or flash frozen tissue from multiple biological

samples were placed on petri dishes and vigorous chopped with a razor blade for two minutes in

~500 uL LB01 buffer (15mM Tris pH 7.5, 2mM EDTA, 0.5mM Spermine, 80mM KCl, 20mM NaCl,

15mM 2-ME, 0.15% TrixtonX-100). Homogenized tissue was then filtered through two layers of

miracloth, stained with DAPI to a final concentration of ~1uM and loaded onto a Beckman Coulter

MoFlo XDP flow cytometer instrument. A total of 120,000 nuclei were sorted for each sample

across four catch tubes (30,000 nuclei each) containing 200 uL LB01. Isolated nuclei were spun

down in a swinging-bucket (5 minutes, 500 rcf) centrifuge resuspended in 10uL LB01, pooled,

and then visualized on a hemocytometer with a fluorescent microscope. Nuclei suspensions were

then spun down (5 minutes, 500 rcf) and resuspended in diluted nuclei buffer (10X Genomics) to

a final concentration of 3,200 nuclei per uL and used as input for scATAC-seq library preparation

(5 uL; 16,000 nuclei total). Samples were kept on ice for all intermittent steps. For B73/Mo17

mixed library, we pooled 8,000 nuclei from both B73 and Mo17 that were independently isolated.

Single-cell ATAC-seq libraries were constructed according to the manufacture's instruction (10X

Genomics, catalog: 1000176). Libraries were sequenced with Illumina NovaSeq 6000 in dual-

index mode with six and 16 cycles for i7 and i5 index, respectively.

In situ hybridizations

3-4mm tassel and ear primordia and young seedlings from the maize B73 inbred line were

dissected and fixed in a cold paraformaldehyde acetic acid solution (4% PFA) for 48 hours.

Following dehydration through a graded ethanol series and clearing of the tissue with a Histo-

clear II solution (Electron Microscopy Sciences), samples were embedded using Paraplast Plus

tissue embedding media (McCormick Scientific). 8mm sections were hybridized at 56°C with

antisense probes labelled with digoxigenin (DIG RNA labeling mix, Roche), and detected using

NBT/BCIP (Roche). Probes were synthesized by *in vitro* transcription (T7 RNA polymerase,

Promega) of PCR products obtained from embryo cDNA or from digested full-length cDNA clones.

The vectors and primers used for probe design are listed in **Supplementary Table 6**.

Raw reads processing

The following data processing was performed using each tissue and/or replicate independently

unless noted otherwise. Raw BCL files were demultiplexed and convert into fastq format using

the default settings of the 10X Genomics tool *cellranger-atac makefastq* (v1.2.0). Partial raw read

processing (adapter/quality trimming, mapping and barcode attachment/correction) was carried

out with *cellranger-atac count* (v1.2.0) using AGPv4 of the maize B73 reference genome[43].

Properly paired, uniquely mapped reads with mapping quality greater than 10 were retained using

samtools view (v1.6; -f 3 -q 10) and by filtering reads with XA tags[44]. Duplicate fragments were

collapsed on a per-nucleus basis using *picardtools* (http://broadinstitute.github.io/picard)

MarkDuplicates (v2.16; BARCODE_TAG=CB REMOVE_DUPLICATES=TRUE). Reads mapping

to mitochondrial and chloroplast genomes were counted for each barcode, then excluded from

downstream analysis. We removed reads representing potential artifacts by excluding alignments

coincident with a blacklist of regions composed of low-complexity and homopolymeric sequences

(*RepeatMasker* v4.07)[45], nuclear sequences with homology (greater than 80% identity and

coverage) to mitochondrial and chloroplast genomes (*BLAST+* v2.7.1)[46], regions exhibiting Tn5

integration bias from Tn5-treated genomic DNA (1-kb windows with greater than 2-fold coverage

over the genome-wide median), and potential collapsed sequences in the reference (1-kb

windows with greater than 2-fold coverage over the genome-wide median using ChIP-seq input).

Genomic Tn5 and ChIP input data were acquired from Ricci, Lu and Ji et al[10]. BAM alignments

were then converted to single base-pair Tn5 integration sites in BED format by adjusting

coordinates of reads mapping to positive and negative strands by +4 and -5, respectively, and

retaining only unique Tn5 integration sites for each distinct barcode. Sequencing saturation was

calculated as the proportion of unique reads relative to the estimated library complexity output by

the *MarkDuplicates* function apart of picardtools.

Cell calling

To identify high-quality nuclei (a term used interchangeably with "barcodes") using the filtered set

of alignments, we implemented heuristic cutoffs for genomic context and sequencing depth

indicative of high-quality nuclei. Specifically, we fit a smoothed spline to the $\log_{10}$ transformed

unique Tn5 integration sites per nucleus (response) against the ordered $\log_{10}$ barcode rank

(decreasing per-nucleus unique Tn5 integration site counts) using the *smooth.spline* function

(spar=0.01) from base R[47]. We then used the fitted values from the smoothed spline model to

estimate the first derivative (slope), taking the local minima within the first 16,000 barcodes as a

potential knee/inflection point (16,000 was selected to match the maximum number of input

nuclei). We set the unique Tn5 library depth threshold to the lesser of 1,000 reads and the

knee/inflection point, excluding all barcodes below the threshold. Spurious integration patterns

throughout the genome can be representative of incomplete Tn5 integration, fragmented/low-

quality nuclei, or poor sequence recovery, among other sources of technical noise. In contrast,

high quality nuclei often demonstrate a strong aggregate accessibility signal near TSSs.

Therefore, we implemented two approaches for estimating signal-noise ratios in our scATAC-seq

data. First, nuclei below two standard deviations from the mean fraction of reads mapping to within

2-kb of TSSs were removed on a per-library basis. Then, we estimated TSS enrichment scores

by calculating the average per-bp coverage of 2-kb windows surrounding TSSs, scaling by the

average per-bp coverage of the first and last 100-bp in the window (background estimate; average

of 1-100-bp and 1901-2000-bp), and smoothing the scaled signal with rolling-means (R package;

Zoo). Per barcode TSS enrichment scores were taken as the maximum signal within 250-bp of

the TSS. Lastly, for each library, we removed any barcode with a proportion of reads mapping to

chloroplast and mitochondrial genomes greater than two standard deviations from the mean of

the library.

Detection of multiplet droplets

To estimate the empirical proportion of doublets present in our data, we demultiplexed the two-

genotype (B73 and Mo17) pooled seedling scATAC-seq sample and assessed the proportion of

barcodes reflecting a mixtures of reads derived from both genotypes. Specifically, B73 and Mo17

whole genome short read resequencing data were acquired from PRJNA338953. Paired-end

reads were quality and adapter trimmed with *fastp* (v0.19.5)[48] and aligned to the B73 v4 maize

reference genome[43] using *BWA mem*[49] with non-default settings (-MT 1). Duplicate reads were

removed using *samtools rmdup*[44] (v1.6). The genomic coordinates of short nucleotide variants

(SNVs; single nucleotide polymorphisms [SNPs] and small insertions/deletions [INDELs]) for both

genotypes were identified using *freebayes*[50] (v1.0.0) with non-default settings (--min-repeat-

entropy 1 --min-alternate-fraction 0.05). Only biallelic SNPs – requiring at least 5 reads per

genotype where B73 and Mo17 were homozygous for reference and alternate nucleotides,

respectively – were retained. Genotypes were called by modeling allele counts as a binomial

distribution with a term accounting for the sequencing error rate, E_t (determined empirically as

the fraction of SNPs failing to match either allele), estimating posterior probabilities via Bayes

theorem, and assigning the genotype (or mixture of genotypes) with the greatest probability (Eq.

1-7). Specifically, the probability to observe k out of n SNPs from B73 can be modeled as a

binomial distribution for each B73 (A_l), Mo17 (A_2), and doublet barcode state (N) (Eq. 1-3):

1. $P(k|A_1) = \binom{n}{k} \times E_t^{n-k} \times (1 - E_t)^k$

2. $P(k|A_2) = \binom{n}{k} \times (1 - E_t)^{n-k} \times E_t^{k}$

3. $P(k|N) = \binom{n}{k} \times (0.5)^k \times (0.5)^{n-k}$

Let $P(A_1|k)$, $P(A_2|k)$, and $P(N|k)$ reflect posterior probabilities for genotypes B73, Mo17, and

doublet barcodes given k allele counts from B73; posterior probabilities can be estimated as

follows (Eq. 4-6):

4. $P(A_1|k) = \frac{P(k|A_1) \times P(A_1)}{\sum_{i=0}^{n} P(k|A_i) \times P(A_i)}$

5. $P(A_2|k) = \frac{P(k|A_2) \times P(A_2)}{\sum_{i=0}^{n} P(k|A_i) \times P(A_i)}$

6. $P(N|k) = \frac{P(k|N) \times P(N)}{\sum_{i=0}^{n} P(k|A_i) \times P(A_i)}$

Finally, the genotype called for each barcode was determined as the event with the greatest

posterior probability (Eq. 7):

7. $\max\{P(A_1|k), P(A_2|k), P(N|k)\}$

In-silico sorting

To provide sufficient sensitivity for peak calling prior to clustering, we followed an *in-silico* sorting

strategy to identify crude clusters of similar cells within each organ[1]. To do so, we generate a

binary matrix representing the presence/absence of Tn5 integration sites in 1-kb windows across

all cells in a given organ. Bins with less than 1% accessible cells and cells with less than 100

accessible bins were removed. This binary matrix was then transformed using the matrix

normalization method term-frequency inverse document-frequency (TF-IDF). Briefly, the TF term

was estimated by weighting binary counts at each bin by the total number of bins containing Tn5

integration sites in a given cell, scaling each cell to sum to 100,000, adding a pseudo-count of

one, and log transforming the resulting values to reduce the effects of outliers in downstream

processing. The IDF term was calculated as the log transformed ratio of the total number of nuclei

to the number of nuclei that were marked as accessible for a given bin. We add a pseudo-count

of one to the inverse frequency term to avoid taking the log of zero. The TF-IDF scaled matrix

was estimated by taking the dot product of the TF and IDF matrices. To enable faster downstream

computation, we kept the top 25,000 bins with the greatest TF-IDF variance across nuclei. The

reduced TF-IDF matrix was denoised with singular value decomposition (SVD), retaining the 2nd

– 11th dimensions (termed Latent Sematic Indexing, LSI). Each row was centered and

standardized, capping the values at $\pm$ 1.5. Crude clusters were visually identified (k=[6..12]) using

ward.D2 hierarchical bi-clustering on the cosine distances of LSI nuclei and bin embeddings.

ACR identification

ACRs were identified by treating each bulk and single-cell ATAC-seq library as a traditional bulk

ATAC-seq library. Aligned reads were filtered by mapping quality greater than 10, and duplicate

reads were removed via *samtools rmdup.* We then identified ACRs for each library by converting

the BAM alignments in BED format, adjusting the coordinates to reflect single-base Tn5

integrations, and running *MACS2*[51] with non-default parameters: --extsize 150 --shift -75 --

nomodel --keep-dup all. A final set of ACRs for comparing bulk and aggregate scATAC-seq

libraries (**Extended Data Fig. 1b**) was constructed by taking the union of ACRs across all

libraries. To leverage the increased sensitivity afforded by cell-type resolved cluster information

while ensuring robust reproducibility in ACR identification, we generated pseudo-replicated bulk

alignments using the LSI-based crude clusters (see above, "*In-silico* sorting"). Pseudo-replicates

were constructed by randomly allocating nuclei from each cluster into two groups, with a third

group composed of all cells from the cluster (cluster bulk). These groupings were used to

concatenate Tn5 integration sites corresponding to the nuclei from each group into three BED

files. ACRs were then identified from the enrichment of Tn5 integration sites from the pseudo-

replicate or cluster bulk aggregates using *MACS2* run with non-default parameters: --extsize 150

--shift -75 --nomodel --keep-dup all. ACRs from both pseudo-replicates and the cluster bulk were

intersected with BEDtools, retaining ACRs on the conditional intersection of all three groupings

(both pseudo-replicates and the cluster bulk) by at least 25% overlap. The remaining ACRs were

then redefined as 500-bp windows centered on the ACR coverage summit. To integrate

information across all clusters, ACRs from each cluster were concatenated into a single master

list. Lastly, overlapping ACRs were filtered recursively to retain the ACR with the greater

normalized kernel Tn5 integration density as previously described[25].

Nuclei clustering

Starting with a binary nucleus x ACR matrix, we first removed ACRs that were accessible in less

than 0.5% of all nuclei, and filtered nuclei less than 50 accessible ACRs. Inspired by recent

developments in modeling single-cell RNA-seq data[52], we developed a regularized quasibinomial

logistic framework that overcomes noise inherent to sparse, binary scATAC-seq data by pooling

information across ACRs while simultaneously removing variation due to technical effects,

particularly those stemming from differences in barcode sequencing depths. First, a subset of

5,000 representative ACRs selected by kernel density sampling of ACR usage (fraction nuclei

that are accessible at a given ACR) were used to model the parameters of each ACR, using ACR

usage as a covariate in a generalized linear model. Specifically, the expected accessibility of an

ACR, y_i, can be estimated with a generalized linear model containing a binomial error distribution

and logit-link function, and an overdispersion term with a quasibinomial probability density

function (Eq. 8).

8. $\mathbb{E}(y_i) \sim \beta_0 + \beta_1 log10(t)$

Where t is a vector of the sums of accessible ACRs across cell j (Eq. 9):

9. $t = \sum_i y_{ij}$

To prevent over-fitting and ensure robust estimates in light of sampling noise, we learned the

global regularized model parameters, including overdispersion, using the representative ACRs by

fitting each parameter against the log_{10} fraction of accessible nuclei via kernel regression,

resulting in smoothed parameter estimates across the spectrum of ACR accessibility penetrance

present in these data. The learned global regularized model parameters were then used to

constrain fitted values across all ACRs for each nucleus with a simple affine transformation. To

account for technical variation among nuclei (variation in barcode log_{10} transformed read-depth,

in particular) we calculated Pearson residuals for each ACR, scaling the residuals by the

regularized dispersion estimate and centering values via mean subtraction, representing

variance-stabilized and read-depth normalized values of accessibility for a nucleus at a given

ACR. We note that this method is amenable to calculating residuals that account for additional

sources of technical variation, including categorical and numeric covariates, that may obscure

biological signal, such as batch effects, proportion of mitochondrial reads, etc.

The dimensionality of the Pearson residual matrix was reduced using singular value

decomposition (SVD) implemented by the R package *irlba*[53], retaining the first 25 left singular

vectors scaled by singular values (hereafter referred to as nuclei embeddings), analogous to

principal components (PCs) on an uncentered matrix. Nuclei embeddings were then standardized

across components and filtered to remove components correlated with barcode read depth

(Spearman's rho > 0.7). We further reduced the dimensionality of the nuclei embedding with

Uniform Manifold Approximation Projection (UMAP) via the R implementation of *umap-learn*

(min_dist = 0.1, k=50, metric="euclidean"). Nuclei were clustered with the *Seurat* v3[54] framework

and Louvain clustering on a k=50 nearest neighborhood graph at a resolution of 0.02 with 100

iterations and 100 random starts. Clusters with aggregated read depths less than 1.5M were

removed. To filter outliers in the UMAP embedding, we estimated the mean distance for each

nucleus with its k (k=50) nearest neighbors and removed nuclei greater than 3 standard deviations

from the mean.

We observed fine-scale heterogeneity within major clusters, thus we repeated our

clustering pipeline for each major cluster independently by partitioning the SVD embedding into

the top 20 components, L2 normalizing nuclei embeddings across components, and projecting

the L2-normalized embeddings into the UMAP space. Subclusters of nuclei were identified by

Louvain clustering on the L2 normalized SVD embedding (resolution set manually, range = 0.6 –

1.0) with 20 nearest neighbors, filtering outlier nuclei more than 2 standard deviations from the

mean distance of 25 nearest neighbors within each cluster.

For analysis of chromatin accessibility across clusters, we assembled a matrix of clusters

by ACRs by aggregating the number of single-base resolution Tn5 integration sites from nuclei

within the same cluster for each ACR, analogous to normalizing by the proportion of reads in

peaks for each cluster. To account for differences in read depth and other technical factors, the

raw counts were transformed with *edgeR's* "cpm" (log=T, prior.count=5) as previously

described[55]. Log-transformed ACR coverage scores were quantile normalized using

"normalize.quantiles" with the R package, *preprocessCore.* Finally, to aid data visualization, we

estimated per ACR Z-scores across clusters by mean subtraction and standardization (identical

to row-wise execution of the R function, "scale").

Identification of co-accessible ACRs

Recent experiments of population-level chromatin accessibility found that pairwise correlations of

accessibility among ACRs recapitulates higher-order chromatin interactions observed in Hi-C and

other chromatin architecture experiments[24]. A similar framework was applied to populations of

single cells, which showed that co-accessible ACRs are typically more conserved and functionally

associated[26]. To identify potentially functional co-accessible ACRs, we applied a recently

developed method, *Cicero*[23], that estimates regularized correlation scores (ranging from -1 to 1)

among nearby ACRs with graphical LASSO to penalize potential interactions by physical

distances. Using the binary nuclei x ACR matrix as input, we subset nuclei by their subcluster IDs

and estimated co-accessibility among ACRs within 500-kb for each of the 92 clusters,

independently. *Cicero* was run by applying a background sample of 100 random regions, and 15

nuclei pseudo-aggregates based on k-nearest-neighbors derived from the UMAP coordinates. To

control the false discovery rate (FDR) of co-accessible ACR calls, we shuffled the nuclei x ACR

matrix such that the total number of reads per ACR and reads per nucleus were identical to the

original matrix. We then repeated co-accessible ACR identification with the shuffled matrix,

keeping the original parameters to *Cicero* unchanged. Empirical FDR cluster-specific cut-offs

were constructed by identifying the minimum positive co-accessibility score in the background

where the FDR < 0.05. Co-accessible links below cluster-specific thresholds were removed. Co-

accessible ACRs passing thresholds were compared with previously published HiC and HiChIP

data sets derived from maize seedling and pistillate inflorescence primordia[10].

Estimation of gene accessibility scores

Chromatin accessibility at TSSs and gene bodies exhibit marked correlation with transcription output in bulk samples (**Extended Data Fig. 3f**). To aid the identification of marker genes underlying distinct cell-types, we used *Cicero* to estimate gene activity scores. *Cicero* models gene activity as a weighted accessibility score that integrates both proximal and distal regulatory elements linked to a single gene by co-accessibility analysis (see above section "Identification of co-accessible ACRs"). Relative gene accessibility scores per nucleus were estimated by taking a weighted average (3:1, gene body score to proximal/distal activity) of the scaled number of reads mapping to gene bodies for each barcode (summing to 1) with the *Cicero* estimate of gene activity derived from ACRs mapping to 1-kb upstream of gene TSSs and their associated distal ACRs linked by co-accessible ACRs passing FDR < 0.05 thresholds (connected ACRs were constrained to a minimum and maximum intervening distance of 1- and 500-kb, respectively). These weighted gene accessibility scores were rescaled such that gene accessibility scores for a given nucleus summed to 1.

Relative gene accessibility scores exhibited a bimodal distribution with relative gene accessibility values near zero resembling low or non-expressed genes. We applied a gaussian mixture-model (two distributions) based scaling step per cluster to reduce noise introduced by genes with low gene accessibility. Briefly, the average gene accessibility across nuclei was fit to a two distribution gaussian mixture model in each cluster using the R package *mclust*. We estimated cluster-specific scaling parameters determined as the 5% quantile of non-zero gene accessibility values of genes from the gaussian distribution with the larger mean, for each cluster. This parameter was then used to scale gene accessibility scores for all genes in each nucleus within the cluster. Scaled gene accessibility scores were rounded to the nearest integer and normalized across all nuclei and clusters using nucleus-specific size factors estimated as the total gene accessibility of a nucleus divided by the exponential of the mean of log-transformed gene

accessibility sums across nuclei. To aid visualization, we smoothed normalized gene accessibility

scores by estimating a diffusion nearest neighbor graph (k=15) using the SVD embedding with 3

steps similar to the *run_MAGIC* function in R package *snapATAC*[56]. Downstream analyzes based

on binarized gene accessibly were conducted by simply converting normalized (non-smoothed)

accessibility scores to 1 for all positive values.

Cell-type annotation

To identify and annotate cell-types for each barcode, we identified marker genes known to localize

to discrete cell-types or domains expected in the sampled tissues/organs based on extensive

review of the literature (**Supplementary Table 2**). To enable gene accessibility comparisons

among clusters, we generated three pseudo-replicates for each cluster by resampling nuclei

within the cluster such that all cluster pseudo-replicates contained the mean number of nuclei

across clusters (number of nuclei per pseudo-replicate = 552) without replacement when possible.

To identify genes with increased accessibly relative to other clusters, we constructed a reference

panel with three pseudo-replicates by uniformly sampling nuclei without replacement from each

organ (number of nuclei per organ = 92), with a total of 552 nuclei per reference panel pseudo-

replicate. Read counts per gene were summed across nuclei within each pseudo-replicate. Using

the *DESeq2* R package, we identified genes with significantly different (FDR < 0.01) accessibility

profiles between each cluster and the reference panel.

The list of significantly differentially accessible genes was filtered to retain the genes on

our list of cell-type specific markers. We initially ranked the top three marker genes in each cluster

by their test statistics. To account for clusters containing small proportions of contaminating nuclei

of a different cell-type, we adjusted the test statistics using a previously described method[1],

effectively scaling marker activity scores by the proportion of nuclei in the cluster that were derived

from an organ in which the marker gene i is an expected cell-type. Clusters where the top three

markers corresponded to the same cell-type were annotated with the consensus cell-type.

As an independent method for cell-type annotation, we devised a resampling and

normalization procedure on the $\log_2$ fold-change values of marker genes to evaluate cell-type

enrichment across all possible cell-types for each cluster, normalizing enrichment scores by

random permutations accounting for different numbers of markers associated with each cell-type.

Briefly, starting with differential gene accessibility information for each cluster, we iterated over all

cell-types, extracting markers associated with the cell-type of interest. Then, we summed the $\log_2$

fold-changes values of all markers and multiplied the sum by the proportion of markers passing

heuristic thresholds (fold-change > 2 and FDR < 0.01). This score was subtracted by the average

of 1,000 random permuted scores from combinations of markers from the remaining cell-types

(selecting the same number of random genes as the cell-type of interest) and divided by the

standard deviation of the permuted scores. Cell-type enrichment scores in each cluster were

scaled from zero to one by dividing each cell-type enrichment score by the maximum scores

across possible cell-types. This approach is effective in normalizing differences arising from

varying numbers of markers specified for each cell-type. Additionally, cell-type annotation scores

for clusters with mixed or unknown identity are approximately equally distributed, thus controlling

ascertainment bias stemming from marker gene selection. Stated differently, an advantage of this

approach is that clusters corresponding to cell-types with few or no markers in the tested list are

left unassigned as their enrichment scores do not deviated significantly from background levels.

Finally, scaled cell-type enrichment scores greater than 0.9 were taken as possible annotations

and intersected with putative cell-type labels from the marker ranking approach described above.

For clusters with ambiguous marker gene labels, we developed a logistic regression

classifier to identify putative cell-types based on whole-genome gene accessibility scores of well-

annotated cells. First, we counted the number of Tn5 integration sites per cell overlapping 2-kb

upstream to 500-bp downstream of each gene. Read counts were transformed by trimmed mean

of M-values (TMM) to enable intra and inter-nucleus comparisons using *edgeR*[57], scaling gene

accessibility scores in each nucleus with counts per million. Next, we estimated cell-type

enrichment scores for each nucleus by calculating the mean accessibility scores of markers for a

given cell-type, subtracting the mean background signal defined as 1,000 sets of averaged

randomly sampled genes (each set had the same number of genes as the number of markers),

divided by the standard deviation of the background signal. Enrichment scores for each nucleus

were transformed into a probability distribution by dividing by the sum of cell-type enrichment

scores. For each nucleus, we compared the top two most likely cell-types, retaining nuclei where

the top predicted cell-type had a two-fold greater probability than the next most likely assignment.

We used these high-confidence cells to train a regularized logistic multinominal classifier with the

R package, *glmnet*. Cell-type classifications with less than 10 nuclei in the training set were

excluded. We used a LASSO L1 penalty to regularize the logistic classifier, modeling the training

set of nuclei as observations and TMM gene accessibility scores as variables. We balanced

observations by weighting by the inverse frequency of cell-types in the training set. The model

was trained with 10-folds and evaluated by testing a 20% hold-out set of nuclei. The predicted

cell-type for each nucleus in the atlas was taken as the cell-type with the greatest probability if

the probability ratio between the best and next best assignment was greater than five-fold,

otherwise labeled as 'unknown'. Using these per-cell assignments, we defined subclusters as the

majority cell-type if greater than 50% of nuclei in the cluster were in agreement, labelling clusters

with two or more majority cell-types as 'mixed' and all other clusters as 'unknown'. All cell-type

labels from these three automated approaches were manually reviewed by careful evaluation with

UMAP gene accessibility score embeddings and cluster aggregated coverages for all marker

genes and refined *ad hoc*.

STARR-seq analysis

Single bp-resolution enhancer activities were available from a previous study[10]. Enhancer activity

(defined as the $\log_2$ ratio between RNA and DNA input fragments scaled per million) for each ACR

was taken as the maximum over the entire ACR. A control set of regions was generated to match

each ACR with the following criteria: (*i*) GC content within 5%, (*ii*) physically constrained to within

50-kb of an ACR, and (*iii*) the same length (500-bp) distribution. The same set of control regions

was used throughout the analysis.

Analysis of differential chromatin accessibility

Next, we implemented a logistic regression framework based on binarized ACR accessibility

scores for assessing the importance of each ACR to cluster membership by estimating the

likelihood ratio between logistic models with, and without a term for cluster membership.

Specifically, for each cluster, we compared binarized ACR accessibility scores to a reference

panel of uniformly sampled nuclei from each organ (111 nuclei from each organ) where the total

number of reference nuclei was set to the average number of nuclei per cluster (n=666). We then

fit two generalized linear logistic regression models (Eq. 10-11), with and without a term for

membership to the cluster of interest.

10. $logit(p_{ij}) = u_i + \alpha_j + \beta_j + \varepsilon_i$

11. $logit(p_{ij}) = u_i + \beta_j + \varepsilon_i$

Where p_{ij} is the probability that ACR i is accessible in nucleus j, u_i is the proportion of nuclei

where ACR i is accessible, α_j is the cluster membership of nucleus j, β_j is the log_{10} number of

accessible ACRs in nucleus j and ε_i is the error term for the i^{th} ACR. We then used a likelihood

ratio test to compare the fits of the two models and estimated the false discovery rate (FDR) using

the Benjamini-Hochberg method to identify ACRs that were significantly differentially accessible

across clusters by conditioning on FDR < 5% and fold-change threshold greater than two. ACRs

meeting these criteria with positive Z-scores in nine or fewer clusters (< 10% of clusters) were

considered as cluster-specific. Analysis of differential gene accessibility was performed as

described in the section titled "Cell-type annotation".

GO gene set enrichment analysis

Gene set enrichment using GO biological process terms was performed using the R package

fgsea. For each cluster, test statistics were multiplied by the sign of the $\log_2$ fold-change value

versus the reference panel. GO terms with gene sets less than 10 and greater than 600 were

excluded from the analysis. GO terms were considered significantly enriched at FDR < 0.05

following 10,000 permutations.

Motif analysis

Motif occurrences were identified genome-wide with *fimo* from the MEME suite toolset[58] using

position weight matrices (PWM) based on DAP-seq data in *A. thaliana* and *Zea mays*[59,60]. To

identity TF motifs associated with cell-type specific ACRs, we ranked the top 2,000 ACRs in each

cell-type by Z-scores derived from CPM normalized accessibility values (see section above

"Nuclei clustering"). As a reference for comparison, we identified 2,000 "constitutive" ACRs that

varied the least and were broadly accessible across clusters. The number of ACRs containing a

specific motif was compared to the frequency of constitutive ACRs harboring the same motif using

a binomial test for each cell-type and motif. To control for multiple testing, we used the Benjamini-

Hochberg method to estimate the FDR, considering tests with FDR < 0.05 as significantly different

between the focal cell-type and constitutively accessible regions. Maize homologs of *A. thaliana*

TFs were identified using protein fasta alignments from BLAST+ v2.10.0 with an E-value cut-off

of 1e-5. Only fasta sequences classified as transcription factors from either species were

considered during alignment. To narrow the list of putative orthologs based on functional similarity

to *A. thaliana* TFs, we filtered alignments with less than 30% identity, removed maize TFs

classified as belonging to a different family, and selected the homolog with the greatest Pearson

correlation coefficient (PCC) with respect to the motif deviation score. Motif deviation scores of

specific TF motifs among nuclei were estimated using *chromVAR*[61] with the non-redundant core

plant PWM database from JASPAR2018. The input matrix for *chromVar* was filtered to retain a

minimum of 50 accessible nuclei per ACR and barcodes with at least 50 accessible ACRs. We

visualized differences in global motif usage per nucleus by projecting deviation scores onto the

UMAP embeddings.

Analysis of cell-type specific selection signatures

Multi-locus allele-frequency differentiation signals between chronologically sampled elite maize

inbred lines were mapped onto ACRs[19], where the selection score for an ACR was taken as the

maximum XP-CLR value within the 500-bp ACR interval. To identity cell-types associated with

increased signatures of selection, the top 2,000 ACRs defined by standardized quantile-scaled

CPM chromatin accessibility (Z-scores, see above "Nuclei clustering") were identified for each

cell-type. The mean XP-CLR scores per-cell type were standardized by the mean and standard

deviation of randomly sampled ACRs (n=2,000) without replacement across 1,000 permutations,

where each permutation estimates the mean XP-CLR scores of a random subset of 2,000 ACRs

from the total list of 165,913 possible ACRs. Enrichment Z-scores were converted into *P*-values

using the R function *pnorm* (log.p=T, lower.tail=F) and used to estimate FDR via the Benjamini-

Hochberg method with the R function *p.adjust* (method="fdr").

Analysis of co-accessible ACRs

To enable comparison with previously identified Hi-C and HiChIP loops[10], we constrained the

distance between co-accessible ACRs to the same range as loops identified in leaf Hi-C and

HiChIP (minimum loop distance = 20-kb). Co-accessible ACRs and Hi-C/HiChIP loops were

considered overlapping if both anchors overlapped by at least 50-bp. We compared motif

composition of co-accessible ACRs by scoring motif occurrence as binary for each ACR and

estimating a Jaccard similarity score on the union of motif sets. Motif similarity scores for co-

accessible ACRs in each cell-type were compared to a null distribution by repeating Jaccard

similarity calculations for non-co-accessible ACR-ACR connections (constraining the null

connections to blocks of 1,000 ACR on the same chromosome with the same ACR-ACR distance

distribution as co-accessible ACRs) across 1,000 permutations. To identity motifs enriched at co-

accessible ACR anchors, we first estimated the proportion of co-accessible ACRs with an identical

motif at both anchors for each motif and cell-type. Then, we constructed the same number of

random ACR-ACR connections as co-accessible ACRs, again estimating the proportion of links

with an identical motif at both anchors, building a null distribution over 1,000 random

permutations. The estimated proportion of co-accessible ACRs with identical motifs at both

anchors for each motif was transformed to a Z-score by subtracting and scaling by the mean and

standard deviation of the null distribution. Z-scores were converted to *P*-values using the R

function, *qnorm* with non-default parameters (log.p=T, lower.tail=F). FDR values were estimated

using *p.adjust* (method="fdr"). Co-accessible motif scores were plotted as heatmaps using

heatmap.2 by subtracting and dividing observed with expected proportions. Rows and columns

were clustered with *hclust* (method="ward.D2").

Pseudotime analysis

Pseudotime trajectories were constructed similar to previous methods[62]. Briefly, nuclei were

ordered based on the principal component space by fitting a continuous trajectory via a smooth

spline on the Euclidean distances of each nuclei to a predefined order of cell-types. For feature

analysis (ACRs, motifs, and TF activity) across pseudotime, nuclei were sorted by ascending

pseudotime. The ACR x nucleus matrix was filtered to retain differentially accessible ACRs (see

section "Analysis of differential accessibility across pseudotime" below) with at least one nucleus

defined as accessible. For each ACR, we fit a generalized additive model with the binary

accessibility scores as the response and a smoothed pseudotime component as the dependent

variable [s(pseudotime, bs="cs")] with a binomial error term and a logit-link function with *gam* from

the *mgcv* R package. Predicted accessibility scores across pseudotime were generated from 500

equally spaced interpolated points covering the range of pseudotime values. Finally, predicted

accessibility scores were mean-centered, standardized and constrained to the range ± 1 for each

ACR. Model specification for motif deviations and TF gene accessibility analysis was similar to

ACR pseudotime analysis with the exception of a Gaussian error distribution, and TF gene

accessibility was normalized by the row maximum rather than rescaling on a ± 1 distribution.

Analysis of differential accessibility across pseudotime

To identity differentially accessible ACRs across pseudotime, we fit normalized accessibility

residuals (Pearson residuals from a generalized linear logistic regression model with $\log_{10}$ number

of accessible ACRs per barcode as the dependent variable, see section "Nuclei Clustering"

above) as the response and pseudotime as the dependent variable using a natural spline with six

degrees of freedom [ns(pseudotime, df=6)] from the R package *splines* for each trajectory. We

took an F-test based approach for hypothesis testing of differential accessibility across

pseudotime by comparing the variance explained by the splined linear model with that of the

residuals normalized by degrees of freedom. *P*-values from the model were used to estimate

Benjamini-Hochberg FDR values with the R function *p.adjust* (method="fdr"), where a FDR

threshold < 0.05 denoted statistical significance for differentially accessible ACRs across

pseudotime. To identify genes and TF motifs with differential accessibility across pseudotime, we

fit the linear splined regression model with the normalized gene accessibility scores and motif

deviations from each nucleus, respectively, similar to the analysis of ACRs.

Aligning pseudotime trajectories between *A. thaliana* and *Z. mays*

To enable comparison of companion cell development between *A. thaliana* and *Z. mays*, we first

identified putative one-to-one orthologs using *OrthoFinder* (v2)[63]. Gene accessibility scores for

10,976 putative orthologs were imputed using a diffusion-based approach[56,64] and scaled from 0

to 1 across pseudotime for barcodes associated with companion cell development in *A. thaliana*

and *Z. mays*. To account for different distributions, pseudotime coverage, and number of

barcodes between species, we used the R package, *cellAlign,* that interpolates, scales, and

weights gene accessibility scores on a fixed set of (n=200) equally spaced points (width

parameter: winSz=0.1) from two trajectories to remove technical biases inherent to each data

set[65]. For each putative ortholog, we performed global alignment of gene accessibility scores

across *A. thaliana* and *Z. mays* pseudotime using the dynamic time warping algorithm with default

settings in *cellAlign.* We then extracted the pseudotime shifts, representing the extent of gene

accessibility deviation at any given point along the trajectory, for each putative ortholog. We

clustered genes into two groups based on pseudotime shifts across companion cell development

using k-means clustering. To identify conserved gene accessibility patterns across pseudotime,

we clustered the normalized distances between *A. thaliana* and *Z. mays* putative orthologs using

a mixture model (G=2) with the R package, *mclust.* The mixture model identified a bimodal

distribution of normalized distances with 0.15 as a natural cut-off for defining conserved

accessibility patterns. Putative orthologs with normalized distances less than the cut-off were

placed in a third group defined as conserved. The above analysis was repeated with TF motif

deviations scores for 440 TF motifs, without the need for ortholog searching as the same TF

position weight matrices were used for both species, affording identical TF motif labels.

DATA AVAILABILITY

Supplementary Data are available on the online versions of this manuscript. Raw and processed

data has been deposited in NCBI GEO database under accession code GSEXXXXXX. Cell-type

resolved data can be viewed through our public Plant Epigenome JBrowse Genome Browser[66]

(URL available upon publication).

CODE AVAILABILITY

Code used throughout the analysis will be made available upon publication.

Extended Data Fig. 2: Cell calling and barcode quality control

a, Comparison of normalized (0-1) read depths at the union of all peaks across bulk and single-cell samples (n=265,992) between replicated libraries, and between bulk and single-cell ATAC-seq assays. **b**, Fragment length distributions for each library. Solid lines indicate the average distribution across cells within the sample. Grey polygons represent the standard deviation across cells in the library. **c**, Enrichment plots centered on 2-kb windows surrounding TSSs for barcodes in each tissue. Grey polygons indicate the standard deviation across cells within the noted tissue. **d**, Density scatter plots of log10 transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites mapping to within 2-kb of transcription start sites (TSSs). Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes. **e**, Knee plots illustrating log10 transformed cellular read depths of log10 ranked barcodes across libraries. **f**, Density scatter plots of log10 transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites derived from organeller sequences (chloroplast and mitocondrial) relative to the total number of unique Tn5 integration sites associated with cognate barcode. Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes.

Extended Data Fig. 3: Clustering metrics and comparison of bulk gene accessibility and expression
a, Parameter regularization of model coefficients (y-axes) with respect to ACR usage (x-axes; proportion of nuclei with at least one Tn5 integration site in an ACR). **b**, Proportion of variance captured by the first 26 PCs. Inset: Spearman's correlation of principal components with cell read depth (log10-transformed). **c**, Number of accessible sites per cell (log10). **d**, Proportion of Tn5 integrations within 2kb of gene TSSs per cell. **e**, Co-localization of nuclei barcodes from different biological replicates for three organs. **f**, Comparison of bulk RNA-seq expression levels (y-axis, log2 CPM) versus aggregate scATAC-seq gene accessibility scores (x-axis, log2 CPM) within an organ.

Extended Data Fig. 4: Cell-type annotation and GO enrichment

a, UMAP embeddings of nuclei barcodes colored by low (grey) to high (dark purple) gene activity values of cell-type specific marker genes. **b**, RNA *in situ* hybridization showing expression of *LOX10* in glume primordia and *GRFTF36* in IM and SMs of staminate inflorescence; *LRR445* in the IM periphery and SPMs and *MYB89* in the IM and suppressed bract primordia of pistillate inflorescence; Zm00001d038453 in ground tissue of SAM and leaf primordia sections. Gene accessibility scores and predicted cell-types are shown on the right. *i*, tassel primordia. *ii*, ear primordia. *iii*, SAM/leaf. Black triangles point to the glume primordia. Red triangles point to suppressed bract primordia. Size bars illustrate 100-um. AM, axillary meristem; BS, bundle sheath; GC, guard cell; GM, ground meristem; GMC, guard mother cell; GP, glume primordia; IM, inflorescence meristem; L1, layer 1; LFM, lower floral meristem; SAM, shoot apical meristem; SBP, suppressed bract primordia; SM, spikelet meristem; SPM; spikelet pair meristem; Stomatal PC, stomatal precursor; UKN, unknown; VEMI, vascular/epidermal meristematic identity; VP, vascular parenchyma; XP, xylem parenchyma. **c,** Proportion of cells within subcluster (column) derived from one of six organs (rows). **d,** Distribution of GO term enrichment across clusters, where the x-axis indicates the number of clusters in which a GO term is significantly enriched.

Extended Data Fig. 5: Chromatin accessibility variation across plant cell-types

a, Row chromatin accessibility Z-scores across cell-types. Cell-type and subcluster labels are located adjacent to the heatmap in identical order as the matrix. The first number after the cell-type label indicates the major cluster number, while the second number represents the subcluster ID. Legend colors: top left triangle, subcluster (corresponds to the sub-cluster annotation on heatmap). bottom right triangle, major cluster (corresponds to the major cluster annotation on heatmap). **b**, Row gene accessibility Z-scores across cell-types. **c**, Proportion of cells derived from one of six organs within each cell-type/subcluster.

Extended Data Fig. 6: Transcription factor motif variation underly dynamic accessible regions
a, Comparison of cell-type specific ACR counts (by cell-type) and the number of unique Tn5 integration sites. **b,** Proportion of distal, genic and proximal ACRs per cell-type. **c,** Ranked TF motif variability across cells, colored by TF family. **d,** Mean TF family motif enrichment (average deviation scores per cell-type per TF family) across cell-types scaled to +/- 1. **e,** Spearman's correlation coefficient (rho) between TF motifs (comparison of motif deviations across all nuclei). Row and column colors represent TF motif families. **f,** TF motif deviations for 440 TF motifs (columns) per nucleus (rows). The TF family for each corresponding motif is denoted by column header colors. Cluster identification (sub-cluster/major cluster) are illustrated as row header colors on the right side of the heatmap. **g,** Relative gene accessibility scores for 2,423 maize transcription factors (columns) for 50,640 nuclei (rows). Nuclei are sorted by sub-cluster cell-type (row color labels). A cell-type annotation color legend applicable to panels b, f, and g where left triangle reflects sub-cluster, right triangle indicates the major cluster.

Extended Data Fig. 7: Co-accessible ACRs reflect in vivo chromatin interactions driven by coordinated TF activity

a, Proportions of co-accessible ACR types, illustrated by toy examples. **b**, Proportion leaf Hi-C, H3K4me3-HiChIP and H3K27me3-HiChIP chromatin loops that overlap co-accessible ACRs from leaf cell-types (clusters with greater than 50% of cell derived from seedlings). **c**, Top, log10 number of connections per ACR per cell-type from all co-accessible ACRs split by genomic context: distal, proximal, and genic. Bottom, log10 number of connections per ACR per cell-type from cell-type specific (purple) and non-specific (grey) co-accessible ACRs, split by genomic context. **d**, Proportion of co-accessible classifications by cell-type for all co-accessible ACRs (top) and cell-type specific co-accessible ACRs (bottom). **e**, Jaccard similarity of motif composition between co-accessible ACR edges by cell-type (colored diamonds) relative to the same number of random ACR-ACR links, permuted 1,000 times for each cell-type (grey boxplots). Box plots represent the interquartile range, grey lines indicate the permuted range. **f**, Heatmaps of observed proportion of co-accessible ACRs with the same motif embedded within link edges subtracted and divided by the expected proportion estimated by 1,000 permutations using random ACR-ACR links with the same number of co-accessible ACRs. **g**, Exemplary motifs enriched in reciprocal co-accessible ACRs for *TCP*, *AP2-EREB*, and *LBD* TF families.

Extended Data Fig. 8: Pseudotime trajectory construction

a, Overview of pseudotime developmental trajectory analysis from four organs: root, seedling, tassel (staminate inflorescence), and ear (pistillate inflorescence). **b,** Endodermis development in crown roots. **c,** Cortex development in crown roots. **d,** Pericycle development in embryonic roots. **e,** Atrichoblast development in crown roots. **f,** Trichoblast development in crown roots. **g,** Trichoblast development in embryonic roots. **h,** Lateral root cap (LRC) development in crown roots. **i,** Lateral root cap (LRC) development in embryonic roots. **j,** Phloem sieve element (SE) development in crown roots. **k,** Companion cell development in crown roots. **l,** Phloem sieve element (SE) development in embryonic roots. **m,** Xylem development in crown roots. **n,** Xylem development in embryonic roots. **o,** Procambial development in crown roots. **p,** Guard cell development in seedling. **q,** Subsidiary cell development in seedlings. **r,** Floral primordia development in staminate inflorescence (tassel). **s,** Floral primordia development in pistillate inflorescence (ear).

Extended Data Fig. 9: *Arabidopsis thaliana* **root cell-type atlas**

a, Knee plots for Arabidopsis thaliana root samples illustrating log10 transformed cellular read depths of log10 ranked barcodes across two biological replicates. **b,** Density scatter plots of log10 transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites derived from organellar sequences (chloroplast and mitochondrial) relative to the total number of unique Tn5 integration sites associated with each barcode from the two biological replicates. Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes. **c,** Density scatter plots of log10 transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites mapping to within 2-kb of transcription start sites (TSSs). Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes. **d,** Average TSS enrichment (normalized read depth adjusted by the two 10 bp windows 1-kb away from TSSs) across 5,001 Arabidopsis thaliana root barcodes (rows). **e,** UMAP (Uniform manifold approximation projection) embeddings of Arabidopsis thaliana root barcodes colored by biological replicate; **f,** total number of accessible chromatin regions (ACRs); **g,** the proportion of Tn5 integration sites within 1-kb of TSSs. **h,** Relative gene accessibility for 27 known cell-type/domain restricted marker genes used to inform cell-type annotation of Arabidopsis thaliana root clusters. **i,** Relative motif deviations for transcription factors with known cell-type specificities.

Extended Data Fig. 10: Dynamic and conserved chromatin accessibility across pseudotime between *Arabidopsis thaliana* and *Zea mays*

a, Pseudotime trajectories for Atrichoblast, Trichoblast, Lateral root cap (LRC), Cortex, Endodermis, Lateral root primordia (LRP), Companion cells (CC), and Xylem development. **b,** Averaged alignments of conserved, shift early *Z. mays*, and shift early *A. thaliana* putative orthologs. **c,** Pseudotime shifts of TF motifs between *A. thaliana* and *Z. mays*, clustered into k-means and conserved groups. **d,** Distributions of motif-motif normalized distances between *Z. mays* and *A. thaliana* for the three groups. **e,** Conserved motifs (n=142) ordered by pseudotime. Heatmaps for *A. thaliana* and *Z. mays* have identical row orders. **f,** Averaged alignments of conserved, shift early *Z. mays*, and shift early *A. thaliana* groups based on motif-motif global alignments from the dynamic time-warping algorithm. **g,** Examples of conserved, shift early *Z. mays*, and shift early *A. thaliana* motifs from both species. **h,** Gene expression Z-scores across *A. thaliana* FAC sorted root cell-types for the TFs recognizing the top four conserved motifs.

 FIGURE LEGENDS

Figure 1: Atlas-scale profiling of single nuclei chromatin accessibility in *Zea mays*

a, Overview of experimental protocol. Homogenized samples release nuclei that are extracted by

flow cytometry. Accessible chromatin is tagged by Tn5 and individual nuclei are distinctly

barcoded in disparate droplets following microfluidic separation, allowing droplet pooling and

library amplification prior to sequencing. **b,** Nuclei similarity clustering as a UMAP embedding

derived from the denoised quasibinomial residuals across all ACRs for each nucleus. UMAP

embedding of nuclei colored by **b,** organ identity and **c,** major cluster identity. **d,** UMAP

embedding of sub-cluster assignments following a second round of clustering within each major

cluster. **e,** Cell-type specific enrichment of gene accessibility for a subset of marker genes

associated with six different cell-types. **f,** Sub-cluster specific chromatin accessibility profiles

surrounding known marker genes for floral primordia, xylem precursors, and L1 epidermal cells.

Bold, circled numbers indicate the cognate major cluster shown in panel **c**. Sub-cluster numeric

identifications are present on the sides of the coverage plots. **g,** Top, gene accessibility for

ZmGRFTF36, an inflorescence and spikelet meristem enriched TF with no previously known cell-

type specificity. Bottom, RNA in situ hybridization of *ZmGRFTF36* in maize B73 staminate (tassel)

primordia. FP, floral primordia; IM, inflorescence meristem; LFM, lower floral meristem; SM,

spikelet meristem; SPM, spikelet pair meristem; TF, transcription factor; UKN, unknown.

Figure 2: Characterization of accessible chromatin regions

a, Distribution of enhancer activity (maximum $\log_2$[RNA/input]) for control regions (n=165,913),

constitutively accessible (n=113,393) and cluster-specific ACRs (n=52,520). The dash red line

indicates the overall mean. Orange lines reflect differences between the group mean and overall

mean. **b,** Distribution of ACR distances to the nearest gene. Inset, distribution of ACR genomic

context. **c,** Relative DNA methylation levels 2-kb flanking ACRs. **d,** Relative enrichment of

polymorphisms after normalizing by mapability for 5-kb regions flanking cell-type specific and

constitutive ACR summits. Smoothed splines are shown as dark lines. **e,** Relative enrichment of

significant GWAS polymorphisms relative to all polymorphisms for 5-kb regions flanking cell-type

specific and constitutive ACR summits. **f,** Enrichment of signatures of selection (XP-CLR) in the

top 2,000 ACRs for all cell-type clusters. The 20 most enriched cell-types are highlighted on the

left. AM, axillary meristem; CC, companion cell; FM, floral meristem; FP, floral primordia; IM,

inflorescence meristem; QC, quiescent center; SM, spikelet meristem; SPM, spikelet pair

meristem. **g,** Average motif coverage for ACRs (n=165,913) and control regions (n=165,913).

Shaded polygon, 95% confidence intervals. **h,** Enrichment of TF motifs in the top 2,000 ACRs

ranked by Z-score for each cell-type compared to the top 2,000 most constitutive ACRs via

binomial tests. FDR was estimated by the Benjamini-Hochberg method. **i,** Comparative heatmaps

for matched TF gene accessibility (bottom) and motif deviation (top) Z-scores across clusters. **j,**

Gene accessibility scores for five maize transcription factors (top) and their associated motif

deviations (bottom).

Figure 3: Co-accessible ACRs reflect *in vivo* chromatin interactions and are established

by cell-type specific TFs. a, Comparison of ear and seedling co-accessible ACRs across nine

cell-types at the genetically mapped *tb1-enhancer* domestication locus (highlighted region). Link

height reflects co-accessibility scores between ACRs across cells in a cluster. **b**, Average

normalized Hi-C signal across 4-kb windows centered on ACRs equally distributed into three

groups (each group: n=54,904) based on the number and strength of participating co-accessible

links. **c**, Co-accessible ACR interaction frequency across clusters. Inset: Single cell-type co-

accessible links (n=1,018,417) for different genomic contexts. **d**, Standardized (row-wise) number

of connections for each ACR (rows) by cell-type (columns). Column color map reflect cell-types

from the legend in Extended Data Fig. 4. Gene-proximal ACRs for a subset of marker genes are

indicated on the right. **e**, Right, chromatin accessibility and co-accessible ACR links surrounding

the *UB2* locus associated with ear row number and tassel branch number quantitative traits. Black

arrow indicates a distal ACR upstream of *UB2* present only in spikelet meristems. Co-accessible

links with an edge within 2-kb of *UB2* are colored pink while remaining links are grey. Link height

represents co-accessibility strength. Left, close-up of accessibility profiles of *UB2*. **f**, Distributions

of average ACR-ACR links across cell-types for ACRs that overlap (purple) and do not overlap

(grey) phenotype-associated genetic variants from maize GWAS. The median of each distribution

is shown as a white horizontal line. Violin plots present the entire range of average number of

connections on a log scale. Hypothesis testing was conducted within the *R* statistical framework

via Wilcoxon rank sum test. **g,** Distributions of average ACR-ACR links cell-types for ACRs with

(blue) and without (grey) enhancer activity ($\log_2$ RNA/input greater than 0). Hypothesis testing

and distribution illustration was performed similarly as panel **f**.

Figure 4: Chromatin accessibility is dynamic across pseudotime and conserved between

Arabidopsis thaliana* and *Zea mays

a, Overview of pseudotime trajectory analysis. Inflorescence development includes trajectory

construction of pistillate and staminate inflorescence, and root development encompasses

embryonic and crown root trajectories. **b,** UMAP embedding of companion cell developmental

trajectory depicting cell-types (left) and pseudotime progression (right). **c,** Relative motif

deviations for 440 TF motifs (left, rows), 402 TF gene accessibility scores (middle, rows), and

relative accessibility of 8,094 ACRs (right, rows) associated with pseudotime (columns). Four

motifs enriched along the trajectory gradient are shown on the left. ACRs, accessible chromatin

regions; TF, transcription factor. **d,** Genome browser screenshots of cell-type specific chromatin

accessibility profiles along the developmental trajectory for quiescent center, procambial/phloem

sieve element precursor, and companion cell at associated marker gene loci. **e,** UMAP

embedding based on whole-genome chromatin accessibility profiles of 4,655 *A. thaliana* root

nuclei. **f,** UMAP embedding of companion cell developmental trajectories in *A. thaliana* depicting

cell-types (left) and pseudotime progression (right). Motifs, TFs, and ACRs with significant

association with pseudotime are shown as heatmaps below. **g,** Per-gene pseudotime shift scores

from alignments between *Z. mays* and *A. thaliana* companion cell development progressions,

clustered by k-means into three groups. Distribution of gene-gene distances from the alignments

are shown on the right, split by k-mean groups. **h,** Exemplary one-to-one homologs between *A.*

thaliana and *Z. mays* for the three groups split by pseudotime shifts. Acronyms: CC, companion

cell; LRC, lateral root cap; LRP, lateral root primordia. QC, quiescent center; SE, sieve elements.

 EXTENDED DATA FIGURE LEGENDS

Extended Data Fig. 1: Construction of single-cell chromatin accessibility profiles from a

diverse set of *Zea mays* organs.

a, Top; Genome browser screenshot of chromatin accessibility from bulk and single-cell

aggregate ATAC-seq experiments. Chromatin accessibility profiles depict the *tb1* locus and the

tb1 enhancer located approximately 67-kb upstream. Bottom; binary accessibility scores from a

random selection of 1,000 individual nuclei from each organ. **b,** Spearman's correlation coefficient

matrix comparing bulk and pseudobulk maize ATAC-seq samples across various organs. Sample

codes are shorthand for assay type, sample, and replicate. For example, s-R1 denotes single cell

assay for seminal root replicate 1. The term b-L2 denotes a bulk-ATAC assay for seedling

replicate 2. Codes are as follows: b, bulk; s, single cell; R, seminal root; C, crown root; E, ear; T,

tassel; A, axillary bud; L, seedling. Numbers represent replicate. **c,** Fragment length distributions

across 56,575 cells. The solid line and grey polygon represent the average and standard

deviation, respectively. **d,** Average TSS enrichment (normalized read depth adjusted by the two

windows 1-kb away from TSSs) across all 56,575 nuclei. Grey polygon denotes the standard

deviation. **e**, Genotype-mixing experimental schematic. **f,** Scatterplot of per cell B73 and Mo17

SNP counts from a mixed-genotype experiment (V1 seedlings) colored by genotype classification.

g, Posterior probabilities of individual barcodes (rows) highlighting the occurrence of cells with

B73, Mo17, and mixed (doublet) genotype identities. **h,** Genome browser screenshot of traditional

bulk ATAC-seq from 7 day old seedling (row 1), single-cell ATAC-seq from B73 seven day old

seedlings (row 2), pooled B73 and Mo17 nuclei (library ID: Seedling 2) single cell ATAC-seq from

7 day old seedlings (row 3), and the genotype-sorted B73 (row 4) and Mo17 (row 5) alignments

after sorting barcodes by genotype calls from the B73-Mo17 scATAC-seq 7 day old seedling

sample (row 3).

1020

Extended Data Fig. 2: Cell calling and barcode quality control.

a, Comparison of normalized (0-1) read depths at the union of all peaks across bulk and single-

cell samples (n=265,992) between replicated libraries, and between bulk and single-cell ATAC-

seq assays. **b,** Fragment length distributions for each library. Solid lines indicate the average

distribution across cells within the sample. Grey polygons represent the standard deviation across

nuclei in the library. **c,** Enrichment plots centered on 2-kb windows surrounding TSSs for barcodes

in each tissue. Grey polygons indicate the standard deviation across cells within the noted tissue.

d, Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5

integration sites mapping to within 2-kb of transcription start sites (TSSs). Dashed red lines

indicate the threshold of two standard deviations from the mean used to filter lower quality

barcodes. **e,** Knee plots illustrating $\log_{10}$ transformed cellular read depths of $\log_{10}$ ranked barcodes

across libraries. **f,** Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the

fraction of Tn5 integration sites derived from organellar sequences (chloroplast and mitochondrial)

relative to the total number of unique Tn5 integration sites associated with cognate barcode.

Dashed red lines indicate the threshold of two standard deviations from the mean used to filter

lower quality barcodes.

1037

Extended Data Fig. 3: Clustering metrics and comparison of bulk gene accessibility and

expression

a, Parameter regularization of model coefficients (y-axes) with respect to ACR usage (x-axes;

proportion of nuclei with at least one Tn5 integration site in an ACR). **b**, Proportion of variance

captured by the first 26 PCs. Inset: Spearman's correlation of principal components with cell read

depth ($\log_{10}$-transformed). **c**, Number of accessible sites per cell ($\log_{10}$). **d**, Proportion of Tn5

integrations within 2-kb of gene TSSs per cell. **e**, Co-localization of nuclei barcodes from different

biological replicates for three organs. **f**, Comparison of bulk RNA-seq expression levels (y-axis,

log$_2$ CPM) versus aggregate scATAC-seq gene accessibility scores (x-axis, log$_2$ CPM) within an

organ.

1048

Extended Data Fig. 4: Cell-type annotation and GO enrichment

a, UMAP embeddings of nuclei barcodes colored by low (grey) to high (dark purple) gene activity

values of cell-type specific marker genes. **b**, RNA *in situ* hybridization showing expression of

LOX10 in glume primordia and *GRFTF36* in IM and SMs of staminate inflorescence; *LRR445* in

the IM periphery and SPMs and *MYB89* in the IM and suppressed bract primordia of pistillate

inflorescence; Zm00001d038453 in ground tissue of SAM and leaf primordia sections. Gene

accessibility scores and predicted cell-types are shown on the right. *i*, tassel primordia. *ii,* ear

primordia. *iii*, SAM/leaf. Black triangles point to the glume primordia. Red triangles point to cryptic

bract primordia. Size bars illustrate 100-um. AM, axillary meristem; BS, bundle sheath; GC, guard

cell; GM, ground meristem; GMC, guard mother cell; GP, glume primordia; IM, inflorescence

meristem; L1, layer 1; LFM, lower floral meristem; SAM, shoot apical meristem; SBP, suppressed

bract primordia; SM, spikelet meristem; SPM; spikelet pair meristem; Stomatal PC, stomatal

precursor; UKN, unknown; VEMI, vascular/epidermal meristematic identity; VP, vascular

parenchyma; XP, xylem parenchyma. **c,** Proportion of cells within subcluster (column) derived

from one of six organs (rows). **d,** Distribution of GO term enrichment across clusters, where the

x-axis indicates the number of clusters in which a GO term is significantly enriched.

1065

Extended Data Fig. 5: Chromatin accessibility variation across plant cell-types

a, Row chromatin accessibility Z-scores across cell-types. Cell-type and subcluster labels are

located adjacent to the heatmap in identical order as the matrix. The first number after the cell-

type label indicates the major cluster number, while the second number represents the subcluster

ID. Legend colors: top left triangle, subcluster (corresponds to the sub-cluster annotation on

heatmap). bottom right triangle, major cluster (corresponds to the major cluster annotation on

heatmap). **b**, Row gene accessibility Z-scores across cell-types. **c**, Proportion of cells derived

from one of six organs within each cell-type/subcluster.

1074

Extended Data Fig. 6: Transcription factor motif variation underly dynamic accessible

regions

a, Comparison of cell-type specific ACR counts (by cell-type) and the number of unique Tn5

integration sites. **b,** Proportion of distal, genic and proximal ACRs per cell-type. **c**, Ranked TF

motif variability across cells, colored by TF family. **d**, Mean TF family motif enrichment (average

deviation scores per cell-type per TF family) across cell-types scaled to +/- 1. **e**, Spearman's

correlation coefficient (rho) between TF motifs (comparison of motif deviations across all nuclei).

Row and column colors represent TF motif families. **f,** TF motif deviations for 440 TF motifs

(columns) per nucleus (rows). The TF family for each corresponding motif is denoted by column

header colors. Cluster identification (sub-cluster/major cluster) are illustrated as row header colors

on the right side of the heatmap.   **g**, Relative gene accessibility scores for 2,423 maize

transcription factors (columns) for 50,640 nuclei (rows). Nuclei are sorted by sub-cluster cell-type

(row color labels). A cell-type annotation color legend applicable to panels **b**, **f**, and **g** where left

triangle reflects sub-cluster, right triangle indicates the major cluster.

1089

Extended Data Fig. 7: Co-accessible ACRs reflect in vivo chromatin interactions and are

driven by coordinated TF activity

a, Proportions of co-accessible ACR types, illustrated by toy examples. **b**, Proportion leaf Hi-C,

H3K4me3-HiChIP and H3K27me3-HiChIP chromatin loops that overlap co-accessible ACRs from

leaf cell-types (clusters with greater than 50% of cell derived from seedlings). **c**, Top, $\log_{10}$ number

of connections per ACR per cell-type from all co-accessible ACRs split by genomic context: distal,

proximal, and genic. Bottom, $\log_{10}$ number of connections per ACR per cell-type from cell-type

specific (purple) and constitutive (grey) co-accessible ACRs, split by genomic context. **d,**

Proportion of co-accessible classifications by cell-type for all co-accessible ACRs (top) and cell-

type specific co-accessible ACRs (bottom). **e**, Jaccard similarity of motif composition between co-

accessible ACR edges by cell-type (colored diamonds) relative to the same number of random

ACR-ACR links, permuted 1,000 times for each cell-type (grey boxplots). Box plots represent the

interquartile range, grey lines indicate the permuted range. **f**, Heatmaps of observed proportion

of co-accessible ACRs with the same motif embedded within link edges subtracted and divided

by the expected proportion estimated by 1,000 permutations using random ACR-ACR links with

the same number of co-accessible ACRs. **g,** Exemplary motifs enriched in reciprocal co-

accessible ACRs for *TCP, AP2-EREB,* and *LBD* TF families.

1107

Extended Data Fig. 8: Pseudotime trajectory construction

a, Overview of pseudotime developmental trajectory analysis from four organs: root, seedling,

tassel (staminate inflorescence), and ear (pistilate inflorescence). **b,** Endodermis development in

crown roots. **c,** Cortex development in crown roots. **d,** Pericycle development in embryonic roots.

e, Atrichoblast development in crown roots. **f,** Trichoblast development in crown roots. **g,**

Trichoblast development in embryonic roots. **h,** Lateral root cap (LRC) development in crown

roots. **i,** Lateral root cap (LRC) development in embryonic roots. **j,** Phloem sieve element (SE)

development in crown roots. **k,** Companion cell development in crown roots. **l,** Phloem sieve

element (SE) development in embryonic roots. **m,** Xylem development in crown roots. **n,** Xylem

development in embryonic roots. **o,** Procambial development in crown roots. **p,** Guard cell

development in seedling. **q,** Subsidiary cell development in seedlings. **r,** Floral primordia

development in staminate inflorescence (tassel). **s,** Floral primordia development in pistilate

inflorescence (ear).

1121

1122 **Extended Data Fig. 9: *Arabidopsis thaliana* root cell-type atlas**

a, Knee plots for *A. thaliana* root samples illustrating $\log_{10}$ transformed cellular read depths of $\log_{10}$ ranked barcodes across two biological replicates. **b,** Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites derived from organellar sequences (chloroplast and mitochondrial) relative to the total number of unique Tn5 integration sites associated with each barcode from the two biological replicates. Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes. **c,** Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites mapping to within 2-kb of transcription start sites (TSSs). Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes. **d,** Average TSS enrichment (normalized read depth adjusted by the two 10-bp windows 1-kb away from TSSs) across 5,001 *A. thaliana* root barcodes (rows). **e,** UMAP (Uniform manifold approximation projection) embeddings of *A. thaliana* root barcodes colored by biological replicate. **f,** total number of accessible chromatin regions (ACRs). **g,** the proportion of Tn5 integration sites within 1-kb of TSSs. **h,** Relative gene accessibility for 27 known cell-type/domain restricted marker genes used to inform cell-type annotation of *A. thaliana* root clusters. **i,** Relative motif deviations for transcription factors with known cell-type specificities.

Extended Data Fig. 10: Dynamic and conserved chromatin accessibility across pseudotime between *Arabidopsis thaliana* and *Zea mays*

a, Pseudotime trajectories for Atrichoblast, Trichoblast, Lateral root cap (LRC), Cortex, Endodermis, Lateral root primordia (LRP), Companion cells (CC), and Xylem development. **b,** Averaged alignments of conserved, shift early *Z. mays*, and shift early *A. thaliana* putative orthologs. **c,** Pseudotime shifts of TF motifs between *A. thaliana* and *Z. mays*, clustered into k-means and conserved groups. **d,** Distributions of motif-motif normalized distances between *Z. mays* and *A. thaliana* for the three groups. **e,** Conserved motifs (n=142) ordered by pseudotime. Heatmaps for *A. thaliana* and *Z. mays* have identical row orders. **f,** Averaged alignments of

1149 conserved, shift early *Z. mays*, and shift early *A. thaliana* groups based on motif-motif global

1150 alignments from the dynamic time-warping algorithm. **g,** Examples of conserved, shift early *Z.*

1151 *mays*, and shift early *A. thaliana* motifs from both species. **h,** Gene expression Z-scores across

1152 *A. thaliana* FAC sorted root cell-types for the TFs recognizing the top four conserved motifs.

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

REFERENCES

1175

1   Cusanovich, D. A. *et al.* A Single-Cell Atlas of In Vivo Mammalian Chromatin Accessibility.
Cell **174**, 1309-1324 e1318, doi:10.1016/j.cell.2018.06.052 (2018).

2   Marand, A. P., Zhang, T., Zhu, B. & Jiang, J. Towards genome-wide prediction and
characterization of enhancers in plants. *Biochim Biophys Acta Gene Regul Mech* **1860**,
131-139, doi:10.1016/j.bbagrm.2016.06.006 (2017).

3   Teale, W. D., Paponov, I. A. & Palme, K. Auxin in action: signalling, transport and the
control of plant growth and development. *Nat Rev Mol Cell Biol* **7**, 847-859,
doi:10.1038/nrm2020 (2006).

4   Wittkopp, P. J. & Kalay, G. Cis-regulatory elements: molecular mechanisms and
evolutionary processes underlying divergence. *Nat Rev Genet* **13**, 59-69,
doi:10.1038/nrg3095 (2011).

5   Lu, Z. *et al.* The prevalence, evolution and chromatin signatures of plant regulatory
elements. *Nat Plants* **5**, 1250-1259, doi:10.1038/s41477-019-0548-z (2019).

6   Andersson, R. & Sandelin, A. Determinants of enhancer and promoter activities of
regulatory elements. *Nat Rev Genet* **21**, 71-87, doi:10.1038/s41576-019-0173-8 (2020).

7   Long, H. K., Prescott, S. L. & Wysocka, J. Ever-Changing Landscapes: Transcriptional
Enhancers in Development and Evolution. *Cell* **167**, 1170-1187,
doi:10.1016/j.cell.2016.09.018 (2016).

8   Chang, Y. M. *et al.* Characterizing regulatory and functional differentiation between maize
mesophyll and bundle sheath cells by transcriptomic analysis. *Plant Physiol* **160**, 165-177,
doi:10.1104/pp.112.203810 (2012).

9   Langdale, J. A., Lane, B., Freeling, M. & Nelson, T. Cell lineage analysis of maize bundle
sheath and mesophyll cells. *Dev Biol* **133**, 128-139, doi:10.1016/0012-1606(89)90304-7
(1989).

10 Ricci, W. A. *et al.* Widespread long-range cis-regulatory elements in the maize genome.
Nat Plants **5**, 1237-1249, doi:10.1038/s41477-019-0547-0 (2019).

11 Oka, R. *et al.* Genome-wide mapping of transcriptional enhancer candidates using DNA
and chromatin features in maize. *Genome Biol* **18**, 137, doi:10.1186/s13059-017-1273-4
(2017).

12 Crisp, P. A. *et al.* Stable unmethylated DNA demarcates expressed genes and their cis-
regulatory space in plant genomes. *Proc Natl Acad Sci U S A*,
doi:10.1073/pnas.2010250117 (2020).

13 Zhao, H. *et al.* Proliferation of Regulatory DNA Elements Derived from Transposable
Elements in the Maize Genome. *Plant Physiol* **176**, 2789-2803, doi:10.1104/pp.17.01467
(2018).

14   Clark, R. M., Wagler, T. N., Quijada, P. & Doebley, J. A distant upstream enhancer at the
maize domestication gene tb1 has pleiotropic effects on plant and inflorescent
architecture. *Nat Genet* **38**, 594-597, doi:10.1038/ng1784 (2006).

15   Noshay, J. M. *et al.* Cis-regulatory elements within TEs can influence expression of nearby
maize genes. *bioRxiv*, 2020.2005.2020.107169, doi:10.1101/2020.05.20.107169 (2020).

16   Rebeiz, M. & Tsiantis, M. Enhancer evolution and the origins of morphological novelty.
Curr Opin Genet Dev **45**, 115-123, doi:10.1016/j.gde.2017.04.006 (2017).

17   Villar, D. *et al.* Enhancer evolution across 20 mammalian species. *Cell* **160**, 554-566,
doi:10.1016/j.cell.2015.01.006 (2015).

18   Wallace, J. G. *et al.* Association mapping across numerous traits reveals patterns of
functional variation in maize. *PLoS Genet* **10**, e1004845,
doi:10.1371/journal.pgen.1004845 (2014).

19   Wang, B. *et al.* Genome-wide selection and genetic improvement during modern maize
breeding. *Nat Genet* **52**, 565-571, doi:10.1038/s41588-020-0616-3 (2020).

20   Gage, J. L., White, M. R., Edwards, J. W., Kaeppler, S. & de Leon, N. Selection Signatures
Underlying Dramatic Male Inflorescence Transformation During Modern Hybrid Maize
Breeding. *Genetics* **210**, 1125-1138, doi:10.1534/genetics.118.301487 (2018).

21   Verweij, W. *et al.* Functionally Similar WRKY Proteins Regulate Vacuolar Acidification in
Petunia and Hair Development in Arabidopsis. *Plant Cell* **28**, 786-803,
doi:10.1105/tpc.15.00608 (2016).

22   Gomez-Mena, C., de Folter, S., Costa, M. M., Angenent, G. C. & Sablowski, R.
Transcriptional program controlled by the floral homeotic gene AGAMOUS during early
organogenesis. *Development* **132**, 429-438, doi:10.1242/dev.01600 (2005).

23   Pliner, H. A. *et al.* Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell
Chromatin Accessibility Data. *Mol Cell* **71**, 858-+, doi:10.1016/j.molcel.2018.06.044
(2018).

24   Gate, R. E. *et al.* Genetic determinants of co-accessible chromatin regions in activated T
cells across humans. *Nat Genet* **50**, 1140-1150, doi:10.1038/s41588-018-0156-2 (2018).

25   Satpathy, A. T. *et al.* Massively parallel single-cell chromatin landscapes of human
immune cell development and intratumoral T cell exhaustion. *Nat Biotechnol* **37**, 925-936,
doi:10.1038/s41587-019-0206-z (2019).

26   Buenrostro, J. D. *et al.* Single-cell chromatin accessibility reveals principles of regulatory
variation. *Nature* **523**, 486-490, doi:10.1038/nature14590 (2015).

27   Peng, Y. *et al.* Chromatin interaction maps reveal genetic regulation for quantitative traits
in maize. *Nat Commun* **10**, 2632, doi:10.1038/s41467-019-10602-5 (2019).

28   Sun, Y. *et al.* 3D genome architecture coordinates trans and cis regulation of differentially
expressed ear and tassel genes in maize. *Genome Biol* **21**, 143, doi:10.1186/s13059-020-
02063-7 (2020).

29   Chuck, G. S., Brown, P. J., Meeley, R. & Hake, S. Maize SBP-box transcription factors
unbranched2 and unbranched3 affect yield traits by regulating the rate of lateral primordia
initiation. *Proc Natl Acad Sci U S A* **111**, 18775-18780, doi:10.1073/pnas.1407401112
(2014).

30   Heger, P., Marin, B., Bartkuhn, M., Schierenberg, E. & Wiehe, T. The chromatin insulator
CTCF and the emergence of metazoan diversity. *Proc Natl Acad Sci U S A* **109**, 17507-
17512, doi:10.1073/pnas.1111941109 (2012).

31   Liu, C., Cheng, Y. J., Wang, J. W. & Weigel, D. Prominent topologically associated
domains differentiate global chromatin packing in rice from Arabidopsis. *Nat Plants* **3**, 742-
748, doi:10.1038/s41477-017-0005-9 (2017).

32   Karaaslan, E. S. *et al.* Marchantia TCP transcription factor activity correlates with three-
dimensional chromatin structure. *Nat Plants*, doi:10.1038/s41477-020-00766-0 (2020).

33   Takacs, E. M. *et al.* Ontogeny of the maize shoot apical meristem. *Plant Cell* **24**, 3219-
3234, doi:10.1105/tpc.112.099614 (2012).

34   Somssich, M., Je, B. I., Simon, R. & Jackson, D. CLAVATA-WUSCHEL signaling in the
shoot meristem. *Development* **143**, 3238-3248, doi:10.1242/dev.133645 (2016).

35   Jiang, K., Zhu, T., Diao, Z., Huang, H. & Feldman, L. J. The maize root stem cell niche: a
partnership between two sister cell populations. *Planta* **231**, 411-424,
doi:10.1007/s00425-009-1059-3 (2010).

36   Wallner, E. S. *et al.* Strigolactone- and Karrikin-Independent SMXL Proteins Are Central
Regulators of Phloem Formation. *Curr Biol* **27**, 1241-1247, doi:10.1016/j.cub.2017.03.014
(2017).

37   Baker, R. F. *et al.* Sucrose Transporter ZmSut1 Expression and Localization Uncover New
Insights into Sucrose Phloem Loading. *Plant Physiol* **172**, 1876-1898,
doi:10.1104/pp.16.00884 (2016).

38   Kajala, K. *et al.* Innovation, conservation and repurposing of gene function in plant root
cell type development. *bioRxiv*, 2020.2004.2009.017285, doi:10.1101/2020.04.09.017285
(2020).

39   Brady, S. M. *et al.* A high-resolution root spatiotemporal map reveals dominant expression
patterns. *Science* **318**, 801-806, doi:10.1126/science.1146265 (2007).

40   You, Y. *et al.* Phloem Companion Cell-Specific Transcriptomic and Epigenomic Analyses
Identify MRF1, a Regulator of Flowering. *Plant Cell* **31**, 325-345,
doi:10.1105/tpc.17.00714 (2019).

41   Jiang, H., Li, H., Bu, Q. & Li, C. The RHA2a-interacting proteins ANAC019 and ANAC055
may play a dual role in regulating ABA response and jasmonate response. *Plant Signaling*
& Behavior **4**, 464-466, doi:10.4161/psb.4.5.8543 (2009).

42   Yamaguchi, M. *et al.* VND-INTERACTING2, a NAC domain transcription factor, negatively
regulates xylem vessel formation in Arabidopsis. *Plant Cell* **22**, 1249-1263,
doi:10.1105/tpc.108.064048 (2010).

43   Jiao, Y. *et al.* Improved maize reference genome with single-molecule technologies.
Nature **546**, 524-527, doi:10.1038/nature22971 (2017).

44   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-
2079, doi:10.1093/bioinformatics/btp352 (2009).

45   AFA Smit, R. H., and P Green. *RepeatMasker Open-4.0* (2013-2015).

46   Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421,
doi:10.1186/1471-2105-10-421 (2009).

47   Team, R. C. *R: A language and environment for statistical computing.*, <http://www.R-
project.org/> (2013).

48   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884-i890, doi:10.1093/bioinformatics/bty560 (2018).

49   Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv e-prints, arXiv:1303.3997 (2013).
<https://ui.adsabs.harvard.edu/abs/2013arXiv1303.3997L>.

50   Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.
arXiv e-prints, arXiv:1207.3907 (2012).
<https://ui.adsabs.harvard.edu/abs/2012arXiv1207.3907G>.

51   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137,
doi:10.1186/gb-2008-9-9-r137 (2008).

52   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
seq data using regularized negative binomial regression. *Genome Biol* **20**, doi:ARTN 296
10.1186/s13059-019-1874-1 (2019).

53   Witten, D. M., Tibshirani, R. & Hastie, T. A penalized matrix decomposition, with
applications to sparse principal components and canonical correlation analysis.
Biostatistics **10**, 515-534, doi:10.1093/biostatistics/kxp008 (2009).

54   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-+,
doi:10.1016/j.cell.2019.05.031 (2019).

55   Corces, M. R. *et al.* The chromatin accessibility landscape of primary human cancers.
Science **362**, doi:10.1126/science.aav1898 (2018).

56   Fang, R. *et al.* SnapATAC: A Comprehensive Analysis Package for Single Cell ATAC-
seq. *bioRxiv*, 615179, doi:10.1101/615179 (2020).

57   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-
140, doi:10.1093/bioinformatics/btp616 (2010).

58   Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif.
Bioinformatics **27**, 1017-1018, doi:10.1093/bioinformatics/btr064 (2011).

59   O'Malley, R. C. *et al.* Cistrome and Epicistrome Features Shape the Regulatory DNA
Landscape. *Cell* **166**, 1598, doi:10.1016/j.cell.2016.08.063 (2016).

60   Galli, M. *et al.* The DNA binding landscape of the maize AUXIN RESPONSE FACTOR
family. *Nat Commun* **9**, 4526, doi:10.1038/s41467-018-06977-6 (2018).

61   Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat
Methods* **14**, 975-978, doi:10.1038/nmeth.4401 (2017).

62   Granja, J. M. *et al.* ArchR: An integrative and scalable software package for single-cell
chromatin accessibility analysis. *bioRxiv*, 2020.2004.2028.066498,
doi:10.1101/2020.04.28.066498 (2020).

63   Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol* **20**, 238, doi:10.1186/s13059-019-1832-y (2019).

64   van Dijk, D. *et al.* Recovering Gene Interactions from Single-Cell Data Using Data
Diffusion. *Cell* **174**, 716-729 e727, doi:10.1016/j.cell.2018.05.061 (2018).

65   Alpert, A., Moore, L. S., Dubovik, T. & Shen-Orr, S. S. Alignment of single-cell trajectories
to compare cellular expression dynamics. *Nat Methods* **15**, 267-270,
doi:10.1038/nmeth.4628 (2018).

66   Hofmeister, B. T. & Schmitz, R. J. Enhanced JBrowse plugins for epigenomics data
visualization. *BMC Bioinformatics* **19**, 159, doi:10.1186/s12859-018-2160-z (2018).
1343