A *cis*-regulatory atlas in maize at single-cell resolution

Alexandre P. Marand[1]*, Zongliang Chen[2], Andrea Gallavotti[2,3], and Robert J. Schmitz[1]*

Affiliations:
[1]Department of Genetics, University of Georgia, Athens, GA 30602, USA
[2]Waksman Institute, Rutgers University, Piscataway, NJ, 08854, USA
[3]Department of Plant Biology, Rutgers University, New Brunswick, NJ, 08901, USA

Correspondence: Alexandre P. Marand (marand@uga.edu), Robert J. Schmitz (schmitz@uga.edu)

Keywords: chromatin, single-cell, epigenomics, maize, gene regulation, ATAC-seq

SUMMARY

Cis-regulatory elements (CREs) encode the genomic blueprints of spatiotemporal gene

expression programs enabling highly specialized cell functions. To identify CREs at cell-type

resolution in *Zea mays*, we implemented single-cell sequencing of Assay for Transposase

Accessible Chromatin (scATAC-seq) in seedlings, embryonic roots, crown roots, axillary buds,

and pistillate and staminate inflorescence. We describe 92 states of chromatin accessibility across

165,913 putative CREs and 52 known cell types. Patterns of transcription factor (TF) motif

accessibility predicted cell identity with high accuracy, uncovered putative non-cell autonomous

TFs, and revealed TF motifs underlying higher-order chromatin interactions. Comparison of maize

and *Arabidopsis thaliana* developmental trajectories identified TF motifs with conserved patterns

of accessibility. Cell type-specific CREs were enriched with enhancer activity, phenotype-

associated genetic variants, and signatures of breeding-era selection. These data, along with

companion software, *Socrates*, afford a comprehensive framework for understanding cellular

heterogeneity, evolution, and *cis*-regulatory grammar of cell-type specification in a major crop.

INTRODUCTION

Global consumption of maize per kilogram per person is expected to increase by 163% by 2050 (CIMMYT 2016). However, climate instability and disease strain are increasingly predicted to lower global maize yields by more than 10% over the same time frame (Tigchelaar et al., 2018; Zhao et al., 2017). As the foundational unit of plants, individual cells are responsible for the synthesis, transportation, and storage of rich primary and secondary metabolites that sequesters carbon and provide human nourishment. However, our understanding of cell-type functions in plants has been precluded by technical limitations imposed by the cell wall and an inability to culture homogenous cell lines, in contrast to mammalian models.

Past studies querying plant cell-type functions employed technically challenging experimental procedures to circumvent these obstacles, such as fluorescence-activated cell sorting (FACS) of GFP-tagged marker proteins, or isolation of nuclei tagged in specific cell types (INTACT) (Birnbaum et al., 2003; Brady et al., 2007; Deal and Henikoff, 2011). Although instrumental to understanding certain cell types, a shortcoming of these methods is the requirement of transgenesis and prior information regarding cell-type specificity for purification, thereby occluding unbiased efforts for discovery of unknown and poorly studied cell types. As a result, molecular profiling on a genome-wide scale of individual cells and cell types in plants have been largely limited to the roots of *Arabidopsis thaliana* and a handful of isolated tissues (Dorrity et al., 2020; Farmer et al., 2020; Jean-Baptiste et al., 2019; Lee et al., 2019; Lopez-Anido et al., 2020; Nelms and Walbot, 2019; Ryu et al., 2019; Shulse et al., 2019). Although critical for driving innovation in biotechnology, a comprehensive organismal cell-type atlas has yet to be realized in any plant species.

Development, differentiation, and response to environment in eukaryotic cells rely on precise spatiotemporal gene expression mediated by *cis*-regulatory elements (CREs) (Andersson and Sandelin, 2020; Cusanovich et al., 2018; Long et al., 2016; Lu et al., 2019; Marand et al., 2017; Teale et al., 2006; Wittkopp and Kalay, 2011). CREs encode DNA binding sites for

transcription factors (TF) that cooperatively dictate transcriptional outcomes (Buchler et al., 2003;

Cheng et al., 2012; Consortium, 2012; Gerstein et al., 2012; Ravasi et al., 2010). In metazoan

genomes, CCCTC-binding factor (CTCF) directs higher-order chromatin interactions that facilitate

spatial proximity of CREs and their target genes (Phillips and Corces, 2009). Plant genomes are

also generalized by higher-order chromatin architecture, yet all plant lineages lack an ortholog to

CTCF (Heger et al., 2012). How cells interpret the *cis*-regulatory code, establish diverse chromatin

contact landscapes, and adopt specialized functions in discrete cell types are essential questions

for understanding the rules governing biology. As a consequence of their centrality in establishing

cell identity and function, a growing body of evidence point to genetic variation of CREs as a major

source of phenotypic innovation, including disease and evolutionary divergence (Rebeiz and

Tsiantis, 2017; Villar et al., 2015). However, it has become increasingly apparent that genetic

variants may only affect CRE activity in a subset of cell types (Hekselman and Yeger-Lotem,

2020). We reasoned that a thorough investigation comprising the full spectrum of evolutionary

changes through both inter- and intraspecies comparisons would be informative for detangling

the cell type-specific contributions towards phenotypic variation.

Here, we describe the construction of a *cis*-regulatory atlas in the historically rich genetic

model and crop species, *Zea mays*. We measure chromatin accessibility and nuclear gene

expression in 72,090 cells across six major maize organs. Model-based normalization of

chromatin accessibility enabled the identification and validation of diverse cell types, many of

which lacked previous genome-wide characterization. We define the *cis*-regulatory combinatorial

grammar underlying cell identity, reveal distinct TFs coordinating higher-order chromatin

interactions, and demonstrate enhancer CREs with increased capacity for interactions as major

contributors to phenotypic variation. Through an evolutionary lens, we uncover CREs and cell

types targeted by modern breeding and evaluate the evolutionary impacts on *cis*-regulatory

specification of cellular development between two highly diverged angiosperms (maize and *A.*

thaliana). Finally, we present the R package "*Socrates*", a unified framework for scATAC-seq pre-

processing, normalization, downstream analysis, and integration with scRNA-seq data as a

streamlined method for single-cell genomic studies.

RESULTS

Assembly and validation of a high-quality *cis*-regulatory atlas in maize

To comprehensively assess *cis*-regulatory variation among cell types in a major crop species, we isolated nuclei using fluorescence-activated nuclei sorting (FANS) and generated single-cell chromatin accessibility profiles using Assay for Transposase Accessible Chromatin (scATAC-seq). Libraries were prepared from six major *Z. mays* L. cultivar B73 organs, including axillary buds, staminate and pistillate inflorescence, whole seedling (composed of stem, leaf, and coleoptile tissues), embryonic root tips, and post-embryonic crown roots as a representative sample of *cis*-regulatory diversity across a suite of maize cell types and tissues (**Figure 1A**, **S1A**, and **S1B**; **Table S1**).

We evaluated several quantitative and qualitative metrics reflective of high quality scATAC-seq data. In aggregate, scATAC-seq libraries were highly correlated between biological replicates and with previously generated organ-matched bulk ATAC-seq data (**Figure S1B**, **S2A**). Individual nuclei exhibited strong enrichment at transcription start sites (TSSs) (**Figure S1D**, **S2B** and **S2C**) and were consistent with the expected distributions of nucleosome-free and nucleosome-protected fragments (**Figure S1E** and **S2D**). By genotyping individual nuclei from a pooled population composed of B73 and Mo17 genotypes, we found 96% (4,944/5,177) were representative of a single genotype, validating our experimental approach (**Figure S1F-S1I**; **STAR Methods**). We identified a total of 56,575 nuclei passing quality filters (range: 4,704 - 18,393 nuclei per organ) with an average of 31,660 unique Tn5 integration sites per nucleus (**Figure S2C**, **S2E** and **S2F**; **Table S2**; **STAR Methods**).

Towards identifying clusters of nuclei resembling cell types, we first identified accessible chromatin regions (ACRs) by *in silico* sorting, resulting in a catalog of 165,913 putative CREs covering ~4% of the maize genome (**Figure S3**; **STAR Methods**). To enable species-agnostic model-based analysis of scATAC-seq data, we developed an R package, termed '*Socrates',* that streamlines data processing, clustering and downstream analysis. At its heart, '*Socrates*'

Figure 1. Atlas-scale cell-type profiling from single nuclei chromatin accessibility in *Zea mays*

(A) Overview of experimental samples, with an example of the cell-type diversity present in seedlings.

(B) Nuclei similarity clustering as a UMAP embedding derived from the denoised quasibinomial Pearson's residuals across all ACRs for each nucleus. UMAP embedding of nuclei colored by organ identity.

(C) UMAP embedding of nuclei colored major cluster identity.

(D) UMAP embedding of sub-cluster assignments following a second round of clustering within each major cluster. Sub-cluster color reflects the organ with the greatest proportion of nuclei in the cluster. See panel (B) for color code.

(E) Cell type-specific enrichment of gene accessibility for a subset of marker genes associated with six different cell types.

(F) Sub-cluster-specific chromatin accessibility profiles surrounding known marker genes for floral primordia, xylem precursors, and L1 epidermal cells. Bold, circled numbers indicate the cognate major cluster shown in panel c. Sub-cluster numeric identifications are present on the sides of the coverage plots.

(G) Top, gene accessibility for *ZmGRFTF36*, an inflorescence and spikelet meristem enriched transcription factor with no previously known cell-type specificity. Bottom, RNA *in situ* hybridization of *ZmGRFTF36* in maize B73 staminate (tassel) primordia. FP, floral primordia; GP, glume primordia; IM, inflorescence meristem; LFM, lower floral meristem; SM, spikelet meristem; SPM, spikelet pair meristem; UKN, unknown.

implements a regularized quasibinomial logistic regression framework to remove unwanted

variation stemming from differences in nuclei read depth or other experimental factors (**Figure**

S4A; **STAR Methods**). Following normalization with *Socrates*, we projected nuclei into a two-

dimensional space using Uniform Manifold Approximation Projection (UMAP), revealing 10 major

clusters unbiased by technical variation (**Figure 1B**, **1C**, and **S4B-S4E**).

Consistent with functional diversification of spatially distinct cells, most major clusters were

generally composed of nuclei from the same organ (**Figure 1B** and **1C**). However, we also found

evidence of common cell identities from nuclei in different organs, such as the co-localization of

pistillate inflorescence, staminate inflorescence, and seedling nuclei in clusters 2, 4, 7 and 8

(**Figure 1B** and **1C**). Apparent heterogeneity within major clusters prompted us to implement a

second round of partitioning for each major grouping, producing a total of 92 sub-clusters

(hereafter referred to as clusters) with an average of 551 nuclei (**Figure 1D**; **STAR Methods**).

Clear reproducibility and mitigation of technical variation in the UMAP embedding justifies

'*Socrates'* as a robust approach for establishing shared cell identities across heterogenous organs

through the removal of technical variation typical of single-cell experiments.

Cell-type annotation and validation by *in situ* hybridization

To annotate clusters with corresponding cell types, we integrated chromatin accessibility

information on a per-gene and nucleus basis as a proxy for gene expression (bulk RNA-seq

versus aggregate scATAC-seq Spearman's correlation coefficient = 0.54-0.58; **Figure S4F**;

STAR Methods). We then (*i*) evaluated differential accessibility among clusters for a manually

curated list of 221 literature-derived known marker genes (**Table S3**), (*ii*) classified cell types of

individual nuclei with a multinomial logistic classifier trained on nuclei with discriminative cell type-

specific signatures, and visually assessed (*iii*) accessibility scores of *a priori* marker genes over

the UMAP embeddings and (*iv*) cluster-aggregated ATAC-seq coverages (**Figure 1E**, **1F**, **S5A**;

Table S4; **STAR Methods**). Patterns of gene accessibility were consistent with *a priori*

information of cell type/domain-specific expression, such as co-localized accessibility of bundle

sheath-specific genes *DICARBOXYLIC ACID TRANSPORTER1* (*DCT2*) and *RIBULOSE*

BISPHOSPHATE CARBOXYLASE SMALL SUBUNIT2 (*SSU2*), and mesophyll-specific genes

MALATE DEHYDROGENASE6 (*MDH6*) and *PYRUVATE DEHYDOGENASE KINASE1* (*PDK1*)*,*

in addition to many other previously established cell type-specific marker genes (**Figure 1E**, **1F**,

and **S5A**) (Chang et al., 2012).

To corroborate predicted cell-type annotations, we performed RNA *in situ* hybridization for

a subset of differentially accessible genes with no prior evidence of cell type-specificity. In all

cases (5/5), *in situ* expression patterns were in line with the predicted localization based on gene

accessibility (**Figure 1G** and **S5B**). Estimates of cell-type proportions within and across organs

were also concordant with prior observations, such as the occurrence of vascular bundle sheath

and parenchymal mesophyll cells within multiple organs, including those derived from stem and

leaf tissues of seedlings and within pistillate and staminate inflorescence (**Figure S5C**; **Table S4**)

(Langdale et al., 1989).

Integration of chromatin accessibility and gene expression from individual nuclei

To evaluate the correspondence between nuclear transcription and chromatin accessibility on a

global scale, we sequenced the transcriptomes of 15,515 nuclei derived from 7-day old seedlings

using single-nucleus RNA-seq (snRNA-seq; **STAR Methods**). We then integrated the

corresponding nuclei with seedling-derived nuclei from scATAC-seq via integrative non-negative

matrix factorization (iNMF; **Figure 2A**; **STAR Methods**) (Welch et al., 2019). Co-embedding

nuclei on the basis of chromatin (n=11,882) and nuclear gene expression (n=15,515) revealed 19

clusters of nuclei with similar genome-wide profiles (**Figure 2B** and **2C**; **STAR Methods**).

Comparison of the two modalities (n=36,322 genes) across clusters revealed a striking

correspondence between the patterns of chromatin accessibility and nuclear transcription

(Spearman's correlation coefficient range across cell types = [0.52-0.69]; **Figure 2D**). The

Figure 2. Concordance between chromatin accessibility and gene expression at single-nuclei resolution

(A) Illustration of iNMF integration of scATAC-seq and snRNA-seq seedling data sets.

(B) UMAP co-embedding of seedling nuclei from scATAC-seq (n=11,882; purple) and snRNA-seq (n=15,515; blue).

(C) Louvain clustering and cell-type annotations for co-embedded seedling nuclei.

(D) Comparison of within-cluster averaged gene accessibility (left) and gene expression (right) between cell types/clusters. Column color legend corresponds to the cell-type colors specified in panel C.

(E) UMAP embeddings displaying per nucleus gene accessibility (top, n=11,882) and gene expression (bottom, n=15,515) values for five cell type-specific marker genes.

(F) Left: Aggregate scATAC-seq tracks across clusters at the *CAH1* locus. Right: Average expression of *CAH1*.

(G) Spearman correlations between clusters based on nuclear transcription (snRNA) and chromatin accessibility (scATAC).

(H) Density scatter plot comparing gene accessibility (x-axis) and expression (y-axis) for each cluster and gene.

(I) Expression (left), chromatin accessibility (middle), and H3K27me3 ChIP-seq meta-profiles (relative reads per million, RPM) of accessible/expressed genes (turquoise; n=19,402), accessible/non-expressed genes (pink; n=6,063) and non-accessible/non-expressed genes (grey, n=4,315).

(J) Top two *de novo* motifs enriched in ACRs within 1-kb of accessible/non-expressed genes (pink, panel H and I).

molecular relationship between chromatin accessibility and gene expression was further

exemplified by marker genes with recognized cell-type specificity, including *DCT2* (bundle

sheath), *CARBONIC ANHYDRASE1* (*CAH1*; mesophyll), *POTASSIUM CHANNEL 3 (KCH3;*

guard cell), *GLYCEROL-3-PHOSPHATE ACYLTRANSFERASE 12* (*GPAT12*; epidermis), and

homolog to *GLABRA3* (*ZmGL3*; trichome) (**Figure 2E** and **2F**). Comparison of aggregated cell-

type profiles indicated a greater extent of variation in chromatin accessibility relative to gene

expression, suggesting chromatin structure provides additional information for dissecting cell-type

heterogeneity (**Figure 2G**).

Despite the strong association between gene accessibility and expression, we observed

a subset of accessible genes lacking evidence of transcription that were highly enriched with

H3K27me3 (**Figure 2H** and **2I**). As we defined gene accessibility to include upstream sequences,

we posited that ACRs associated with accessible and silenced genes might contain Polycomb

Response Elements (PREs) directing transcriptional silencing via deposition of H3K27me3 by the

Polycomb Repression Complex (PRC). *De novo* motif analysis of 15,073 ACRs within 1-kb of

accessible and silenced genes (n=6,063) identified several enriched motifs, including a CNN-

repeat (E-value < 2.0e-738, 83% of ACRs, 12,858/15,073) and a CTGCAG palindromic motif (E-

value < 2.4e-205, 80% of ACRs, 12,014/15,703) (**Figure 2J**, **STAR Methods**). A query with

experimentally established TF binding sites revealed a significant (FDR < 4.09e-3) overlap

between the CNN-repeat motif and sequences recognized by BASIC PENTACYSTEINE1

(BPC1), a BARLEY B RECOMBINANT-BASIC PENTACYSTEINE (BBR-BPC) family TF

previously associated with PREs and H3K27me3-mediated silencing in *A. thaliana* (Xiao et al.,

2017) (**STAR Methods**). Taken together, we establish gene accessibility as a robust proxy for

transcription and suggest the activity of PREs as a possible explanation for imperfect correlations

between gene accessibility and expression.

To comprehensively investigate the extent of gene accessibility variation, we performed

differentially accessibility hypothesis testing for each gene model across cell types (**STAR**

Methods). After multiple test correction and heuristic thresholding (FDR < 0.05, log fold-change

> 2), we identified 74% (28,625/38,752) of genes with significant differential accessibility in at

least one cell type, with an average of 2,768 differentially accessible genes per cluster (**Figure**

S6A; **Table S5**). Marker-agnostic gene set enrichment analysis (GSEA) of Gene Ontology (GO)

terms exemplified prior information regarding specific cell-type functions, with enriched terms

such as "root hair cell development" in root epidermal initials, "regulation of stomatal closure" in

subsidiary cells, and "malate transmembrane transport" for mesophyll cells (**Figure S6D**). Distinct

cell types were generalized by highly specific GO annotations, as most (>51%) GO terms were

identified in only a handful of cell types (five or fewer), implicating chromatin accessibility

dynamics as underling the signature hallmarks of cell-type identity and function (**Figure S6D**). In

summary, we identified 52 cell types for 83% (76/92) of scATAC-seq clusters, capturing nearly all

major expected cell types in the profiled organs and suggesting the existence of novel

uncharacterized cell types present in these data (**Table S4**).

Characterization of *cis*-regulatory variation

Deconvolution of nuclei into discrete cell types provides an opportunity to identify CREs encoding

cell identity. To this end, we implemented a generalized linear model to catalog ACRs with

discrete patterns of chromatin accessibility across cell types (**STAR Methods**). In total, 52,520

ACRs (31%) were differentially accessible (FDR < 0.05, fold-change > 2) and restricted to one or

a handful of clusters, with an average of 2,826 per-cluster (**Figure S6B**, **S7A**, and **S7B**). Similar

to the reported functions of transcriptional enhancers in metazoan genomes, we found cluster-

specific ACRs were associated with significantly greater enhancer activity based on Self-

Transcribing Active Regulatory Region sequencing (STARR-seq) of maize leaf protoplasts (Ricci

et al., 2019) relative to controls (n=165,913) and non-specific ACRs (n=113,393; **Figure 3A**;

STAR Methods). Deconvolution of chromatin accessibility by cell type revealed accessible sites

primarily located distal to genic regions (>2-kb from any gene) compared to previously published

Figure 3. Characterization of accessible chromatin regions

(A) Distribution of enhancer activity (maximum log2[RNA/input]) for control regions (n=165,913), non-specific (n=113,393) and cluster-specific ACRs (n=52,520). The dash red line indicates the overall mean. Orange lines reflect differences between the group median and overall mean.

(B) Distribution of ACR distances to the nearest gene. Inset, distribution of ACR genomic context.

(C) Relative DNA methylation levels 2-kb flanking ACRs.

(D) Relative enrichment of polymorphisms after normalizing by mappability for 5-kb regions flanking cell type-specific and non-specific ACR summits. Smoothed splines are shown as dark lines.

(E) Relative enrichment of significant GWAS polymorphisms relative to all polymorphisms for 5-kb regions flanking cell type-specific and non-specific ACR summits.

(F) Enrichment of signatures of selection (XP-CLR) in the top 2,000 ACRs for all cell-type clusters. The 20 most enriched cell types are highlighted on the left. AM, axillary meristem; CC, companion cell; FM, floral meristem; FP, floral primordia; IM, inflorescence meristem; QC, quiescent center; SM, spikelet meristem; SPM, spikelet pair meristem.

(G) Aggregate scATAC-seq tracks for seven floral cell types and a random assortment of 10 non-floral cell types at *ZMM29* and *ZMM18* loci. CEI, cortex/endodermis initials; GC, guard cell; GMC, guard mother cell; PP/PSEP, pre-procambial/phloem sieve element precursor.

bulk-level experiments (**Figure 3B**) (Ricci et al., 2019). Notably, 30% (22,456/73,791) of distal

ACRs overlapped with LTR transposons, including the major maize domestication locus

TEOSINTE-BRANCHED 1-enhancer (*tb1*-enhancer), and were generally devoid of DNA

methylation (**Figure 3B**, **3C**, and **S1**) (Crisp et al., 2020; Oka et al., 2020; Oka et al., 2017). These

findings are consistent with transposable elements playing a prominent role in CRE evolution of

the maize genome (Clark et al., 2006; Noshay et al., 2020; Zhao et al., 2018).

Sequence variation underlying CREs contribute to disease emergence and phenotypic

innovation over evolutionary timescales (Rebeiz and Tsiantis, 2017; Villar et al., 2015). In contrast

to broadly accessible chromatin regions, analysis of extant genetic variation in maize revealed

lower polymorphism rates within cell type-specific ACRs (**Figure 3D**). However, of the genetic

variants embedded within ACRs, those within cell type-specific ACRs were more frequently

associated with phenotypic variation (Wallace et al., 2014) (**Figure 3E**). To investigate the

contribution of domestication and selection in distinct cell-type contexts, we assessed the relative

enrichment of selection signatures from chronologically sampled elite inbred maize lines within

cell type-specific ACRs (**STAR Methods**) (Wang et al., 2020). Of the 21 cell types with significant

(FDR < 0.01) selection signature enrichment, 57% (12) correspond to staminate and pistillate cell

types, such as spikelet meristems, spikelet pair meristems, inflorescence meristems, floral

meristems, and floral primordia (**Figure 3F**). For example, a single block encompassing two

adjacent class B floral-organ morphology loci, *ZEA MAYS MADS 29* (*ZMM29*) and *ZMM18*,

exhibited inflorescence, spikelet, and floral meristem and primordia-specific ACRs at both TSSs

(**Figure 3G**). These findings indicate that modern maize breeding resulted in the selection of

alleles containing floral-specific ACRs associated with agronomically favorable inflorescence

architecture (Gage et al., 2018).

Variation in transcription factor activities underlies cell identity

Differential TF binding has been proposed as a key driver of differential gene expression

signatures underlying diverse cell identities. In line with recognized *cis*-regulatory function, ACRs

were highly enriched with putative TF binding sites relative to control (n=165,913) and flanking

regions, and were strongly depleted within ACR summits, consistent with TF-bound sequence

occluding Tn5 integration (**Figure 4A**). Using the top 2,000 differential ACRs for each cell type,

we found 84% of TF motifs (475/568) were enriched (binomial test: FDR < 0.05; **STAR Methods**)

in at least one cell type, with a median of 43 enriched TF motifs per cell type (**Figure 4B**).

Next, we hypothesized that the relative accessibilities of motif across ACRs in a single

nucleus could be used to elucidate the regulatory rules governing discrete cell states (**Figure**

S7C-**S7F**). Comparison of TF gene accessibility with the relative accessibility of their sequence-

specific binding sites revealed strikingly similar patterns across cell types (median Pearson's

correlation coefficient across cell types = 0.45), establishing synchronized chromatin accessibility

of *cis* and *trans* cell-autonomous factors as major determinants of cell identity (**Figure 4C**, **4D**,

and **S7G**). Assessment of enriched TFs and their cognate motifs identified several known cell

type-specific regulators – including *WRKY* family TFs in root epidermal progenitors and

trichoblasts (Verweij et al., 2016), *G2-like1* in parenchymal mesophyll (Chang et al., 2012), and

AGAMOUS-like and *SEPALLATA* (Gomez-Mena et al., 2005) TFs in floral primordia – as well as

previously uncharacterized TFs with new potential roles as cell-type regulators (**Figure 4C** and

4D; **Table S6** and **S7**). To determine the utility of TF motif signatures for discerning cell identity,

we trained a neural network (NN) on patterns of TF motif accessibility underlying various cell

types. The NN model achieved an overall accuracy of 0.94 and an average sensitivity and

specificity of 0.93 and 0.99, respectively, indicating that patterns of motif accessibility enable

highly predictive classifications of diverse cell states (**Figure 4E**).

Past developmental genetic studies have described a handful of mobile TFs capable of

influencing the identities of neighboring cells. As a proxy for non-cell autonomous activity, we

searched for TFs with increased motif accessibility in cell types lacking expression (and

Figure 4. Combinatorial accessibility of transcription factor motifs and genes contribute to distinct cell identities.

(A) Average motif coverage for all ACRs (n=165,913) and control regions (n=165,913). Shaded polygon, 95% confidence intervals.

(B) Enrichment of TF motifs in the top 2,000 ACRs ranked by Z-score for each cell type compared to the top 2,000 most constitutive ACRs via binomial tests. FDR was estimated by the Benjamini-Hochberg method.

(C) Comparative heatmaps for matched TF gene accessibility (bottom) and motif deviation (top) Z-scores across clusters.

(D) Gene accessibility scores for five maize transcription factors (top) and their associated motif deviations (bottom).

(E) Comparison of predicted vs. reference cell-type annotations from a neural-network multinomial classifier trained on combinatorial motif deviation scores.

(F) Co-embedded seedling nuclei gene accessibility, RNA expression, and motif accessibility for *ZmDOF36*.

accessibility) of the cognate TF. Of 279 TFs, we identified 20 with putative non-autonomous

activity, including at least four TFs, *PHLOEM EARLY DOF1 (PEAR1), TEOSINTE*

BRANCHED1/CYCLOIDEA/PROLIFERATING CELL NUCLEAR ANTIGEN FACTOR4 (*TCP4*),

TCP5, TCP14, and *ETHYLENE RESPONSE FACTOR 018* (*ERF018*), with predicted or known

cell-cell mobility (Miyashima et al., 2019; Nag et al., 2009; Savaldi-Goldstein et al., 2007;

Tatematsu et al., 2008). For example, *PEAR1* was recently described as a mobile DOF TF

expressed in the procambium functioning to promote radial growth in the vasculature of *A.*

thaliana. Consistent with predicted mobility, the maize *PEAR1* homolog, *ZmDOF36*, was largely

expressed in procambial and protophloem cells, while its target motif was enriched in procambial,

bundle sheath, phloem parenchyma, meta/protophloem, xylem, and epidermal cell types (**Figure**

4F; **Table S8**). These results indicate robust inference of CRE and TF activity at the level of single

nuclei and reveal TF dynamics central to *cis*-regulatory specification of diverse cell states.

Coordinated dynamic chromatin accessibility recapitulates *in vivo* chromatin interactions

Correlated changes in chromatin accessibility of nearby loci represent putative physical chromatin

interactions with regulatory potential (Buenrostro et al., 2015; Cusanovich et al., 2018; Gate et

al., 2018; Pliner et al., 2018; Satpathy et al., 2019). We identified 3.8 million (M) ACR-ACR

linkages (hereafter referred to as co-accessible ACRs) with significantly correlated patterns of

chromatin accessibility across cell types, capturing known gene-to-CRE physical interactions for

gene loci such as *tb1,* maize *RELATED TO AP2.7* (*ZmRAP2.7*), and *BENZOXAZINLESS 1* (*BX1*)

(empirical FDR < 0.05; **Figure 5A**, **S8**, **S9A**; **STAR Methods**) (Clark et al., 2006; Peng et al.,

2019; Ricci et al., 2019; Salvi et al., 2007; Sun et al., 2020; Zheng et al., 2015). To assess the

broad interactive potential of co-accessible ACRs *in vivo*, we compared co-accessible ACRs from

seedling cell types with maize seedling chromatin conformation capture data, recovering more

than 78% (3,313/4,265), 57% (37,712/65,691) and 44% (17,108/38,567) of chromatin loops from

Hi-C, H3K4me3-HiChIP and H3K27me3-HiChIP experiments, respectively (**Figure S9B**). Hi-

Figure 5. Co-accessible ACRs reflect *in vivo* chromatin interactions and are established by co-accessible TFs.
(A) Comparison of ear and seedling co-accessible ACRs across nine cell types at the genetically mapped *tb1-enhancer* domestication locus (highlighted region). Link height reflects co-accessibility scores between ACRs across cells in a cluster.
(B) Average normalized Hi-C signal across 4-kb windows centered on ACRs equally distributed into three groups (each group: n=54,904) based on the number and strength of participating co-accessible links.
(C) Co-accessible ACR interaction frequency across clusters. Inset: Single cell-type co-accessible links (n=1,018,417) for different genomic contexts. D, distal; P, proximal; G, genic. Example: D-P indicates co-accessible ACRs where one edge is distal (> 2-kb from any gene) and the other is proximal (< 2-kb from any gene).
(D) Standardized (row-wise) number of connections for each ACR (rows) by cell type (columns). Column color map reflect cell types from the legend in Figure S6. Gene-proximal ACRs for a subset of marker genes are indicated on the right.
(E) Right, chromatin accessibility and co-accessible ACR links surrounding the *UB2* locus associated with ear row number and tassel branch number quantitative traits. Black arrow indicates a distal ACR upstream of *UB2* present only in spikelet meristems. Co-accessible links with an edge within 2-kb of *UB2* are colored pink while remaining links are grey. Link height represents co-accessibility strength. Left, close-up of accessibility profiles of *UB2*.
(F) Distributions of average ACR-ACR links across cell types for ACRs that overlap (purple) and do not overlap (grey) phenotype-associated genetic variants from maize GWAS. The median of each distribution is shown as a white horizontal line. Violin plots present the entire range of average number of connections on a log scale. Hypothesis testing was conducted within the *R* statistical framework via Wilcoxon rank sum test.
(G) Distributions of average ACR-ACR links across cell types for ACRs with (blue) and without (grey) enhancer activity (log$_2$ RNA/input greater than 0). Hypothesis testing and distribution illustration was performed similarly as panel F.
(H) Motifs ranked by the average co-accessibility enrichment over background across all cell types.
(I) Exemplary motifs enriched in reciprocal co-accessible ACRs for *TCP*, *AP2-EREB*, and *LBD* TF families.

C/HiChIP is a direct reflection of the proportion of cells exhibiting a particular interaction, as

ubiquitously interacting loci dominate rarer cell type-specific contexts that are frequently missed

by loop-calling algorithms. To determine the relative predictability of *in vivo* interactions using co-

accessible ACRs, we estimated the interaction strength of each ACR by integrating correlative

scores across cell types (**STAR Methods**). ACRs classified by interaction strength recapitulated

expected *in vivo* chromatin interaction frequencies, where even the weakest class of co-

accessible ACRs were associated with elevated interaction frequencies relative to flanking and

randomized control regions (**Figure 5B**).

Cataloguing the usage of co-accessible ACRs across cell types identified more than 27%

(~1M in total) that were unique to a single type, 49% of which were classified as distal-genic or

distal-proximal, and an average of 11,069 distinct links per cell type (**Figure 5C**). Consistent with

regulatory models where a single gene can interact with multiple distant loci, proximal ACRs,

rather than distal or genic ACRs, were associated with the greatest number of links on average

(Wilcoxon rank sum test: P < 2.2e-308; **Figure S9C**). Highlighting long-range "hub" interactions

as key contributors towards cell identity, cell type-specific co-accessible ACRs were associated

with greater number of links per site (Wilcoxon rank sum test: P < 2.2e-308) and a greater

proportion of links involving distal ACRs (Chi-squared test: P < 2.2e-308; **Figure S9C** and **S9D**).

Furthermore, the interactive capacity of any given ACR strongly depended on the cell-type context

(**Figure 5D**). For example, *UNBRANCHED 2* (*UB2*) – a major ear row number and tassel branch

number quantitative trait locus (Chuck et al., 2014) – demonstrated preferential accessibility in

spikelet meristems that coincided with the greatest number of *UB2* proximal to distal ACR

interactions, including a cell type-specific ACR located upstream approximately 150-kb (**Figure**

5E). We posited that ACRs with expanded interactive capacity resemble enhancers with the

potential to influence organismal phenotypes. Indeed, ACRs with enhancer activity and co-

localization with phenotype-associated genetic variants from GWAS were associated with a

significantly greater number of ACR-ACR connections (Wilcoxon rank sum: P < 2.2e-308; **Figure**

5F and **5G**). These results highlight the occurrence of diverse cell type-specific regulatory

configurations among distal enhancer ACRs and their target genes and implicate genetic variants

perturbing highly interactive distal enhancers as major contributors towards phenotypic variation.

The structural protein, CCCTC-binding factor (CTCF) plays an important role in metazoan

genome organization and is notably absent in plant lineages (Heger et al., 2012). In search of an

orthogonal factor in maize, we hypothesized that higher-order chromatin structure captured by

co-accessible ACRs may be driven by TFs recognizing similar sequence motifs embedded within

interacting accessible regions. Comparison of motif occurrences between co-accessible link

edges indicated that ACRs in co-accessible links are more similar to one another than randomly

linked ACRs (empirical: P < 1e-4; **Figure S9E**). Furthermore, we identified several cases where

co-accessible ACR edges were reciprocally enriched for the same TF motif in both cell type-

specific and non-specific co-accessible links (FDR < 0.05; **Figure S9F**; **STAR Methods**). Ranking

motifs by the average enriched across cell types, we identified *TCP*, *APETALA2*/*ETHYLENE-*

RESPONSIVE ELEMENT BINDING PROTEINS (*AP2-EREBP*) and *LATERAL ORGAN*

BOUNDARIES DOMAIN (*LBD*) motifs that were not only broadly associated with co-accessible

edges, but also exhibited strikingly similar GC-rich palindromic binding sites (FDR < 0.05; **Figure**

5H, **5I** and **S9F**). A role in chromatin organization is supported by previous research

demonstrating *TCP* motif overrepresentation in topologically associated domain-like (TAD)

boundaries in *Oryza sativa* and *Marchantia polymorpha,* and the distal edges of chromatin loops

in *Z. mays* (Karaaslan et al., 2020; Liu et al., 2017; Peng et al., 2019; Sun et al., 2020). Consistent

with these past studies, our results implicate independently evolved TF families with CTCF-like

function capable of organizing higher-order chromatin architecture through DNA-protein

interactions.

Dynamic chromatin accessibility specifies cell developmental trajectories

The apical domains of maize enclose a pool undifferentiated meristematic stem cells that give

continuous rise to all differentiated cell types (Somssich et al., 2016; Takacs et al., 2012). To

define a *cis*-regulatory catalog of temporal cell fate progressions, we ordered nuclei along pseudo-

temporal trajectories for 18 developmental continuums, reflecting meristematic to differentiated

cell states; identifying ACRs, TF loci, and TF motifs with significant variation across pseudotime

(**Figure 6A** and **S10**; **Table S9-S12**; **STAR Methods**). To showcase the power of trajectory

construction to characterize a relatively understudied process, we focused our analysis on root

phloem companion cell (PCC) development (**Figure 6B**). We identified 8,004 ACRs, 440 TF

motifs, 7,955 genes, and 402 TF loci that were differentially accessible across the PCC

pseudotime trajectory (**Figure 6C**; **STAR Methods**). Several known meristem and phloem

developmental genes including *AT-RICH INTERACTIVE DOMAIN-CONTAINING 8* (*ARID8*)

(Jiang et al., 2010), *SUPPRESSOR OF MAX2 1-LIKE3* (*ZmSMXL3*) (Wallner et al., 2017), *and*

SUCROSE TRANSPORTER 1 (*ZmSUT1*) (Baker et al., 2016), were identified among the top

differentially accessible genes throughout PCC development (**Figure 6D**).

Past studies of root cell fate decisions have focused on the role of cell cycle in establishing

patterns of asymmetric cell division, with quiescent center/meristematic cells dividing much slower

than cells in the rapidly dividing transition and elongation zones (Ten Hove and Heidstra, 2008).

To investigate the contribution of cell-cycling to PCC development, we annotated nuclei using *a*

priori compiled list of known cell-cycle marker genes (**STAR Methods**) (Nelms and Walbot, 2019).

As consequence of slower DNA replication and consistent with previous reports, the majority of

QC and meristem/initial-like nuclei were in S-phase, while differentiated companion cells largely

presented as G1 (**Figure 6E**). Ordering nuclei by PCC pseudotime indicated sequential

progression of cycle stages within each cell type, revealing the cell cycle context preceding cell

fate transitions along the PCC trajectory (**Figure 6F**). Furthermore, evaluation of global

accessibility across pseudotime illustrated steady decrease in chromatin accessibility throughout

PCC development (**Figure 6G**). Thus, cell-cycling and cell fate transitions in the context of PCC

Figure 6. Chromatin accessibility is dynamic across pseudotime

(A) Overview of pseudotime trajectory analysis. Inflorescence development was performed for both pistillate and staminate inflorescence.

(B) UMAP embedding of companion cell developmental trajectory depicting cell types (left) and pseudotime progression (right).

(C) Relative motif deviations for 440 TF motifs (left, rows), 402 TF gene accessibility scores (middle, rows), and relative accessibility of 8,094 ACRs (right, rows) associated with pseudotime (columns). Four motifs enriched along the trajectory gradient are shown on the left. ACRs, accessible chromatin regions; TF, transcription factor.

(D) Genome browser screenshots of cell type-specific chromatin accessibility profiles along the developmental trajectory for quiescent center (QC), pre-procambial/phloem sieve element precursor (PP/PSEP), and phloem companion cell (PCC) at associated marker gene loci.

(E) Proportion of cells at various stages of the cell-cycle in QC, PP/PSEP, and PCC annotated clusters.

(F) Top: Cell state ordered by pseudotime. Middle: Proportion of nuclei with the corresponding cell-type annotation ordered by pseudotime. Bottom: proportion of nuclei with various cell-cycle stage annotations ordered by pseudotime. Nuclei were binned into 250 blocks.

(G) Left: Average fraction of ACRs that are accessible across pseudotime. The grey polygon indicates standard deviation. Red windows indicate cell state transitions. Right: heatmap of relative accessibility (relative to the row maximum) for each ACR (rows) across pseudotime (columns). Nuclei were binned into 250 blocks ordered on pseudotime.

development accompany global decreases in chromatin accessibility, a consequence we posit is

associated with acquisition of more specialized functions in PCCs relative to their meristematic

progenitors.

Evolutionary innovation in root development

Despite nearly 150M years of divergence, monocot and eudicot angiosperm species exhibit

remarkable phenotypic similarity with core organs such as seeds, roots, and shoots functionally

maintained. To explore the degree of regulatory conservation in angiosperm root development,

we profiled chromatin accessibility in 4,655 nuclei from 7-day old embryonic root tissues in the

eudicot model species *A. thaliana*, integrated with previously generated *A. thaliana* root scRNA-

seq data (n=12,606), and constructed eight *cis*-regulatory pseudotime trajectories encompassing

vascular, dermal and ground development (**Figure 7A**-**7C**, **S11** and **S12A**; **Table S13-S17**).

Maintaining focus on PCC development, we first validated the utility of the integrated data

sets by visualizing gene expression and accessibility of known marker genes representative of

QC (*WUSCHEL-RELATED HOMEOBOX 5, WOX5*), procambial (*WUSCHEL-RELATED*

HOMEOBOX 4, WOX4) and PCC (*SUCROSE TRANSPORTER 2, SUC2*) cell types (**Figure 7D**

and **7E**). Next, we aligned *Z. mays* and *A. thaliana* PCC trajectories using a time-warping

algorithm to enable direct comparison of gene accessibility dynamics in a common space.

Consistent with recent comparative analysis of vascular development in *O. sativa*, *A. thaliana,*

and *Solanum lycopersicum* (Kajala et al., 2020), only 206 out of 10,976 putative orthologs were

significantly associated (FDR < 0.01) with PCC pseudotime in both species, indicating that the

majority PCC trajectories-associated genes are unique to each lineage (97% *Z. mays*, 83% *A.*

thaliana). However, of the 206 PCC-associated orthologs, ~50% (102/206) exhibited similar

patterns of gene accessibility across pseudotime (**Figure 7F**, **7G**, **S12B**). Several putative

orthologs with matching gene accessibility patterns have been previously associated with PCC

Figure 7. Dynamic and conserved *cis*-regulation in *A. thaliana* and *Z. mays* phloem companion cell development.
(A) UMAP embedding based on whole-genome chromatin accessibility profiles of 4,655 *A. thaliana* root nuclei.
(B) UMAP embedding of companion cell developmental trajectories in *A. thaliana* depicting cell types (left) and pseudotime progression (right). Motifs, TFs, and ACRs with significant association with pseudotime are shown as heatmaps below.
(C) UMAP embedding of integrated scRNA-seq and scATAC-seq cell and nuclei profiles derived from *A. thaliana* roots.
(D) Marker gene expression levels for individual scRNA-seq cells for quiescent center (QC; *WOX5*),
pre-procambial/phloem SE precursors (PP/PSEP; *WOX4*), and phloem companion cells (PCC; *SUC2*).
(E) Pseudobulk chromatin accessibility pile-ups for clusters labeled as QC, PP/PSEP, and PCC across three marker genes associated with each cell type, respectively.
(F) Per-gene pseudotime shift scores from alignments between *Z. mays* and *A. thaliana* companion cell development progressions, clustered by k-means into three groups.
(G) Distribution of gene-gene distances from the alignments, split by k-mean groups.
(H) Exemplary one-to-one homologs between *A. thaliana* and *Z. mays* for the three groups split by pseudotime shifts.
Acronyms: LRC, lateral root cap; LRP, lateral root primordia. SE, sieve elements.

development, such as *SCARECROW-LIKE8* (*SCL8*), which displays an increasing expression

gradient in nascent to mature *A. thaliana* PCCs in a previously generated *A. thaliana* root cell-

type gene expression atlas (**Figure 7H**) (Brady et al., 2007). The remaining homologs (n=104)

exhibited differential accessibility patterns clustered into two groups that reflect changes in the

timing of gene accessibility across the PCC trajectory, underscoring putatively functional novelty

in PCC development between *Z. mays* and *A. thaliana*.

470   To understand the extent of innovation in *cis* regulation along the pseudotime continuum,

we aligned *Z. mays* and *A. thaliana* TF motif accessibility profiles associated with PCC

progression (**Figure S12C**, **S12D**). Of the 440 motifs, 142 demonstrated highly conserved *cis-*

regulatory dynamics between species (**Figure S12E**-**S12G**). Indeed, TFs recognizing the top four

motifs ranked by normalized distances (**STAR Methods**) included *HOMEOBOX25* (*HB25*),

HOMEOBOX18 (*HB18*), *NAC DOMAIN CONTAINING PROTEIN 55* (*NAC055*)*, and *NAC*

DOMAIN CONTAINING PROTEIN 83 (*NAC083*) that have been previously implicated in

regulation of hormonal responses and vascular development (Jiang et al., 2009; Yamaguchi et

al., 2010; You et al., 2019). Gene expression profiles of these TFs from published root cell-type

resolved data in *A. thaliana* was largely restricted to maturing procambial and companion cells,

consistent with motif accessibility dynamics in both *Z. mays* and *A. thaliana* ontogenies (**Figure**

S12H) (Brady et al., 2007)*. These finding signify a high degree of conservation in the *cis-*

regulatory specification of PCC development between *Z. mays* and *A. thaliana* despite an

analogous lack of concordance in accessibility dynamics among orthologous genes.

DISCUSSION

Here, we describe 92 distinct cell-states from a suite of *Z. mays* organs, many of which previously lacked prior genome-wide characterization. Although we provide evidence supported by RNA *in situ* hybridization and snRNA-seq for the use of chromatin accessibility as a robust proxy of gene expression, the cell-type annotations should be considered preliminary. We anticipate that as single-cell methods become more widely adopted, including same-cell multi-modal experiments, these cell-type classifications, as well as those that were left unannotated, will become refined. An important consideration is that the sparse and binary nature of current scATAC-seq protocols require cell/nuclei aggregation by cell type for downstream analyses. Thus, additional gains in experimental procedures, particularly in reads per cell together with comprehensive transcriptome profiling within the same cell, will be necessary to fully investigate heterogeneity of cells within the same type. We anticipate that the application of same-cell multimodal techniques will open the door to better establish the molecular relationships among chromatin accessibility, gene expression, and cellular heterogeneity. Advancements in computational tools for comparing cell-type atlases and for data integration will play a key role in enabling higher resolution analyses than is currently possible.

Notwithstanding the technical challenges of single-cell experiments, our results represent a landmark advance for appreciating variation in cell-type functions established by diverse *cis*-regulatory grammar. We defined the TFs, CREs, and other loci that discretize cell-type identities and the sequential trajectories for an array of developmental ontogenies. Evaluation of motif accessibility variation alone was sufficient to predict cell identity with a high degree of accuracy, sensitivity and specificity. Querying patterns of motif accessibility relative to TF gene expression highlighted transcriptional regulators with putative non-cell autonomous activity, a particularly exciting result as identification of candidate non-cell autonomous factors previously relied exclusively on transgenic approaches to illuminate both transcript and protein localizations. Towards uncovering major regulators of global chromatin structure in species that lack CTCF,

516   analysis of co-accessible ACRs successfully linked distal ACRs to their target genes and provided

517   CTCF-like candidate TFs putatively orchestrating higher-order chromatin interactions that have

518   been posited by orthogonal approaches. Further dissection of candidate regulatory genes and

519   regions promises to be a fruitful endeavor for precise engineering of spatiotemporal patterns of

520   gene expressions.

521   With an evolutionary perspective, we reveal that floral cell type-specific ACRs have been

the historical targets of modern agronomic selection in maize. While subject to considerable

sequence constraint, both by artificial and natural selection, genetic variation that does exist within

cell type-specific ACRs is highly enriched with significant phenotypic variation. These findings

point to abundant genetic variation capable of large phenotypic effects present within extant maize

germplasm and present an ideal launchpad towards allele-mining for crop improvement. To

understand the extent of cis-regulatory evolution in two highly diverged species, we constructed

cell-type resolved chromatin accessibility profiles in the eudicot species *A. thaliana*. To our

surprise, comparison of established orthologs indicated that a majority of genes involved in cell-

type development were unique to each lineage. This finding contrasted with the observation of

greater conservation of *cis*-regulatory elements involved in PCC development, as a greater

proportion of TF motifs exhibited consistent spatiotemporal progressions among the two species.

Viewed collectively, the maize *cis*-regulatory atlas presents an ideal framework for understanding

the basis of cell heterogeneity, cis-regulatory control of gene transcription, and the foundation for

future crop improvement efforts through targeted genome editing, synthetic biology approaches,

and traditional allele-mining.

AUTHOR CONTRIBUTIONS

APM and RJS designed the research. APM and ZC performed the experiments. APM, ZC, AG and RSJ analyzed the data. APM and RJS wrote the manuscript.

DECLARATION OF INTERESTS

RJS is a co-founder of REquest Genomics, LLC, a company that provides epigenomic services. APM, ZC and AG declare no competing interests.

ACKNOWLEDGEMENTS

We would like to acknowledge Dr. Zefu Lu for training in nuclei extraction, Dr. William Ricci for providing inflorescence samples, Julie Nelson from the University of Georgia flow cytometry core for helping sort nuclei, Noah Workman and Julia Portocarrero from the Georgia Genomics and Bioinformatics Core for library preparation, the Georgia Advanced Computing Resource Center for providing access to computing resources, and Tyler Earp for assistance with data management. This study was funded by support from the National Science Foundation (IOS-1856627 and IOS-1802848), the Pew Charitable Trusts and the UGA Office of Research to RJS, the National Science Foundation (IOS-1916804) to AG, and an NSF Postdoctoral Fellowship in Biology (DBI-1905869) to APM. AG and RJS are also supported by an NSF Collaborative award to support this research (IOS-2026554 and IOS-2026561). RJS is a Pew Scholar in the Biomedical Sciences, supported by The Pew Charitable Trusts.

CORRESPONDENCE

Correspondence and requests for materials should be addressed to Robert J. Schmitz (schmitz@uga.edu) and Alexandre P. Marand (marand@uga.edu).

Figure S1: Evaluation and quality control of maize scATAC-seq
(**A**) Genome browser screenshot of chromatin accessibility from bulk and aggregated single-cell ATAC-seq experiments. Chromatin accessibility profiles depict the *tb1* locus and the *tb1* enhancer located approximately 67kb upstream.
(**B**) Binary accessibility scores from a random selection of 1,000 individual nuclei from each organ.
(**C**) Spearman's rho matrix comparing bulk ATAC-seq and aggregate scATAC-seq samples across various organs. Sample codes are short hand for assay type, sample, and replicate. For example, s-R1 denotes single cell assay for seminal root replicate 1. The term b-L2 denotes a bulk-ATAC assay for seedling replicate 2. Codes are as follows: b, bulk; s, single cell; R, seminal root; C, crown root; E, ear; T, tassel; A, axillary bud; L, seedling. Numbers represent replicate.
(**D**) Average TSS enrichment (normalized read depth adjusted by the two 10 bp windows 1kb away from TSSs) across all 56,575 cells. Grey polygon denotes the standard deviation.
(**E**) Fragment length distributions across 56,575 cells. The solid line and grey polygon represent the average and standard deviation, respectively.
(**F**) Genotype-mixing experimental schematic.
(**G**) Scatterplot of per cell B73 and Mo17 SNP counts from a mixed-genotype experiment (V1 seedlings) colored by genotype classification.
(**H**) Posterior probabilities of individual barcodes (rows) highlighting the occurence of cells with B73, Mo17, and mixed (doublet) genotype identities.
(**I**) Genome browser screenshot of traditional bulk ATAC-seq from 7 day old seedling (row 1), single-cell ATAC-seq from B73 seven day old seedlings (row 2), pooled B73 and Mo17 nuclei (library ID: Seedling 2) single cell ATAC-seq from 7 day old seedlings (row 3), and the genotype-sorted B73 (row 4) and Mo17 (row 5) alignments after sorting barcodes by genotype calls from the B73-Mo17 scATAC-seq 7 day old seedling sample (row 3).

Figure S2. Cell calling and barcode quality control

(**A**) Comparison of normalized (0-1) read depths at the union of all peaks across bulk and single-cell samples (n=265,992) between replicated libraries, and between bulk and single-cell ATAC-seq assays.

(**B**) Enrichment plots centered on 2-kb windows surrounding TSSs for barcodes in each tissue. Grey polygons indicate the standard deviation across cells within the noted tissue.

(**C**) Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites mapping to within 2-kb of transcription start sites (TSSs). Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes.

(**D**) Fragment length distributions for each library. Solid lines indicate the average distribution across cells within the sample. Grey polygons represent the standard deviation across cells in the library.

(**E**) Knee plots illustrating $\log_{10}$ transformed cellular read depths of $\log_{10}$ ranked barcodes across libraries.

(**F**) Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites derived from organeller sequences (chloroplast and mitochondrial) relative to the total number of unique Tn5 integration sites associated with cognate barcode. Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes.

Figure S3. *In silico* sorting via Latent Semantic Indexing

Standardized LSI accessibility scores (2nd - 11th dimensions) capped at ±1.5 for:

(**A**) Axillary buds.

(**B**) Crown roots.

(**C**) Embryonic roots.

(**D**) Seedlings.

(**E**) Ear (pistillate inflorescence).

(**F**) Tassel (staminate inflorescence).

Figure S4. Clustering metrics and comparison of bulk gene accessibility and expression

(**A**) Parameter regularization of model coefficients (y-axes) with respect to ACR usage (x-axes; proportion of nuclei with at least one Tn5 integration site in an ACR).

(**B**) Proportion of variance captured by the first 26 PCs. Inset: Spearman's correlation of principal components with cell read depth (log_{10}-transformed).

(**C**) Number of accessible sites per cell (log_{10}).

(**D**) Proportion of Tn5 integrations within 2kb of gene TSSs per cell.

(**E**) Co-localization of nuclei barcodes from different biological replicates for three organs.

(**F**) Comparison of bulk RNA-seq expression levels (y-axis, log_2 CPM) versus aggregate scATAC-seq gene accessibility scores (x-axis, log_2 CPM) within an organ.

Figure S5. Cell-type annotation and GO enrichment

(**A**) UMAP embeddings of nuclei barcodes colored by low (grey) to high (dark purple) gene activity values of cell-type specific marker genes.

(**B**) RNA *in situ* hybridization showing expression of *LOX10* in glume primordia and *GRFTF36* in IM and SMs of staminate inflorescence; *LRR445* in the IM periphery and SPMs and *MYB89* in the IM and suppressed bract primordia of pistillate inflorescence; Zm00001d038453 in ground tissue of SAM and leaf primordia sections. Gene accessibility scores and predicted cell-types are shown on the right. *i*, tassel primordia. *ii,* ear primordia. *iii*, SAM/leaf. Black triangles point to the glume primordia. Red triangles point to suppressed bract primordia. Size bars illustrate 100-um. AM, axillary meristem; BS, bundle sheath; GC, guard cell; GM, ground meristem; GMC, guard mother cell; GP, glume primordia; IM, inflorescence meristem; L1, layer 1; LFM, lower floral meristem; SAM, shoot apical meristem; SBP, suppressed bract primordia; SM, spikelet meristem; SPM; spikelet pair meristem; Stomatal PC, stomatal precursor; UKN, unknown; VEMI, vascular/epidermal meristematic identity; VP, vascular parenchyma; XP, xylem parenchyma.

(**C**) Proportion of cells within subcluster (column) derived from one of six organs (rows).

Figure S6. Chromatin accessibility variation across plant cell-types
(**A**) Row gene accessibility Z-scores across cell-types. Cell-type and subcluster labels are located adjacent to the heatmap in identical order as the matrix. The first number after the cell-type label indicates the major cluster number, while the second number represents the subcluster ID. Legend colors: top left triangle, subcluster (corresponds to the sub-cluster annotation on heatmap). bottom right triangle, major cluster (corresponds to the major cluster annotation on heatmap).
(**B**) Row ACR chromatin accessibility Z-scores across cell-types.
(**C**) Proportion of cells derived from one of six organs within each cell-type/subcluster.
(**D**) Distribution of GO term enrichment across clusters, where the x-axis indicates the number of clusters in which a GO term is significantly enriched.

Figure S7. Transcription factor motif variation underly dynamic accessible regions

(**A**) Comparison of cell-type specific ACR counts (by cell-type) and the number of unique Tn5 integration sites.

(**B**) Proportion of distal, genic and proximal ACRs per cell-type.

(**C**) Ranked TF motif variability across cells, colored by TF family.

(**D**) Mean TF family motif enrichment (average deviation scores per cell-type per TF family) across cell-types scaled to +/- 1.

(**E**) Spearman's correlation coefficient (rho) between TF motifs (comparison of motif deviations across all nuclei). Row and column colors represent TF motif families.

(**F**) TF motif deviations for 440 TF motifs (columns) per nucleus (rows). The TF family for each corresponding motif is denoted by column header colors. Cluster identification (sub-cluster/major cluster) are illustrated as row header colors on the right side of the heatmap.

(**G**) Relative gene accessibility scores for 2,423 maize transcription factors (columns) for 50,640 nuclei (rows). Nuclei are sorted by sub-cluster cell-type (row color labels). A cell-type annotation color legend applicable to panels **B**, **F**, and **G** where left triangle reflects sub-cluster, right triangle indicates the major cluster.

Figure S8: Capture of known long-range chromatin loops in maize

(**A**) Co-accessible ACRs at the *ZmRAP.2* locus in maize, with root-specific expression patterns, across eight root-derived (left) and eight above-ground (right) cell-types. The height of the loops reflect the strength of the co-accessibility. Pseudobulk chromatin accessibility tracks are shown under co-accessible ACR linkages for each cell-type.

(**B**) Co-accessible ACRs at the *BX1* locus in maize, predominantly expressed in seedling tissue, across eight seedling-derived (left) and eight non-seedling derived cell-types. The height of the loops reflect the strength of the co-accessibility. Pseudobulk chromatin accessibility tracks are shown under co-accessible ACR linkages for each cell-type.

Figure S9. Co-accessible ACRs reflect in vivo chromatin interactions driven by coordinated TF activity

(**A**) Proportions of co-accessible ACR types, illustrated by toy examples.

(**B**) Proportion leaf Hi-C, H3K4me3-HiChIP and H3K27me3-HiChIP chromatin loops that overlap co-accessible ACRs from leaf cell-types (clusters with greater than 50% of cell derived from seedlings).

(**C**) Top, log10 number of connections per ACR per cell-type from all co-accessible ACRs split by genomic context: distal, proximal, and genic. Bottom, $\log_{10}$ number of connections per ACR per cell-type from cell-type specific (purple) and non-specific (grey) co-accessible ACRs, split by genomic context.

(**D**) Proportion of co-accessible classifications by cell-type for all co-accessible ACRs (top) and cell-type specific co-accessible ACRs (bottom).

(**E**) Jaccard similarity of motif composition between co-accessible ACR edges by cell-type (colored diamonds) relative to the same number of random ACR-ACR links, permuted 1,000 times for each cell-type (grey boxplots). Box plots represent the interquartile range, grey lines indicate the permuted range.

(**F**) Heatmaps of observed proportion of co-accessible ACRs with the same motif embedded within link edges subtracted and divided by the expected proportion estimated by 1,000 permutations using random ACR-ACR links with the same number of co-accessible ACRs.

Figure S10. Pseudotime trajectory construction

(**A**) Overview of pseudotime developmental trajectory analysis from four organs: root, seedling, tassel (staminate inflorescence), and ear (pistillate inflorescence).

(**B**) Endodermis development in crown roots.

(**C**) Cortex development in crown roots.

(**D**) Pericycle development in embryonic roots.

(**E**) Atrichoblast development in crown roots.

(**F**) Trichoblast development in crown roots.

(**G**) Trichoblast development in embryonic roots.

(**H**) Lateral root cap (LRC) development in crown roots.

(**I**) Lateral root cap (LRC) development in embryonic roots.

(**J**) Phloem sieve element (SE) development in crown roots.

(**K**) Companion cell development in crown roots.

(**L**) Phloem sieve element (SE) development in embryonic roots.

(**M**) Xylem development in crown roots.

(**N**) Xylem development in embryonic roots.

(**O**) Procambial development in crown roots.

(**P**) Guard cell development in seedling.

(**Q**) Subsidiary cell development in seedlings.

(**R**) Floral primordia development in staminate inflorescence (tassel).

(**S**) Floral primordia development in pistillate inflorescence (ear).

Figure S11. *Arabidopsis thaliana* root cell-type atlas

(**A**) Knee plots for Arabidopsis thaliana root samples illustrating $\log_{10}$ transformed cellular read depths of $\log_{10}$ ranked barcodes across two biological replicates.

(**B**) Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites derived from organellar sequences (chloroplast and mitochondrial) relative to the total number of unique Tn5 integration sites associated with each barcode from the two biological replicates. Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes.

(**C**) Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites mapping to within 2-kb of transcription start sites (TSSs). Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes.

(**D**) Average TSS enrichment (normalized read depth adjusted by the two 10 bp windows 1-kb away from TSSs) across 5,001 Arabidopsis thaliana root barcodes (rows).

(**E**) UMAP (Uniform manifold approximation projection) embeddings of Arabidopsis thaliana root barcodes colored by biological replicate;

(**F**) total number of accessible chromatin regions (ACRs);

(**G**) the proportion of Tn5 integration sites within 1-kb of TSSs.

(**H**) Relative gene accessibility for 27 known cell-type/domain restricted marker genes used to inform cell-type annotation of Arabidopsis thaliana root clusters.

(**I**) Relative motif deviations for transcription factors with known cell-type specificities.

Figure S12. Dynamic and conserved chromatin accessibility across pseudotime between *Arabidopsis thaliana* and *Zea mays*

(**A**) Pseudotime trajectories for Atrichoblast, Trichoblast, Lateral root cap (LRC), Cortex, Endodermis, Lateral root primordia (LRP), Companion cells (CC), and Xylem development.

(**B**) Averaged alignments of conserved, shift early *Z. mays*, and shift early *A. thaliana* putative orthologs.

(**C**) Pseudotime shifts of TF motifs between *A. thaliana* and *Z. mays*, clustered into k-means and conserved groups.

(**D**) Distributions of motif-motif normalized distances between *Z. mays* and *A. thaliana* for the three groups.

(**E**) Conserved motifs (n=142) ordered by pseudotime. Heatmaps for *A. thaliana* and *Z. mays* have identical row orders.

(**F**) Averaged alignments of conserved, shift early *Z. mays*, and shift early *A. thaliana* groups based on motif-motif global alignments from the dynamic time-warping algorithm.

(**G**) Examples of conserved, shift early *Z. mays*, and shift early *A. thaliana* motifs from both species.

(**H**) Gene expression Z-scores across *A. thaliana* FAC sorted root cell-types for the TFs recognizing the top four conserved motifs.

 MAIN FIGURE LEGENDS

Figure 1. Atlas-scale cell type profiling from single nuclei chromatin accessibility in *Zea*

mays

(A) Overview of experimental samples, with an example of the cell type diversity present in

seedlings.

(B) Nuclei similarity clustering as a UMAP embedding derived from the denoised quasibinomial

Pearson's residuals across all ACRs for each nucleus. UMAP embedding of nuclei colored by

organ identity.

(C) UMAP embedding of nuclei colored major cluster identity.

(D) UMAP embedding of sub-cluster assignments following a second round of clustering within

each major cluster. Sub-cluster color reflects the organ with the greatest proportion of nuclei in

the cluster. See panel (B) for color code.

(E) Cell type-specific enrichment of gene accessibility for a subset of marker genes associated

with six different cell types.

(F) Sub-cluster-specific chromatin accessibility profiles surrounding known marker genes for floral

primordia, xylem precursors, and L1 epidermal cells. Bold, circled numbers indicate the cognate

major cluster shown in panel c. Sub-cluster numeric identifications are present on the sides of the

coverage plots.

(G) Top, gene accessibility for *ZmGRFTF36*, an inflorescence and spikelet meristem enriched

transcription factor with no previously known cell type-specificity. Bottom, RNA *in situ*

hybridization of *ZmGRFTF36* in maize B73 staminate (tassel) primordia. FP, floral primordia; GP,

glume primordia; IM, inflorescence meristem; LFM, lower floral meristem; SM, spikelet meristem;

SPM, spikelet pair meristem; UKN, unknown.

Figure 2. Concordance between chromatin accessibility and gene expression at single-

nuclei resolution

(A) Illustration of iNMF integration of scATAC-seq and snRNA-seq seedling data sets.

(B) UMAP co-embedding of seedling nuclei from scATAC-seq (n=11,882; purple) and snRNA-

seq (n=15,515; blue).

(C) Louvain clustering and cell-type annotations for co-embedded seedling nuclei.

(D) Comparison of within-cluster averaged gene accessibility (left) and gene expression (right)

between cell types/clusters. Column color legend corresponds to the cell-type colors specified in

panel C.

(E) UMAP embeddings displaying per nucleus gene accessibility (top, n=11,882) and gene

expression (bottom, n=15,515) values for five cell type-specific marker genes.

(F) Left: Aggregate scATAC-seq tracks across clusters at the *CAH1* locus. Right: Average

expression of *CAH1*.

(G) Spearman correlations between clusters based on nuclear transcription (snRNA) and

chromatin accessibility (scATAC). (H) Density scatter plot comparing gene accessibility (x-axis)

and expression (y-axis) for each cluster and gene.

(I) Expression (left), chromatin accessibility (middle), and H3K27me3 ChIP-seq meta-profiles

(relative reads per million, RPM) of accessible/expressed genes (turquoise; n=19,402),

accessible/non-expressed genes (pink; n=6,063) and non-accessible/non-expressed genes

(grey, n=4,315).

(J) Top two *de novo* motifs enriched in ACRs within 1-kb of accessible/non-expressed genes

(pink, panel H and I).

Figure 3. Characterization of accessible chromatin regions

(A) Distribution of enhancer activity (maximum log2[RNA/input]) for control regions (n=165,913),

non-specific (n=113,393) and cluster-specific ACRs (n=52,520). The dash red line indicates the

overall mean. Orange lines reflect differences between the group median and overall mean.

(B) Distribution of ACR distances to the nearest gene. Inset, distribution of ACR genomic context.

(C) Relative DNA methylation levels 2-kb flanking ACRs.

(D) Relative enrichment of polymorphisms after normalizing by mappability for 5-kb regions

flanking cell type-specific and non-specific ACR summits. Smoothed splines are shown as dark

lines.

(E) Relative enrichment of significant GWAS polymorphisms relative to all polymorphisms for 5-

kb regions flanking cell type-specific and non-specific ACR summits.

(F) Enrichment of signatures of selection (XP-CLR) in the top 2,000 ACRs for all cell-type clusters.

The 20 most enriched cell types are highlighted on the left. AM, axillary meristem; CC, companion

cell; FM, floral meristem; FP, floral primordia; IM, inflorescence meristem; QC, quiescent center;

SM, spikelet meristem; SPM, spikelet pair meristem.

(G) Aggregate scATAC-seq tracks for seven floral cell types and a random assortment of 10 non-

floral cell types at *ZMM29* and *ZMM18* loci. CEI, cortex/endodermis initials; GC, guard cell; GMC,

guard mother cell; PP/PSEP, pre-procambial/phloem sieve element precursor.

Figure 4. Combinatorial accessibility of transcription factor motifs and genes contribute to

distinct cell identities.

(A) Average motif coverage for all ACRs (n=165,913) and control regions (n=165,913). Shaded

polygon, 95% confidence intervals.

(B) Enrichment of TF motifs in the top 2,000 ACRs ranked by Z-score for each cell type compared

to the top 2,000 most constitutive ACRs via binomial tests. FDR was estimated by the Benjamini-

Hochberg method.

(C) Comparative heatmaps for matched TF gene accessibility (bottom) and motif deviation (top)

Z-scores across clusters.

(D) Gene accessibility scores for five maize transcription factors (top) and their associated motif

deviations (bottom).

(E) Comparison of predicted vs. reference cell-type annotations from a neural-network

multinomial classifier trained on combinatorial motif deviation scores.

(F) Co-embedded seedling nuclei gene accessibility, RNA expression, and motif accessibility for

ZmDOF36.

Figure 5. Co-accessible ACRs reflect *in vivo* chromatin interactions and are established by

co-accessible TFs.

(A) Comparison of ear and seedling co-accessible ACRs across nine cell types at the genetically

mapped *tb1-enhancer* domestication locus (highlighted region). Link height reflects co-

accessibility scores between ACRs across cells in a cluster.

(B) Average normalized Hi-C signal across 4-kb windows centered on ACRs equally distributed

into three groups (each group: n=54,904) based on the number and strength of participating co-

accessible links.

(C) Co-accessible ACR interaction frequency across clusters. Inset: Single cell-type co-accessible

links (n=1,018,417) for different genomic contexts. D, distal; P, proximal; G, genic. Example: D-P

indicates co-accessible ACRs where one edge is distal (> 2-kb from any gene) and the other is

proximal (< 2-kb from any gene).

(D) Standardized (row-wise) number of connections for each ACR (rows) by cell type (columns).

Column color map reflect cell types from the legend in Figure S6. Gene-proximal ACRs for a

subset of marker genes are indicated on the right.

(E) Right, chromatin accessibility and co-accessible ACR links surrounding the *UB2* locus

associated with ear row number and tassel branch number quantitative traits. Black arrow

indicates a distal ACR upstream of *UB2* present only in spikelet meristems. Co-accessible links

with an edge within 2-kb of *UB2* are colored pink while remaining links are grey. Link height

represents co-accessibility strength. Left, close-up of accessibility profiles of *UB2*.

(F) Distributions of average ACR-ACR links across cell types for ACRs that overlap (purple) and

do not overlap (grey) phenotype-associated genetic variants from maize GWAS. The median of

each distribution is shown as a white horizontal line. Violin plots present the entire range of

average number of connections on a log scale. Hypothesis testing was conducted within the *R*

statistical framework via Wilcoxon rank sum test.

(G) Distributions of average ACR-ACR links across cell types for ACRs with (blue) and without

(grey) enhancer activity ($\log_2$ RNA/input greater than 0). Hypothesis testing and distribution

illustration was performed similarly as panel F.

(H) Motifs ranked by the average co-accessibility enrichment over background across all cell

types.

(I) Exemplary motifs enriched in reciprocal co-accessible ACRs for *TCP*, *AP2-EREB*, and *LBD* TF

families.

Figure 6. Chromatin accessibility is dynamic across pseudotime

(A) Overview of pseudotime trajectory analysis. Inflorescence development was performed for

both pistillate and staminate inflorescence.

(B) UMAP embedding of companion cell developmental trajectory depicting cell types (left) and

pseudotime progression (right).

(C) Relative motif deviations for 440 TF motifs (left, rows), 402 TF gene accessibility scores

(middle, rows), and relative accessibility of 8,094 ACRs (right, rows) associated with pseudotime

(columns). Four motifs enriched along the trajectory gradient are shown on the left. ACRs,

accessible chromatin regions; TF, transcription factor.

(D) Genome browser screenshots of cell type-specific chromatin accessibility profiles along the

developmental trajectory for quiescent center (QC), pre-procambial/phloem sieve element

precursor (PP/PSEP), and phloem companion cell (PCC) at associated marker gene loci.

(E) Proportion of cells at various stages of the cell-cycle in QC, PP/PSEP, and PCC annotated

clusters.

(F) Top: Cell state ordered by pseudotime. Middle: Proportion of nuclei with the corresponding

cell-type annotation ordered by pseudotime. Bottom: proportion of nuclei with various cell-cycle

stage annotations ordered by pseudotime. Nuclei were binned into 250 blocks.

(G) Left: Average fraction of ACRs that are accessible across pseudotime. The grey polygon

indicates standard deviation. Red windows indicate cell state transitions. Right: heatmap of

relative accessibility (relative to the row maximum) for each ACR (rows) across pseudotime

(columns). Nuclei were binned into 250 blocks ordered on pseudotime.

Figure 7. Dynamic and conserved *cis*-regulation in *A. thaliana* and *Z. mays* phloem

companion cell development.

(A) UMAP embedding based on whole-genome chromatin accessibility profiles of 4,655 *A.*

thaliana root nuclei.

(B) UMAP embedding of companion cell developmental trajectories in *A. thaliana* depicting cell

types (left) and pseudotime progression (right). Motifs, TFs, and ACRs with significant association

with pseudotime are shown as heatmaps below.

(C) UMAP embedding of integrated scRNA-seq and scATAC-seq cell and nuclei profiles derived

from *A. thaliana* roots.

(D) Marker gene expression levels for individual scRNA-seq cells for quiescent center (QC;

WOX5), pre-procambial/phloem SE precursors (PP/PSEP; *WOX4*), and phloem companion cells

(PCC; *SUC2*).

(E) Pseudobulk chromatin accessibility pile-ups for clusters labeled as QC, PP/PSEP, and PCC

across three marker genes associated with each cell type, respectively.

(F) Per-gene pseudotime shift scores from alignments between *Z. mays* and *A. thaliana*

companion cell development progressions, clustered by k-means into three groups.

(G) Distribution of gene-gene distances from the alignments, split by k-mean groups.

(H) Exemplary one-to-one homologs between *A. thaliana* and *Z. mays* for the three groups split

by pseudotime shifts. Acronyms: LRC, lateral root cap; LRP, lateral root primordia. SE, sieve

elements.

 SUPPLEMENTAL FIGURE LEGENDS

Figure S1: Evaluation and quality control of maize scATAC-seq

(**A**) Genome browser screenshot of chromatin accessibility from bulk and aggregated single-cell

ATAC-seq experiments. Chromatin accessibility profiles depict the *tb1* locus and the *tb1* enhancer

located approximately 67kb upstream.

(**B**) Binary accessibility scores from a random selection of 1,000 individual nuclei from each organ.

(**C**) Spearman's rho matrix comparing bulk ATAC-seq and aggregate scATAC-seq samples

across various organs. Sample codes are shorthand for assay type, sample, and replicate. For

example, s-R1 denotes single cell assay for seminal root replicate 1. The term b-L2 denotes a

bulk-ATAC assay for seedling replicate 2. Codes are as follows: b, bulk; s, single cell; R, seminal

root; C, crown root; E, ear; T, tassel; A, axillary bud; L, seedling. Numbers represent replicate.

(**D**) Average TSS enrichment (normalized read depth adjusted by the two 10 bp windows 1kb

away from TSSs) across all 56,575 cells. Grey polygon denotes the standard deviation.

(**E**) Fragment length distributions across 56,575 cells. The solid line and grey polygon represent

the average and standard deviation, respectively.

(**F**) Genotype-mixing experimental schematic.

(**G**) Scatterplot of per cell B73 and Mo17 SNP counts from a mixed-genotype experiment (V1

seedlings) colored by genotype classification.

(**H**) Posterior probabilities of individual barcodes (rows) highlighting the occurrence of cells with

B73, Mo17, and mixed (doublet) genotype identities.

(**I**) Genome browser screenshot of traditional bulk ATAC-seq from 7-day old seedling (row 1),

single-cell ATAC-seq from B73 seven day old seedlings (row 2), pooled B73 and Mo17 nuclei

(library ID: Seedling 2) single cell ATAC-seq from 7 day old seedlings (row 3), and the genotype-

sorted B73 (row 4) and Mo17 (row 5) alignments after sorting barcodes by genotype calls from

the B73-Mo17 scATAC-seq 7 day old seedling sample (row 3).

Figure S2. Cell calling and barcode quality control

(**A**) Comparison of normalized (0-1) read depths at the union of all peaks across bulk and single-cell samples (n=265,992) between replicated libraries, and between bulk and single-cell ATAC-seq assays.

(**B**) Enrichment plots centered on 2-kb windows surrounding TSSs for barcodes in each tissue. Grey polygons indicate the standard deviation across cells within the noted tissue.

(**C**) Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites mapping to within 2-kb of transcription start sites (TSSs). Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes.

(**D**) Fragment length distributions for each library. Solid lines indicate the average distribution across cells within the sample. Grey polygons represent the standard deviation across cells in the library.

(**E**) Knee plots illustrating $\log_{10}$ transformed cellular read depths of $\log_{10}$ ranked barcodes across libraries.

(**F**) Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5 integration sites derived from organeller sequences (chloroplast and mitochondrial) relative to the total number of unique Tn5 integration sites associated with cognate barcode. Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes.

Figure S3. *In silico* sorting via Latent Semantic Indexing

Standardized LSI accessibility scores (2^{nd} - 11^{th} dimensions) capped at ± 1.5 for:

(**A**) Axillary buds.

(**B**) Crown roots.

(**C**) Embryonic roots.

(**D**) Seedlings.

(**E**) Ear (pistillate inflorescence).

(**F**) Tassel (staminate inflorescence).

Figure S4. Clustering metrics and comparison of bulk gene accessibility and expression

(**A**) Parameter regularization of model coefficients (y-axes) with respect to ACR usage (x-axes;

proportion of nuclei with at least one Tn5 integration site in an ACR).

(**B**) Proportion of variance captured by the first 26 PCs. Inset: Spearman's correlation of principal

components with cell read depth (log_{10}-transformed).

(**C**) Number of accessible sites per cell (log_{10}).

(**D**) Proportion of Tn5 integrations within 2-kb of gene TSSs per cell.

(**E**) Co-localization of nuclei barcodes from different biological replicates for three organs.

(**F**) Comparison of bulk RNA-seq expression levels (y-axis, log_2 CPM) versus aggregate scATAC-

seq gene accessibility scores (x-axis, log_2 CPM) within an organ.

Figure S5. Cell-type annotation and GO enrichment

(**A**) UMAP embeddings of nuclei barcodes colored by low (grey) to high (dark purple) gene activity

values of cell type-specific marker genes.

(**B**) RNA *in situ* hybridization showing expression of *LOX10* in glume primordia and *GRFTF36* in

IM and SMs of staminate inflorescence; *LRR445* in the IM periphery and SPMs and *MYB89* in

the IM and suppressed bract primordia of pistillate inflorescence; Zm00001d038453 in ground

tissue of SAM and leaf primordia sections. Gene accessibility scores and predicted cell types are

shown on the right. *i*, tassel primordia. *ii,* ear primordia. *iii*, SAM/leaf. Black triangles point to the

glume primordia. Red triangles point to suppressed bract primordia. Size bars illustrate 100-um.

AM, axillary meristem; BS, bundle sheath; GC, guard cell; GM, ground meristem; GMC, guard

mother cell; GP, glume primordia; IM, inflorescence meristem; L1, layer 1; LFM, lower floral

meristem; SAM, shoot apical meristem; SBP, suppressed bract primordia; SM, spikelet meristem;

SPM; spikelet pair meristem; Stomatal PC, stomatal precursor; UKN, unknown; VEMI,

vascular/epidermal meristematic identity; VP, vascular parenchyma; XP, xylem parenchyma.

(**C**) Proportion of cells within subcluster (column) derived from one of six organs (rows).

Figure S6. Chromatin accessibility variation across plant cell types

(**A**) Row gene accessibility Z-scores across cell types. Cell type and subcluster labels are located

adjacent to the heatmap in identical order as the matrix. The first number after the cell-type label

indicates the major cluster number, while the second number represents the subcluster ID.

Legend colors: top left triangle, subcluster (corresponds to the sub-cluster annotation on

heatmap). bottom right triangle, major cluster (corresponds to the major cluster annotation on

heatmap).

(**B**) Row ACR chromatin accessibility Z-scores across cell types.

(**C**) Proportion of cells derived from one of six organs within each cell type/subcluster.

(**D**) Distribution of GO term enrichment across clusters, where the x-axis indicates the number of

clusters in which a GO term is significantly enriched.

Figure S7. Transcription factor motif variation underly dynamic accessible regions

(**A**) Comparison of cell type-specific ACR counts (by cell-type) and the number of unique Tn5

integration sites.

(**B**) Proportion of distal, genic and proximal ACRs per cell type.

(**C**) Ranked TF motif variability across cells, colored by TF family.

(**D**) Mean TF family motif enrichment (average deviation scores per cell type per TF family) across

cell-types scaled to +/- 1.

(**E**) Spearman's correlation coefficient (rho) between TF motifs (comparison of motif deviations

across all nuclei). Row and column colors represent TF motif families.

(**F**) TF motif deviations for 440 TF motifs (columns) per nucleus (rows). The TF family for each

corresponding motif is denoted by column header colors. Cluster identification (sub-cluster/major

cluster) are illustrated as row header colors on the right side of the heatmap.

(**G**) Relative gene accessibility scores for 2,423 maize transcription factors (columns) for 50,640

nuclei (rows). Nuclei are sorted by sub-cluster cell type (row color labels).

Figure S8: Capture of known long-range chromatin loops in maize

(**A**) Co-accessible ACRs at the *ZmRAP.2* locus in maize, with root-specific expression patterns,

across eight root-derived (left) and eight above-ground (right) cell types. The height of the loops

reflects the strength of co-accessibility. Pseudobulk chromatin accessibility tracks are shown

under co-accessible ACR linkages for each cell-type.

(**B**) Co-accessible ACRs at the *BX1* locus in maize, predominantly expressed in seedling tissue,

across eight seedling-derived (left) and eight non-seedling derived cell types. The height of the

loops reflects the strength of co-accessibility. Pseudobulk chromatin accessibility tracks are

shown under co-accessible ACR linkages for each cell-type.

Figure S9. Co-accessible ACRs reflect in vivo chromatin interactions driven by

coordinated TF activity

(**A**) Proportions of co-accessible ACR types, illustrated by toy examples.

(**B**) Proportion leaf Hi-C, H3K4me3-HiChIP and H3K27me3-HiChIP chromatin loops that overlap

co-accessible ACRs from leaf cell types (clusters with greater than 50% of cell derived from

seedlings).

(**C**) Top, log10 number of connections per ACR per cell type from all co-accessible ACRs split by

genomic context: distal, proximal, and genic. Bottom, $\log_{10}$ number of connections per ACR per

cell type from cell type-specific (purple) and non-specific (grey) co-accessible ACRs, split by

genomic context.

(**D**) Proportion of co-accessible classifications by cell type for all co-accessible ACRs (top) and

cell type-specific co-accessible ACRs (bottom).

(**E**) Jaccard similarity of motif composition between co-accessible ACR edges by cell type (colored

diamonds) relative to the same number of random ACR-ACR links, permuted 1,000 times for each

cell-type (grey boxplots). Box plots represent the interquartile range, grey lines indicate the

permuted range.

(**F**) Heatmaps of observed proportion of co-accessible ACRs with the same motif embedded

within link edges subtracted and divided by the expected proportion estimated by 1,000

permutations using random ACR-ACR links with the same number of co-accessible ACRs.

Figure S10. Pseudotime trajectory construction

(**A**) Overview of pseudotime developmental trajectory analysis from four organs: root, seedling,

tassel (staminate inflorescence), and ear (pistilate inflorescence).

(**B**) Endodermis development in crown roots.

(**C**) Cortex development in crown roots.

(**D**) Pericycle development in embryonic roots.

(**E**) Atrichoblast development in crown roots.

(**F**) Trichoblast development in crown roots.

(**G**) Trichoblast development in embryonic roots.

(**H**) Lateral root cap (LRC) development in crown roots.

(**I**) Lateral root cap (LRC) development in embryonic roots.

(**J**) Phloem sieve element (SE) development in crown roots.

(**K**) Companion cell development in crown roots.

(**L**) Phloem sieve element (SE) development in embryonic roots.

(**M**) Xylem development in crown roots.

(**N**) Xylem development in embryonic roots.

(**O**) Procambial development in crown roots.

(**P**) Guard cell development in seedling.

(**Q**) Subsidiary cell development in seedlings.

(**R**) Floral primordia development in staminate inflorescence (tassel).

(**S**) Floral primordia development in pistillate inflorescence (ear).

Figure S11. *Arabidopsis thaliana* root cell type atlas

(**A**) Knee plots for Arabidopsis thaliana root samples illustrating $\log_{10}$ transformed cellular read

depths of $\log_{10}$ ranked barcodes across two biological replicates.

(**B**) Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5

integration sites derived from organellar sequences (chloroplast and mitochondrial) relative to the

total number of unique Tn5 integration sites associated with each barcode from the two biological

replicates. Dashed red lines indicate the threshold of two standard deviations from the mean used

to filter lower quality barcodes.

(**C**) Density scatter plots of $\log_{10}$ transformed barcode read depths (x-axis) by the fraction of Tn5

integration sites mapping to within 2-kb of transcription start sites (TSSs). Dashed red lines

indicate the threshold of two standard deviations from the mean used to filter lower quality

barcodes.

(**D**) Average TSS enrichment (normalized read depth adjusted by the two 10 bp windows 1-kb

away from TSSs) across 5,001 *Arabidopsis thaliana* root barcodes (rows).

(**E**) UMAP (Uniform manifold approximation projection) embeddings of *Arabidopsis thaliana* root

barcodes colored by biological replicate.

(**F**) total number of accessible chromatin regions (ACRs);

(**G**) the proportion of Tn5 integration sites within 1-kb of TSSs.

(**H**) Relative gene accessibility for 27 known cell-type/domain restricted marker genes used to

inform cell-type annotation of Arabidopsis thaliana root clusters.

(**I**) Relative motif deviations for transcription factors with known cell-type specificities.

Figure S12. Dynamic and conserved chromatin accessibility across pseudotime between

Arabidopsis thaliana* and *Zea mays

(**A**) Pseudotime trajectories for Atrichoblast, Trichoblast, Lateral root cap (LRC), Cortex,

Endodermis, Lateral root primordia (LRP), Companion cells (CC), and Xylem development.

(**B**) Averaged alignments of conserved, shift early *Z. mays*, and shift early *A. thaliana* putative

orthologs.

(**C**) Pseudotime shifts of TF motifs between *A. thaliana* and *Z. mays*, clustered into k-means and

conserved groups.

(**D**) Distributions of motif-motif normalized distances between *Z. mays* and *A. thaliana* for the three

groups.

(**E**) Conserved motifs (n=142) ordered by pseudotime. Heatmaps for *A. thaliana* and *Z. mays*

have identical row orders.

(**F**) Averaged alignments of conserved, shift early *Z. mays*, and shift early *A. thaliana* groups

based on motif-motif global alignments from the dynamic time-warping algorithm.

(**G**) Examples of conserved, shift early *Z. mays*, and shift early *A. thaliana* motifs from both

species.

(**H**) Gene expression Z-scores across *A. thaliana* FAC sorted root cell-types for the TFs

recognizing the top four conserved motifs.

STAR*METHODS

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Bob Schmitz (schmitz@uga.edu).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

Raw and processed data has been deposited in NCBI GEO database under accession code

GSE155178. Code used throughout the analysis can be found in the following GitHub repository:

https://github.com/plantformatics/maize_single_cell_cis_regulatory_atlas. We also released an R

package for pre-processing, normalization, clustering, and other downstream analytical steps into

streamlined toolkit of scATAC-seq data can be found in the following GitHub repository:

https://github.com/plantformatics/Socrates.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Growth conditions

For libraries derived from seedlings, kernels from genotypes B73 and Mo17 were obtained from

USDA National Plant Germplasm System (https://npgsweb.ars-grin.gov) and sown in Sungro

Horticulture professional growing mix (Sungro Horticulture Canada Ltd.). Soil was saturated with

tap water and placed under a 50/50 mixture of 4100K (Sylvania Supersaver Cool White Delux

F34CWX/SS, 34W) and 3000K (GE Ecolux w/ starcoat, F40CX30ECO, 40W) lighting. Seedlings

were grown under a photoperiod of 16 hours of light, eight hours of dark. The temperature was

approximately 25°C during light hours with a relative humidity of approximately 54%.

Maize seedlings

Above ground seedling tissues were harvested between 8 and 9 AM six days (V1-stage) after

sowing. We used both fresh (B73/Mo17 pooled) and flash frozen (B73 only) seedling tissue to

construct scATAC-seq libraries (**Table S1**).

Maize roots

Maize root samples were obtained as follows: B73 kernels were sterilized with 70% EtOH

treatment for 5 minutes. After removing the ethanol solution, kernels were suspended with 50%

bleach for 30 minutes, followed by five washes with autoclaved Milli-Q water. Sterilized kernels

were then sown onto mesh plates with half strength MS (Phytotech laboratories, catalog: M519)

media and wrapped in Millipore tape. Plates were incubated in a Percival growth chamber with a

photoperiod of 16 hours of light, eight hours of dark. The growth chamber temperature was set to

25°C with a relative humidity of approximately 60%. Apical root tips (bottom 2 cm) of seminal and

primary root samples were harvested six days (V1-stage) after sowing between 8 and 9 am.

Crown root samples (21 days after sowing) were derived from the three developmental zones of

greenhouse grown B73 plants between 8 and 9 am and rinsed with sterile water 3 times.

Maize Inflorescence

Data generated from young inflorescence (ear and tassel primordia) were derived from B73 maize

grown in the greenhouse. Inflorescence primordia were extracted from shoots harvested

approximately one month (V7-stage, 2-4 mm) after sowing, between 8 and 9 AM. Inflorescence

primordia between three and eight millimeters from the base to the apical tip were placed in sterile

water and used for nuclei isolation.

1000

1001 ***Maize axillary buds***

1002 Axillary buds (~30 samples per library) were taken from B73 maize plants grown in the

1003 greenhouse at approximately the same developmental stage (V7) as tassel and ear primordia.

1004

1005 ***Arabidopsis roots***

1006 Seven-day old *A. thaliana* roots were prepared similarly as for maize with the exception of deriving

1007 nuclei from whole roots.

1008

METHOD DETAILS

Single cell ATAC-seq library preparation

Each library was prepared by mixing at least three independent biological samples (3-4 seedlings,

3 tassel or ear primordia, 12-14 root tips, 12-14 crown root samples, ~30 axillary buds, and 100-

200 *A. thaliana* whole roots). One scATAC-seq library (B73 seedling) was derived from flash

frozen tissue (liquid nitrogen, followed by 7-day -80°C storage), while the remaining libraries were

constructed with freshly harvested tissue (**Table S1**).

To isolate individual plant nuclei, fresh or flash frozen tissue from multiple biological

samples were placed on petri dishes and vigorous chopped with a No. 2 razor blade for two

minutes in ~500 uL LB01 buffer (15mM Tris pH 7.5, 2mM EDTA, 0.5mM Spermine, 80mM KCl,

20mM NaCl, 15mM 2-ME, 0.15% TrixtonX-100). Homogenized tissue was then filtered through

two layers of miracloth, stained with DAPI to a final concentration of ~1uM and loaded onto a

Beckman Coulter MoFlo XDP flow cytometer instrument. A total of 120,000 nuclei were sorted for

each sample across four catch tubes (30,000 nuclei each) containing 200 uL LB01. Isolated nuclei

were spun down in a swinging-bucket (5 minutes, 500 rcf) centrifuge resuspended in 10uL LB01,

pooled, and then visualized on a hemocytometer with a fluorescent microscope. Nuclei

suspensions were then spun down (5 minutes, 500 rcf) and resuspended in diluted nuclei buffer

(10X Genomics) to a final concentration of 3,200 nuclei per uL and used as input for scATAC-seq

library preparation (5 uL; 16,000 nuclei total). Samples were kept on ice for all intermittent steps.

For B73/Mo17 mixed library, we pooled 8,000 nuclei from both B73 and Mo17 that were

independently isolated. Single-cell ATAC-seq libraries were constructed according to the

manufacture's instruction (10X Genomics, catalog: 1000176). Libraries were sequenced with

Illumina NovaSeq 6000 in dual-index mode with eight and 16 cycles for i7 and i5 index,

respectively.

1033

Single nuclei RNA-seq library preparation

We prepared snRNA-seq libraries from two biological replicates, each composed of three

independent 7-day old B73 seedlings. Seedlings were vigorously chopped with a No. 2 razor

blade on a petri dish in 500 uL of nuclei isolation buffer (Phosphate-Buffered Saline [PBS;

ThermoFisher], 500U SUPERase RNase inhibitor [Invitrogen], 1mM 1,4-Dithiothreitol [DTT;

Millipore Sigma], and 0.05% Triton X-100 [Millipore Sigma]). Homogenized tissue in nuclei

isolation buffer was filtered through a 40-um cell strainer (pluriSelect) and spun at 500 rcf for 5

minutes. The supernatant was discarded, followed by two more wash (500 uL nuclei isolation

buffer) and centrifugation steps (500 rcf for 5 minutes), discarding the supernatant and

resuspending in 10 uL nuclei isolation buffer lacking Triton X-100. The concentration of nuclei in

solution was estimated on a hemocytometer under a fluorescent microscope and adjusted to

2,000 nuclei per uL with nuclease-free water. Single-nuclei RNA-seq libraries were prepared from

a total of 16,000 nuclei per library following the manufactures instructions for the Single Cell Gene

Expression 3' V3 library kit (10X Genomics, catalog: 1000269). Libraries were sequenced on an

Illumina NovaSeq 6000 in dual-index mode.

1049

In situ hybridizations

3-4mm tassel and ear primordia and young seedlings from the maize B73 inbred line were

dissected and fixed in a cold paraformaldehyde acetic acid solution (4% PFA) for 48 hours.

Following dehydration through a graded ethanol series and clearing of the tissue with a Histo-

clear II solution (Electron Microscopy Sciences), samples were embedded using Paraplast Plus

tissue embedding media (McCormick Scientific). 8mm sections were hybridized at 56°C with

antisense probes labelled with digoxigenin (DIG RNA labeling mix, Roche), and detected using

NBT/BCIP (Roche). Probes were synthesized by *in vitro* transcription (T7 RNA polymerase,

Promega) of PCR products obtained from embryo cDNA or from digested full-length cDNA clones.

The vectors and primers used for probe design are listed in **Table S18**.

1060

QUANTIFICATION AND STATISTICAL ANALYSIS

scATAC-seq raw reads processing

The following data processing was performed using each tissue and/or replicate independently

unless noted otherwise. Raw BCL files were demultiplexed and convert into fastq format using

the default settings of the 10X Genomics tool *cellranger-atac makefastq* (v1.2.0). Partial raw read

processing (adapter/quality trimming, mapping and barcode attachment/correction) was carried

out with *cellranger-atac count* (v1.2.0) using AGPv4 of the maize B73 reference genome (Jiao et

al., 2017). Properly paired, uniquely mapped reads with mapping quality greater than 10 were

retained using *samtools view* (v1.6; -f 3 -q 10) and by filtering reads with XA tags (Li et al., 2009).

Duplicate fragments were collapsed on a per-nucleus basis using *picardtools*

(http://broadinstitute.github.io/picard) *MarkDuplicates* (v2.16; BARCODE_TAG=CB

REMOVE_DUPLICATES=TRUE). Reads mapping to mitochondrial and chloroplast genomes

were counted for each barcode, then excluded from downstream analysis. We removed reads

representing potential artifacts by excluding alignments coincident with a blacklist of regions

composed of low-complexity and homopolymeric sequences (*RepeatMasker* v4.07) (AFA Smit,

2013-2015), nuclear sequences with homology (greater than 80% identity and coverage) to

mitochondrial and chloroplast genomes (*BLAST+* v2.7.1) (Camacho et al., 2009), regions

exhibiting Tn5 integration bias from Tn5-treated genomic DNA (1-kb windows with greater than

2-fold coverage over the genome-wide median), and potential collapsed sequences in the

reference (1-kb windows with greater than 2-fold coverage over the genome-wide median using

ChIP-seq input). Genomic Tn5 and ChIP input data were acquired from Ricci, Lu and Ji et al.

BAM alignments were then converted to single base-pair Tn5 integration sites in BED format by

adjusting coordinates of reads mapping to positive and negative strands by +4 and -5,

respectively, and retaining only unique Tn5 integration sites for each distinct barcode. Sequencing

saturation was calculated as the proportion of unique reads relative to the estimated library

complexity output by the *MarkDuplicates* function apart of picardtools.

1087

Cell calling

To identify high-quality nuclei (a term used interchangeably with "barcodes") using the filtered set

of alignments, we implemented heuristic cutoffs for genomic context and sequencing depth

indicative of high-quality nuclei. Specifically, we fit a smoothed spline to the $\log_{10}$ transformed

unique Tn5 integration sites per nucleus (response) against the ordered $\log_{10}$ barcode rank

(decreasing per-nucleus unique Tn5 integration site counts) using the *smooth.spline* function

(spar=0.01) from base R (Team, 2013). We then used the fitted values from the smoothed spline

model to estimate the first derivative (slope), taking the local minima within the first 16,000

barcodes as a potential knee/inflection point (16,000 was selected to match the maximum number

of input nuclei). We set the unique Tn5 library depth threshold to the lesser of 1,000 reads and

the knee/inflection point, excluding all barcodes below the threshold. Spurious integration patterns

throughout the genome can be representative of incomplete Tn5 integration, fragmented/low-

quality nuclei, or poor sequence recovery, among other sources of technical noise. In contrast,

high quality nuclei often demonstrate a strong aggregate accessibility signal near TSSs.

Therefore, we implemented two approaches for estimating signal-noise ratios in our scATAC-seq

data. First, nuclei below two standard deviations from the mean fraction of reads mapping to within

2-kb of TSSs were removed on a per-library basis. Then, we estimated TSS enrichment scores

by calculating the average per-bp coverage of 2-kb windows surrounding TSSs, scaling by the

average per-bp coverage of the first and last 100-bp in the window (background estimate; average

of 1-100-bp and 1901-2000-bp), and smoothing the scaled signal with rolling-means (R package;

Zoo). Per barcode TSS enrichment scores were taken as the maximum signal within 250-bp of

the TSS. Lastly, for each library, we removed any barcode with a proportion of reads mapping to

chloroplast and mitochondrial genomes greater than two standard deviations from the mean of

the library.

1112

Detection of multiplet droplets

To estimate the empirical proportion of doublets present in our data, we demultiplexed the two-

genotype (B73 and Mo17) pooled seedling scATAC-seq sample and assessed the proportion of

barcodes reflecting a mixtures of reads derived from both genotypes. Specifically, B73 and Mo17

whole genome short read resequencing data were acquired from PRJNA338953. Paired-end

reads were quality and adapter trimmed with *fastp* (v0.19.5) (Chen et al., 2018) and aligned to

the B73 v4 maize reference genome (Jiao et al., 2017) using *BWA mem* (Li, 2013) with non-

default settings (-MT 1). Duplicate reads were removed using *samtools rmdup* (Li et al., 2009)

(v1.6). The genomic coordinates of short nucleotide variants (SNVs; single nucleotide

polymorphisms [SNPs] and small insertions/deletions [INDELs]) for both genotypes were

identified using *freebayes* (Garrison and Marth, 2012) (v1.0.0) with non-default settings (--min-

repeat-entropy 1 --min-alternate-fraction 0.05). Only biallelic SNPs – requiring at least 5 reads per

genotype where B73 and Mo17 were homozygous for reference and alternate nucleotides,

respectively – were retained. Genotypes were called by modeling allele counts as a binomial

distribution with a term accounting for the sequencing error rate, E_t (determined empirically as

the fraction of SNPs failing to match either allele), estimating posterior probabilities via Bayes

theorem, and assigning the genotype (or mixture of genotypes) with the greatest probability (Eq.

1-7). Specifically, the probability to observe k out of n SNPs from B73 can be modeled as a

binomial distribution for each B73 (A_l), Mo17 (A_2), and doublet barcode state (N) (Eq. 1-3):

1132

1133   1. $P(k|A_1) = \binom{n}{k} \times E_t^{n-k} \times (1 - E_t)^k$

1134

1135   2. $P(k|A_2) = \binom{n}{k} \times (1 - E_t)^{n-k} \times E_t^{k}$

1136

1137   3. $P(k|N) = \binom{n}{k} \times (0.5)^k \times (0.5)^{n-k}$

1138

1139   Let $P(A_1|k)$, $P(A_2|k)$, and $P(N|k)$ reflect posterior probabilities for genotypes B73, Mo17, and

1140   doublet barcodes given k allele counts from B73; posterior probabilities can be estimated as

1141   follows (Eq. 4-6):

1142

1143   4. $P(A_1|k) = \dfrac{P(k|A_1) \times P(A_1)}{\sum_{i=0}^{n} P(k|A_i) \times P(A_i)}$

1144

1145   5. $P(A_2|k) = \dfrac{P(k|A_2) \times P(A_2)}{\sum_{i=0}^{n} P(k|A_i) \times P(A_i)}$

1146

1147   6. $P(N|k) = \dfrac{P(k|N) \times P(N)}{\sum_{i=0}^{n} P(k|A_i) \times P(A_i)}$

1148

1149   Finally, the genotype called for each barcode was determined as the event with the greatest

1150   posterior probability (Eq. 7):

1151

1152   7. $\max\{P(A_1|k), P(A_2|k), P(N|k)\}$

1153

In silico sorting

To provide sufficient sensitivity for peak calling prior to clustering, we followed an *in-silico* sorting

strategy to identify crude clusters of similar cells within each organ (Cusanovich et al., 2018). To

do so, we generate a binary matrix representing the presence/absence of Tn5 integration sites in

1-kb windows across all cells in a given organ. Bins with less than 1% accessible cells and cells

with less than 100 accessible bins were removed. This binary matrix was then transformed using

the matrix normalization method term-frequency inverse document-frequency (TF-IDF). Briefly,

the TF term was estimated by weighting binary counts at each bin by the total number of bins

containing Tn5 integration sites in a given cell, scaling each cell to sum to 100,000, adding a

pseudo-count of one, and log transforming the resulting values to reduce the effects of outliers in

downstream processing. The IDF term was calculated as the log transformed ratio of the total

number of nuclei to the number of nuclei that were marked as accessible for a given bin. We add

a pseudo-count of one to the inverse frequency term to avoid taking the log of zero. The TF-IDF

scaled matrix was estimated by taking the dot product of the TF and IDF matrices. To enable

faster downstream computation, we kept the top 25,000 bins with the greatest TF-IDF variance

across nuclei. The reduced TF-IDF matrix was denoised with singular value decomposition (SVD),

retaining the 2^{nd} – 11^{th} dimensions (termed Latent Semantic Indexing, LSI). Each row was

centered and standardized, capping the values at $\pm$ 1.5. Crude clusters were visually identified

using ward.D2 hierarchical bi-clustering on the cosine distances of LSI nuclei and bin

embeddings.

1174

ACR identification

ACRs were identified by treating each bulk and single-cell ATAC-seq library as a traditional bulk

ATAC-seq library. Aligned reads were filtered by mapping quality greater than 10, and duplicate

reads were removed via *samtools rmdup.* We then identified ACRs for each library by converting

the BAM alignments in BED format, adjusting the coordinates to reflect single-base Tn5

integrations, and running *MACS2* (Zhang et al., 2008) with non-default parameters: --extsize 150

--shift -75 --nomodel --keep-dup all. A final set of ACRs for comparing bulk and aggregate

scATAC-seq libraries (**Figure S1**) was constructed by taking the union of ACRs across all

libraries. To leverage the increased sensitivity afforded by cell-type resolved cluster information

while ensuring robust reproducibility in ACR identification, we generated pseudo-replicated bulk

alignments using the LSI-based crude clusters (see above, "*In-silico* sorting"). Pseudo-replicates

were constructed by randomly allocating nuclei from each cluster into two groups, with a third

group composed of all cells from the cluster (cluster bulk). These groupings were used to

concatenate Tn5 integration sites corresponding to the nuclei from each group into three BED

files. ACRs were then identified from the enrichment of Tn5 integration sites from the pseudo-

replicate or cluster bulk aggregates using *MACS2* run with non-default parameters: --extsize 150

--shift -75 --nomodel --keep-dup all. ACRs from both pseudo-replicates and the cluster bulk were

intersected with BEDtools, retaining ACRs on the conditional intersection of all three groupings

(both pseudo-replicates and the cluster bulk) by at least 25% overlap. The remaining ACRs were

then redefined as 500-bp windows centered on the ACR coverage summit. To integrate

information across all clusters, ACRs from each cluster were concatenated into a single master

list. Lastly, overlapping ACRs were filtered recursively to retain the ACR with the greater

normalized kernel Tn5 integration density as previously described (Satpathy et al., 2019).

1198

Nuclei clustering

Starting with a binary nucleus x ACR matrix, we first removed ACRs that were accessible in less

than 0.5% of all nuclei, and filtered nuclei less than 50 accessible ACRs. Inspired by recent

developments in modeling single-cell RNA-seq data (Hafemeister and Satija, 2019), we

developed a regularized quasibinomial logistic framework that overcomes noise inherent to

sparse, binary scATAC-seq data by pooling information across ACRs while simultaneously

removing variation due to technical effects, particularly those stemming from differences in

barcode sequencing depths. First, a subset of 5,000 representative ACRs selected by kernel

density sampling of ACR usage (fraction nuclei that are accessible at a given ACR) were used to

model the parameters of each ACR, using ACR usage as a covariate in a generalized linear

model. Specifically, the expected accessibility of an ACR, y_i, can be estimated with a generalized

linear model containing a binomial error distribution and logit-link function, and an overdispersion

term with a quasibinomial probability density function (Eq. 8).

1212

1213   8. $\mathbb{E}(y_i) \sim \beta_0 + \beta_1 log10(t)$

1214

1215   Where t is a vector of the sums of accessible ACRs across cell j (Eq. 9):

1216

1217   9. $t = \sum_i y_{ij}$

1218

To prevent over-fitting and ensure robust estimates in light of sampling noise, we learned the

global regularized model parameters, including overdispersion, using the representative ACRs by

fitting each parameter against the $\log_{10}$ fraction of accessible nuclei via kernel regression,

resulting in smoothed parameter estimates across the spectrum of ACR accessibility penetrance

present in these data. The learned global regularized model parameters were then used to

constrain fitted values across all ACRs for each nucleus with a simple affine transformation. To

account for technical variation among nuclei (variation in barcode $\log_{10}$ transformed read-depth,

in particular) we calculated Pearson residuals for each ACR, scaling the residuals by the

regularized dispersion estimate and centering values via mean subtraction, representing

variance-stabilized and read-depth normalized values of accessibility for a nucleus at a given

ACR. We note that this method is amenable to calculating residuals that account for additional

sources of technical variation, including categorical and numeric covariates, that may obscure

biological signal, such as batch effects, proportion of mitochondrial reads, etc.

The dimensionality of the Pearson residual matrix was reduced using singular value

decomposition (SVD) implemented by the R package *irlba* (Witten et al., 2009), retaining the first

25 left singular vectors scaled by singular values (hereafter referred to as nuclei embeddings),

analogous to principal components (PCs) on an uncentered matrix. Nuclei embeddings were then

standardized across components and filtered to remove components correlated with barcode read

depth (Spearman's rho > 0.7). We further reduced the dimensionality of the nuclei embedding

with Uniform Manifold Approximation Projection (UMAP) via the R implementation of *umap-learn*

(min_dist = 0.1, k=50, metric="euclidean"). Nuclei were clustered with the *Seurat* v3 (Stuart et al.,

2019) framework and Louvain clustering on a k=50 nearest neighborhood graph at a resolution

of 0.02 with 100 iterations and 100 random starts. Clusters with aggregated read depths less than

1.5M were removed. To filter outliers in the UMAP embedding, we estimated the mean distance

for each nucleus with its k (k=50) nearest neighbors and removed nuclei greater than 3 standard

deviations from the mean.

We observed fine-scale heterogeneity within major clusters, thus we repeated our

clustering pipeline for each major cluster independently by partitioning the SVD embedding into

the top 20 components, L2 normalizing nuclei embeddings across components, and projecting

the L2-normalized embeddings into the UMAP space. Subclusters of nuclei were identified by

Louvain clustering on the L2 normalized SVD embedding (resolution set manually, range = 0.6 –

1.0) with 20 nearest neighbors, filtering outlier nuclei more than 2 standard deviations from the

mean distance of 25 nearest neighbors within each cluster.

For analysis of chromatin accessibility across clusters, we assembled a matrix of clusters

by ACRs by aggregating the number of single-base resolution Tn5 integration sites from nuclei

within the same cluster for each ACR, analogous to normalizing by the proportion of reads in

peaks for each cluster. To account for differences in read depth and other technical factors, the

raw counts were transformed with *edgeR's* "cpm" (log=T, prior.count=5) as previously described

(Corces et al., 2018). Log-transformed ACR coverage scores were quantile normalized using

"normalize.quantiles" with the R package, *preprocessCore.* Finally, to aid data visualization, we

estimated per ACR Z-scores across clusters by mean subtraction and standardization (identical

to row-wise execution of the R function, "scale").

1261

Identification of co-accessible ACRs

Recent experiments of population-level chromatin accessibility found that pairwise correlations of

accessibility among ACRs recapitulates higher-order chromatin interactions observed in Hi-C and

other chromatin architecture experiments (Gate et al., 2018). A similar framework was applied to

populations of single cells, which showed that co-accessible ACRs are typically more conserved

and functionally associated (Buenrostro et al., 2015). To identify potentially functional co-

accessible ACRs, we applied a recently developed method, *Cicero* (Pliner et al., 2018), that

estimates regularized correlation scores (ranging from -1 to 1) among nearby ACRs with graphical

LASSO to penalize potential interactions by physical distances. Using the binary nuclei x ACR

matrix as input, we subset nuclei by their subcluster IDs and estimated co-accessibility among

ACRs within 500-kb for each of the 92 clusters, independently. *Cicero* was run by applying a

background sample of 100 random regions, and 15 nuclei pseudo-aggregates based on k-

nearest-neighbors derived from the UMAP coordinates. To control the false discovery rate (FDR)

of co-accessible ACR calls, we shuffled the nuclei x ACR matrix such that the total number of

reads per ACR and reads per nucleus were identical to the original matrix. We then repeated co-

accessible ACR identification with the shuffled matrix, keeping the original parameters to *Cicero*

unchanged. Empirical FDR cluster-specific cut-offs were constructed by identifying the minimum

positive co-accessibility score in the background where the FDR < 0.05. Co-accessible links below

cluster-specific thresholds were removed. Co-accessible ACRs passing thresholds were

compared with previously published HiC and HiChIP data sets derived from maize seedling and

pistillate inflorescence primordia (Ricci et al., 2019).

1283

Estimation of gene accessibility scores

Chromatin accessibility at TSSs and gene bodies exhibit marked correlation with transcription

output in bulk samples (**Extended Data Fig. 3f**). To aid the identification of marker genes

underlying distinct cell-types, we used *Cicero* to estimate gene activity scores. *Cicero* models

gene activity as a weighted accessibility score that integrates both proximal and distal regulatory

elements linked to a single gene by co-accessibility analysis (see above section "Identification of

co-accessible ACRs"). Relative gene accessibility scores per nucleus were estimated by taking a

weighted average (3:1, gene body score to proximal/distal activity) of the scaled number of reads

mapping to gene bodies for each barcode (summing to 1) with the *Cicero* estimate of gene activity

derived from ACRs mapping to 1-kb upstream of gene TSSs and their associated distal ACRs

linked by co-accessible ACRs passing FDR < 0.05 thresholds (connected ACRs were constrained

to a minimum and maximum intervening distance of 1- and 500-kb, respectively). These weighted

gene accessibility scores were rescaled such that gene accessibility scores for a given nucleus

summed to 1.

Relative gene accessibility scores exhibited a bimodal distribution with relative gene

accessibility values near zero resembling low or non-expressed genes. We applied a gaussian

mixture-model (two distributions) based scaling step per cluster to reduce noise introduced by

genes with low gene accessibility. Briefly, the average gene accessibility across nuclei was fit to

a two distribution gaussian mixture model in each cluster using the R package *mclust*. We

estimated cluster-specific scaling parameters determined as the 5% quantile of non-zero gene

accessibility values of genes from the gaussian distribution with the larger mean, for each cluster.

This parameter was then used to scale gene accessibility scores for all genes in each nucleus

within the cluster. Scaled gene accessibility scores were rounded to the nearest integer and

normalized across all nuclei and clusters using nucleus-specific size factors estimated as the total

gene accessibility of a nucleus divided by the exponential of the mean of log-transformed gene

accessibility sums across nuclei. To aid visualization, we smoothed normalized gene accessibility

scores by estimating a diffusion nearest neighbor graph (k=15) using the SVD embedding with 3

steps similar to the *run_MAGIC* function in R package *snapATAC* (Fang et al., 2020).

Downstream analyzes based on binarized gene accessibility were conducted by simply

converting normalized (non-smoothed) accessibility scores to 1 for all positive values.

1314

Cell-type annotation

To identify and annotate cell types for each barcode, we identified marker genes known to localize

to discrete cell types or domains expected in the sampled tissues/organs based on extensive

review of the literature (**Table S2**). To enable gene accessibility comparisons among clusters, we

generated three pseudo-replicates for each cluster by resampling nuclei within the cluster such

that all cluster pseudo-replicates contained the mean number of nuclei across clusters (number

of nuclei per pseudo-replicate = 552) without replacement when possible. To identify genes with

increased accessibility relative to other clusters, we constructed a reference panel with three

pseudo-replicates by uniformly sampling nuclei without replacement from each organ (number of

nuclei per organ = 92), with a total of 552 nuclei per reference panel pseudo-replicate. Read

counts per gene were summed across nuclei within each pseudo-replicate. Using the *DESeq2* R

package, we identified genes with significantly different (FDR < 0.01) accessibility profiles

between each cluster and the reference panel.

The list of significantly differentially accessible genes was filtered to retain the genes on

our list of cell type specific markers. We initially ranked the top three marker genes in each cluster

by their test statistics. To account for clusters containing small proportions of contaminating nuclei

of a different cell type, we adjusted the test statistics using a previously described method

(Cusanovich et al., 2018), effectively scaling marker activity scores by the proportion of nuclei in

the cluster that were derived from an organ in which the marker gene i is an expected cell type.

Clusters where the top three markers corresponded to the same cell type were annotated with

the consensus cell type.

As an independent method for cell-type annotation, we devised a resampling and

normalization procedure on the $\log_2$ fold-change values of marker genes to evaluate cell-type

enrichment across all possible cell types for each cluster, normalizing enrichment scores by

random permutations accounting for different numbers of markers associated with each cell type.

Briefly, starting with differential gene accessibility information for each cluster, we iterated over all

cell types, extracting markers associated with the cell type of interest. Then, we summed the $\log_2$

fold-changes values of all markers and multiplied the sum by the proportion of markers passing

heuristic thresholds (fold-change > 2 and FDR < 0.01). This score was subtracted by the average

of 1,000 random permuted scores from combinations of markers from the remaining cell types

(selecting the same number of random genes as the cell type of interest) and divided by the

standard deviation of the permuted scores. Cell-type enrichment scores in each cluster were

scaled from zero to one by dividing each cell-type enrichment score by the maximum scores

across possible cell types. This approach is effective in normalizing differences arising from

varying numbers of markers specified for each cell type. Additionally, cell-type annotation scores

for clusters with mixed or unknown identity are approximately equally distributed, thus controlling

ascertainment bias stemming from marker gene selection. Stated differently, an advantage of this

approach is that clusters corresponding to cell types with few or no markers in the tested list are

left unassigned as their enrichment scores do not deviate significantly from background levels.

Finally, scaled cell-type enrichment scores greater than 0.9 were taken as possible annotations

and intersected with putative cell-type labels from the marker ranking approach described above.

For clusters with ambiguous marker gene labels, we developed a logistic regression

classifier to identify putative cell types based on whole-genome gene accessibility scores of well-

annotated cells. First, we counted the number of Tn5 integration sites per cell overlapping 2-kb

upstream to 500-bp downstream of each gene. Read counts were transformed by trimmed mean

of M-values (TMM) to enable intra and inter-nucleus comparisons using *edgeR* (Robinson et al.,

2010), scaling gene accessibility scores in each nucleus with counts per million. Next, we

estimated cell-type enrichment scores for each nucleus by calculating the mean accessibility

scores of markers for a given cell type, subtracting the mean background signal defined as 1,000

sets of averaged randomly sampled genes (each set had the same number of genes as the

number of markers), divided by the standard deviation of the background signal. Enrichment

scores for each nucleus were transformed into a probability distribution by dividing by the sum of

cell-type enrichment scores. For each nucleus, we compared the top two most likely cell types,

retaining nuclei where the top predicted cell type had a two-fold greater probability than the next

most likely assignment. We used these high-confidence cells to train a regularized logistic

multinominal classifier with the R package, *glmnet*. Cell-type classifications with less than 10

nuclei in the training set were excluded. We used a LASSO L1 penalty to regularize the logistic

classifier, modeling the training set of nuclei as observations and TMM gene accessibility scores

as variables. We balanced observations by weighting by the inverse frequency of cell types in the

training set. The model was trained with 10-folds and evaluated by testing a 20% hold-out set of

nuclei. The predicted cell type for each nucleus in the atlas was taken as the cell type with the

greatest probability if the probability ratio between the best and next best assignment was greater

than five-fold, otherwise labeled as 'unknown'. Using these per-cell assignments, we defined

subclusters as the majority cell type if greater than 50% of nuclei in the cluster were in agreement,

labelling clusters with two or more majority cell types as 'mixed' and all other clusters as

'unknown'. All cell-type labels from these three automated approaches were manually reviewed

by careful evaluation with UMAP gene accessibility score embeddings and cluster aggregated

coverages for all marker genes and refined *ad hoc*.

1383

Cell-cycle annotation

Cell cycle annotation was performed similarly as cell-type annotation. Briefly, we acquired cell-

cycle marker genes from Nelms et al. 2019, selecting 35 markers at random for each cell stage

(Nelms and Walbot, 2019). The rationale behind selecting equivalent numbers of markers per

stage was to prevent biasing cell cycle annotations to cycle stages with more markers, while 35

markers was the minimum gene count across all stages (mitosis). For each stage, we subset the

nuclei by gene accessibility (TMM) matrix by the cognate stage, and summed accessibility scores

for each nucleus. This cell-cycle stage score was then standardized using the mean and standard

deviation of 1,000 permutation of 35 random cell-cycle stage genes, excluding the focal stage. Z-

scores corresponding to each cell-cycle stage were converted into probabilities using the R

function *pnorm*. Per nucleus posterior cell-cycle probabilities were estimated using Bayes

theorem with each cell-cycle stage prior probability set to 0.2 (1/5, for five stages: G1, G1/S, S,

G2/M, M). The cell-cycle stage with the maximum probability was selected as the most likely cell

stage. Nuclei with multiple cell-cycle annotations with equal maximum probability were considered

"ambiguous".

1399

snRNA-seq data processing

Raw fastq files from each snRNA-seq seedling library (across two biological replicates) were

processed with *cellranger count* v4.0 to align reads to AGPv4 of the maize B73 reference genome

(Jiao et al., 2017). BAM files were filtered to remove multiple mapping reads using a mapping

quality filter selecting reads with MQ greater than or equal to 30. The number of nuclear,

organeller and transcript-derived unique molecular identifiers (UMIs) reads for each barcode were

tabulated from the filtered BAM file. Barcodes with less than 1,000 total UMIs and less than 500

genes with at least one UMI were removed. We then estimated the Z-score distributions for the

proportion of mitochondrial, chloroplast, nuclear, and transcript derived UMIs across barcodes.

Barcodes above 1 standard deviation (Z-score less than 1) from the mean proportion of UMIs

derived from mitochondrial and chloroplast genomes were removed. Likewise, barcodes below 1

standard deviation from the mean proportion of UMIs derived from the nuclear genome were

removed.

1413

Integration of scATAC-seq and snRNA-seq data

To integrate scATAC-seq and snRNA-seq data into a shared embedding, we input gene

accessibility scores and gene expression values from all seedling-derived nuclei passing quality

filters described above using *liger* with the function *createLiger* (Welch et al., 2019). Each data

set was normalized, subset by highly variable genes, and scaled using the functions *normalize,*

selectGenes, and *scaleNotCenter,* sequentially with default arguments. An integrated non-

negative matrix factorization (iNMF) embedding was constructed from the gene by nuclei

scATAC-seq and snRNA-seq matrices using *optimizeALS* with default settings (k=20, lambda=5).

The iNMF embedding was quantile normalized with *quantile_norm* and non-default settings

(do.center=FALSE). Louvain clusters from the normalized iNMF embedding were identified at a

resolution of 0.25 with *louvainCluster.* To visualize the integrated assays, we used *runUMAP* with

non-default settings (n_neighbors = 20, min_dist = 0.01). Differentially accessible and expressed

genes per cluster were identified using *runWilcoxon* requiring FDR less than 0.05 and a $\log_2$ fold

change greater 0.25 using the integrated embedding (both gene accessibility and expression

across all co-embedded nuclei), gene accessibility in isolation (scATAC-seq nuclei only), and

gene expression in isolation (snRNA-seq nuclei only). Differentially accessible ACRs from the

normalized (with *liger* function *normalize*) sparse ACR by nuclei matrix were identified using

identical heuristic thresholds as for gene expression and accessibility.

To impute ACR accessibility in snRNA-seq derived-nuclei and gene expression values in

scATAC-seq nuclei, we ran *imputeKNN* from the *liger* package using either the scATAC-seq or

snRNA-seq nuclei as reference cells. We then used the imputed gene expression and ACR

accessibility matrices, constrained to only differentially accessible ACRs (n=55,939), to identify

significantly associated gene-to-peak linkages with the *liger* function *linkGenesAndPeaks* with

non-default settings (dist = 'spearman', alpha = 0.05). To remove potential false positives, we

shuffled the imputed ACR and gene nuclei matrices and repeated gene-to-peak linkage

identification using the same arguments. We then estimated FDR empirically over a grid of a 100

possible correlation values in both the negative and positive directions by identifying correlation

cut-offs that removed 95% of gene-to-peak linkages from the shuffled matrices. We then filtered

the non-shuffled gene-to-peak linkages according to the thresholds identified from the empirical

FDR estimates.

1444

STARR-seq analysis

Single bp-resolution enhancer activities were available from a previous study (Ricci et al., 2019).

Enhancer activity (defined as the $\log_2$ ratio between RNA and DNA input fragments scaled per

million) for each ACR was taken as the maximum over the entire ACR. A control set of regions

was generated to match each ACR with the following criteria: (*i*) GC content within 5%, (*ii*)

physically constrained to within 50-kb of an ACR, and (*iii*) the same length (500-bp) distribution.

The same set of control regions was used throughout the analysis.

1452

Analysis of differential chromatin accessibility

Next, we implemented a logistic regression framework based on binarized ACR accessibility

scores for assessing the importance of each ACR to cluster membership by estimating the

likelihood ratio between logistic models with, and without a term for cluster membership.

Specifically, for each cluster, we compared binarized ACR accessibility scores to a reference

panel of uniformly sampled nuclei from each organ (111 nuclei from each organ) where the total

number of reference nuclei was set to the average number of nuclei per cluster (n=666). We then

fit two generalized linear logistic regression models (Eq. 10-11), with and without a term for

membership to the cluster of interest.

1462

1463   10. $logit(p_{ij}) = u_i + \alpha_j + \beta_j + \varepsilon_i$

1464   11. $logit(p_{ij}) = u_i + \beta_j + \varepsilon_i$

1465

Where p_{ij} is the probability that ACR i is accessible in nucleus j, u_i is the proportion of nuclei

where ACR i is accessible, α_j is the cluster membership of nucleus j, β_j is the $\log_{10}$ number of

accessible ACRs in nucleus j and ε_i is the error term for the i^{th} ACR. We then used a likelihood

ratio test to compare the fits of the two models and estimated the false discovery rate (FDR) using

the Benjamini-Hochberg method to identify ACRs that were significantly differentially accessible

across clusters by conditioning on FDR < 5% and fold-change threshold greater than two. ACRs

meeting these criteria with positive Z-scores in nine or fewer clusters (< 10% of clusters) were

considered as cluster-specific. Analysis of differential gene accessibility was performed as

described in the section titled "Cell-type annotation".

1475

GO gene set enrichment analysis

Gene set enrichment using GO biological process terms was performed using the R package

fgsea. For each cluster, test statistics were multiplied by the sign of the $\log_2$ fold-change value

versus the reference panel. GO terms with gene sets less than 10 and greater than 600 were

excluded from the analysis. GO terms were considered significantly enriched at FDR < 0.05

following 10,000 permutations.

1482

Motif analysis

Motif occurrences were identified genome-wide with *fimo* from the MEME suite toolset (Grant et

al., 2011) using position weight matrices (PWM) based on DAP-seq data in *A. thaliana* and *Zea*

mays (Galli et al., 2018; O'Malley et al., 2016). To identity TF motifs associated with cell type-

specific ACRs, we ranked the top 2,000 ACRs in each cell type by Z-scores derived from CPM

normalized accessibility values (see section above "Nuclei clustering"). As a reference for

comparison, we identified 2,000 "constitutive" ACRs that varied the least and were broadly

accessible across clusters. The number of ACRs containing a specific motif was compared to the

frequency of constitutive ACRs harboring the same motif using a binomial test for each cell type

and motif. To control for multiple testing, we used the Benjamini-Hochberg method to estimate

the FDR, considering tests with FDR < 0.05 as significantly different between the focal cell type

and constitutively accessible regions. Maize homologs of *A. thaliana* TFs were identified using

protein fasta alignments from BLAST+ v2.10.0 with an E-value cut-off of 1e-5. Only fasta

sequences classified as transcription factors from either species were considered during

alignment. To narrow the list of putative orthologs based on functional similarity to *A. thaliana*

TFs, we filtered alignments with less than 30% identity, removed maize TFs classified as

belonging to a different family, and selected the homolog with the greatest Pearson correlation

coefficient (PCC) with respect to the motif deviation score. Motif deviation scores of specific TF

motifs among nuclei were estimated using *chromVAR* (Schep et al., 2017) with the non-redundant

core plant PWM database from JASPAR2018. The input matrix for *chromVar* was filtered to retain

a minimum of 50 accessible nuclei per ACR and barcodes with at least 50 accessible ACRs. We

visualized differences in global motif usage per nucleus by projecting deviation scores onto the

UMAP embeddings. To determine if patterns of TF motif accessibility from individual nuclei could

be used to predict cell-type annotations, we constructed a neural network for multinomial

classification using the R package, *caret* (Kuhn, 2008) (method="multinom") using 80% of nuclei

to train, 10-fold cross-validation, averaging error terms across 10 iterations. The nuclei in the 20%

withheld group were used to test the model. Sensitivity, specificity and accuracy of the model was

evaluated using the function confusionMatrix from *caret*.

To identity *de novo* motifs enriched in accessible but non-transcribed genes, we selected

ACRs (n=15,576) within 1-kb of genes that were accessible (ATAC log2 TPM > 1.5) and non-

expressed (mRNA log2 TPM < 1) in at least 10 clusters. We then constructed a set of control

regions by randomly sampling ACRs within 1-kb of genes expressed (mRNA log2 TPM > 1) and

accessible (ATAC log2 TPM > 1.5) in at least 10 clusters (n=15,576). *De novo* motif identification

was conducted using the discriminative motif discovery workflow of MEME-ChIP (v5.1.1) with

default settings (Machanick and Bailey, 2011). Comparison of *de novo* motifs with experimentally

identified motifs was performed using TOMTOM from the MEME-suite toolkit (Gupta et al., 2007).

1519

Analysis of cell type-specific selection signatures

Multi-locus allele-frequency differentiation signals between chronologically sampled elite maize

inbred lines were mapped onto ACRs (Wang et al., 2020), where the selection score for an ACR

was taken as the maximum XP-CLR value within the 500-bp ACR interval. To identify cell types

associated with increased signatures of selection, the top 2,000 ACRs defined by standardized

quantile-scaled CPM chromatin accessibility (Z-scores, see above "Nuclei clustering") were

identified for each cell type. The mean XP-CLR scores per-cell type were standardized by the

mean and standard deviation of randomly sampled ACRs (n=2,000) without replacement across

1,000 permutations, where each permutation estimates the mean XP-CLR scores of a random

subset of 2,000 ACRs from the total list of 165,913 possible ACRs. Enrichment Z-scores were

converted into *P*-values using the R function *pnorm* (log.p=T, lower.tail=F) and used to estimate

FDR via the Benjamini-Hochberg method with the R function *p.adjust* (method="fdr").

1532

Analysis of co-accessible ACRs

To enable comparison with previously identified Hi-C and HiChIP loops (Ricci et al., 2019), we

constrained the distance between co-accessible ACRs to the same range as loops identified in

leaf Hi-C and HiChIP (minimum loop distance = 20-kb). Co-accessible ACRs and Hi-C/HiChIP

loops were considered overlapping if both anchors overlapped by at least 50-bp. We compared

motif composition of co-accessible ACRs by scoring motif occurrence as binary for each ACR and

estimating a Jaccard similarity score on the union of motif sets. Motif similarity scores for co-

accessible ACRs in each cell type were compared to a null distribution by repeating Jaccard

similarity calculations for non-co-accessible ACR-ACR connections (constraining the null

connections to blocks of 1,000 ACR on the same chromosome with the same ACR-ACR distance

distribution as co-accessible ACRs) across 1,000 permutations. To identify motifs enriched at co-

accessible ACR anchors, we first estimated the proportion of co-accessible ACRs with an identical

motif at both anchors for each motif and cell type. Then, we constructed the same number of

random ACR-ACR connections as co-accessible ACRs, again estimating the proportion of links

with an identical motif at both anchors, building a null distribution over 1,000 random

permutations. The estimated proportion of co-accessible ACRs with identical motifs at both

anchors for each motif was transformed to a Z-score by subtracting and scaling by the mean and

standard deviation of the null distribution. Z-scores were converted to *P*-values using the R

function, *qnorm* with non-default parameters (log.p=T, lower.tail=F). FDR values were estimated

using *p.adjust* (method="fdr"). Co-accessible motif scores were plotted as heatmaps using

heatmap.2 by subtracting and dividing observed with expected proportions. Rows and columns

were clustered with *hclust* (method="ward.D2").

1555

Pseudotime analysis

Pseudotime trajectories were constructed similar to previous methods (Granja et al., 2020).

Briefly, nuclei were ordered based on the principal component space by fitting a continuous

trajectory via a smooth spline on the Euclidean distances of each nuclei to a predefined order of

cell types. For feature analysis (ACRs, motifs, and TF activity) across pseudotime, nuclei were

sorted by ascending pseudotime. The ACR x nucleus matrix was filtered to retain differentially

accessible ACRs (see section "Analysis of differential accessibility across pseudotime" below)

with at least one nucleus defined as accessible. For each ACR, we fit a generalized additive model

with the binary accessibility scores as the response and a smoothed pseudotime component as

the dependent variable [s(pseudotime, bs="cs")] with a binomial error term and a logit-link function

with *gam* from the *mgcv* R package. Predicted accessibility scores across pseudotime were

generated from 500 equally spaced interpolated points covering the range of pseudotime values.

Finally, predicted accessibility scores were mean-centered, standardized and constrained to the

range ±1 for each ACR. Model specification for motif deviations and TF gene accessibility analysis

was similar to ACR pseudotime analysis with the exception of a Gaussian error distribution, and

TF gene accessibility was normalized by the row maximum rather than rescaling on a ±1

distribution.

1573

Analysis of differential accessibility across pseudotime

To identity differentially accessible ACRs across pseudotime, we fit normalized accessibility

residuals (Pearson residuals from a generalized linear logistic regression model with $\log_{10}$ number

of accessible ACRs per barcode as the dependent variable, see section "Nuclei Clustering"

above) as the response and pseudotime as the dependent variable using a natural spline with six

degrees of freedom [ns(pseudotime, df=6)] from the R package *splines* for each trajectory. We

took an F-test based approach for hypothesis testing of differential accessibility across

pseudotime by comparing the variance explained by the splined linear model with that of the

residuals normalized by degrees of freedom. *P*-values from the model were used to estimate

Benjamini-Hochberg FDR values with the R function *p.adjust* (method="fdr"), where a FDR

threshold < 0.05 denoted statistical significance for differentially accessible ACRs across

pseudotime. To identify genes and TF motifs with differential accessibility across pseudotime, we

fit the linear splined regression model with the normalized gene accessibility scores and motif

deviations from each nucleus, respectively, similar to the analysis of ACRs.

1588

A. thaliana scATAC-seq processing

scATAC-seq data derived from *A. thaliana* root nuclei were processed similarly to the scATAC-

seq data derived from maize nuclei. Specifically, we processed raw fastq files using *cellranger-*

atac, filtered multi-mapped reads (MQ less than 10 and the presence XA:Z: tags), removed PCR

duplicates by barcode, filtered barcodes by proportion of Tn5 integration sites mapping to

organeller genomes above 1 standard deviation from the mean, and removed barcodes with less

than 1000 unique Tn5 integration sites. We used *in silico* sorting to group nuclei by similarity,

identify ACRs, estimate residuals with regularized quasibinomial regression from the binary ACR

by nuclei matrix, and reduced dimensions with SVD (singular values = 50) similarly as for maize

nuclei. We coded library sequence depth per nucleus as a covariate using the *dplyr* function *ntile*

with n=3 and removed additional technical variance with *Harmony* using the SVD matrix as input

with non-default settings for a weak correction (tau=3, nclust=15, max.iter.harmony=30, theta=0,

lambda=10) (Korsunsky et al., 2019). Nuclei were clustered with Louvain clustering (resolution =

1) in the *Harmony* corrected embedding, and project into an additionally reduced space with

UMAP (n_neighbors=15, min_dist=0.1).

1604

Aligning pseudotime trajectories between A. thaliana and Z. mays

To enable comparison of companion cell development between *A. thaliana* and *Z. mays*, we first

identified putative one-to-one orthologs using *OrthoFinder* (v2) (Emms and Kelly, 2019). Gene

accessibility scores for 10,976 putative orthologs were imputed using a diffusion-based approach

(Fang et al., 2020; van Dijk et al., 2018) and scaled from 0 to 1 across pseudotime for barcodes

associated with companion cell development in *A. thaliana* and *Z. mays*. To account for different

distributions, pseudotime coverage, and number of barcodes between species, we used the R

package, *cellAlign,* that interpolates, scales, and weights gene accessibility scores on a fixed set

of (n=200) equally spaced points (width parameter: winSz=0.1) from two trajectories to remove

technical biases inherent to each data set(Alpert et al., 2018). For each putative ortholog, we

performed global alignment of gene accessibility scores across *A. thaliana* and *Z. mays*

pseudotime using the dynamic time warping algorithm with default settings in *cellAlign.* We then

extracted the pseudotime shifts, representing the extent of gene accessibility deviation at any

given point along the trajectory, for each putative ortholog. We clustered genes into two groups

based on pseudotime shifts across companion cell development using k-means clustering. To

identify conserved gene accessibility patterns across pseudotime, we clustered the normalized

distances between *A. thaliana* and *Z. mays* putative orthologs using a mixture model (G=2) with

the R package, *mclust.* The mixture model identified a bimodal distribution of normalized

distances with 0.15 as a natural cut-off for defining conserved accessibility patterns. Putative

orthologs with normalized distances less than the cut-off were placed in a third group defined as

conserved. The above analysis was repeated with TF motif deviations scores for 440 TF motifs,

without the need for ortholog searching as the same TF position weight matrices were used for

both species, affording identical TF motif labels.

1628

1629   **ADDITIONAL RESOURCES**

1630   Cell-type resolved data can be viewed through our public Plant Epigenome JBrowse Genome

1631   Browser   (Hofmeister   and   Schmitz,   2018)

1632   (http://epigenome.genetics.uga.edu/PlantEpigenome/index.html) by selecting either the *Z. mays*

1633   or *A. thaliana* Genome Browser links, followed by the scATAC_celltypes tab in the tracks panel.

1634

1635

1636

1637

1638

1639

1640

1641

REFERENCES

AFA Smit, R.H., and P Green (2013-2015). RepeatMasker Open-4.0.

Alpert, A., Moore, L.S., Dubovik, T., and Shen-Orr, S.S. (2018). Alignment of single-cell
trajectories to compare cellular expression dynamics. Nat Methods *15*, 267-270.

Andersson, R., and Sandelin, A. (2020). Determinants of enhancer and promoter activities of
regulatory elements. Nat Rev Genet *21*, 71-87.

Baker, R.F., Leach, K.A., Boyer, N.R., Swyers, M.J., Benitez-Alfonso, Y., Skopelitis, T., Luo, A.,
Sylvester, A., Jackson, D., and Braun, D.M. (2016). Sucrose Transporter ZmSut1 Expression and
Localization Uncover New Insights into Sucrose Phloem Loading. Plant Physiol *172*, 1876-1898.

Birnbaum, K., Shasha, D.E., Wang, J.Y., Jung, J.W., Lambert, G.M., Galbraith, D.W., and Benfey,
P.N. (2003). A gene expression map of the Arabidopsis root. Science *302*, 1956-1960.

Brady, S.M., Orlando, D.A., Lee, J.Y., Wang, J.Y., Koch, J., Dinneny, J.R., Mace, D., Ohler, U.,
and Benfey, P.N. (2007). A high-resolution root spatiotemporal map reveals dominant expression
patterns. Science *318*, 801-806.

Buchler, N.E., Gerland, U., and Hwa, T. (2003). On schemes of combinatorial transcription logic.
Proc Natl Acad Sci U S A *100*, 5136-5141.

Buenrostro, J.D., Wu, B., Litzenburger, U.M., Ruff, D., Gonzales, M.L., Snyder, M.P., Chang, H.Y.,
and Greenleaf, W.J. (2015). Single-cell chromatin accessibility reveals principles of regulatory
variation. Nature *523*, 486-490.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden,
T.L. (2009). BLAST+: architecture and applications. BMC Bioinformatics *10*, 421.

Chang, Y.M., Liu, W.Y., Shih, A.C., Shen, M.N., Lu, C.H., Lu, M.Y., Yang, H.W., Wang, T.Y.,
Chen, S.C., Chen, S.M.*, et al.* (2012). Characterizing regulatory and functional differentiation
between maize mesophyll and bundle sheath cells by transcriptomic analysis. Plant Physiol *160*,
165-177.

Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics *34*, i884-i890.

Cheng, C., Alexander, R., Min, R., Leng, J., Yip, K.Y., Rozowsky, J., Yan, K.K., Dong, X., Djebali,
S., Ruan, Y.*, et al.* (2012). Understanding transcriptional regulation by integrative analysis of
transcription factor binding data. Genome Res *22*, 1658-1667.

Chuck, G.S., Brown, P.J., Meeley, R., and Hake, S. (2014). Maize SBP-box transcription factors
unbranched2 and unbranched3 affect yield traits by regulating the rate of lateral primordia
initiation. Proc Natl Acad Sci U S A *111*, 18775-18780.

Clark, R.M., Wagler, T.N., Quijada, P., and Doebley, J. (2006). A distant upstream enhancer at
the maize domestication gene tb1 has pleiotropic effects on plant and inflorescent architecture.
Nat Genet *38*, 594-597.

Consortium, E.P. (2012). An integrated encyclopedia of DNA elements in the human genome.
Nature *489*, 57-74.

Corces, M.R., Granja, J.M., Shams, S., Louie, B.H., Seoane, J.A., Zhou, W., Silva, T.C.,
Groeneveld, C., Wong, C.K., Cho, S.W.*, et al.* (2018). The chromatin accessibility landscape of
primary human cancers. Science *362*.

Crisp, P.A., Marand, A.P., Noshay, J.M., Zhou, P., Lu, Z., Schmitz, R.J., and Springer, N.M.
(2020). Stable unmethylated DNA demarcates expressed genes and their cis-regulatory space in
plant genomes. Proc Natl Acad Sci U S A.

Cusanovich, D.A., Hill, A.J., Aghamirzaie, D., Daza, R.M., Pliner, H.A., Berletch, J.B., Filippova,
G.N., Huang, X., Christiansen, L., DeWitt, W.S.*, et al.* (2018). A Single-Cell Atlas of In Vivo
Mammalian Chromatin Accessibility. Cell *174*, 1309-1324 e1318.

Deal, R.B., and Henikoff, S. (2011). The INTACT method for cell type-specific gene expression
and chromatin profiling in Arabidopsis thaliana. Nat Protoc *6*, 56-68.

Dorrity, M.W., Alexandre, C., Hamm, M., Vigil, A.-L., Fields, S., Queitsch, C., and Cuperus, J.
(2020). The regulatory landscape of Arabidopsis thaliana roots at single-cell resolution. bioRxiv,
2020.2007.2017.204792.

Emms, D.M., and Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol *20*, 238.

Fang, R., Preissl, S., Li, Y., Hou, X., Lucero, J., Wang, X., Motamedi, A., Shiau, A.K., Zhou, X.,
Xie, F.*, et al.* (2020). SnapATAC: A Comprehensive Analysis Package for Single Cell ATAC-seq.
bioRxiv, 615179.

Farmer, A., Thibivilliers, S., Ryu, K.H., Schiefelbein, J., and Libault, M. (2020). The impact of
chromatin remodeling on gene expression at the single cell level in Arabidopsis
thaliana. bioRxiv, 2020.2007.2027.223156.

Gage, J.L., White, M.R., Edwards, J.W., Kaeppler, S., and de Leon, N. (2018). Selection
Signatures Underlying Dramatic Male Inflorescence Transformation During Modern Hybrid Maize
Breeding. Genetics *210*, 1125-1138.

Galli, M., Khakhar, A., Lu, Z., Chen, Z., Sen, S., Joshi, T., Nemhauser, J.L., Schmitz, R.J., and
Gallavotti, A. (2018). The DNA binding landscape of the maize AUXIN RESPONSE FACTOR
family. Nat Commun *9*, 4526.

Garrison, E., and Marth, G. (2012). Haplotype-based variant detection from short-read
sequencing. In arXiv e-prints, pp. arXiv:1207.3907.

Gate, R.E., Cheng, C.S., Aiden, A.P., Siba, A., Tabaka, M., Lituiev, D., Machol, I., Gordon, M.G.,
Subramaniam, M., Shamim, M.*, et al.* (2018). Genetic determinants of co-accessible chromatin
regions in activated T cells across humans. Nat Genet *50*, 1140-1150.

Gerstein, M.B., Kundaje, A., Hariharan, M., Landt, S.G., Yan, K.K., Cheng, C., Mu, X.J., Khurana,
E., Rozowsky, J., Alexander, R.*, et al.* (2012). Architecture of the human regulatory network
derived from ENCODE data. Nature *489*, 91-100.

Gomez-Mena, C., de Folter, S., Costa, M.M., Angenent, G.C., and Sablowski, R. (2005).
Transcriptional program controlled by the floral homeotic gene AGAMOUS during early
organogenesis. Development *132*, 429-438.

Granja, J.M., Corces, M.R., Pierce, S.E., Bagdatli, S.T., Choudhry, H., Chang, H.Y., and
Greenleaf, W.J. (2020). ArchR: An integrative and scalable software package for single-cell
chromatin accessibility analysis. bioRxiv, 2020.2004.2028.066498.

Grant, C.E., Bailey, T.L., and Noble, W.S. (2011). FIMO: scanning for occurrences of a given
motif. Bioinformatics *27*, 1017-1018.

Gupta, S., Stamatoyannopoulos, J.A., Bailey, T.L., and Noble, W.S. (2007). Quantifying similarity
between motifs. Genome Biol *8*, R24.

Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of single-cell
RNA-seq data using regularized negative binomial regression. Genome Biol *20*.

Heger, P., Marin, B., Bartkuhn, M., Schierenberg, E., and Wiehe, T. (2012). The chromatin
insulator CTCF and the emergence of metazoan diversity. Proc Natl Acad Sci U S A *109*, 17507-
17512.

Hekselman, I., and Yeger-Lotem, E. (2020). Mechanisms of tissue and cell-type specificity in
heritable traits and diseases. Nat Rev Genet *21*, 137-150.

Hofmeister, B.T., and Schmitz, R.J. (2018). Enhanced JBrowse plugins for epigenomics data
visualization. BMC Bioinformatics *19*, 159.

Jean-Baptiste, K., McFaline-Figueroa, J.L., Alexandre, C.M., Dorrity, M.W., Saunders, L., Bubb,
K.L., Trapnell, C., Fields, S., Queitsch, C., and Cuperus, J.T. (2019). Dynamics of Gene
Expression in Single Root Cells of Arabidopsis thaliana. Plant Cell *31*, 993-1011.

Jiang, H., Li, H., Bu, Q., and Li, C. (2009). The RHA2a-interacting proteins ANAC019 and
ANAC055 may play a dual role in regulating ABA response and jasmonate response. Plant
Signaling & Behavior *4*, 464-466.

Jiang, K., Zhu, T., Diao, Z., Huang, H., and Feldman, L.J. (2010). The maize root stem cell niche:
a partnership between two sister cell populations. Planta *231*, 411-424.

Jiao, Y., Peluso, P., Shi, J., Liang, T., Stitzer, M.C., Wang, B., Campbell, M.S., Stein, J.C., Wei,
X., Chin, C.S.*, et al.* (2017). Improved maize reference genome with single-molecule
technologies. Nature *546*, 524-527.

Kajala, K., Shaar-Moshe, L., Mason, G.A., Gouran, M., Rodriguez-Medina, J., Kawa, D., Pauluzzi,
G., Reynoso, M., Canto-Pastor, A., Lau, V.*, et al.* (2020). Innovation, conservation and
repurposing of gene function in plant root cell type development. bioRxiv,
2020.2004.2009.017285.

Karaaslan, E.S., Wang, N., Faiss, N., Liang, Y., Montgomery, S.A., Laubinger, S., Berendzen,
K.W., Berger, F., Breuninger, H., and Liu, C. (2020). Marchantia TCP transcription factor activity
correlates with three-dimensional chromatin structure. Nat Plants.

Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner,
M., Loh, P.R., and Raychaudhuri, S. (2019). Fast, sensitive and accurate integration of single-cell
data with Harmony. Nat Methods *16*, 1289-1296.

Kuhn, M. (2008). Building Predictive Models in R Using the caret Package. 2008 *28*, 26.

Langdale, J.A., Lane, B., Freeling, M., and Nelson, T. (1989). Cell lineage analysis of maize
bundle sheath and mesophyll cells. Dev Biol *133*, 128-139.

Lee, L.R., Wengier, D.L., and Bergmann, D.C. (2019). Cell-type-specific transcriptome and
histone modification dynamics during cellular reprogramming in the Arabidopsis stomatal lineage.
Proc Natl Acad Sci U S A *116*, 21914-21924.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
In arXiv e-prints, pp. arXiv:1303.3997.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., and Genome Project Data Processing, S. (2009). The Sequence Alignment/Map
format and SAMtools. Bioinformatics *25*, 2078-2079.

Liu, C., Cheng, Y.J., Wang, J.W., and Weigel, D. (2017). Prominent topologically associated
domains differentiate global chromatin packing in rice from Arabidopsis. Nat Plants *3*, 742-748.

Long, H.K., Prescott, S.L., and Wysocka, J. (2016). Ever-Changing Landscapes: Transcriptional
Enhancers in Development and Evolution. Cell *167*, 1170-1187.

Lopez-Anido, C.B., Vatén, A., Smoot, N.K., Sharma, N., Guo, V., Gong, Y., Anleu Gil, M.X.,
Weimer, A.K., and Bergmann, D.C. (2020). Single-Cell Resolution of Lineage Trajectories in the
Arabidopsis Stomatal Lineage and Developing Leaf. bioRxiv, 2020.2009.2008.288498.

Lu, Z., Marand, A.P., Ricci, W.A., Ethridge, C.L., Zhang, X., and Schmitz, R.J. (2019). The
prevalence, evolution and chromatin signatures of plant regulatory elements. Nat Plants *5*, 1250-
1259.

Machanick, P., and Bailey, T.L. (2011). MEME-ChIP: motif analysis of large DNA datasets.
Bioinformatics *27*, 1696-1697.

Marand, A.P., Zhang, T., Zhu, B., and Jiang, J. (2017). Towards genome-wide prediction and
characterization of enhancers in plants. Biochim Biophys Acta Gene Regul Mech *1860*, 131-139.

Miyashima, S., Roszak, P., Sevilem, I., Toyokura, K., Blob, B., Heo, J.O., Mellor, N., Help-Rinta-
Rahko, H., Otero, S., Smet, W.*, et al.* (2019). Mobile PEAR transcription factors integrate
positional cues to prime cambial growth. Nature *565*, 490-494.

Nag, A., King, S., and Jack, T. (2009). miR319a targeting of TCP4 is critical for petal growth and
development in Arabidopsis. Proc Natl Acad Sci U S A *106*, 22534-22539.

Nelms, B., and Walbot, V. (2019). Defining the developmental program leading to meiosis in
maize. Science *364*, 52-56.

Noshay, J.M., Marand, A.P., Anderson, S.N., Zhou, P., Guerra, M.K.M., Lu, Z., O'Connor, C.,
Crisp, P.A., Hirsch, C.N., Schmitz, R.J.*, et al.* (2020). Cis-regulatory elements within TEs can
influence expression of nearby maize genes. bioRxiv, 2020.2005.2020.107169.

O'Malley, R.C., Huang, S.C., Song, L., Lewsey, M.G., Bartlett, A., Nery, J.R., Galli, M., Gallavotti,
A., and Ecker, J.R. (2016). Cistrome and Epicistrome Features Shape the Regulatory DNA
Landscape. Cell *166*, 1598.

Oka, R., Bliek, M., Hoefsloot, H.C.J., and Stam, M. (2020). In plants distal regulatory sequences
overlap with unmethylated rather than low-methylated regions, in contrast to mammals. bioRxiv,
2020.2003.2024.005678.

Oka, R., Zicola, J., Weber, B., Anderson, S.N., Hodgman, C., Gent, J.I., Wesselink, J.J., Springer,
N.M., Hoefsloot, H.C.J., Turck, F.*, et al.* (2017). Genome-wide mapping of transcriptional
enhancer candidates using DNA and chromatin features in maize. Genome Biol *18*, 137.

Peng, Y., Xiong, D., Zhao, L., Ouyang, W., Wang, S., Sun, J., Zhang, Q., Guan, P., Xie, L., Li,
W.*, et al.* (2019). Chromatin interaction maps reveal genetic regulation for quantitative traits in
maize. Nat Commun *10*, 2632.

Phillips, J.E., and Corces, V.G. (2009). CTCF: master weaver of the genome. Cell *137*, 1194-
1211.

Pliner, H.A., Packer, J.S., McFaline-Figueroa, J.L., Cusanovich, D.A., Daza, R.M., Aghamirzaie,
D., Srivatsan, S., Qiu, X.J., Jackson, D., Minkina, A.*, et al.* (2018). Cicero Predicts cis-Regulatory
DNA Interactions from Single-Cell Chromatin Accessibility Data. Mol Cell *71*, 858-+.

Ravasi, T., Suzuki, H., Cannistraci, C.V., Katayama, S., Bajic, V.B., Tan, K., Akalin, A., Schmeier,
S., Kanamori-Katayama, M., Bertin, N.*, et al.* (2010). An atlas of combinatorial transcriptional
regulation in mouse and man. Cell *140*, 744-752.

Rebeiz, M., and Tsiantis, M. (2017). Enhancer evolution and the origins of morphological novelty.
Curr Opin Genet Dev *45*, 115-123.

Ricci, W.A., Lu, Z., Ji, L., Marand, A.P., Ethridge, C.L., Murphy, N.G., Noshay, J.M., Galli, M.,
Mejia-Guerra, M.K., Colome-Tatche, M.*, et al.* (2019). Widespread long-range cis-regulatory
elements in the maize genome. Nat Plants *5*, 1237-1249.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics *26*, 139-140.

Ryu, K.H., Huang, L., Kang, H.M., and Schiefelbein, J. (2019). Single-Cell RNA Sequencing
Resolves Molecular Relationships Among Individual Plant Cells. Plant Physiol *179*, 1444-1456.

Salvi, S., Sponza, G., Morgante, M., Tomes, D., Niu, X., Fengler, K.A., Meeley, R., Ananiev, E.V.,
Svitashev, S., Bruggemann, E.*, et al.* (2007). Conserved noncoding genomic sequences
associated with a flowering-time quantitative trait locus in maize. Proc Natl Acad Sci U S A *104*,
11376-11381.

Satpathy, A.T., Granja, J.M., Yost, K.E., Qi, Y., Meschi, F., McDermott, G.P., Olsen, B.N.,
Mumbach, M.R., Pierce, S.E., Corces, M.R.*, et al.* (2019). Massively parallel single-cell chromatin

landscapes of human immune cell development and intratumoral T cell exhaustion. Nat
Biotechnol *37*, 925-936.

Savaldi-Goldstein, S., Peto, C., and Chory, J. (2007). The epidermis both drives and restricts
plant shoot growth. Nature *446*, 199-202.

Schep, A.N., Wu, B., Buenrostro, J.D., and Greenleaf, W.J. (2017). chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. Nat Methods *14*,
975-978.

Shulse, C.N., Cole, B.J., Ciobanu, D., Lin, J., Yoshinaga, Y., Gouran, M., Turco, G.M., Zhu, Y.,
O'Malley, R.C., Brady, S.M.*, et al.* (2019). High-Throughput Single-Cell Transcriptome Profiling of
Plant Cell Types. Cell Rep *27*, 2241-2247 e2244.

Somssich, M., Je, B.I., Simon, R., and Jackson, D. (2016). CLAVATA-WUSCHEL signaling in the
shoot meristem. Development *143*, 3238-3248.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y.H.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data.
Cell *177*, 1888-+.

Sun, Y., Dong, L., Zhang, Y., Lin, D., Xu, W., Ke, C., Han, L., Deng, L., Li, G., Jackson, D.*, et al.*
(2020). 3D genome architecture coordinates trans and cis regulation of differentially expressed
ear and tassel genes in maize. Genome Biol *21*, 143.

Takacs, E.M., Li, J., Du, C., Ponnala, L., Janick-Buckner, D., Yu, J., Muehlbauer, G.J., Schnable,
P.S., Timmermans, M.C., Sun, Q.*, et al.* (2012). Ontogeny of the maize shoot apical meristem.
Plant Cell *24*, 3219-3234.

Tatematsu, K., Nakabayashi, K., Kamiya, Y., and Nambara, E. (2008). Transcription factor
AtTCP14 regulates embryonic growth potential during seed germination in Arabidopsis thaliana.
Plant J *53*, 42-52.

Teale, W.D., Paponov, I.A., and Palme, K. (2006). Auxin in action: signalling, transport and the
control of plant growth and development. Nat Rev Mol Cell Biol *7*, 847-859.

Team, R.C. (2013). R: A language and environment for statistical computing. (R Foundation for
Statistical Computing, Vienna, Austria.).

Ten Hove, C.A., and Heidstra, R. (2008). Who begets whom? Plant cell fate determination by
asymmetric cell division. Curr Opin Plant Biol *11*, 34-41.

Tigchelaar, M., Battisti, D.S., Naylor, R.L., and Ray, D.K. (2018). Future warming increases
probability of globally synchronized maize production shocks. Proc Natl Acad Sci U S A *115*,
6644-6649.

van Dijk, D., Sharma, R., Nainys, J., Yim, K., Kathail, P., Carr, A.J., Burdziak, C., Moon, K.R.,
Chaffer, C.L., Pattabiraman, D.*, et al.* (2018). Recovering Gene Interactions from Single-Cell Data
Using Data Diffusion. Cell *174*, 716-729 e727.

Verweij, W., Spelt, C.E., Bliek, M., de Vries, M., Wit, N., Faraco, M., Koes, R., and Quattrocchio,
F.M. (2016). Functionally Similar WRKY Proteins Regulate Vacuolar Acidification in Petunia and
Hair Development in Arabidopsis. Plant Cell *28*, 786-803.

Villar, D., Berthelot, C., Aldridge, S., Rayner, T.F., Lukk, M., Pignatelli, M., Park, T.J., Deaville,
R., Erichsen, J.T., Jasinska, A.J.*, et al.* (2015). Enhancer evolution across 20 mammalian species.
Cell *160*, 554-566.

Wallace, J.G., Bradbury, P.J., Zhang, N., Gibon, Y., Stitt, M., and Buckler, E.S. (2014).
Association mapping across numerous traits reveals patterns of functional variation in maize.
PLoS Genet *10*, e1004845.

Wallner, E.S., Lopez-Salmeron, V., Belevich, I., Poschet, G., Jung, I., Grunwald, K., Sevilem, I.,
Jokitalo, E., Hell, R., Helariutta, Y.*, et al.* (2017). Strigolactone- and Karrikin-Independent SMXL
Proteins Are Central Regulators of Phloem Formation. Curr Biol *27*, 1241-1247.

Wang, B., Lin, Z., Li, X., Zhao, Y., Zhao, B., Wu, G., Ma, X., Wang, H., Xie, Y., Li, Q.*, et al.* (2020).
Genome-wide selection and genetic improvement during modern maize breeding. Nat Genet *52*,
565-571.

Welch, J.D., Kozareva, V., Ferreira, A., Vanderburg, C., Martin, C., and Macosko, E.Z. (2019).
Single-Cell Multi-omic Integration Compares and Contrasts Features of Brain Cell Identity. Cell
177, 1873-1887 e1817.

Witten, D.M., Tibshirani, R., and Hastie, T. (2009). A penalized matrix decomposition, with
applications to sparse principal components and canonical correlation analysis. Biostatistics *10*,
515-534.

Wittkopp, P.J., and Kalay, G. (2011). Cis-regulatory elements: molecular mechanisms and
evolutionary processes underlying divergence. Nat Rev Genet *13*, 59-69.

Xiao, J., Jin, R., Yu, X., Shen, M., Wagner, J.D., Pai, A., Song, C., Zhuang, M., Klasfeld, S., He,
C.*, et al.* (2017). Cis and trans determinants of epigenetic silencing by Polycomb repressive
complex 2 in Arabidopsis. Nat Genet *49*, 1546-1552.

Yamaguchi, M., Ohtani, M., Mitsuda, N., Kubo, M., Ohme-Takagi, M., Fukuda, H., and Demura,
T. (2010). VND-INTERACTING2, a NAC domain transcription factor, negatively regulates xylem
vessel formation in Arabidopsis. Plant Cell *22*, 1249-1263.

You, Y., Sawikowska, A., Lee, J.E., Benstein, R.M., Neumann, M., Krajewski, P., and Schmid, M.
(2019). Phloem Companion Cell-Specific Transcriptomic and Epigenomic Analyses Identify
MRF1, a Regulator of Flowering. Plant Cell *31*, 325-345.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C.,
Myers, R.M., Brown, M., Li, W.*, et al.* (2008). Model-based analysis of ChIP-Seq (MACS).
Genome Biol *9*, R137.

Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D.B., Huang, Y., Huang, M., Yao, Y., Bassu, S.,
Ciais, P.*, et al.* (2017). Temperature increase reduces global yields of major crops in four
independent estimates. Proc Natl Acad Sci U S A *114*, 9326-9331.

Zhao, H., Zhang, W., Chen, L., Wang, L., Marand, A.P., Wu, Y., and Jiang, J. (2018). Proliferation
of Regulatory DNA Elements Derived from Transposable Elements in the Maize Genome. Plant
Physiol *176*, 2789-2803.

Zheng, L., McMullen, M.D., Bauer, E., Schon, C.C., Gierl, A., and Frey, M. (2015). Prolonged
expression of the BX1 signature enzyme is associated with a recombination hotspot in the
benzoxazinoid gene cluster in Zea mays. J Exp Bot *66*, 3917-3930.
1906