IL-1R1 dependent signals improve clearance of cytosolic virulent mycobacteria

in vivo.

Running title: In vivo Cytosolic mycobacteria are cleared by IL-1R1

Authors / Affiliations

Sanne van der Niet[1], Maaike van Zon[2], Karin de Punder[2], Anita Grootemaat[1], Sofie Rutten[1], Simone J.C.F.M. Moorlag[2], Diane Houben[2], Astrid van der Sar[3], Wilbert Bitter[3], Roland Brosch[4], Rogelio Hernandez Pando[5], Maria T. Pena[6], Peter J. Peters[2], Eric A. Reits[1], Katrin D. Mayer-Barber[7] and Nicole N. van der Wel[1,2]*

Contact information

[1] Electron Microscopy Centre Amsterdam, Amsterdam University Medical Centre AMC, the Netherlands, [2] Netherlands Cancer Institute, The Netherlands, [3] Amsterdam University Medical Centre VUMC, The Netherlands, [4] Unit for Integrated Mycobacterial Pathogenomics, CNRS UMR 3525, Paris, France, [5] National Institute of Medical Sciences and Nutrition, Mexico, [6] Department of Health and Human Services, Health Resources and Services Administration, Healthcare Systems Bureau, National Hansen's Disease Programs, Baton Rouge, LA, USA, [7] Inflammation and Innate Immunity Unit, Laboratory of Clinical Immunology and Microbiology, National Institute of Allergy and Infectious Diseases (NIAID), National Institutes of Health (NIH), Bethesda, USA.

*Corresponding author: Nicole van der Wel

email address: n.n.vanderwel@amsterdamumc.nl

Abstract

Mycobacterium tuberculosis infections claim more than a million lives each year and better treatments or vaccines are required. A crucial pathogenicity factor is translocation from the phago-lysosomes to the cytosol upon phagocytosis by macrophages. The translocation from the phago-lysosome into the cytosol is an ESX-1 dependent process as previously shown *in vitro*. Here we show that *in vivo,* mycobacteria also translocate to the cytosol but mainly when host immunity is compromised. We observed only low numbers of cytosolic bacilli in mice, armadillo, zebrafish and patient material infected with *M. tuberculosis, M. marinum* or *M. leprae.* In contrast, when innate or adaptive immunity was compromised, as in SCID or IL-1R1 deficient mice, a significant number of cytosolic *M. tuberculosis* bacilli were detected in lungs of infected mice. Taken together, *M. tuberculosis* infection is controlled by adaptive immune responses as well as IL-1R1-mediated signals that result in clearance of cells containing cytosolic mycobacteria *in vivo.*

Importance

For decades, *Mycobacterium tuberculosis* is one of the deathliest pathogens known. Despite infecting approximately one third of the human population, no effective treatment or vaccine is available. A crucial pathogenicity factor is the subcellular localization, as *M. tuberculosis* can translocate from the phago-lysosome to the cytosol in macrophages. The situation *in vivo* is more complicated. In this study we establish that high level cytosolic escape of mycobacteria can indeed occur *in vivo*, but mainly when host resistance is compromised. The IL-1 pathway is crucial for the control of the number of cytosolic mycobacteria. The establishment that immune signals result in clearance of cells containing cytosolic mycobacteria, connects two important fields:

46 cell-biology and immunology which is vital for the understanding of the pathology of

47 *M. tuberculosis*.

49 **Keywords:** *Mycobacterium tuberculosis*, *Mycobacterium leprae*, *Mycobacterium*

50 *marinum*, cytosolic localization, IL-1 receptor 1, phagosome, lysosome, phago-

51 lysosomal fusion.

Introduction

Mycobacterium tuberculosis (*Mtb*) is not only one of the most deadliest pathogens in history, but it continues to claim an estimated 1.5 million human lives per year (1) and is a big threat for the future as multidrug resistant strains are arising with no effective treatment nor vaccine available. The treatment success rate is just 56% and in approximately 6.2% of the cases, infection is caused by extensive drug resistant *Mtb* (1). Furthermore, *Mtb* is the leading cause of death among HIV infected patients. In patients with AIDS, tuberculosis can thrive because their immune system is impaired by CD4$^+$T cell loss, which secondarily affects many other immune compartments (reviewed in (2)).

For an effective response to *Mtb* infections, both the innate and the adaptive immune system are important. The development of an active pulmonary *Mtb* infection is related to a disordered immune balance, which results in the inability of the host to keep the infection under control (3). The first immune response is the innate response (reviewed in (4)) including a series of cells that come in contact with *Mtb* such as alveolar macrophages. Alveolar macrophages provide a nutritionally permissive niche (5) and are critical for dissemination of the bacteria in the lung, spreading the infection from the alveoli to the interstitium (6). Here *Mtb* infects other cell types like neutrophils, monocyte derived macrophages and dendritic cells (DC). Since DCs present antigens via MHC class I and II to T cells, they function as a connection between the innate and adaptive immune system (7, 8). Upon antigen presentation to T cells, CD4$^+$T cells produce IFNγ, which is involved in the enhancement of macrophage killing and plays an important role in granuloma formation (9). Indeed in mice (10) and humans (11) loss of IFNγ or its receptors, acting as single non-redundant factor, leads to TB disease. In order to spread to new individuals, *Mtb* needs to cause pulmonary lesions

(12). Inflammation driven by IL-1 contributes to host resistance to *Mtb* (13, 14). Mice

that lack the IL-1R1, IL-1α or IL-1β display high susceptibility to *Mtb* infection (15) with

uncontrolled bacterial replication in the lungs, again demonstrating a non-redundant

role of one key host pathway.

We hypothesized that bacterial translocation from the phagosome to the cytosol might

also be regulated by IL-1. *In vitro, Mtb* can translocate from the phago-lysosome to the

cytosol (16–23) in an ESX-1 dependent manner (16, 21, 22). This system is

responsible for the secretion of a number of proteins, including EsxA (ESAT-6) and

EsxB (CFP-10) (reviewed in (24)). When this secretion system is not present, as is the

case for *M. bovis* BCG, cytosolic localization is abrogated in *in vitro* macrophage

systems, rendering the bacteria restricted in a membrane enclosed phago-lysosome.

Reintroducing the extended *esx-1* locus in BCG allowed the translocation to the cytosol

and increased virulence providing clear evidence for an essential role of this Type VII

secretion system in escape (16, 25–28). In addition, it is shown that virulent *Mtb* can

form cords in the cytosol and not in the phagosome in human lymphatic endothelial

cells *in vitro* (23). This cording is dependent on the ESX-1 secretion system and PDIM

glycolipids. The formation of cords in the cytosol rather than in phagosomes suggests

a permissive environment for bacterial replication in the cytosol. When *Mtb* is present

in the cytosol its bacterial DNA is sensed by cyclic guanosine monophosphate-AMP

synthase (cGAS) (29–32). This detection is dependent on the presence of a functioning

ESX-1 system, suggesting that it is dependent on pathogen-induced cytosolic

localization. Cytosolic bacteria co-localize with cytosolic ubiquitin, while this is not the

case when the mycobacteria are present in the phagosome (16). Another factor

involved in the cytosolic localization is Rv3167c, which regulates the escape of *Mtb*

from the phagosome, since a mutant unable to produce this protein (MtbΔRv3167c)

displayed increased cytosolic escape (33). Other virulent factors involved in escape from the phagosome are PDIM glycolipids located on the outer membrane of *Mtb* (19, 25, 34). Mycobacterial strains that lack PDIM are less capable of damaging the phagosomal membrane, resulting in less *Mtb* in the cytosol of THP-1 macrophages. Recently it was shown that phagosomal rupture causes activation of NLRP3-dependent IL-1β release and pyroptosis, a programmed form of cell death (35) facilitating the spread of bacteria to other cells.

While most studies focused on *in vitro* experiments using cultured macrophages, the sub-cellular localization of *Mtb* and the factors affecting cytosolic localization and pathogenesis are less intensively studied *in vivo*. When macrophages purified from bronchoalveolar lavages from TB infected patients were analyzed by electron microscopy, it was found that *Mtb* is primarily localized in phagosome-like compartments (36, 37). The phagosomal localization does not affect the ability of *Mtb* to proliferate, and it is long known that when *Mtb* is located in phagosomes it is able to arrest its maturation (38, 39) up to 5-7 days (40). More recently, it was shown *in vivo* that *Mtb* is able to translocate to the cytosol as early as three hours post infection using a Förster resonance energy transfer (FRET)-based detection system (41). This study also demonstrated that the pH of the lysosomes plays a role in the cytosolic localization; when the lysosome is more acidic, less *Mtb* is present in the cytosol at 3 days post infection (41).

To examine whether activation via adaptive or innate immunity pathways would affect the capability of mycobacteria to translocate into the cytoplasm *in vivo,* we tested the subcellular localization of different mycobacterial species in zebrafish, armadillo and mice models as well as in patient material. In adult zebrafish and zebrafish embryos we used *M. marinum*, a close homologue of *Mtb* that is also known to escape the

phago-lysosome in an ESX-1 dependent manner (42). *M. leprae* is also known to

escape to the cytosol (22), probably using a similar mechanism, although some of the

members of ESX-1 system (like esxC, esxG, esxS) are pseudogenes, it has functional

ESAT-6 (esxA) and CFP-10 (esxB), the two most important components needed for

cytosolic escape. We used both skin biopsies of leprosy patients and the armadillo

model for *M. leprae* since, the armadillo model is known to exhibit the entire clinical

spectrum of leprosy (43). For *Mtb* both SCID mice that lack both T and B cells, and IL-

1R1 knockout mice were compared to determine their ability to limit cytosolic escape

in vivo in infected cells. Here we show that while cytosolic localization is limited by both

innate and adaptive immunity, IL-1 seems to be a key effector pathway in controlling

cytosolic translocation of *Mtb*.

Results

The pH of the phagosome and lysosome does not affect cytosolic localization of *M. marinum*.

Mtb blocks maturation and acidification of the phago-lysosome, promoting its intracellular survival (38, 39, 44–46). Phagosomal acidification is essential for increased activity of the lysosomal digestive process and thus for degradation of its content (47). *Mtb* partially avoids acid mediated killing by blocking fusion between lysosomes and phagosomes (36, 39) and the secretion of antacid known as 1-tuberculosinyladenosine (TbAd) (48). In addition, it is shown that the mycobacterial cell wall plays a role in the resistance to acidic environments (reviewed in (49)). We hypothesized that cytosolic escape is a fourth mechanism to avoid lysosome mediated killing. To determine whether these mechanisms are interdependent, we examine if the phagosomal pH affects translocation from the phagosome to the cytosol. To exclude the effect of TbAd, we utilized *M. marinum*, which does not express TbAd (50) in THP-1 cells. The acidity of the phagosome and the lysosome was measured using lysotracker and by incubation with (N-(3-((2,4-dinitrophenyl)amino)propyl)-N-(3-aminopropyl) methylamine (DAMP), a weak basic amine that will be taken up by acidic organelles in live cells (36). After fixation and sample preparation, the DAMP was visualized by Transmission Electron Microscopy (TEM) using immuno-gold labelling with αDNP antibody conjugated to a gold particle. The more acidic the phagosome or lysosome, the more DAMP was present, thus resulting in a higher label density in acidic organelles. As expected, upon *M. marinum* infection of THP-1 cells, low amounts of DAMP labelling were observed surrounding cytosolic *M. marinum* (Fig. S1A/1A') while more labeling was detected in *M. marinum* containing phagosomes (Fig. S1A/1A"). We

next blocked acidification with 10nM Concanamycin B (ConB), an inhibitor of vacuolar

ATPases which prevents acidification of endosomes and lysosomes. When cells were

treated with ConB and infected with *M. marinum,* both a lower label density was

measured and the lysotracker imaged with fluorescence microscopy confirm that the

pH is less acidic in the phagosome/lysosome when treated with ConB as already well

described (36, 51) (Fig. S1B). We next examined whether a raising pH would affect

the translocation efficiency of *M. marinum* to the cytosol. The percentage of cytosolic

bacteria in THP-1 cells treated with ConB or no treatment control was determined both

at 24h and 48h after infection with *M. marinum,* which is the known timeframe for

escape (16) (Fig. 1A,B). The number of bacteria present in CD63 labeled

compartments (phago-lysosomal), membrane enclosed but not CD63 positive

compartments (phagosomes) and the number of bacteria in the cytosol were counted

(Fig. S2A). At both 24h and 48h after infection no difference in the percentage cytosolic

bacteria was detected for untreated or ConB treated cells. This indicates that a higher

pH has no effect on *M. marinum* translocation to the cytosol. In conclusion, a raised

lysosomal pH has no effect on the percentage of bacteria translocating to the cytosol

in THP-1 cells.

***M. marinum* translocates to the cytosol in embryonic but not adult zebrafish.**

After studying the effect of the pH on the ability of *M. marinum to* translocate to the

cytoplasm *in vitro*, we next focussed on the subcellular localisation *in vivo*. To address

whether either adaptive or innate immunity was required to limit bacterial cytosolic

escape we used a zebrafish model combined with *M. marinum* infection. Among other

age-related differences, zebrafish larvae do not have a fully developed adaptive

immune system in contrast to adult zebrafish (52). Zebrafish embryos and adult

zebrafish were infected with *M. marinum* or *M. marinum* Tn::ESX5 mutant and tissue

was fixed for TEM analysis. For analysis of adult zebrafish a specific *M. marinum*

Tn::ESX5 mutant was used as this infection was shown to cause hypervirulence (53)

and thus large amounts of bacteria can be detected *in vivo* using TEM. The following

conditions were analysed using TEM: whole zebrafish embryo day 9 and the spleen of

adult zebra fish at day 11 (Fig. 2A,B). In zebrafish embryos, we have previously

demonstrated that injected *M. marinum* with fluorescent tags is present in phagocytic

cells in close proximity of blood vessels and endothelial cells of these blood vessels

(54). Here we injected the zebrafish with untagged *M. marinum* and detected bacteria

in similar phagocytic and endothelial cells, close to or part of the blood vessels and in

phagocytic cells spread through the tissue. In these cells, the percentage of cytosolic

M. marinum was determined by counting the number of mycobacteria in a membrane

enclosed compartment and in the cytoplasm. Discrimination between phagosome and

phago-lysome based on immuno-gold labelling with lysosomal markers available in

humans or mice (CD63, LAMP1) was not possible as no lysosomal nor cellular markers

are available for immuno-gold labelling in zebrafish. Alternatively, actin antibodies were

used as a cellular cytoskeleton marker. In embryos infected for 9 days 32% of *M.*

marinum was present in the cytosol (Fig. 2C and Fig. S2B). Noteworthy, ESX-1

mutants remain restricted in the lumen of the blood vessels and not able to pass the

basal lamina of the blood vessel wall (54). As only part of the ESX-1 mutant bacteria

is intracellular and the majority restricted in the lumen of the vessels, the ESX-1

mutants were not included in this study. Moreover, there is ample literature on the ESX-

1 localisation in membrane enclosed compartments (16, 22, 25–28, 35). Out of the four

adult zebrafish analyzed, one was discarded as no bacteria could be deteced. In the

other three fish ample bacteria were detected and catagorized as phagosomal or

cytosolic based on the presence of membrane. In adult zebrafish less than five percent

of the bacteria were found in the cytosol (Fig. 2C). Thus, adult but not embryo zebrafish

were able to limit bacterial cytosolic escape. This outcome suggests the presence of

an adaptive immune system promotes bacterial containement in the phago-lysosome,

although other age related factors could be at play.

Cytosolic localization of *M. leprae* is restrained in both armadillo and patient skin

To examine whether cytosolic localization can only be detected in early, innate stages

of infections, we used another model organism; *M. leprae*, the causative agent for

leprosy. Like *Mtb*, these bacteria have been shown to translocate to the cytosol in an

in vitro model (22). We studied early (day 3) and late stage (day 21) infections in

armadillos and in addition, skin biopsies were taken from 4 lepromatous leprosy (LL)

patients with a well-established infection. The skin of the abdomen of armadillos was

infected with both unviable (irradiated) as well as viable *M. leprae.* At the site of

infection, loss of pigment was visible which increased during infection progression (Fig.

3A and B). No differences were observed in loss of pigment between the sites infected

with irradiated *M. leprae* and viable *M. leprae*. At the border of pigment loss, armadillo

skin biopsies infected with viable *M. leprae* were fixed for TEM analysis, to determine

if the localization of the bacteria is cytosolic versus phagosomal. Immuno-gold labelling

against *M. leprae* specific anti-Cell Wall Protein was used to verify that these are

indeed mycobacteria. Similar to the localization of *M. marinum* in adult zebrafish, the

majority of the bacteria were present in membrane enclosed phagosomes and

cytosolic bacteria were only occasionally detected at both day 3 (Fig. 3C) and 21 after

infection. In addition, skin biopsies of 4 different lepromatous leprosy patients taken

from the border of the infection as defined by the depigmentation line and were

analyzed using TEM and immuno-gold labelling for *M. leprae* specific anti-Cell Wall

Protein and lysosomal markers such as Cathepsin D (Fig. 4), LAMP1, and CD63.

Worth mentioning is that based on the ultrastructure of the cells, and more specifically

the ultrastructure of the cell nucleus or the presence of multiple lysosomes, most

bacteria reside in macrophage-like cells. From the 4 patients, individual bacteria were

detected and classified for their subcellular localization based on the presence of

surrounding membrane and 2 or more gold particles detecting Cathepsin D (Fig. S2C).

Similar to the armadillo and adult zebrafish, the percentage of cytosolic mycobacteria

was low (1.1%), while over 700 bacilli were assessed. In conclusion, in both armadillo

and human skin, a low percentage of cytosolic *M. leprae* bacteria is present at all

measured stages of infection.

Immunocompetent BALB/c, but not SCID mice can contain *M. tuberculosis*

inside phagosomes of infected pulmonary cells.

After showing that neither *M. leprae* nor *M. marinum* translocate in high numbers to

the cytosol even at later stages of infection, we next wanted to study the subcellular

localization of *M. tuberculosis* longitudinally in mice. To do so BALB/c mice were

infected with the virulent *Mtb* strain H37Rv and lung tissue was fixed for EM analysis

at day 2, 7, 21, 45 and 120 after infection. Based on the ultra-structure of the nucleus

and/or the high number of lysosomes, the intracellular bacteria were detected in

macrophage-like cells. These samples were immunogold labelled for lysosomes using

LAMP1 or Cathapsin D and imaged using TEM (Fig. 5). As for *M. marinum* in THP1

cells and *M. leprae* in skin, the number of bacteria present in LAMP-1 labeled phago-

lysosomes, membrane enclosed but not LAMP-1 labelled phagosomes and the

number of bacteria in the cytosol were counted (Table 1S). We found that cytosolic localization was highest at day 7 of infection, with still less than five percent of *Mtb* detectable in the cytosol and most bacteria residing in a membrane enclosed compartment. To determine if patient derived strains behave similar to H37Rv, BALB/c mice were infected with two additional *Mtb* strains from lineage 2, often referred to as Beijing family *Mtb* (1998-1500 ancient Beijing) and multi drug resistant strain (2002-0230 Beijing). EM analysis was done at 21, 45 and 120 dpi and demonstrated also low amounts cytosolic bacteria (Table S1). Thus, as shown previously in immune-competent mice, different strains of *Mtb* reside primarily inside membrane enclosed compartments and not inside the cytosol.

To directly address a potential contribution of the adaptive immune system we next quantified *Mtb* cytosolic translocation in SCID mice. SCID mice lack functional T and B cells, key mediators of adaptive immunity in vertebrates and succumb rapidly to *Mtb* infections. SCID mice were infected with *Mtb* strain H37Rv via aerosol and sacrificed 21 days after infection. Lungs were then fixed and processed for EM analysis and sections labelled for LAMP1 using immuno-gold labelling to indicate lysosomes and phago-lysosomes (Table S1). We detected a 10-fold increase in *Mtb* cytosolic translocation in SCID compared to BALB/c mice, arguing that T and B cells are required for optimal clearance of cells with cytosolic bacilli.

Mtb is preferentially located in the cytosol of infected pulmonary cells in IL-1R1 deficient mice

IL-1 is a potent innate inflammatory cytokine critically required for resistance against bacterial infections. Mice deficient in the IL-1 pathway, such as IL-1α and IL-1β are

highly susceptible to *Mtb* infection with increased mortality and bacterial growth in lung

and spleen and development of necrotic granulomatous lesions that more closely

resemble human necrotic lesions (14, 15, 55–57). To investigate the contribution of

this innate immune pathway in the prevention of mycobacterial cytosolic escape, we

infected *Il1r1-/-* and B6 WT mice with *Mtb* and 4 weeks after infection fixed lung tissue

and processed it for TEM analysis. As Mtb is normally difficult to find, first fluorescence

microscopy was performed and tissues were sectioned at 200 nm and stained for

detection of nuclei and mycobacteria (Fig. 6 A,B) to be able to select for the infected

area (54). Whereas in B6 mice only small spots of *Mtb* are detected (like for BALB/c

mice), infected *Il1r1-/-* tissue is heavily labelled and thus heavily infected. Ultrathin

sections were labelled for LAMP1 and CD63 using immuno-gold labelling to indicate

lysosomes and phago-lysosomes for TEM analysis. As before, difference between

phago-lysosomal, phagosomal and cytosolic *Mtb* was determined based on the

presence of immuno-gold labelling and a membrane surrounding the bacteria (Fig.

6C,D and Fig. S2D). *Mtb* was found in the phago-lysosome, the phagosome and in the

cytosol. Strikingly, we detected a 14-fold increase in the number of *Mtb* translocated

to the cytosol in *Il1r1-/-* deficient lungs compared to WT B6 lungs infected with *Mtb*.

While only 2.5% of *Mtb* bacilli were located in the cytosol in the lungs of infected WT

B6 mice, over 30% of the *Mtb* bacilli in the lung were able to escape to the cytosol in

the absence of IL-1 signaling. Thus, IL-1 dependent signals are required to control

cytosolic localization of *Mtb.*

Discussion

The cytosolic localization of mycobacteria has been debated ever since the first description in the late 1980s (18, 20, 58). Our immuno-gold TEM analysis of *Mtb*, BCG and mutant strains in 2007 (22) restarted the discussion and several studies have now confirmed the presence of extra-phagosomal *Mtb* bacilli (16, 21, 41). In addition, the release of DNA into the cytosol was described and is, like escape from the phagosomal compartment, dependent on the ESX-1 secretion system (29, 31, 32). In this study, we showed that *in vivo,* and with an intact immune response *Mtb* is mainly present in membrane bound compartments and not in the cytosol of infected cells. This also holds true for the pathogenic mycobacteria, *M. leprae* and *M. marinum.*

Here, we showed that raising the lysosomal pH did not affect cytosolic localization of *M. marinum.* In THP-1 cells, *M. marinum* can escape irrespective of the raised pH of the lysosome. As well-known already, maturation of the *Mtb* phagosome is altered (38, 44, 45) and our data suggest that the pH of phagosomes has no effect on translocation. Simeone and collegues (2015) (41) showed *in vivo* that when phagosomal acidification was blocked using bafilomycin, an induction of mycobacterial access to the cytosol is detected. Therefore, we hypothesized that raising the phagosomal and lysosomal pH by ConB would result in increased cytosolic localization of *M. marinum.* However, we found at a cellular level, no difference in the percentage of cytosolic *M. marinum.* Our cell-culture derived data merely demonstrate that mycobacteria can translocate irrespective of the pH, and thus manipulation of the pH by ConB did not change the subcellular localization in cells under *in vitro* conditions. Importantly, *in vivo,* as studied here and by Simeone et al., (2015) (41), subcellular localization of mycobacteria is a highly complex and dynamic process.

After inhalation in the lung, *Mtb* is initially taken up by alveolar macrophages and spreads among innate immune cells while delaying the initiating adaptive immunity (59–61). Importantly, both innate and adaptive immunity are critically important for optimal host resistance against *Mtb*. Here, we show that while the lysosomal pH itself has a minimal role, effective adaptive and innate host immunity play a critical role in regulating the number of cells with cytosolic mycobacteria. The delicate balance between bacterial replication and containment by the host immune response is lost when immunity is compromised by either lack of T and B cells in SCID mice, or even more dramatically, in the absence of innate IL-1R1 signaling. Of note, the profound increase in IL-1R1 deficient mice exceeds moderate but significant increase in SCID mice, even though the former likely has a broader set of immunological defects. This may argue that despite increased susceptibility to *Mtb*, certain specific immune pathways like IL-1 may be more directly and preferentially involved in regulating clearance of cells containing cytosolic bacteria. Activation of cytosolic immunosurveillance pathways, including those linked to IL-1, promote rapid clearance by the immune cells attracted to infected cells.

Cytosolic immunosurveillance pathways are triggered by pathogens directly or by pathogenic products entering the cytosol, and are often linked to anti-viral immunity and type I IFN induction as well as inflammasome activation. Inflammasomes are cytosolic signaling complexes that ultimately lead to cytolytic cell death and IL-1 family cytokine processing. It is now established that bacterial DNA can translocate to the cytosol, in an ESX-1 dependent manner, where it is detected by cytosolic DNA sensors such as cGAS and AIM2 (27, 29, 31, 32) cGAS in turn synthesizes a second messenger (cGAMP) which activates the STimulator of IFN Genes (STING) and type I IFN signaling and expression of IFN-α and β (reviewed in (62)). When cytosolic DNA

is detected by AIM-2, the NRLP3 inflammasome is activated and leads to proteolytic cleavage of IL-1β (27, 29, 31, 32). Thus translocation of *Mtb* to the cytosol triggers a cascade of responses resulting at a cellular level in cell death (22, 63, 64). The connection between translocation, cellular damage and cell death has recently been studied in more detail; plasma membrane lysis caused by Mtb triggers inflammasome activation, IL-1β release and pyroptosis, a form of programmed necrosis (35). Beckwith et al (35) demonstrated that ESX-1-mediated phagosomal damage is a requirement for NLRP3 activation and although their live cell imaging demonstrated that inflammasome activation by *Mtb* is independent of lysosomal damage, others have demonstrated that active cathepsin release from ruptured phagolysosomes is a trigger of NLRP3 inflammasome activation (64). Taken together, after rupture of the phagolysomes by *Mtb*, cell death is induced and with an intact immune response, via the IL-1 signaling, these cells are cleared. This explains the fact that in skin biopsies of leprosy patients, armadillo, lung of BalbC mice, adult zebrafish the majority of the mycobacteria is detected in phagolysosomes.

We have previously shown that IL-1 and type I IFNs exhibit potent cross regulation important for host resistance against *Mtb* with excessive type I IFN induction in the absence of IL-1 signaling (14, 15, 65). The elevated type I IFNs expression in the absence of IL-1 contributed to the increased susceptibility of the *Il1r1-/-* mice, as mice doubly deficient in IL-1R1 and IFNAR1 displayed increased resistance (14). Our new findings here of increased cytosolic *Mtb* in *Il1r1-/-* mice provide a possible molecular explanation for the increased type I IFN production previously reported. In this context, our recent study demonstrated that infected cells themselves do not need to express IL-1R1 *in vivo* to mediate host resistance and that IL-1R1 expression coordinates immune responses in multiple cells types (13). Along these lines, it has been proposed

that IL-1R1 on non-immune cells was required for the ability of infected alveolar

macrophage to leave the airway to establish infection in the interstitial lung space (6).

Thus, cytosolic containment of bacilli in infected cells, may not require direct cell-

autonomous antimicrobial signaling pathways but may be the result of dynamic cellular

interactions between infected cells and cells of both non-hematopoietic and bone

marrow origin.

Overall, this study establishes that high level cytosolic escape of mycobacteria can

indeed occur *in vivo,* but mainly when host resistance is compromised. When the

complete adaptive immune system including B and T cells is abrogated, like in

zebrafish embryos and SCID mice, a substantial percentage of mycobacteria is

detected in the cytosol as compared to immunocompetent hosts. Strikingly, the highest

proportion cytosolic *Mtb* was observed in mice lacking IL-1 signaling. This argues that

the IL-1 pathway is crucial for the control of the number of cytosolic mycobacteria and

likely IL-1 mediated resistance to *Mtb*.

Methods

Bacteria

M. marinum E11 strain was grown on Middlebrook 7H10 plates supplemented with

OADC. A single colony was inoculated into 7H9 liquid medium (BD) supplemented with

10% ADC and 0.05% Tween 80 and incubated with shaking at 30°C and grown to an

OD600 of 0.6-1. Before infection the bacteria were centrifuged at 750 rpm to remove

clumps, leaving a bacterial suspension.

Inhibiting acidification

Inhibition of acidification of phagosomes/lysosomes in THP-1 cells treated with Concanamycin B (ConB) (Alexis Biochemicals: 380098C100) was assessed by confocal microscopy. THP-1 macrophages were grown in Roswell Park Memorial Institute (RPMI)-1640 medium supplemented with 10% FCS and ConB was added to the cells in different concentrations 1h and 24h before fixation. The concentration 10nM of ConB strongly inhibited acidification and did not affect THP-1 cell viability after a 48h incubation period. THP-1 cells were washed 3 times with RPMI-1640 10% FCS medium and kept as a control or pretreated with 10nM ConB for 1 hour. Cells were infected with *M. marinum* (MOI 10:1) in the presence or absence of the acidification inhibitor. After an incubation time of 1 hour at 32°C, cells were washed 3 times with culture medium without antibiotics and with or without acidification inhibitors to remove extracellular bacteria. After washing, the cells were further incubated in culture medium with or without ConB for 24 or 48 hours at 32°C prior to fixation. Fixed samples were prepared for cryo-immunogold microscopy, sectioned for EM and immuno-gold labeled with CD63. To assess the effect on the amount of cytosolic *M. marinum*, from 3 independent experiments, 100-200 randomly chosen bacteria from one grid were counted and for each bacterium it was determined if it was cytosolic or resided in a phago-lysosome.

DAMP assay for measuring lysosomal acidification

Luminal acidification in lysosomes was measured via the probe, DAMP [3-(2,4-dinitroanilino)-3′-amino-N-methyldipropylamine], incubated at 30 µM for 30 min to allow accumulation in acidic compartments. DAMP was quantified by immuno-gold staining using anti-DNP.

Mouse infections

IL1r1⁻/⁻ mice were purchased from Jackson Laboratories (JAX 3018) and backcrossed to C57BL/6 control mice from Taconic Farms (Hudson, NY) for 11 generations. Male and female mice, 8-12 weeks of age were infected via the aerosol route with *Mtb* H37Rv (100-200 CFU/mouse) as previously described (13) and sacrificed 27 days later. In short, mice were infected using a whole-body inhalation system (Glas- Col; Terre Haute, IN) exposing the mice to aerosolized *Mtb*. Lungs were perfusion-fixed in 4% paraformaldehyde and 0.4% glutaraldehyde overnight. After fixation, tissues were transferred to storage buffer containing 0.5 % paraformaldehyde. All animals were maintained in an Association for Assessment and Accreditation of Laboratory Animal Care (AALAC)-accredited BSL2 or BSL3 facilities at the National Institutes of Health (NIH) and experiments performed in compliance with an animal study proposal approved by the National Institute of Immunology Allergy and Infectious Diseases Animal Care and Use Committee.

SCID Mice

Severe combined immunodeficiency (SCID) mice were purchased from Charles River and were aerosol infected with *M. tuberculosis* H37Rv for 21 days, when mice were sacrificed and lungs fixed by perfusion fixation as described below.

BALB/c, B6 mice

Pathogen-free male BALB/c mice, 6-8 weeks old, were anaesthetized with sevoflurane vapours (Abbott Laboratories, Abbott Park, IL, USA) and 100 µl of PBS with 2.5×10^5 viable H37Rv bacilli or either Beijing clinical isolates were inoculated intra-tracheally using a stainless steel cannula. Groups of 15 animals were then maintained in cages fitted with microisolators in a BSL-3 biosecurity level facility. Following infection, three

mice were euthanized by exsanguination under anesthesia with pentobarbital at days

2, 7, 21, 45 and 120 of infection. Lung tissues were fixed by perfusion fixation as

described below.

Zebrafish

Zebrafish embryos and adult zebrafish (Danio rerio) were microinjected with *M.*

marinum strain E11 as described by (53) at 30 °C. In short, zebrafish embryos were

infected 28h post infection with 100 CFU *M. marinum* wild type through micro-injection

in the caudal vein. Adult zebrafish were anaesthetized in 0.02% MS-222 (Sigma) and

injected intra-peritoneally with $2x10^4$ *M. marinum* Tn::ESX-5. The embryos were

incubated for 6 or 9 days and fixed as described below. For classification of the

subcellular localization, 3 embryos of 9 days incubation were used. Three adult fish

were infected with E11 or E11 ESX5 mutant (53) and sacrificed at day 11 when the

spleen was dissected out and fixed as described below.

Armadillo

Male Armadillos were intradermally inoculated in the abdomen with either 1×10^7 live

or irradiated *M. leprae* (0.1 ml). After, 3, 11 and 21 days post-inoculation, 6 mm punch

biopsies were taken from the boarder of the depigmented area at the inoculation sites

and fixed for 2 hours as described below. For classification of the subcellular

localization, 1-2 punches were used from day 3 and day 21. As no lysosomal markers

suited for immuno-TEM analysis were known, anti-Cell Wall Protein (CWP) was used

to immuno-label the leprosy bacteria.

Human samples

Leprosy skin biopsies were taken at the boarder of the depigmented lesions with

written consent from 4 different lepromatous leprosy patients. Materials were directly

incubated in EM grade fixatives and transported to the EM lab in fixatives.

Fixation of tissues

All tissues were fixed in a combination of 4% paraformaldehyde and 0.4%

glutaraldehyde in 0.2M PHEM (with 240mM PIPES, 100mM HEPES, 8mM MgCl2 and

40mM EGTA) buffer. Fixation for at least 2 hours in fixative containing glutaraldehyde

is essential to kill mycobacteria. After fixation, tissues were transferred to storage

buffer containing 0.5% paraformaldehyde in 0.2M PHEM buffer.

Embedding and sectioning

After fixation, tissues were washed in phosphate buffered saline (PBS), to remove

fixatives. The required structures were dissected out using a razor blade and cut into

1-3 mm^2 sized blocks. Lung tissue was embedded in increasing percentages gelatin

(2%, 5% and 12% in 0.1M phosphate buffer) and incubated at 37 °C. After the removal

from liquid gelatin, blocks were incubated overnight in 2.3M sucrose at 4 °C. Then

blocks were snap frozen and stored in liquid nitrogen till sectioned. After trimming at -

100 °C, semi thin sectioning was performed for analysis with fluorescence microscopy

or ultra-thin sectioning at -120 °C was performed using a diamond Diatome cryo-

immuno knife on a Leica Ultracut UC6. Sections are picked up with a loop filled with a

1:1 mixture of 2,3 M sucrose and 1% tylose (Metylcellulose, G1095) in milliQ water and

placed on a copper, formvar coated grid or glass slides. Grids with sections were stored

at 4 °C till immuno-labeled. Fluorescence Microscopy was used to search the region

of infection as described in (54, 66). In short, semi-thin sections (200-300nm) of the

whole sample were labelled with Hoechst 33342 (Thermo Fisher) to indicate the nuclei

of the tissue and anti-Cell Wall protein labelling was used to indicate mycobacteria.

Immuno-gold labelling

Grids with cryo-sections were incubated on 2% gelatin in 0.1M phosphate buffer plates

at 37 °C for 30 min. Thereafter, grids were washed with PBS/0.02M glycine, blocked

with 1% BSA and incubated for 45 minutes with primary antibody. Various antibodies

were used on different tissues: for human skin and sputum: Cathapsin-B (Zymed, clone

1C11), Lysosome Associated Membrane Protein 1 and 2 (LAMP1 and LAMP2)

(Pharmingen, H4A3 and H4B3), Cluster of Differentiation CD63 (clone 435 Sanquin).

For Zebrafish anti-actin (Sigma AC-15), and tested but without specific labeling: anti-

LAMP1 (Pharmingen, H4A3 and H4B3, Abcam ab67283), anti CD63 (clone 435

Sanquin). For *M. leprae*, anti Cell Wall Protein (C188, a kind gift form John Spencer

and Patrick Brennan Colorado State University). DAMP (N-(3-((2,4-Dinitrophenyl)

Amino)propyl)-N-(3-Aminopropyl) Methylamine, Dihydrochloride) detection was

performed using anti dinitrophenol (DNP) (Polyclonal Anti-DNP, Oxford Biomedical

Research). All antibodies were diluted in 1% BSA in PBS. After washing in PBS/0.02M

glycine and blocking in 0.1% BSA in PBS/0.02M glycine, grids are incubated on

bridging antibody when primary antibody was monoclonal or goat origin. After washing

and blocking, grids are incubated on protein A gold diluted in 1% BSA in PBS. To

remove unbound gold, grids were washed with PBS and fixed using 1% glutaraldehyde

in PBS, then washed with MiliQ and contrasted with uranyl acetate and methylcellulose

at pH 4.

Immuno-fluorescence labelling

Semi-thin sections (200-300nm) placed on glass were washed with PBS/0.02M

Glycine and incubated with primary antibody anti Cell Wall Protein for 45 minutes.

Then, the sections were washed with PBS and incubated with secondary antibody

goat-anti-rabbit Alexa 488 (Mol. Probes, A32731) for 20 minutes and Hoechst 33342

for 5 minutes (Thermo Fisher, H3570). After washing with PBS, samples were mounted

with Vectashield. The sections were imaged using a Leica DM6 wide-field microscope

and images were analyzed using FIJI.

Statistical analysis and subcellular classification

Statistical analysis was performed using Graphpad-Prism 8.0 software. In the legends,

the average is given with the standard deviation and n indicate the number of bacteria

localized in a specific subcellular compartment. Significance was determined by using

unpaired T test and defined in the graphs as $P < 0.05$ = *, $P < 0.01$ = ** and $P < 0.001$ =

***. The number of biological samples, and bacteria counted in mice are listed in Table

S1.

Classification of the subcellular localization of bacteria was performed blindfolded and

by 2 individual counters to establish inter-counter reproducibility. Bacteria are classified

as cytosolic when 1/3 or less of the bacteria or bacterial cluster is surrounded by a

visible membrane and 2 or less gold particles detecting lysosomal markers are present.

Bacteria are classified as phagosomal when 1/3 or more of the bacteria or bacterial

cluster is surrounded by a visible membrane and 2 or less gold particles detecting

lysosomal markers is present and phago-lysosomal when 1/3 or more of the bacteria

or bacterial cluster is surrounded by a visible membrane and 3 or more gold particles

detecting lysosomal markers are present.

Ethics statement

SCID mouse infections were performed in agreement with European and French guidelines (Directive 86/609/CEE and Decree 87–848 of 19 October 1987). The experiments received the approval by the Institut Pasteur Safety Committee (Protocol 11.245) and the ethical approval by local ethical committees "Comité National de Réflexion Ethique sur l'Expérimentation Animale N° 59 (CNREEA)".

Adult zebrafish of the local Free University of Amsterdam (VU) line were handled in compliance with the local animal welfare regulations and approved by the local animal welfare commission (IvD) of the VU/Amsterdam University Medical Centre. For zebrafish embryo experiments that are performed within the grace period (i.e. first 6 days) no special permission is allowed, since these experiments fall under animal experimentation law according to the EU Animal Protection Directive 2010/63/EU.

BALB/c mouse infections were approved by the Institutional Ethics Committee of Animals Experimentation of the National Institute of Medical Sciences and Nutrition Salvador Zubirán in accordance with the guidelines of the Mexican national regulations on Animal Care and Experimentation NOM 062-ZOO-1999

Experiments using armadillos were performed in accordance with USPHS Policy on the Humane Care and Use of Laboratory Animals and the USDA Animal and Plant Health Inspection Service. The Institutional Animal Care and Use Committee reviewed and approved the protocol.

Acknowledgements

We would like to thank Wikky Tigchelaar – Gutter, Pekka Kujala, Hans Janssen for EM analysis, Gidado Mustapha, Nigeria for leprosy skin biopsies. We are also grateful to

Wafa Frigui and Alexandre Pawlik for help with infection experiments. We would like

to thank Branch Moody for critically reading the manuscript and the helpful comments.

RB acknowledges support by ANR-10-LABX-62-IBEID. SvdN PJP and NvdW

acknowledge NIH grant no AI116604 and Netherlands Leprosy Relief. The NIH, NIAID

funded the armadillo studies through the Interagency Agreement No. AAI15006 with

the Health Resources and Services Administration, Healthcare Systems Bureau,

National Hansen's Disease Program.

Author contributions

580 SvdN, ER, KDMB and NvdW wrote the manuscript, SvdN, MvZ, KdP, AG, SR, SM,

581 DH, PP performed EM analysis AvdS, WB, RB, RHP, MP performed infection

582 experiments, KDMB performed and designed experiments and NvdW conceived and

583 designed experiments.

Declaration of interests

585 The authors declare no competing interests.

References

1.   World Health Organization. 2019. Global Tuberculosis Report 2019: Fact sheet

https://doi.org/10.1017/CBO9781107415324.004.

2.   Doitsh G, Greene WC. 2016. Dissecting how CD4 T cells are lost during HIV

infection. Cell Host Microbe 19:280–291.

3.   Queval CJ, Brosch R, Simeone R. 2017. The Macrophage : A Disputed

Fortress in the Battle against Mycobacterium tuberculosis. Front Microbiol.

4.   Lerner TR, Borel S, Gutierrez MG. 2015. The innate immune response in

human tuberculosis. Cell Microbiol 17:1277–1285.

5.   Huang L, Nazarova E V., Tan S, Liu Y, Russell DG. 2018. Growth of

Mycobacterium tuberculosis in vivo segregates with host macrophage

metabolism and ontogeny. J Exp Med 215:1135–1152.

6.   Cohen SB, Gern BH, Delahaye JL, Adams KN, Plumlee CR, Winkler JK,

Sherman DR, Gerner MY, Urdahl KB. 2018. Alveolar Macrophages Provide an

Early Mycobacterium tuberculosis Niche and Initiate Dissemination. Cell Host

Microbe 24:439-446.e4.

7.   Marino S, Pawar S, Fuller CL, Reinhart TA, Flynn JL, Kirschner DE. 2004.

Dendritic Cell Trafficking and Antigen Presentation in the Human Immune

Response to Mycobacterium tuberculosis . J Immunol 173:494–506.

8.   Sayes F, Blanc C, Ates LS, Deboosere N, Orgeur M, Le Chevalier F, Gröschel

MI, Frigui W, Song OR, Lo-Man R, Brossier F, Sougakoff W, Bottai D, Brodin

P, Charneau P, Brosch R, Majlessi L. 2018. Multiplexed Quantitation of

Intraphagocyte Mycobacterium tuberculosis Secreted Protein Effectors. Cell

Rep 23:1072–1084.

9.   Muruganandah V, Sathkumara HD, Navarro S, Kupz A. 2018. A Systematic

Review: The Role of Resident Memory T Cells in Infectious Diseases and Their

Relevance for Vaccine Development. Front Immunol 9.

10.   Flynn JAL, Chan J, Triebold KJ, Dalton DK, Stewart TA, Bloom BR. 1993. An

essential role for interferon γ in resistance to mycobacterium tuberculosis

infection. J Exp Med 178:2249–2254.

11.   Dupuis S, Döffinger R, Picard C, Fieschi C, Altare F, Jouanguy E, Abel L,

Casanova JL. 2000. Human interferon-g-mediated immunity is a genetically

controlled continuous trait that determines the outcome of mycobacterial

invasion. Immunol Rev 178:129–137.

12.   Guirado E, Mbawuike U, Keiser TL, Arcos J, Azad AK, Wang SH, Schlesinger

LS. 2015. Characterization of host and microbial determinants in individuals

with latent tuberculosis infection using a human granuloma model. MBio 6:1–

13.

13.   Bohrer AC, Tocheny C, Assmann M, Ganusov V V., Mayer–Barber KD. 2018.

Cutting Edge: IL-1R1 Mediates Host Resistance to Mycobacterium tuberculosis

by Trans -Protection of Infected Cells . J Immunol 201:1645–1650.

14.   Mayer-Barber KD, Andrade BB, Oland SD, Amaral EP, Barber DL, Gonzales J,

Derrick SC, Shi R, Pavan Kumar N, Wei W, Yuan X, Zhang G, Cai Y, Babu S,

Catalfamo M, Salazar AM, Via LE, Barry III CE, Sher A. 2014. Host-directed

therapy of tuberculosis based on interleukin-1 and type I interferon crosstalk.

Nature 118:6072–6078.

15.   Mayer-Barber KD, Andrade BB, Barber DL, Hieny S, Feng CG, Caspar P,

633   White S, Gordon S, Sher A. 2011. Innate and adaptive interferons suppress IL-

634   1α and IL-1β production by distinct pulmonary myeloid subsets during

635   Mycobacterium tuberculosis infection. Immunity 36:1023–1034.

636   16.   Houben D, Demangel C, van Ingen J, Perez J, Baldeón L, Abdallah AM,

637   Caleechurn L, Bottai D, van Zon M, de Punder K, van der Laan T, Kant A,

638   Bossers-de Vries R, Willemsen P, Bitter W, van Soolingen D, Brosch R, van

639   der Wel N, Peters PJ. 2012. ESX-1-mediated translocation to the cytosol

640   controls virulence of mycobacteria. Cell Microbiol 14:1287–1298.

641   17.   Lerner TR, Carvalho-Wodarz CDS, Repnik U, Russell MRG, Borel S, DIedrich

642   CR, Rohde M, Wainwright H, Collinson LM, Wilkinson RJ, Griffiths G, Gutierrez

643   MG. 2016. Lymphatic endothelial cells are a replicative niche for

644   Mycobacterium tuberculosis. J Clin Invest 126:1093–1108.

645   18.   Leake ES, Myrvik QN, Wright MJ, Carolina N. 1984. Phagosomal Membranes

646   of Mycobacterium bovis BCG-Immune Alveolar Macrophages Are Resistant to

647   Disruption by Mycobacterium tuberculosis H37Rv. Infect Immun 45:443–446.

648   19.   Lerner TR, Queval CJ, Fearns A, Repnik U, Griffiths G, Gutierrez MG. 2018.

649   Phthiocerol dimycocerosates promote access to the cytosol and intracellular

650   burden of Mycobacterium tuberculosis in lymphatic endothelial cells. BMC Biol

651   1–13.

652   20.   Mcdonough KA, Kress Y, Bloom BR. 1993. Pathogenesis of Tuberculosis :

653   Interaction of Mycobacterium tuberculosis with Macrophages. Infect Immun

654   61:2763–2773.

21.   Simeone R, Bobard A, Lippmann J, Bitter W, Majlessi L, Brosch R, Enninga J.

2012. Phagosomal rupture by Mycobacterium tuberculosis results in toxicity

and host cell death. PLoS Pathog 8.

22.   van der Wel N, Hava D, Houben D, Fluitsma D, van Zon M, Pierson J, Brenner

M, Peters PJ. 2007. M . tuberculosis and M . leprae Translocate from the

Phagolysosome to the Cytosol in Myeloid Cells. Cell 1287–1298.

23.   Lerner TR, Queval CJ, Lai RP, Russell M, Fearns A, Greenwood DJ, Collins L,

Wilkinson RJ, Gutierrez MG. 2020. Mycobacterium tuberculosis cords in the

cytosol of live lymphatic endothelial cells to evade host immune surveillance.

JCI Insight.

24.   Vaziri F, Brosch R. 2019. ESX / Type VII Secretion Systems — An Important

Way Out for Mycobacterial Proteins. Microbiol Spectr 7(4):PSIB-.

25.   Augenstreich J, Arbues A, Simeone R, Haanappel E, Wegener A, Sayes F, Le

Chevalier F, Chalut C, Malaga W, Guilhot C, Brosch R, Astarie-Dequeker C.

2017. ESX-1 and phthiocerol dimycocerosates of Mycobacterium tuberculosis

act in concert to cause phagosomal rupture and host cell apoptosis. Cell

Microbiol 19:1–19.

26.   Gröschel MI, Sayes F, Shin SJ, Frigui W, Pawlik A, Orgeur M, Canetti R,

Honoré N, Simeone R, van der Werf TS, Bitter W, Cho SN, Majlessi L, Brosch

R. 2017. Recombinant BCG Expressing ESX-1 of Mycobacterium marinum

Combines Low Virulence with Cytosolic Immune Signaling and Improved TB

Protection. Cell Rep 18:2752–2765.

27.   Kupz A, Zedler U, Stäber M, Perdomo C, Dorhoi A, Brosch R, Kaufmann SHE.

2016. ESAT-6 – dependent cytosolic pattern recognition drives noncognate

tuberculosis control in vivo. J Clin Invest 126:2109–2122.

28.   Pym AS, Brodin P, Majlessi L, Brosch R, Demangel C, Williams A, Griffiths KE,

681   Marchal G, Leclerc C, Cole S. 2003. Recombinant BCG exporting ESAT-6

682   confers enhanced protection against tuberculosis. Nat Med 9:533–539.

683   29.   Collins AC, Cai H, Li T, Franco LH, Li XD, Nair VR, Scharn CR, Stamm CE,

684   Levine B, Chen ZJ, Shiloh MU. 2015. Cyclic GMP-AMP Synthase Is an Innate

685   Immune DNA Sensor for Mycobacterium tuberculosis. Cell Host Microbe

686   17:820–828.

687   30.   Majlessi L, Brosch R. 2015. Mycobacterium tuberculosis Meets the Cytosol:

688   The Role of cGAS in Anti-mycobacterial Immunity. Cell Host Microbe 17:733–

689   735.

690   31.   Wassermann R, Gulen MF, Sala C, Perin SG, Lou Y, Rybniker J, Schmid-

691   Burgk JL, Schmidt T, Hornung V, Cole ST, Ablasser A. 2015. Mycobacterium

692   tuberculosis Differentially Activates cGAS- and Inflammasome-Dependent

693   Intracellular Immune Responses through ESX-1. Cell Host Microbe 17:799–

694   810.

695   32.   Watson RO, Bell SL, MacDuff DA, Kimmey JM, Diner EJ, Olivas J, Vance RE,

696   Stallings CL, Virgin HW, Cox JS. 2015. The cytosolic sensor cGAS detects

697   Mycobacterium tuberculosis DNA to induce type I interferons and activate

698   autophagy. Cell Host Microbe 17:811–819.

699   33.   Srinivasan L, Gurses SA, Hurley BE, Miller JL, Karakousis PC, Briken V. 2016.

700   Identification of a Transcription Factor That Regulates Host Cell Exit and

701   Virulence of Mycobacterium tuberculosis. PLoS Pathog 12:1–29.

702   34.   Quigley J, Hughitt VK, Velikovsky CA, Mariuzza RA, El-Sayed NM, Briken V.

703   2017. The cell wall lipid PDIM contributes to phagosomal escape and host cell

exit of Mycobacterium tuberculosis. MBio 8:1–12.

35.   Beckwith KS, Beckwith MS, Ullmann S, Sætra RS, Kim H, Marstad A, Åsberg

SE, Strand TA, Haug M, Niederweis M, Stenmark HA, Flo TH. 2020. Plasma

membrane damage causes NLRP3 activation and pyroptosis during

Mycobacterium tuberculosis infection. Nat Commun 11.

36.   Mwandumba HC, Russell DG, Nyirenda MH, Anderson J, White SA, Molyneux

ME, Squire SB. 2004. Mycobacterium tuberculosis Resides in Nonacidified

Vacuoles in Endocytically Competent Alveolar Macrophages from Patients with

Tuberculosis and HIV Infection 1. J Immunol.

37.   Russell DG, Mwandumba HC, Rhoades EE. 2002. Mycobacterium and the

coat of many lipids. J Cell Biol 158:421–426.

38.   Armstrong JA, D'Arcy Hart P. 1971. Response of cultured macrophages to

mycobacterium tuberculosis, with observations on fusion of lysosomes with

phagosomes. J Exp Med 134:713–740.

39.   Sturgill-Koszycki S, Schlesinger PH, Chakraborty P, Haddix P., Collins H., Fok

A., Allen R., Gluck S., Russell DG. 1994. Lack of acidification in

Mycobacterium phagosomes produced by exclusion of the vesicular proton-

ATPase. Science (80-) 263:678–681.

40.   Sundaramurthy V, Korf H, Singla A, Scherr N, Nguyen L, Ferrari G, Landmann

R, Huygen K, Pieters J. 2017. Survival of Mycobacterium tuberculosis and

Mycobacterium bovis BCG in lysosomes in vivo. Microbes Infect 19:515–526.

41.   Simeone R, Sayes F, Song O, Gröschel MI, Brodin P. 2015. Cytosolic Access

of Mycobacterium tuberculosis : Critical Impact of Phagosomal Acidification

Control and Demonstration of Occurrence In Vivo. PLOS Pathog 1–24.

42.   Stamm LM, Morisaki JH, Gao LY, Jeng RL, McDonald KL, Roth R, Takeshita

S, Heuser J, Welch MD, Brown EJ. 2003. Mycobacterium marinum Escapes

from Phagosomes and Is Propelled by Actin-based Motility. J Exp Med

198:1361–1368.

43.   Sharma R, Lahiri R, Scollard DM, Pena M, Williams DL, Adams LB, Figarola J,

Truman RW. 2013. The armadillo: A model for the neuropathy of leprosy and

potentially other neurodegenerative diseases. DMM Dis Model Mech 6:19–24.

44.   Clemens DL, Lee BY, Horwitz MA. 2000. Mycobacterium tuberculosis and

Legionella pneumophila phagosomes exhibit arrested maturation despite

acquisition of Rab7. Infect Immun 68:5154–5166.

45.   Russell DG, Vanderven B, Glennie S, Mwandumba H, Heyderman R. 2009.

The macrophage marches on its phagosome: dynamic assays of phagosome

function. Nat Rev Immunol 9:594–600.

46.   Wong D, Bach H, Sun J, Hmama Z, Av-gay Y. 2011. Mycobacterium

tuberculosis protein tyrosine to inhibit phagosome acidification. PNAS

108:19371–19376.

47.   Vieira O V., Botelho RJ, Grinstein S. 2002. Phagosome maturation: aging

gracefully. Biochem J 366:689–704.

48.   Buter J, Cheng T-Y, Ghanem M, Grootemaat AE, Raman S, Feng X, Plantijn

AR, Ennis T, Wang J, Cotton RN, Layre E, Ramnarine AK, Mayfield JA, Young

DC, Jezek Martinot A, Siddiqi N, Wakabayashi S, Botella H, Calderon R,

Murray M, Ehrt S, Snider BB, Reed MB, Oldfield E, Tan S, Rubin EJ, Behr MA,

van der Wel NN, Minnaard AJ, Moody DB. 2019. Mycobacterium tuberculosis

releases an antacid that remodels phagosomes. Nat Chem Biol 15:889–899.

49. Vandal OH, Nathan CF, Ehrt S. 2009. Acid resistance in Mycobacterium

tuberculosis. J Bacteriol 191:4714–4721.

50. Young DC, Layre E, Pan S, Tapley A, Adamson J, Wu Z, Buter J, Minnaard AJ,

Coscolla M, Gagneux S, Copin R, Ernst JD, Bishai W, Snider BB, Moody DB.

2015. chemical markers of Mycobacterium tuberculosis infection. Chem Biol

22:516–526.

51. Crowle AJ, Dahl R, Ross E, May MH. 1991. Evidence that vesicles containing

living, virulent Mycobacterium tuberculosis or Mycobacterium avium in cultured

human macrophages are not acidic. Infect Immun 59:1823–1831.

52. Langenau DM, Ferrando AA, Traver D, Kutok JL, Hezel JD, Kanki JP, Zon LI,

Look AT, Trede NS. 2004. In vivo tracking of T cell development , ablation ,

and engraftment in transgenic zebrafish. PNAS 101:7369–7374.

53. Weerdenburg EM, Abdallah AM, Mitra S, De Punder K, Van der Wel NN, Bird

S, Appelmelk BJ, Bitter W, Van der Sar AM. 2012. ESX-5-deficient

Mycobacterium marinum is hypervirulent in adult zebrafish. Cell Microbiol

14:728–739.

54. van Leeuwen LM, Boot M, Kuijl C, Picavet DI, van Stempvoort G, van der Pol

SMA, de Vries HE, van der Wel NN, van der Kuip M, van Furth AM, van der

Sar AM, Bitter W. 2018. Mycobacteria employ two different mechanisms to

cross the blood–brain barrier. Cell Microbiol 20:1–17.

55. Juffermans NP, Florquin S, Camoglio L, Verbon A, Kolk AH, Speelman P,

Deventer SJH Van. 2000. Interleukin-1 Signaling Is Essential for Host Defense during Murine Pulmonary Tuberculosis. J Infect Dis 902–908.

56.   Mayer-Barber KD, Barber DL, Shenderov K, White SD, Wilson MS, Cheever A, Kugler D, Hieny S, Caspar P, Núñez G, Schlueter D, Flavell RA, Sutterwala FS, Sher A. 2010. Cutting Edge: Caspase-1 Independent IL-1β Production Is Critical for Host Resistance to Mycobacterium tuberculosis and Does Not Require TLR Signaling In Vivo . J Immunol 184:3326–3330.

57.   Yamada H, Mizuno S, Horai R, Iwakura Y, Sugawara I. 2000. Protective role of interleukin-1 in mycobacterial infection in IL-1 alpha/beta double knockout mice. Lab Invest 80:759–767.

58.   Myrvik QN, Leake ES, Wright MJ. 1984. Disruption of phagosomal membranes by the H37Rv strain of Mycobacterium tuberculosis. A correlate of virulence. Am Rev Respir Dis 129:322–328.

59.   Rothchild AC, Olson GS, Nemeth J, Amon LM, Mai D, Gold ES, Diercks AH, Aderem A. 2019. Alveolar macrophages generate a non-canonical NRF2-driven transcriptional response to Mycobacterium tuberculosis in vivo. Sci Immunol 4.

60.   Samstein M, Schreiber HA, Leiner IM, Sušac B, Glickman MS, Pamer EG. 2013. Essential yet limited role for CCR2+ inflammatory monocytes during Mycobacterium tuberculosis-specific T cell priming. Elife https://doi.org/10.7554/eLife.01086.

61.   Mayer-Barber KD, Barber DL. 2015. Innate and adaptive cellular immune responses to Mycobacterium tuberculosis infectionCold Spring Harbor Perspectives in Medicine.

62.   Chen Q, Sun L, Chen ZJ. 2016. Regulation and function of the cGAS-STING

pathway of cytosolic DNA sensing. Nat Immunol 17:1142–1149.

63.   Abdallah AM, Bestebroer J, Savage NDL, de Punder K, van Zon M, Wilson L,

Korbee CJ, van der Sar AM, Ottenhoff THM, van der Wel NN, Bitter W, Peters

PJ. 2011. Mycobacterial Secretion Systems ESX-1 and ESX-5 Play Distinct

Roles in Host Cell Death and Inflammasome Activation. J Immunol 187:4744–

4753.

64.   Amaral EP, Riteau N, Moayeri M, Maier N, Mayer–Barber KD, Pereira RM,

Lage SL, Kubler A, Bishai WR, Império-Lima MR, Sher A, Andrade BB. 2018.

Lysosomal Cathepsin Release Is Required for NLRP3-Inflammasome

Activation by Mycobacterium tuberculosis in Infected Macrophages. Front

Microbiol 9.

65.   Ji DX, Yamashiro LH, Chen KJ, Mukaida N, Kramnik I, Darwin KH, Vance RE.

2019. Type I interferon-driven susceptibility to Mycobacterium tuberculosis is

mediated by interleuking-1 receptor antagonist IL-1Ra. Nat Microbiol 4:2128–

2135.

66.   van der Wel NN, Fluitsma DM, Dascher CC, Brenner MB, Peters PJ. 2005.

Subcellular localization of mycobacteria in tissues and detection of lipid

antigens in organelles using cryo-techniques for light and electron microscopy.

Curr Opin Microbiol 8:323–330.

Main Fig. titles + Legends

Legend Fig. 1. The pH of the phago-lysosome does not affect cytosolic

translocation of *M. marinum*. A) Electron micrograph of a THP-1 cell infected with

M. marinum for 24 hour in the presence of ConB showing *M. marinum* in the cytosol

without membrane and CD63 labeling. CD63 immunolabelling indicated by 10 nm gold

particles is present on the multivesicular lysosome in the left top corner. A') schematic

representation of micrograph in A with in blue lines host membranes, black dots CD63

labelling indicated by 10 nm gold particles, orange bacteria and bar represents 200

nm. B) Quantification of the percentage of *M. marinum* in the cytosol 24 and 48 hours

after infection using immuno-gold labelling for CD63 (see also Fig. S2A). Cells were

treated with ConB to raise lysosomal pH, which did not affect the percentage of

cytosolic *M. marinum*. Error bars represent the standard deviation of 3 experiments.

Fig. 2

Legend Fig. 2: Cytosolic localization of *M. marinum* in zebrafish is abundant

when the adaptive immune system is not yet developed. Embryo and adult

zebrafish infected with *M. marinum* were analysed using TEM. A) Cytosolic *M.*

marinum in zebrafish embryo tissue. B) Phagosomal *M. marinum* ΔESX5 in adult

zebrafish tissue. A') and B') Schematic representation of A and B, with black dots

indicating actin immuno-gold labelling, orange lines *M. marinum* and blue lines

phagosomal and host membranes, green lines mitochondria. C) Quantification of the

percentage of cytosolic *M. marinum* at embryo day 9 and in spleen adult zebrafish at

day 11 (see also Fig. S2B). Error bars indicate standard deviation between 3 different

zebrafish embryos and 3 adult fish, n represents the total number of bacteria counted.

Legend Fig. 3: Restrained cytosolic *M. leprae* localization in armadillo skin biopsies. Live *M. leprae* injected in the skin of the armadillo at the red circle indicated by the white arrow text box. The skin was observed at A) 3 days after infection and B) 21 days after infection. At infection sites loss of pigmentation was detected. C) TEM image of infected armadillo skin biopsy 3 days after infection with viable *M. leprae*. Immuno-gold labelling using αCWP to indicate *M. leprae*. C') enlargement of boxed area in C, *M. leprae* enclosed by host membranes and thus phagosomal. C''), enlargement of lower boxed area in C, *M. leprae* was not enclosed by host membranes and thus cytosolic. C'''), C'''') Schematic representation of C' and C'' with in orange *M. leprae* and the blue lines indicate host membranes and green mitochondrial membranes. Total number of bacteria detected at day 3 is 47 and at day 21 is 45.

855 **Fig. 4.**

Legend Fig. 4. Restrained cytosolic localization of *M. leprae* in human skin biopsies. Leprosy patient skin biopsies were analyzed using TEM. Immuno-gold labelling for Cathepsin-D was used to label lysosomes and phagolysosomes. A) TEM image of patient skin biopsy with cytosolic *M. leprae*, the mycobacteria was not enclosed with host membranes. A') Schematic representation of A, orange indicate *M. leprae,* blue lines indicate host membranes, and green mitochondria. B) *M. leprae* present in the phagolysosome, the mycobacterium is enclosed by host membranes, immunogold labelled for lysosomal marker Cathepsin-D. B') Schematic representation of B, black dots indicate Cathepsin-D labelling, orange indicate *M. leprae* and the blue lines indicate phagolysosomal membranes. C) Quantification of the average percentage of *M. leprae* present in the cytosol, error bar indicates standard deviation from 4 different patients, n represents the total number of intracellular bacteria (see also Fig. S2C).

Fig. 5.

Legend Fig. 5. Early cytosolic localization of *Mtb* in SCID mouse lungs. A,B)

Sections of SCID mice lung tissue infected with *Mtb* H37Rv for 21 days were labelled

for LAMP1 using immuno-gold labelling, to indicate lysosomes. A'B') schematic

representation with in orange *Mtb*, blue lines host membranes, black dots gold particles

indicating LAMP1 decorated lysosomes and green mitochondria. A) cross section of

Mtb surrounded by cellular membranes thus phagosomal. B) *Mtb* is present in the

cytosol, without cellular membranes surrounding the bacteria, scale bar indicates 200

nm. Information number of imaged bacteria and mice see Table S1.

Fig. 6.

Legend Fig. 6. *Mtb* preferentially localize to the cytosol in *Il1r1-/-* mice.

Fluorescence microscopy of 200 nm sections stained with DAPI for nuclei (white) and

anti-cell wall protein to detect *Mtb* (green) in granuloma in lung tissue of A) WT B6

mice and B) *Il1r1-/-* mice infected with *Mtb* for 28 days. C) Immuno-gold labelling using

LAMP1 indicate lysosomal membranes in *Il1r1-/-* mice imaged using TEM. C')

schematic representation of C, *Mtb* is depicted in orange, host membranes in blue and

host nucleus in pink. D) Quantification of the localization of *Mtb* in B6 and *Il1r1-/-* lungs,

here presenting cytosolic localization (see also Fig. S2D). The analysis included 897

(WT) or 618 (*Il1r1-/-*) bacteria and error bars indicate standard deviation based on in

multiple granulomas of 2 WT B6 and 2 *Il1r1-/-* mice.