The composition of the microbiota in the full-term fetal gut and amniotic fluid: a bovine caesarean section study

Aleksi Husso*, Leen Lietaer*, Tiina Pessa-Morikawa, Thomas Grönthal, Jan Govaere, Ann Van Soom, Antti Iivanainen**, Geert Opsomer**, Mikael Niku**

*First authors contributed equally

** supervised the study

Corresponding author: Mikael Niku, mikael.niku@helsinki.fi

Abstract

The fetal gut microbiome is being investigated for its putative effect on fetal development and maturation of the immune system. In this environment the low amounts of bacteria and bacterial DNA makes them highly challenging to accurately isolate and identify, and sensitive to contamination issues. The aims of the current study were to assess the bacterial load and characterize the bacterial composition of the amniotic fluid and meconium of full-term calves, leading to a better knowledge of prenatal bacterial seeding of the fetal intestine. Amniotic fluid and rectal meconium samples were collected during and immediately after elective caesarean section, performed in 25 Belgian Blue cow-calf couples. The samples were analyzed by qPCR, bacterial culture using GAM agar and 16S rRNA gene amplicon sequencing. To minimize the effects of contaminants, we included numerous technical controls and stringently filtered the 16S rRNA gene sequencing data to exclude putative contaminant sequences. The meconium samples contained a significantly higher amount of bacterial DNA than the negative controls and 5 of the samples contained culturable bacteria. In the amniotic fluid, the amount of bacterial DNA was not significantly different from the negative controls and no colonies were obtained

via cultures. Bacterial sequences were identified in both sample types and were primarily of phyla Proteobacteria, Firmicutes, Bacteroidetes and Actinobacteria, with some individual variation. We conclude that most calves encounter *in utero* maternal-fetal transmission of bacterial DNA, but the amount of bacterial DNA is low and viable bacteria rare.

Introduction

Characterizing the very first intestinal bacteria is essential for a better understanding of the co-development of the newborn and its intestinal microbiome. Host-microbiome interactions enable early-life education and maturation of the immune system during the so-called *window of opportunity*. However, the timing of the microbial colonization of the mammalian gut is still unclear (Perez-Munoz et al., 2017; Korpela and de Vos, 2018; Liu et al., 2019). Vertical as well as environmental transmission of bacteria occur during and after birth and seed the neonatal gastrointestinal (GI) tract (Funkhouser and Bordenstein, 2013; Korpela and de Vos, 2018). Multiple recent studies in humans and mice have reported the identification of microbial DNA or culturable bacteria in the mammalian fetal environment, while others interpret such observations as intrauterine infections or contamination (Aagaard et al., 2014; Lauder et al., 2016; Eisenhofer et al., 2018; de Goffau et al., 2019; Seferovic et al., 2019; Al Alam et al., 2020; Hockney et al., 2020; Olomu et al., 2020; Rackaityte et al., 2020; Theis et al., 2020a, 2020b). The newborn's GI tract is a low-microbial-biomass environment, implying multiple challenges for performing reliable microbiome analyses (Glassing et al., 2016; Eisenhofer et al., 2018; Stinson et al., 2018). Besides the technical limitations inherent to DNA sequencing, there are practical and ethical concerns while collecting samples in humans. First-pass meconium has been used as a proxy to assess the fetal intestinal microbiome, although samples are often collected hours after birth and even breastfeeding, suggesting potential postnatal effects on the microbial composition of the meconium (Collado et al., 2016; Liu et al., 2019;

Stinson et al., 2019). Mammalian animal models can overcome several of these ethical and practical difficulties, providing insights on the fetal gastrointestinal microbiome.

In the current study, the microbial composition of the full-term fetal gut and the corresponding amniotic fluid was assessed in Belgian Blue cow-calf couples. In this double-muscled beef breed, caesarean sections (C-sections) are performed on a routine basis during the very early stages of parturition, while fetal membranes are still intact, rendering this breed a highly suitable model for full-term gestation microbiome studies.

Besides the potential role of the cow model for research, there is also a significant interest in the composition and the development of the calf's intestinal microbiome. Gut health and growth performance during the first weeks of life are main drivers for cost effective livestock rearing (Thornton Philip, 2010; Lorenz et al., 2011). Few studies have been conducted on the intestinal microbiome in vaginally born neonatal calves. These have reported low numbers of bacteria, mostly belonging to the phyla Proteobacteria, Firmicutes, Actinobacteria and Bacteroidetes. Shared microbiota was found between calf meconium and the maternal vaginal vestibulum (Alipour et al., 2018; Yeoman et al., 2018; Klein-Jöbstl et al., 2019). To the best of our knowledge, no microbiome studies have been performed on calves born by C-section yet, and no studies are available to assess the association between the microbiome of the calf's meconium and the microbiome of the amniotic fluid.

Our aims were to assess the bacterial load and characterize the bacterial composition of the amniotic fluid and meconium of full-term calves. To this end, samples were collected during elective C-section, and analyzed by qPCR, bacterial culture under carefully controlled and validated conditions, and 16S rRNA gene amplicon sequencing. To minimize reagent and environmental bacterial DNA contaminants, we included numerous technical controls and

stringently filtered the 16S rRNA gene sequencing data to exclude putative contaminant sequences.

Materials and methods

All experimental procedures were approved by the institutional ethics and animal welfare committee of the Faculty of Veterinary Medicine (EC2018/002 - University of Ghent, Belgium). The cows' owners were informed about the study and gave their written consent.

Study design

The sampling of this study was performed at the teaching hospital of the Department of Reproduction, Obstetrics, and Herd Health of the Faculty of Veterinary Medicine in Ghent (Belgium), where pregnant Belgian Blue beef cows of different herds (parity between 1 and 5) were housed for an elective C-section.

In total 25 Belgian Blue cows and their calves (7 males and 18 females) were sampled from November 2017 until March 2019. The cows were housed in tie-stalls at the facility for 9.5 d ± 5.8 (mean ± standard deviation) prior to C-section and had ad libitum access to hay and water. During this period, rectal temperature was measured twice daily, and calving indicators such as udder distension, teat filling, pelvic ligament relaxation, vaginal discharge, vulvar edema, and behavioral changes were monitored every two hours by graduate veterinary students.

Prior to elective C-section, in cows that exhibited a drop in temperature, cervical dilation was assessed by manual palpation. In short, the vulvar region was cleaned with iodine soap and water. A gloved hand was inserted vaginally and the opening of the portio vaginalis cervicis was estimated. For this study, elective C-section was performed when the cow had a minimal cervical dilation of eight centimeters, with no rupture of the fetal membranes prior to surgery.

All cows were healthy according to their vital parameters (heart rate, temperature, respiratory rate), and there was no clinical evidence of intrauterine infection or contamination.

Sampling

Prior to surgery, the cows were restrained in a standing position in a cattle crush. Surgery was done as described by Kolkman et al. (2007). Briefly, the surgical area (left flank) was washed and disinfected, an abdominal incision was made, and part of the uterus was exteriorized for uterotomy. The allantoic sac was opened up to expose the intact amniotic sac. Amniotic fluid was aspirated through the amniotic membrane, using a sterile 16 G needle (Agani, Terumo Europe, Hamburg, Germany) and sterile 20 ml syringe (B. Braun, Melsungen, Germany). Within one hour after sampling, the retrieved volume was aliquoted, under a laminar-flow hood, into 2 sterile 15 ml tubes (188271, Cellstar, Greiner bio-one, Frickenhausen, Germany). In the first tube, 6 ml of amniotic fluid was dissolved in 3 ml of glycerol (≥99 %, G2025, Sigma-Aldrich (Merck), Overijse, Belgium) and subsequently stored at -80 °C to be used in culture experiments. In the second tube, 12 ml of amniotic fluid was stored at -80 °C to be used for bacterial DNA extraction.

Meconium samples were acquired directly from the calves' rectum, immediately after birth (no more than 30 minutes). Until the moment of sampling, calves laid on a clean, concrete floor, with access to neither the mother, nor colostrum. The perineum of the calf was dried with a clean paper towel and disinfected with 70% ethanol. A sterile double-guarded equine uterine culture swab (Har-vet, 17705, Spring Valley, USA; or Minitube, 17214/2950, Tiefenbach, Germany) was gently introduced in the rectum and the swab was exposed. Samples were taken in duplicate, one stored immediately at -80 °C with no additives, and the other stored at -80 °C in a sterile 2 ml cryovial containing 1 ml of a 30 % glycerol stock solution. This stock solution was prepared by diluting glycerol (≥99 %, G2025, Sigma-Aldrich (Merck), Overijse, Belgium),

in ultra-pure, nuclease-free water (W4502, Sigma-Aldrich (Merck), Overijse, Belgium) to a final 30 % concentration.

Negative field controls were processed in the surgery room, using the same sampling procedures and disposables. In total, 13 empty, sterile double-guarded equine uterine culture swabs (7 Har-vet swabs and 6 Minitube swabs) were included for the meconium sampling, 2 of the Har-vet swabs stored in 1ml of a 30 % glycerol stock solution, the others with no additives. Additionally, 12 negative controls were included for the amniotic fluid sampling, aspirating ultra-pure, nuclease-free water (W4502, Sigma-Aldrich (Merck), Overijse, Belgium) instead of amniotic fluid. For 8 of them, the ultra-pure, nuclease-free water was stored with no additives, and for 4 of them, 6 ml ultra-pure, nuclease-free water was dissolved in 3 ml of glycerol.

All samples were shipped on dry ice to the laboratory of the Department of Veterinary Biosciences of the Faculty of Veterinary Medicine in (Helsinki, Finland), for further processing.

Culture

Validation of the culture media

One of our aims was to define a single bacterial culture medium, capable of sustaining a majority of the calf intestinal core microbiota. Consequently, this single medium could be used for culturing our meconium and amniotic fluid samples, both low-biomass samples, avoiding further splitting or dilution of the samples. We first tested GAM "Nissui" (Gifu Anaerobic Medium Agar, Code 05420, HyServe, Germany), YCFA (Yeast extract, Casitone and Fatty Acid), LB (Lysogeny Broth, BD) and BB (Trypticase Soy Agar supplemented with Bovine Blood 211043, Tammer BioLab Oy, Tampere, Finland) media for the ability to sustain calf

intestinal core microbiota, by plating each type of medium anaerobically with feces of a healthy 7 day old calf (Lopez-Siles et al., 2012; Alipour et al., 2018). After 7 days of growth, the mixed culture was extracted from the plate, 16S rRNA gene amplicon sequenced, and compared to the core microbiota observed previously in fecal samples of the young calves (Alipour et al., 2018).

Sample culturing

Bacterial culture on GAM-agar plates was performed for 24 meconium samples (2 corresponding negative controls) and 24 amnion samples (4 negative controls). The frozen samples were first transferred to an anaerobic workstation (Ruskinn Concept Plus), mixed thoroughly and plated, using aseptic technique, at +37 °C. After this, the samples were transferred to a laminar flow cabinet and plated in aerobic conditions at +37 °C. The culture plates were checked for growth daily for 14 days. All visible bacterial colonies were subcultured until pure isolates were obtained. Fresh cultures were then identified using MALDI-TOF (Bruker Microflex LT) at the Central laboratory of the Faculty of Veterinary Medicine (Helsinki, Finland).

Samples were prepared using MALDI Biotyper MSP Identification Standard Method v 1.1. Mass spectra were analyzed in a mass/charge range from 2000 to 20 000 Da with MBT Compass v4.1 on flexControl v3.4 (Bruker Daltonik GmbH) using BDAL-7311 as the reference library. The Bruker Bacterial Test Standard (RUO) (Bruker Daltonik GmbH) was used for instrument calibration. If the identification confidence score was <2.00, further identification was done with 16S rRNA gene amplicon Sanger sequencing at the Institute of Biotechnology (University of Helsinki, Finland).

DNA extraction

DNA from the meconium samples (N = 25) and corresponding negative controls (N = 11) were extracted using ZymoBIOMICS DNA Miniprep Kit (Zymo Research, Irvine, CA, USA) according to the manufacturer's instructions, with minor modifications to the protocol as described previously (Alipour et al., 2018; Husso et al., 2020). ZymoBIOMICS™ Microbial Community Standard and an in-house fecal standard were processed with the meconium samples and no-template-controls were included in every batch.

For the amniotic fluid samples (N = 23) and the corresponding negative controls (N = 8), 2 ml of each sample was first centrifuged in a microcentrifuge at $16100 \times g$ 10 min in +4 °C. Most of the supernatant was removed and 750 µl of ZymoBIOMICS Lysis Solution and 19 µl of proteinase K (D3001-2-20/D3001-2-5, Zymo Research) were added to the remaining 200 µl of the samples and incubated for 30 min at +55 °C. After these steps, the same protocol as applied for the meconium samples was followed.

All manipulations of the tubes during the process were performed in a laminar flow cabinet, and the workplace, instruments and pipettes were cleaned routinely with 10 % bleach. Certified DNA, RNase, DNase and PCR inhibitor free tubes (STARLAB International, Germany) and Nuclease-free Water (Ambion, Thermo Fisher Scientific, USA) were used for DNA extraction and downstream analyses. All extracted DNA was stored at -80 °C.

Quantitative PCR

The bacterial 16S rRNA gene copy numbers in the meconium, amnion and negative control samples were determined using quantitative PCR. The analyses were performed as described previously (Alipour et al., 2018; Husso et al., 2020), with the exception that the PCR master

mix was first treated using a dsDNAse based decontamination kit (Enzo Life Sciences, Farmingdale, New York). The amplification was performed using the Bio-Rad CFX96 instrument (Bio-Rad, Hercules, California) and universal eubacteria probe and primers (Nadkarni et al., 2002). A standard series and negative controls were included in every run. The data were analyzed using the Bio-Rad CFX Maestro software. The results were normalized to the averages of absolute 16S rRNA gene copy numbers in the negative controls included in each run.

Library preparation and 16S rRNA gene amplicon sequencing

The V3-V4 region of the 16S rRNA gene amplicons was sequenced using the Illumina MiSeq platform in the DNA core facility of the University of Helsinki, as described previously (Alipour et al., 2018; Husso et al., 2020). In total, 23 amniotic fluid samples and 23 meconium samples of the same cow-calf couple were sequenced.

The ZymoBIOMICS Microbial Community Standard (Zymo Research, USA) and the in-house adult cow fecal standard were preamplified with 12 cycles and all other sample types, including negative controls, with 21 cycles.

The observed composition and abundances for the commercial standard matched the expected composition provided by the manufacturer (data not shown).

Bioinformatics

The detailed bioinformatics pipeline is described in the Supplementary materials. Briefly, the read quality was first inspected with FastQC and MultiQC (Andrews, 2011; Ewels et al., 2016). Leftover primes and spacers were then trimmed with Cutadapt v1.10 (Martin, 2011). A mapping file was created for QIIME2 and validated with Keemei (Rideout et al., 2016). The

FASTQ-files were imported to QIIME2 v2019.4, where the DADA2 plugin was used to denoise and quality filter the reads, call ASVs and generate a feature table (Callahan et al., 2016; Bolyen et al., 2019). A naïve Bayes classifier was trained in QIIME2 against SILVA v132 99 % database, extracted to only include the V3-V4 region and used to assign taxonomy to ASVs (Quast et al., 2013; Bokulich et al., 2018). Sequences derived from chloroplasts or mitochondria were removed and singletons were filtered out, leaving only bacteria with at least phylum-level identification.

The processed data was *in silico* filtered to remove ASVs which represented probable contaminants, as described previously (Husso et al., 2020). Briefly, an ASV was removed if its prevalence in actual samples was ≤2× its prevalence in instrument controls (DNA extracted from empty sampling instruments only handled aseptically in a laminar flood cabinet) or its prevalence in field controls (empty sampling instruments exposed to the surgery room environment), and if its mean relative abundance in actual samples was ≤10× its mean abundance in instrument controls or its mean abundance in field controls. The filtering was performed separately for meconium and amnion samples. If less than 500 reads remained after the decontamination, as was the case with six meconium samples, the sample was removed from further analyses.

Statistics

The 16S rRNA gene qPCR results from samples and negative controls were compared using two-tailed Mann-Whitney U test in IBM SPSS Statistics 25.

Shannon diversity indices were calculated for genus-level data with the R package phyloseq (McMurdie and Holmes, 2013). Kruskal-Wallis rank sum test was used to compare the bacterial community structure of both sample types in RStudio (R Studio Team, 2020).

The PCoA figures were plotted using ASV and genus-level data and Bray-Curtis distances with the R package phyloseq (McMurdie and Holmes, 2013). Permutational analysis of variance (PERMANOVA) and multivariate homogeneity of group dispersions (betadisper) were calculated using the R package vegan, using 9999 permutations (Oksanen et al., 2019).

DESeq2 was used to explore differentially abundant ASVS by calculating differential expression between sample groups (Love et al., 2014).

LEfSe (Linear discriminant analysis Effect Size) web application was used to determine taxons most likely to explain the differences between the sample types (Segata et al., 2011).

An ecologically organized heatmap of the top 40 most abundant ASVs was created with the R package phyloseq (Rajaram and Oono, 2010; McMurdie and Holmes, 2013).

Spearman correlations and their Bonferroni corrected p-values were calculated using R-package Hmisc 4.3.0 (Harrel, 2019).

Differences were considered significant at $P < 0.05$.

Results

Quantification of bacterial 16S rRNA in amniotic fluid and meconium samples by qPCR

The 16S rRNA gene copy numbers in amniotic fluid and meconium samples were assessed by qPCR (Fig 1). To minimize variation between qPCR runs, the copy numbers were normalized to the averages of the negative controls as explained in Methods.

In the amniotic fluid samples, there was no significant difference (P = 0.104) between the normalized 16S rRNA gene copy numbers of the samples (N = 23; mean = 2.4; SD = 2.1) and the negative controls (N = 8; mean = 1; SD = 0.53).

However, for the meconium samples there was a significant difference (P < 0.001) between the normalized 16S copy numbers of the samples (N = 25; mean = 9.9; SD = 27) and the negative controls (N = 11; mean = 1; SD = 0.86), indicating more bacterial DNA in the samples than in the negative controls.

Figure 1 Boxplots representing 16S rRNA gene copy numbers (fold change vs. negative controls) for amniotic fluid (n = 23, negative controls n = 8) and meconium samples (n = 25, negative controls n = 11). Samples were collected during elective C-section in Belgian Blue cow-calf pairs. The copy number

of each sample was normalized to the average copy number of the corresponding negative controls and log10 transformed. The midline of the box is the median, with the upper and lower limits of the box being the third and first quartile. The whiskers extend up to 1.5 times the interquartile range from the box to the furthest datapoint within that distance.

Decontamination of the 16S rRNA gene amplicon sequencing data

We performed a stringent filtering of the 16S rRNA gene sequencing data to remove the amplicon sequence variants (ASVs) potentially stemming from reagent contaminants and environmental bacterial DNA (Table 1).

The decontamination was based on the prevalence and relative abundance of each ASV in the samples and negative controls (see Methods).

Table 1 Mean read counts (standard deviation) detected in meconium and amniotic fluid samples and their negative controls. Only sequences with at least phylum-level identification were retained for further analyses. If less than 500 reads remained after in silico decontamination, as in six meconium samples, the sample was removed from further analyses.

Sample type	Raw		Processed		Decontaminated	
	N	Reads	N	Reads	N	Reads
Meconium	23	92266 (36235)	23	57082 (25543)	17	4189 (8931)
Meconium control	11	126800 (21164)	11	84023 (14585)	-	-
Amnion	23	127797 (22376)	23	83279 (15120)	23	3338 (2922)
Amnion control	6	136488 (12663)	6	90802 (8998)	-	-

Microbial composition of amniotic fluid and meconium samples

The microbiota of both sample types consisted primarily of Proteobacteria, Firmicutes, Bacteroidetes and Actinobacteria phyla, with individual variation (Fig 2). In both sample types,

the inter-individual variation increased at lower taxonomic levels (Supplementary Table 1 and Fig 5). The most abundant bacterial genera in meconium were *Delftia*, *Staphylococcus* and *Clostridium sensu stricto 1*, while the most prevalent genera were *Delftia*, *Acinetobacter*, unclassified *Burkholderiaceae*, *Staphylococcus* and *Corynebacterium 1* (Supplementary Table 1). In the amniotic fluid samples, *Staphylococcus*, *Streptococcus, Delftia, Sphingomonas* and *Enterococcus* were the most abundant genera, and *Delftia*, *Streptococcus*, *Staphylococcus, Sphingomonas* and *Acinetobacter* were the most prevalent genera (Supplementary Table 1). The observed composition and abundances for the commercial standard matched the expected composition provided by the manufacturer.

Figure 2 Microbiota composition in bovine meconium and amniotic fluid samples collected during elective C-section, adult feces, and a commercial community composition standard. The main colors indicate the bacterial phyla. Within phyla, the shades indicate bacterial genera. The lightest shade of each phylum shows the combined abundance of the least abundant genera (with a maximum of <0.5% of total).

The alpha diversity (Shannon index) at the genus level was not significantly different between the meconium and amniotic fluid samples (Fig 3, P = 0.889).

Figure 3 Shannon diversity index showing the bacterial community structure (at the genus level) of amniotic fluid and meconium samples, collected during elective C-section in Belgian Blue cow-calf pairs. Boxplot as in Fig 1.

Comparison of amniotic fluid and meconium microbiota composition

The microbiota composition was significantly different for amniotic fluid and meconium samples at the ASV level (PERMANOVA, $P < 0.001$, $R2 = 0.049$, Fig. 4 A), and at the genus level (PERMANOVA, $P < 0.001$, $R2 = 0.062$, Fig. 4 B).

Figure 4 Comparison of the bacterial community structure of amniotic fluid and meconium. 4 A: PCoA on Bray-Curtis distances based on ASV level data. Figure 4 B: PCoA on Bray- Curtis distances based on genus level data. Colors and shapes indicate the sample types.

A heatmap analysis of the 40 most abundant ASVs shows the difference between the sample types in more detail (Fig 5).

Figure 5 Ecologically organized heatmap (NMDS, Bray) of the 40 most abundant ASVs in bovine meconium and amniotic fluid samples collected during elective C-section, sorted by sample type. The taxon names are presented as genus level identifications. Color scale indicates relative abundance and is a log transformation with base 4.

According to the DESeq2 analysis, three ASVs were more abundant in meconium than in amniotic fluid: one *Staphylococcus* ASV (log2 Fold Change = 25.084, P_{adj} < 0.001), one *Rubrobacter* ASV (log2 Fold Change = 8.337, P_{adj} < 0.001), and one *Clostridium* ASV (log2 Fold Change = 8.055, P_{adj} = 0.004).

In amniotic fluid, one *Sphingomonas* ASV (log2 Fold Change = -8.420, P_{adj} < 0.001) and one *Staphylococcus* ASV (log2 Fold Change = -23.438, P_{adj} < 0.001) were more abundant than in meconium.

By LefSe (Linear discriminant analysis effect size) analysis with all taxonomic levels, meconium samples had a greater relative abundance of *Clostridiales* and *Rubrobacter* (LDA score (log10) > 4.0) than the amniotic fluid, but a lower relative abundance of Bacillales and *Corynebacteriales* (LDA score (log10) < -4.0).

Correlations between meconium and amniotic fluid from the same animal

We calculated Spearman rank correlations for the microbiota between meconium and amniotic fluid samples, taking into account ASVs that were present >100 times in all of the samples. Average correlations (ρ) at ASV (N = 17, ρ_{avg} = -0.0201, P_{avg} < 0.05) and genus level (N = 17, ρ_{avg} = 0.1685, P_{avg} < 0.05) were very weak and statistically not significant. All p-values were Bonferroni corrected for multiple testing.

Bacterial cultures

Based on our testing of multiple media, GAM (Gifu Anaerobic Medium) agar was chosen for the bacterial cultures, to assess if there were live bacteria in the samples stored in glycerol (see Methods). The GAM medium was able to sustain the largest amount of bacterial core genera found in calf feces, and additionally some unique genera not found on any other plate type (Alipour et al., 2018). All amniotic samples and their negative controls were culture negative, both in anaerobic and aerobic conditions. Five out of 24 meconium samples and 1 out of 2 negative controls were culture positive on GAM agar. The results for each sample are presented in detail in Table 2. Both gram negative and gram-positive strains were isolated from the samples.

Table 2 Culture positive meconium samples and identified bacteria. If the identification confidence score was <2.00 for MALDI-TOF, further identification was done with 16S rRNA gene amplicon Sanger sequencing to achieve genus level identification for all isolates.

Sample	Aerobic strains	Anaerobic strains
Negative control 1	*Staphylococcus* sp. (Gram+), *Kocuria* sp. (Gram+)	*Rothia* sp. (Gram+)
Meconium 5	*Roseomonas* sp. (Gram-)	-
Meconium 10	*Acinetobacter* sp. (Gram-)	*Streptococcus* sp. (Gram+), *Fusobacterium* sp. (Gram-)
Meconium 12	-	*Cutibacterium* sp. (Gram+)
Meconium 16	*Kocuria* sp. (Gram+)	-
Meconium 21	*Achromobacter* sp. (Gram-)	-

The Sanger sequence from the *Roseomonas* sp. isolate acquired from a meconium sample was a 100% identical match to the 16S rRNA gene amplicon sequencing data from the same sample. In addition, the Sanger sequences of *Staphylococcus* sp., *Kocuria* sp., *Roseomonas* sp.*,* and *Acinetobacter* sp. isolates matched the 16S rRNA gene amplicon sequencing data when compared to all meconium samples.

Three bacterial genera could be identified in the negative controls: *Rothia* sp., *Staphylococcus* sp. and *Kocuria* sp.. Of these, only *Kocuria* sp. was also identified from actual meconium samples.

Discussion

Vertical transmission of microbes during, and perhaps already before birth is a universal phenomenon across the whole animal kingdom and plays an important role in shaping host immunity and fitness (Funkhouser and Bordenstein, 2013; Vandenplas et al., 2020). Despite the advances in our understanding of the assembly and lifelong effects of the perinatal microbiota, the in utero exposure of the fetus to maternal bacteria and bacterial components remains poorly understood (Perez-Munoz et al., 2017; Korpela and de Vos, 2018; Liu et al., 2019).

In the present study, we sampled calves immediately after they were born by elective C-section. The elective C-section is a routine procedure in Belgian Blue beef cattle, since the pelvis of the dam does not allow vaginal birth of the relatively large double-muscled calf (Kolkman et al., 2007). We performed the surgery during the very early stages of parturition, when both fetal membranes (amniotic and allantochorionic membrane) were still intact, allowing us to sample the full-term fetus and the intrauterine environment without exposure to the vaginal microbes and minimal exposure to the environmental microbiota. Meconium samples were collected rectally, immediately after birth. The relatively large size of the calves permits the insertion of a double-guarded sampling device into the rectum, minimizing the risk of contamination.

In contrast, in most human studies, first-pass meconium is collected from diapers (Hansen et al., 2015; Collado et al., 2016; Liu et al., 2019; Stinson et al., 2019). Because the first-pass meconium is often collected several hours after birth, microbes from the environment and the mother (especially during breastfeeding) have time to colonize the intestine. This usually impedes the use of human meconium samples to research the potential intrauterine transmission of bacteria to the neonatal gastrointestinal tract (Dornelles et al., 2020). In the

current study, amniotic fluid samples were collected *in vivo*, while the amniotic membrane was still intact. Also in humans, amniotic fluid has been sampled in women during pre-labor C-section (Collado et al., 2016).

Bacterial culture is traditionally used to determine the viable population in a bacterial community. However, standard culture-dependent techniques are known to be inadequate to identify all bacteria. Capturing a broad bacterial diversity would require the use of many different selective media and multiple specific growing conditions (Lau et al., 2016; Lagier et al., 2018). In the current study, one of our aims was to define a single bacterial culture medium, capable of sustaining a majority of the calf intestinal core microbiota. We first tested four different culture media: GAM "Nissui" (Gifu Anaerobic Medium Agar), mostly used for anaerobic bacteria, YCFA (Yeast extract, Casitone and Fatty Acid), containing volatile fatty acids typically supporting the growth of several gut bacteria, LB (Lysogeny Broth) as a general media, and BB (Trypticase Soy Agar supplemented with Bovine Blood) supporting fastidious bacteria with blood enrichment. The GAM medium was found the most optimal for both aerobic and anaerobic culturing, although unprobable to support the entire bacterial diversity in the meconium and amniotic fluid samples.

Bacteria were successfully cultured from 5 out of 24 meconium samples, representing both gram-negative and gram-positive bacterial genera, known to live in mammalian hosts. *Roseomonas* and *Achromobacter* species have been isolated from a wide variety of environmental sources but are also known to act as opportunistic pathogens (Reverdy et al., 1984; Rihs et al., 1993). Some *Acinetobacter* and *Streptococcus* species have non-exclusively been associated with mammals (Krzyściak et al., 2013; Touchon et al., 2014). *Fusobacteria* species are typically closely related to mucous membranes, especially the oral cavities, while *Cutibacterium* and *Kocuria* species have been isolated mostly from the skin (Hofstad, 1992;

Savini et al., 2010; Scholz and Kilian, 2016). One of our negative controls yielded *Kocuria* colonies. This was the only taxon shared between the actual meconium samples and the negative controls, suggesting low-level contamination.

Although we detected cultivable bacteria in some of the meconium samples, most of the samples were culture negative. This indicates that the presence of live bacteria in the fetal intestine is rare, in contrast to the significant amount of bacterial DNA. Individual differences in amounts of viable bacteria retrieved from meconium were also described in a recent dog study. Interestingly, puppies born without detectable meconium microbiota were shown to have a slower growth rate than those in which meconium microbiota were detected (Pipan et al., 2020).

Both the mucosal surface of the intestine as well as the amniotic fluid contain bacteriostatic substances, such as lactoferrin and salivary scavenger and agglutinin (SALSA), which are able to suppress bacterial growth and viability (Reichhardt and Meri, 2016; Lisowska-Myjak et al., 2019). In human meconium, SALSA amounts to 10 % of all proteins, highlighting its potential role in antimicrobial defense (Reichhardt et al., 2014). Lactoferrin, SALSA and other similar substances may explain the small number of viable bacteria observed in our study, while still being able to find a significant amount of bacterial DNA relative to the negative controls.

The culture-independent techniques were performed based on 16S rRNA. Since contamination is a major challenge in the analysis of low-bacterial biomass samples, we processed several types of negative controls alongside our biological samples and applied rigorous *in silico* filtering of potential contaminants (Glassing et al., 2016; Eisenhofer et al., 2018). Many similar studies are published without the use of any or with a very limited number of contamination controls, which makes our approach more reliable, especially for assessing low-bacterial biomass environments.

We observed a small but significant amount of bacterial DNA in the meconium samples by 16S qPCR, suggesting a prenatal transmission of bacterial DNA to the fetal intestine. This is consistent with previous studies in cattle, in which calves born at term *per vias naturales*, were sampled (Mayer et al., 2012; Alipour et al., 2018; Klein-Jöbstl et al., 2019). A profile dominated by the phyla Proteobacteria, Firmicutes, Bacteroidetes and Actinobacteria was identified in the meconium, in agreement with previous studies with 16s rRNA gene amplicon sequencing (Alipour et al., 2018; Yeoman et al., 2018; Klein-Jöbstl et al., 2019; Husso et al., 2020). Additionally, some of the most prevalent genera observed in meconium, such as *Sphingomonas*, *Acinetobacter*, *Pseudomonas*, *Ruminococcus* and *Stenotrophomonas* have also been previously described in neonatal calves (Klein-Jöbstl et al., 2019).

In contrast to the meconium, amniotic fluid samples were not distinguishable from negative controls by 16S qPCR and were all culture negative. A recent caesarean section study in sheep stated amniotic fluid sterility during the third trimester of pregnancy (Malmuthuge and Griebel, 2018). In another study however, a low-biomass microbiome in the pregnant bovine uterus was described (Alipour et al., 2018).

Using 16S rRNA gene amplicon sequencing, we found a bacterial signature in the amniotic fluid, dominated by the phyla Proteobacteria, Firmicutes, Bacteroidetes and Actinobacteria, in accordance with previous studies (Karstrup et al., 2017; Moore et al., 2017). Deeper taxonomic identification, up to the family or genus level, revealed no further agreements with these studies.

From a comparative point of view between mammalian species, the amniotic fluid sampled by Collado et al. (2016) in women during pre-labor C-section was found to contain a distinct, low diversity, low-biomass microbiome, predominated by Proteobacteria (*Enterobacteriaceae*). Despite this and other recent studies, the presence of microbial DNA and culturable bacteria in the human placenta and fetal environment is still under debate (de Goffau et al., 2019).

Comparing the results of the 16S rRNA gene sequencing for meconium and amniotic fluid, the compositions were similar at the phylum. However, they presented a significantly different microbial composition at ASV level: genera *Staphylococcus*, *Rubrobacter* and *Clostridium* were relatively more abundant in meconium samples than in amniotic fluid samples according to DESeq2 analysis. Complementing LefSe analysis using all taxonomic levels also found *Clostridiales* and *Rubrobacter* taxons relatively more abundant in the meconium.

Very weak and statistically non-significant correlations were found between the amniotic fluid and meconium of the correspondent calf in the current study, conversely to earlier results in humans (Collado et al., 2016; He et al., 2020). In ruminants, a synepitheliochorial placentation occurs, containing triple fetal membranes: the fetus is enclosed by the allantoic membrane, in addition to chorion and amnion. These layers form a distinct barrier between the maternal endometrium and the amniotic cavity, holding the fetus. Small-molecular microbial metabolites likely still penetrate the placenta, whereas larger components and possibly even live bacteria could be translocated by leukocytes (Perez et al., 2007; Gomez de Agüero et al., 2016).

The amniotic fluid is an aggregation of fetal excretions and epithelial cells, continuously swallowed by the fetus (Bongso and Basrur, 1976; Rossi et al., 2014). Along the fetal GI tract, this fluid is accumulated, concentrated and retained to form the meconium, generally only expelled after birth. The process of accumulation and concentration may explain the observed difference in bacterial composition between the meconium and amniotic fluid. The extremely

low levels of bacteria in the amniotic fluid probably decrease the accuracy of analysis. When concentrated in the meconium, the amount of bacterial DNA could however reach the limit of detection and be clearly separated from the negative controls.

It is also possible that the microbial seeding occurs hematogenously via the placental and umbilical blood vessels to the fetal internal organs, surpassing the amniotic cavity. In general, the limited taxonomic resolution of 16S gene amplicon sequencing does not allow actual tracing of bacteria between samples.

Bacteria can enter the uterus ascending from the vagina, migration via the fallopian tubes or hematogenous spread (Gonçalves et al., 2002; Jeon et al., 2017; Baker et al., 2018; Romero et al., 2019; Brunham and Paavonen, 2020; Cappelletti et al., 2020). These bacteria or bacterial metabolites can reach the fetal membranes, allantoic and amniotic fluid or seed the fetus via transplacental migration, although this relocation is limited by the barrier imposed by the triple fetal membranes. However, it is unclear how microbes subsequently reach the fetal intestine.

Although the mechanics are still unknown, dam to fetus transmission of bacteria has been observed in mice: a specifically labelled *Enterococcus faecium* strain orally inoculated in pregnant mice, could be isolated in the meconium of their pre-labour, offspring delivered by C-section (Jimenez et al., 2008). However, evidence of vertical transfer of probiotic bacterial strains from mother to infant during pregnancy remains inconclusive (Moore et al., 2020). With the increasing prenatal and intrapartum use of antibiotics in humans, potential alterations in the microbiota of the offspring still continues to be of high interest (Dierikx et al., 2020).

In conclusion, we found that the meconium of full-term calves delivered by elective caesarean section in most cases contains a small amount of diverse bacterial DNA and possibly rare culturable bacteria. In the amniotic fluid, bacteria were not observed by 16S qPCR or culturing, but a microbial DNA profile was distinguishable from controls by amplicon sequencing. Based

on these results, calves encounter *in utero* maternal-fetal transmission of bacterial components, but the prenatal acquisition of live bacteria is likely not physiologically significant.

Data availability

The raw 16S rRNA gene amplicon sequencing dataset is fully available in the NCBI Sequence Read Archive (SRA) with the accession number PRJNA643145 after journal acceptance.

Acknowledgements

We thank the owners of the cows for allowing sampling of their animals, and all clinicians and graduate veterinary students for helping with the monitoring of the animals, performing the surgery and technical assistance during sampling. Additionally, we thank Kirsi Lahti for expert technical assistance in the laboratory.

Author contributions

MN, LL, AI and GO contributed to conception and design of the study. AH performed the bioinformatics, statistical analyses and bacterial culture. AH and TG performed MALDI-TOF analyses. LL and JG organized the sampling and collected the samples. AH and LL wrote the manuscript. All authors contributed to manuscript revision, read, and approved the submitted version

Ethics declaration

All experimental procedures were approved by the institutional ethics and animal welfare committee of the Faculty of Veterinary Medicine (EC2018/002 - University of Ghent,

Belgium). The cows' owners were informed about the study and gave their written consent for their cows' participation.

References

Aagaard, K., Ma, J., Antony, K. M., Ganu, R., Petrosino, J., and Versalovic, J. (2014). The placenta harbors a unique microbiome. *Sci Transl Med* 6, 237ra65. doi:10.1126/scitranslmed.3008599.

Al Alam, D., Danopoulos, S., Grubbs, B., Ali, N. A. B. M., MacAogain, M., Chotirmall, S. H., et al. (2020). Human Fetal Lungs Harbor a Microbiome Signature. *Am. J. Respir. Crit. Care Med.* doi:10.1164/rccm.201911-2127LE.

Alipour, M. J., Jalanka, J., Pessa-Morikawa, T., Kokkonen, T., Satokari, R., Hynönen, U., et al. (2018). The composition of the perinatal intestinal microbiota in cattle. *Sci. Rep.* 8, 10437. doi:10.1038/s41598-018-28733-y.

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.

Baker, J. M., Chase, D. M., and Herbst-Kralovetz, M. M. (2018). Uterine Microbiota: Residents, Tourists, or Invaders? *Front. Immunol.* 9. doi:10.3389/fimmu.2018.00208.

Bokulich, N. A., Kaehler, B. D., Rideout, J. R., Dillon, M., Bolyen, E., Knight, R., et al. (2018). Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* 6, 90. doi:10.1186/s40168-018-0470-z.

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., et al. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* 37, 852–857. doi:10.1038/s41587-019-0209-9.

Bongso, T. A., and Basrur, P. K. (1976). Foetal fluids in cattle. *Can. Vet. J.* 17, 38.

Brunham, R. C., and Paavonen, J. (2020). Reproductive system infections in women: lower genital tract syndromes. *Pathog. Dis.*

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* 13, 581. doi:10.1038/nmeth.3869.

Cappelletti, M., Presicce, P., and Kallapur, S. G. (2020). Immunobiology of Acute Chorioamnionitis. *Front. Immunol.* 11.

Collado, M. C., Rautava, S., Aakko, J., Isolauri, E., and Salminen, S. (2016). Human gut colonisation may be initiated in utero by distinct microbial communities in the placenta and amniotic fluid. *Sci Rep* 6, 23129. doi:10.1038/srep23129.

de Goffau, M. C., Lager, S., Sovio, U., Gaccioli, F., Cook, E., Peacock, S. J., et al. (2019). Human placenta has no microbiome but can contain potential pathogens. *Nature*. doi:10.1038/s41586-019-1451-5.

Dierikx, T. H., Visser, D. H., Benninga, M. A., van Kaam, A., de Boer, N. K. H., de Vries, R., et al. (2020). The influence of prenatal and intrapartum antibiotics on intestinal microbiota colonisation in infants: A systematic review. *J. Infect.*

Dornelles, L. V., Procianoy, R. S., Roesch, L. F., Corso, A. L., Dobbler, P. T., Mai, V., et al. (2020). Meconium microbiota predicts clinical early-onset neonatal sepsis in preterm neonates. *J. Matern. Fetal Neonatal Med.*, 1–9.

Eisenhofer, R., Minich, J. J., Marotz, C., Cooper, A., Knight, R., and Weyrich, L. S. (2018). Contamination in Low Microbial Biomass Microbiome Studies: Issues and Recommendations. *Trends Microbiol.* doi:10.1016/j.tim.2018.11.003.

Ewels, P., Magnusson, M., Lundin, S., and Kaller, M. (2016). MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* 32, 3047–8. doi:10.1093/bioinformatics/btw354.

Funkhouser, L. J., and Bordenstein, S. R. (2013). Mom knows best: the universality of maternal microbial transmission. *PLoS Biol* 11, e1001631. doi:10.1371/journal.pbio.1001631.

Glassing, A., Dowd, S. E., Galandiuk, S., Davis, B., and Chiodini, R. J. (2016). Inherent bacterial DNA contamination of extraction and sequencing reagents may affect interpretation of microbiota in low bacterial biomass samples. *Gut Pathog.* 8, 24.

Gomez de Agüero, M., Ganal-Vonarburg, S. C., Fuhrer, T., Rupp, S., Uchimura, Y., Li, H., et al. (2016). The maternal microbiota drives early postnatal innate immune development. *Science* 351, 1296. doi:10.1126/science.aad2571.

Gonçalves, L. F., Chaiworapongsa, T., and Romero, R. (2002). Intrauterine infection and prematurity. *Ment. Retard. Dev. Disabil. Res. Rev.* 8, 3–13.

Hansen, R., Scott, K. P., Khan, S., Martin, J. C., Berry, S. H., Stevenson, M., et al. (2015). First-pass meconium samples from healthy term vaginally-delivered neonates: an analysis of the microbiota. *PloS One* 10, e0133320.

Harrel, F. (2019). *Hmisc: Harrell Miscellaneous*. Available at: https://CRAN.R-project.org/package=Hmisc.

He, Q., Kwok, L. Y., Xi, X., Zhong, Z., Ma, T., Xu, H., ... & Zhang, H. (2020). The meconium microbiota shares more features with the amniotic fluid microbiota than the maternal fecal and vaginal microbiota. *Gut Microbes*, *12*(1), 1794266.

Hockney, R., Waring, G. J., Taylor, G., Cummings, S. P., Robson, S. C., Orr, C. H., et al. (2020). Fetal membrane bacterial load is increased in histologically confirmed inflammatory chorioamnionitis: A retrospective cohort study. *Placenta* 91, 43–51. doi:10.1016/j.placenta.2020.01.006.

Hofstad, T. (1992). "The genus Fusobacterium," in *The prokaryotes* (Springer), 4114–4126.

Husso, A., Jalanka, J., Alipour, M. J., Huhti, P., Kareskoski, M., Pessa-Morikawa, T., et al. (2020). The composition of the perinatal intestinal microbiota in horse. *Sci. Rep.* 10, 1–12.

Oksanen J, F. G. B., Michael Friendly, Roeland Kindt, Pierre Legendre, Dan McGlinn, Peter R. Minchin, R. B. O'Hara, Gavin L. Simpson, Peter Solymos, M. Henry, H. Stevens, Eduard Szoecs and Helene Wagner (2019). vegan: Community Ecology Package. R package version

2.5-4. Available at: https://CRAN.R-project.org/package=vegan.

Jeon, S. J., Cunha, F., Vieira-Neto, A., Bicalho, R. C., Lima, S., Bicalho, M. L., et al. (2017). Blood as a route of transmission of uterine pathogens from the gut to the uterus in cows. *Microbiome* 5, 1–13.

Jimenez, E., Marin, M. L., Martin, R., Odriozola, J. M., Olivares, M., Xaus, J., et al. (2008). Is meconium from healthy newborns actually sterile? *Res Microbiol* 159, 187–93. doi:10.1016/j.resmic.2007.12.007.

Karstrup, C. C., Klitgaard, K., Jensen, T. K., Agerholm, J. S., and Pedersen, H. G. (2017). Presence of bacteria in the endometrium and placentomes of pregnant cows. *Theriogenology* 99, 41–47. doi:10.1016/j.theriogenology.2017.05.013.

Klein-Jöbstl, D., Quijada, N. M., Dzieciol, M., Feldbacher, B., Wagner, M., Drillich, M., et al. (2019). Microbiota of newborn calves and their mothers reveals possible transfer routes for newborn calves' gastrointestinal microbiota. *PloS One* 14.

Kolkman, I., De Vliegher, S., Hoflack, G., Van Aert, M., Laureyns, J., Lips, D., et al. (2007). Protocol of the caesarean section as performed in daily bovine practice in Belgium. *Reprod. Domest. Anim.* 42, 583–589.

Korpela, K., and de Vos, W. M. (2018). Early life colonization of the human gut: microbes matter everywhere. *Curr. Opin. Microbiol.* 44, 70–78.

Krzyściak, W., Pluskwa, K. K., Jurczak, A., and Kościelniak, D. (2013). The pathogenicity of the Streptococcus genus. *Eur. J. Clin. Microbiol. Infect. Dis.* 32, 1361–1376.

Lagier, J.-C., Dubourg, G., Million, M., Cadoret, F., Bilen, M., Fenollar, F., et al. (2018). Culturing the human microbiota and culturomics. *Nat. Rev. Microbiol.* doi:10.1038/s41579-018-0041-0.

Lau, J. T., Whelan, F. J., Herath, I., Lee, C. H., Collins, S. M., Bercik, P., et al. (2016). Capturing the diversity of the human gut microbiota through culture-enriched molecular profiling. *Genome Med* 8, 72. doi:10.1186/s13073-016-0327-7.

Lauder, A. P., Roche, A. M., Sherrill-Mix, S., Bailey, A., Laughlin, A. L., Bittinger, K., et al. (2016). Comparison of placenta samples with contamination controls does not provide evidence for a distinct placenta microbiota. *Microbiome* 4, 29. doi:10.1186/s40168-016-0172-3.

Lisowska-Myjak, B., Skarżyńska, E., Wojdan, K., and Nasierowska-Guttmejer, A. (2019). Protein and peptide profiles in neonatal meconium. *J. Obstet. Gynaecol. Res.* 45, 556–564.

Liu, C.-J., Liang, X., Niu, Z.-Y., Jin, Q., Zeng, X.-Q., Wang, W.-X., et al. (2019). Is the delivery mode a critical factor for the microbial communities in the meconium? *EBioMedicine* 49, 354–363. doi:10.1016/j.ebiom.2019.10.045.

Lopez-Siles, M., Khan, T. M., Duncan, S. H., Harmsen, H. J., Garcia-Gil, L. J., and Flint, H. J. (2012). Cultured representatives of two major phylogroups of human colonic Faecalibacterium prausnitzii can utilize pectin, uronic acids, and host-derived substrates for growth. *Appl Env. Microbiol* 78, 420–428.

Lorenz, I., Fagan, J., and More, S. J. (2011). Calf health from birth to weaning. II. Management of diarrhoea in pre-weaned calves. *Ir. Vet. J.* 64, 9.

Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550.

Malmuthuge, N., and Griebel, P. J. (2018). Fetal environment and fetal intestine are sterile during the third trimester of pregnancy. *Vet. Immunol. Immunopathol.* 204, 59–64.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 17. doi:10.14806/ej.17.1.200.

Mayer, M., Abenthum, A., Matthes, J. M., Kleeberger, D., Ege, M. J., Hölzel, C., et al. (2012). Development and genetic influence of the rectal bacterial flora of newborn calves. *Vet. Microbiol.* 161, 179–185. doi:10.1016/j.vetmic.2012.07.023.

McMurdie, P. J., and Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLOS ONE* 8, e61217. doi:10.1371/journal.pone.0061217.

Moore, R. L., Geraghty, A. A., Feehily, C., Saldova, R., Murphy, E. F., Van Sinderen, D., et al. (2020). Can a probiotic supplement in pregnancy result in transfer to the neonatal gut: A systematic review. *Acta Obstet. Gynecol. Scand.*

Moore, S. G., Ericsson, A. C., Poock, S. E., Melendez, P., and Lucy, M. C. (2017). Hot topic: 16S rRNA gene sequencing reveals the microbiome of the virgin and pregnant bovine uterus. *J. Dairy Sci.* 100, 4953–4960.

Nadkarni, M. A., Martin, F. E., Jacques, N. A., and Hunter, N. %J M. (2002). Determination of bacterial load by real-time PCR using a broad-range (universal) probe and primers set. 148, 257–266.

Olomu, I. N., Pena-Cortes, L. C., Long, R. A., Vyas, A., Krichevskiy, O., Luellwitz, R., et al. (2020). Elimination of "kitome" and "splashome" contamination results in lack of detection of a unique placental microbiome. *BMC Microbiol.* 20, 1–19.

Perez, P. F., Doré, J., Leclerc, M., Levenez, F., Benyacoub, J., Serrant, P., et al. (2007). Bacterial imprinting of the neonatal immune system: lessons from maternal cells? *Pediatrics* 119, e724–e732.

Perez-Munoz, M. E., Arrieta, M. C., Ramer-Tait, A. E., and Walter, J. (2017). A critical assessment of the "sterile womb" and "in utero colonization" hypotheses: implications for research on the pioneer infant microbiome. *Microbiome* 5, 48. doi:10.1186/s40168-017-0268-4.

Pipan, M. Z., Kajdič, L., Kalin, A., Plavec, T., and Zdovc, I. (2020). Do newborn puppies have their own microbiota at birth? Influence of type of birth on newborn puppy microbiota. *Theriogenology*.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* 41, D590-6. doi:10.1093/nar/gks1219.

R Studio Team (2020). RStudio: Integrated Development for R. *RStudio Support*. Available at: http://support.rstudio.com/hc/en-us/articles/206212048

Rackaityte, E., Halkias, J., Fukui, E. M., Mendoza, V. F., Hayzelden, C., Crawford, E. D., et al.

(2020). Viable bacterial colonization is highly limited in the human intestine in utero. *Nat. Med.* 26, 599–607.

Rajaram, S., and Oono, Y. (2010). NeatMap-non-clustering heat map alternatives in R. *BMC Bioinformatics* 11, 1–9.

Reichhardt, M. P., Jarva, H., de Been, M., Rodriguez, J. M., Quintana, E. J., Loimaranta, V., et al. (2014). The salivary scavenger and agglutinin in early life: diverse roles in amniotic fluid and in the infant intestine. *J. Immunol.* 193, 5240–5248.

Reichhardt, M. P., and Meri, S. (2016). SALSA: a regulator of the early steps of complement activation on mucosal surfaces. *Front. Immunol.* 7, 85.

Reverdy, M. E., Freney, J., Fleurette, J., Coulet, M., Surgot, M., Marmet, D., et al. (1984). Nosocomial colonization and infection by Achromobacter xylosoxidans. *J. Clin. Microbiol.* 19, 140–143.

Rideout, J. R., Chase, J. H., Bolyen, E., Ackermann, G., Gonzalez, A., Knight, R., et al. (2016). Keemei: cloud-based validation of tabular bioinformatics file formats in Google Sheets. *Gigascience* 5, 27. doi:10.1186/s13742-016-0133-6.

Rihs, J. D., Brenner, D. J., Weaver, R. E., Steigerwalt, A. G., Hollis, D. G., and Yu, V. L. (1993). Roseomonas, a new genus associated with bacteremia and other human infections. *J. Clin. Microbiol.* 31, 3275–3283.

Romero, R., Gomez-Lopez, N., Winters, A. D., Jung, E., Shaman, M., Bieda, J., et al. (2019). Evidence that intra-amniotic infections are often the result of an ascending invasion–a molecular microbiological study. *J. Perinat. Med.* 47, 915–931.

Rossi, B., Merlo, B., Colleoni, S., Iacono, E., Tazzari, P. L., Ricci, F., et al. (2014). Isolation and in vitro characterization of bovine amniotic fluid derived stem cells at different trimesters of pregnancy. *Stem Cell Rev. Rep.* 10, 712–724.

Savini, V., Catavitello, C., Masciarelli, G., Astolfi, D., Balbinot, A., Bianco, A., et al. (2010). Drug sensitivity and clinical impact of members of the genus Kocuria. *J. Med. Microbiol.* 59, 1395–1402.

Scholz, C. F., and Kilian, M. (2016). The natural history of cutaneous propionibacteria, and reclassification of selected species within the genus Propionibacterium to the proposed novel genera Acidipropionibacterium gen. nov., Cutibacterium gen. nov. and Pseudopropionibacterium gen. nov. *Int. J. Syst. Evol. Microbiol.* 66, 4422–4432.

Seferovic, M. D., Pace, R. M., Carroll, M., Belfort, B., Major, A. M., Chu, D. M., et al. (2019). Visualization of microbes by 16S in situ hybridization in term and preterm placentas without intraamniotic infection. *Am. J. Obstet. Gynecol.* 221, 146.e1-146.e23. doi:10.1016/j.ajog.2019.04.036.

Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W. S., et al. (2011). Metagenomic biomarker discovery and explanation. *Genome Biol.* 12, 1–18.

Stinson, L. F., Boyce, M. C., Payne, M. S., and Keelan, J. A. (2019). The not-so-sterile womb: evidence that the human fetus is exposed to bacteria prior to birth. *Front. Microbiol.* 10, 1124.

Stinson, L. F., Keelan, J. A., and Payne, M. S. (2018). Comparison of Meconium DNA Extraction Methods for Use in Microbiome Studies. *Front Microbiol* 9, 270. doi:10.3389/fmicb.2018.00270.

Theis, K. R., Romero, R., Greenberg, J. M., Winters, A. D., Garcia-Flores, V., Motomura, K., et al. (2020a). No Consistent Evidence for Microbiota in Murine Placental and Fetal Tissues. *mSphere* 5, e00933-19. doi:10.1128/mSphere.00933-19.

Theis, K. R., Romero, R., Winters, A. D., Jobe, A. H., and Gomez-Lopez, N. (2020b). Lack of evidence for microbiota in the placental and fetal tissues of rhesus macaques. *mSphere* 5.

Thornton Philip, K. (2010). Livestock Production: Recent Trends, Future Prospects. Nairobi Kenya, doi: 10.1098/rstb. 2010.0134 Phil. *Trans R Soc B 27 Sept. 2010 Vol 365 No 1554 2853* 2867.

Touchon, M., Cury, J., Yoon, E.-J., Krizova, L., Cerqueira, G. C., Murphy, C., et al. (2014). The genomic diversification of the whole Acinetobacter genus: origins, mechanisms, and consequences. *Genome Biol. Evol.* 6, 2866–2882.

Vandenplas, Y., Carnielli, V. P., Ksiazyk, J., Luna, M. S., Migacheva, N., Mosselmans, J. M., et al. (2020). Factors affecting early-life intestinal microbiota development. *Nutrition*, 110812.

Yeoman, C. J., Ishaq, S. L., Bichi, E., Olivo, S. K., Lowe, J., and Aldridge, B. M. (2018). Biogeographical Differences in the Influence of Maternal Microbial Sources on the Early Successional Development of the Bovine Neonatal Gastrointestinal tract. *Sci Rep* 8, 3197. doi:10.1038/s41598-018-21440-8.