The composition of the microbiota in the full-term fetal gut and amniotic fluid: a bovine caesarean section study

Aleksi Husso*, Leen Lietaer*, Tiina Pessa-Morikawa, Thomas Grönthal, Jan Govaere, Ann Van Soom, Antti Iivanainen**, Geert Opsomer**, Mikael Niku**

*First authors contributed equally

** supervised the study

Corresponding author: Mikael Niku, mikael.niku@helsinki.fi

Abstract

The fetal development of the intestinal immune system is stimulated by the maternal microbiota, but it is still unclear whether viable bacteria exist in the healthy fetus. Analysis of such low microbial biomass environments are challenging due to contamination issues. The aims of the current study were to assess the bacterial load and characterize the bacterial composition of the amniotic fluid and meconium of full-term calves, leading to a better knowledge of prenatal bacterial seeding of the fetal intestine. Amniotic fluid and rectal meconium samples were collected during and immediately after elective caesarean section, performed in 25 Belgian Blue cow-calf couples. The samples were analyzed by qPCR, bacterial culture using GAM agar and 16S rRNA gene amplicon sequencing. To minimize the effects of contaminants, we included multiple technical controls and stringently filtered the 16S rRNA gene sequencing data to exclude putative contaminant sequences. The meconium samples contained a significantly higher amount of bacterial DNA than the negative controls and 5 of 24 samples contained culturable bacteria. In the amniotic fluid, the amount of bacterial DNA was not significantly different from the negative controls and all samples were culture negative.

25   Bacterial sequences were identified in both sample types and were primarily of phyla

26   Proteobacteria, Firmicutes, Bacteroidetes and Actinobacteria, with some individual variation.

27   We conclude that most calves encounter *in utero* maternal-fetal transmission of bacterial DNA,

28   but the amount of bacterial DNA is low and viable bacteria are rare.

Introduction

Characterizing the very first intestinal bacteria is essential for a better understanding of the co-

development of the newborn and its intestinal microbiome. Host-microbiome interactions

enable early-life education and maturation of the immune system during a *window of*

opportunity. The use of prenatal and intrapartum antibiotics may disturb this process (Dierikx

et al., 2020). However, the timing of the microbial colonization of the mammalian gut is still

unclear (Perez-Munoz et al., 2017; Korpela and de Vos, 2018; Liu et al., 2019; Guzman et al.,

2020). Vertical as well as environmental transmission of bacteria occur during and after birth

and seed the neonatal gastrointestinal (GI) tract (Funkhouser and Bordenstein, 2013; Korpela

and de Vos, 2018). Transmission of an orally inoculated *Enterococcus faecium* strain in

pregnant mice to the meconium of their fetuses has been described (Jimenez et al., 2008).

Recent studies in humans, mice and cattle have reported the identification of microbial DNA

or culturable bacteria in the mammalian fetal environment (Aagaard et al., 2014; Severovic et

al., 2019; Al Alam et al., 2020; Rackaityte et al., 2020; Guzman et al., 2020). Nevertheless,

others interpret such observations as intrauterine infections or contamination (Lauder et al.,

2016; Eisenhofer et al., 2018; de Goffau et al., 2019; Hockney et al., 2020; Olomu et al., 2020;

Theis et al., 2020a, 2020b). The newborn's GI tract is a low microbial biomass environment,

implying multiple challenges for performing reliable microbiome analyses (Glassing et al.,

2016; Eisenhofer et al., 2018; Stinson et al., 2018). Besides the technical limitations inherent

to DNA sequencing, there are practical and ethical concerns while collecting samples in

humans. First-pass meconium has been used as a proxy to assess the fetal intestinal

microbiome, although samples are often collected hours after birth and even breastfeeding,

suggesting potential postnatal effects on the microbial composition of the meconium (Collado

et al., 2016; Liu et al., 2019; Stinson et al., 2019). Mammalian animal models can overcome

several of these ethical and practical difficulties, providing insights in the fetal gastrointestinal

microbiome.

In the current study, the microbial composition of the full-term fetal gut and the corresponding

amniotic fluid was assessed in Belgian Blue cow-calf couples. In this double-muscled beef

breed, caesarean sections (C-sections) are performed on a routine basis during the very early

stages of parturition, while fetal membranes are still intact, rendering this breed a highly

suitable model for full-term gestation microbiome studies.

Besides the potential role of the cow model for research, there is also a significant interest in

the composition and the development of the calf's intestinal microbiome. Gut health and

growth performance during the first weeks of life are main drivers for cost effective livestock

rearing (Thornton Philip, 2010; Lorenz et al., 2011). Few studies have been conducted on the

intestinal microbiome in vaginally born neonatal calves. These have reported low numbers of

bacteria, mostly belonging to the phyla Proteobacteria, Firmicutes, Actinobacteria and

Bacteroidetes (Alipour et al., 2018; Klein-Jöbstl et al., 2019; Guzman et al., 2020). Shared

microbiota were found between calf meconium and the maternal vaginal vestibulum (Alipour

et al., 2018; Yeoman et al., 2018; Klein-Jöbstl et al., 2019). To the best of our knowledge, no

microbiome studies have yet been performed on calves born by C-section, and no studies are

available to assess the association between the microbiome of the calf's meconium and the

microbiome of the corresponding amniotic fluid.

Our aims were to assess the bacterial load and characterize the bacterial composition of the

amniotic fluid and meconium of full-term neonatal calves. To this end, samples were collected

during elective C-section, and analyzed by qPCR, bacterial culture under carefully controlled

and validated conditions, and 16S rRNA gene amplicon sequencing. To minimize reagent and

environmental bacterial DNA contaminants, we included multiple technical controls and

stringently filtered the 16S rRNA gene sequencing data to exclude putative contaminant

sequences.

Materials and methods

80 All experimental procedures were approved by the institutional ethics and animal welfare

81 committee of the Faculty of Veterinary Medicine (EC2018/002 - Ghent University, Belgium).

82 The cows' owners were informed about the study and gave their written consent.

Study design

The sampling of this study was performed at the teaching hospital of the Department of

Reproduction, Obstetrics, and Herd Health of the Faculty of Veterinary Medicine in Ghent

(Belgium), where pregnant Belgian Blue beef cows of different herds (parity between 1 and 5)

were housed for an elective C-section.

In total 25 Belgian Blue cows and their calves (7 males and 18 females) were sampled from

November 2017 until March 2019. The cows were housed in tie-stalls at the facility for 9.5 d

± 5.8 (mean ± standard deviation) prior to C-section and had ad libitum access to hay and water.

During this period, rectal temperature was measured twice daily, and calving indicators such

as udder distension, teat filling, pelvic ligament relaxation, vaginal discharge, vulvar edema,

and behavioral changes were monitored every two hours by graduate veterinary students.

Prior to elective C-section, in cows that exhibited a drop in temperature, cervical dilation was

assessed by manual palpation. The vulvar region was cleaned with iodine soap and water. A

gloved hand was inserted vaginally and the opening of the portio vaginalis cervicis was

estimated. For the present study, elective C-section was performed when the cow had a minimal

cervical dilation of eight centimeters, with no rupture of the fetal membranes prior to surgery.

All cows were healthy according to their vital parameters (heart rate, temperature, respiratory

rate) and there was no clinical evidence of intrauterine infection or contamination.

Sampling

Prior to surgery, the cows were restrained in a standing position in a surgery crush specifically

designed for cattle. C-section procedure was done as described by Kolkman et al. (2007).

Briefly, the surgical area (left flank) was washed and disinfected, an abdominal incision was

made, and part of the uterus was exteriorized for uterotomy. The allantoic sac was opened up

to expose the intact amniotic sac. Amniotic fluid was aspirated through the amniotic membrane,

using a sterile 16 G needle (Agani, Terumo Europe, Hamburg, Germany) and sterile 20 ml

syringe (B. Braun, Melsungen, Germany). Within one hour after sampling, the retrieved

volume was aliquoted, under a laminar-flow hood, into 2 sterile 15 ml tubes (188271, Cellstar,

Greiner bio-one, Frickenhausen, Germany). In the first tube, 6 ml of amniotic fluid was

dissolved in 3 ml of glycerol (≥99 %, G2025, Sigma-Aldrich (Merck), Overijse, Belgium) and

subsequently stored at -80 °C to be used in culture experiments. In the second tube, 12 ml of

amniotic fluid was stored at -80 °C to be used for bacterial DNA extraction.

Meconium samples were acquired directly from the calves' rectum, immediately after birth (no

more than 30 minutes). Until the moment of sampling, calves laid on a clean concrete floor,

with access to neither the dam, nor colostrum. The perineum of the calf was dried with a clean

paper towel and disinfected with 70 % ethanol. A sterile double-guarded equine uterine culture

swab (Har-vet, 17705, Spring Valley, USA; or Minitube, 17214/2950, Tiefenbach, Germany)

was gently introduced in the rectum and the swab was exposed. Samples were taken in

duplicate, one stored immediately at -80 °C with no additives, and the other stored at -80 °C in

a sterile 2 ml cryovial containing 1 ml of a 30 % glycerol solution, prepared by diluting glycerol

(≥99 %, G2025, Sigma-Aldrich (Merck), Overijse, Belgium), in ultra-pure, nuclease-free water

(W4502, Sigma-Aldrich (Merck), Overijse, Belgium) to a final 30 % concentration.

Negative field controls were processed in the surgery room, using the same sampling

procedures and disposables. In total, 16 empty, sterile double-guarded equine uterine culture

swabs (10 Har-vet swabs and 6 Minitube swabs) were included for the meconium sampling, 5

of the Har-vet swabs stored in 1 ml of a 30 % glycerol stock solution, the others with no

additives. Additionally, 12 negative controls were included for the amniotic fluid sampling,

aspirating ultra-pure, nuclease-free water (W4502, Sigma-Aldrich (Merck), Overijse, Belgium)

instead of amniotic fluid. For 8 of them, the ultra-pure, nuclease-free water was stored with no

additives, and for 4 of them, 6 ml ultra-pure, nuclease-free water was dissolved in 3 ml of

glycerol.

All samples were shipped on dry ice to the laboratory of the Department of Veterinary

Biosciences of the Faculty of Veterinary Medicine in Helsinki (Finland) for further processing.

Culture

Validation of culture media

One of our aims was to define a single bacterial culture medium, capable of sustaining a

majority of the calf intestinal core microbiota. Consequently, this single medium could be used

for culturing the meconium and amniotic fluid samples, avoiding further splitting or dilution

of the low biomass samples. We first tested GAM "Nissui" (Gifu Anaerobic Medium Agar,

Code 05420, HyServe, Germany), YCFA (Yeast extract, Casitone and Fatty Acid), LB

(Lysogeny Broth, BD) and BB (Trypticase Soy Agar supplemented with Bovine Blood 211043,

Tammer BioLab Oy, Tampere, Finland) media for the ability to sustain calf intestinal core

microbiota, by plating each type of medium anaerobically with feces of a healthy 7 day old calf

(Lopez-Siles et al., 2012; Alipour et al., 2018). After 7 days of growth, mixed cultures were

extracted from the plates, 16S rRNA gene amplicon sequenced, and compared to the core

microbiota previously observed in fecal samples of young calves (Alipour et al., 2018).

Sample culturing

Bacterial culture on GAM-agar plates was performed for 24 meconium samples (5

corresponding negative controls) and 24 amnion samples (4 negative controls). The frozen

samples were first transferred to an anaerobic workstation (Ruskinn Concept Plus), mixed

thoroughly and plated, using an aseptic technique, at +37 °C. After this, the samples were

transferred to a laminar flow cabinet and plated in aerobic conditions at +37 °C. The culture

plates were checked for growth daily for 14 days. All visible bacterial colonies were

subcultured until pure isolates were obtained. Fresh cultures were then identified using

MALDI-TOF (Bruker Microflex LT) at the Central laboratory of the Faculty of Veterinary

Medicine (Helsinki, Finland).

Samples were prepared using MALDI Biotyper MSP Identification Standard Method v 1.1.

Mass spectra were analyzed in a mass/charge range from 2000 to 20 000 Da with MBT

Compass v4.1 on flexControl v3.4 (Bruker Daltonik GmbH) using BDAL-7311 as the

reference library. The Bruker Bacterial Test Standard (RUO) (Bruker Daltonik GmbH) was

used for instrument calibration. If the identification confidence score was <2.00, further

identification was done with 16S rRNA gene amplicon Sanger sequencing at the Institute of

Biotechnology (University of Helsinki, Finland).

DNA extraction

DNA from the meconium samples (N = 25) and corresponding negative controls (N = 11) were extracted using ZymoBIOMICS DNA Miniprep Kit (Zymo Research, Irvine, CA, USA) according to the manufacturer's instructions, with minor modifications to the protocol as described previously (Alipour et al., 2018; Husso et al., 2020). ZymoBIOMICS™ Microbial Community Standard and an in-house fecal standard were processed with the meconium samples and no-template-controls were included in every batch.

For the amniotic fluid samples (N = 23) and the corresponding negative controls (N = 8), 2 ml of each sample was first centrifuged in a microcentrifuge at 16100 × g 10 min in +4 °C. Most of the supernatant was removed and 750 µl of ZymoBIOMICS Lysis Solution and 19 µl of proteinase K (D3001-2-20/D3001-2-5, Zymo Research) were added to the remaining 200 µl of the samples and incubated for 30 min at +55 °C. After these steps, the same protocol as applied for the meconium samples was followed.

All manipulations of the tubes during the process were performed in a laminar flow cabinet, and the workplace, instruments and pipettes were cleaned routinely with 10 % bleach. Certified DNA, RNase, DNase and PCR inhibitor free tubes (STARLAB International, Germany) and Nuclease-free Water (Ambion, Thermo Fisher Scientific, USA) were used for DNA extraction and downstream analyses. All extracted DNA was stored at -80 °C.

Quantitative PCR

The bacterial 16S rRNA gene copy numbers in the meconium, amnion and negative control samples were determined using quantitative PCR. The analyses were performed as described previously (Alipour et al., 2018; Husso et al., 2020), with the exception that the PCR master

mix was first treated using a dsDNAse based decontamination kit (Enzo Life Sciences,

Farmingdale, New York). The amplification was performed using the Bio-Rad CFX96

instrument (Bio-Rad, Hercules, California) and universal eubacteria probe and primers

(Nadkarni et al., 2002). A standard series and negative controls were included in every run.

The data were analyzed using the Bio-Rad CFX Maestro software. The results were normalized

to the averages of absolute 16S rRNA gene copy numbers in the negative controls included in

each run.

Library preparation and 16S rRNA gene amplicon sequencing

The V3-V4 region of the 16S rRNA gene amplicons was sequenced using the Illumina MiSeq

platform in the DNA core facility of the University of Helsinki, as described previously

(Alipour et al., 2018; Husso et al., 2020). In total, 23 amniotic fluid samples and 23 meconium

samples of the same cow-calf couple were sequenced.

The ZymoBIOMICS Microbial Community Standard (Zymo Research, USA) and the in-house

adult cow fecal standard were pre-amplified with 12 cycles and all other sample types,

including negative controls, with 21 cycles.

The observed composition and abundances for the commercial standard matched the expected

composition provided by the manufacturer (data not shown).

Bioinformatics

205 The detailed bioinformatics pipeline is described in the Supplementary materials. Briefly, the

206 read quality was first inspected with FastQC and MultiQC (Andrews, 2011; Ewels et al., 2016).

207 Leftover primes and spacers were then trimmed with Cutadapt v1.10 (Martin, 2011).

208 A mapping file was created for QIIME2 and validated with Keemei (Rideout et al., 2016). The

FASTQ-files were imported to QIIME2 v2019.4, where the DADA2 plugin was used to denoise and quality filter the reads, call ASVs and generate a feature table (Callahan et al., 2016; Bolyen et al., 2019). A naïve Bayes classifier was trained in QIIME2 against SILVA v132 99 % database, extracted to only include the V3-V4 region and used to assign taxonomy to ASVs (Quast et al., 2013; Bokulich et al., 2018). Sequences derived from chloroplasts or mitochondria were removed and singletons were filtered out, leaving only bacteria with at least phylum-level identification.

The processed data was *in silico* filtered to remove ASVs which represented probable contaminants, as described previously (Husso et al., 2020). Briefly, an ASV was removed if its prevalence in actual samples was ≤2× its prevalence in instrument controls (DNA extracted from empty sampling instruments only handled aseptically in a laminar flood cabinet) or its prevalence in field controls (empty sampling instruments exposed to the surgery room environment), and if its mean relative abundance in actual samples was ≤10× its mean abundance in instrument controls or its mean abundance in field controls. The filtering was performed separately for meconium and amnion samples. If less than 500 reads remained after the decontamination, as was the case for six meconium samples, the sample was removed from further analyses.

Statistics

The 16S rRNA gene qPCR results from samples and negative controls were compared using two-tailed Mann-Whitney U test in IBM SPSS Statistics 25.

Shannon diversity indices were calculated for genus-level data with the R package phyloseq (McMurdie and Holmes, 2013). The Kruskal-Wallis rank sum test was used to compare the bacterial community structure of both sample types in RStudio (R Studio Team, 2020).

The PCoA figures were plotted using ASV and genus-level data and Bray-Curtis distances with

the R package phyloseq (McMurdie and Holmes, 2013). Permutational analysis of variance

(PERMANOVA) and multivariate homogeneity of group dispersions (betadisper) were

calculated using the R package vegan, using 9999 permutations (Oksanen et al., 2019).

DESeq2 was used to explore differentially abundant ASVS by calculating differential

expression between sample groups (Love et al., 2014).

The LEfSe (Linear discriminant analysis Effect Size) web application was used to identify the

taxons most likely explaining the differences between the sample types (Segata et al., 2011).

An ecologically organized heatmap of the top 40 most abundant ASVs was created with the R

package phyloseq (Rajaram and Oono, 2010; McMurdie and Holmes, 2013).

Spearman correlations and their Bonferroni corrected p-values were calculated using R-

package Hmisc 4.3.0 (Harrel, 2019).

Differences were considered significant at $P < 0.05$.

Results

Quantification of bacterial 16S rRNA in amniotic fluid and meconium samples by qPCR

248 The 16S rRNA gene copy numbers in amniotic fluid and meconium samples were assessed by

249 qPCR (Fig. 1). To minimize variation between qPCR runs, the copy numbers were normalized

250 to the averages of the negative controls in the same run as explained in Materials and Methods.

In the amniotic fluid samples, there was no significant difference (P = 0.104) between the

normalized 16S rRNA gene copy numbers of the samples (N = 23; mean = 2.4; SD = 2.1) and

the negative controls (N = 8; mean = 1; SD = 0.53).

In the meconium samples, the 16S rRNA gene copy number (N = 25; mean = 9.9; SD = 27)

was significantly higher (P < 0.001) than in the negative controls (N = 11; mean = 1; SD =

0.86).

Figure 1. 16S rRNA gene copy numbers (fold change vs. negative controls) for amniotic fluid (n = 23,

negative controls n = 8) and meconium samples (n = 25, negative controls n = 11). Samples were

collected during elective C-section in Belgian Blue cow-calf pairs. The copy number of each sample

was normalized to the average copy number of the corresponding negative controls and log10

263 *transformed. The midline of the box is the median, with the upper and lower limits of the box being the*
264 *third and first quartile. The whiskers extend up to 1.5 times the interquartile range from the box to the*
265 *furthest data point within that distance.*

Decontamination of the 16S rRNA gene amplicon sequencing data

We performed a stringent filtering of the 16S rRNA gene sequencing data to remove the
amplicon sequence variants (ASVs) potentially stemming from reagent contaminants and
environmental bacterial DNA (Table 1).

The decontamination was based on the prevalence and relative abundance of each ASV in the
samples and negative controls (see Materials and Methods).

Table 1. Mean read counts (standard deviation) detected in meconium and amniotic fluid samples and
their negative controls. Only sequences with at least phylum-level identification were retained for
further analyses. If less than 500 reads remained after in silico decontamination, as in six meconium
samples, the sample was removed from further analyses.

Sample type	Raw		Processed		Decontaminated	
	N	Reads	N	Reads	N	Reads
Meconium	23	92266 (36235)	23	57082 (25543)	17	4189 (8931)
Meconium control	11	126800 (21164)	11	84023 (14585)	-	-
Amnion	23	127797 (22376)	23	83279 (15120)	23	3338 (2922)
Amnion control	6	136488 (12663)	6	90802 (8998)	-	-

Microbial composition of amniotic fluid and meconium samples

The microbiota of both sample types consisted primarily of Proteobacteria, Firmicutes,
Bacteroidetes and Actinobacteria phyla, with individual variation (Fig. 2). In both sample
types, the inter-individual variation increased at lower taxonomic levels (Supplementary Table

1 and Fig. 3). The most abundant bacterial genera in meconium were *Delftia, Staphylococcus*

and *Clostridium sensu stricto 1*, while the most prevalent genera were *Delftia, Acinetobacter*,

unclassified *Burkholderiaceae, Staphylococcus* and *Corynebacterium 1* (Supplementary Table

1). In the amniotic fluid samples, *Staphylococcus, Streptococcus, Delftia, Sphingomonas* and

Enterococcus were the most abundant genera, and *Delftia, Streptococcus, Staphylococcus,*

Sphingomonas and *Acinetobacter* were the most prevalent genera (Supplementary Table 1).

The observed composition and abundances for the commercial standard matched the expected

composition provided by the manufacturer.

290
Figure 2. Microbiota composition in bovine meconium and amniotic fluid samples collected during

elective C-section, adult feces, and a commercial community composition standard. The main colors

indicate the bacterial phyla. Within phyla, the shades indicate bacterial genera. The lightest shade of

each phylum shows the combined abundance of the least abundant genera (with a maximum of <0.5%

of total).

A heatmap analysis of the 40 most abundant ASVs shows the difference between the sample

types in more detail (Fig. 3).

Figure 3. Ecologically organized heatmap (NMDS, Bray) of the 40 most abundant ASVs in bovine meconium and amniotic fluid samples collected during elective C-section, sorted by sample type. The taxon names are presented as genus level identifications. Color scale indicates relative abundance and is a log transformation with base 4.

305 The alpha diversity (Shannon index) at the genus level was not significantly different between

306 the meconium and amniotic fluid samples (Fig. 4, P = 0.889).

309 *Figure 4. Shannon diversity index showing the bacterial community structure (at genus level) of*

310 *amniotic fluid and meconium samples, collected during elective C-section in Belgian Blue cow-calf*

311 *pairs. Boxplot as in Fig 1.*

312
313
314
315

Comparison of amniotic fluid and meconium microbiota composition

The microbiota composition was significantly different for amniotic fluid and meconium

samples at the ASV (PERMANOVA, P < 0.001, R2 = 0.049, Fig. 5 A), and genus level

(PERMANOVA, P < 0.001, R2 = 0.062, Fig. 5 B).

Figure 5. Comparison of the bacterial community structure of amniotic fluid and meconium. 5 A: PCoA
on Bray-Curtis distances based on ASV level data. Figure 5 B: PCoA on Bray- Curtis distances based
on genus level data. Colors and shapes indicate the sample types.

According to the DESeq2 analysis, three ASVs were more abundant in meconium than in

amniotic fluid: one *Staphylococcus* ASV (log2 Fold Change = 25.084, P_{adj} < 0.001), one

Rubrobacter ASV (log2 Fold Change = 8.337, P_{adj} < 0.001), and one *Clostridium* ASV (log2

Fold Change = 8.055, P_{adj} = 0.004).

In amniotic fluid, one *Sphingomonas* ASV (log2 Fold Change = -8.420, $P_{adj} < 0.001$) and one

Staphylococcus ASV (log2 Fold Change = -23.438, $P_{adj} < 0.001$) were more abundant than in

meconium.

By LefSe (Linear discriminant analysis effect size) analysis including all taxonomic levels,

meconium samples had a greater relative abundance of *Clostridiales* and *Rubrobacter* (LDA

score (log10) > 4.0) than the amniotic fluid, but a lower relative abundance of Bacillales and

Corynebacteriales (LDA score (log10) < -4.0).

Correlations between meconium and amniotic fluid from the same animal

338   We calculated Spearman rank correlations for the microbiota between meconium and amniotic

339   fluid samples, taking into account ASVs that were present >100 times in all of the samples.

340   Average correlations (ρ) at ASV (N = 17, ρ_{avg} = -0.0201, $P_{avg} < 0.05$) and genus level (N = 17,

341   ρ_{avg} = 0.1685, $P_{avg} < 0.05$) were very weak and biologically not significant. All p-values were

342   Bonferroni corrected for multiple testing.

Bacterial cultures

Based on our testing of multiple media, the GAM (Gifu Anaerobic Medium) agar was selected

to assess if there were live bacteria in the samples stored in glycerol (see Materials and

Methods), since it was able to sustain the largest number of bacterial core genera found in calf

feces, including some additional unique genera not found on any other plate type (Alipour et

al., 2018).

All amniotic samples and their negative controls were culture negative, both in anaerobic and

aerobic conditions. Five out of 24 meconium samples and 1 out of 5 negative controls were

culture positive on GAM agar. The results for each sample are presented in detail in Table 2.

Both gram negative and gram-positive strains were isolated from the samples.

Table 2. Culture positive meconium samples and identified bacteria. If the identification confidence
score was <2.00 for MALDI-TOF, further identification was done with 16S rRNA gene amplicon Sanger
sequencing to achieve genus level identification for all isolates.

Sample	Aerobic strains	Anaerobic strains
Negative control 1	*Staphylococcus* sp. (Gram+), *Kocuria* sp. (Gram+)	*Rothia* sp. (Gram+)
Meconium 5	*Roseomonas* sp. (Gram-)	-
Meconium 10	*Acinetobacter* sp. (Gram-)	*Streptococcus* sp. (Gram+), *Fusobacterium* sp. (Gram-)
Meconium 12	-	*Cutibacterium* sp. (Gram+)
Meconium 16	*Kocuria* sp. (Gram+)	-
Meconium 21	*Achromobacter* sp. (Gram-)	-

The Sanger sequence from the *Roseomonas* sp. isolate acquired from a meconium sample was

a 100 % identical match to the 16S rRNA gene amplicon sequencing data from the same

sample. In addition, the Sanger sequences of *Staphylococcus* sp., *Kocuria* sp., *Roseomonas* sp.,

and *Acinetobacter* sp. isolates matched the 16S rRNA gene amplicon sequencing data when

compared to all meconium samples.

Three bacterial genera could be identified in the negative controls: *Rothia* sp., *Staphylococcus*

sp. and *Kocuria* sp.. Of these, only *Kocuria* sp. was also identified from the actual meconium

samples.

Discussion

Vertical transmission of microbes during, and perhaps already before birth is a universal

phenomenon across the whole animal kingdom and plays an important role in shaping host

immunity and fitness (Funkhouser and Bordenstein, 2013; Vandenplas et al., 2020). Despite

the advances in our understanding of the assembly and lifelong effects of the perinatal

microbiota, the *in utero* exposure of the fetus to maternal bacteria and bacterial components

remains poorly understood (Perez-Munoz et al., 2017; Korpela and de Vos, 2018; Liu et al.,

2019).

In the present study, we sampled amniotic fluid during, and the corresponding calves

immediately after they were born by elective C-section. The elective C-section is a routine

procedure in Belgian Blue beef cattle, since the pelvis of the dam does not allow vaginal birth

of the relatively large double-muscled calf (Kolkman et al., 2007). We performed the surgery

during the very early stages of parturition, when both fetal membranes (amniotic and

allantochorionic membrane) were still intact, allowing us to sample the full-term fetus and the

intrauterine environment without exposure to the vaginal microbes and minimal exposure to

the environmental microbiota. Meconium samples were collected rectally. The relatively large

size of the calves permits the insertion of a double-guarded sampling device into the rectum,

minimizing the risk of contamination.

In contrast, in most human studies first-pass meconium was collected from diapers (Hansen et

al., 2015; Collado et al., 2016; Liu et al., 2019; Stinson et al., 2019). Because the first-pass

meconium is often collected several hours after birth, microbes from the environment and the

mother (especially during breastfeeding) have time to colonize the intestine. Hence, this

sampling procedure should be seriously questioned for researching the eventual intrauterine

transmission of bacteria to the neonatal gastrointestinal tract (Dornelles et al., 2020). In the

current study, amniotic fluid samples were collected *in vivo*, while the amniotic membrane was

still intact, as has been done in humans during pre-labor C-section (Collado et al., 2016).

Bacterial culture is traditionally used to determine the viable population in a bacterial

community. However, standard culture-dependent techniques are known to be inadequate to

identify all bacteria. Capturing a broad bacterial diversity would require the use of many

different selective media and multiple specific growing conditions (Lau et al., 2016; Lagier et

al., 2018). In the current study, one of our aims was to define a single bacterial culture medium,

capable of sustaining the majority of the calf intestinal core microbiota. We first tested four

different culture media: GAM "Nissui" (Gifu Anaerobic Medium Agar), mostly used for

anaerobic bacteria, YCFA (Yeast extract, Casitone and Fatty Acid), containing volatile fatty

acids typically supporting the growth of several gut bacteria, LB (Lysogeny Broth) as a general

media, and BB (Trypticase Soy Agar supplemented with Bovine Blood) supporting fastidious

bacteria by blood enrichment. The GAM medium was found optimal for both aerobic and

anaerobic culturing.

Bacteria were successfully cultured from 5 out of 24 meconium samples, representing both

gram-negative and gram-positive bacterial genera, known to live in mammalian hosts.

Roseomonas and *Achromobacter* species are known to act as opportunistic pathogens, but have

also been isolated from a wide variety of environmental sources (Reverdy et al., 1984; Rihs et

al., 1993). Various *Acinetobacter* and *Streptococcus* species have physiological functions in

their mammalian hosts, but the genera also include pathogenic and/or environmental species

(Krzyściak et al., 2013; Touchon et al., 2014). *Fusobacteria* species are typically closely related

to mucous membranes, especially the oral cavities, while *Cutibacterium* and *Kocuria* species

have most often been isolated from the skin (Hofstad, 1992; Savini et al., 2010; Scholz and

Kilian, 2016). One of our negative controls yielded *Kocuria* colonies. This was the only taxon

shared between the actual meconium samples and the negative controls, suggesting low-level

contamination from the sampling environment.

Although we detected cultivable bacteria in some of the meconium samples, most of the

samples were culture negative. This indicates that the presence of live bacteria in the fetal

intestine is rare, in contrast to the prevalent bacterial DNA. Individual differences in amounts

of viable bacteria retrieved from meconium were also described in a recent dog study.

Interestingly, puppies born without detectable meconium microbiota were shown to have a

slower growth rate than those in which meconium microbiota were detected (Pipan et al.,

2020). Both the mucosal surface of the intestine as well as the amniotic fluid contain

bacteriostatic substances, such as lactoferrin and salivary scavenger and agglutinin (SALSA),

which are able to suppress bacterial growth and viability (Reichhardt and Meri, 2016;

Lisowska-Myjak et al., 2019). In human meconium, SALSA amounts to 10 % of all proteins,

highlighting its potential role in antimicrobial defense (Reichhardt et al., 2014). Lactoferrin,

SALSA and other similar substances may explain the small number of viable bacteria observed

in our study, in contrast to the significant amount of detected bacterial DNA.

The culture-independent techniques were performed based on the 16S rRNA gene. Since

contamination is a major challenge in the analysis of low bacterial biomass samples, we

processed several types of negative controls alongside our biological samples, treated qPCR

mastermix with dsDNAse, and applied rigorous *in silico* filtering of potential contaminants

(Glassing et al., 2016; Eisenhofer et al., 2018). Many similar studies are published without or

with a very limited number of contamination controls, which makes our approach more

reliable, especially for assessing low bacterial biomass environments.

We observed a small but significant amount of bacterial DNA in the meconium samples by

16S qPCR, suggesting a prenatal transmission of bacterial DNA to the fetal intestine. This is

consistent with previous studies in cattle, in which calves born at term *per vias naturales*, were

sampled (Mayer et al., 2012; Alipour et al., 2018; Klein-Jöbstl et al., 2019). A profile

dominated by the phyla Proteobacteria, Firmicutes, Bacteroidetes and Actinobacteria was

identified in the meconium, in agreement with previous studies using 16S rRNA gene amplicon

sequencing (Alipour et al., 2018; Yeoman et al., 2018; Klein-Jöbstl et al., 2019; Husso et al.,

2020; Guzman et al., 2020). Additionally, some of the most prevalent genera observed in

meconium in our study, such as *Sphingomonas*, *Acinetobacter*, *Pseudomonas*, *Ruminococcus*

and *Stenotrophomonas,* have also been previously described in neonatal calves (Klein-Jöbstl

et al., 2019).

In contrast to the meconium, there was no significant difference between the 16S rRNA gene

copy numbers of amniotic fluid samples and the negative controls, and all were culture

negative. A recent caesarean section study in sheep stated amniotic fluid sterility during the

third trimester of pregnancy (Malmuthuge and Griebel, 2018). In other studies however, a low

biomass microbiome in the pregnant bovine uterus was described (Moore et al., 2017; Guzman

et al., 2020).

Using 16S rRNA gene amplicon sequencing, we found a bacterial signature in the amniotic

fluid, dominated by the phyla Proteobacteria, Firmicutes, Bacteroidetes and Actinobacteria, in

accordance with previous studies (Karstrup et al., 2017; Moore et al., 2017; Guzman et al.,

2020). Deeper taxonomic identification, down to the family or genus level, revealed no further

agreements with these studies.

From a comparative point of view between mammalian species, the amniotic fluid sampled by

Collado et al. (2016) in women during pre-labor C-section was found to contain a distinct, low

diversity, low biomass microbiome, predominated by Proteobacteria (*Enterobacteriaceae*).

Despite this and other recent studies, the presence of microbial DNA and culturable bacteria in

the human placenta and fetal environment is still under debate (de Goffau et al., 2019).

The microbial DNA signatures in meconium and amniotic fluid were similar at the phylum

level. However, they presented a significantly different microbial composition at genus and

ASV level: genera *Staphylococcus*, *Rubrobacter* and *Clostridium* were relatively more

abundant in meconium samples than in amniotic fluid samples according to DESeq2 analysis.

Complementing this with the LefSe analysis using all taxonomic levels, also found

Clostridiales and *Rubrobacter* taxons relatively more abundant in the meconium. The

microbial compositions in the bovine amniotic fluid and meconium of the corresponding

neonate did not show a significant correlation, contrary to earlier studies in humans (Collado

et al., 2016; He et al., 2020).

Our analysis suggests that the bacterial DNA found in meconium does not necessarily originate

from the amniotic fluid. The maternal microbial components may be translocated via the

placental and umbilical blood vessels directly to the fetal internal organs. Microbial

macromolecules and even live bacteria may be transported by leukocytes via these blood

vessels (Perez et al., 2007).

A recent bovine study revealed differing microbial communities between the amniotic fluid

and the meconium (Guzman et al., 2020). As the fetus swallows the amniotic fluid, which is

concentrated and retained in the intestine as meconium, together with epithelial cells and

intestinal secretions, it is unexpected that the two would harbor completely different bacteria.

However, the amniotic fluid microbiota may fluctuate dynamically over time, while meconium

represents a more stable collection of substances accumulated over the gestation period.

Moreover, fetal excretion in the amniotic fluid is limited in cattle. Meconium is usually

expelled only after birth, and fetal urine largely accumulates in the allantoic cavity, between

chorion and amnion, rather than in the amniotic cavity (Bongso and Basrur, 1976).

The less permeable synepitheliochorial placenta evidenced by three distinct layers may restrict

the translocation of bacteria and their components from the dam to the fetus in cattle, in

comparison to humans. The very low levels of bacterial DNA in the bovine amniotic fluid limit

the sequencing depth and thus the precision of the compositional analyses. Our strict data

decontamination protocol furthermore minimized the artefactual similarity arising from shared

contaminants.

In conclusion, we found that the meconium of full-term calves delivered by elective caesarean

section in most cases contains a small amount of diverse bacterial DNA and eventually

although rather rare, culturable bacteria. In the amniotic fluid, bacteria were not observed by

16S qPCR or culturing, but a microbial DNA profile was distinguishable from controls by

amplicon sequencing. Based on these results, bacterial components are translocated to the fetus

in utero, but the prenatal acquisition of live bacteria is likely not physiologically significant.

Data availability

The raw 16S rRNA gene amplicon sequencing dataset is fully available in the NCBI Sequence

Read Archive (SRA) with the accession number PRJNA643145 after journal acceptance.

Acknowledgements

502 We thank the owners of the cows for allowing sampling of their animals, and all clinicians and
503 graduate veterinary students for helping with the monitoring of the animals, performing the
504 surgery and sampling. Additionally, we thank Kirsi Lahti for expert technical assistance in the
505 laboratory.

Author contributions

507 MN, LL, AI and GO contributed to conception and design of the study. AH performed the
508 bioinformatics, statistical analyses and bacterial culture. AH and TG performed MALDI-TOF
509 analyses. LL and JG organized the sampling and collected the samples. AH and LL wrote the
510 manuscript. All authors contributed to manuscript preparation, read, and approved the
511 submitted version.

Ethics declaration

514 All experimental procedures were approved by the institutional ethics and animal welfare
515 committee of the Faculty of Veterinary Medicine (EC2018/002 - Ghent University, Belgium).
516 The cows' owners were informed about the study and gave their written consent for their cows'
517 participation.

References

Aagaard, K., Ma, J., Antony, K. M., Ganu, R., Petrosino, J., and Versalovic, J. (2014). The placenta harbors a unique microbiome. *Sci Transl Med* 6, 237ra65. doi:10.1126/scitranslmed.3008599.

Al Alam, D., Danopoulos, S., Grubbs, B., Ali, N. A. B. M., MacAogain, M., Chotirmall, S. H., et al. (2020). Human Fetal Lungs Harbor a Microbiome Signature. *Am. J. Respir. Crit. Care Med.* doi:10.1164/rccm.201911-2127LE.

Alipour, M. J., Jalanka, J., Pessa-Morikawa, T., Kokkonen, T., Satokari, R., Hynönen, U., et al. (2018). The composition of the perinatal intestinal microbiota in cattle. *Sci. Rep.* 8, 10437. doi:10.1038/s41598-018-28733-y.

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data. Available at: *http://www.bioinformatics.babraham.ac.uk/projects/fastqc*

Baker, J. M., Chase, D. M., and Herbst-Kralovetz, M. M. (2018). Uterine Microbiota: Residents, Tourists, or Invaders? *Front. Immunol.* 9. doi:10.3389/fimmu.2018.00208.

Bokulich, N. A., Kaehler, B. D., Rideout, J. R., Dillon, M., Bolyen, E., Knight, R., et al. (2018). Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* 6, 90. doi:10.1186/s40168-018-0470-z.

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., et al. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* 37, 852–857. doi:10.1038/s41587-019-0209-9.

Bongso, T. A., and Basrur, P. K. (1976). Foetal fluids in cattle. *Can. Vet. J.* 17, 38.

Brunham, R. C., and Paavonen, J. (2020). Reproductive system infections in women: lower genital tract syndromes. *Pathog. Dis. 78, 5.*

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* 13, 581. doi:10.1038/nmeth.3869.

Cappelletti, M., Presicce, P., and Kallapur, S. G. (2020). Immunobiology of Acute Chorioamnionitis. *Front. Immunol.* 11. doi: 10.3389/fimmu.2020.00649.

Collado, M. C., Rautava, S., Aakko, J., Isolauri, E., and Salminen, S. (2016). Human gut colonisation may be initiated in utero by distinct microbial communities in the placenta and amniotic fluid. *Sci Rep* 6, 23129. doi:10.1038/srep23129.

de Goffau, M. C., Lager, S., Sovio, U., Gaccioli, F., Cook, E., Peacock, S. J., et al. (2019). Human placenta has no microbiome but can contain potential pathogens. *Nature*. doi:10.1038/s41586-019-1451-5.

Dierikx, T. H., Visser, D. H., Benninga, M. A., van Kaam, A., de Boer, N. K. H., de Vries, R., et al. (2020). The influence of prenatal and intrapartum antibiotics on intestinal microbiota colonisation in infants: A systematic review. *J. Infect.*

doi:https://doi.org/10.1016/j.jinf.2020.05.002.

Dornelles, L. V., Procianoy, R. S., Roesch, L. F., Corso, A. L., Dobbler, P. T., Mai, V., et al. (2020).
Meconium microbiota predicts clinical early-onset neonatal sepsis in preterm neonates. *J.*
Matern. Fetal Neonatal Med., 1–9.

Eisenhofer, R., Minich, J. J., Marotz, C., Cooper, A., Knight, R., and Weyrich, L. S. (2018).
Contamination in Low Microbial Biomass Microbiome Studies: Issues and
Recommendations. *Trends Microbiol.* doi:10.1016/j.tim.2018.11.003.

Ewels, P., Magnusson, M., Lundin, S., and Kaller, M. (2016). MultiQC: summarize analysis results
for multiple tools and samples in a single report. *Bioinformatics* 32, 3047–8.
doi:10.1093/bioinformatics/btw354.

Funkhouser, L. J., and Bordenstein, S. R. (2013). Mom knows best: the universality of maternal
microbial transmission. *PLoS Biol* 11, e1001631. doi:10.1371/journal.pbio.1001631.

Glassing, A., Dowd, S. E., Galandiuk, S., Davis, B., and Chiodini, R. J. (2016). Inherent bacterial
DNA contamination of extraction and sequencing reagents may affect interpretation of
microbiota in low bacterial biomass samples. *Gut Pathog.* 8, 24.

Gomez de Agüero, M., Ganal-Vonarburg, S. C., Fuhrer, T., Rupp, S., Uchimura, Y., Li, H., et al.
(2016). The maternal microbiota drives early postnatal innate immune development. *Science*
351, 1296. doi:10.1126/science.aad2571.

Gonçalves, L. F., Chaiworapongsa, T., and Romero, R. (2002). Intrauterine infection and prematurity.
Ment. Retard. Dev. Disabil. Res. Rev. 8, 3–13.

Guzman, C.E., Wood, J.L., Egidi, E. et al. (2020) A pioneer calf foetus microbiome. *Sci.*
Rep. 10, 17712.

Hansen, R., Scott, K. P., Khan, S., Martin, J. C., Berry, S. H., Stevenson, M., et al. (2015). First-pass
meconium samples from healthy term vaginally-delivered neonates: an analysis of the
microbiota. *PloS One* 10, e0133320.

Harrel, F. (2019). *Hmisc: Harrell Miscellaneous*. Available at: https://CRAN.R-
project.org/package=Hmisc.

He, Q., Kwok, L. Y., Xi, X., Zhong, Z., Ma, T., Xu, H., ... & Zhang, H. (2020). The meconium
microbiota shares more features with the amniotic fluid microbiota than the maternal fecal
and vaginal microbiota. *Gut Microbes*, *12*(1), 1794266.

Hockney, R., Waring, G. J., Taylor, G., Cummings, S. P., Robson, S. C., Orr, C. H., et al. (2020).
Fetal membrane bacterial load is increased in histologically confirmed inflammatory
chorioamnionitis: A retrospective cohort study. *Placenta* 91, 43–51.
doi:10.1016/j.placenta.2020.01.006.

Hofstad, T. (1992). "The genus Fusobacterium," in *The prokaryotes* (Springer), 4114–4126.

Husso, A., Jalanka, J., Alipour, M. J., Huhti, P., Kareskoski, M., Pessa-Morikawa, T., et al. (2020).
The composition of the perinatal intestinal microbiota in horse. *Sci. Rep.* 10, 1–12.

Jeon, S. J., Cunha, F., Vieira-Neto, A., Bicalho, R. C., Lima, S., Bicalho, M. L., et al. (2017). Blood
as a route of transmission of uterine pathogens from the gut to the uterus in cows. *Microbiome*
5, 1–13.
625

626 Jimenez, E., Marin, M. L., Martin, R., Odriozola, J. M., Olivares, M., Xaus, J., et al. (2008). Is
627   meconium from healthy newborns actually sterile? *Res Microbiol* 159, 187–93.
628   doi:10.1016/j.resmic.2007.12.007.
629

630 Karstrup, C. C., Klitgaard, K., Jensen, T. K., Agerholm, J. S., and Pedersen, H. G. (2017). Presence of
631   bacteria in the endometrium and placentomes of pregnant cows. *Theriogenology* 99, 41–47.
632   doi:10.1016/j.theriogenology.2017.05.013.
633

634 Klein-Jöbstl, D., Quijada, N. M., Dzieciol, M., Feldbacher, B., Wagner, M., Drillich, M., et al. (2019).
635   Microbiota of newborn calves and their mothers reveals possible transfer routes for newborn
636   calves' gastrointestinal microbiota. *PloS One* 14(8): e0220554.
637

638 Kolkman, I., De Vliegher, S., Hoflack, G., Van Aert, M., Laureyns, J., Lips, D., et al. (2007). Protocol
639   of the caesarean section as performed in daily bovine practice in Belgium. *Reprod. Domest.*
640   *Anim.* 42, 583–589.
641

642 Korpela, K., and de Vos, W. M. (2018). Early life colonization of the human gut: microbes matter
643   everywhere. *Curr. Opin. Microbiol.* 44, 70–78.
644

645 Krzyściak, W., Pluskwa, K. K., Jurczak, A., and Kościelniak, D. (2013). The pathogenicity of the
646   Streptococcus genus. *Eur. J. Clin. Microbiol. Infect. Dis.* 32, 1361–1376.
647

648 Lagier, J.-C., Dubourg, G., Million, M., Cadoret, F., Bilen, M., Fenollar, F., et al. (2018). Culturing
649   the human microbiota and culturomics. *Nat. Rev. Microbiol.* doi:10.1038/s41579-018-0041-0.
650

651 Lau, J. T., Whelan, F. J., Herath, I., Lee, C. H., Collins, S. M., Bercik, P., et al. (2016). Capturing the
652   diversity of the human gut microbiota through culture-enriched molecular profiling. *Genome*
653   *Med* 8, 72. doi:10.1186/s13073-016-0327-7.
654

655 Lauder, A. P., Roche, A. M., Sherrill-Mix, S., Bailey, A., Laughlin, A. L., Bittinger, K., et al. (2016).
656   Comparison of placenta samples with contamination controls does not provide evidence for a
657   distinct placenta microbiota. *Microbiome* 4, 29. doi:10.1186/s40168-016-0172-3.
658

659 Lisowska-Myjak, B., Skarżyńska, E., Wojdan, K., and Nasierowska-Guttmejer, A. (2019). Protein and
660   peptide profiles in neonatal meconium. *J. Obstet. Gynaecol. Res.* 45, 556–564.
661

662 Liu, C.-J., Liang, X., Niu, Z.-Y., Jin, Q., Zeng, X.-Q., Wang, W.-X., et al. (2019). Is the delivery
663   mode a critical factor for the microbial communities in the meconium? *EBioMedicine* 49,
664   354–363. doi:10.1016/j.ebiom.2019.10.045.
665

666 Lopez-Siles, M., Khan, T. M., Duncan, S. H., Harmsen, H. J., Garcia-Gil, L. J., and Flint, H. J.
667   (2012). Cultured representatives of two major phylogroups of human colonic
668   Faecalibacterium prausnitzii can utilize pectin, uronic acids, and host-derived substrates for
669   growth. *Appl Env. Microbiol* 78, 420–428.
670

671 Lorenz, I., Fagan, J., and More, S. J. (2011). Calf health from birth to weaning. II. Management of
672   diarrhoea in pre-weaned calves. *Ir. Vet. J.* 64, 9.
673

675 Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion
676   for RNA-seq data with DESeq2. *Genome Biol.* 15, 550.
677

678 Malmuthuge, N., and Griebel, P. J. (2018). Fetal environment and fetal intestine are sterile during the
679   third trimester of pregnancy. *Vet. Immunol. Immunopathol.* 204, 59–64.
680

681 Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
682   *EMBnet.journal* 17. doi:10.14806/ej.17.1.200.
683

684 Mayer, M., Abenthum, A., Matthes, J. M., Kleeberger, D., Ege, M. J., Hölzel, C., et al. (2012).
685   Development and genetic influence of the rectal bacterial flora of newborn calves. *Vet.*
686   *Microbiol.* 161, 179–185. doi:10.1016/j.vetmic.2012.07.023.
687

688 McMurdie, P. J., and Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive
689   Analysis and Graphics of Microbiome Census Data. *PLOS ONE* 8, e61217.
690   doi:10.1371/journal.pone.0061217.
691

692 Moore, R. L., Geraghty, A. A., Feehily, C., Saldova, R., Murphy, E. F., Van Sinderen, D., et al.
693   (2020). Can a probiotic supplement in pregnancy result in transfer to the neonatal gut: A
694   systematic review. *Acta Obstet. Gynecol. Scand*. doi:10.1111/aogs.13899.
695

696 Moore, S. G., Ericsson, A. C., Poock, S. E., Melendez, P., and Lucy, M. C. (2017). Hot topic: 16S
697   rRNA gene sequencing reveals the microbiome of the virgin and pregnant bovine uterus. *J.*
698   *Dairy Sci.* 100, 4953–4960.
699

700 Nadkarni, M. A., Martin, F. E., Jacques, N. A., and Hunter, N. %J M. (2002). Determination of
701   bacterial load by real-time PCR using a broad-range (universal) probe and primers set. 148,
702   257–266.
703

704 Oksanen J, F. G. B., Michael Friendly, Roeland Kindt, Pierre Legendre, Dan McGlinn, Peter R.
705   Minchin, R. B. O'Hara, Gavin L. Simpson, Peter Solymos, M. Henry, H. Stevens, Eduard
706   Szoecs and Helene Wagner (2019). vegan: Community Ecology Package. R package version
707   2.5-4. Available at: https://CRAN.R-project.org/package=vegan.
708

709 Olomu, I. N., Pena-Cortes, L. C., Long, R. A., Vyas, A., Krichevskiy, O., Luellwitz, R., et al. (2020).
710   Elimination of "kitome" and "splashome" contamination results in lack of detection of a
711   unique placental microbiome. *BMC Microbiol.* 20, 1–19.
712

713 Perez, P. F., Doré, J., Leclerc, M., Levenez, F., Benyacoub, J., Serrant, P., et al. (2007). Bacterial
714   imprinting of the neonatal immune system: lessons from maternal cells? *Pediatrics* 119,
715   e724–e732.
716

717 Perez-Munoz, M. E., Arrieta, M. C., Ramer-Tait, A. E., and Walter, J. (2017). A critical assessment of
718   the "sterile womb" and "in utero colonization" hypotheses: implications for research on the
719   pioneer infant microbiome. *Microbiome* 5, 48. doi:10.1186/s40168-017-0268-4.
720

721 Pipan, M. Z., Kajdič, L., Kalin, A., Plavec, T., and Zdovc, I. (2020). Do newborn puppies have their
722   own microbiota at birth? Influence of type of birth on newborn puppy microbiota.
723   *Theriogenology*. doi:10.1016/j.theriogenology.2020.04.014.
724

725 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2013). The SILVA
726   ribosomal RNA gene database project: improved data processing and web-based tools.
727   *Nucleic Acids Res* 41, D590-6. doi:10.1093/nar/gks1219.

R Studio Team (2020). RStudio: Integrated Development for R. *RStudio Support*. Available at:
    http://support.rstudio.com/hc/en-us/articles/206212048

Rackaityte, E., Halkias, J., Fukui, E. M., Mendoza, V. F., Hayzelden, C., Crawford, E. D., et al.
    (2020). Viable bacterial colonization is highly limited in the human intestine in utero. *Nat.*
    Med. 26, 599–607.

Rajaram, S., and Oono, Y. (2010). NeatMap-non-clustering heat map alternatives in R. *BMC*
    Bioinformatics 11, 1–9.

Reichhardt, M. P., Jarva, H., de Been, M., Rodriguez, J. M., Quintana, E. J., Loimaranta, V., et al.
    (2014). The salivary scavenger and agglutinin in early life: diverse roles in amniotic fluid and
    in the infant intestine. *J. Immunol.* 193, 5240–5248.

Reichhardt, M. P., and Meri, S. (2016). SALSA: a regulator of the early steps of complement
    activation on mucosal surfaces. *Front. Immunol.* 7, 85.

Reverdy, M. E., Freney, J., Fleurette, J., Coulet, M., Surgot, M., Marmet, D., et al. (1984).
    Nosocomial colonization and infection by Achromobacter xylosoxidans. *J. Clin. Microbiol.*
    19, 140–143.

Rideout, J. R., Chase, J. H., Bolyen, E., Ackermann, G., Gonzalez, A., Knight, R., et al. (2016).
    Keemei: cloud-based validation of tabular bioinformatics file formats in Google Sheets.
    Gigascience 5, 27. doi:10.1186/s13742-016-0133-6.

Rihs, J. D., Brenner, D. J., Weaver, R. E., Steigerwalt, A. G., Hollis, D. G., and Yu, V. L. (1993).
    Roseomonas, a new genus associated with bacteremia and other human infections. *J. Clin.*
    Microbiol. 31, 3275–3283.

Romero, R., Gomez-Lopez, N., Winters, A. D., Jung, E., Shaman, M., Bieda, J., et al. (2019).
    Evidence that intra-amniotic infections are often the result of an ascending invasion–a
    molecular microbiological study. *J. Perinat. Med.* 47, 915–931.

Rossi, B., Merlo, B., Colleoni, S., Iacono, E., Tazzari, P. L., Ricci, F., et al. (2014). Isolation and in
    vitro characterization of bovine amniotic fluid derived stem cells at different trimesters of
    pregnancy. *Stem Cell Rev. Rep.* 10, 712–724.

Savini, V., Catavitello, C., Masciarelli, G., Astolfi, D., Balbinot, A., Bianco, A., et al. (2010). Drug
    sensitivity and clinical impact of members of the genus Kocuria. *J. Med. Microbiol.* 59,
    1395–1402.

Scholz, C. F., and Kilian, M. (2016). The natural history of cutaneous propionibacteria, and
    reclassification of selected species within the genus Propionibacterium to the proposed novel
    genera Acidipropionibacterium gen. nov., Cutibacterium gen. nov. and
    Pseudopropionibacterium gen. nov. *Int. J. Syst. Evol. Microbiol.* 66, 4422–4432.

Seferovic, M. D., Pace, R. M., Carroll, M., Belfort, B., Major, A. M., Chu, D. M., et al. (2019).
    Visualization of microbes by 16S in situ hybridization in term and preterm placentas without
    intraamniotic infection. *Am. J. Obstet. Gynecol.* 221, 146.e1-146.e23.
    doi:10.1016/j.ajog.2019.04.036.

781   Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W. S., et al. (2011).
782   Metagenomic biomarker discovery and explanation. *Genome Biol.* 12, 1–18.

784   Stinson, L. F., Boyce, M. C., Payne, M. S., and Keelan, J. A. (2019). The not-so-sterile womb:
785   evidence that the human fetus is exposed to bacteria prior to birth. *Front. Microbiol.* 10, 1124.

787   Stinson, L. F., Keelan, J. A., and Payne, M. S. (2018). Comparison of Meconium DNA Extraction
788   Methods for Use in Microbiome Studies. *Front Microbiol* 9, 270.
789   doi:10.3389/fmicb.2018.00270.

791   Theis, K. R., Romero, R., Greenberg, J. M., Winters, A. D., Garcia-Flores, V., Motomura, K., et al.
792   (2020a). No Consistent Evidence for Microbiota in Murine Placental and Fetal Tissues.
793   *mSphere* 5, e00933-19. doi:10.1128/mSphere.00933-19.

795   Theis, K. R., Romero, R., Winters, A. D., Jobe, A. H., and Gomez-Lopez, N. (2020b). Lack of
796   evidence for microbiota in the placental and fetal tissues of rhesus macaques. *mSphere* 5. doi:
797   10.1128/mSphere.00210-20.

799   Thornton Philip, K. (2010). Livestock Production: Recent Trends, Future Prospects. Nairobi Kenya,
800   doi: 10.1098/rstb. 2010.0134 Phil. *Trans R Soc B 27 Sept. 2010 Vol 365 No 1554 2853* 2867.

802   Touchon, M., Cury, J., Yoon, E.-J., Krizova, L., Cerqueira, G. C., Murphy, C., et al. (2014). The
803   genomic diversification of the whole Acinetobacter genus: origins, mechanisms, and
804   consequences. *Genome Biol. Evol.* 6, 2866–2882.

806   Vandenplas, Y., Carnielli, V. P., Ksiazyk, J., Luna, M. S., Migacheva, N., Mosselmans, J. M., et al.
807   (2020). Factors affecting early-life intestinal microbiota development. *Nutrition*, 110812.

809   Yeoman, C. J., Ishaq, S. L., Bichi, E., Olivo, S. K., Lowe, J., and Aldridge, B. M. (2018).
810   Biogeographical Differences in the Influence of Maternal Microbial Sources on the Early
811   Successional Development of the Bovine Neonatal Gastrointestinal tract. *Sci Rep* 8, 3197.
812   doi:10.1038/s41598-018-21440-8.