Title: No evidence of *Toxoplasma gondii* infection in urban and rural squirrels

Riikka P Kinnunen[1*], Chloé Schmidt[1], Adrián Hernández-Ortiz[2], Colin J Garroway[1] *

[1] Department of Biological Sciences, Biological Sciences Building, University of Manitoba, Winnipeg, MB, Canada R3T 2N2

[2] Department of Veterinary Microbiology, Western College of Veterinary Medicine, University of Saskatchewan, 52 Campus Drive, Saskatoon, SK, Canada S7N 5B4

Correspondence

Riikka Kinnunen

email: kinnuner@myumanitoba.ca

Colin Garroway

email: colin.garroway@umanitoba.ca

Abstract

Wildlife in cities can act as reservoirs for parasites that can infect domestic pets and have implications for human health. For example, the Coccidian parasite *Toxoplasma gondii* can infect all mammalian species, with felids acting as the only definitive hosts. Rodents, such as squirrels, could play an important role in urban infection dynamics of *T. gondii*, as squirrels occur at higher densities in cities than in natural environments and regularly share their territories with cats. In urban and suburban areas squirrels can encounter infectious oocysts shed in cat feces in contaminated soil or in the food they eat. They are also preyed upon by cats making them a potentially important species for completing the *T. gondii* life cycle in cities. We hypothesized that due to increased exposure to cats, urban squirrels would be more susceptible to *T. gondii* infection relative to squirrels in more natural areas. We investigated this using molecular and serological methods on samples collected from American red squirrels (*Tamiasciurus hudsonicus*), eastern grey squirrels (*Sciurus carolinensis*), northern flying squirrels (*Glaucomys sabrinus*), and least chipmunks (*Tamias minimus*) in and around the city of Winnipeg, Manitoba, Canada. We tested a total of 230 tissue samples from 46 squirrels for *T. gondii* DNA using quantitative PCR and supplemented these data with analyses of 13 serum samples from grey squirrels (*Sciurus carolinensis*) testing for *T. gondii* antibodies by indirect ELISA. We found no evidence of *T. gondii* infection in any squirrel. This suggests that squirrels may not be important intermediate hosts of *T. gondii* in cities and do not need to be considered as sources of infection to cats.

Introduction

Wildlife in cities can act as reservoirs for parasites that can infect domestic pets and have implications for human health [1]. Indeed, previous work has found increased levels of wildlife parasitism in cities compared to rural areas [2–5]. This increased level of parasitism could reflect both the increased population density of urban host species and higher within

and between species contact rates in response to resource provisioning in cities relative to

rural areas [6,7]. Contrasting this pattern, in some cases the overall species richness and

diversity of parasites can be reduced in cities as urbanization can lead to a decline in host

species richness. With reduced access to hosts, parasites with one or a few host species may

then also become extirpated [7]. Parasites that spread through direct contact or oral-fecal

routes are likely to be favored in urban areas, [7] but general knowledge underlying the ways

host-pathogen interactions operate in cities is still limited [1].

The Coccidian parasite *Toxoplasma gondii* is an interesting parasite within the context of

urban ecosystems. Domestic cats—numerous in cities as pets [8,9]—and other Felidae are the

only known definitive hosts of *T. gondii* [10]. When infected, cats can shed millions of

infectious *T. gondii* oocysts into the environment daily in their feces for a duration of one to

two weeks, making them an important part of the parasite's life cycle [11,12]. Humans can

acquire infection from cats by accidentally ingesting oocysts, for example when cleaning cat

litter or not washing hands after gardening [13]. Other mammals can act as intermediate hosts

for *T. gondii*, acting as reservoirs for the parasite and, in some cases, as possible sources of

infection for cats in cities. Cats and intermediate hosts can become infected by consuming

carcasses infected by *T. gondii* tissue cysts, or by coming into contact with oocyst

contaminated soil, water, or food. A significant proportion of the human population globally

is infected with *T. gondii,* but most healthy people do not experience symptoms of infection

[14]. However, immunocompromised people and pregnant women require medical

intervention to avoid serious health issues [14–16]. Consequently, further knowledge of *T.*

gondii infection dynamics in cities is needed.

Urbanization can increase the risk of an animal being exposed to *T. gondii* [4,17,18]. Many

squirrel species (Sciuridae) are ubiquitous in cities, where they are commonly found at much

higher densities than in natural environments [19]. In urban and suburban areas squirrels

regularly share their territories with domestic cats and collect and cache their food in

backyards and gardens where they can encounter infectious oocysts shed in cat feces in

contaminated soil or in the food they eat. This may make urban squirrels particularly

susceptible to parasite infection compared to their rural counterparts. After being infected

squirrels act as intermediate hosts for the parasite, and the parasite can remain within the host

body in tissue cysts for the rest of the host's life. The infection can be asymptomatic or

develop into the disease toxoplasmosis [20,21]. Many *T. gondii* strains isolated from nature

are of low virulence, leading to subclinical toxoplasmosis that does not kill the animal, but

can make prey—such as a squirrel—susceptible to predation by cats [22], thus enabling the

parasite to complete its life cycle [20,22]. Squirrels may thus act as a source of infection to

cats, in a similar way to other prey species [23]. As such squirrels may play a role in *T. gondii*

population and infection dynamics in cities.

Our aim in this paper was to survey the prevalence of *T. gondii* infection in squirrel

(Sciuridae) populations in and around the city of Winnipeg, Manitoba, Canada. We

specifically asked whether squirrel species (American red squirrel, *Tamiasciurus hudsonicus*;

eastern grey squirrel, *Sciurus carolinensis*; northern flying squirrel, *Glaucomys sabrinus*; and

least chipmunk, *Tamias minimus*) are important intermediate hosts of *T. gondii* and whether

T. gondii infection is more common in a city than in more natural habitats. We hypothesized

that, due to high population densities of both squirrels and cats in cities [9,19], urban squirrels

may act as intermediate hosts of *T. gondii* and that urban squirrels will have a higher

prevalence of *T. gondii* infection than rural squirrels. We tested these hypotheses by using

90   molecular and serological methods on samples collected from four squirrel species in and

91   around the city of Winnipeg, Manitoba, Canada.

Materials and Methods

To investigate *T. gondii* prevalence in squirrels in chronic and acute stages we followed

recommendations from previous studies and used complementary molecular and serological

methods [24]. Quantitative PCR was done on a wide-ranging sample from four squirrel

species (American red squirrel; eastern grey squirrel; northern flying squirrel; and least

chipmunk) and serological testing on focal study populations of grey squirrels in and around

the city of Winnipeg, Manitoba, Canada (Fig. 1). Winnipeg is the largest city in the province

of Manitoba with a population of 778,489 and a total land area of 464,33 km^2 [25]. Winnipeg

lies 239 meters above sea level and has high seasonal climatic variation, with temperature

varying from the extremes ranging between −24 °C to −33 °C between January to March to

around +30 °C to +35 °C between June to September [26].

Collection of Carcass Samples

106 To investigate *T. gondii* prevalence in squirrels during the chronic stage we collected the

107 carcasses of 25 red squirrels, 16 grey squirrels, four flying squirrels, and one chipmunk from

108 trappers, wildlife rehabilitation centers, and pest control companies during the years 2017 to

109 2019 to collect tissue samples for PCR (Fig. 1). We did not obtain the carcasses close enough

110 to death to collect usable serological samples.

Molecular Methods

Upon necropsy, we collected the entire liver, spleen, brain, heart, kidneys, and lungs, and

stored tissues separately at −20 °C until further analysis. As *T. gondii* is a cyst-forming

parasite, the detection probability of *T. gondii* can differ between organs [28]. Consequently,

we tested multiple samples per individual from two to six different organs to maximize the

probability of detecting the parasite.

Cell lysis.—Cell lysis was done by first adding 3 ball bearings to each 2 mL screw-cap

tube containing 0.6 mL of ATL buffer, and adding 100 mg of frozen tissue or pipetting 0.1

mL of sample (if liquid such as a thawed brain) to the tube. We placed the samples in a

BeadBeater for 3 minutes after which they were quickly centrifuged. We added 70 µL of

Proteinase K to the ATL lysate. We incubated the lysate at +56 °C for 1-3 hours during which

the tubes were intermittently inverted several times. We centrifuged the lysate quickly and

added 0.6 mL of AL buffer. We inverted the tubes several times and incubated them at +70

°C for 10-30 minutes in a dry block. We then again inverted the tubes intermittently several

times. The samples were then centrifuged for 3 minutes at 10,000 x *g*. After the completion of

the cell lysis, we used 100 µL of the ATL/ProtK/AL lysate to continue the extraction.

Nucleic acid extraction and real-time PCR.—We did nucleic acid extraction for *T.*

gondii using 5X MagMAX viral Isolation kit (Applied Biosystems AMB1836-5). Primers and

probes for real-time PCR were designed according to the protocol of De Craeye et al. (2011)

[29] (see Table S1 for primers and probes). PCR product size was 106 bp. We used cellular

r18S (ribosomal RNA gene) as an internal control of all PCRs. Real-time PCR was conducted

using TaqMan Fast Advanced Master Mix (Applied Biosystem) in Applied Biosystems™

7500 Real-Time PCR System. Each PCR reaction had concentrations of 10 µM of T2 /F

primer, 10 µM of T3/R primer, and 5 µM of probe in the master mix. Thermo cycling

Program: Initial denaturation and activation of the Taq polymerase at +95 °C for 2 minutes,

followed by 45 cycles at +95 °C for 5 seconds and +60 °C for 33 seconds. We analyzed the

results using 7500 System SDS Software.

Serum collection

To investigate *T. gondii* prevalence in squirrels during the acute stage we conducted live

trapping of grey squirrels in one urban and one rural site between 5[th] June and 1[st] August 2019

(Fig. 1). The urban site is located in the city of Winnipeg, Manitoba, Canada, and consists of

an ~10 ha park located on the University of Manitoba campus and a suburban neighborhood

next to the park. The study site is bordered by the Red River and two major highways with

high amounts of car traffic. The rural site is a ~34 ha forest patch next to an active honey-

farm, near the twin cities Morden and Winkler in southern Manitoba (49°24'01.1"N,

98°00'29.2"W), bordered by agricultural land. We used live traps (Tomahawk Live Trap Co.,

Tomahawk, WI, USA) to capture grey squirrels at the study sites. Between 20-40 traps in total

were set to the urban site and between 80-100 traps to the rural site each trapping day. Traps

were baited with peanut butter and checked regularly. The number of traps was different for

the two sites to ensure approximately the same number of squirrels was captured at both sites.

Traps were placed in sheltered locations under vegetation cover or covered with canvas to

provide shade and to calm the animals when in the trap. After capture, squirrels were handled

in a canvas capture bag and we recorded the weight (g), body and tail length (cm), skull width

(cm), age (adult or juvenile), reproductive status, and sex for each individual. We collected a

minimum of 500 µL of blood from the femoral vein of each grey squirrel and stored the

sample on ice until processing. Each squirrel was pit-tagged between the shoulder blades with

passive integrated transponder (PIT) tags. All efforts were made to minimize suffering. We

then released the squirrels at the place of capture. Our protocol (Protocol Number: f16-003)

was approved by the University of Manitoba animal care and use committee following

Canadian Council on Animal Care guidelines.

Serological Methods

We collected a minimum of 500 µL of blood from 15 individual grey squirrels and stored

samples on ice. All samples were processed within 12 hours. We centrifuged the collected

blood samples at 3500 rpm for 15 minutes and froze the serum at −20 °C until used for

testing.

*Enzyme-linked immunosorbent assay.—*We used enzyme-linked immunosorbent

assays (ELISA) to detect serum antibodies (IgG) against *T. gondii.* As species-specific

conjugates are not available for squirrels, we used a commercially available ELISA kit for

testing the samples (Multi-species ID Screen Toxoplasmosis Indirect kit, IDVet, Grabels,

France) following the manufacturer's instructions. We read the optical density values at

450 nm in a spectrophotometer and calculated results using these values and kit controls

expressed as S/P (Sample to Positive Ratio**)** percentage (S/P%). We considered samples with

S/P% less or equal to 40% negative; samples with S/P% between 40 and 50% doubtful or

inconclusive; and samples with an S/P% higher than 50% positive, following the kit's

protocol.

Results

We tested a total of 230 tissue samples from 46 squirrels from four squirrel species for *T.*

gondii DNA using quantitative PCR (Table 1). Twenty-six of the carcasses were from urban

locations within Winnipeg and 20 were from rural locations between 30 and 250 km from

Winnipeg (Fig. 1). We had 25 American red squirrels (*Tamiasciurus hudsonicus*); 16 eastern

grey squirrels (*Sciurus carolinensis*); four northern flying squirrels (*Glaucomys sabrinus*); and

one least chipmunk (*Tamias minimus*). *T. gondii* DNA was not detected in any of the 230

tissue samples (liver; heart; brain; lung; spleen; kidney).

We also tested a total of 13 (out of 15) samples of blood sera from grey squirrels for *T. gondii*

antibodies (IgG) by indirect ELISA. Two of the collected samples did not have enough

volume for testing. These results were negative—no *T. gondii* antibodies were detected in any

sample (see Table S2).

Discussion

We hypothesized that, due to higher population densities of both squirrels and cats in cities

[9,19], urban squirrels may act as important intermediate hosts of *T. gondii* and have a higher

prevalence of *T. gondii* infection than rural squirrels. However, we found no evidence of *T.*

gondii infection in squirrels.

T. gondii has been found in many wild animals including on farms and in natural areas (e.g.,

gray fox (*Urocyon cinereoargenteus*) [30–32], red fox (*Vulpes vulpes*) [33]). However,

studies investigating *T. gondii* infection dynamics in cities are still relatively rare [4,17, but

see e.g., 18,34–37] considering the parasite can infect most mammalian species, including

domestic cats and humans. Rodents can act as intermediate hosts for the parasite and as

sources of infection to cats if consumed when infected by *T. gondii*. Squirrels are among the

many species capable of getting a *T. gondii* infection, but our knowledge is limited, and

studies have mostly focused on acute, fatal cases with little information existing on host-

pathogen dynamics during chronic, latent infection [21].

The prevalence of *T. gondii* in Sciurids in urban areas is not well known. An earlier study

from Guelph, Ontario, Canada found no evidence of infection across nine locations within and

around the city from eastern grey squirrels (n= 16, the number of urban captures not

specified) and chipmunks (*Tamias striatus*, n= 6) using the Sabin-Feldman dye test [30]. In

natural areas, toxoplasmosis has been found in the eastern grey squirrel [20,38–40], western

grey squirrel (*Sciurus griseus*) [41], and Eurasian red squirrel (*Sciurus vulgaris*) [12,21].

However, *T. gondii* prevalence in squirrels has generally been low (indirect hemagglutination

test 1 of 265 positive individuals [42]; PCR 3 of 19 [21]; Sabin-Feldman dye test 2 of 11

[40]).

Using serological techniques with bioassay and PCR together can give a more reliable

estimation of infection rate [24], as *T. gondii* is a cyst-forming parasite, and the distribution of

T. gondii tissue cysts can be uneven and vary between organs [28]. This can lead to detection

difficulties using PCR-based methods [43]. The sensitivity and specificity of antibody

detection methods can also vary, which can lead to misinterpretation of results and possible

false negatives and positives [44]. The multi-species ELISA kit we used has been successfully

used to detect *T. gondii* antibodies in wildlife [45,46], has high sensitivity and specificity

compared to other serological tests such as the modified agglutination test [47], and does not

cross-react with other coccidian parasites—a factor known to limit the specificity of

serological assays [48]. As we sampled several different organs and tissues per individual by

PCR and used serological methods as an additional test to survey *T. gondii* prevalence in

squirrels our results are less likely to be false negatives.

T. gondii prevalence in Manitoba, in general, is not well known and no previous survey of *T.*

gondii prevalence in squirrels exists from the province. Serological testing in 1981 reported

that of 55,527 pregnant women 129 showed signs of a recent *T. gondii* infection (Sekla et al.

1981). The same study also reported that 19 of 72 cats and one polar bear (*Ursus maritimus*)

tested positive for *T. gondii* but results from 28 other species were all negative. An earlier

study from Manitoba in 1976 found that *T. gondii* prevalence in pregnant women in urban

areas was 8.17 %, and 6.29 % in rural areas [50]. The prevalence of *T. gondii* can be high in

domestic sheep, pigs, and cattle [e.g., 30,51, however, see 52], yet studies from Manitoba or

neighboring Saskatchewan have mostly found low prevalence [53,54].

The extreme climatic variations in the region may decrease the viability or infectivity of

oocysts and tissue cysts in carcasses, therefore decreasing overall *T. gondii* prevalence in the

area [53]. The temperature in Winnipeg between June to September can reach extremes of

+30 °C to +35 °C [26] with daily average temperatures varying from approximately +20 °C to

+13 °C [55]. *T. gondii* oocysts are highly resistant to environmental variation but sporulation

(i.e. infectivity) is dependent on fixed temperatures and factors such as soil moisture that can

influence the time oocysts can survive at high temperatures [56]. Additionally, winters in

Winnipeg can be cold and windy with extreme temperatures ranging between −24 °C and −33

°C between January to March. Oocysts cannot sporulate and become infective after exposure

to −21 °C for 1 day or −6 °C for 7 days [57]. After sporulation, oocysts can withstand lower

temperatures better, being able to survive at −21 °C for 28 days [57], yet oocysts can not

sporulate if the conditions are unfavorable [56]. It is thus possible that the combination of hot

summers and cold winters reduces the viability of oocysts in Manitoba. Contact rates between

domestic cats and *T. gondii* intermediate hosts may also be lower during the winter if people

keep their pets indoors in cold weather. Winnipeg has an average yearly precipitation of

521 mm, but the years 2018 and 2019 were unusually dry in southern Manitoba [58]. This

could have influenced oocyst survival in the area during data collection, as oocysts survive

better in moist than in dry conditions [57,59].

We note that *T. gondii* antibodies have been found in skunks (*Mephitis mephitis*) and

raccoons (*Procyon lotor*) from Saskatchewan (modified agglutination test: average

seroprevalence in skunks 15.6%; in raccoons 20.8% in 1999, 12.5% in 2000) and Manitoba

(in skunks 28%; in raccoons 27.5%) [60], which implies that although climatic variation may

lower the prevalence of *T. gondii* in the prairie provinces of Canada, *T. gondii* is still present

in the area. It is also possible that our negative findings are the result of infected individuals

dying from the disease. Nonetheless, our results in conjunction with previous studies suggest

that squirrels are not important intermediate hosts of *T. gondii* and likely do not act as

reservoirs for *T. gondii* in cities. This knowledge is important as squirrels tend to occur at

higher densities in cities than in more natural habitats which creates the possibility for

increased parasite transmission between squirrels and cats, and further, cats and humans.

When wildlife parasites are a human health concern, such as with *T. gondii*, then management

actions are warranted. Our results suggest that squirrels do not need to be considered as

sources of infection to cats. Consequently, no management efforts are needed in cities with

abundant squirrel populations. In general, our results shed light on the prevalence of *T. gondii*

in Sciurids in urban areas. This is important as *T. gondii* infection dynamics are still relatively

unknown in cities.

Acknowledgments

We want to thank Neil Pople, Amanda Salo, Niaz Rahim, and other members of the Manitoba

Veterinary Diagnostic Services Laboratory for their helpful guidance and data collection. We

thank Mitchell Green, Kyle Lefort, and Paul O'Brien for their help with necropsies, and

Constance Finney and Emily Jenkins for their excellent suggestions regarding the methods.

We want to give special thanks to Vera and Phil Froese for giving their permission to work on

their land.

References

1.   Mackenstedt U, Jenkins D, Romig T. The role of wildlife in the transmission of

290   parasitic zoonoses in peri-urban and urban areas. Int J Parasitol Parasites Wildl

291   2015;4:71–9. doi:10.1016/j.ijppaw.2015.01.006.

292   2.   Deplazes P, Hegglin D, Gloor S, Romig T. Wilderness in the city: The urbanization of

293   Echinococcus multilocularis. Trends Parasitol 2004;20:77–84.

294   doi:10.1016/j.pt.2003.11.011.

295   3.   Reperant LA, Hegglin D, Tanner I, Fischer C, Deplazes P. Rodents as shared indicators

296   for zoonotic parasites of carnivores in urban environments. Parasitology

297   2009;136:329–37. doi:10.1017/S0031182008005428.

298   4.   Lehrer EW, Fredebaugh SL, Schooley RL, Mateus-Pinilla NE. Prevalence of

299   antibodies to Toxoplasma gondii in woodchucks across an urban-rural gradient. J Wildl

300   Dis 2010;46:977–80. doi:10.7589/0090-3558-46.3.977.

301   5.   Giraudeau M, Mousel M, Earl S, McGraw K. Parasites in the city: Degree of

302   urbanization predicts poxvirus and coccidian infections in house finches (Haemorhous

303   mexicanus). PLoS One 2014;9. doi:10.1371/journal.pone.0086747.

304   6.   Gliwicz J, Goszczynski J, Luniak M. Characteristic features of animal populations

305   under synurbanization - the case of the blackbird and of the striped field mouse.

306   Memorab Zool 1994;49:237–44.

307   7.   Bradley CA, Altizer S. Urbanization and the ecology of wildlife diseases. Trends Ecol

308   Evol 2007;22:95–102. doi:10.1016/j.tree.2006.11.001.

309   8.   Baker PJ, Molony SE, Stone E, Cuthill IC, Harris S. Cats about town: Is predation by

310   free-ranging pet cats Felis catus likely to affect urban bird populations? Ibis (Lond

311   1859) 2008;150:86–99. doi:10.1111/j.1474-919X.2008.00836.x.

312   9.   Sims V, Evans KL, Newson SE, Tratalos JA, Gaston KJ. Avian assemblage structure

313   and domestic cat densities in urban environments. Divers Distrib 2008.

doi:10.1111/j.1472-4642.2007.00444.x.

10.   Elmore SA, Jones JL, Conrad PA, Patton S, Lindsay DS, Dubey JP. Toxoplasma

gondii: epidemiology, feline clinical aspects, and prevention. Trends Parasitol

2010;26:190–6. doi:10.1016/j.pt.2010.01.009.

11.   Dubey JP. Oocyst shedding by cats fed isolated bradyzoites and comparison of

infectivity of bradyzoites of the VEG strain Toxoplasma gondii to cats and mice. J

Parasitol 2001;87:215–9. doi:10.2307/3285204.

12.   Fayyad A, Kummerfeld M, Davina I, Wohlsein P, Beineke A, Baumgärtner W, et al.

Fatal systemic Toxoplasma gondii infection in a red squirrel (Sciurus vulgaris), a

Swinhoe's striped squirrel (Tamiops swinhoei) and a New World porcupine

(Erethizontidae sp.). J Comp Pathol 2016;154:263–7. doi:10.1016/j.jcpa.2016.02.002.

13.   Centers for Disease Control and Prevention. Toxoplasmosis 2019. Available from:

https://www.cdc.gov/parasites/toxoplasmosis/.

14.   Tenter AM, Heckeroth AR, Weiss LM. Toxoplasma gondii: From animals to humans.

Int J Parasitol 2000;30:1217–58. doi:10.1016/S0020-7519(00)00124-7.

15.   Dixon BR. Prevalence and control of toxoplasmosis - a Canadian perspective. Food

Control 1992;3:68–75. doi:10.1016/0956-7135(92)90034-8.

16.   Hill DE, Chirukandoth S, Dubey JP. Biology and epidemiology of Toxoplasma gondii

in man and animals. Anim Heal Res Rev 2005;6:41–61. doi:10.1079/ahr2005100.

17.   Conrad PA, Miller MA, Kreuder C, James ER, Mazet J, Dabritz H, et al. Transmission

of Toxoplasma: Clues from the study of sea otters as sentinels of Toxoplasma gondii

flow into the marine environment. Int J Parasitol 2005;35:1155–68.

doi:10.1016/j.ijpara.2005.07.002.

18.   Ballash GA, Dubey JP, Kwok OCHH, Shoben AB, Robison TL, Kraft TJ, et al.

Seroprevalence of Toxoplasma gondii in White-tailed deer (Odocoileus virginianus)

and free-roaming cats (Felis catus) across a suburban to urban gradient in northeastern

Ohio. Ecohealth 2015;12:359–67. doi:10.1007/s10393-014-0975-2.

19. Parker TS, Nilon CH. Gray squirrel density, habitat suitability, and behavior in urban

parks. Urban Ecosyst 2008. doi:10.1007/s11252-008-0060-0.

20. Dubey JP, Hodgin EC, Hamir AN. Acute Fatal Toxoplasmosis in Squirrels (Sciurus

carolensis) with Bradyzoites in Visceral Tissues. J Parasitol 2006;92:658–9.

doi:10.1645/GE-749R.1.

21. Jokelainen P, Nylund M. Acute fatal Toxoplasmosis in three Eurasian Red Squirrels

(Sciurus vulgaris) caused by genotype II of Toxoplasma gondii. J Wildl Dis

2012;48:454–7. doi:10.7589/0090-3558-48.2.454.

22. Dubey JP, Frenkel JK. Experimental toxoplasma infection in mice with strains

producing oocysts. J Parasitol 1973;59:505–12. doi:10.2307/3278784.

23. Afonso E, Thulliez P, Pontier D, Gilot-Fromont E. Toxoplasmosis in prey species and

consequences for prevalence in feral cats: Not all prey species are equal. Parasitology

2007;134:1963–71. doi:10.1017/S0031182007003320.

24. Galeh TM, Sarvi S, Montazeri M, Moosazadeh M, Nakhaei M, Shariatzadeh SA, et al.

Global status of Toxoplasma gondii seroprevalence in rodents: A systematic review

and meta-analysis. Front Vet Sci 2020;7:461. doi:10.3389/fvets.2020.00461.

25. Statistics Canada. Winnipeg, CY, Manitoba. Census Profile. 2016 Census. Statistics

Canada Catalogue no. 98-316-X2016001. 2016. Available from:

https://www12.statcan.gc.ca.

26. Environment Canada. Canadian Climate Normals 1981–2010 Station Data for

Winnipeg. 2020. Available from:

https://climate.weather.gc.ca/climate_normals/results_1981_2010_e.html.

27. Kahle D, Wickham H. ggmap: Spatial visualization with ggplot2. R J 2013;5:144–61.

doi:10.32614/rj-2013-014.

28. Elmore SA, Huyvaert KP, Bailey LL, Iqbal A, Su C, Dixon BR, et al. Multi-scale

occupancy approach to estimate Toxoplasma gondii prevalence and detection

probability in tissues: an application and guide for field sampling. Int J Parasitol

2016;46:563–70. doi:10.1016/j.ijpara.2016.04.003.

29. De Craeye S, Speybroeck N, Ajzenberg D, Dardé ML, Collinet F, Tavernier P, et al.

Toxoplasma gondii and Neospora caninum in wildlife: Common parasites in Belgian

foxes and Cervidae? Vet Parasitol 2011;178:64–9. doi:10.1016/j.vetpar.2010.12.016.

30. Tizard IR, Harmeson J, Lai CH. The prevalence of serum antibodies to Toxoplasma

gondii in Ontario mammals. Can J Comp Med Rev Can Med Comp 1978;42:177–83.

31. Smith DD, Frenkel JK. Prevalence of antibodies to Toxoplasma gondii in wild

mammals of Missouri and east central Kansas: biologic and ecologic considerations of

transmission. J Wildl Dis 1995;31:15–21. doi:10.7589/0090-3558-31.1.15.

32. Lindsay DS, Weston JL, Little SE. Prevalence of antibodies to Neospora caninum and

Toxoplasma gondii in gray foxes (Urocyon cinereoargenteus) from South Carolina. Vet

Parasitol 2001;97:159–64. doi:10.1016/S0304-4017(01)00390-9.

33. Wanha K, Edelhofer R, Gabler-Eduardo C, Prosl H. Prevalence of antibodies against

Neospora caninum and Toxoplasma gondii in dogs and foxes in Austria. Vet Parasitol

2005;128:189–93. doi:10.1016/j.vetpar.2004.11.027.

34. Frenkel JK, Hassanein KM, Hassanein RS, Brown E, Thulliez P, Quintero- Nunez R.

Transmission of Toxoplasma gondii in Panama City, Panama: A five-year prospective

cohort study of children, cats, rodents, birds, and soil. Am J Trop Med Hyg

1995;53:458–68. doi:10.4269/ajtmh.1995.53.458.

35. Dubey JP, Dennis PM, Verma SK, Choudhary S, Ferreira LR, Oliveira S, et al.

Epidemiology of toxoplasmosis in white tailed deer (Odocoileus virginianus):

Occurrence, congenital transmission, correlates of infection, isolation, and genetic

characterization of Toxoplasma gondii. Vet Parasitol 2014;202:270–5.

doi:10.1016/j.vetpar.2014.01.006.

36. Mercier A, Garba M, Bonnabau H, Kane M, Rossi JP, Dardé ML, et al. Toxoplasmosis

seroprevalence in urban rodents: A survey in Niamey, Niger. Mem Inst Oswaldo Cruz

2013;108:399–407. doi:10.1590/S0074-0276108042013002.

37. Murphy RG, Williams RH, Hughes JM, Hide G, Ford NJ, Oldbury DJ. The urban

house mouse (Mus domesticus) as a reservoir of infection for the human parasite

Toxoplasma gondii: An unrecognised public health issue? Int J Environ Health Res

2008;18:177–85. doi:10.1080/09603120701540856.

38. Jacobs L, Stanley AM, Herman CM. Prevalence of Toxoplasma antibodies in rabbits,

squirrels, and raccoons collected in and near the Patuxent Wildlife Research Center. J

Parasitol 1962;48:550. doi:10.2307/3274906.

39. Walton BC, Walls KW. Prevalence of toxoplasmosis in wild animals from Fort

Stewart, Georgia, as indicated by serological tests and mouse inoculation. Am J Trop

Med Hyg 1964;13:530–3. doi:10.4269/ajtmh.1964.13.530.

40. Smith DD, Frenkel JK. Prevalence of Antibodies To Toxoplasma Gondii in Wild

Mammals of Missouri and East Central Kansas: Biologic and Ecologic Considerations

of Transmission. J Wildl Dis 1995;31:15–21. doi:10.7589/0090-3558-31.1.15.

41. Soave OA, Lennette EH. Naturally acquired toxoplasmosis in the gray squirrel, sciurus

griseus, and its bearing on the laboratory diagnosis of rabies. J Lab Clin Med

1959;53:163–6. doi:10.5555/uri:pii:0022214359900642.

42. Burridge MJ, Bigler WJ, Forrester DJ, Hennemann JM. Serologic survey for

Toxoplasma gondii in wild animals in Florida. J Am Vet Med Assoc 1979;175:964–7.

43. Opsteegh M, Langelaar M, Sprong H, den Hartog L, De Craeye S, Bokken G, et al.

Direct detection and genotyping of Toxoplasma gondii in meat samples using magnetic

capture and PCR. Int J Food Microbiol 2010;139:193–201.

doi:10.1016/j.ijfoodmicro.2010.02.027.

44.   Gilbert AT, Fooks AR, Hayman DTS, Horton DL, Müller T, Plowright R, et al.

Deciphering serology to understand the ecology of infectious diseases in wildlife.

Ecohealth 2013;10:298–313. doi:10.1007/s10393-013-0856-0.

45.   Roqueplo C, Halos L, Cabre O, Davoust B. Toxoplasma gondii in wild and domestic

animals from New Caledonia. Parasite 2011;18:345–8.

doi:10.1051/parasite/2011184345.

46.   Sharma R, Parker S, Elkin B, Mulders R, Branigan M, Pongracz J, et al. Risk factors

and prevalence of antibodies for Toxoplasma gondii in diaphragmatic fluid in

wolverines (Gulo gulo) from the Northwest Territories, Canada. Food Waterborne

Parasitol 2019;15:e00056. doi:10.1016/j.fawpar.2019.e00056.

47.   Sharma R, Parker S, Al-Adhami B, Bachand N, Jenkins E. Comparison of tissues

(heart vs. brain) and serological tests (MAT, ELISA and IFAT) for detection of

Toxoplasma gondii in naturally infected wolverines (Gulo gulo) from the Yukon,

Canada. Food Waterborne Parasitol 2019;15:e00046.

doi:10.1016/j.fawpar.2019.e00046.

48.   Hirota J, Nishi H, Matsuda H, Tsunemitsu H, Shimizu S. Cross-reactivity of Japanese

encephalitis virus-vaccinated horse sera in serodiagnosis of West Nile Virus. J Vet Med

Sci 2010;72:369–72. doi:10.1292/jvms.09-0311.

49.   Sekla, L, Stackiw W, Rodgers S. A serosurvey of toxoplas-mosis in Manitoba. Can J

Public Heal 1981;72:111–7.

50.   Shettigara PT, Choi NW, Abu-Zeid HAH. Prevalence of toxoplasmosis in Manitoba

province: a seroepidemiologic study of 23 146 prenatal sera. Am J Epidemiol

1976;104:340–340.

51.   Fayer R. Toxoplasmosis update and public health implications. Can Vet J

441   1981;22:344–52.

442   52.   Poljak Z, Dewey CE, Friendship RM, Martin SW, Christensen J, Ojkic D, et al. Pig and

443   herd level prevalence of Toxoplasma gondii in Ontario finisher pigs in 2001, 2003, and

444   2004. Can J Vet Res 2008;72:303–10.

445   53.   Nation PN, Allen JR. Antibodies to Toxoplasma gondii in Saskatchewan cats, sheep

446   and cattle. Can Vet J 1976;17:308–10.

447   54.   Smith HJ. Seroprevalence of anti-Toxoplasma IgG in Canadian swine. Can J Vet Res

448   1991;55:380–1.

449   55.   Environment Canada. Monthly Data Report for 2007 for Winnipeg The Forks,

450   Manitoba. 2020. Available from:

451   https://climate.weather.gc.ca/climate_data/monthly_data_e.html.

452   56.   Dubey JP, Miller NL, Frenkel JK. The toxoplasma gondii oocyst from cat feces. J Exp

453   Med 1970;132:636–62. doi:10.1084/jem.132.4.636.

454   57.   Frenkel JK, Ruiz A, Chinchilla M. Soil survival of Toxoplasma oocysts in Kansas and

455   Costa Rica. Am J Trop Med Hyg 1975;24:439–43. doi:10.4269/ajtmh.1975.24.439.

456   58.   Government of Manitoba. Manitoba Drought Management Strategy. 2020. Available

457   from: https://www.gov.mb.ca/sd/water/drought_condition/index.html.

458   59.   Lélu M, Villena I, Dardé ML, Aubert D, Geers R, Dupuis E, et al. Quantitative

459   estimation of the viability of Toxoplasma gondii oocysts in soil. Appl Environ

460   Microbiol 2012;78:5127–32. doi:10.1128/AEM.00246-12.

461   60.   Hwang YT, Pitt JA, Quirk TW, Dubey JP. Seroprevalence of Toxoplasma gondii in

462   mesocarnivores of the Canadian prairies. J Parasitol 2007;93:1370–3. doi:10.1645/GE-

463   1319.1.

Tables and Figures

Table 1. The number of individuals and the number of tissue samples from each squirrel

species from rural and urban sampling locations tested for *Toxoplasma gondii* DNA using

quantitative PCR. The common and scientific names of species are listed, as are the total

numbers of PCR samples tested for each species, and the total number of individuals and

samples in each sampling location.

Species	Scientific name	#squirrels	#squirrels	#samples	#samples	Total PCR
		Rural	Urban	Rural	Urban	
Red squirrel	*Tamiasciurus hudsonicus*	12	13	72	77	149
Grey squirrel	*Sciurus carolinensis*	4	12	8	44	52
Least chipmunk	*Tamias minimus*	0	1	0	5	5
Northern flying squirrel	*Glaucomys sabrinus*	4	0	24	0	24
Total		**20**	**26**	**104**	**126**	**230**

Figure 1. Map showing the sampling locations of squirrel (Sciuridae) carcasses (circles) used

to collect tissue samples for quantitative PCR detection of *Toxoplasma gondii*, and the urban

and rural study sites for blood collection (triangles) from grey squirrels (*Sciurus carolinensis*)

for serological testing of *Toxoplasma gondii* antibodies. The map on the left shows the urban

and rural locations of squirrels from in and around the city of Winnipeg, Manitoba, Canada,

and the map on the right is a close-up of the urban locations within the city perimeter. The

map was created using an R package *ggmap* [27]. Map tiles by Stamen Design, under CC BY

3.0. Data by OpenStreetMap, under ODbL.

483