Restructured mitochondrial-nuclear interaction in *P. falciparum* dormancy and persister survival of artemisinin exposure

Sean V. Connelly[1], Javier Manzella-Lapeira[2], Zoë C. Levine[1], Joseph Brzostowski[2], Ludmila Krymskaya[2], Rifat S. Rahman[1], Juliana M. Sá[1], Thomas E. Wellems[1*]

[1] Laboratory of Malaria and Vector Research (LMVR), National Institute of Allergy and Infectious Diseases (NIAID), National Institutes of Health (NIH), Bethesda, MD

[2] Laboratory of Immunogenetics (LIG), National Institute of Allergy and Infectious Diseases (NIAID), National Institutes of Health (NIH), Bethesda, MD

*Corresponding author

TEW: twellems@niaid.nih.gov

ABSTRACT

Artemisinin and its semi-synthetic derivatives (ART) are fast acting, potent antimalarials; however, their use in malaria treatment is frequently confounded by recrudescences from bloodstream *Plasmodium* parasites that enter into and later reactivate from a dormant persister state. Here we show that the mitochondria of dihydroartemisinin (DHA)-exposed persisters are dramatically altered and enlarged relative to the mitochondria of young, actively replicating ring forms. Persister forms exhibit restructured mitochondrial-nuclear associations and an altered metabolic state consistent with stress from reactive oxygen species. New contacts between the mitochondria and nuclei may support communication pathways of mitochondrial retrograde signaling, resulting in transcriptional changes in the nucleus as a survival response. Further characterization of the organellar interactions and metabolic dependencies of persisters may suggest strategies to combat recrudescences of malaria after treatment.

Introduction

Artemisinin, from the Chinese natural Qinghaosu remedy, provides the basis for first line treatments of malaria worldwide, particularly for the deadliest form caused by *Plasmodium falciparum* [1]. Artemisinin and its semi-synthetic derivatives (collectively abbreviated here as ART) are among the best antimalarials for rapid clearance of parasitemia and resolution of illness [2, 3]. Yet, despite these invaluable qualities, lasting cures from ART treatment are not assured: since the time of its introduction, frequent recrudescences have been reported after ART monotherapy, necessitating the utilization of a partner drug for their prevention [4]. Various studies have identified dormant persister forms from erythrocytic-stage populations that give rise to these recrudescences [5-8]. Much remains to be understood about the nature of these persisters and how they develop from only a small fraction (estimated less than ~1% [5, 7]) of the circulating ART treated parasite population.

Although dormant hypnozoites in the liver stage cycle of certain *Plasmodium* spp. were well known, evidence was not reported until 1995 that dormant parasites could be found in erythrocytes after drug and toxin exposure. Nakazawa *et al.* [9] demonstrated the presence of a small number of the dormant parasites following destruction of the great majority of actively replicating parasites by D-sorbitol treatments or exposure to pyrimethamine. Importantly, the dormant parasites were able to survive these exposures for several days, after which the dormant parasites gave rise to recrudescent populations that were as susceptible to the

treatments as the original populations. Further, no relationship could be

demonstrated between the incidence or timing of recrudescence and drug

concentration to which the original population was exposed [10].

Microscopic images of dormant forms after ART exposure were presented by

LaCrue *et al.* [6] and Tucker *et al.* [8]: in Giemsa-stained blood films, these forms

were distinguished from actively replicating ring stages and pyknotic parasites by

their small rounded appearance with magenta colored chromatin and condensed

blue cytoplasm. The importance of mitochondrial membrane potential to persister

viability was demonstrated by Rhodamine 123 staining and outgrowth

experiments [11]. Although their metabolic pathways were found to be largely

downregulated after ART exposure, dormant parasites demonstrated

upregulation of some enzymes involved in fatty acid biosynthesis, pyruvate

metabolism, and the isoprenoid pathway [12, 13]. Changes in the expression of

cyclin and cyclin dependent kinase (CDK) genes have also been reported with

dormancy and reactivation of persister forms [14].

Recent advances in methods for isolation of persister parasites, imaging of their

subcellular organelles, and metabolic activity analysis offer approaches for new

insights into these important *Plasmodium* stages. The application of the AiryScan

principle in laser scanning microscopy provides increased resolution for organelle

level imaging of individual cells, as compared to standard confocal microscopy,

with improved signal-to-noise ratio from 32 detectors that collect emitted light

usually excluded [15]. Modern methods of fluorescence lifetime imaging (FLIM)

with exogenous or endogenous (autofluorescent) fluorophores enable non-

invasive characterization of the metabolic status of cells. In the technique of time

domain autofluorescence FLIM, a laser scanning microscope (LSM) with a

pulsed multiphoton laser excites the endogenous fluorophore and records the

distribution of photons that are emitted over time from each pixel in an image [16,

17]. The metabolic coenzyme NADH has specific mean fluorescence lifetimes

based on its free or protein bound state, allowing detection of metabolic changes

and providing insight on the redox status of tissues [18]. Phasor analysis of raw

FLIM data provides a 2D graphical approach that does not require fitting to a

decay model, and it correlates each pixel in an image to a position in phasor

space [19]. For example, metabolic characterization through the phasor

approach was applied to germ cells, bacteria and primary keratinocyte cultures to

assay the response of these cells to metabolic perturbations [20-22].

Here we compare the structural and metabolic phenotypes of persisters to those

of actively replicating ring forms in two *P. falciparum* lines: GB4, an African clone

of Ghanaian origin; and 803, a clonal line from Cambodia. These two lines differ

by the *pfk13*-determined survivals of early (0–3 h post invasion) ring-stage

parasites exposed to a 6 hour pulse of physiological level DHA (in ring-stage

survival assays, or RSAs) [23]. Nevertheless, after exposure to these pulses of

DHA, GB4 and 803 populations both develop persister parasite forms that show

changes in the morphological features and volumes of mitochondria as well as

similar tight associations between the mitochondria and nuclei. In

autofluorescence FLIM-phasor analysis of GB4 and 803 persisters relative to

actively replicating ring forms, metabolic shifts are consistent with damage

produced by DHA in the mitochondria, altered electron transport chain function,

and increased free NADH and reactive oxygen species (ROS). These effects are

discussed in context of what is known from findings with other biological systems

and insights they may offer into the persister survival mechanism.

Results

Prolongation of the intraerythrocytic cycle occurs in *P. falciparum* lines

exposed to DHA, irrespective of K13 mutation and ring survival phenotype

For the experiments of this report, four lines were chosen based on K13

genotype. Two of these lines were utilized in a genetic cross that differed in DHA

ring survival phenotypes: GB4 with Kelch propeller wild type K13, and 803 with

the K13 C580Y propeller polymorphism [23]. The other two lines are an isogenic

pair differing at the K13 locus: 76H10 with the C580Y polymorphism, and

76H10^{C580Rev} with an engineered C580 wild-type (revertant) codon. Assays of

ring-stage survival were performed with these four lines as previously described

[24]. Zero-to-three hour ring-stage parasites were exposed for 6 hours to control

vehicle (0.1% DMSO) or 700 nM DHA/0.1% DMSO treatment at $t = 0$ hours and

percentage survivals were counted at $t = 72$ hours (Figure 1A). In these RSAs,

K13 C580Y parasites exhibited a higher proportion of parasites that survived

DHA treatment than C580 parasites, a phenotype previously linked to C580Y

inheritance [23].

Stage distributions were assessed among the survivors in the control and DHA-

treated parasites of each line at t = 72 hours (Figure 1B). At this timepoint, all

four parasite lines exhibited a higher proportion of ring-stage parasites after DHA

than after control vehicle exposure, suggesting that the peak beginning of the 2^{nd}

cell cycle had been delayed until 52-56 hours after DHA treatment (Figure 1C).

These findings are consistent with a previous report of ART exposure inducing a

delay in the erythrocytic cycle of *P. falciparum* [25]. In another study,

transcriptomic data of *P. falciparum* isolates from patients with acute malaria

likewise showed evidence for decelerated ring-stage development after ART

treatment [26].

A fluorescence sensor of mitochondrial potential identifies viable

erythrocytic-stage GB4 and 803 parasites after DHA treatment

Peatey *et al.* [11] described the use of Rhodamine 123, a mitochondrial potential

marker, with Fluorescence Activated Cell Sorting (FACS) to identify and isolate

viable parasites after DHA treatment of synchronized ring stages. Among these

viable parasites were two distinct populations: (1) dormant persister forms; and

(2) actively growing stages that continued to progress through the cell cycle and

could be removed by sequential captures in magnetic separation columns. We

therefore adopted magnetic capture to enrich for persister forms after DHA

exposure. GB4 or 803 parasites at 0 – 3 hours ring-stage (post-invasion) were

placed in cultures at timepoint $t = 0$ hours with control vehicle (0.1% DMSO) or

700 nM DHA/0.1% DMSO. Control or DHA-treated samples were washed twice

at $t = 6$ hours and placed back into culture. At $t = 30$ hours, the DHA-treated

samples were passed through a magnetically activated cell sorting (MACS)

column to deplete mature-stage, hemozoin-containing parasites that had

survived drug treatment, reducing but not completely removing the contribution of

these stages from reinvasion into the next cycle. To isolate and compare the

populations of control vehicle- and DHA-exposed GB4 and 803 parasite

populations at the beginning of the 2nd erythrocytic-stage cycle, we performed

FACS at $t = 50$ hours with SYBR Green I (SG) as a DNA dye and MitoTracker

Deep Red FM (MT) as a mitochondrial membrane potential marker.

Representative FACS dot plots of the GB4 and 803 samples at $t = 50$ hours are

shown in Figure 2. In each experiment, uninfected erythrocytes that lack SG and

MT signals present in the lower left quadrant of the plot. In the upper left

quadrant of the plot, the gate region labeled "PYK" counts pyknotic parasites that

lack mitochondrial potential whereas, in the upper right quadrant of the plot, the

gate region labeled MT+ counts parasites that exhibit mitochondrial activity. In

the MT+ gate region, increasing SG and MT signals moving upward and to the

right arise from the progression of young ring stages to mature stage parasites,

as has been described with fluorescent markers in previous studies [27]. Clusters

of ring and late stage schizonts are evident in the control populations of both

GB4 and 803. Referring to the timeline (Fig. 1C), the clusters of schizonts derive from the end of the first erythrocytic-stage cycle while the more numerous ring stages derive from the beginning of the second cycle. FACS results from the DHA-treated parasites show large numbers of pyknotic parasites that lack mitochondrial potential in the "PYK" region, as expected from drug killing. Surviving parasites, however, are evident in both the GB4 and 803 populations, with greater counts of 803, consistent with their higher RSA survival levels compared to GB4 parasites (Fig. 1A). Thus, in the DHA-treated 803 MT+ distribution, the presence of trophozoites from the delay in the first intraerythrocytic cycle contributes to the appearance of a more continual representation of developmental stages across the intensity spectrum.

While the SG+MT+ subsets of the GB4 and 803 control populations contained predominantly ring-stage forms, the DHA-treated parasites after passage through the magnetic column at $t = 30$ hrs were enriched for persister forms [11].Thus, FACS at $t = 50$ hours identified a distribution of persisters as well as rings and mature stages that escaped removal by the single magnetic column. From this distribution, we were able to use the criteria of size and fluorescence intensity to identify and individually study these different forms.

AiryScan Imaging Reveals Organelle Level Changes in GB4 and 803 Persisters

Isolated persister and ring-stage forms were imaged by high-resolution AiryScan

Microscopy (ASM) and compared for fluorescent mitochondrial and nuclear

volumes, the shapes of these volumes as indicators of organellar morphology,

and the mean distances between these volumes as indicators of relative mito-

nuclear positioning.

Figure 3 shows representative surface renderings of the volumes of nuclear DNA

and mitochondrial fluorescence regions in three separate experiments with

control or DHA-treated GB4 and 803 parasites. Examples of distance

determinations between the centers of volume are shown in Figure 3 – Figure

Supplement 1. In the control GB4 and 803 samples, the nuclear DNA (green)

and mitochondrial (red) volumes of the ring forms were clearly separated from

one another. Nuclear DNA volumes presented with an oblate or boxy

appearance and were larger in volume than the mitochondria. The mitochondrial

volumes exhibited an oblate appearance, consistent with a recent account of

localized mitochondrial regions in ring-stage parasites [28]. Long tubular

structures [29] were not characteristic of the MT-stained mitochondria in this

study.

After DHA treatment of the synchronized GB4 parasites followed by magnetic

column separation at $t = 30$ hrs, actively growing parasites were mostly

eliminated, and surviving persisters greatly predominated in the DHA-treated

GB4 population captured for analysis at $t = 50$ hours. Figure 3A (right panel)

presents the nuclear and mitochondrial morphologies of six parasites from this

population: one exhibited a smooth oblate mitochondrion separate from the

nucleus as is characteristic of an actively growing ring-stage form, whereas the

five others had distinctly different appearances, with rumpled and corrugated

mitochondria in close approximation to the nuclei. The mitochondria partially

enwrapped the nuclei and were often enlarged compared to the control ring-

stage parasites. Contact sites appeared to be present between the nuclear DNA

and mitochondria as another potential feature of the persister phenotype.

Because single column separation at $t = 30$ hrs did not completely remove

actively replicating parasites from the DHA-treated 803 parasites (Figure 2,

bottom right panel; three daily column separations would have been required for

more complete isolation of persisters [7]), the parasites imaged by ASM at $t = 50$

hours comprised a mixed population of persister and ring-stage forms. In

comparison to the DHA-treated GB4 population, images from the DHA-treated

803 population showed more frequent examples of cells with smooth surfaced

mitochondria separate from the nuclei, similar to those of the control 803

population (Figure 3B, right panel). Other images of DHA-treated 803 parasites

showed rumpled, corrugated mitochondria in close approximation to nuclei as

observed for the DHA-treated GB4 persisters (Figure 3B, right panel). These

findings confirmed the presence at $t = 50$ hours of a mixed population of

persisters and actively replicating ring stages in the DHA-treated 803 sample,

consistent with the results shown in Figure 2.

Population measures of mito-nuclear distance and organelle volumes in

persisters and actively replicating parasites

To quantify the phenotypes of persisters *vs.* actively replicating ring-stage

parasites, mito-nuclear distances, mitochondrial volumes, and nuclear volumes

were determined from parasites of the control vehicle- and DHA-treated GB4 and

803 populations. For GB4 parasites (Figure 4A, left), a mito-nuclear median

distance and interquartile range (IQR) of 0.56 (0.32 – 0.98) µm was obtained

from the control, untreated population of ring-stages (n = 167), whereas a much

smaller median distance of 0.19 (0.11 – 0.28) µm was obtained from the

population of DHA-treated GB4 parasites that predominantly consisted of

persisters (n = 167) (p < 0.0001, unpaired t-test). For 803 parasites (Figure 4B,

left), the control vehicle-treated population of ring-stages presented a mito-

nuclear median distance and interquartile range (IQR) of 0.39 (0.25 – 0.53) µm (n

= 190), while the median distance of 0.31 (0.17 – 0.49) µm (n = 158) in DHA-

treated 803 parasites was not significantly smaller (p = 0.50, unpaired t-test). The

insignificance of this difference between the control and DHA-treated 803

populations is consistent with presence of mixed persisters and surviving active

ring-stage forms in the DHA-treated 803 population.

Mitochondrial volume comparisons of control and DHA-exposed GB4 or 803

parasites revealed an increase in the mitochondrial volume after DHA treatment.

In control GB4 parasites (Figure 4A, right), the median mitochondrial volume and

IQR was 0.26 (0.20 – 0.34) μm^3, which was much smaller than the post DHA

treatment mitochondrial median volume of 1.30 (0.64 – 1.9) μm^3, representing a

5-fold increase (p < 0.0001, unpaired t-test). In control 803 parasites, the median

mitochondrial volume was 0.31 (0.20 – 0.43) μm^3 while the DHA-exposed

median volume was 0.54 (0.23 – 1.5) μm^3 (Figure 4B, right) (p < 0.0001,

unpaired t-test). Compared to the results from GB4 parasites, the less dramatic

increase of median mitochondrial volume in DHA-exposed 803 parasites at $t = 50$

hours may be explained by its mixed population of persisters, with large

mitochondrial volumes, plus actively replicating ring forms, with smaller

mitochondrial volumes. These findings with GB4 and 803 parasites are

consistent with the increased mitochondrial volumes that have been reported

from other studies of ART-treated *Plasmodium* and *Toxoplasma gondii* [30, 31].

For GB4 parasites, the median nuclear DNA volume in the control population

was 1.1 (0.85 – 1.4) μm^3, whereas DHA-exposed parasites presented a smaller

median volume of 0.83 (0.57 – 1.1) μm^3 (Figure 4 – Figure Supplement 1) (p =

0.0012, unpaired t-test). In control 803 parasites, the median nuclear DNA

volume was 1.8 (1.2 – 2.2) μm^3 and in DHA-treated 803, a slightly larger median

nuclear DNA volume was observed, 2.1 (1.5 – 2.8) μm^3 (Figure 4 – Figure

Supplement 1) (p < 0.0001, unpaired t-test).

Isolation and Characterization of Persister Parasites 5 and 11 Days after

DHA Treatment

After examining the morphological characteristics of persister mitochondria and

quantifying their close apposition to nuclei at $t = 50$ hours, we looked for the

presence of these features in persisters 1 – 2 weeks old, before their

recrudescence in culture. Accordingly, we followed the *in vitro* cultures of

synchronized 803 and GB4 parasites after treatment with 700 nM DHA for 6

hours and three successive 5% D-Sorbitol treatments at 24, 48 and 72 hours, as

described previously [32]. Figure 4C presents the resulting dormancy and

recrudescence time courses, beginning from parasitemias of 2% upon DHA

exposure, through the dormancy period, to recrudescence of the 803 and GB4

parasitemias 16 and 20 days later. Samples of the GB4 and 803 persister

populations were sorted, and MT-positive subsets were collected at day 5 and

day 11 for imaging and analysis.

Representative ASM volume images of GB4 and 803 parasites are shown in

Figure 4D for day 5 and Figure 4E for day 11. On day 5, persister morphology

was observed in the 803 line, as evidenced by the large and rumpled

mitochondrial volumes enwrapping the nucleus (Figure 4D). On day 11, this

persister morphology continued to characterize the GB4 and 803 parasites and

was similar to the morphology observed earlier, at $t = 50$ hours and at 5 days.

However, 803 parasites of the day 11 population were also found with smaller

oblate mitochondria separate from the nucleus, *i.e.* with features that resembled

the actively growing ring-stage forms of the original 803 control group (Figure 4E,

compared with Figure 3B). Such evidence for actively growing ring-stage forms

among persisters in the 11-day 803 population is consistent with waking from

dormancy at the beginning of the recrudescence curve shown in Figure 4C.

In support of the presence of active ring stages wakened from 803 persisters, the

median mito-nuclear distance of the 803 population at 11 days (0.24 (0.15 –

0.39) µm) was greater than that of the GB4 population (0.20 (0.13 – 0.28) µm) (p

= 0.0187, unpaired t-test) (Figure 4B, left and Figure 4A, left). Before the waking

events, ASM of the 803 parasites was performed on day 5 after DHA treatment

and the small median distance typical of a pure persisters population was

observed (0.18 (0.087 – 0.26) µm) (Figure 4B). On day 11, the median

mitochondrial volume of GB4 persisters (2.1 (1.6 – 2.7) µm^3) was larger than that

of 803 parasites (0.71 (0.24 – 1.6) µm^3) (p < 0.0001, unpaired t-test) (Figure 4A,

right and Figure 4B, right). Consistent with the persister phenotype, a larger

median mitochondrial volume of 1.5 (1.2 – 1.8) µm^3 was observed in the 803

persister parasites on day 5 *vs.* day 11 (Figure 4B, right) (p < 0.0001, unpaired t-

test). In terms of nuclear volume on day 11 (Figure 4 – Figure Supplement 1), the

GB4 persister median fluorescence volume of 1.5 (1.1 – 1.9) µm^3 was

insignificantly different from the median volume of persister 803 parasites (1.6

(1.2 – 2.2) µm^3; p = 0.09), unpaired t-test). However, on day 5, persister 803

parasites had a smaller nuclear volume of 1.2 (1.1 – 1.5) µm^3 than on day 11 (p <

0.0001, unpaired t-test).

Metabolic Phenotyping of GB4 and 803 Parasites at *t* = 50 Hours

FLIM as a metabolic imaging technique detects the state of native fluorophores,

particularly nicotinamide adenine dinucleotide (NADH). NADH, a metabolic

coenzyme can be used as an autofluorescence indicator of the redox state of

cells [33], because this molecule has characteristic emission spectrum in its

enzyme bound vs. free state [34]. These two states have characteristic

fluorescence lifetimes in FLIM, with enzyme bound NADH having a longer

fluorescence lifetime and free NADH having a shorter fluorescence lifetime [18].

To study parasites for autofluorescence FLIM, control and DHA-treated GB4 or

803 parasites were stained only with MT to avoid spectral overlap with the broad

SG spectrum [35], and MT fluorescence-positive parasites were sorted by FACS,

allowed to adhere to a poly-lysine-coated PDMS microwell, and imaged in the

DIC and MT emission channels of the confocal microscope (Figure 5A). At least

ten million photon events were counted while limiting the time of exposure of

these parasites to 750 nm excitation. Comparison of intensity images in the same

field of view before and after FLIM acquisition confirmed little or no

photobleaching. To process each field of view, a masking strategy was used to

isolate the photon distributions for only ring-stage or persister forms, and the

FLIM data from each pixel were collected in the region of interest.

Phasor analysis of the raw autofluorescence FLIM data [20] was utilized to

assess the ratio of bound vs. free NADH in a cell or organism (Figure 5A).

Figure 5B presents FLIM-phasor comparisons of actively replicating control *vs.*

DHA-treated populations of GB4 and 803 parasites at $t = 50$ hours. In three

separate comparisons performed on different days, the phasor space

distributions of the data from the individual parasites were plotted, and asterisks

were used to mark the mean phasor positions of the control and DHA-treated

parasite distributions. In the GB4 comparisons (Figure 5B, left panel), the phasor

distribution of the DHA-treated population was shifted to the right of the untreated

actively growing control ring stages, indicative of a relatively greater level of free

NADH and diminished metabolic state of the persister phenotype (control: (0.75,

0.22), DHA: (0.80, 0.19); $p < 0.0001$, Pillai trace). In the 803 results (Figure 5B,

right panel), the DHA-treated population had only a higher S coordinate than the

control group (control: (0.63, 0.14), DHA: (0.63, 0.16); $p < 0.05$, Pillai trace). Little

or no shift to the right in the phasor position of the DHA-treated 803 population

was consistent with the presence of metabolically active ring forms along with

dormant persisters in this population at $t = 50$ hours.

Metabolic Phenotypes of GB4 and 803 Persisters Prior to Recrudescence *in*

vitro

Autofluorescence FLIM and phasor analysis were used to also examine the

metabolic states of the DHA/sorbitol treated GB4 and 803 parasites on day 5

(Figure 5C, left panel) and day 11 of dormancy (Figure 5C, right panel). On these

days, the mean coordinates of the GB4 and 803 distributions were located on the

right side of the phasor plot. Although the coordinates of the mean phasor

positions from each assessment may be affected by particulars of the culture

conditions and experimental manipulations, we note the large shift of the mean

803 signal from its position at $t = 50$ hours, consistent with the higher level of free

NADH in the purer population of persisters (DHA $t = 50$ hours coordinates (0.63,

0.16) *vs.* DHA day 5 coordinates (0.77, 0.21); $p < 0.0001$, Pillai Trace; compare

Figures 5B, right panel and 5C, left panel). In the 803 line, the calculated mean

position of the distribution at day 11 trended to the left of the position at day 5

(DHA day 5 coordinates (0.77, 0.21) *vs.* DHA day 11 coordinates (0.74, 0.22); p

= 0.05, Pillai Trace; compare Figure 5C, right panel and 5C, left panel). This

trend suggests that a relatively greater level of bound NADH was developing on

day 11, consistent with the aforementioned presence of active ring forms

recrudescing from the 803 persister population.

Discussion

Subsequent to early proposals that dormant erythrocytic stages are responsible

for recrudescences of *P. falciparum* infection after treatment with artemisinin

drugs [5], *in vitro* as well as *in vivo* studies continued to build evidence for the

importance of these parasite forms in failures of the artemisinins to achieve

complete cure [6-8, 23, 32, 36, 37]. Dormant parasites were found to be small

and to have characteristic features that distinguish them from pyknotic or ring-

stage forms in erythrocytes: in Giemsa-stained thin blood smears, these features

include a magenta nucleus bordered by a small region of blue cytoplasm [8, 32,

37]. Mitochondrial activity is vital to the survival of dormant forms and their return

to active propagation; this activity has enabled the identification and separation of

these forms by fluorescent markers specific for mitochondrial membrane

potential [11]. In the present study, we have used FACS sorting, ASM, and

autofluorescence FLIM-phasor analysis to characterize signature changes in the

mitochondria of persisters that develop after exposure of *P. falciparum*-infected

erythrocytes to DHA.

Remarkable morphological differences distinguish the mitochondria of DHA-

induced persisters from the mitochondria of actively growing ring-stage parasites.

The mitochondria of persisters have a more rumpled and corrugated appearance

that appears within 50 hours of DHA exposure, whereas smooth and oblate

mitochondria are characteristic of the actively replicating ring-stage parasites.

Along with their irregular shapes, the fluorescence volumes of persister

mitochondria are often large, up to 5-fold greater than the mitochondrial volumes

of ring stages. These differences are accompanied by a much reduced average

separation between the mitochondria and nuclei in persisters; indeed, many ASM

images suggest partial enwrapment of the nuclei by the mitochondria with

contact between these organelles. Studies of the ultrastructural effects of

artemisinins on various *Plasmodium* species *in vivo* have likewise found

mitochondrial enlargement in electron microscopy images [30, 38-40]; among

other apicomplexans, mitochondrial enlargement is reported in *Toxoplasma*

gondii following artemisinin drug pressure [31].

P. falciparum persisters maintain mitochondrial enlargement in tight

approximation with the nucleus throughout their period of dormancy. Open

questions remain regarding the molecular pathways underlying these changes

and how they might be induced upon ART exposure. Reactive oxygen species

(ROS) are induced by artemisinin in isolated *Plasmodium* and yeast

mitochondria, and agents that reduce electron transport chain (ETC) activity

reduce artemisinin-induced ROS production and resulting membrane damage

[41]. Depolarization of mitochondrial membrane potential accompanies the

disruption of mitochondrial function by artemisinin in yeast [42]. In human tumor

cell lines, activation of the artemisinin endoperoxide bridge by mitochondrial

heme is thought to promote generation of ROS by the electron transport chain,

leading to mitochondrial dysfunction and cellular apoptosis [43]. Artemisinin-

resistant lines of *Toxoplasma gondii* selected *in vitro* provided further evidence

for the involvement mitochondrial pathways, as demonstrated by amplifications of

mitochondrial cytochrome b and cytochrome c oxidase I genes and mutations of

a DegP2 ortholog in the mitochondrial proteome [31]. Another study identified the

same DegP2 ortholog in a genetic screen, genetic disruption facilitated survival

of *T. gondii* exposed to lethal concentrations of DHA*,* and deletion of this region

in *P. falciparum* resulted in higher survival in the RSA [44].

Interestingly, in breast cancer cells and mouse embryonic fibroblasts the

responses to mitochondrial stress include formation of tethers between the

nucleus and Nuclear Associated Mitochondria (NAM) [45]. ROS production after

exposure of THP-1 derived macrophages to the fungal toxin altertoxin II was

followed by relocation of mitochondria to perinuclear regions [46]. In other

systems of cancer cells, ROS may both promote and result from dysfunctional

mitochondrial communications with cell nuclei and organelles including

endoplasmic reticulum and peroxisomes, facilitating tumorigenesis [47]. Similarly,

we speculate, ROS effects and oxidative stress may be involved in triggering the

development and survival of *P. falciparum* persisters through the morphological

changes of mitochondria and their repositioning to tight associations with nuclei.

Mitochondrial retrograde response (MRR) signaling provides communications

from mitochondria to nuclei that promote cellular adaptations under conditions of

stress [48]. In studies of MRR in breast cancer, contact microdomains between

the two organelles involve cholesterol redistribution and may promote a survival

response [45]. The MRR pathway is conserved among species, as exemplified

by nuclear protein CLK-1 in *Caenorhabditis elegans* and its homolog in human

cells COQ7, which serves as a regulator of the MRR pathway that limits ROS

production [49]. In *Saccharomyces cerevisiae*, nuclear gene *Nonmitochondrial*

Citrate Synthase 2 (CIT2) transcription was increased after inhibition of

mitochondrial respiration [50], and mitochondrial pathways were activated in

response to ER stress [51]. In these studies, mitochondria and the MRR were

found to serve as barometers of cellular damage, capable of changing gene

expression pathways to facilitate survival. While different mechanisms of

endoperoxide activation and target impairment by carbon-centered or oxygen-

centered radicals have been proposed in the damage of malaria parasites by

ART, rapidly killing is thought to result from depolarization of the parasite

mitochondrial and plasma membranes upon exposure to ROS, initiated by iron

bioactivation of ART or by iron-dependent oxidative stress [52]. In the present

work, changes in mitochondrial morphology, reduced mito-nuclear distances, and

metabolic consequences induced by DHA exposure may be tied to gene

expression responses involved in persister formation as a survival mechanism.

These responses may relate to an innate growth bistability phenomenon whereby

a fraction of the drug-exposed population switches into the metabolic quiescence

of persister forms, as we have discussed elsewhere [53].

Our findings from autofluorescence FLIM-phasor analysis indicate an increased

ratio amount of free NADH in persisters relative to actively replicating ring-stage

parasites. This measure relates to the redox state of the cells. In other systems,

human breast cell lines exposed to potassium cyanide (KCN) treatment or

exposed to hypoxic conditions, autofluorescence FLIM revealed a higher amount

of free NADH due to inhibition of cellular respiration [54]. Increased

concentrations of free NADH in the mitochondria have been shown to promote

increased ROS production [55]. When assaying ART activity in tumor cell lines,

ETC function in the mitochondria was found to be essential for apoptosis through

ROS production [43]. Taken together, these findings suggest that activation of

the endoperoxide bridge of artemisinin by mitochondrial heme leads to altered

ETC function, increased free NADH, and increased ROS species with

concomitant mitochondrial damage.

After DHA treatment, a small fraction (less than ~1%; [5, 7]) of the parasite

population survives killing by entering the metabolically quiescent state of

dormancy. In this state, persisters may avoid damage by mitochondrial signaling

and modulation of essential pathways of gene expression. How is it that this

small fraction of the parasite population becomes persisters while the larger

fraction dies away? The answer may be a stochastic one that lies in systems of

gene control. In *P. falciparum*, feedback loops could rapidly flip the growth state

of some cells to dormancy, producing persister forms that can survive DHA

treatment while the actively replicating parasites succumb to the toxic effects of

the drug. Understanding these feedback loops and the metabolic and functional

features of persisters may lead to new chemotherapeutic approaches that can

prevent clinical recrudescences.

Materials and Methods

Parasite Cultivation

Human erythrocytes were obtained weekly from Virginia Blood Services

(Richmond, VA). Blood was washed upon arrival with filtered RPMI 1640 media

(containing 25 mM HEPES and 50 µg/mL hypoxanthine) (KD Medical, Columbia,

MD), and stored at 50% hematocrit in a 4˚C refrigerator for use within a week

from processing. *P. falciparum* GB4 [56] and 803 [57] parasites were grown in

complete culture media (cRPMI) containing 1% Albumax II (Life Technologies,

CA, USA), 0.21% sodium bicarbonate (KD Medical, Columbia, MD), and 20

µg/mL gentamicin (KD Medical, Columbia, MD) in RPMI-1640 (KD Medical,

Columbia, MD), at 5% hematocrit, 37˚C, and a gas mixture containing 90% N_2,

5% CO_2, and 5% O_2. To monitor parasitemia, thin blood films were prepared on

glass slides, fixed with methanol and stained for 15 minutes with 20% Giemsa

staining (Sigma-Aldrich, St. Louis, MO). Using bright-field microscopy and an

100× oil objective, an estimated 1,000 erythrocytes were counted, and the

number of parasitized cells was used to estimate percentage parasitemia. In

recrudescence experiments, media was changed every other day and fresh

erythrocytes were added every four days. When media was changed, blood films

were made and used to monitor parasitemia.

Ring-stage Survival Assay (RSA)

803 and GB4 cultures were tightly synchronized to obtained 0 – 3 hour ring

stages through two successive 5% D-sorbitol treatments 46 hours apart [24].

Each sorbitol treatment lasted 10 minutes at room temperature. Immediately

following the second sorbitol treatment, cultures were adjusted to 2% parasitemia

and 5% hematocrit in a total volume of 10 mL in a T25 flask (Thermo Fisher

Scientific, Waltham, MA). Thin blood smears were prepared just before drug

exposure. Cultures were then treated at a final concentration of 0.1% DMSO or

700 nM DHA/0.1% DMSO by addition of 10 µl of DMSO or 10 µl of 700 µM DHA

for 6 hours. After 6 hours of incubation, cells were washed and returned to

culture in drug-free cRPMI. Sixty-six to ninety hours post DHA treatment, thin

blood smears were prepared, fixed with 100% methanol, and stained with 20%

Giemsa (Sigma-Aldrich, St. Louis, MO). Slides identifications were blinded from

the investigators while parasitemia counts were obtained.

Dihydroartemisinin Treatment and Magnetic Removal of Mature Stages

803 and GB4 parasites were synchronized and treated with the control vehicle or

DHA as above. Thirty hours after the initiation of treatment, the DHA treated

culture was passed over a magnetically-activated cell sorting LD column (Miltenyi

Biotec, Auburn, CA) to deplete the culture of parasites that escaped the DHA

treatment and progressed past ring-stage. The same methodology as Teuscher

et al. [7] was applied to the use of the LD column. Pelleted cells from culture

were suspended in 2 mL of cold cRPMI and passed three times through the

column over a period of 3 – 5 hours. Afterwards, the column was washed with 30

mL of cRPMI. Eluates were combined and pelleted at 2,500 rpm for 5 minutes,

resuspended in 10 mL of cRPMI, and returned to culture. As the parasitized

erythrocytes approached maturity, the cultures were placed on a rotator in the

37˚C incubator to minimize the occurrence of multiply-infected relative to singly-

infected cells.

Dihydroartemisinin Recrudescence Assay

Tightly synchronized 803 and GB4 parasites, at approximately 2% ring-stage

parasitemia, were treated with 700 nM of DHA for 6 hours. At 24, 48 and 72

hours post DHA treatment, the cells of both cultures were treated with 10 mL of

5% D-Sorbitol for 30 minutes at 37°C and washed twice with cRPMI. This

treatment removed mature stage parasites that survived the DHA treatment.

Parasitemia was monitored by microscopy while media was changed on

alternate days and fresh erythrocytes were added every four days until

recrudescence.

Fluorescence Associated Cell Sorting (FACS)

At t = 50 hours, cells from 1 mL of each culture were pelleted and washed three

times with 1ml of 1× HBSS (Gibco, Waltham, MA) containing 2% fetal bovine

serum (FBS; Gibco, Waltham, MA). Washed samples were stained with 0.4× SG

(Invitrogen, Waltham, MA) for DNA content and 0.1 µM MT (Invitrogen, Waltham,

MA) for mitochondrial potential [58]. Samples for FLIM analysis were stained only

with 0.1 µM MT to avoid SG spectral overlap (bleedthrough) with the endogenous

FLIM signal [35]. Control 0.1% DMSO vehicle-treated and DHA-treated samples

were filtered through a 35 µm filter (Corning, Corning, NY) immediately before

sorting.

Cells were sorted by using the BD FACS Aria Fusion, and gating was performed

using the BD FACSDiva™ software (BD Biosciences, Franklin Lakes, NJ). For

ASM, samples underwent two sorting steps: 1) isolation of infected from

uninfected erythrocytes and 2) separation of parasite populations by DNA and

mitochondrial signals. The first step enriched infected cells by collecting 1 million

SG positive cells. Enrichment removes the vast majority of uninfected erythrocyte

allowing for more accurate sorting of live and dead parasites in the second step.

A highly pure sorting modality (4-way purity sorting for FACS Aria Fusion) was

chosen for the second round of sorting. Enriched samples were collected in

polystyrene round-bottom 5 mL tubes (Corning, Corning, NY) with 1 mL of 1×

HBSS containing 2% FBS, pelleted at 2500 rpm for 5 minutes in the 5 mL

polystyrene tube, resuspended, and sorted into two populations for each sample

(Figure 2). Parasites in the left gate were used for a Giemsa-stained thin blood

smear while parasites in the right gate were used for ASM. All four samples were

collected in polystyrene round-bottom 5 mL tubes (Corning, Corning, NY) with 1

mL of 1× HBSS/2% FBS.

For autofluorescence FLIM, samples underwent a single step sorting. The MT

positive cells were collected in polystyrene round-bottom 5 mL tubes (Corning,

Corning, NY,) with 1 mL of 1× HBSS/2% FBS.

Data analyses were conducted using FCS Express (De Novo Software,

Pasadena, CA).

AiryScan Microscopy Analysis

A 10 μm x 10 μm polydimethylsiloxane (PDMS) stencil microwell (Alvéole, Paris,

France) was placed inside each well of a Lab-Tek 8 well chamber (Thermo

Fisher Scientific, Waltham, MA). Ten μl of 0.01% poly-lysine (Sigma-Aldrich, St.

Louis, MO) was added to the microwell and incubated for 30 minutes at room

temperature to allow poly-lysine to adhere to the slide. Afterwards, the microwell

was washed twice with 10 µL of 1× HBSS to remove unbound poly-lysine. The

DMSO and DHA-treated parasites positive for SG/MT were transferred to 1.5 mL

tubes, spun down at 2,500 rpm for 3 minutes and resuspended in 20 µL of

remaining wash buffer. Ten µL of cell suspension was added directly to the

center of the poly-lysine coated microwell and incubated for 30 minutes at room

temperature in the dark. Lastly, 200 µL of a 1:4 dilution of Intracellular (IC)

Fixation Buffer (Invitrogen, Waltham, MA) in cell culture water (Gibco, Waltham,

MA) was added to the chamber. The chamber was covered with a plastic

membrane and stored at 4˚C for imaging within the following 24 hours. Parasites

were imaged using the Zeiss LSM 880 with AiryScan. Afterwards, the AiryScan

processed images were analyzed using Imaris 9.1.2 and volumes were adjusted.

In the majority of cells, the small and low level mitochondrial DNA SG

fluorescence was clearly distinct and excluded in the analysis. In a few cells,

mitochondrial DNA fluorescence was not visible or was merged during

thresholding of the DNA volume. Mitochondrial DNA regions were excluded if

possible. The volumes of the mitochondrial and DNA volume were calculated as

well as the distance between these volumes.

Autofluorescence FLIM and Phasor Analysis

When NADH undergoes multi-photon excitation, photons are released over time

as NADH returns to its ground state, establishing a pattern of fluorescence decay

that can be detected over a period of nanoseconds. The lifetime of decay differs

for NADH in its free (0.4 nanoseconds) or protein bound state (1 nanosecond),

which allows for the quantification of the redox state of the organism or tissue

under study [18]. In an early example of use of this observation, Chance *et al.*

[34] demonstrated increases of the characteristic emission spectrum from NADH

in rat organs under anoxic conditions; further, this spectrum matched the NADH

emission spectrum from isolated mitochondria. With recent technical and

computation advances, fluorescence emission data can translated into phasor

space, a fit-free method that allows for the direct transform of each pixel of an

image with a fluorescence decay to its corresponding phasor position [19]. Each

fluorescence decay from each pixel undergoes a Fourier transform and is

represented as a vector object with two Cartesian coordinates, one coordinate as

the real part of the Fourier transform (G) and while the other as the imaginary

part (S) [59]. The phasor plane is bounded by a "universal semicircle" that ranges

from 0 to 1.0 on the G axis (cosine transform) and 0 to 0.5 on the S axis (sine

transform). Single exponential decays fall along the boundary of the universal

semicircle while multi-exponential decays fall within the semicircle, where they

are subject to the law of phasor addition [60]. Autofluorescence FLIM of

endogenous fluorophores, such as free and protein bound NADH, typically yields

curves of multi-exponential decay and phasor coordinates within the universal

semicircle. Greater relative amounts of free NADH shift the vector positions to

the right side of the graph (shorter lifetimes) whereas greater amounts of protein-

bound NADH shift the vector positions to the left (longer lifetimes) [61]. Through

the phasor-FLIM approach, the effects of different treatments on metabolism can

be conveniently visualized by shifts of the vector position on the phasor plot [62].

DHA-treated or control parasites stained with MT were seeded onto a microwell

in an 8-well chamber coated with 0.01% poly-lysine. A confocal image was

acquired using a Zeiss LSM 780 for each field of view with two channels:

Differential Interference Contrast (DIC) and MT emission (633nm excitation and

emission collected 640 – 740 nm range). Next, emission data were collected

from the given field of view with the integrated Coherent Chameleon II

Ti:Sapphire femtosecond-pulsed laser tuned to 750 nm. Photons for lifetime data

fitting were collected using a Becker & Hickl TCSPC DCC-100 control card with

two HCM-130 GaAsP hybrid detectors. A 525 nm dichroic separated the

emission into the detectors and a 480/40 bandpass filter was used with the short

wavelength detector. Each field of view was scanned until at least ten million

photon events were collected. Phasor values were calculated for each pixel with

the SPCImage software. The resulting files were exported, and a custom

MATLAB algorithm used the signal from the MT confocal images to threshold the

mitochondria, obtain the mean phasor values, and run statistical tests.

Outgrowth Using Sorted Parasitized Cells

Outgrowth experiments were used to investigate parasite viability after FACS.

For this purpose, control and DHA-treated samples at $t = 50$ hours were obtained

through three gates: SG/MT Dim (SG/MT^D), SG/MT Intermediate (SG/MT^I) and

SG/MT Bright (SG/MT^B) (Supplementary File 1). Each population was placed into

a polystyrene tube with 1× HBSS/2% FBS at a final concentration of 1 mL. One

hundred µL was taken from these samples for a blood film, and the cells in the

samples were pelleted at 2,500 rpm for 5 minutes, resuspended in 180 µL of

cRPMI, and transferred to a 96 well plate. Twenty µL of a 50% erythrocyte stock

were added to each well to obtain a 5% hematocrit culture. Media were changed

on alternate days and 150 µL of 3.3% hematocrit solution were added every 4 or

5 days to replenish erythrocytes lost to sampling. The parasitemia in each well

was monitored through thick and thin smears, using 1 µL of parasitized

erythrocytes for each smear, and the time it took for the parasitemia to reach

0.5% was recorded.

Parasite viability was also verified for both control or DHA-treated GB4 and 803

parasites, utilizing the SG and MT stained samples (Supplementary Files 1 and

2).

Statistical Analysis

AiryScan datasets were compared by unpaired t-tests calculated with Graphpad

Prism version 8.4.3. Autofluorescence FLIM-phasor datasets were compared

through Pillai trace calculations in MATLAB version 2019b. Summarized statistics

are shown in Supplementary File 3.

Acknowledgements

We thank Tom Moyer, Teresa Hawley, and David Stephany in the Flow

Cytometry Section of the Research Technologies Branch for their advice on

FACS sorting, and Susan Pierce for supporting experiments at the Laboratory of

Immunogenetics, NIAID. This work was funded by the NIAID Division of

Intramural Research.

Competing interests

No competing interests declared.

References

1. World Health Organization (2015) Guidelines for the treatment of malaria,
third edition. Geneva: World Health Organization. Report No.:
9789241549127. Available at:
https://apps.who.int/iris/handle/10665/162441. Accessed 21 May 2020.
2. Dondorp A, Nosten F, Stepniewska K, Day N, & White N (2005)
Artesunate versus quinine for treatment of severe falciparum malaria: a
randomised trial. *Lancet* 366(9487):717-725.
3. Dondorp AM*, et al.* (2010) Artesunate versus quinine in the treatment of
severe falciparum malaria in African children (AQUAMAT): an open-label,
randomised trial. *Lancet* 376(9753):1647-1657.
4. Li GQ, Arnold K, Guo XB, Jian HX, & Fu LC (1984) Randomised
comparative study of mefloquine, qinghaosu, and pyrimethamine-
sulfadoxine in patients with falciparum malaria. *Lancet* 2(8416):1360-
1361.
5. Hoshen MB, Na-Bangchang K, Stein WD, & Ginsburg H (2000)
Mathematical modelling of the chemotherapy of Plasmodium falciparum
malaria with artesunate: postulation of 'dormancy', a partial cytostatic
effect of the drug, and its implication for treatment regimens. *Parasitology*
121 (Pt 3):237-246.
6. LaCrue AN, Scheel M, Kennedy K, Kumar N, & Kyle DE (2011) Effects of
artesunate on parasite recrudescence and dormancy in the rodent malaria
model Plasmodium vinckei. *PLoS One* 6(10):e26689.

7.   Teuscher F*, et al.* (2010) Artemisinin-induced dormancy in Plasmodium
739   falciparum: duration, recovery rates, and implications in treatment failure.
740   *J Infect Dis* 202(9):1362-1368.
741   8.   Tucker MS, Mutka T, Sparks K, Patel J, & Kyle DE (2012) Phenotypic and
742   genotypic analysis of in vitro-selected artemisinin-resistant progeny of
743   Plasmodium falciparum. *Antimicrob Agents Chemother* 56(1):302-314.
744   9.   Nakazawa S, Kanbara H, & Aikawa M (1995) Plasmodium falciparum:
745   recrudescence of parasites in culture. *Exp Parasitol* 81(4):556-563.
746   10.   Nakazawa S, Maoka T, Uemura H, Ito Y, & Kanbara H (2002) Malaria
747   parasites giving rise to recrudescence in vitro. *Antimicrob Agents*
748   *Chemother* 46(4):958-965.
749   11.   Peatey CL*, et al.* (2015) Mitochondrial membrane potential in a small
750   subset of artemisinin-induced dormant Plasmodium falciparum parasites
751   in vitro. *J Infect Dis* 212(3):426-434.
752   12.   Chen N*, et al.* (2014) Fatty acid synthesis and pyruvate metabolism
753   pathways remain active in dihydroartemisinin-induced dormant ring stages
754   of Plasmodium falciparum. *Antimicrob Agents Chemother* 58(8):4773-
755   4781.
756   13.   Duvalsaint M & Kyle DE (2018) Phytohormones, isoprenoids, and role of
757   the apicoplast in recovery from dihydroartemisinin-induced dormancy of
758   Plasmodium falciparum. *Antimicrob Agents Chemother* 62(3).
759   14.   Gray KA*, et al.* (2016) Correlation between cyclin dependent kinases and
760   artemisinin-induced dormancy in Plasmodium falciparum in vitro. *PLoS*
761   *One* 11(6):e0157906.
762   15.   Korobchevskaya K, Lagerholm BC, Colin-York H, & Fritzsche M (2017)
763   Exploring the potential of Airyscan microscopy for live cell imaging.
764   *Photonics* 4(3):41.
765   16.   Becker W*, et al.* (2004) Fluorescence lifetime imaging by time-correlated
766   single-photon counting. *Microsc Res Tech* 63(1):58-66.
767   17.   Berezin MY & Achilefu S (2010) Fluorescence lifetime measurements and
768   biological imaging. *Chem Rev* 110(5):2641-2684.
769   18.   Lakowicz JR, Szmacinski H, Nowaczyk K, & Johnson ML (1992)
770   Fluorescence lifetime imaging of free and protein-bound NADH. *Proc Natl*
771   *Acad Sci U S A* 89(4):1271-1275.
772   19.   Digman MA, Caiolfa VR, Zamai M, & Gratton E (2008) The phasor
773   approach to fluorescence lifetime imaging analysis. *Biophys J* 94(2):L14-
774   16.
775   20.   Stringari C*, et al.* (2011) Phasor approach to fluorescence lifetime
776   microscopy distinguishes different metabolic states of germ cells in a live
777   tissue. *Proc Natl Acad Sci U S A* 108(33):13582-13587.
778   21.   Bhattacharjee A, Datta R, Gratton E, & Hochbaum AI (2017) Metabolic
779   fingerprinting of bacteria by fluorescence lifetime imaging microscopy. *Sci*
780   *Rep* 7(1):3743.
781   22.   Pouli D*, et al.* (2016) Imaging mitochondrial dynamics in human skin
782   reveals depth-dependent hypoxia and malignant potential for diagnosis.
783   *Sci Transl Med* 8(367):367ra169.

23. Sá JM*, et al.* (2018) Artemisinin resistance phenotypes and K13
inheritance in a Plasmodium falciparum cross and Aotus model. *Proc Natl*
Acad Sci U S A 115(49):12513-12518.
24. Kite WA, Melendez-Muniz VA, Moraes Barros RR, Wellems TE, & Sa JM
(2016) Alternative methods for the Plasmodium falciparum artemisinin
ring-stage survival assay with increased simplicity and parasite stage-
specificity. *Malar J* 15:94.
25. Klonis N*, et al.* (2011) Artemisinin activity against Plasmodium falciparum
requires hemoglobin uptake and digestion. *Proc Natl Acad Sci U S A*
108(28):11405-11410.
26. Mok S*, et al.* (2015) Drug resistance. Population transcriptomics of human
malaria parasites reveals the mechanism of artemisinin resistance.
Science 347(6220):431-435.
27. Malleret B*, et al.* (2011) A rapid and robust tri-color flow cytometry assay
for monitoring malaria parasite development. *Sci Rep* 1:118.
28. Scarpelli PH*, et al.* (2019) Melatonin activates FIS1, DYN1, and DYN2
Plasmodium falciparum related-genes for mitochondria fission:
Mitoemerald-GFP as a tool to visualize mitochondria structure. *J Pineal*
Res 66(2):e12484.
29. van Dooren GG*, et al.* (2005) Development of the endoplasmic reticulum,
mitochondrion and apicoplast during the asexual life cycle of Plasmodium
falciparum. *Mol Microbiol* 57(2):405-419.
30. Maeno Y*, et al.* (1993) Morphologic effects of artemisinin in Plasmodium
falciparum. *Am J Trop Med Hyg* 49(4):485-491.
31. Rosenberg A, Luth MR, Winzeler EA, Behnke M, & Sibley LD (2019)
Evolution of resistance in vitro reveals mechanisms of artemisinin activity
in Toxoplasma gondii. *Proc Natl Acad Sci U S A*.
32. Breglio KF, Rahman RS, Sa JM, Roberts DJ, & Wellems TE (2018) Kelch
mutations in Plasmodium falciparum protein K13 do not modulate
dormancy after artemisinin exposure and sorbitol selection in vitro.
Antimicrob Agents Chemother 62(5).
33. Chance B, Schoener B, Oshino R, Itshak F, & Nakase Y (1979) Oxidation-
reduction ratio studies of mitochondria in freeze-trapped samples. NADH
and flavoprotein fluorescence signals. *J Biol Chem* 254(11):4764-4771.
34. Chance B, Legallais V, & Schoener B (1962) Metabolically linked changes
in fluorescence emission spectra of cortex of rat brain, kidney and adrenal
gland. *Nature* 195:1073-1075.
35. Zipper H, Brunner H, Bernhagen J, & Vitzthum F (2004) Investigations on
DNA intercalation and surface binding by SYBR Green I, its structure
determination and methodological implications. *Nucleic Acids Res*
32(12):e103.
36. Ittarat W*, et al.* (2003) Recrudescence in artesunate-treated patients with
falciparum malaria is dependent on parasite burden not on parasite
factors. *Am J Trop Med Hyg* 68(2):147-152.
37. Chavchich M, Van Breda K, Rowcliffe K, Diagana TT, & Edstein MD
(2016) The spiroindolone KAE609 does not induce dormant ring stages in

830   Plasmodium falciparum parasites. *Antimicrob Agents Chemother*
831   60(9):5167-5174.

38.   Ellis DS*, et al.* (1985) The chemotherapy of rodent malaria, XXXIX.
Ultrastructural changes following treatment with artemisinine of
Plasmodium berghei infection in mice, with observations of the localization
of [3H]-dihydroartemisinine in P. falciparum in vitro. *Ann Trop Med*
Parasitol 79(4):367-374.

39.   Kawai S, Kano S, & Suzuki M (1993) Morphologic effects of artemether on
Plasmodium falciparum in Aotus trivirgatus. *Am J Trop Med Hyg*
49(6):812-818.

40.   Jiang JB, Jacobs G, Liang DS, & Aikawa M (1985) Qinghaosu-induced
changes in the morphology of Plasmodium inui. *Am J Trop Med Hyg*
34(3):424-428.

41.   Wang J*, et al.* (2010) Artemisinin directly targets malarial mitochondria
through its specific mitochondrial activation. *PLoS One* 5(3):e9582.

42.   Li W*, et al.* (2005) Yeast model uncovers dual roles of mitochondria in
action of artemisinin. *PLoS Genet* 1(3):e36.

43.   Mercer AE, Copple IM, Maggs JL, O'Neill PM, & Park BK (2011) The role
of heme and the mitochondrion in the chemical and molecular
mechanisms of mammalian cell death induced by the artemisinin
antimalarials. *J Biol Chem* 286(2):987-996.

44.   Harding CR*, et al.* (2020) Genetic screens reveal a central role for heme
metabolism in artemisinin susceptibility. *Nat Commun* 11(1):4813.

45.   Desai R*, et al.* (2020) Mitochondria form contact sites with the nucleus to
couple pro-survival retrograde response. *bioRxiv*:445411.

46.   Del Favero G, Hohenbichler J, Mayer RM, Rychlik M, & Marko D (2020)
Mycotoxin altertoxin II induces lipid peroxidation connecting mitochondrial
stress response to NF-kappaB inhibition in THP-1 macrophages. *Chem*
Res Toxicol 33(2):492-504.

47.   Xia M*, et al.* (2019) Communication between mitochondria and other
organelles: a brand-new perspective on mitochondria in cancer. *Cell*
Biosci 9:27.

48.   Butow RA & Avadhani NG (2004) Mitochondrial signaling: the retrograde
response. *Mol Cell* 14(1):1-15.

49.   Monaghan RM*, et al.* (2015) A nuclear role for the respiratory enzyme
CLK-1 in regulating mitochondrial stress responses and longevity. *Nat Cell*
Biol 17(6):782-792.

50.   Liao XS, Small WC, Srere PA, & Butow RA (1991) Intramitochondrial
functions regulate nonmitochondrial citrate synthase (CIT2) expression in
Saccharomyces cerevisiae. *Mol Cell Biol* 11(1):38-46.

51.   Hijazi I, Knupp J, & Chang A (2020) Retrograde signaling mediates an
adaptive survival response to endoplasmic reticulum stress in
Saccharomyces cerevisiae. *J Cell Sci* 133(6).

52.   Antoine T*, et al.* (2014) Rapid kill of malaria parasites by artemisinin and
semi-synthetic endoperoxides involves ROS-dependent depolarization of
the membrane potential. *J Antimicrob Chemother* 69(4):1005-1016.

53. Wellems TE, Sa JM, Su XZ, Connelly SV, & Ellis AC (2020) 'Artemisinin Resistance': Something new or old? Something of a misnomer? *Trends Parasitol*.

54. Bird DK*, et al.* (2005) Metabolic mapping of MCF10A human breast cells via multiphoton fluorescence lifetime imaging of the coenzyme NADH. *Cancer Res* 65(19):8766-8773.

55. Murphy MP (2009) How mitochondria produce reactive oxygen species. *Biochem J* 417(1):1-13.

56. Hayton K*, et al.* (2008) Erythrocyte binding protein PfRH5 polymorphisms determine species-specific pathways of Plasmodium falciparum invasion. *Cell Host Microbe* 4(1):40-51.

57. Amaratunga C*, et al.* (2012) Artemisinin-resistant Plasmodium falciparum in Pursat province, western Cambodia: a parasite clearance rate study. *Lancet Infect Dis* 12(11):851-858.

58. Amaratunga C, Neal AT, & Fairhurst RM (2014) Flow cytometry-based analysis of artemisinin-resistant Plasmodium falciparum in the ring-stage survival assay. *Antimicrob Agents Chemother* 58(8):4938-4940.

59. Dalbey RE, Weiel J, Perkins WJ, & Yount RG (1984) Resolution of multiple fluorescence lifetimes in heterogeneous systems by phase-modulation fluorometry. *J Biochem Biophys Methods* 9(3):251-266.

60. Jameson DM, Gratton E, & Hall RD (1984) The measurement and analysis of heterogeneous emissions by multifrequency phase and modulation fluorometry. *Applied spectroscopy reviews* 20(1):55-106.

61. Ranjit S, Malacrida L, Jameson DM, & Gratton E (2018) Fit-free analysis of fluorescence lifetime imaging data using the phasor approach. *Nat Protoc* 13(9):1979-2004.

62. Aguilar-Arnal L*, et al.* (2016) Spatial dynamics of SIRT1 and the subnuclear distribution of NADH species. *Proc Natl Acad Sci U S A* 113(45):12715-12720.

914
915
916
Figure 1. Intraerythrocytic cycle delay exhibited by DHA-treated *P. falciparum*
parasites t = 72 hours after DHA incubation. GB4, 803, 76H10[C580Rev] and 76H10
parasite lines underwent a standard Ring-stage Survival Assay (RSA) procedure,
in which cultures at approximately 2% parasitemia were treated with control 0.1%
DMSO (vehicle) or 700 nM DHA at t = 0 hours. After 6 hours of incubation, the
samples were washed and returned to culture. At t = 72 hours, the stage
distributions were counted and percentage survivals were calculated. (**A**) Ring-
stage Survivals. At the t = 72 hour timepoint, the percent survival was calculated
for each strain through dividing the number of surviving DHA-treated parasites by
the number of control vehicle-treated parasites. After DHA treatment, GB4 and
76H10[C580Rev] parasites (C580 K13 propeller wild type) survived at lower levels
than the C580Y propeller mutant 803 and 76H10 parasites (0.93 +/- 0.16 (mean
+/- SEM, N=5) and 0.69 +/- 0.14 (mean +/- SEM, N=6) *vs.* 10.84 +/- 1.32 (mean
+/- SEM, N=6) and 9.50 +/- 1.09 (mean +/- SEM, N=8), respectively). (**B**) Stage
distributions in control vehicle-treated or DHA-treated GB4, 803, 76H10[C580Rev]
and 76H10 parasite lines at t = 72 hours. The stage distributions were counted
from Giemsa stained thin blood smears. (**C**) Timeline of intraerythrocytic delay.
Bell curves represent the age distributions of control vehicle-treated or DHA-
treated K13 C580 (GB4, 76H10[C580Rev]) or C580Y (803, 76H10) parasite lines.

The dotted lines at $t = 50$ and at $t = 72$ hours indicate the collection timepoints for
FACS and RSA determinations, respectively.
938

940
941
Figure 2. Representative dot plots with gating strategy for AiryScan microscopy
analysis of control or DHA-treated GB4 and 803 parasites at t = 50 hours. The
experimental procedure is shown in Figure 2 – Figure Supplement 1. In each
experiment, the SG positive parasites were separated into two populations based
on MT fluorescence intensity. The DHA treatments killed most parasites, as
demonstrated by larger populations in the pyknotic gate (PYK). Parasites from
the MT+ gate were used for ASM. SG = SYBR Green I, MT = MitoTracker Deep
Red FM
950
951

953
Figure 3. Airyscan microscopy images of control or DHA-treated GB4 and 803
parasites at $t = 50$ hours. Individual panels show processed ASM images of the
mitochondrial (red) and nuclear (green) fluorescences from six representative
parasites in each treatment group separated by FACS. (**A**) GB4 parasites from
control 0.1% DMSO (vehicle) treated samples presented smooth, oblate
mitochondrial and nuclear volumes. Images in the control group are consistent
with young ring-stage parasites in the culture at $t = 50$ hours. DHA-treated
parasites show the distinctly different morphologies of persisters, with rumpled
and corrugated mitochondrial volumes that are in close approximation to the
nuclei. (**B**) Images of the control 803 parasites at t = 50 hours also have the
smooth, oblate appearance of mitochondria separated from the nucleus, as
expected of actively replicating young parasite ring forms. Images of DHA-
treated 803 parasites, in comparison to those of the treated GB4 population, are
consistent with a more mixed population of cells, many with the rumpled,
corrugated mitochondria closely apposed to nuclei and others with the smooth,
oblate mitochondria separate from the nuclei.
970
971

Figure 4. Quantifications of mito-nuclear distances and mitochondrial volumes of GB4 and 803 parasites at $t = 50$ hours, day 5 and day 11 post DHA treatment. Medians and interquartile ranges (IQR) are indicated over the distributions of mito-nuclear distances and mitochondrial volumes. (**A**) Scatterplots of the quantifications for each group of GB4 parasites exposed to 0.1% DMSO (control) or 700 nM DHA. (**B**) Scatterplots of the quantifications for each group of 803 parasites exposed to 0.1% DMSO (control) or 700 nM DHA. Datasets labeled control and DHA were obtained from samples $t = 50$ hours according to the schematic outline in Figure 1 – Figure Supplement 1. (**C**) Datasets on day 5 and day 11 were collected from the populations of GB4 and 803 parasites after DHA exposure and three daily sorbitol selections. GB4 parasites recrudesced to above 2% parasitemia by day 20 while 803 parasites recrudesced on day 16. In the mito-nuclear distance scatterplots, five outliers are beyond the bounds of the y-axis in the GB4 Control (2.8 µm, 2.5 µm, 2.2 µm, 2.7 µm, and 6.8 µm), one outlier in the 803 Control (3.7 µm), and three outliers in the 803 DHA results (2.6 µm, 2.0 µm, 5.5 µm). (**D**) Representative ASM images of DHA-treated 803 parasites on day 5 post treatment. Shown are six examples of persisters with rumpled, corrugated mitochondria in close approximation to the nuclei. (**E**) Surface images of DHA-treated GB4 and 803 parasites at day 11. Six representative parasites from each line are shown. Persister morphologies were evident in each parasite line, with the mitochondria in close approximation to the

995   nuclei. In the DHA-treated GB4 sample, large mitochondria enwrapping the
996   nuclei were observed, similar to the persister morphology shown in Figure 3. In
997   the DHA-treated 803 sample, persister morphology was apparent as well, but
998   parasites with smooth oblate mitochondria that did not enwrap nuclei were
999   frequently observed also, indicating an awakening from dormancy.
1000
1001
1002

Figure 5. Metabolic phenotyping of untreated control *vs.* DHA-treated GB4 and 803
parasites. (**A**) Schematic flow of FLIM-phasor data collection and analysis. MT+
parasites were sorted and seeded into a 10 μm x 10 μm polydimethylsiloxane (PDMS)
stencil microwell covered with 0.01% poly-lysine. The region of interest (ROI) in a
parasite is established by the boundary of MT fluorescence. This ROI mask is then
applied to the lifetime datafile represented by its FLIM intensity image. Next, Fourier
transformation of the FLIM data is performed on the data from each pixel and the
frequency domain results are averaged to represent a data point on the 2D phasor plot.
Each position has coordinates G (from 0 to 1) and S (from 0 to 0.5); single exponential
decays fall upon the semicircle whereas complex multiexponential decays, such as
occur with fluorescent lifetimes of metabolic coenzymes, fall within the semicircle. A shift
to the right indicates increased free NADH whereas a to the left indicates increased
enzyme bound NADH. (**B**) Autofluorescence FLIM-phasor graphs from the GB4 and 803
parasite populations at *t* = 50 hours. The phasor distribution of untreated control GB4
parasites has a mean coordinate of (0.7542, 0.2150) whereas the distribution of DHA-
treated parasites has a mean coordinate of (0.7971, 0.1879) (p < 0.0001, Pillai trace).
This shift after DHA treatment is toward increased free NADH, consistent with the
quiescent metabolic state of persisters. The distribution of untreated 803 parasites at *t* =
50 hours (right) has a mean coordinate of (0.6341, 0.1400) and the phasor distribution of
DHA-treated parasites has a mean coordinate of (0.6270, 0.1570). The distribution of
DHA-treated 803 parasites is slightly shifted upward relative to control (p < 0.05, Pillai
trace), but not to the left or right, consistent with the presence of actively replicating 803
parasites that survived ring-stage exposure to DHA. (**C**) Autofluorescence FLIM-phasor

graphs for DHA-treated GB4 and 803 on day 5 and day 11. In the phasor distributions
from day 5 (left), DHA-exposed GB4 and 803 parasites have mean coordinates of (0.72,
0.21) and (0.77, 0.21), respectively. Consistent with depletion of the many actively
replicating parasites from the 803 population by three daily sorbitol treatments, the 803
phasor distribution is shifted right with quiescence and increased levels of free NADH in
persisters. The varied genetic backgrounds and baseline metabolic states of African
GB4 and Southeast Asian 803 parasites help to explain different positions of the 803
and GB4 coordinates ($p < 0.0001$, Pillai trace).
1036
1037

1038

1039
1040
Figure 2 – Figure Supplement 1. Schematic of GB4 and 803 parasite
preparations for Airyscan and FLIM-phasor analysis. GB4 or 803 parasites were
synchronized through two sorbitol treatments, 46 hours apart, to obtain 0 – 3
hour rings. At the start of the experiment (t = 0 hours), the GB4 or 803
populations of 0 – 3 hour rings were treated at approximately 2% parasitemia
with either 700 nM DHA or control 0.1% DMSO (vehicle) for 6 hours, washed and
returned to culture. At t = 30 hours, the DHA-treated cultures were passed
through a depletion column to remove mature-stage parasites that continued to
actively grow after DHA treatment. Both the control and DHA-treated cultures
were placed on a shaking incubator overnight to facilitate invasion of the control
culture and to maintain the experimental conditions in each flask. At t = 50 hours,
1mL of each culture were stained with either SG plus MT for AiryScan
microscopy or MT only for FLIM-phasor autofluorescence analysis.
1054
1055
1056
1057

1058
Figure 3 – Figure Supplement 1. Example ASM images showing distance
between centroids in control or DHA-exposed GB4 and 803 parasites at $t = 50$
hours. The center of image mass was marked in Imaris using surfaces formed
from SYBR Green I and MitoTracker Deep Red FM fluorescence regions. (**A**)
GB4 parasites were exposed to control vehicle 0.1% DMSO (left panel) or 700
nM DHA (right panel). In GB4 parasites after DHA treatment, the distance
between the green and red fluorescence regions was substantially reduced in the
DHA-treated parasites. (**B**) In 803 parasites, similar distances were observed
between the control (left panel) or DHA-treated (right panel) parasites. Mean
distances were calculated between each voxel (volumetric pixel, a discrete
subset of the volume with a specific 3D coordinate) and averaged for each
parasite.

1081
1082
1083 **Figure 4 – Figure Supplement 1.** DNA volumes of control or DHA-treated GB4
1084 and 803 parasites at t = 50 hours, and of the DHA-treated parasites at days 5
1085 and 11 in the recrudescence studies.

Supplementary File 1. Isolation of three stages of P. falciparum-infected red
blood cells. Representative dot plots are shown for outgrowth experiments of
control or DHA-treated GB4 and 803 parasites at t = 50 hours. Control or DHA-
treated GB4 and 803 parasites were stained with SYBR Green (SG) and
MitoTracker Deep Red (MT) and sorted using the BD FACS Aria Fusion. Labels
identify dimly MT-fluorescent pyknotic cells (SG/MT^D), a ring-enriched population
of intermediate fluorescence (SG/MT^I), and brightly fluorescent late stages
(SG/MT^B).
1099
Supplementary File 2. Results of outgrowth experiments from the fluorescent
GB4 and 803 parasites isolated by FACS and collected in 1xHBSS/2% FBS.
Cells were pelleted at 2500 rpm for 5 minutes and returned to culture (200 µl
volume, 5% HCT) a 96 well plate. Thin and thick Giemsa stained blood smears
were monitored microscopically for parasite population growth. Rows in green
indicate growth; the rightmost column lists the days to 0.5% parasitemia in
culture.
1107
1108 **Supplementary File 3.** Statistic results from ASM and autofluorescence FLIM-
1109 phasor analysis. Significance testing of the ASM data was by unpaired t-test
1110 calculations. In the results from autofluorescence FLIM-phasor data, mean
1111 phasor distribution coordinates were assessed for significant differences by
1112 Pillai's trace calculations between the treatment groups.
1113
1114