Lyl-1 regulates primitive macrophages and microglia development

Shoutang Wang[1, †, *], Deshan Ren[1, ††, *], Anna-Lila Kaushik[1, †††], Gabriel Matherat[1, ††††], Yann

Lécluse[2], Dominik Filipp[3], William Vainchenker[1], Hana Raslova[1], Isabelle Plo[1], Isabelle Godin[1]

[1] Gustave Roussy, INSERM UMR1287, Villejuif; Université Paris-Saclay, France.

[2] PFIC, lUMS AMMICa (US 23 INSERM / UMS 3655 CNRS; Gustave Roussy, Villejuif, France

[3] Laboratory of Immunobiology, Institute of Molecular Genetics of the Czech Academy of Sciences,

Prague, Czech Republic.

Present addresses: [†] Department of Pathology and Immunology, Washington University School of

Medicine, St. Louis, MO, 63110, USA; [††] Medical school of Nanjing university, Model Animal

Research Center, Nanjing University, Nanjing 210093, China; [†††] Plasseraud IP, 33064 Bordeaux,

France; [††††] Agence Nationale pour la Recherche, Paris, France

*** Equal contribution;

Short Title: Lyl-1 in macrophage/microglia ontogeny

Corresponding author:

Isabelle Godin (ORCID: 0000-0001-8577-8388)

INSERM U1287; Institut Gustave Roussy-PR1; 114, rue Edouard Vaillant; 94805 VILLEJUIF Cedex;

France

Email address: Isabelle.Godin@gustaveroussy.fr ;

Phone: (33) 142 11 41 43; Fax: (33) 142 11 52 40

Key points:

1- Yolk sac primitive macrophage progenitors and microglia/Border Associated macrophages

express Lyl-1.

2- Lyl-1-deficiency impairs primitive macrophage and microglia development and leads to the up-

regulation of gene sets related to embryo patterning and neuro-development.

Abstract:

During ontogeny, macrophages (MΦ) populations emerge in the Yolk Sac (YS) via two distinct

progenitor waves, prior to hematopoietic stem cell development. MΦ-progenitors from the

primitive/"early EMP" and transient-definitive/"late EMP" waves both contribute to various

resident-MΦ populations in the developing embryonic organs. Identifying factors that modulates

early stages of MΦ-progenitor development may lead to a better understanding of defective

function of specific resident-MΦ subsets.

Here we show that primitive macrophage (MΦ^{Prim}) progenitors in the YS express Lyl-1, a bHLH

transcription factor related to SCL/Tal-1. Transcriptomic analysis of YS MΦ-progenitors indicated

that MΦ^{Prim} progenitors present at embryonic day (E) 9 are clearly distinct from those present at

later stages. Disruption of Lyl-1 basic helix-loop-helix domain led initially to an early increased

emergence of MΦ^{Prim} progenitors, and later to their defective differentiation. These defects were

associated with a disrupted expression of gene sets related to embryonic patterning and

neurodevelopment. Lyl-1-deficiency also induced a reduced production of mature MΦ/microglia

in the early brain, as well as a transient reduction of the microglia pool at midgestation and in the

newborn.

We thus identify Lyl-1 as a critical regulator of MΦ^{Prim} and microglia development, which

disruption may impair resident-MΦ function during organogenesis.

 INTRODUCTION

Amongst the components of the transcription factor network that regulate hematopoietic cells

features, *Tal-1*, *Lmo2*, *Runx1* and *Gata2* stand out as major regulators of hematopoietic progenitor

development.[1, 2] *Tal-1*, *Lmo2* and *Gata-2* belong to a transcriptional complex, which also includes

the basic helix-loop-helix (bHLH) transcription factor (TF) lymphoblastic leukemia-derived

sequence 1 (*Lyl-1*). Unlike its paralog *Tal-1*, which is mandatory for the specification of all

hematopoietic progenitors[3, 4], Lyl-1 roles during developmental hematopoiesis remains poorly

characterized. We analyzed these functions at the onset of YS hematopoietic development using

Lyl-1$^{LacZ/LacZ}$ mutant mice.[5]

During ontogeny, hematopoietic progenitors are generated in three successive and overlapping

waves.[6, 7] The emergence of the Hematopoietic Stem Cells (HSC) that will maintain lifelong

hematopoiesis in the adult occurs at mid-gestation in the third and definitive hematopoietic wave.

HSC generated in the aorta region immediately migrate to the fetal liver (FL) where they mature

and amplify before homing to the bone marrow before birth.[8, 9] Prior to HSC generation, the

production of blood cells relies on two hematopoietic waves provided by the YS. This HSC-

independent hematopoiesis comprises first the primitive hematopoietic wave, with the transient

production of progenitors with embryonic specific features: From Embryonic day (E) 7.00, the YS

produces monopotent progenitors for erythrocytes, megakaryocytes and macrophages (MΦ)[10, 11],

along with bipotent Erythro-Megakaryocytic (EMk) progenitors[12], in a *Myb*-independent

pathway.[13, 14] The second YS wave, called transient-definitive, provides for a limited duration

progenitors (mostly erythro-myeloid) that seed the FL and produce a hematopoietic progeny that

displays definitive/adult differentiation features. Erythro-myeloid cell production in this wave

occurs in a *Myb*-dependent pathway, through the progressive differentiation of erythro-myeloid

progenitors (EMP) in a pathway similar to the adult one.[6] As primitive and transient definitive YS

waves both produce cells from erythro-myeloid lineages, they are also termed respectively "early

EMP" and "late EMP".[15, 16]

Considering the MΦ lineage, fate-mapping approaches aimed at determining the embryonic

origin of resident-MΦs indicated that most tissues harbor resident-MΦs of diverse origins (YS, FL

and adult bone marrow)[15, 17, 18], which complicates the characterization of wave-dependent

functions of the various subsets. However, these fate-mapping analyses established that, contrary

to others tissue, brain MΦs (microglia and Border Associated MΦ (BAM)) develop only from YS-

derived MΦ-progenitors[14, 16, 19, 20, 21], confirming a model we previously put forward.[22] Due to the

coexistence of two waves in the YS, the origin of microglia has been debated (reviewed in [6, 15, 18,

[23]). An origin of microglia from MΦ-progenitors from the primitive/"early EMP" wave was

supported by microglia labelling following an early (E7.0-E7.5) CRE-mediated induction of Runx1[19]

and by the intact microglia pool in *Myb*-deficient mice.[14, 21] The origin of microglia was also

attributed to the primitive wave in zebrafish embryos, since in MΦ[Prim] progenitors arise in this

species from a location distinct from other hematopoietic progenitors.[24, 25, 26] Finally, the normal

microglia development in mice lacking Kit ligand, leading to an impaired EMP development and

the depletion of resident-MΦs in the skin, lung and liver supports this model.[27]

We here show that, at the early stages of YS hematopoiesis, *Lyl-1* expression characterizes

primitive MΦ-progenitors. Through RNA-seq. analyses, it appears that these primitive MΦ-

progenitors harbor an immune-modulatory phenotype, while those produce at a later stage favor

the inflammatory signaling which promotes the emergence of HSC in the third and definitive

hematopoietic wave. [28]

Our results also indicate that in the brain, Lyl-1 is expressed in the entire microglia/BAM cell

population at the onset of brain colonization and appeared to regulate microglia/BAM

development.

Altogether, these data point to Lyl-1 as a major regulator of early embryonic MΦ-progenitors

development and advocate for further analyses to more precisely delineate Lyl-1 function during

the development of resident-MΦs in homeostatic and pathological contexts.

RESULTS AND DISCUSSION

Lyl-1 expression marks MΦ^Prim progenitors from the early YS

Lyl-1 being expressed in the YS from the onset of YS hematopoiesis[29], we first explored its

function by characterizing the clonogenic potential of WT, *Lyl-1^WT/LacZ* and *Lyl-1^LacZ/LacZ* YS. E8-YS

were maintained in organ culture for 1 day (E8 OrgD1-YS), allowing only the development of

primitive and transient-definitive progenitors.[30, 31] Compared to WT, the production of MΦ

colonies was increased in *Lyl-1^WT/LacZ* and *Lyl-1^LacZ/LacZ* OrgD1-YS. Otherwise, the clonogenic

potential and colony distribution were unmodified ***(Figure 1A)***.

Using FACS-Gal assay, we noticed that the entire MΦ-progenitor population (Kit$^+$CD45$^+$CD11b$^+$)

expressed Lyl-1 at E9. In contrast, two MΦ-progenitor subsets, discriminated by FDG/Lyl-1

expression, were present after E9.5 ***(Figure 1B)***. Since after E9.5, the YS harbors both MΦ^Prim and

transient-definitive (MΦ^T-Def) MΦ–progenitors, and as these progenitors subsets cannot be

discriminated by phenotype,[11] we investigated the known features discriminating these two

waves: the origin from monopotent progenitors for MΦ^Prim progenitors[10, 11] and the *Myb*-

dependent[14, 32] differentiation of MΦ^T-Def progenitors from EMP, via the production of granulo-

monocytic (GM-), then granulocyte (G-) and MΦ-progenitors.[6]

At E8 (0-5 somites (S)), the YS only harbors MΦ^Prim progenitors, harboring a CD11b$^+$CD31$^+$

phenotype.[11] At this stage, all MΦ-progenitors expressed FDG/Lyl-1 ***(Figure 1C)***. Most FDG$^+$/Lyl-1$^+$

CD11b and CD31cells (69.27%±0.33%) from *Lyl-1^WT/LacZ* E8-YS reliably produced MΦ colonies

(72.78±9.65%; n=3) in clonogenic assays, amounting 1-4 MΦ progenitors per YS, a value consistent

with previously published data.[10, 11]

Lyl-1 expression by MΦ^Prim progenitors was strengthened by RT-qPCR comparison of *Myb*

expression: Both E9-YS MΦ^Prim progenitors and FDG$^+$/Lyl-1$^+$ progenitors from E10-YS expressed low

Myb levels strengthening their primitive status, while FDG⁻/Lyl-1⁻ progenitors from E10-YS

progenitors displayed *Myb* levels similar to lineage-negative Sca1⁺cKit⁺ progenitors from E12-FL

(Figure 1D).

The differentiation potential of FDG⁺/Lyl-1⁺ and FDG⁻/Lyl-1⁻ fractions of Ter119⁻Kit⁺CD45⁺CD11b⁺

myeloid progenitors isolated from E10-YS also pointed to Lyl-1 expression by MΦ^Prim progenitors

(Figure 1E): Similar to WT E9-YS MΦ^Prim progenitors, E10 FDG⁺/Lyl-1⁺ progenitors appeared

monopotent, as they nearly exclusively produced MΦ colonies. In contrast, E10 FDG⁻/Lyl-1⁻

myeloid progenitors produced GM, G and MΦ colonies, a feature typical of transient-definitive

progenitors [6]. Overall, these data together suggested that Lyl-1 may mark MΦ^Prim progenitors from

the earliest wave.

Distinct features of WT MΦ-progenitors at E9 and E10

The distinction between E9 and E10 MΦ-progenitors was confirmed in RNA-seq. analysis of

CD45⁺CD11b⁺Kit⁺ MΦ-progenitors sorted at E9 (MΦ^Prim progenitors) and E10 (MΦ^Prim and MΦ^T-Def

progenitors). Principal Component Analysis separated E9 and E10 MΦ-progenitors according to

stage and genotypes *(Figure 2A)*. E9 and E10 WT MΦ-progenitors differed by the expression of

726 genes, 176 were up-regulated at E9 and 550 at E10. Considering the coexistence of MΦ^Prim

and MΦ^T-Def progenitors in E10-YS, differentially expressed genes (DEGs) found at E10 may reflect

wave-specific differences or stage-dependent changes related to MΦ^Prim progenitor maturation.

Overlapping the identified DEGs to the EMP and E10.25-E10.5 MΦs signatures obtained by

Mass *et al.*[33] confirmed that WT E9 MΦ^Prim progenitors were distinct from these two populations,

since none of the 176 upregulated at E9 belonged to these signatures. Comparatively, about 5%

of the genes up-regulated at E10 belonged to the EMP and MΦ signatures *(Figure 2B)*. A similar

separation was observed in GSEA analyses *(Supplemental figure 1B)*. These observations suggest

that within E10 MΦ-progenitors some, likely the MΦ^{T-Def} ones, retain part of the EMPs signature.

In a WT context, E9 MΦ^{Prim} progenitors differed from E10 MΦ-progenitors by their TF

repertoire. Genes regulating erythroid development (*Gata1, Gata2, Klf1*) and globin genes,

embryonic (*Hbb-bh1, Hba-x, Hbb-y*) and definitive (*Hba-a2, Hba-a1, Hbb-bt*) were enriched at E10

(Figure 2B), while *Spi1*/PU.1 was highly expressed compared to *Gata1* at both stages *(Figure 2C)*.

The lower expression level at E9 of erythroid genes and of genes involved in granulo-monocytic

(*Mpo, Csf2r/GM-CSF receptors, Cebp, Jun*) and megakaryocytic development (*Pf4, TPO signaling*)

(Figure 2B; Supplemental table 1) sustains the monopotent/primitive status of E9 MΦ

progenitors, and suggests that MΦ^{T-Def} progenitors may retain the expression of genes that

characterize their EMP ancestor.

IPA and GSEA analyses indicated that E9 MΦ^{Prim} progenitors were more active in Eicosanoid

signaling than E10 progenitors *(Figure 2D)*. They were also enriched in type I interferon (IFN) β and

type II IFNγ signaling *(Figure 2E)* and in MHC-II related genes, especially *Cd74* (top 1 IPA network)

(Figure 2F; supplemental figure 1C). Cytometry analyses confirmed a low, but significant,

enrichment of MHC-II expression at E9, compared to E10 *(Supplemental figure 1D)*.

Comparatively, E10 MΦ-progenitors were more active in inflammatory signaling *(Figure 2D, E;*

supplemental figure 1C, E-G; supplemental table 1), and metabolically active *(Supplemental table*

1). The complement cascade and phagocytosis also prevailed at E10 *(Supplemental figure 1H-I)*.

Altogether, the signature for WT E9 MΦ^{Prim} progenitors points to an immuno-modulatory

function, while E10 MΦ-progenitors appear involved in phagocytosis and inflammatory signaling.

Interestingly, inflammatory signaling has been revealed as a key factor favoring embryonic HSC

emergence (reviewed in[34]). The source of inflammatory signals was further identified as MΦ

progenitors expressing *Mrc1*/CD206[35], a marker up-regulated in E10 WT MΦ-progenitors

compared to E9 *(Figure 2B)*.

Lyl-1 deficiency impacts embryonic development

When evaluating the effect of *Lyl-1*-deficiency at the earliest stage of MΦ^{Prim} development,

clonogenic assays pointed to an increased production of MΦ-progenitors in Mutant E8-YS

compared to WT *(Figure 3A)*, concordant with our first observation *(Figure 1A)*. At this stage, the

increased size of the initial MΦ-progenitor pool appeared to results from an elevated commitment

of mesodermal/pre-hematopoietic cells to a MΦ fate, rather than from a defective differentiation

(Supplemental figure 2 A-D and related information). The high increase of *Itga2b*/CD41

expression level in E9 *Lyl-1$^{LacZ/LacZ}$* MΦ^{Prim} progenitors *(Figure 3B)* may reflect this elevated

commitment. Lyl-1 expression in YS mesoderm[29], where it cannot substitute for *Tal-1* mandatory

function for the generation of hematopoietic progenitors[4], was already established[3]. Recently, Lyl-

1 was identified as a regulator of mesoderm cell fate[36] and of the maintenance of primitive

erythroid progenitors.[37]

The TF network that controls developmental hematopoiesis[2] was also modified *(Figure 3C)*:

beside the expected reduction of *Lyl-1* expression, the expression of *Lmo2*, a *Lyl-1* target[38], was

down-regulated, while *Tal-1* up-regulation might reflect some compensatory function.[3] The

consequences were apparent in GSEA analyses: both pathways and GO terms uncovered an up-

regulation of signaling pathways involved in embryo patterning (Wnt, Hox and Smad) in *Lyl-

1$^{LacZ/LacZ}$* MΦ-progenitors, as well as a highly modified collagen, integrin and cadherin usage

(Supplemental table 2). Accordingly, developmental trajectories were affected *(Figure 3D)*, with

the up-regulation in E9 *Lyl-1$^{LacZ/LacZ}$* MΦ-progenitors of gene sets related to "anterior-posterior

pattern specification" and "anatomical structure formation involved in morphogenesis", notably

skeletal and nervous system development.

GSEA and KEGG comparison of *Lyl-1^{LacZ/LacZ}* and WT MΦ-progenitors at E10 highlighted another

patterning modification, namely the down-regulation of gene sets involved in heart development

(Supplemental figure 2E; supplemental table 3), which might stem from a defective MΦ

development. The heart harbors three resident-MΦ subsets, two of which originate from the YS.[39]

Amongst the features that distinguish WT E9 MΦ^{Prim} progenitors from E10 MΦ-progenitors, the

enriched expression of MHC-II *(Figure 2F)* and poor expression of phagocytosis-related genes

(Supplemental figure 1I) at E9 also characterize one of the two YS-derived CCR2^{-} resident-MΦ

subsets in the heart.[39, 40] Therefore, a function for Lyl-1^{+} MΦ^{Prim} progenitors in heart development

may be considered. This observation reinforces the need to better characterize the contributions

of MΦ-progenitors from both primitive and transient-definitive waves to tissues harboring YS-

derived resident-MΦs.

The patterning defects highlighted in defective MΦ^{Prim}, might be responsible, at least in part,

for the significantly decreased litter size and increased perinatal lethality observed in *Lyl-1^{LacZ/LacZ}*

mice compared to WT *(Supplemental figure 2F)*, which indicates the requirement for Lyl-1 during

various developmental processes.

Defective MΦ^{Prim} development in Lyl-1 mutant YS

The analysis of *Lyl-1* expression in A1-A2-A3 subsets from *Cx3cr1^{WT/GFP}* YS indicated that *Lyl-1* is

expressed throughout MΦ-progenitor differentiation, with levels decreasing from A1 to A3

(Supplemental figure 3A). We monitored the distribution of A1-A2 and A3 MΦ subsets

(Supplemental figure 2A) at E10-YS, when all three subsets are present, using the *Cx3cr1^{WT/GFP}:Lyl-*

1^{LacZ/LacZ} strain. While the size of the whole MΦ population was not overtly modified, *Lyl-1-*

deficiency impacted the subset distribution, with increased A1 and reduced A2 and A3 pool sizes

(Figure 4A). Lyl-1 appears to regulate MΦ-progenitor differentiation towards mature MΦs. This

defect could result from the altered cytokine signaling uncovered in E9 mutant progenitors

through GSEA and IPA analyses *(Figure 4B; supplemental figure 3B)*. A limited or delayed

differentiation of E9 MΦ^{Prim} progenitors was supported by the down-regulated *Spi1*/PU.1 signaling

pathway *(Supplemental figure 3C; Supplemental table 4B)* and the decreased expression of

Ptprc/CD45, *Csfr1, Itgam*/CD11b and CD33 *(Figure 4C)*.

Within the MΦ lineage, Lyl-1 function during normal development would initially consist to

restrict the size of the MΦ^{Prim} progenitor pool and/or the duration of its production, which is

transient[6], as indicated by the maintenance of the intermediate mesoderm to MΦ-progenitor pool

observed in *Lyl-1$^{LacZ/LacZ}$* E8-YS. Indeed, the increased size of the MΦ-progenitor pool in E8-E9 YS

appears independent from the defective/delayed differentiation of MΦ-progenitors observed at

E10, since this process starts after E9.5.[10, 11, 15] Subsequently, the increased size of MΦ^{Prim}

progenitor pool in E10 *Lyl-1$^{LacZ/LacZ}$* YS likely results from a defective/delayed differentiation

mediated by a defective cytokine signaling, implying that during normal development, Lyl-1 would

promote the differentiation of MΦ^{Prim} progenitors.

Lyl-1$^{LacZ/LacZ}$ MΦ-progenitors were also deficient in the IFN signaling that characterize E9 MΦ^{Prim}

progenitors, notably *Irf8*, a factor involved in YS-MΦ and microglia development[21, 41] *(Figure 4D)*.

Compared to WT, MΦ-progenitors from *Lyl-1$^{lacZ/lacZ}$* E9-YS up-regulated the LXR/RXR activation

pathway *(Figure 4E)* and metabolic pathways, some enriched WT MΦ-progenitors at E10

(Butanoate, steroid) *(Supplemental table 1)*, and other not (Fructose/mannose, fatty acid) *(Figure*

3D). They were also less active in inflammatory signaling pathways, particularly through NFκb, a

factor known to interact with *Lyl-1*[42], and in TLR signaling *(Figure 4D; supplemental figure 3B, D-*

E; supplemental table 4B). Overlapping the DEGs identified in *Lyl-1$^{LacZ/LacZ}$* MΦ-progenitors at E9

and E10 identified the core signature of *Lyl-1*-deficiency, independent of the maturation occurring

between these stages *(Figure 4F)*.

Unfortunately, the co-existence of $M\Phi^{Prim}$ and $M\Phi^{T-Def}$ progenitors in E10-YS complicates the

attribution of gene expression changes to a stage-dependent maturation of $M\Phi^{Prim}$ progenitors or

to a signature specific to $M\Phi^{T-Def}$ progenitors. However, most pathways favored by E10 progenitors

were insensitive to *Lyl-1*-deficiency, except TLR signaling pathway that was down-regulated in E9

$Lyl\text{-}1^{LacZ/LacZ}$ progenitors, compared to WT.

Lyl-1-expressing MΦ-progenitors contribute to the fetal liver and brain

The FL[9] and brain[19, 22] are colonized as early as E9 by YS-derived resident-MΦ progenitors. We

evaluated the contribution of Lyl-1-expressing $M\Phi^{Prim}$ progenitors to these rudiments at E10

(Figure 5A). While E10-YS comprised $FDG^+/Lyl\text{-}1^+$ and $FDG^-/Lyl\text{-}1^-$ MΦ-progenitors and mature

$(F4/80^+)$ MΦ subsets *(Figure 1B)*, the brain from the same embryos essentially harbored FDG^+/Lyl-

1^+ MΦ-progenitors and MΦs. In contrast, both $FDG^+/Lyl\text{-}1^+$ and $FDG^-/Lyl\text{-}1^-$ MΦ-progenitors were

present in E10-FL, as in E10-YS, and MΦ-progenitors were more abundant in mutant FL than in WT

(Figure 5B).

We next focused on brain MΦs during the colonization stage, which lasts until E11.[43] At this

stage, microglia and perivascular, meningeal and choroid plexus MΦσ, collectively referred to as

BAMs, are all located in the brain mesenchyme and therefore undistinguishable.[16, 44] FACS-Gal

assay demonstrated that the whole $F4/80^+$ microglia/BAM expressed Lyl-1 throughout the

settlement period *(Figure 5C)*. The presence of $FDG^+/Lyl\text{-}1^+F4/80^+$ microglia/BAM at early stage of

brain colonization suggests that MΦs could participate to this step.

$Lyl\text{-}1^+$ $M\Phi^{Prim}$ progenitors and early microglia/BAM shared similar features, such as an early

appearance timing and low level of *Myb* expression *(Figure 5D)*, concordant with a *Myb-*

independent development of microglia.[14, 21] *Lyl-1* was also similarly expressed in A1-A2 and A3 MΦ

subsets from the YS and brain *(Figure 5E; supplemental figure 3A)*. *Lyl-1*-deficiency impacted the

distribution of MΦs subsets in E10 *Cx3cr1$^{WT/GFP}$:Lyl-1$^{LacZ/LacZ}$* brain: an increased A1 and a reduced

A3 pool size indicated that Lyl-1 regulates MΦ-progenitor differentiation in both YS and brain

(Figure 5F).

The proximity between YS MΦ^{Prim} progenitors and microglia was also apparent in RNA-seq.

data: E9 WT MΦ^{Prim} progenitors expressed significantly lower *Mrc1*/CD206 and higher *Sall3* levels

than E10 MΦ-progenitors, and a slightly increased *Sall1* level *(Figure 5G)*, a transcriptomic pattern

that characterizes microglia.[33, 43, 45] This partial bias toward a microglia signature suggests that the

first stage of microglia development program is already initiated in MΦ^{Prim} /"early

EMP"progenitors in E9-YS.

Lyl-1 inactivation impairs microglia development at two development stages

Having defined Lyl-1 implication during microglia/BAM settlement in the brain, we turned to

later development stages. Cytometry and database analyses[43] confirmed the continuous

expression of Lyl-1 in CD45^{low} microglia until adulthood *(Supplementary figure 4A)*. *LYL-1*

expression was also reported in microglia from healthy murine and human adults.[46, 47, 48, 49] We

examined the impact of *Lyl-1*-deficiency on microglia pool size during development. Microglia

quantification pointed to E12 as the first step impacted*.* The arrested increase of microglia pool in

Lyl-1$^{LacZ/LacZ}$ brain at E12 *(Figure 6A)* resulted from a reduced proliferation *(Figure 6B)* rather than

an increased apoptosis *(Supplementary figure 4B)*. Moreover, *Lyl-1*-deficiency provoked

morphological changes in E12 *Cx3cr1$^{WT/GFP}$:Lyl-1$^{LacZ/LacZ}$*, compared to *Cx3cr1$^{WT/GFP}$* microglia,

namely a reduced number and extent of ramifications *(Figure 6C; supplementary figure 4C-D)*.

From E14, the microglia pool size returned to levels similar to WT *(Figure 6A)*, probably due to the

highly reduced apoptosis level in *Lyl-1^{LacZ/LacZ}* microglia at E14 *(Supplementary figure 4B)*.

P0-P3 was identified as a second stage altered in *Lyl-1^{LacZ/LacZ}* microglia. At birth, the cellularity

of *Lyl-1^{LacZ/LacZ}* brain was significantly decreased compared to WT *(Figure 6D)*, which was not the

case at earlier stages *(Supplementary figure 4E)*. CD11b$^+$ cells recovery was also reduced (WT:

$140.96 \pm 0.91 \times 10^3$, n=9; *Lyl-1^{LacZ/LacZ}*: $87.18 \pm 0.37 \times 10^3$, n=9). Consequently, *Lyl-1*-deficiency triggered

a nearly 2-fold reduction of the microglia population *(Figure 6D)*. This perinatal reduction of

microglia appeared transient, since no difference with WT brain was observed in the adult

(Supplementary figure 4F). Transient decreases of microglia pool size, such as those we observed

at E12 and P0-P3 in *Lyl-1^{LacZ/LacZ}* mutant, have been reported to occur during normal development

in postnatal weeks 2-3[50], but also in *Cx3cr1* mutant mice during the 1st postnatal week[51]. This

indicates a highly dynamic control of the microglia pool size during key steps of neural

development that seems preserved in *Lyl-1* mutant, with the exception of the E12 and P0-P3 time-

points. At this later stage, the reduction of brain cellularity in *Lyl-1^{LacZ/LacZ}* mice points to Lyl-1 as a

possible regulator of the trophic function of microglia on brain cells[52, 53].

The identification of E12 and P0-P3 as key stages for Lyl-1 function in microglia development

was confirmed by RT-qPCR analyses of the expression of genes essential for MΦs (*Spi1/*PU.1, *Csf1r*,

Mafb) and/or microglia (*Runx1*, *Cx3cr1*, *Irf8*) development and function, of known regulators of

developmental hematopoiesis (*Tal-1, Lmo2, Runx1*) and related factors (*Tcf3/*E2A, *Tcf4/*E2.2)

(Figure 6E-F; supplementary figure 4G). Time-course analyses highlighted the down-regulation of

Csf1r, *Irf8* and *Lmo2* in *Lyl-1^{LacZ/LacZ}* microglia at E12 and P0-P3, while *Cx3cr1* was only decreased at

E12 *(Figure 6G)*. Note that *Lyl-1* expression was unmodified in *Cx3cr1^{GFP/GFP}* mutants *(Figure 6G)*.

Interestingly, *Cx3cr1*, as well as *Irf8* and *Lmo2*, belong to potential Lyl-1 target genes.[2]

Mafb expression levels in *Lyl-1*$^{LacZ/LacZ}$ microglia transiently decreased at P0-P3 and later returned

back to WT expression levels *(Figure 6H)*. As Mafb represses resident-MΦ self-renewal[54], the

recovery of a normal amount of microglia after birth may stem from this transient decrease.

Spi1/PU.1, *Tcf3*/E2A and *Tcf4*/E2.2 expression levels were unmodified in *Lyl-1*$^{LacZ/LacZ}$ microglia,

while *Runx1* expression was only affected after birth. *Tal-1* expression was decreased at E14 and

increased after birth, suggesting that this *Lyl-1* paralog[3] does not compensate *Lyl-1*-deficiency

during embryonic stages, but may do so at postnatal stages *(Supplementary figure 4G)*.

Remarkably, RNA-seq. results indicated that some genes deregulated in *Lyl-1*$^{LacZ/LacZ}$ microglia at

E12 and P0-P3 were also down-regulated in *Lyl-1*$^{LacZ/LacZ}$ MΦ-progenitors at E9 (*Csfr1:*

Supplementary figure 3C; *Lmo2:* ***Figure 3C***; *Irf8:* ***Figure 4D***; *Cx3cr1:* ***Figure 6I***). These deregulations

were transient, however in both locations and stages they coincided with a defective MΦ/microglia

differentiation.

Other genes enriched in microglia (*Fcrls*, *Mef2c*, *Maf*)[55] or involved in the maintenance of

microglia homeostasis (*P2ry12*)[56] were also expressed in E9 *Lyl-1*$^{LacZ/LacZ}$ MΦ-progenitors at a lower

level than in the WT, except *Lpr8* and *Aif-1*/Iba1 *(Figure 6I)*. These deregulations highlight again

shared features between MΦ^{Prim} progenitors and microglia/neural development which became

apparent upon *Lyl-1* inactivation considering the large number of neural signaling pathways up-

regulated in E9 *Lyl-1*$^{LacZ/LacZ}$ MΦ-progenitors *(Figure 3D)* and the relationship of the DEGs enriched

in E9 MΦ^{Prim} progenitors with brain formation and neuro-development uncovered in IPA analysis

(Supplementary figure 4H).

Based on the gene expression pattern of *Lyl-1*-deficient microglia and the signature of MΦ^{Prim}

progenitors in the early YS, a contribution of *Lyl-1*-deficiency to neurodevelopmental disorders may

be considered. Synaptic pruning and neural maturation, which characterize the perinatal phase of

microglia development[43], might be impaired in *Lyl-1*$^{LacZ/LacZ}$ embryos considering the defects

observed at P0-P3, the later key developmental stage regulated by Lyl-1. Indeed, *Lyl-1* deregulation

has been observed in datasets reporting pathological models of brain myeloid cells (http://research-

pub.gene.com/BrainMyeloidLandscape/#Mouse-gene/Mouse-gene/17095/geneReport.html)[57], as

well as in human neuropathies[58, 59], including the 19p13.13 micro-deletion neuro-developmental

disabilities.[60] However, since Lyl-1 is expressed in endothelial cells, including in the brain[61], a

contribution of *LYL-1*-deficient endothelial cells to these diseases must be considered.

Altogether, our findings reveal Lyl-1 as a key factor regulating the production and differentiation

of YS MΦ-progenitors and the development of microglia. Lyl-1 is the least studied partners of the

transcription factor complex that regulates developmental hematopoiesis. The development of

more appropriate models is required to precise Lyl-1 functions in microglia and determine its role

in the development of other resident-MΦs populations.

MATERIALS AND METHODS

Mice. The following mouse strains were used: C57BL/6 (Charles Rivers Laboratories), called wild type (WT); *Lyl-1*^{LacZ,5}; *Cx3cr1*^{GFP,62}; *Cx3cr1*$^{GFP/GFP}$:*Lyl-1*$^{LacZ/LacZ}$ double knock-in strain (breeding schemes: **supplemental information**). Experiments were conducted in compliance with French/European laws, under authorized project *#2016-030-5798*, approved by officially accredited local institutional animal (committee n°26) and French "Ministère de la Recherche" ethics boards.

Tissues. E7.5-E10.5 YS and E10-FL were dissected as described.[9, 63] For cytometry analyses, whole E9-E11 brains were prepared as described.[22] After E12, microglia were recovered following Percoll (Sigma) separation, as described.[64] The number of microglia per brain was estimated by reporting the percentage of CD11b$^+$CD45^{low}F4/80$^+$ microglia to the cellularity of the corresponding sample.

Tissue culture. YS explants, maintained for 1 day in organ culture in plates containing OptiMEM+Glutamax, 1% Penicillin-streptomycin, 0.1% β-mercaptoethanol (ThermoFisher) and 10% fetal calf serum (Hyclone), are referred to as OrgD1-YS.

In clonogenic assays, YS suspension or sorted cells were plated in triplicate (3x10^3 or 100-150 cells/mL) in Methocult® M3234 (StemCell Technologies) always supplemented the cytokines listed in **supplemental table 5**. Colonies were scored at day 5 for primitive erythroblasts and day 7 for other progenitors.

Flow cytometry. Cells, stained with antibodies listed in **supplemental table 6**, for 30 min. on ice, were acquired (Canto II) or sorted (FACS-Aria III or Influx, BD Biosciences) and analyzed using FlowJo (Treestar) software.

The β-Galactosidase substrate fluorescein di-β-galactopyranoside (FDG; Molecular probe), was

used as reporter for Lyl-1 expression in FACS-Gal assay.[65,66] For apoptosis analysis, microglia were

immune-stained and incubated with Annexin V-FITC. 7AAD was added before acquisition. For

proliferation assays, pregnant females (12 gestational days) were injected with BrdU (10μM) and

sacrificed 2 hours later. Microglia were isolated, immune-stained and prepared according to kit

instruction (BD Pharmingen 552598). BrdU incorporation was revealed using anti-BrdU-APC.

Brain imaging. To assess microglia morphology, the midbrain was dissected from $Cx3cr1^{WT/GFP}$:Lyl-

$1^{WT/WT}$ and $Cx3cr1^{WT/GFP}$:Lyl-1$^{LacZ/LacZ}$ embryos, immune-labeled and image stacks collected using

Leica SP8 confocal microscope *(See Supplemental information)*.

RT-qPCR analyses. RNA was extracted using Trizol. After cDNA synthesis (SuperScript™ VILO™

Master-Mix reverse transcriptase, ThermoFisher), Real Time (RT)-PCR was performed (SYBR

Premix Ex TaqII, Takara Bio). Reference genes were *Actin*, *Hprt* and *Tubulin*. Gene expressions were

normalized to the values obtained for E10-YS MΦ-progenitors, E12 WT Lin$^-$Sca$^+$cKit$^+$ (LSK)

progenitors or E12 WT microglia. Primers are listed in *supplemental table 7*.

RNA-seq. MΦ-progenitors (CD45$^+$CD11b$^+$Kit$^+$) were sorted from E9 (MΦ^{Prim} progenitors) and E10

(MΦ^{Prim} + MΦ^{T-Def} progenitors) YS from WT or *Lyl-1$^{LacZ/LacZ}$* embryos. See *supplemental information*

for sample processing, RNA-sequencing and analysis protocols. RNA-seq. data (accession number

E-MTAB-9618) were deposited in EMBL-EBI ArrayExpress database (www.ebi.ac.uk/arrayexpress).

Data were analyzed using Ingenuity® Pathway Analysis (IPA, QIAGEN)[67], Gene set enrichment

analysis (GSEA)[68, 69], Morpheus and Venny softwares.

Statistical analysis. Statistical tests were performed using Prism 7 (GraphPad). Statistical

significance is indicated by p-values and/or as $*p<0.05$, $**p<0.01$, $***p<0.001$ and $****p<0.0001$.

Acknowledgements.

The authors thank Julien Bertrand for critical reading of the manuscript. We are grateful to the

staff of the facilities at Gustave Roussy, the animal facility (PFEP, UMS AMMICa UMS 3655/US23,

directed by P. Gonin), the imaging facility (PFIC, UMS AMMICa UMS 3655/US23, directed by C.

Laplace-Builhe), the genomic facility directed by N. Droin, the bioinformatics facility (G. Meurice),

directed by M. Deloger.

This work was supported by fundings from Institut National de la Santé et de la Recherche

Médicale to W. Vainchenker, I. Plo and H. Raslova, from Centre National de la Recherche

Scientifique and Université de Paris-Saclay to I. Godin, from grants INCA PLBio to I. Plo, "Ligue

Nationale contre le Cancer" Certified Team to H. Raslova, "Association pour la Recherche sur le

Cancer" (n°4878) to I. Godin, Gustave Roussy (TA DERE 17) to D. Ren, Grant Agency of the Czech

Republic (GACR n°19-23154S) to D. Filipp and from fellowships from "Association pour la

Recherche sur le Cancer" to A.-L. Kaushik; "Société Française d'Hématologie" to S. Wang and

Chinese Scholarship Council fellowships to S. Wang and D. Ren.

Author Information.

The authors declare no competing financial interests.

Correspondence should be addressed to I.G. (Isabelle.Godin@gustaveroussy.fr).

REFERENCES

1. Pina, C. & Enver, T. Differential contributions of haematopoietic stem cells to foetal and adult haematopoiesis: insights from functional analysis of transcriptional regulators. *Oncogene* **26**, 6750-6765 (2007).

2. Wilson, N.K. *et al.* Combinatorial Transcriptional Control In Blood Stem/Progenitor Cells: Genome-wide Analysis of Ten Major Transcriptional Regulators. *Cell Stem Cell* **7**, 532-544 (2010).

3. Curtis, D.J., Salmon, J.M. & Pimanda, J.E. Concise Review: Blood Relatives: Formation and regulation of hematopoietic stem cells by the basic helix-loop-helix transcription factors stem cell leukemia and lymphoblastic leukemia-derived sequence 1. *Stem Cells* **30**, 1053-1058 (2012).

4. Porcher, C., Chagraoui, H. & Kristiansen, M.S. SCL/TAL1: a multifaceted regulator from blood development to disease. *Blood* **129**, 2051-2060 (2017).

5. Capron, C. *et al.* The SCL relative LYL-1 is required for fetal and adult hematopoietic stem cell function and B-cell differentiation. *Blood* **107**, 4678-4686 (2006).

6. McGrath, K.E., Frame, J.M. & Palis, J. Early hematopoiesis and macrophage development. *Seminars in Immunology* **27**, 379-387 (2015).

7. Palis, J. Hematopoietic stem cell-independent hematopoiesis: emergence of erythroid, megakaryocyte, and myeloid potential in the mammalian embryo. *FEBS Lett* **590**, 3965-3974 (2016).

8. Cumano, A. & Godin, I. Ontogeny of the hematopoietic system. *Annu Rev Immunol* **25**, 745-785 (2007).

9. Kieusseian, A., Brunet de la Grange, P., Burlen-Defranoux, O., Godin, I. & Cumano, A. Immature hematopoietic stem cells undergo maturation in the fetal liver. *Development* **139**, 3521-3530 (2012).

10. Palis, J., Robertson, S., Kennedy, M., Wall, C. & Keller, G. Development of erythroid and myeloid progenitors in the yolk sac and embryo proper of the mouse. *Development* **126**, 5073-5084 (1999).

11. Bertrand, J.Y. *et al.* Three pathways to mature macrophages in the early mouse yolk sac. *Blood* **106**, 3004-3011 (2005).

12. Tober, J. *et al.* The megakaryocyte lineage originates from hemangioblast precursors and is an integral component both of primitive and of definitive hematopoiesis. *Blood* **109**, 1433-1441 (2007).

13. Sumner, R., Crawford, A., Mucenski, M. & Frampton, J. Initiation of adult myelopoiesis can occur in the absence of c-Myb whereas subsequent development is strictly dependent on the transcription factor. *Oncogene* **19**, 3335-3342 (2000).

14. Schulz, C. *et al.* A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science* **336**, 86-90 (2012).

15. Hoeffel, G. & Ginhoux, F. Fetal monocytes and the origins of tissue-resident macrophages. *Cell Immunol* **330**, 5-15 (2018).

16. Utz, S.G. *et al.* Early Fate Defines Microglia and Non-parenchymal Brain Macrophage Development. *Cell* **181**, 557-573 e518 (2020).

17. Ginhoux, F. & Guilliams, M. Tissue-Resident Macrophage Ontogeny and Homeostasis. *Immunity* **44**, 439-
449 (2016).

18. Mass, E. Delineating the origins, developmental programs and homeostatic functions of tissue-resident
macrophages. *Int Immunol* **30**, 493-501 (2018).

19. Ginhoux, F. *et al.* Fate Mapping Analysis Reveals That Adult Microglia Derive from Primitive Macrophages.
Science **330**, 841-845 (2010).

20. Gomez Perdiguero, E. *et al.* Tissue-resident macrophages originate from yolk-sac-derived erythro-myeloid
progenitors. *Nature* **518**, 547-551 (2015).

21. Kierdorf, K. *et al.* Microglia emerge from erythromyeloid precursors via Pu.1- and Irf8-dependent
pathways. *Nat Neurosci* **16**, 273-280 (2013).

22. Alliot, F., Godin, I. & Pessac, B. Microglia derive from progenitors, originating from the yolk sac, and which
proliferate in the brain. *Brain Res Dev Brain Res* **117**, 145-152 (1999).

23. Wu, Y. & Hirschi, K.K. Tissue-Resident Macrophage Development and Function. *Frontiers in Cell and*
Developmental Biology **8** (2021).

24. Herbomel, P., Thisse, B. & Thisse, C. Zebrafish early macrophages colonize cephalic mesenchyme and
developing brain, retina, and epidermis through a M-CSF receptor-dependent invasive process. *Dev Biol*
238, 274-288 (2001).

25. Ferrero, G. *et al.* Embryonic Microglia Derive from Primitive Macrophages and Are Replaced by cmyb-
Dependent Definitive Microglia in Zebrafish. *Cell Reports* **24**, 130-141 (2018).

26. Wittamer, V. & Bertrand, J.Y. Yolk sac hematopoiesis: does it contribute to the adult hematopoietic
system? *Cell Mol Life Sci* (2020).

27. Azzoni, E. *et al.* Kit ligand has a critical role in mouse yolk sac and aorta-gonad-mesonephros
hematopoiesis. *EMBO reports* (2018).

28. Espin-Palazon, R. *et al.* Proinflammatory signaling regulates hematopoietic stem cell emergence. *Cell* **159**,
1070-1085 (2014).

29. Giroux, S. *et al.* lyl-1 and tal-1/scl, two genes encoding closely related bHLH transcription factors, display
highly overlapping expression patterns during cardiovascular and hematopoietic ontogeny. *Gene Expr*
Patterns **7**, 215-226 (2007).

30. Cumano, A., Dieterlen-Lievre, F. & Godin, I. Lymphoid potential, probed before circulation in mouse, is
restricted to caudal intraembryonic splanchnopleura. *Cell* **86**, 907-916 (1996).

31. Cumano, A., Ferraz, J.C., Klaine, M., Di Santo, J.P. & Godin, I. Intraembryonic, but not yolk sac
hematopoietic precursors, isolated before circulation, provide long-term multilineage reconstitution.
Immunity **15**, 477-485 (2001).

32. Hoeffel, G. *et al.* C-Myb+ Erythro-Myeloid Progenitor-Derived Fetal Monocytes Give Rise to Adult Tissue-
Resident Macrophages. *Immunity* **42**, 665-678 (2015).

33. Mass, E. *et al.* Specification of tissue-resident macrophages during organogenesis. *Science* **353** (2016).

34. Espin-Palazon, R., Weijts, B., Mulero, V. & Traver, D. Proinflammatory Signals as Fuel for the Fire of
Hematopoietic Stem Cell Emergence. *Trends in Cell Biology* **28**, 58-66 (2018).

35. Mariani, S.A. *et al.* Pro-inflammatory Aorta-Associated Macrophages Are Involved in Embryonic
Development of Hematopoietic Stem Cells. *Immunity* (2019).

36. Palpant, N.J. *et al.* Chromatin and Transcriptional Analysis of Mesoderm Progenitor Cells Identifies HOPX
as a Regulator of Primitive Hematopoiesis. *Cell Reports* **20**, 1597-1608 (2017).

37. Chiu, S.K. *et al.* Shared roles for Scl and Lyl1 in murine platelet production and function. *Blood* **134**, 826-
835 (2019).

38. McCormack, M.P. *et al.* Requirement for Lyl1 in a model of Lmo2-driven early T-cell precursor ALL. *Blood*
122, 2093-2103 (2013).

39. Epelman, S. *et al.* Embryonic and Adult-Derived Resident Cardiac Macrophages Are Maintained through
Distinct Mechanisms at Steady State and during Inflammation. *Immunity* **40**, 91-104 (2014).

40. Leid, J. *et al.* Primitive Embryonic Macrophages are Required for Coronary Development and Maturation.
Circulation Research **118**, 1498 (2016).

41. Hagemeyer, N. *et al.* Transcriptome-based profiling of yolk sac-derived macrophages reveals a role for Irf8
in macrophage maturation. *EMBO J* **35**, 1730-1744 (2016).

42. Ferrier, R. *et al.* Physical interaction of the bHLH LYL1 protein and NF-kappaB1 p105. *Oncogene* **18**, 995-
1005 (1999).

43. Matcovitch-Natan, O. *et al.* Microglia development follows a stepwise program to regulate brain
homeostasis. *Science* **353**, aad8670 (2016).

44. Goldmann, T. *et al.* Origin, fate and dynamics of macrophages at central nervous system interfaces. *Nature*
Immunology **17**, 797-805 (2016).

45. Lavin, Y. *et al.* Tissue-Resident Macrophage Enhancer Landscapes Are Shaped by the Local
Microenvironment. *Cell* **159**, 1312-1326 (2014).

46. Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells
of the cerebral cortex. *J Neurosci* **34**, 11929-11947 (2014).

47. Zeisel, A. *et al.* Brain structure. Cell types in the mouse cortex and hippocampus revealed by single-cell
RNA-seq. *Science* **347**, 1138-1142 (2015).

48. Wehrspaun, C.C., Haerty, W. & Ponting, C.P. Microglia recapitulate a hematopoietic master regulator
network in the aging human frontal cortex. *Neurobiol Aging* **36**, 2443 e2449-2443 e2420 (2015).

49. Bennett, M.L. *et al.* New tools for studying microglia in the mouse and human CNS. *Proceedings of the*
National Academy of Sciences **113**, E1738-E1746 (2016).

50. Zhan, Y. *et al.* Deficient neuron-microglia signaling results in impaired functional brain connectivity and
social behavior. *Nat Neurosci* **17**, 400-406 (2014).

51. Paolicelli, R.C. *et al.* Synaptic pruning by microglia is necessary for normal brain development. *Science* **333**,
1456-1458 (2011).

52. Antony, J.M., Paquin, A., Nutt, S.L., Kaplan, D.R. & Miller, F.D. Endogenous microglia regulate development

of embryonic cortical precursor cells. *J Neurosci Res* **89**, 286-298 (2011).

53. Ueno, M. *et al.* Layer V cortical neurons require microglial support for survival during postnatal

development. *Nat Neurosci* **16**, 543-551 (2013).

54. Soucie, E.L. *et al.* Lineage-specific enhancers activate self-renewal genes in macrophages and embryonic

stem cells. *Science* **351** (2016).

55. Crotti, A. & Ransohoff, R.M. Microglial Physiology and Pathophysiology: Insights from Genome-wide

Transcriptional Profiling. *Immunity* **44**, 505-515 (2016).

56. Bisht, K., Sharma, K. & Tremblay, M.-È. Chronic stress as a risk factor for Alzheimer's disease: Roles of

microglia-mediated synaptic remodeling, inflammation, and oxidative stress. *Neurobiology of Stress* **9**, 9-

21 (2018).

57. Friedman, B.A. *et al.* Diverse Brain Myeloid Expression Profiles Reveal Distinct Microglial Activation States

and Aspects of Alzheimer's Disease Not Evident in Mouse Models. *Cell Reports* **22**, 832-847 (2018).

58. Colangelo, V. *et al.* Gene expression profiling of 12633 genes in Alzheimer hippocampal CA1: transcription

and neurotrophic factor down-regulation and up-regulation of apoptotic and pro-inflammatory signaling.

J Neurosci Res **70**, 462-473 (2002).

59. Thomas, D.M., Francescutti-Verbeem, D.M. & Kuhn, D.M. Gene expression profile of activated microglia

under conditions associated with dopamine neuronal damage. *Faseb J* **20**, 515-517 (2006).

60. Nimmakayalu, M. *et al.* Apparent germline mosaicism for a novel 19p13.13 deletion disrupting NFIX and

CACNA1A. *Am J Med Genet A* **161A**, 1105-1109 (2013).

61. Pirot, N. *et al.* LYL1 activity is required for the maturation of newly formed blood vessels in adulthood.

Blood **115**, 5270-5279 (2010).

62. Jung, S. *et al.* Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green

fluorescent protein reporter gene insertion. *Mol Cell Biol* **20**, 4106-4114 (2000).

63. Bertrand, J.Y., Giroux, S., Cumano, A. & Godin, I. Hematopoietic stem cell development during mouse

embryogenesis. *Methods Mol Med* **105**, 273-288 (2005).

64. Mildner, A. *et al.* Microglia in the adult brain arise from Ly-6ChiCCR2+ monocytes only under defined host

conditions. *Nat Neurosci* **10**, 1544-1553 (2007).

65. Fiering, S.N. *et al.* Improved FACS-Gal: flow cytometric analysis and sorting of viable eukaryotic cells

expressing reporter gene constructs. *Cytometry* **12**, 291-301 (1991).

66. Guo, W. & Wu, H. Detection of LacZ expression by FACS-Gal analysis. *Nature Protocol Exchange* (2008).

67. Krämer, A., Green, J., Pollard, J., Jr. & Tugendreich, S. Causal analysis approaches in Ingenuity Pathway

Analysis. *Bioinformatics* **30**, 523-530 (2014).

68. Mootha, V.K. *et al.* PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately

downregulated in human diabetes. *Nat Genet* **34**, 267-273 (2003).

69. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting

genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550 (2005).

 FIGURE LEGENDS

Figure 1: Lyl-1 expression marks MΦ^{Prim} progenitors in the early YS

A. Lyl-1-deficiency leads to an increased production of MΦ-progenitors in the early YS: Left: Clonogenic

potential of E8 OrgD1-YS cells: production of MΦ-progenitors (CFU-M) in WT and *Lyl-1$^{LacZ/LacZ}$* OrgD1-YS

(n=3-5, 3-6 YS per sample; mean $\pm$ s.e.m.; Unpaired, two-tailed *t*-Test). The size of the MΦ colonies and the

cell morphology were similar for the 3 genotypes (data not shown). Right: distribution of other progenitors

with a myeloid potential (EMP and GM) in WT, *Lyl-1$^{WT/LacZ}$* and *Lyl-1$^{LacZ/LacZ}$* E8 OrgD1-YS.

B. Lyl-1 expression in MΦ-progenitors: FACS-Gal assay, using the β-Gal fluorescent substrate FDG was used

as a reporter for Lyl-1 expression. While all MΦ-progenitors in E9-YS (left panel) expressed FDG/Lyl-1, E9.5

and E10-YS (middle panels) harbored two MΦ-progenitor subsets discriminated by their FDG/Lyl-1

expression. FDG$^+$/Lyl-1$^+$ and FDG$^-$/Lyl-1$^-$ mature MΦs (CD11b$^+$F4/80$^+$) also coexisted in E10-YS (right). The

contour plots in WT samples indicate the level of non-specific background β-Gal activity/FDG labeling in WT

samples. Representative profiles of 3 independent samples, each consisting of 3-4 YS *(See the gating*

strategy in **Supplemental figure 1A**).

C. MΦ^{Prim} progenitors express Lyl-1. Upper panel: Flow cytometry profiles of WT (left) and *Lyl-1$^{WT/LacZ}$*

(middle left) E8-YS (0-3S). CD11b$^+$CD31$^-$ MΦs (top gate) correspond to maternal MΦs present in E8-YS.[11] All

CD11b$^+$CD31$^+$ MΦ-progenitors (lower gate) displayed FDG/Lyl-1 expression.

D. RT-qPCR quantification of *Myb* expression levels: Kit$^+$CD45$^+$CD11b$^+$ progenitors were sorted from WT

E9-YS, WT and *Lyl-1$^{WT/LacZ}$* E10-YS, and from FDG/Lyl-1 positive and negative fractions of MΦ-progenitors

from *Lyl-1$^{WT/LacZ}$* E10-YS. Lin$^-$Sca$^+$Kit$^+$ (LSK) progenitors from WT E12-FL were used as positive control.

FDG$^+$/Lyl-1$^+$ MΦ-progenitors from E10-YS expressed *Myb$^{Low/Neg}$* levels similar to MΦ^{Prim} progenitors from E9-

YS. The FDG$^-$/Lyl-1$^-$ fraction expressed significantly higher *Myb* levels, similar to LSK cells from E12-FL. *Myb*

expression levels, shown on a Log^2 scale, were normalized to the mean expression value obtained for WT

E10-YS, considered as 1 (Unpaired, two-tailed *t*-Test).

E. FDG/Lyl-1 positive and negative myeloid progenitors produce a distinct progeny: Clonogenic assays

characterization of the type of progenitors produced by myeloid progenitors (Ter119$^-$Kit$^+$CD45$^+$CD11b$^+$)

sorted from WT and *Lyl-1$^{WT/LacZ}$* E9-YS (<18 S; n=7) and E10-YS (n=15) in 3 independent experiments. At E10,

myeloid progenitors from *Lyl-1$^{WT/LacZ}$* YS were subdivided into FDG/Lyl-1 negative (n=15) and positive (n=12)

fractions (5 independent experiments). Samples were biological replicates comprising 6-8 YS. 100 to 150

Kit$^+$CD45$^+$CD11b$^+$ cells per condition were platted in triplicate. All samples produced few non-myeloid

contaminants, such as EMk and EMP in similar, non-significant amounts. FDG$^+$/Lyl-1$^+$ progenitors essentially

produced MΦ colonies, while FDG$^-$/Lyl-1$^-$ progenitors produced also GM and G colonies.

Figure 2: Distinct features of WT MΦ-progenitors at E9 and E10

A. Differentially expressed genes in MΦ-progenitors (Kit⁺CD45⁺CD11b⁺) sorted from WT and *Lyl-1^LacZ/LacZ* YS at E9 and E10. Upper panel: Unsupervised principal component analysis (PCA) plot positioned E9 and E10 MΦ-progenitors in two distinct groups, followed by segregation of WT and *Lyl-1^LacZ/LacZ* samples. Lower panel: Volcano plot of E9 WT vs E10 WT MΦ-progenitors. Red and green dots indicate genes with statistically significant changes in expression level. (*p*-value <0.05, absolute fold change≥2.0) (NDE: not deregulated genes; DE-Up: up-regulated genes; DE-Down: down-regulated genes).

B. Upper panel: Venn diagram comparing DEGs in E9 WT versus E10 WT MΦ-progenitors to the EMP (Left) or MΦ signatures defined by Mass et al.[33] (GEO accession number GSE81774). The number and percentage of DEGs common to the EMP or MΦ signatures is shown.

Lower panel: Expression profiles of the overlapping genes identified in the Venn diagram (Heatmap displays transformed log2-expression values; Unpaired *t*-Test, two-tailed). Note the higher expression at E10 of genes involved in erythroid (Globins: Pink arrow; Transcription factors: green arrow), and megakaryocytic and granulocytic-related genes (blue arrow), and of *Mrc1*/CD206 (Asterisk).

C. Relative expression levels of *Gata1* and *Spi1/*PU.1, indicated by their relative Transcripts per million kilo-bases (TPM).

D. Enriched Pathways in E9 and E10 WT MΦ-progenitors with absolute z-score ≥2, from QIAGEN's Ingenuity® Pathway Analysis (IPA). Bars: minus log of the p-value of each canonical pathway; Orange line: threshold p-value of 0.05. Ratio: genes detected/genes per pathway.

E. Expression profiles of DEGs related to IFNγ and IFNβ response, identified by g:Profiler. (Heatmap displays transformed log2-expression values; unpaired t-Test, two-tailed).

F. Expression profiles of DEGs related to MHC-II complex (Heatmap displays transformed log2-expression values; unpaired *t*-Test, two-tailed).

G. Expression profiles of DEGs related to cytokine signaling (Heatmap displays transformed log2-expression values; unpaired *t*-Test, two-tailed).

Figure 3: Lyl-1 regulates the production of E8 MΦ^Prim progenitors.

A. In clonogenic assays, the number of MΦ colonies obtained from E8-YS (0-3S) was increased in *Lyl-1^WT/LacZ* and *Lyl-1^LacZ/LacZ* compared to WT. The majority of the 25-30 colonies per YS were Ery^P (60 to 80% in each 3 genotypes). Other progenitors were occasionally and randomly detected in WT and mutant samples (less than one EMP (0.81%±0.66; n=3) and/or GM progenitor per E8-YS), confirming that the assay was performed at a time when MΦ^T-Def progenitors were absent. (n=3-5, 5-10 YS per sample; plots show mean ± s.e.m.; Unpaired, two-tailed *t*-Test).

B. Relative expression levels of the CD41 coding gene *Itg2b* in WT and *Lyl-1^LacZ/LacZ* MΦ-progenitors at E9 (unpaired *t*-Test, two-tailed).

C. Relative expression levels of TF regulating hematopoietic progenitor emergence in *Lyl-1^{LacZ/LacZ}* MΦ-

progenitors compared to WT at E9 (unpaired, two-tailed *t*-Test).

D. GSEA pathways (Top; FDR q-Value <0.29) and GO terms (Bottom; FDR q-value <0.01) enriched in E9 *Lyl-*

1^{LacZ/LacZ} compared to E9 WT MΦ-progenitors. Highlighted are the pathways specifically related to embryo

patterning (blue) and to the development of skeletal (green) and nervous systems (yellow). Pink arrows

point to changes related to metabolic pathways.

Figure 4: Defective differentiation of Lyl-1^{LacZ/LacZ} MΦ-progenitors from E10-YS.

A. Distribution of A1-A2 and A3 MΦ subsets in E10-YS from *Cx3cr1^{WT/GFP}:Lyl-1^{WT/WT}*, *Cx3cr1^{WT/GFP}:Lyl-1^{WT/LacZ}*

and *Cx3cr1^{WT/GFP}: Lyl-1^{LacZ/LacZ}* embryos. While the size of the whole MΦ population is similar in the three

genotypes (Top panel), *Lyl-1* deficiency leads to a modified distribution of the MΦ subsets (middle and

lower panel) with an increased size of the A1 subset and a reduced A3 pool (5-12 independent analyses, 6-

8 YS per sample. Plots show mean ± s.e.m.; Unpaired, two-tailed *t*-Test).

B. GSEA pathway indicates a deficit in Jak1-Stat signaling in *Lyl-1^{LacZ/LacZ}* MΦ-progenitors compared to WT

at E9 (NES: normalized enrichment score; FDR: false discovery rate).

C. Relative expression levels (read counts) of hematopoietic markers in WT and *Lyl-1^{LacZ/LacZ}* MΦ-

progenitors at E9 (unpaired *t*-Test, two-tailed).

D. Top 1 GSEA pathway indicates that the IFN signaling pathway (left) which characterize E9 MΦ^{Prim}

progenitors, and particularly *Irf8* (right), is defective in *Lyl-1^{LacZ/LacZ}* MΦ-progenitors (NES: normalized

enrichment score; FDR: false discovery rate).

E. From the 53 canonical pathways identified by IPA in the DEGs, 9 were enriched with an absolute Z score

≥ 1. Bars: minus log of the *p*-value of each canonical pathway; Orange line: threshold p-value of 0.05. Ratio:

genes detected/genes per pathway.

F. Upper panel: Venn diagram comparing the DEGs in E9 *Lyl-1^{LacZ/LacZ}* vs E9 WT to those in E10 *Lyl-1^{LacZ/LacZ}* vs

E10 WT MΦ-progenitors. Lower panel: Expression profiles of the DEGs common to both stages identified

by the Venn comparison (Heatmap displays transformed log2-expression values; unpaired *t*-Test, two-

tailed).

Figure 5: Contribution of Lyl-1-expressing MΦ-progenitors to the fetal liver and brain.

A. Left panel: All MΦ-progenitors from E10-Brain (Left) expressed Lyl-1, contrary to the corresponding YS

(Figure 1B right) which harbor both FDG^+/Lyl-1^+ and FDG^-/Lyl-1^- subsets. MΦ-progenitors from E10 *Lyl-*

1^{LacZ/LacZ} FL (right) harbored both FDG^+/Lyl-1^+ and FDG^-/Lyl-1 MΦ-progenitor subsets. Right panel: in E10-

brain, mature MΦs (CD11b^+F4/80^+ gate) were all FDG^+/Lyl-1^+. The contour plots in WT samples tTop panel)

indicate the level of non-specific background β-Gal activity/FDG labeling in WT samples. Representative

profiles of 3 independent samples, each consisting of 3-4 E10-Brain or 8-12 E10-FL.

B. Quantification of cKit⁺CD45⁺CD11b⁺ MΦ-progenitors in E10-FL (plots show mean ± s.e.m.; Unpaired, two-tailed *t*-Test).

C. Lyl-1 marks the entire F4/80+ microglia/BAM population from the onset of brain colonization. FDG/Lyl-1 expression in F4/80+ microglia/BAM from the brain of *Lyl-1$^{WT/LacZ}$* and *Lyl-1$^{LacZ/LacZ}$* embryos at E9 to E11. The rare CD11b⁺ F4/80$^{low-neg}$ cells present in the brain at E9 are FDG⁺/Lyl-1⁺ (Top Panel)**.** Grey histograms indicate non-specific background β-Gal activity/FDG levels in WT samples.

D. MΦ-progenitors from E10-brain express *Myb* levels similar to E9-YS MΦ^{Prim} progenitors. RT-qPCR quantification of *Myb* expression levels in Kit⁺CD45⁺CD11b⁺ MΦ-progenitors sorted from WT E9-YS and from WT and *Lyl-1$^{WT/LacZ}$* brain at E10. Lin⁻Sca⁺cKit⁺ (LSK) progenitors from WT E12-FL were used as positive control. *Myb* expression levels, shown on a Log² scale, were normalized to the mean expression value obtained for WT E10-YS, considered as 1 (Unpaired, two-tailed *t*-Test).

E. Heatmap expression profile of the genes that mark the development of tissue resident-MΦs in WT E9 and E10 MΦ-progenitors (Heatmap displays transformed log2-expression values; Unpaired, two-tailed *t*-Test).

F. RT-qPCR analyses of *Lyl-1* expression in A1 to A3 MΦ subsets isolated from *Cx3cr1$^{WT/GFP}$* brain at E10. *Lyl-1* is expressed by the 3 subsets, with levels decreasing with differentiation. Expression levels were normalized to the mean value obtained for *Cx3cr1$^{WT/GFP}$* YS A1 progenitors (n=3).

G. Defective differentiation of brain MΦ-progenitor in Lyl-1 mutant embryos. Distribution of A1-A2 and A3 MΦ subsets in E10 brain from *Cx3cr1$^{WT/GFP}$:Lyl-1$^{WT/WT}$*, *Cx3cr1$^{WT/GFP}$:Lyl-1$^{WT/LacZ}$* and *Cx3cr1$^{WT/GFP}$:Lyl-1$^{LacZ/LacZ}$* embryos. The size of the whole MΦ population was similar in the three genotypes (Top panel), but Lyl-1 deficiency modified the distribution of the MΦ subsets (middle and lower panel) with an increased size of the A1 subset and a reduced A3 pool (5-12 independent analyses, 6-8 brains per sample. Plots show mean ± s.e.m.; Unpaired, two-tailed t Test).

Figure 6: Lyl-1 deficiency leads to transient reductions of the microglia pool at E12 and P0-P3.

A. Quantification of the microglia population in E12 and E14 brain showing the decreased size of the microglia pool at E12 and its recovery to a normal pool size at E14. Plots show mean ± s.e.m.; Two tailed, unpaired *t*-test.

B. The decreased microglia pool at E12 may result from a reduced proliferation, as shown by the two folds decrease (right) of BrdU-labeled cells in *Lyl-1$^{LacZ/LacZ}$* (middle) compared to WT (left) brains. Plots show mean±s.e.m.; Two tailed, unpaired *t*-test.

C. At E12, *Cx3cr1$^{WT/GFP}$:Lyl-1$^{LacZ/LacZ}$* microglia displayed a reduced number and extent of ramifications compared to their *Cx3cr1$^{WT/GFP}$* counterpart. Bottom: Microglia morphology was classified into subtypes depending on the number of main ramifications (A: none, B: 2, C: 3 and D:>3). Top: Microglia deprived of ramifications predominated in *Lyl-1*-deficient microglia. 65 and 61 cells were respectively acquired from

the midbrain of E12 *Cx3cr1*^{WT/GFP} and *Cx3cr1*^{WT/GFP}:*Lyl-1*^{LacZ/LacZ} embryos (for each genotype, brains from 12

embryos were acquired in 3 independent experiments). Microglia were identified by *Cx3cr1*-driven GFP

expression and F4/80-APC immuno-staining. Bar=10μm. Plots show mean±s.e.m.; Two tailed, unpaired *t*-

test.

D. In *Lyl-1*^{LacZ/LacZ} newborns, the cellularity of the brain was consistently lower than in WT (left), and so was

the estimated microglia number (right). Plots show mean ± s.e.m.; Two tailed, unpaired *t*-test.

E. Kinetic evolution of *Lyl-1* expression levels in WT microglia from embryonic stages to adulthood. An

increased expression of *Lyl-1* from embryonic stages to adulthood was also inferred from timeline RNA-seq.

data.[43] (GEO accession number GSE79812).

F. Quantitative RT-PCR analyses also point to E12 and P0 as key development stages regulated by Lyl-1.

CD11b⁺F4/80⁺CD45^{low} microglia were isolated at sequential development stages. Bar graphs show the

kinetic of expression of genes modified in *Lyl-1*^{LacZ/LacZ} microglia (arrowheads), normalized to the mean

expression value in WT E12 microglia (n=3). Error bars indicate s.e.m. Two tailed, unpaired *t*-test.

G. *Cx3Cr1* and *Lyl-1* expression in mutant microglia. The expression level of *Cx3CR1*, analyzed as in F, was

decreased in Lyl-1 mutants at E12 (left), while *Lyl-1* expression levels were unmodified in *CX3CR1*^{GFP/GFP}

microglia at E12 and in newborns (right).

H. *Mafb* expression in mutant microglia. *Mafb* expression level, analyzed as in F, was reduced in the

microglia of *Lyl-1*^{LacZ/LacZ} newborns.

I. The expression of genes enriched in microglia and/or essential for their function are deregulated in *Lyl-*

1^{LacZ/LacZ} MΦ-progenitors at E9. Relative expression levels (read counts) in WT and *Lyl-1*^{lacZ/lacZ} MΦ-

progenitors from E9-YS (Unpaired, two-tailed *t*-Test).

Figure 1

A

%MΦP in E8 Ord1-YS

- Wild type
- *Lyl-1*$^{WT/LacZ}$
- *Lyl-1*$^{LacZ/LacZ}$

P = 0.037
P = 0.40

%GMP and EMP in OrgD1-YS

GMP   EMP

B

Kit$^+$CD45$^+$CD11b$^+$ gate

E9-YS (18-20S)	E9.5-YS (20-25S)	E10 (28-32S) YS

CD11b$^+$ F4/80$^+$ gate

E10 (28-32S) YS MΦ

Wild type: 2.41% 2.16% 2.28% 5.59%

Lyl-1$^{LacZ/LacZ}$: 75.70% 69.00% 54.7% 38.90%

Kit — FDG

CD11b — FDG

C

Wild type   *Lyl-1*$^{WT/LacZ}$

CD11b / CD31
CD11b+CD31+-1 0.323
CD11b+CD31+ 0.119
CD11b+CD31+-1 0.328
CD11b+CD31+ 0.543

CD45 / FDG

% Max / FDG
FDG+-1 1.69

CD31 / CD11b

D

- E9 WT YS
- E10 *Lyl-1*$^{WT/LacZ}$ YS FDG+
- E10 *Lyl-1*$^{WT/LacZ}$ YS FDG-
- E12 WT FL LSK

Myb relative expression levels

p=0.0023
n.s.
p=0.029
FDG+ FDG-
p=0.427

E

Lyl-1$^{WT/LacZ}$

WT   FDG+ FDG- WT

- GMP
- GP
- MΦP

% Myeloid colonies

E9   E10

Figure 2

Figure 3

D

E9 *Lyl-1*^{LacZ/LacZ} vs E9 WT

Figure 4

Figure 5

Figure 6