Novel mechanisms of type IV pilus regulation in *Acinetobacter baylyi*

Courtney K. Ellison[1,2*], Triana N. Dalia[3], Joshua W. Shaevitz[1*], Zemer Gitai[2*], and Ankur B. Dalia[3*]

1. Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ
2. Department of Molecular Biology, Princeton University, Princeton, NJ
3. Department of Biology, Indiana University, Bloomington, IN

*Correspondence to: c.ellison@princeton.edu, shaevitz@princeton.edu, zgitai@princeton.edu, ankdalia@indiana.edu

Bacteria employ extracellular appendages called type IV pili (T4P) to interact with their environment. T4P are essential for diverse microbial behaviors including DNA uptake, surface sensing, virulence, protein secretion, and twitching motility[1]. While T4P have been studied extensively, our understanding of these nanomachines largely comes from work on a few model species. Here, we develop *Acinetobacter baylyi* as a new model organism to study T4P and uncover several unreported mechanisms of T4P regulation. First, using recently-developed T4P-labeling methods[2,3], we demonstrate that *A. baylyi* T4P are synthesized on one side of the cell body along the long axis of the cell, and we uncover that this pattern is dependent on components of a conserved chemosensory pathway. Second, we overturn the current dogma that T4P extension occurs through the action of a single, highly conserved ATP-hydrolyzing motor (ATPase) called PilB by showing that T4P synthesis in *A. baylyi* is dependent on two partially redundant and phylogenetically distinct motors, PilB and PilB2. Third, we uncover a small protein inhibitor of T4P synthesis that specifically inhibits PilB but not PilB2 activity. Together, these results demonstrate novel mechanisms of T4P regulation, which have broad implications for the unexplored diversity of T4P biology in microbial species.

T4P are thin, proteinaceous appendages that are broadly distributed throughout Bacteria and Archaea[4]. T4P are composed primarily of major pilin protein subunits that are polymerized or depolymerized through the activity of ATPases to mediate fiber extension and retraction, respectively[5]. T4P are subdivided into subcategories (generally T4aP, T4bP, and T4cP) based on protein homology and pilus function, with T4aP being the best-characterized. In the T4aP systems found in Gram-negative organisms, polymerization and depolymerization of pilins occurs through interactions of the extension ATPase PilB or retraction ATPase PilT with the

integral inner membrane platform protein PilC (Supplemental figure 1a). The growing fiber

spans an alignment complex from the inner membrane through the periplasm composed of

PilNOP to exit through the PilQ outer membrane secretin pore. Dynamic cycles of T4P

extension and retraction are critical for the diverse processes that these structures mediate

including twitching motility[6], surface sensing[2,7], virulence[8,9], and DNA uptake[10,11].

Acinetobacter species like A. baumannii and A. nosocomialis have recently become an

urgent medical threat due to their prevalence in hospital-acquired infections and their capacity to

acquire antibiotic resistance genes, a process that is achieved in part by T4aP-mediated DNA

uptake[12,13]. Acinetobacter baylyi is the most naturally transformable species reported to date[14],

with up to 50% of cells undergoing natural transformation in ideal settings. To study A. baylyi

T4P, we applied a recently-developed labeling method[2,3] by targeting the major pilin, ComP, for

cysteine substitution and subsequent labeling with thiol-reactive maleimide dyes. Maleimide-

labeling of the functional $comP^{T129C}$ strain (Supplemental figure 1b) revealed an unusual pattern

of pilus synthesis, with T4P appearing only on one side of the cell body in a line along the long

axis of the cell (Figure 1a). Deletion of the retraction ATPase, PilT, resulted in hyperpiliated cells

exhibiting the same localization of T4P along the long axis of the cell. Cells with T4P facing the

microscope objective show T4P localize linearly along the side of the cell where they are

synthesized (Figure 1a). T4P localization has previously been shown to be polar or peritrichous

in other species, and to our knowledge, a long-axis localization pattern has not previously been

reported for any bacterial surface appendage or nanomachine.

Studies on the polarly localized T4P in Pseudomonas aeruginosa, a species closely

related to the Acinetobacter clade and one of the major model systems used to study T4P[1],

suggest that the chemosensory Pil-Chp system may play a role in controlling which pole of the

cell extends T4P[15]. The Pil-Chp system shares homology to flagellar chemotaxis systems, and

in P. aeruginosa, the methyl-accepting chemotaxis protein homologue, PilJ, is thought to signal

to the CheA homologue, ChpA. The CheW homologue, PilI is thought to act as a scaffold for Pil-

Chp signaling, leading to phosphoryl transfer from ChpA to the CheY homologues, PilG and

PilH to affect T4P synthesis. To determine whether the Pil-Chp system plays a role in controlling

where T4P are synthesized in A. baylyi, we made in-frame deletions of all Pil-Chp component

genes (Figure 1b). Deletion of pilG, pilI, and chpA all abrogated T4P localization and resulted in

the production of peritrichous T4P. However, deletions of pilH and, more surprisingly, pilJ did

not affect T4P patterning, suggesting that another inner membrane component may initiate

ChpA phosphoryl transfer in A. baylyi (Figure 1c, d). In P. aeruginosa, the scaffold protein FimV

interacts with the accessory protein FimL to recruit PilG to T4P machinery sites[16]. Although A.

baylyi lacks a FimV homologue, deletion of *fimL* also resulted in peritrichous pilus localization

(Figure 1c, d). In *P. aeruginosa*, T4P machinery is localized to both poles, but PilG is thought to

control which cell pole extends T4P[17,18]. However, an mCherry fusion to the outer membrane

secretin, PilQ, in *A. baylyi* demonstrated the same long-axis localization pattern as labeled T4P,

revealing a unique function for the Pil-Chp pathway in *A. baylyi* in regulating positioning of T4P

machinery and not simply regulating pilus dynamic activity (Supplemental figure 2).

Because PilG shares homology to two-component system response regulators, which

drive physiological changes downstream of an input signal, we hypothesized that PilG may

control the long-axis positioning of T4P machines. Consistent with this, we found that PilG-

mRuby3 localizes to the base of extended T4P (Supplemental figure 3a). Deletions of *pilI* and

chpA resulted in aberrant PilG-mRuby3 localization, mirroring T4P patterning in those mutants

(Figure 1e). The *fimL* mutant displayed cytoplasmic PilG signal, consistent with FimL playing a

role in PilG localization (Figure 1e). Notably, PilG-mRuby3 fluorescence intensity was

diminished in mutants with aberrant localization, suggesting a role for the Pil-Chp pathway in

controlling PilG levels (Supplemental figure 3b). The percent of cells making T4P in Pil-Chp

mutants was largely unchanged, suggesting an additional divergence from *P. aeruginosa* where

the Pil-Chp pathway also controls T4P production[19] (Supplemental figure 4a).

The Pil-Chp system is homologous to flagellar chemotaxis systems, so we suspected

that an external signal may feed into the Pil-Chp pathway to control T4P localization. Outer

membrane proteins in Gram-negative bacteria interface with the extracellular environment, and

we hypothesized that an outer membrane protein could influence the Pil-Chp system to regulate

T4P localization. To test this hypothesis, we deleted several outer membrane proteins

containing OmpA-like domains, and found that deletion of the major *ompA* gene, but not others,

resulted in peritrichous T4P localization (Figure 1c, d, Supplemental figure 5). Consistent with

an external signal feeding into the Pil-Chp pathway through OmpA, PilG-mRuby3 localization in

the *ompA* mutant displayed peritrichous localization similar to localization of labeled T4P (Figure

1d, e).

Because *A. baylyi* T4P participate in DNA uptake, we tested whether Pil-Chp mutants

with peritrichous localization were impacted for natural transformation. Surprisingly,

transformation was unaffected in mutants lacking Pil-Chp components (Supplemental figure 4b).

However, incubation of cells with fluorescently-labeled DNA resulted in co-localization of DNA

with T4P, which is consistent with the essential role of T4P-DNA binding during natural

transformation in *Vibrio cholerae*[11] (Figure 2a, Supplemental figure 4b). We reasoned that

natural transformation could be used to screen for other factors that regulate T4P in *A. baylyi*.

To that end, we performed a high throughput transposon-sequencing screen (Tn-seq) to identify

genes required for natural transformation. In this screen, loss of T4P-related genes resulted in

negative selection, indicating that they were critical for natural transformation as expected

(Figure 2b). To validate our Tn-seq results, we made in-frame deletions of representative genes

from each T4P-encoding operon, as well as genes known to be essential for natural

transformation that act downstream of T4P. Mutations in the T4P platform protein gene *pilC*, the

pilus regulatory gene *pilY1*, the outer membrane secretin gene *pilQ* and the retraction ATPase

gene *pilT* all showed a marked reduction in natural transformation, with most mutants exhibiting

transformation rates below our limit of detection (Figure 2c). T4P-labeling of *pilC*, *pilY1*, and *pilQ*

mutants revealed no visible T4P fibers (Figure 2d). Mutations in genes that act downstream of

T4P, including the periplasmic DNA-binding protein gene *comEA*, the inner membrane DNA-

transporter protein gene *comEC*, and the DNA-recombination helicase gene *comM* likewise

resulted in a reduction in transformation frequency, however, these strains still produced T4P

fibers as expected (Figure 2c, d).

117   We next sought to determine whether any of the previously uncharacterized genes from

our Tn-seq screen reduced natural transformation in *A. baylyi* by affecting T4P activity. Tn-seq

revealed that mutations in *pilB*, the canonical extension ATPase gene that is typically co-

transcribed with *pilC* and the pre-pilin peptidase, *pilD*, resulted in reduced natural transformation

(Figure 2b). However, we also found an additional *pilB* homologue, which we named *pilB2*, that

likewise exhibited lower rates of transformation (Figure 2b). Surprisingly, deletion of the

canonical *pilB* did not ablate transformation as would be expected based on homology to other

T4P[10] (Figure 3a). Mutation of *pilB2* only partially reduced transformation; however, natural

transformation was undetectable in the *pilB pilB2* double mutant (Figure 3a). T4P labeling in

pilB, *pilB2*, and *pilB pilB2* mutants revealed that all three mutants were defective in T4P

synthesis while *pilT* was intact. Strikingly, a *pilT* deletion, which prevents T4P retraction[20],

revealed that *pilB* mutant populations had similar numbers of cells producing T4P while *pilB2*

mutants were highly defective in T4P synthesis (Figure 3b, c). The *pilB pilB2 pilT* triple mutant

produced no detectable T4P fibers (Figure 3b, c). Together, these results indicate that both PilB

and PilB2 are essential for efficient T4P extension in *A. baylyi*, with PilB2 playing the dominant

role. Phylogenetic analysis of PilB2 homologues found among Gammaproteobacteria[4,21], which

include members of the PilT/VirB11 family of secretion ATPases, revealed that PilB2 clusters

with a group of proteins that are phylogenetically distinct from the canonical PilB ATPase

(Figure 3d, Supplemental figure 6). PilB and PilB2 proteins are as divergent from one another

as PilB and the type II secretion system proteins XcpR/GspE, or PilB and the MshE motors that

drive mannose-sensitive haemagglutinin (MSHA) pilus synthesis, suggesting that PilB2 evolved

as a functionally divergent class of proteins that is distinct from canonical PilB extension motors.

These results suggest that proteins that cluster with PilB2 may play similar functions in other

bacterial species and that multiple extension ATPases may be a common feature of diverse

T4P.

In addition to revealing the importance of PilB2 in T4P synthesis, our Tn-seq results also

uncovered an uncharacterized transcriptional regulator belonging to the XRE-family of

transcriptional repressors that is critical for natural transformation (Figure 2b, Figure 4a).

Deletion of this regulator, named *cpiR* for competence pilus inhibition repressor, resulted in a

100-fold reduction in transformation (Figure 4b). We hypothesized that CpiR repressed a factor

that inhibits natural transformation. Transcriptional repressors are often transcribed immediately

adjacent to the genes they repress. We thus deleted the upstream gene, named *cpiA* for

competence pilus inhibition actuator (Figure 4a), and found that transformation frequency was

restored in the *cpiR cpiA* double mutant background (Figure 4b). Expression of *cpiA* under the

control of an IPTG-inducible promoter (P_{tac}) was sufficient to reduce transformation frequency

even in the presence of *cpiR*, further suggesting that CpiR represses *cpiA* transcription.

Furthermore, we found that CpiR was sufficient to repress the *cpiA* promoter (using a P_{cpiA}-GFP

reporter) in the heterologous host *Vibrio cholerae*, which suggests that CpiR is a direct

repressor of P_{cpiA} (Figure 4c). Finally, we found that CpiA protein is only produced in a *cpiR*

mutant (Figure 4d). These results demonstrate that under lab conditions, CpiR represses *cpiA*

transcription to allow for high rates of natural transformation; however, in the absence of CpiR,

CpiA is produced and natural transformation is inhibited ~100-fold.

We next sought to determine the mechanism of transformation inhibition by CpiA.

Labeling of T4P in a *cpiR* mutant (when CpiA is expressed) revealed that cells were deficient in

T4P synthesis (Figure 4e). This defect was dependent on *cpiA* because piliation was restored in

the *cpiR cpiA* double mutant (Figure 4e). Deletion of *pilT* restored T4P synthesis in the *cpiR*

mutant (i.e. the *cpiR pilT* double mutant) (Figure 4e, f). This phenotype was reminiscent of the

restoration of piliation observed in the *pilB pilT* double mutant, and we hypothesized that CpiA

may regulate natural transformation by inhibiting PilB.

To test this hypothesis, we made *pilB cpiR* and *pilB2 cpiR* double mutants. T4P

synthesis was unaffected in *pilB cpiR pilT* (where PilB2 is the sole extension ATPase present),

while no T4P were detected in *pilB2 cpiR pilT* (where PilB is the sole extension ATPase

present) (Figure 4e, f). The defect in T4P synthesis in the *pilB2 cpiR pilT* strain was restored

when *cpiA* was deleted in this background, demonstrating that CpiA acts as a PilB inhibitor to

control natural transformation (Figure 4e, f). Transformation frequency assays corroborated

these results by demonstrating that deletion of *cpiR* only inhibited transformation in the *pilB2*

mutant background (where PilB is the sole extension ATPase) and not the *pilB* mutant

background (where PilB2 is the sole extension ATPase) (Figure 4g).

Environmental conditions play an integral role in regulating how bacteria respond to and

interact with their surroundings. The novel mechanisms of T4P regulation identified here are

likely a reflection of how environmental conditions influence bacterial physiology. Many bacteria

use chemotaxis systems to sense and respond to changes in environmental signals[22]. By

homology, an external cue likely controls T4P localization through the Pil-Chp pathway. Indeed,

the dependence of T4P localization on the outer membrane protein OmpA supports this idea, as

OmpA is a juncture for intracellular components and extracellular media. These data

demonstrate a novel role for OmpA in addition to its role in cell envelope integrity and virulence

as shown in other *Acinetobacter* species[23], highlighting its versatility in adapting to variable

environmental contexts. Future work will involve identifying the external signals that feed into

the Pil-Chp pathway through OmpA to control localization as well as the molecular mechanism

underlying T4P localization and the role it plays in the physiology of *Acinetobacter* species.

The acquisition of the additional extension motor PilB2 and the PilB-specific inhibitory

protein CpiA also likely resulted from a need to modulate T4P synthesis under different

environmental conditions. Bacteria generally limit T4P synthesis to environments where they

provide a selective advantage. For example, in *V. cholerae*, the production of T4a competence

pili requires chitin and quorum sensing[24], which are environmental conditions where cells are

likely to encounter other bacterial cells for horizontal gene transfer; while the T4b toxin-

coregulated (TCP) pili required for intestinal colonization are produced in response to host-

specific cues[25]. In *A. baylyi*, both PilB and PilB2 are required for T4P extension and,

consequently, efficient natural transformation. While CpiA is not expressed under laboratory

conditions (due to CpiR repression), it is possible that different environmental conditions

derepress *cpiA* to decrease pilus activity.

These results establish *A. baylyi* as a new model system to study T4P. By uncovering

several unknown mechanisms of T4P regulation, these studies underscore the unexplored

versatility and diversity of these ubiquitous systems and further highlight the need for new model

species to study T4P biology.

Methods

<u>Bacterial strains and culture conditions</u>

Acinetobacter baylyi strain ADP1 was used throughout this study. For a list of strains

used throughout, see Supplementary Table 1. *A. baylyi* cultures were grown at 30 °C in Miller

lysogeny broth (LB) medium and on agar supplemented with kanamycin (50 μg/mL),

spectinomycin (60 μg/mL), gentamycin (30 μg/mL), and/or chloramphenicol (30 μg/mL),

rifampicin (30 μg/mL), and/or streptomycin (10 μg/mL) as appropriate.

<u>Construction of mutant strains</u>

Mutants in *A. baylyi* were made using natural transformation as described previously.

Briefly, mutant constructs were made by splicing-by-overlap (SOE) PCR to stich 1) ~3 kb of the

homologous region upstream of the gene of interest, 2) the mutation where appropriate (for

deletion by allelic replacement with an antibiotic resistance cassette, or the fusion protein), and

3) the downstream region of homology. For a list of primers used to generate mutants in this

study, see Supplementary Table 2. The upstream region was amplified using F1 + R1 primers,

and the downstream region was amplified using F2 + R2 primers. All antibiotic resistance (AbR)

cassettes were amplified with ABD123 (ATTCCGGGGATCCGTCGAC) and ABD124

(TGTAGGCTGGAGCTGCTTC). Fusion proteins were amplified using the primers indicated in

Supplementary Table 2. In-frame deletions were constructed using F1 + R1 primer pairs to

amplify the upstream region and F2 + R2 primer pairs to amplify the downstream region with

~20 bp homology to the remaining region of the downstream region built into the R1 primer and

~20 bp homology to the upstream region built into the F2 primer. SOE PCR reactions were

performed using a mixture of the upstream and downstream regions, and middle region where

appropriate using F1 + R2 primers. SOE PCR products were added with 50 μl of overnight-

grown culture to 450 μl of LB in 2 ml round-bottom microcentrifuge tubes (USA Scientific) and

grown at 30 °C rotating on a roller drum for 1-3 hours. For AbR-constructs, transformants were

diluted out to 10^{-6} and dribble plated on LB and LB + AbR. For in-frame deletions and protein

fusion constructs, after the three-hour incubation, cells were incubated with 10 μl of DNaseI

(New England Biolabs) for 5 min at room temperature before 450 μl of LB was added.

Transformations were grown for an additional hour before cells were diluted and 100 μl of 10^{-6}

dilution was plated on LB plates. In-frame deletions were confirmed by PCR using primers ~150

bp up and downstream of the introduced mutation, and fusions were confirmed by sequencing.

Mutants in *V. cholerae* were made by chitin-induced natural transformation exactly as

previously described[26,27]. SOE products were generated exactly as described above. The

chromosomally integrated P*BAD* ectopic expression construct is described in detail in ref [28].

 Natural transformation assays

 Assays were performed exactly as previously described[29]. Briefly, strains were grown

 overnight in LB broth at 30°C rolling. Then, ~10^8 cells were subcultured into fresh LB medium

 and 100 ng of tDNA was added. In this study, a ΔACIAD1551::Spec^R PCR product (with 3kb

 arms of homology) was used as the tDNA. Reactions were incubated with end-over-end rotation

 at 30°C for 5 hours and then plated for quantitative culture on spectinomycin plates (to quantify

 transformants) and on plain LB plates (to quantify total viable counts). Data are reported as the

 transformation frequency, which is defined as the (CFU/mL of transformants) / (CFU/mL of total

 viable counts).

 Tn-seq Analysis

 Transposon mutant libraries of *A. baylyi* were generated by electroporation of cells with

 pDL1093, a vector that allows for Kan^R mini-Tn10 mutagenesis[30,31], and plating on kanamycin

 plates. Approximately 100,000 Tn mutants were scraped off of plates and pooled to generate

 the input transposon mutant library. This mutant library was then subjected to natural

 transformation in three separate replicates exactly as described above using three different

 sources of tDNA: a ΔACIAD1551::Spec^R PCR product, an RpsL K43R (Sm^R) PCR product, or

 an RpoB PCR product from a spontaneous rifampicin resistant mutant. Following incubation

 with tDNA, reactions were split and outgrown in selective medium overnight (i.e. with the

 appropriate antibiotic added to select for transformants = "transformant pool") or grown

 overnight in nonselective medium (i.e. in plain LB medium = "total pool"). Sequencing libraries of

 the transposon-genomic junctions were generated for the Illumina platform using HTM-PCR

 exactly as previously described[31,32]. Sequencing data were mapped to the *V. cholerae* N16961

 genome[33] and the relative abundance of Tn insertions in all samples was determined on the

 Galaxy platform[34]. Comparative analyses for Tn-seq were performed treating the "transformant

 pool" as the output and the "total pool" as the input to determine genes that were over and

 underrepresented following selection for transformants. Gene fitness was only assessed if a

 gene contained at least 1 transposon insertion in at least two out of the three replicates of the

 "total pool" samples. Also, the normalized abundance of insertions within a gene had to be

 greater than 0.01 in the "total pool" when averaged across all three replicates (where 1 is the

 expected normalized abundance for a "neutral" gene). These cutoffs allowed us to assess the

 phenotype of ~83% of the genes (2747/3310) in the *A. baylyi* ADP1 genome. For visualization,

 the relative abundance of Tn insertions within each gene is plotted from the "transformant pool"

 relative to the "total pool".

<u>Pilin labeling, imaging, and quantification</u>

Pilin labeling was performed as described previously with some changes[2,3]. Briefly, 100 µl of overnight-grown cultures was added to 900 µl of LB in 1.5 ml microcentrifuge tube, and cells were grown at 30 °C rotating on a roller drum for 70 min. Cells were then centrifuged at 18,000 x g for 1 min and then resuspended in 50 µl of LB before labeling with 25 µg/ml of AlexaFluor488 C_5-maleimide (AF488-mal) (ThermoFisher) for 15 min at room temperature. Labeled cells were centrifuged, washed once with 100 µl of LB without disrupting the pellet, and resuspended in 5-20 µl LB. Cell bodies were imaged using phase contrast microscopy while labeled pili were imaged using fluorescence microscopy on a Nikon Ti-2 microscope using a Plan Apo 100X objective, an FITC filter cube for pili, an mCherry filter cube for mRuby3 and mCherry fusions, a Hamamatsu ORCAFlash4.0 camera, and Nikon NIS Elements Imaging Software. Cell numbers and the percent of cells making pili were quantified manually using ImageJ[35]. Cells were considered to have "mislocalized pili" if at least two pili were on opposing sides of the cell. All imaging was performed under 1% UltraPure agarose (Invitrogen) pads made with phosphate-buffered saline (PBS) solution.

<u>Western blotting</u>

Western blot analysis was performed exactly as previously described[36]. Briefly, cells from overnight cultures were concentrated to an OD_{600} of 20. Then, cells were lysed on a FastPrep-24 classic instrument at 4°C, and then clarified by centrifugation. Lysates were then mixed with an equal volume of 2× SDS-PAGE sample buffer (220 mM Tris, pH 6.8, 25% glycerol, 1.2% SDS, 0.02% bromophenol blue, and 5% β-mercaptoethanol) and boiled. Proteins were separated on a 15% SDS polyacrylamide gel by SDS electrophoresis, electrophoretically transferred to a polyvinylidene difluoride (PVDF) membrane, and probed with rabbit polyclonal α-FLAG antibodies (Sigma Aldrich) or mouse monoclonal α-RpoA (BioLegend) primary antibodies. Blots were then incubated with α-rabbit or α-mouse horseradish peroxidase (HRP)-conjugated secondary antibodies, developed using Pierce ECL Western blotting substrate (Thermo Fisher), and imaged on a Protein Simple FluorChem E instrument.

<u>Measuring GFP fluorescence in reporter strains</u>

V. cholerae reporter strains harboring P_{cpiA}-GFP were grown to late log in LB medium with or without 0.2% arabinose as indicated. Cells were then washed once in instant ocean medium (7g/L; Aquarium Systems) and transferred to a 96-well plate. Fluorescence was then

determined on a Biotek H1M plate reader with monochromater set to 500 nm for excitation and

540 nm for emission.

<u>Fluorescent DNA binding/uptake</u>

A ~7 kb PCR product was fluorescently labeled as described previously[11] using the Cy3

LabelIT kit (Mirus Biosciences) as per manufacturer recommendations. For the parent strain, 1

µl (100 ng) of Cy3-DNA was added to 900 µl of LB in a 1.5 ml microcentrifuge tube along with

100 µl of overnight-grown cultures, and cells were grown at 30°C rotating on a roller drum for 70

min. Pili were then labeled as described above.

To visualize DNA-binding in the *pilT* mutant, cells were grown as described above, but 1

µl (100 ng) of Cy3-DNA was added to cells along with AF488-mal and incubated for 25 min

before washing. Cells were imaged using the same microscopy setup described above, using a

dsRed filter cube to image Cy3-DNA.

<u>Phylogenetic analysis</u>

PilB2 homologues were identified by BLAST using 43 manually selected

Gammaproteobacteria genomes[37] through Integrated Microbial Genomes and Microbiomes

online resources (IMG)[38] resulting in 209 protein sequences. The FtsK protein from *A. baylyi*

ADP1 was added and used as an outgroup. Through the NGPhylogeny.fr server[39], sequences

were aligned using the default parameters of MAFFT[40], and phylogenetic tree construction was

performed on the aligned sequences using the default parameters of FastTree software[41,42] set

to perform 100 bootstraps[43]. The resulting tree was visualized using the Interactive Tree of Life

(iTOL) visualization software[44], from which trees were exported for publication.

Acknowledgements: We would like to thank E. Geisinger for *A. baylyi* ADP1 ATCC33305

wildtype strain, and we would like to thank K. Hummels and B. Bratton for helpful suggestions

on phylogenetic analysis. CKE is a Damon Runyon Fellow supported by the Damon Runyon

Cancer Research Foundation (DRG-2385-20). This work was supported in part by the National

Science Foundation, through the Center for the Physics of Biological Function (PHY-1734030).

This work was supported by grant R35GM128674 from the National Institutes of Health

awarded to ABD and the National Institutes of Health Pioneer Award 1DP1AI124669-01

awarded to ZG.

Author contributions: CKE and ABD designed and coordinated the overall study. CKE, TND,

and ABD performed the experiments. All authors analyzed and interpreted data. CKE and ABD

wrote the manuscript with help from ZG and JWS.

Competing interests: The authors declare no competing interests.

Data availability statement: The data that support the findings of this study are available from

the corresponding authors upon request.

References

1. Burrows, L. L. Pseudomonas aeruginosa twitching motility: type IV pili in action. *Annu Rev Microbiol* **66**, 493–520 (2012).

2. Ellison, C. K. C. K. *et al.* Obstruction of pilus retraction stimulates bacterial surface sensing. *Science.* **358**, 535–538 (2017).

3. Ellison, C. K., Dalia, T. N., Dalia, A. B. & Brun, Y. V. Real-time microscopy and physical perturbation of bacterial pili using maleimide-conjugated molecules. *Nat. Protoc.* **14**, (2019).

4. Denise, R., Abby, S. S. & Rocha, E. P. C. Diversification of the type IV filament superfamily into machines for adhesion, protein secretion, DNA uptake, and motility. *PLOS Biol.* **17**, e3000390 (2019).

5. Craig, L., Forest, K. T. & Maier, B. Type IV pili: dynamics, biophysics and functional consequences. *Nat. Rev. Microbiol.* 1 (2019). doi:10.1038/s41579-019-0195-4

6. Merz, A. J., So, M. & Sheetz, M. P. Pilus retraction powers bacterial twitching motility. *Nature* **407**, 98–102 (2000).

7. Snyder, R. A. *et al.* Surface sensing stimulates cellular differentiation in Caulobacter crescentus. *Proc. Natl. Acad. Sci.* (2020). doi:10.1073/PNAS.1920291117

8. Siryaporn, A. *et al.* Surface attachment induces Pseudomonas aeruginosa virulence. *Proc. Natl. Acad. Sci.* **111**, 16860–16865 (2014).

9. Persat, A., Inclan, Y. F., Engel, J. N., Stone, H. A. & Gitai, Z. Type IV pili mechanochemically regulate virulence factors in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci.* **112**, 7563–7568 (2015).

10. Seitz, P. & Blokesch, M. DNA-uptake machinery of naturally competent Vibrio cholerae. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 17987–92 (2013).

11. Ellison, C. K. *et al.* Retraction of DNA-bound type IV competence pili initiates DNA uptake during natural transformation in Vibrio cholerae. *Nat. Microbiol.* **3**, 773–780 (2018).

12. Harding, C. M. *et al.* Acinetobacter baumannii strain M2 produces type IV Pili which play a role in natural transformation and twitching motility but not surface-associated motility. *MBio* **4**, (2013).

13. Godeux, A. S. *et al.* Fluorescence-based detection of natural transformation in drug-resistant acinetobacter baumannii. *J. Bacteriol.* **200**, (2018).

14. Metzgar, D. *et al.* Acinetobacter sp. ADP1: An ideal model organism for genetic analysis and genome engineering. *Nucleic Acids Res.* **32**, 5780–5790 (2004).

15. Jain, R., Sliusarenko, O. & Kazmierczak, B. I. Interaction of the cyclic-di-GMP binding protein FimX and the Type 4 pilus assembly ATPase promotes pilus assembly. *PLOS Pathog.* **13**, e1006594 (2017).

16. Inclan, Y. F. *et al.* A scaffold protein connects type IV pili with the Chp chemosensory system to mediate activation of virulence signaling in Pseudomonas aeruginosa. *Mol. Microbiol.* **101**, 590–605 (2016).

17. Bertrand, J. J., West, J. T. & Engel, J. N. Genetic analysis of the regulation of type IV pilus function by the Chp chemosensory system of Pseudomonas aeruginosa. *J. Bacteriol.* **192**, 994–1010 (2010).

18. Carter, T. *et al.* The type IVa pilus machinery is recruited to sites of future cell division. *MBio* **8**, (2017).

19. Buensuceso, R. N. C. *et al.* Cyclic AMP-independent control of twitching motility in Pseudomonas aeruginosa. *J. Bacteriol.* **199**, (2017).

20. Chlebek, J. L. *et al.* PilT and PilU are homohexameric ATPases that coordinate to retract type IVa pili. *PLOS Genet.* **15**, e1008448 (2019).

21. McCallum, M., Tammam, S., Khan, A., Burrows, L. L. & Howell, P. L. The molecular mechanism of the type IVa pilus motors. *Nat. Commun.* **8**, 15091 (2017).

22.   Wadhams, G. H. & Armitage, J. P. Making sense of it all: Bacterial chemotaxis. *Nature*
Reviews Molecular Cell Biology **5**, 1024–1037 (2004).

23.   Gaddy, J. A., Tomaras, A. P. & Actis, L. A. The Acinetobacter baumannii 19606 OmpA
protein plays a role in biofilm formation on abiotic surfaces and in the interaction of this
pathogen with eukaryotic cells. *Infect. Immun.* **77**, 3150–3160 (2009).

24.   Meibom, K. L., Blokesch, M., Dolganov, N. A., Wu, C.-Y. & Schoolnik, G. K. Chitin
Induces Natural Competence in Vibrio cholerae. *Science (80-.).* **310**, 1824–1827 (2005).

25.   Schuhmacher, D. A. & Klose, K. E. Environmental signals modulate ToxT-dependent
virulence factor expression in Vibrio cholerae. *J. Bacteriol.* **181**, 1508–1514 (1999).

26.   Dalia, A. B., McDonough, E. & Camilli, A. Multiplex genome editing by natural
transformation. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 8937–42 (2014).

27.   Dalia, A. B. Natural Cotransformation and Multiplex Genome Editing by Natural
Transformation (MuGENT) of Vibrio cholerae. in *Methods in molecular biology (Clifton,*
N.J.) **1839**, 53–64 (2018).

28.   Dalia, T. N., Chlebek, J. L. & Dalia, A. B. A modular chromosomally integrated toolkit for
ectopic gene expression in Vibrio cholerae. *bioRxiv* 2020.03.02.973776 (2020).
doi:10.1101/2020.03.02.973776

29.   Nero, T. M. *et al.* ComM is a hexameric helicase that promotes branch migration during
natural transformation in diverse Gram-negative species. *Nucleic Acids Res.* **46**, 6099–
6111 (2018).

30.   Shull, L. M. & Camilli, A. Transposon sequencing of vibrio cholerae in the infant rabbit
model of cholera. in *Methods in Molecular Biology* **1839**, 103–116 (Humana Press Inc.,
2018).

31.   Mcdonough, E., Lazinski, D. W. & Camilli, A. Identification of in vivo regulators of the
Vibrio choleraexds gene using a high-throughput genetic selection. *Mol. Microbiol.* **92**,
302–315 (2014).

32.   Lazinski, D. W. & Camilli, A. Homopolymer tail-mediated ligation PCR: A streamlined and
highly efficient method for DNA cloning and library construction. *Biotechniques* **54**, 25–34
(2013).

33.   Heidelberg, J. F. *et al.* DNA sequence of both chromosomes of the cholera pathogen
Vibrio cholerae. *Nature* **406**, 477–483 (2000).

34.   Afgan, E. *et al.* The Galaxy platform for accessible, reproducible and collaborative
biomedical analyses: 2016 update. *Nucleic Acids Res.* **44**, W3–W10 (2016).

35.   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of
image analysis. *Nat. Methods* **9**, 671–675 (2012).

36.   Klancher, C., Newman, J., Ball, A., van Kessel, J. & Dalia, A. Species-specific quorum
sensing represses the chitobiose utilization locus in Vibrio cholerae. *Appl. Environ.*
Microbiol. **86**, (2020).

37.   Mukherjee, S. *et al.* Genomes OnLine database (GOLD) v.7: Updates and new features.
Nucleic Acids Res. **47**, D649–D659 (2019).

38.   Chen, I. M. A. *et al.* IMG/M v.5.0: An integrated data management and comparative
analysis system for microbial genomes and microbiomes. *Nucleic Acids Res.* **47**, D666–
D677 (2019).

39.   Lemoine, F. *et al.* NGPhylogeny.fr: New generation phylogenetic services for non-
specialists. *Nucleic Acids Res.* **47**, W260–W265 (2019).

40.   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

41.   Price, M. N., Dehal, P. S. & Arkin, A. P. Fasttree: Computing large minimum evolution
trees with profiles instead of a distance matrix. *Mol. Biol. Evol.* **26**, 1641–1650 (2009).

42.   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 - Approximately maximum-likelihood
trees for large alignments. *PLoS One* **5**, e9490 (2010).

43.   Lemoine, F. *et al.* Renewing Felsenstein's phylogenetic bootstrap in the era of big data.
Nature **556**, 452–456 (2018).
44.   Letunic, I. & Bork, P. Interactive Tree of Life (iTOL) v4: Recent updates and new
developments. *Nucleic Acids Res.* **47**, W256–W259 (2019).
456

Figure 1. *A. baylyi* T4aP are produced in a line along the long axis of the cell body in a Pil-Chp

dependent manner. (**a**) Representative images of indicated strains labeled with AF488-mal. (**b**)

Schematic of the *A. baylyi* components of the Pil-Chp system. Deletion of tan proteins alters

T4P localization, while deletion of the gray proteins does not. Numbers in parentheses are the

ACIAD gene designations for each component. OM, outer membrane; IM, inner membrane. (**c**)

Percent of piliated cells in *pilT* mutants with mislocalized T4P in indicated strains. Cells were

determined to have "mislocalized pili" if they had at least two pili on opposing sides of the long

axis of the cell body. Cells that only made one pilus were counted in the total number of piliated

cells but were not counted as "mislocalized." Each data point represents an independent,

biological replicate and bar graphs indicate the mean ± SD. For each biological replicate, a

minimum of 70 total cells were assessed. (**d**) Representative images of indicated strains labeled

with AF488-mal with background fluorescence subtracted. (**e**) Representative images of

indicated PilG-mRuby3 strains with background fluorescence subtracted and fluorescence

channel lookup tables adjusted to show PilG-mRuby3 localization. Scale bars, 2 μm.

Figure 2. Tn-seq reveals factors important for natural transformation and T4P synthesis. (**a**) Representative images of cells with AF488-mal labeled pili incubated with fluorescently-labeled DNA. (**b**) Visual representation of Tn-seq screen showing the relative abundance of each gene in the "total pool" of transposon mutants compared to the "transformant pool" recovered after natural transformation. Known T4P regulatory genes are outlined in red. *pilB2* is outlined in green, and *cpiR* is outlined in cyan. (**c**) Natural transformation assays of indicated strains. Each data point represents a biological replicate and bar graphs indicate the mean ± SD. LD, limit of detection. (**d**) Representative images of strains from natural transformation assays shown in **c** labeled with AF488-mal. Scale bars, 2 μm.

491
492

Figure 3. PilB2 is a phylogenetically distinct PilB homologue that is required for efficient T4P extension in *A. baylyi*. (**a**) Natural transformation assays of indicated strains. Each data point represents a biological replicate and bar graphs indicate the mean ± SD. LD, limit of detection. (**b**) Percent of piliated cells in *pilT* mutant populations of indicated strains. Each data point represents an independent, biological replicate and bar graphs indicate the mean ± SD. For each biological replicate, a minimum of 70 total cells were assessed. ND, no pili detected. (**c**) Representative images of indicated strains labeled with AF488-mal with background fluorescence subtracted. Scale bars, 2 μm. (**d**) A rooted phylogeny of PilB2 homologues found among Gammaproteobacteria. Branches are colored according to protein annotations in IMG, and nodes with boostrap values greater than or equal to 70% are indicated by black circles. Black stars are at the tips of branches representing *A. baylyi* proteins with indicated protein names.

Figure 4. *A. baylyi* T4P extension is regulated by the PilB-inhibitor protein, CpiA, whose
expression is controlled by the transcriptional repressor, CpiR. (**a**) Schematic *cpiRA* locus
organization. (**b**) Natural transformation assays of the indicated strains performed with or
without 100 μM IPTG added as indicated. Each data point represents a biological replicate and
bar graphs indicate the mean ± SD. (**c**) Fluorescence intensity of *V. cholerae* strains harboring a
P$_{cpiA}$-GFP reporter and a P$_{BAD}$-*cpiR* ectopic expression construct as indicated when grown with
or without 0.2% arabinose as indicated. Each data point represents a biological replicate and
bar graphs indicate the mean ± SD. (**d**) Western blot showing CpiA-FLAG production in the
indicated strains. RpoA was detected as a loading control. (**e**) Representative images of
indicated strains labeled with AF488-mal with background fluorescence subtracted. Scale bars,

524   2 µm. (**f**) Percent of piliated cells in *pilT* mutant populations of indicated strains. Each data point

525   represents an independent, biological replicate and bar graphs indicate the mean ± SD. For

526   each biological replicate, a minimum of 70 total cells were assessed. ND, no pili detected. (**g**)

527   Natural transformation assays of indicated strains. Each data point represents a biological

528   replicate and bar graphs indicate the mean ± SD. LD, limit of detection.

Supplementary Materials for

Novel mechanisms of type IV pilus regulation in *Acinetobacter baylyi*

Courtney K. Ellison[1,2*], Triana N. Dalia[3], Joshua W. Shaevitz[1*], Zemer Gitai[2*], and Ankur B. Dalia[3*]

1. Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ
2. Department of Molecular Biology, Princeton University, Princeton, NJ
3. Department of Biology, Indiana University, Bloomington, IN

*Correspondence to: c.ellison@princeton.edu, shaevitz@princeton.edu, zgitai@princeton.edu, ankdalia@indiana.edu

This file includes:

Supplemental figure 1. The strain for labeling T4P in *A. baylyi* is fully functional. (a) Schematic of assembled T4P components found in *A. baylyi*. Numbers in parentheses are ACIAD numbers associated with each component. OM, outer membrane; PG, peptidoglycan; IM, inner membrane. (b) Natural transformation assays of indicated strains. Each data point represents a biological replicate and bar graphs indicate the mean ± SD. The transformation frequency of the Δ*pilQ* strain was below the limit of detection, indicated by LD.

Supplemental figure 2. PilQ-mCherry localizes to the long-axis of the cell dependent on components of the Pil-Chp pathway. (a) Schematic of *pilQ-mCherry* gene locus. Attempts to make *pilQ-mCherry* by natural transformation without a selective marker were unsuccessful, and we thus added a gentamycin resistance cassette downstream of *pilQ*. Because *pilQ* is in an operon with the essential gene *aroK* immediately downstream, we added a constitutively active P_{tac} promoter to drive *aroK* expression. (b) Natural transformation assays of indicated strains. The *pilQ*-GmR control strain contains the same GmR-P_{tac} marker downstream of PilQ present in *pilQ-mCherry*, but PilQ is untagged in this background. Each data point represents a biological replicate and bar graphs indicate the mean ± SD. The transformation frequency of the Δ*pilQ* strain was below the limit of detection, indicated by LD. (c) Representative images of indicated PilQ-mCherry strains with background fluorescence subtracted. Scale bars, 2 μm.

Supplemental figure 3. PilG-mRuby3 localizes at the base of T4P, and its expression is reduced in Pil-Chp mutants. (a) Representative image of PilG-mRuby3 Δ*pilT* strain labeled with AF488-mal with background fluorescence subtracted. (b) Representative images of PilG-mRuby3 localization in indicated strains with background fluorescence subtracted. Images in (b) are identical to images shown in Figure 1e, except with altered fluorescence lookup tables (LUT). Normalized fluorescence images have the same fluorescence LUT across strains, while adjusted fluorescence images have LUTs adjusted for protein visibility to show PilG-mRuby3 localization patterns. Scale bars, 2 μm.

Supplemental figure 4. Pil-Chp mutants are unimpaired in T4P synthesis and are not required for natural transformation. (a) Percent of piliated cells in *pilT* mutant populations of indicated strains. Each data point represents an independent, biological replicate and bar graphs indicate the mean ± SD. For each biological replicate, a minimum of 70 total cells were assessed. (b) Natural transformation assays of indicated strains. Each data point represents a biological replicate and bar graphs indicate the mean ± SD. The transformation frequency of the Δ*comP* strain was below the limit of detection, indicated by LD.

Supplemental figure 5. Other OmpA domain-containing proteins do not affect pilus machinery localization. Representative images of indicated PilQ-mCherry strains with background fluorescence subtracted. Scale bars, 2 μm.

Supplemental figure 6. PilB2 clusters with a group of other GspE-like proteins that are distinct from PilB proteins. A rooted phylogeny of PilB2 homologues shown in Figure 3d with labels. Labels indicate IMG GeneID, annotated protein name, species, protein size. Nodes with boostrap values greater than or equal to 70% are indicated by black circles.

Supplementary table 1. Bacterial strains used in this study.

CE#	ZG#	TND/ SAD#	Strain name in manuscript	Genotype	Figure ('S' denotes supplemental figure)
CE1		SAD631	Wildtype (WT)	*Acinetobacter baylyi* ADP1	1A, S1B
CE100		SAD2537	Pil-cys or Parent	ADP1 *comP*^{T129C}	1A, 2A, 2C, 2D, 3A, 4B, 4D, S1B, S2B, S4B
CE329	ZG1709		Δ*pilT*	ADP1 Δ*pilT::spec*	1A
CE317		TND0779	Pil-cys Δ*pilT* or Parent	ADP1 *comP*^{T129C} Δ*pilT::spec*	1A, 1C, 2A, 2C, 3B, S4A
CE2	ZG1693		Δ*pilG*	ADP1 *comP*^{T129C} Δ*pilG*	S4B
CE3	ZG1694		Δ*pilH*	ADP1 *comP*^{T129C} Δ*pilH*	S4B
CE8	ZG1695		Δ*pilI*	ADP1 *comP*^{T129C} Δ*pilI*	S4B
CE9	ZG1696		Δ*pilJ*	ADP1 *comP*^{T129C} Δ*pilJ*	S4B
CE10	ZG1697		Δ*chpA*	ADP1 *comP*^{T129C} Δ*chpA*	S4B
CE122	ZG1698		Δ*ompA*	ADP1 *comP*^{T129C} Δ*ompA::kan*	S4B
CE233	ZG1699		Δ*fimL*	ADP1 *comP*^{T129C} Δ*fimL::kan*	S4B
		TND0114	Δ*comP*	ADP1 Δ*comP::kan*	3A, S4B
CE6	ZG1668		Δ*pilG* Δ*pilT*	ADP1 *comP*^{T129C} Δ*pilG* Δ*pilT::kan*	1C, 1D, S4A
CE7	ZG1669		Δ*pilH* Δ*pilT*	ADP1 *comP*^{T129C} Δ*pilH* Δ*pilT::kan*	1C, 1D, S4A
CE18	ZG1670		Δ*pilI* Δ*pilT*	ADP1 *comP*^{T129C} Δ*pilI* Δ*pilT::kan*	1C, 1D, S4A
CE19	ZG1671		Δ*pilJ* Δ*pilT*	ADP1 *comP*^{T129C} Δ*pilJ* Δ*pilT::kan*	1C, 1D, S4A
CE20	ZG1672		Δ*chpA* Δ*pilT*	ADP1 *comP*^{T129C} Δ*chpA* Δ*pilT::kan*	1C, 1D, S4A
CE312	ZG1675		Δ*ompA* Δ*pilT*	ADP1 *comP*^{T129C} Δ*ompA::kan* Δ*pilT::spec*	1C, 1D, S4A
CE315	ZG1676		Δ*fimL* Δ*pilT*	ADP1 *comP*^{T129C} Δ*fimL::kan* Δ*pilT::spec*	1C, 1D, S4A
CE256	ZG1681		PilQ-mCherry or Parent	ADP1 *comP*^{T129C} *pilQ::pilQ-mCherry, gent,* P$_{tac}$- *aroK*	S2B, S2C
CE331	ZG1682		PilQ-GmR	ADP1 *comP*^{T129C} *pilQ::pilQ, gent,* P$_{tac}$-*aroK*	S2B

CE320	ZG1700		Δ*pilG*	ADP1 *comP*^{T129C} *pilQ::pilQ-mCherry*, gent, P$_{tac}$- *aroK* Δ*pilG*	S2C
CE321	ZG1701		Δ*pilH*	ADP1 *comP*^{T129C} *pilQ::pilQ-mCherry*, gent, P$_{tac}$- *aroK* Δ*pilH*	S2C
CE322	ZG1702		Δ*pilI*	ADP1 *comP*^{T129C} *pilQ::pilQ-mCherry*, gent, P$_{tac}$- *aroK* Δ*pilI*	S2C
CE323	ZG1703		Δ*pilJ*	ADP1 *comP*^{T129C} *pilQ::pilQ-mCherry*, gent, P$_{tac}$- *aroK* Δ*pilJ*	S2C
CE324	ZG1704		Δ*chpA*	ADP1 *comP*^{T129C} *pilQ::pilQ-mCherry*, gent, P$_{tac}$- *aroK* Δ*chpA*	S2C
CE325	ZG1705		Δ*ompA*	ADP1 *comP*^{T129C} *pilQ::pilQ-mCherry*, gent, P$_{tac}$- *aroK* Δ*ompA::kan*	S2C
CE326	ZG1706		ΔACIAD0653	ADP1 *comP*^{T129C} *pilQ::pilQ-mCherry*, gent, P$_{tac}$- *aroK* Δ ACIAD0653::*kan*	S5
CE327	ZG1707		ΔACIAD0898	ADP1 *comP*^{T129C} *pilQ::pilQ-mCherry*, gent, P$_{tac}$- *aroK* Δ ACIAD0898::*kan*	S5
CE328	ZG1708		Δ*fimL*	ADP1 *comP*^{T129C} *pilQ::pilQ-mCherry*, gent, P$_{tac}$- *aroK* Δ*fimL::kan*	S2C
CE17	ZG1686		Parent	ADP1 *comP*^{T129C} *pilG::pilG-mRuby3*	1E, S3B
CE40	ZG1687		Δ*chpA*	ADP1 *comP*^{T129C} *pilG::pilG-mRuby3* Δ*chpA*	1E, S3B
CE127	ZG1688		Δ*ompA*	ADP1 *comP*^{T129C} *pilG::pilG-mRuby3* Δ*ompA::kan*	1E, S3B
CE257	ZG1689		Δ*fimL*	ADP1 *comP*^{T129C} *pilG::pilG-mRuby3* Δ*fimL::chlor*	1E, S3B
CE261	ZG1690		Δ*pilI*	ADP1 *comP*^{T129C} *pilG::pilG-mRuby3* Δ*pilI*	1E, S3B
CE341	ZG1691		Δ*pilH*	ADP1 *comP*^{T129C} *pilG::pilG-mRuby3* Δ*pilH*	1E, S3B

CE342	ZG1692		ΔpilJ	ADP1 comP^T129C pilG::pilG-mRuby3 ΔpilJ	1E, S3B
CE25	ZG1736		PilG-mRuby3 ΔpilT	ADP1 comP^T129C pilG::pilG-mRuby3 ΔpilT::kan	S3A
CE150	ZG1680		ΔpilC	ADP1 comP^T129C ΔpilC	2C, 2D
CE16	ZG1678		ΔpilY1	ADP1 comP^T129C ΔpilY1	2C, 2D
CE66	ZG1679		ΔpilQ	ADP1 comP^T129C ΔpilQ	2C, 2D, S1B, S2B
CE286		TND2544	ΔcomEA	ADP1 comP^T129C ΔcomEA::kan	2C, 2D
CE287		TND2546	ΔcomEC	ADP1 comP^T129C ΔcomEC::kan	2C, 2D
CE289		TND2556	ΔcomM	ADP1 comP^T129C ΔcomM::kan	2C, 2D
CE22	ZG1683		ΔpilB	ADP1 comP^T129C ΔpilB	3A, 3C, 4G
CE67	ZG1685		ΔpilB2	ADP1 comP^T129C ΔpilB2	3A, 3C, 4G
CE54	ZG1684		ΔpilB ΔpilB2	ADP1 comP^T129C ΔpilB ΔpilB2::kan	3A, 3C, 4G
CE310	ZG1673		ΔpilB ΔpilT	ADP1 comP^T129C ΔpilB ΔpilT::spec	3B, 3C, 4F
CE311	G1674		ΔpilB2 ΔpilT	ADP1 comP^T129C ΔpilB2 ΔpilT::spec	3B, 3C, 4F
CE316	ZG1677		ΔpilB ΔpilB2 ΔpilT	ADP1 comP^T129C ΔpilB ΔpilB2::kan ΔpilT::spec	3B, 3C
CE263		TND1201	ΔcpiR	ADP1 comP^T129C ΔcpiR::kan	4B, 4E
CE264		SAD2792	ΔcpiR ΔcpiA	ADP1 comP^T129C ΔcpiRA::kan	4B, 4E
		TND2260	P_tac-cpiA	ADP1 comP^T129C igACIAD0096-ACIAD0098::Cm^R-P_tac-cpiA	4B
		TND2093	P_cpiA-GFP Parent	V. cholerae E7946 Sm^R ΔlacZ::SpecR-P_cpiA-gfp	4C
		TND2126	P_cpiA-GFP P_BAD-cpiR	V. cholerae E7946 Sm^R ΔlacZ::SpecR-P_cpiA-gfp ΔVCA0692::Carb^R-P_BAD-cpiR	4C
		TND1335	cpiA-FLAG	ADP1 comP^T129C cpiA-3X FLAG	4D

		TND1284	*cpiA*-FLAG Δ*cpiR*	ADP1 *comP*^{T129C} *cpiA*-3X FLAG Δ*cpiR*::*kan*	4D
CE282		SAD2808	Δ*pilB* Δ*cpiR*	ADP1 *comP*^{T129C} Δ*pilB* Δ*cpiR*::*kan*	4E, 4G
CE283		SAD2806	Δ*pilB2* Δ*cpiR*	ADP1 *comP*^{T129C} Δ*pilB2* Δ*cpiR*::*kan*	4E, 4G
CE284		SAD2809	Δ*pilB* Δ*cpiR* Δ*cpiA*	ADP1 *comP*^{T129C} Δ*pilB* Δ*cpiRA*::*kan*	4E
CE285		SAD2807	Δ*pilB2* Δ*cpiR* Δ*cpiA*	ADP1 *comP*^{T129C} Δ*pilB2* Δ*cpiRA*::*kan*	4E, 4G
CE265		SAD2645	Δ*cpiR* Δ*pilT*	ADP1 *comP*^{T129C} Δ*cpiR*::*kan* Δ*pilT*::*spec*	4E, 4F
CE266		SAD2810	Δ*cpiR* Δ*cpiA* Δ*pilT*	ADP1 *comP*^{T129C} Δ*cpiRA*::*kan* Δ*pilT*::*spec*	4E
CE302		SAD2822	Δ*pilB* Δ*cpiR* Δ*pilT*	ADP1 *comP*^{T129C} Δ*pilB* Δ*cpiR*::*kan* Δ*pilT*::*spec*	4E, 4F
CE303		SAD2820	Δ*pilB2* Δ*cpiR* Δ*pilT*	ADP1 *comP*^{T129C} Δ*pilB2* Δ*cpiR*::*kan* Δ*pilT*::*spec*	4E, 4F
CE304		SAD2823	Δ*pilB* Δ*cpiR* Δ*cpiA* Δ*pilT*	ADP1 *comP*^{T129C} Δ*pilB* Δ*cpiRA*::*kan* Δ*pilT*::*spec*	4E
CE305		SAD2821	Δ*pilB2* Δ*cpiR* Δ*cpiA* Δ*pilT*	ADP1 *comP*^{T129C} Δ*pilB2* Δ*cpiRA*::*kan* Δ*pilT*::*spec*	4E, 4F

CE strains available upon request from CKE, ZG strains available upon request from ZG, TND or SAD strains available upon request from ABD.

Supplementary table 2. Primers used for strain construction.

Primer name	Primer sequence 5' → 3' (overlapping regions underlined, point mutations in bold)	Description
DOG0175	AAACTATCCAGATAAGGGAAAGC	comP F1
CEdalia87	GAAGTTGTCGTACATTTCCA**ACA**TACAGCGCCACTAGCATATG	comPT129C R1
CEdalia88	CATATGCTAGTGGCGCTGTA**TGT**TGGAAATGTACGACAACTTC	comPT129C F2
DOG0178	AGGATCTGTAATGACGGGTTGAG	comP R2
DOG0176	<u>GTCGACGGATCCCCGGAAT</u>CATAAAATTTCTCCACCAATGTTG	ΔcomP R1
DOG0177	<u>GAAGCAGCTCCAGCCTACA</u>TGATAGTAGTACTATATGGCTTTAAAAG	ΔcomP F2
ABD123	ATTCCGGGGATCCGTCGAC	AbR cassette F
ABD124	TGTAGGCTGGAGCTGCTTC	AbR cassette R
BBC1934	GTTACAAAGTCAGGGACGTAAAG	ΔpilT F1
BBC1935	<u>GTCGACGGATCCCCGGAAT</u>ATCCATATTTTCCCCGAAGATCG	ΔpilT R1
BBC1936	<u>GAAGCAGCTCCAGCCTACA</u>TAAGAATAACGCTACTCGATCTG	ΔpilT F2
BBC1939	GGTATTCAGATTGATCGTCAGTTAG	ΔpilT R2
CE31	GCTGTTTGATTTTATCCAGAACCTTG	ΔpilG F1
CE61	<u>TGACGTGATTACGAATTGCGGT</u>GATTTTGATCACTTTGGATTTGTCTTCC	ΔpilG R1
CE62	<u>GGAAGACAAATCCAAAGTGATCAAAATC</u>ACCGCAATTCGTAATCACGTCA	ΔpilG F2
CE34	GCGCTGTAATTACAATAGTGTTCGGA	ΔpilG R2 and pilG-mRuby3 R2
CE37	GGTGGTACGCAGTTGTGGTAAAAT	ΔpilH F1 and pilG-mRuby3 F1
CE63	<u>ATCATCCGTATGAATATATTGCTGGAT</u>TGAATCATCAACAATCAAAATACGTGCCA	ΔpilH R1
CE64	<u>TGGCACGTATTTTGATTGTTGATGATTCA</u>ATCCAGCAATATATTCATACGGATGAT	ΔpilH F2
CE40	GGCAACTCAATTCAAACTGATGTACTT	ΔpilH R2
CE101	TTCAACGCTCATTCGTTGCCAT	ΔpilI F1
CE102	<u>ATCAATATTGTGAAGCATTCATAAAATCTGT</u>CAAACCAGCAGATGTGGTTGC	ΔpilI R1
CE103	<u>GCAACCACATCTGCTGGTTTG</u>ACAGATTTTATGAATGCTTCACAATATTGAT	ΔpilI F2
CE104	CAGTTGACTGGTTACTGGTTCTGC	ΔpilI R2
CE107	GGTTTTTCCAGTTTTTAATGCTGCC	ΔpilJ F1

CE108	<u>TCAGATGCAAAATCAAGCTTATACGT</u>CGCTTTATTTTTTTTCCGATTGATTGA	ΔpilJ R1
CE109	<u>TCAATCAATCGGAAAAAAAATAAAGC</u>GACGTATAAGCTTGATTTTGCATCTGA	ΔpilJ F2
CE110	CCGAACAGTACGCTGCAATC	ΔpilJ R2
CE29	AATATCTTGTCGCCGATTCATTGAAA	ΔchpA F1
CE113	<u>GGTTTTGGCTATGTTTAATCGCTTT</u>CGTCTCTATAAGATGATTTACTTGTGCA	ΔchpA R1
CE114	<u>TGCACAAGTAAATCATCTTATAGAGACG</u>AAAGCGATTAAACATAGCCAAAACC	ΔchpA F2
CE115	CTGTATTTTGATGCAATGTTGGAGATT	ΔchpA R2
CE502	AATGGTGGGCCGTGAAAAAGC	ΔompA F1
CE503	<u>GTCGACGGATCCCCGGAAT</u>AGTAGCAAGTGCAATACGACTCAATTTc	ΔompA R1
CE504	<u>GAAGCAGCTCCAGCCTACA</u>GTTCAACCTGGTCAACAAGCTCC	ΔompA F2
CE505	CCAGATCAGATACCAGATCAGGTGATTAT	ΔompA R2
CE735	GTGCACTGGGGATATCTGCACT	ΔfimL F1
CE736	<u>GTCGACGGATCCCCGGAAT</u>ATCGAAGCGAATAGAAACTTGTGAAGAC	ΔfimL R1
CE737	<u>GAAGCAGCTCCAGCCTACA</u>AACGCCTCCCAATTTCAATCAACG	ΔfimL F2
CE738	TGCTAACAGAATTGTATGTGTCTACGCC	ΔfimL R2
CE590	ATGAGAACAGATTTTAACGTTTTTACGACTGG	PilQ-mCherry up F
CE196	<u>GAACCAGCAGCTGAACCAGCTGAACC</u>ATGATTTCCAACAATAGTGTCATTAACTATTCG	PilQ-mCherry up R
CE197	TTCAGCTGCTGGTTCAGGTGAATTTATGGTTTCCAAGGGCGAGGA	PilQ-mCherry mCherry region F
CE198	<u>TTACTTGAATTAAAACTATA</u>TTATTTGTACAGCTCATCCATGCCA	PilQ-mCherry mCherry region R
CE637	<u>TATAGTTTTAATTCAAGTAATT</u>CAATAGGTGACTCCATTCCGGGGATCCGTCGAC	PilQ-mCherry AbR region F1
CE593	<u>TCGTTGCTGCTGCGTAACAT</u>GCGAAACTATCCTCATCCTGTCTCTTG	PilQ-mCherry AbR region R1
CE594	<u>CAGGATGAGGATAGTTTCGC</u>ATGTTACGCAGCAGCAACGATG	PilQ-mCherry AbR region F2
CE595	<u>GGAACTTCAAGATCCCCTTA</u>TTAGGTGGCGGTACTTGGGTC	PilQ-mCherry GmR region R2

CE596	<u>GACCCAAGTACCGCCACCTAA</u>TAAGGGGATCTTGAAGTTCCTATTCCG	PilQ-mCherry Ptac region F
CE336	GCTTAATTACCTCCTAATTGAATTCCTAGGC	PilQ-mCherry Ptac region R
CE638	<u>GGAATTCAATTAGGAGGTAATTAAGC</u>TTGCCAAGCAAAGAGTTTGACACC	PilQ-mCherry down F
CE592	TCTAATGCCGCTGCACTGTCAC	PilQ-mCherry down R
CE546	CCAGATCTCTACGCATTTCACCG	ΔACIAD0653 F1
CE547	<u>GTCGACGGATCCCCGGAAT</u>GACCCCAAGAATAGATATTTTGATTGAGTTGT	ΔACIAD0653 R1
CE548	<u>GAAGCAGCTCCAGCCTACA</u>CGCGTTGCAATTGTAATCGTTCC	ΔACIAD0653 F2
CE549	GTAGAAGGGTCTCATGTTGCGGG	ΔACIAD0653 R2
CE552	CACAATACGTCCCAGCAATTGAAAC	ΔACIAD0898 F1
CE553	<u>GTCGACGGATCCCCGGAAT</u>GACTAGTGCACGCATTGCATGC	ΔACIAD0898 R1
CE554	<u>GAAGCAGCTCCAGCCTACA</u>GAGATTAGCATTTACGAACGTCAATAAGCG	ΔACIAD0898 F2
CE555	CGACACATGTTTGCTCTGCATCC	ΔACIAD0898 R2
CE146	<u>TTCACCTGAACCAGCAGCTGAACCAGCTGAACC</u>TACACTGACGTGATTACGAATTGCG	pilG-mRuby3 R1
CE147	TGGTTCAGCTGCTGGTTCAGGTGAATTTATGGTATCTAAGGGTGAAGAATTGATTAAGG	mRuby3 F
CE148	TTACTTATATAATTCATCCATTCCACCCCCAAG	mRuby3 R
CE149	<u>GGGTGGAATGGATGAATTATATAAGTAA</u>CACGATATTGTTTTGAGGTCTTTATGGCAC	pilG-mRuby3 F2
CE389	TCAGGAAGTTGCCTGAACCTGA	ΔpilC F1
CE390	<u>ACAGAGCCCATTTGGAAAATTGG</u>TGGCATTACTTGCGTTTTTTTTGCT	ΔpilC R1
CE391	<u>AGCAAAAAAACGCAAGTAATGCCA</u>CCAATTTTCCAAATGGGCTCTGT	ΔpilC F2
CE392	TTGAAAAGCTGGCTGGCCTTC	ΔpilC R2
CE136	TTGGTCGATAATGCAGATCAGCT	ΔpilY1 F1
CE137	<u>GGGGTGCTTTGCTCATACCATTTTAA</u>GAGATACTGAGTGTATATCGATTTTTTCAT	ΔpilY1 R1
CE138	<u>ATGAAAAAATCGATATACACTCAGTATCTC</u>TTAAAATGGTATGAGCAAAGCACCCC	ΔpilY1 F2

CE139	GCAGGGATATCAGATTTTAACTGTGC	ΔpilY1 R2
CE85	AATAAGAGCGAATAGCAAACTAATTGTCG	ΔpilQ F1
CE340	<u>TTTCCAACAATAGTGTCATTAACTATTC</u>GAGTCGTAAAAACGTTAAAATCTGTTCTCAT	ΔpilQ R1
CE341	<u>ATGAGAACAGATTTTAACGTTTTTACGACT</u>CGAATAGTTAATGACACTATTGTTGGAAA	ΔpilQ F2
CE88	AGCGTATAACGATCAATTACTTCGC	ΔpilQ R2
BBC1891	AATCACTTGAAGTGCAGATTCG	ΔcomEA F1
BBC3451	<u>GTCGACGGATCCCCGGAAT</u>ATGACGTATTGACATGAATTAACC	ΔcomEA R1
BBC3452	<u>GAAGCAGCTCCAGCCTAC</u>ATAAGAGGCTGAGTCATGCTCAAAATAC	ΔcomEA F2
BBC1894	ATCTTCACCTTCTCCAAACTGG	ΔcomEA R2
DOG0180	AGTATTTGAGTTCGGGTATTATTGC	ΔcomEC F1
DOG0181	<u>GTCGACGGATCCCCGGAAT</u>CATGGCAATGCCAGCAATC	ΔcomEC R1
DOG0182	<u>GAAGCAGCTCCAGCCTAC</u>AAAGGGAGCGATTCATTTTAAGTTTG	ΔcomEC F2
DOG0183	TTACTCAAGCCGTTACAGTTCTG	ΔcomEC R2
DOG0140	GTTGCTGCATTTGTTCGATCTG	ΔcomM F1
DOG0141	<u>GTCGACGGATCCCCGGAAT</u>CATACTATTATTGTTCCATTATGGTGC	ΔcomM R1
DOG0142	<u>GAAGCAGCTCCAGCCTAC</u>ATATCGCAGTGAACATAGCTAAAA	ΔcomM F2
DOG0143	ATCAGTGGTTGGGAAGGTG	ΔcomM R2
CE23	TTATGACTATCAATCTCAAGCACATGCT	ΔpilB F1
CE93	<u>ATTCACTGGTTACACGATTAATTTCCTG</u>AAACTTAGGTGGTGTTGTAAATGCTG	ΔpilB R1
CE94	<u>CAGCATTTACAACACCACCTAAGTTT</u>CAGGAAATTAATCGTGTAACCAGTGAAT	ΔpilB F2
CE26	CTTTCATGGCGTGTTTGTCGTG	ΔpilB R2
CE210	AAGCCAACCATAATCCGATTGTGAC	ΔpilB2 F1
CE328	<u>TGGGACCACCCTTAAGACTTCTT</u>CCCATTGTGTATCTATTTCAAAATGATATGTCAT	ΔpilB2 R1 (in frame)
CE329	<u>ATGACATATCATTTTGAAATAGATACACAATGG</u>AAGAAGTCTTAAGGGTGGTCCCA	ΔpilB2 F2 (in frame)
CE213	CTCACGCCACCCAATCAAGAC	ΔpilB2 R2
CE240	<u>GTCGACGGATCCCCGGAAT</u>TTATTTCTCCCCCACACCATTCAC	ΔpilB2 R1 (AbR disruption)

CE241	<u>GAAGCAGCTCCAGCCTACA</u>TAAAAGGTTAGTTTAGATCTTAAGATTCACCTCAT	ΔpilB2 F2 (AbR disruption)
DOG0130	CCTCTCCACTCGTTTCTAAAGAAC	ΔcpiR or ΔcpiA or cpiA-3X FLAG F1
DOG0131	<u>GTCGACGGATCCCCGGAAT</u>CATAAGCCGATACTATCTGGATGG	ΔcpiR R1
DOG0132	<u>GAAGCAGCTCCAGCCTACA</u>GAAGTTCAATGTCAGTTAAAACATC	ΔcpiR F2
DOG0133	GACGTTGATTACGTAAATTTAATTCG	ΔcpiR or ΔcpiA or cpiA-3X FLAG R2
BBC921	<u>GTCGACGGATCCCCGGAAT</u>TTGATCACTTACTAATTTCATTATCATTTAAATG	ΔcpiA R1
BBC922	<u>GAAGCAGCTCCAGCCTACA</u>TAAAAAAGGACTGATTTCTCAGCCC	ΔcpiA F2
BBC2565	TCCACCACTTCCACCTGCATTTTCATGAAACACTTGCTGC	cpiA-3X FLAG R1
BBC2570	GCAGGTGGAGCAGGTGGATAAAAAAGGACTGATTTCTCAGCCC	cpiA-3X FLAG F2
BBC2272	GCAGGTGGAAGTGGTGGAGATTATAAAGACCATGATGGTGACTACAAGGATCACGACATTGATTATAAGGATGACGATGACAAAGCAGGTGGAGCAGGTGGA	3X FLAG MIDDLE F
BBC2273	TCCACCTGCTCCACCTGCTTTGTCATCGTCATCCTTATAATCAATGTCGTGATCCTTGTAGTCACCATCATGGTCTTTATAATCTCCACCACTTCCACCTGC	3X FLAG MIDDLE R
DOG0229	<u>CAATTTCACACAGGATCCCGGG</u>AGGAGGTAACGTAATGATAATGAAATTAGTAAGTGATCAAATTC	Amplify *cpiA* for P*tac*-*cpiA* construct F
DOG0230	<u>TGTAGGCTGGAGCTGCTTC</u>TTAATTTTCATGAAACACTTGCTGC	Amplify *cpiA* for P*tac*-*cpiA* construct R
BBC3072	<u>CACCATTATGATGGCAATCGTATGA</u>TTCGAAGTATGTGAAAAATCTGC	Amplify P*cpiA* for P*cpiA*-GFP F
BBC3074	<u>CCCGGGATCCTGTGTGAAATTG</u>ACTAGCAGTATTGTATTACAAAACTTTTG	Amplify P*cpiA* for P*cpiA*-GFP R
BBC3075	<u>CAATTTCACACAGGATCCCGGGAGGAGG</u>TCGGCTTATGGATATAGGTCGC	Amplify *cpiR* for P_{BAD}-*cpiR* construct F
BBC2560	<u>TGTAGGCTGGAGCTGCTTC</u>TTACTTTTTAGTGTCTATCAAAGTAAAAATGAGATG	Amplify *cpiR* for P_{BAD}-*cpiR* construct R