Mapping single-cell atlases throughout Metazoa unravels cell type evolution

Alexander J. Tarashansky[1], Jacob M. Musser[2,§], Margarita Khariton[1,§], Pengyang Li[1],

Detlev Arendt[2,3], Stephen R. Quake[1,4,5], Bo Wang[1,6*]

[1]Department of Bioengineering, Stanford University, Stanford, CA, USA.

[2]European Molecular Biology Laboratory, Developmental Biology Unit, Heidelberg,

Germany.

[3]Centre for Organismal Studies, University of Heidelberg, Heidelberg, Germany.

[4]Department of Applied Physics, Stanford University, Stanford, CA, USA.

[5]Chan Zuckerberg Biohub, San Francisco, CA, USA.

[6]Department of Developmental Biology, Stanford University School of Medicine, Stanford,

CA, USA.

[§]These authors contributed equally to this work.

*Correspondence: wangbo@stanford.edu.

Abstract

Comparing single-cell transcriptomic atlases from diverse organisms can elucidate the

origins of cellular diversity and assist the annotation of new cell atlases. Yet,

comparison between distant relatives is hindered by complex gene histories and

diversifications in expression programs. Previously, we introduced the self-assembling

manifold (SAM) algorithm to robustly reconstruct manifolds from single-cell data

(Tarashansky et al., 2019). Here, we build on SAM to map cell atlas manifolds across

species. This new method, SAMap, identifies homologous cell types with shared

expression programs across distant species within phyla, even in complex examples

where homologous tissues emerge from distinct germ layers. SAMap also finds many

genes with more similar expression to their paralogs than their orthologs, suggesting

paralog substitution may be more common in evolution than previously appreciated.

Lastly, comparing species across animal phyla, spanning mouse to sponge, reveals

ancient contractile and stem cell families, which may have arisen early in animal

evolution.

Introduction

There is much ongoing success in producing single-cell transcriptomic atlases to investigate the cell type diversity within individual organisms (Regev et al., 2017). With the growing diversity of cell atlases across the tree of life (Briggs et al., 2018; Cao et al., 2019; Fincher et al., 2018; Hu et al., 2020; Musser et al., 2019; Plass et al., 2018; Siebert et al., 2019; Wagner et al., 2018), a new frontier is emerging: the use of cross-species cell type comparisons to unravel the origins of cellular diversity and uncover species-specific cellular innovations (Arendt et al., 2019; Shafer, 2019). Further, these comparisons promise to accelerate cell type annotation and discovery by transferring knowledge from well-studied model organisms to under-characterized animals.

However, recent comparative single-cell analyses are mostly limited to species within the same phylum (Baron et al., 2016; Geirsdottir et al., 2019; Sebé-Pedrós et al., 2018; Tosches et al., 2018). Comparisons across longer evolutionary distances and across phyla are challenging for two major reasons. First, gene regulatory programs diversify during evolution, diminishing the similarities in cell type specific gene expression patterns. Second, complex gene evolutionary history causes distantly related organisms to share few one-to-one gene orthologs (Nehrt et al., 2011), which are often relied upon for comparative studies (Briggs et al., 2018; Shafer, 2019). This effect is compounded by the growing evidence suggesting that paralogs may be more functionally similar than orthologs across species, due to differential gain (neo-functionalization), loss (non-functionalization), or partitioning (sub-functionalization) events among paralogs (Nehrt et al., 2011; Prince & Pickett, 2002; Stamboulian et al., 2020; Studer & Robinson-Rechavi, 2009).

Here, we present the Self-Assembling Manifold mapping (SAMap) algorithm to enable

mapping single-cell transcriptomes between phylogenetically remote species. SAMap

relaxes the constraints imposed by sequence orthology, using expression similarity

between mapped cells to infer the relative contributions of homologous genes, which in

turn refines the cell type mapping. In addition, SAMap uses a graph-based data

integration technique to identify reciprocally connected cell types across species with

greater robustness than previous single-cell data integration methods (Haghverdi et al.,

2018; Hie et al., 2019; Polański et al., 2019; Stuart et al., 2019).

Using SAMap, we compared seven whole-body cell atlases from species spanning animal

phylogeny, which have divergent transcriptomes and complex molecular homologies

(**Figure 1A-B** and **Supplementary Table 1**). We began with well-characterized cell types

in developing frog and fish embryos. We found broad concordance between

transcriptomic signatures and ontogenetic relationships, which validated our mapping

results, yet also detected striking examples of homologous cell types emerging from

different germ layers. We next extended the comparison to animals from the same phylum

but with highly divergent body plans, using a planarian flatworm and a parasitic blood

fluke, and found one-to-one homologies even between cell subtypes. Comparing all

seven species from sponge to mouse, we identified densely interconnected cell type

families broadly shared across animals, including contractile and stem cells, along with

their respective gene expression programs. Lastly, we noticed that homologous cell types

often exhibit differential expression of orthologs and similar expression of paralogs,

suggesting that the substitution and swapping of paralogs in cell types may be more

common in evolution than previously appreciated. Overall, our study represents an

important step towards analyzing the evolutionary origins of specialized cell types and

their associated gene expression programs in animals.

Results

The SAMap algorithm

SAMap iterates between two modules. The first module constructs a gene-gene bipartite

graph with cross-species edges connecting homologous gene pairs, initially weighted by

protein sequence similarity (**Figure 1C**). In the second module, SAMap uses the gene-

gene graph to project the two single-cell transcriptomic datasets into a joint, lower-

dimensional manifold representation, from which each cell's mutual cross-species

neighbors are linked to stitch the cell atlases together (**Figure 1D**). Then, using the joint

manifold, the expression correlations between homologous genes are computed and

used to reweight the edges in the gene-gene homology graph in order to relax SAMap's

initial dependence on sequence similarity. The new homology graph is used as input to

the subsequent iteration of SAMap, and the algorithm continues until convergence,

defined as when the cross-species mapping does not significantly change between

iterations (**Figure 1E**).

This algorithm overcomes several challenges inherent to mapping single-cell

transcriptomes between distantly related species. First, complex gene evolutionary

history often results in many-to-many homologies with convoluted functional relationships

(Briggs et al., 2018; Nehrt et al., 2011). SAMap accounts for this by using the full

homology graph to project each dataset into both its own and its partner's respective

principal component (PC) spaces, constructed by the SAM algorithm, which we previously

developed to robustly and sensitively identify cell types (Tarashansky et al., 2019). The

resulting within- and cross-species projections are concatenated to form the joint space.

For the cross-species projections, we translate each species' features into those of its

partner, with the expression for individual genes imputed as the weighted average of their

homologs specified in the gene-gene bipartite graph. Iteratively refining the homology

graph to only include positively correlated gene pairs prunes the many-to-many

homologies to only include genes that are expressed in the same mapped cell types.

Second, frequent gene losses and the acquisitions of new genes result in many cell type

gene expression signatures being species-specific, limiting the amount of information that

is comparable across species. Restricting the analysis of each dataset to only include

genes that are shared across species would result in a decreased ability to resolve cell

types and subtypes with many species-specific gene signatures. SAMap solves this

problem by constructing the joint space through the concatenation of within- and cross-

species projections, thus encoding all genes from both species.

Lastly, the evolution of expression programs gradually diminishes the similarity between

homologous cell types. To account for this effect, SAMap links cell types across species

while tolerating their differences. Cells are mapped by calculating each of their k mutual

nearest cross-species neighbors in the combined projection. To establish more robust

mutual connectivity, we integrate information from each cell's local, within-species

neighborhood (**Figure 1D**), overcoming the inherent stochasticity of cross-species

correlations. Two cells are thus defined as mutual nearest cross-species neighbors when

their respective neighborhoods have mutual connectivity. It is important to note that the

magnitude of connections is not directly calculated from their expression similarity,

allowing cell types with diverged expression profiles to be tightly linked if they are among

each other's closest cross-species neighbors.

Paralog substitutions are prevalent between homologous cell types in frog and fish

We first applied SAMap to the *Xenopus* and zebrafish atlases, which both encompass

embryogenesis until early organogenesis (Briggs et al., 2018; Wagner et al., 2018).

Previous analysis had linked cell types between these two organisms by matching

ontogeny, thereby providing a reference for comparison. SAMap produced a combined

manifold with a high degree of cross-species alignment while maintaining high resolution

for distinguishing cell types in each species (**Figure 2A**). We measured the mapping

strength between cell types by calculating an alignment score (edge width in **Figure 2B**

and color map in **Figure 2C**), defined as the average number of mutual nearest cross-

species neighbors of each cell relative to the maximum possible number of neighbors.

SAMap revealed broad agreement between transcriptomic similarity and developmental

ontogeny, linking 26 out of 27 expected pairs based on previous annotations (**Figure 2B**

and **Supplementary Table 2**) (Briggs et al., 2018). The only exception is the embryonic

kidney (pronephric duct/mesenchyme), potentially indicating that their gene expression

programs have significantly diverged. In addition, SAMap succeeded in drawing parallels

between the development of homologous cell types and matched time points along

several cell lineages (**Figure 2C**). While the concordance was consistent across cell

types, we noticed that the exact progression of developmental timing can vary, suggesting

that SAMap can quantify heterochrony with cell type resolution.

SAMap also weakly linked several closely related cell types with different ontogeny. For

example, optic cells from both species are also connected to eye primordium, frog skeletal

muscles to fish presomitic mesoderm, and frog hindbrain to fish forebrain/midbrain.

Notable exceptions also included mapped secretory cell types that differ in their

developmental origin and even arise from different germ layers (black edges in **Figure**

2B). They are linked through a large set of genes including conserved transcription factors

(e.g., *foxa1* (Dubaissi et al., 2014), *grhl* (Miles et al., 2017)) and proteins involved in

vesicular protein trafficking (**Figure 2 – figure supplement 1**). This observation supports

the notion that cell types may be transcriptionally and evolutionarily related despite having

different developmental origins (Arendt et al., 2016).

To benchmark SAMap performance, we used Eggnog (Huerta-Cepas et al., 2019) to

define one-to-one vertebrate orthologs between fish and frog and fed these gene pairs

as input to several broadly used single-cell data integration methods, Seurat (Stuart et

al., 2019), Liger (Welch et al., 2019), Harmony (Korsunsky et al., 2019), Scanorama (Hie

et al., 2019), and BBKNN (Polański et al., 2019). We found that they failed to map the two

atlases, yielding minimal alignment between them (**Figure 2D** and **Figure 2 – figure**

supplement 2). We also compared the results when restricting SAMap to using the one-

to-one orthologs instead of the full homology graph. Even when removing the many-to-

many gene homologies and the iterative refinement of the homology graph, we identified

similar, albeit weaker, cell type mappings. This suggests that, at least for the frog and fish

comparison, SAMap's performance is owed in large part to its robust, atlas stitching

approach.

The key benefit of using the full homology graph is to enable the systematic identification

of gene paralogs that exhibit greater similarity in expression across species than their

corresponding orthologs. These events are expected to arise as the result of gene

duplications followed by diversification of the resulting in-paralogs (Studer & Robinson-

Rechavi, 2009). In addition, genetic compensation by transcriptional adaptation, where

loss-of-function mutations are balanced by upregulation of related genes with similar

sequences (El-Brolosy et al., 2019), could also result in this signature. In total, SAMap

selected 8,286 vertebrate orthologs and 7,093 eukaryotic paralogs, as enumerated by

Eggnog, for manifold alignment. Among these, 565 genes have markedly higher

expression correlations (correlation difference > 0.3) with their paralogs than their

orthologs (**Figure 2E** and **Figure 2 – figure supplement 3**), and 209 genes have

orthologs that are either completely absent or lowly-expressed with no cell-type specificity

(**Supplementary Table 3**), suggesting that they may have lost their functional roles at

some point and were compensated for by their paralogs. We term these events as

"paralog substitutions". SAMap linked an additional 297 homologous pairs not previously

annotated by orthology or paralogy, but which exhibited sequence similarity and high

expression correlations (>0.5 Pearson correlation). These likely represent unannotated

orthologs/paralogs or isofunctional, distantly related homologs (Gabaldón & Koonin,

2013). These results illustrate the potential of SAMap in leveraging single-cell gene

expression data for pruning the networks of homologous genes to identify evolutionary

substitution of paralogs and, more generally, identify non-orthologous gene pairs that may

perform similar functions in the cell types within which they are expressed.

Homologous cell types between two flatworm species with divergent body plans

To test if we can identify homologous cell types in animals with radically different body

plans, we mapped the cell atlases of two flatworms, the planarian *Schmidtea*

mediterranea (Fincher et al., 2018), and the trematode *Schistosoma mansoni*, which we

collected recently (Li et al., 2020). They represent two distant lineages within the same

phylum but have remarkably distinct body plans and autecology (Laumer et al., 2015;

Littlewood & Waeschenbach, 2015). While planarians live in freshwater and are known

for their ability to regenerate (Reddien, 2018), schistosomes live as parasites in humans.

The degree to which cell types are conserved between them is unresolved, given the vast

phenotypic differences caused by the transition from free-living to parasitic habits

(Laumer et al., 2015).

SAMap revealed broad cell type homology between schistosomes and planarians. The

schistosome had cells mapped to the planarian stem cells, called neoblasts, as well as

most of the differentiated tissues: neural, muscle, intestine, epidermis, parenchymal,

protonephridia, and *cathepsin*[+] cells, the latter of which consists of cryptic cell types that,

until now, have only been found in planarians (Fincher et al., 2018) (**Figure 3A**). These

mappings are supported by both known cell type specific marker genes and numerous

homologous transcriptional regulators (**Figure 3B** and **Figure 3 – figure supplement 1**).

We next determined if cell type homologies exist at the subtype level. For this, we

compared the neoblasts, as planarian neoblasts are known to comprise populations of

pluripotent cells and tissue-specific progenitors (Fincher et al., 2018; Zeng et al., 2018).

By mapping the schistosome neoblasts to a planarian neoblast atlas (Zeng et al., 2018),

we found that the schistosome has a population of neoblasts (ε-cells (Wang et al., 2018))

that cluster with the planarian's pluripotent neoblasts, both expressing a common set of

TFs (e.g., *soxp2, unc4*, *pax6a*, *gcm1*) (**Figure 3C-D**). The ε-cells are closely associated

with juvenile development and lost in adult schistosomes (Wang et al., 2018), indicating

pluripotent stem cells may be a transient population restricted to their early developmental

stages. This is consistent with the fact that, whereas schistosomes can heal wounds, they

have limited regenerative ability (Wendt & Collins, 2016). SAMap also linked other

schistosome neoblast populations with planarian progenitors, including two populations

of schistosome neoblasts (denoted as μ (Tarashansky et al., 2019) and μ') to planarian

muscle progenitors, all of which express *myoD*, a canonical master regulator of

myogenesis (Scimone et al., 2017). These likely represent early and late muscle

progenitors, respectively, as μ-cells do not yet express differentiated muscle markers

such as *troponin*, whereas μ'-cells do (**Figure 3 – figure supplement 2).**

Cell type families spanning the animal tree of life

To compare cell types across broader taxonomic scales, we extended our analysis to

include juvenile freshwater sponge (*Spongilla lacustris*) (Musser et al., 2019), adult *Hydra*

(*Hydra vulgaris*) (Siebert et al., 2019), and mouse (*Mus musculus*) embryogenesis

(Pijuan-Sala et al., 2019) atlases. In total, SAMap linked 1,051 cross-species pairs of cell

types, defined by the annotations used in each respective study. 95% of the cell type

pairs are supported by at least 40 enriched gene pairs, and 87% are supported by more

than 100 gene pairs, indicating that SAMap does not spuriously connect cell types with

limited overlap in expression profiles (**Figure 4A**).

We next extended the notion of cell type pairs to cell type trios, as mapped cell types gain

additional support if they share transitive relationships to other cell types through

independent mappings, forming cell type triangles among species. The transitivity of a

cell type pair (edge) or a cell type (node) can be quantified as the fraction of triads to

which they belong that form triangles (**Figure 4B**). The majority (81%) of cell type pairs

have non-zero transitivity independent of alignment score and the number of enriched

gene pairs (**Figure 4 – figure supplement 1-2**). Cell type pairs with fewer than 40

enriched gene pairs tend to have lower (<0.4) transitivity (**Figure 4 – figure supplement**

2). In addition, 16% of mapped cell type pairs have zero edge transitivity but non-zero

node transitivity, meaning that at least one of the cell types connects to only a single

member of an interconnected cell type group (**Figure 4C**). Such edges may be of lower

confidence as they should connect to other members of the same group and are thus

excluded from downstream analysis.

Among the interconnected groups of cell types, we identified families of contractile cells

and neural cells (**Figure 4D**). Both cell type families are highly transitive compared to the

overall graph transitivity (bootstrap p-value < 1×10^{-5}), meaning that their constituent cell

types have more transitive edges within the group than outside the group (**Figure 4E**). In

addition, the dense, many-to-many connections within the contractile and neural families

are each supported by at least 40 enriched gene pairs (**Figure 4F**). Consistent with the

nerve net hypothesis suggesting a unified origin of neural cell types (Tosches & Arendt,

2013), the neural family includes vertebrate brain tissues, both bilaterian and cnidarian

neurons, cnidarian nematocytes that share the excitatory characteristics of neurons (Weir

et al., 2020), and *Spongilla* choanocytes and apopylar cells, both of which are not

considered as neurons but have been shown to express postsynaptic-like scaffolding

machinery (Musser et al., 2019; Wong et al., 2019). The contractile family includes

myocytes in bilaterian animals, *Hydra* myoepithelial cells that are known to have

contractile myofibrils (Buzgariu et al., 2015), and sponge pinacocytes and

myopeptidocytes, both of which have been implicated to play roles in contractility (Musser

et al., 2019; Sebé-Pedrós et al., 2018). In contrast to the families encompassing all seven

species, we also found a fully interconnected group that contains invertebrate pluripotent

stem cells, including planarian and schistosome neoblasts, *Hydra* interstitial cells, and

sponge archeocytes (Alié et al., 2015). The lack of one-to-one connections across phyla

is in keeping with recent hypotheses that ancestral cell types diversified into families of

cell types after speciation events (Arendt et al., 2016, 2019). Our findings thus suggest

that these cell type families diversified early in animal evolution.

Transcriptomic signatures of cell type families

The high interconnectedness between cell types across broad taxonomic scales suggests

that they should share ancestral transcriptional programs (Arendt et al., 2016; Tosches

et al., 2018). SAMap identified broad transcriptomic similarity between bilaterian and non-

bilaterian contractile cells that extends beyond the core contractile apparatus. It links a

total of 23,601 gene pairs, connecting 5,471 unique genes, which are enriched in at least

one contractile cell type pair. Performing functional enrichment analysis on these genes,

we found cytoskeleton and signal transduction functions to be enriched (p-value < 10^{-3})

based on the KOG functional classifications (Tatusov et al., 2003) assigned by Eggnog

(**Figure 5A**). These genes include orthology groups spanning diverse functional roles in

contractile cells, including actin regulation, cell adhesion and stability, and signaling

(**Figure 5B** and **Supplementary Table 4**), indicating that contractile cells were likely

multifunctional near the beginning of animal evolution.

We also identified several transcriptional regulators shared among contractile cells

(**Figure 5B**). Previously known core regulators involved in myocyte specification (Brunet

et al., 2016) were enriched only in bilaterian (e.g., *myod*, and *tcf4/E12*) or vertebrate

contractile cells (e.g., *mef2*). In contrast, we found homologs of Muscle Lim Protein (*Csrp*)

and Forkhead Box Group 1 (Larroux et al., 2008) enriched in contractile cells from all

seven species. The Fox proteins included FoxC, which is known to regulate cardiac

muscle identity in vertebrates (Brunet et al., 2016) and is contractile-specific in all species

except schistosome and *Spongilla*. Notably, we also identified FoxG orthologs to be

enriched in three of the four invertebrates (**Figure 5 – figure supplement 1**), suggesting

that FoxG may play an underappreciated role in contractile cell specification outside

vertebrates.

For the family of invertebrate stem cells, we identified 3,343 genes that are enriched in at

least one cell type pair and observed significant enrichment (p-value < 10^{-3}) of genes

involved in translational regulation such as RNA processing, translation, and post-

translational modification (**Figure 5C**). These genes form 979 orthology groups, 17% of

which are enriched in all cell types of this family (**Supplementary Table 4**). Importantly,

other stem cell populations in *Hydra* and planarian lineage-restricted neoblasts have

significantly reduced expression of these genes (**Figure 5D**). These results suggest that

SAMap identified a large, deeply conserved gene module specifically associated with

multipotency.

Discussion

Cell types evolve as their gene expression programs change either as integrated units or

via evolutionary splitting that results in separate derived programs. While this notion of

coupled cellular and molecular evolution has gained significant traction in the past years,

systematically comparing cell type-specific gene expression programs across species

has remained a challenging problem. Here, we map single-cell atlases between

evolutionarily distant species in a manner that accounts for the complexity of gene

evolution. SAMap aligns cell atlases in two mutually reinforcing directions, mapping both

the genes and the cells, with each feeding back into the other. This method allows us to

identify one-to-one cell type concordance between animals in the same phylum, whereas

between phyla, we observe interconnected cell types forming distinct families. These

findings support the notion that cell types evolve via hierarchical diversification (Arendt et

al., 2019), resulting in cell type families composed of evolutionarily related cell types

sharing a regulatory gene expression program that originated in their common ancestor.

One-to-one cell type homologies should exist only if no further cell type diversification has

occurred since the speciation.

In parallel, SAMap systematically identifies instances where paralogs exhibit greater

expression similarity than orthologs across species. Paralog substitution likely occurs due

to differential loss or retention of cell type-specific expression patterns of genes that were

duplicated in the common ancestor (Studer & Robinson-Rechavi, 2009). Alternatively,

paralog substitutions could arise due to compensating upregulation of paralogs following

a loss-of-function mutation acquired by an ortholog (El-Brolosy et al., 2019). While the

analysis presented here focuses on comparisons between two species, incorporating

multiple species into a single analysis that also accounts for their phylogenetic

relatedness could enable determining the specific order of paralog substitutions,

associated cell type diversification events, and the mechanism by which they arose.

However, this would require cell atlases that consistently sample key branching points

along the tree of life. Nevertheless, identifying lineage-specific paralog substitution

signatures should be accessible in extensively studied vertebrate single-cell atlases, as

the vertebrate clade is where existing data and knowledge are most concentrated.

Besides applications in evolutionary biology, we anticipate SAMap can catalyze the

annotation of new cell atlases from non-model organisms, which often represents a

substantial bottleneck requiring extensive manual curation and prior knowledge. Its ability

to use the existing atlases to inform the annotation of cell types in related species will

keep improving as more datasets become available to better sample the diversity of cell

types. Moreover, our approach allows leveraging existing and forthcoming single-cell

gene expression data to predict functionally similar gene homologs, which can serve as

guideposts for mechanistic molecular studies.

Materials and Methods

Data and Code Availability

- The source code for SAMap is publicly available at Github (https://github.com/atarashansky/SAMap), along with the code to perform the analysis and generate the figures.

- The datasets analyzed in this study are detailed in **Supplementary Table 1** with their accessions, and annotations provided.

The SAMap Algorithm

The SAMap algorithm contains three major steps: preprocessing, mutual nearest neighborhood alignment, and gene-gene correlation initialization. The latter two are repeated for three iterations, by default, to balance alignment performance and computational runtime. SAMap runs up to one hour on an average desktop computer for 200,000 total cells.

1. Preprocessing.

1.1. Generate gene homology graph via reciprocal BLAST.

We first construct a gene-gene bipartite graph between two species by performing reciprocal BLAST of their respective transcriptomes using *tblastx*, or proteomes using *blastp*. *tblastn* and *blastx* are used for BLAST between proteome and transcriptome. When a pair of genes share multiple High Scoring Pairs (HSPs), which are local regions

of matching sequences, we use the HSP with the highest bit score to measure homology.

Only pairs with E-value < 10^{-6} are included in the graph.

Here we define similarity using BLAST, though SAMap is compatible with other protein

homology detection methods (e.g. HMMER (Eddy, 2008)) or orthology inference tools

(e.g. OrthoClust (Yan et al., 2014) and Eggnog (Huerta-Cepas et al., 2019)). While each

of these methods has known strengths and limitations, BLAST is chosen for its broad

usage, technical convenience, and compatibility with low-quality transcriptomes.

We encode the BLAST results into two triangular adjacency matrices, A and B, each

containing bit scores in one BLAST direction. We combine A and B to form a gene-gene

adjacency matrix G. After symmetrizing G, we remove edges that only appear in one

direction: $G = Recip(\frac{1}{2}[(A + B) + (A + B)^T]) \in \Re^{m_1+m_2 \times m_1+m_2}$, where $Recip$ only keeps

reciprocal edges, and m_1 and m_2 are the number of genes of the two species,

respectively. To filter out relatively weak homologies, we also remove edges where $G_{ab} <$

$0.25 \max_b(G_{ab})$. Edge weights are then normalized by the maximum edge weight for each

gene and transformed by a hyperbolic tangent function to increase discriminatory power

between low and high edge weights, $\hat{G}_{ab} = 0.5 + 0.5 \, tanh \, (10G_{ab}/\max_b(G_{ab}) - 5)$.

1.2. Construct manifolds for each cell atlas separately using the SAM algorithm.

The scRNAseq datasets are normalized such that each cell has a total number of raw

counts equal to the median size of single-cell libraries. Gene expressions are then log-

normalized with the addition of a pseudocount of 1. Genes expressed (i.e., $log_2(D + 1) >$

1) in greater than 96% of cells are filtered out. SAM is run using the following parameters:

preprocessing = 'StandardScaler', *weight_PCs = False*, *k = 20*, and *npcs = 150*. A detailed

description of parameters is provided previously (Tarashansky et al., 2019). SAM outputs

N_1 and N_2, which are directed adjacency matrices that encode *k*-nearest neighbor graphs

for the two datasets, respectively.

SAM only includes the top 3,000 genes ranked by SAM weights and the first 150 principal

components (PCs) in the default mode to reduce computational complexity. However,

downstream mapping requires PC loadings for all genes. Thus, in the final iteration of

SAM, we run PCA on all genes and take the top 300 PCs. This step generates a loading

matrix for each species i, $L_i \in \Re^{300 \times m_i}$.

2. Mutual nearest neighborhood alignment.

2.1. Transform feature spaces between species.

For the gene expression matrices $Z_i \in \Re^{n_i \times m_i}$, where n and m are the number of cells

and genes respectively, we first zero the expression of genes that do not have an edge

in $\hat{G}$ and standardize the expression matrices such that each gene has zero mean and

unit variance, yielding $\tilde{Z}_i$. $\hat{G}$ represents a bipartite graph in the form of $\hat{G} =$

$\begin{bmatrix} 0_{m_1,m_1} & H \in \Re^{m_1 \times m_2} \\ H^T \in \Re^{m_2 \times m_1} & 0_{m_2,m_2} \end{bmatrix}$, where $0_{m,m}$ is $m \times m$ zero matrix and H is the biadjacency

matrix. Letting $H_1 = H$ and $H_2 = H^T$ encoding directed edges from species 1 to 2 and 2

to 1, respectively, we normalize the biadjacency matrix H_i such that each row sums to 1:

$\hat{H}_i = SumNorm(H_i) \in \Re^{m_i \times m_j}$, where the $SumNorm$ function normalizes the rows to sum

to 1. The feature spaces can be transformed between the two species via weighted

averaging of gene expression, $\tilde{Z}_{ij} = \tilde{Z}_i \hat{H}_i$.

2.2. Project single-cell gene expressions into a joint PC space.

We project the expression data from two species into a joint PC space (Barkas et al.,

2019), $P_i = \tilde{Z}_i {L_i}^T$ and $P_{ij} = \tilde{Z}_{ij} {L_j}^T$. We then horizontally concatenate the principal

components P_i and P_{ij} to form $\hat{P}_i \in \Re^{n_i \times 600}$.

2.3. Calculate k-nearest cross-species neighbors for all cells.

Using the joint PCs, $\hat{P}_i$, we identify for each cell the k-nearest neighbors in the other

dataset using cosine similarity ($k = 20$ by default). Neighbors are identified using the

hnswlib library, a fast approximate nearest-neighbor search algorithm (Malkov &

Yashunin, 2020). This outputs two directed biadjacency matrices $C_i \in \Re^{n_i \times n_j}$ for $(i,j) =$

$(1,2)$ *or* $(2,1)$ with edge weights equal to the cosine similarity between the PCs.

2.4. Apply the graph-coarsening mapping kernel to identify cross-species mutual nearest

neighborhoods.

To increase the stringency and confidence of mapping, we only rely on cells that are

mutual nearest cross-species neighbors, which are typically defined as two cells

reciprocally connected to one another (Haghverdi et al., 2018). However, due to the noise

in cell-cell correlations and stochasticity in the kNN algorithms, cross-species neighbors

are often randomly assigned from a pool of cells that appear equally similar, decreasing

the likelihood of mutual connectivity between individual cells even if they have similar

expression profiles. To overcome this limitation, we integrate information from each cell's

local neighborhood to establish more robust mutual connectivity between cells across

species. Two cells are thus defined as mutual nearest cross-species neighbors when their

respective neighborhoods have mutual connectivity.

Specifically, the nearest neighbor graphs N_i calculated in step 1.2 are used to calculate

the neighbors of cells t_i hops away along outgoing edges: $\bar{N}_i = N_i^{t_i}$, where $\bar{N}_i$ are

adjacency matrices that contain the number of paths connecting two cells t_i hops away,

for $i = 1\ or\ 2$. t_i determines the length-scale over which we integrate incoming edges for

species i. Its default value is 2 if the dataset size is less than 20,000 cells and 3 otherwise.

However, cells within tight clusters may have spurious edges connecting to other parts of

the manifold only a few hops away. To avoid integrating neighborhood information outside

this local structure, we use the Leiden algorithm (Traag et al., 2019) to cluster the graph

and identify a local neighborhood size for each cell (the resolution parameter is set to 3

by default). If cell a belongs to cluster c_a, then its neighborhood size is $l_a = |c_a|$. For each

row a in $\bar{N}_i$ we only keep the l_a geodesically closest cells, letting the pruned graph be

denoted as $\widehat{N}_i$.

Edges outgoing from cell a_i in species i are encoded in the corresponding row in the

adjacency matrix: C_{i,a_i}. We compute the fraction of the outgoing edges from each cell that

target the local neighborhood of a cell in the other species: $\tilde{C}_{i,a_ib_j} = \sum_{c \in X_{j,b_j}} C_{i,a_ic}$, where

X_{j,b_j} is the set of cells in the neighborhood of cell b_j in species j and $\tilde{C}_{i,a_ib_j}$ is the fraction

of outgoing edges from cell a_i in species i targeting the neighborhood of cell b_j in species

j.

To reduce the density of $\tilde{C}_i$ so as to satisfy computational memory constraints, we remove

edges with weight less than 0.1. Finally, we apply the mutual nearest neighborhood

criterion by taking the element-wise, geometric mean of the two directed bipartite graphs:

$\tilde{C} = \sqrt{\tilde{C}_1 \circ \tilde{C}_2}$. This operation ensures that only bidirectional edges are preserved, as small

edge weights in either direction results in small geometric means.

2.5. Assign the k-nearest cross-species neighborhoods for each cell and update edge

weights in the gene homology graph.

Given the mutual nearest neighborhoods $\tilde{C} \in \mathfrak{R}^{n_1 \times n_2}$, we select the k nearest

neighborhoods for each cell in both directions to update the directed biadjacency matrices

C_1 and C_2: $C_1 = KNN(\tilde{C}, k)$ and $C_2 = KNN(\tilde{C}^T, k)$, with $k = 20$ by default.

2.6. Stitch the manifolds.

We use C_1 and C_2 to combine the manifolds N_1 and N_2 into a unified graph. We first weight

the edges in N_1 and N_2 to account for the number of shared cross-species neighbors by

computing the one-mode projections of C_1 and C_2. In addition, for cells with strong cross-

species alignment, we attenuate the weight of their within-species edges. For cells with

little to no cross-species alignment, their within-species are kept the same to ensure that

the local topological information around cells with no alignment is preserved.

Specifically, we use N_1 and N_2 to mask the edges in the one-mode projections, $\widetilde{N}_1 =$

$U(N_1) \circ (Norm(C_1)Norm(C_2))$ and $\widetilde{N}_2 = U(N_2) \circ (Norm(C_2)Norm(C_1))$, where $U(E)$ sets

all edge weights in graph E to 1 and $Norm$ normalizes the outgoing edges from each cell

to sum to 1. The minimum edge weight is set to be 0.3 to ensure that neighbors in the

original manifolds with no shared cross-species neighbors still retain connectivity: $\widetilde{N}_{1,ij} =$

$min(0.3, \widetilde{N}_{1,ij})$ and $\widetilde{N}_{2,ij} = min(0.3, \widetilde{N}_{2,ij})$ for all edges (i,j). We then scale the within-

species edges from cell i by the total weight of its cross-species edges: $\widetilde{N}_{1,i} = (1 -$

$\frac{1}{k}\sum_{j=1}^{n_2} C_{1,ij})\widetilde{N}_{1,i}$ and $\widetilde{N}_{2,i} = (1 - \frac{1}{k}\sum_{j=1}^{n_1} C_{2,ij})\widetilde{N}_{2,i}$. Finally, the within- and cross-species

graphs are stitched together to form the combined nearest neighbor graph N: $N = [\widetilde{N}_1 \oplus$

$C_1] \oplus [C_2 \oplus \widetilde{N}_2]$. The overall alignment score between species 1 and 2 is defined as $S =$

$\frac{1}{n_1+n_2}(\sum_{i=1}^{n_1}\sum_{j=1}^{n_2} C_{1,ij} + \sum_{i=1}^{n_2}\sum_{j=1}^{n_1} C_{2,ij})$.

3. Gene-gene correlation initialization.

3.1. Update edge weights in the gene-gene bipartite graph with expression correlations.

To compute correlations between gene pairs, we first transfer expressions from one

species to the other: $\bar{Z}_{i,n_i m_j} = C_{i,n_i} Z_{j,m_j}$, where $\bar{Z}_{i,n_i m_j}$ is the imputed expressions of gene

m_j from species j for cell n_i in species i, and C_{i,n_i} is row n_i of the biadjacency matrix

encoding the cross-species neighbors of cell n_i in species i, all for $(i,j) = (1,2)$ and $(2,1)$.

We similarly use the manifolds constructed by SAM to smooth the within-species gene

expressions using kNN averaging: $\bar{Z}_{j,m_j} = N_{j,m_j} Z_{j,m_j}$, where N_j is the nearest-neighbor

graph for species j. We then concatenate the within- and cross-species gene expressions

such that the expression of gene m_j from species j in both species is $\bar{Z}_{m_j} = \bar{Z}_{i,m_j} \oplus \bar{Z}_{j,m_j}$.

For all gene pairs in the unpruned homology graph generated in step 1.1., $\hat{G}$, we compute

their correlations, $\hat{G}_{ab} := \theta(0)Corr(\bar{Z}_a, \bar{Z}_b)$, where $\theta(0)$ is a Heaviside step function

centered at 0 to set negative correlations to zero. We then use the expression correlations

to update the corresponding edge weights in $\hat{G}$, which are again normalized through

$\hat{G}_{ab} = 0.5 + 0.5\,tanh\,(10\hat{G}_{ab}/\max_{b}(\hat{G}_{ab}) - 5).$

Annotation of cell atlases

To annotate the primary zebrafish and *Xenopus* cell types, the cell subtype annotations

provided by the original publications (Briggs et al., 2018; Wagner et al., 2018) are

coarsened using a combination of the manual matching and developmental hierarchies.

For example, as "heart - mature", "heart - hoxd9a", "heart", and "heart field" in zebrafish

are all manually matched to "cardiac mesoderm" in *Xenopus*, we label these cells as

"heart". In cases where the matching is insufficient to coarsen the annotations, we use

the provided developmental trees to name a group of terminal cell subtypes by their

common ontogenic ancestor. The annotations provided by their respective studies were

used to label the cells in the *Spongilla*, *Hydra*, and mouse atlases. To annotate the

schistosome cells, we used known marker genes to annotate the main schistosome tissue

types (Li et al., 2020). Annotations for all single cells in all datasets are provided in

Supplementary Table 1.

Visualization

The combined manifold N is embedded into 2D projections using UMAP implemented in

the scanpy package (Wolf et al., 2018) by *scanpy.tl.umap* with the parameter *min_dist* =

0.1. The sankeyD3 package (https://rdrr.io/github/fbreitwieser/sankeyD3/man/sankeyD3-

package.html) in R is used to generate the sankey plots. Edge thickness corresponds to

the alignment score between mapped cell types. The alignment score between cell types

a and b is defined as $s_{ab} = \frac{1}{|c_a|+|c_b|}(\sum_{i \in c_a} \sum_{j \in c_b} C_{1,ij} + \sum_{i \in c_b} \sum_{j \in c_a} C_{2,ij})$, where c_a and c_b

are the set of cells in cell types a and b, respectively. Cell type pairs with alignment score

less than z are filtered out. By default, z is set to be 0.1. Cell types that did not cluster

properly in their respective manifolds were omitted from the sankey plot. In the zebrafish-

Xenopus comparison, we excluded heart, germline, and olfactory placode cells from both

species because they did not cluster in the *Xenopus* atlas. Similarly, the iridoblast,

epiphysis, *nanog+*, apoptotic-like, and forerunner cells were excluded because they did

not cluster in the zebrafish atlas.

The network graphs in **Figure 4D** are generated using the *networkx* package

(https://networkx.github.io) in python. To focus on densely connected cell type groups,

we filter out cell type pairs with alignment score less than 0.05.

Identification of gene pairs that drive cell type mappings

We define g_1 and g_2 to contain SAMap-linked genes from species 1 and 2, respectively.

Note that a gene may appear multiple times as SAMap allows for one-to-many homology.

Let $X_{a_1 b_2}$ denote the set of all cells with cross species edges between cell types a_1 and

b_2. We calculate the average standardized expression of all cells from species i that are

in $X_{a_1 b_2}$: $Y_{i,g_i} = \frac{1}{|\{x, x \in X_{a_1 b_2}\}|} \sum_{x \in X_{a_1 b_2}} \tilde{Z}_{i,x,g_i}$, where $\tilde{Z}_{i,x,g_i} \in \Re^{|g_i|}$ is the standardized

expression of genes g_i in cell x. The correlation between Y_{1,g_1} and Y_{2,g_2} can be written as

$Corr(Y_{1,g_1}, Y_{2,g_2}) = \sum_{j=1}^{|g_1|} S(Y_{1,g_1})_j \circ S(Y_{2,g_2})_j$, where $S(Z)$ standardizes vector Z to have

zero mean and unit variance. We use the summand to identify gene pairs that contribute

most positively to the correlation. We assign each gene pair a score: $h_g = T(S(Y_{1,g_1})) \circ$

$T(S(Y_{2,g_2}))$, where $T(Z)$ sets negative values in vector Z to zero in order to ignore lowly-

expressed genes. To be inclusive, we begin with the top 1,000 gene pairs according to

h_g and filter out gene pairs in which one or both of the genes are not differentially

expressed in their respective cell types (p-value $> 10^{-2}$), have less than 0.2 SAM weight,

or are expressed in fewer than 5% of the cells in the cluster. The differential expression

of each gene in each cell type is calculated using the Wilcoxon rank-sum test

implemented in the *scanpy* function *scanpy.tl.rank_genes_groups*.

Orthology group assignment

We used the Eggnog mapper (v5.0) (Huerta-Cepas et al., 2019) to assign each gene to

an orthology group with default parameters. For the zebrafish-to-*Xenopus* mapping,

genes are considered paralogs if they map to the same eukaryotic orthology group and

orthologs if they map to the same vertebrate orthology group. For the pan-species

analysis, we group genes from all species with overlapping orthology assignments. In

Figure 5B, each column corresponds to one of these groups. As each group may contain

multiple genes from each species, we present the expression of the gene with the highest

enrichment score per species. All gene names and corresponding orthology groups are

reported in **Supplementary Table 4**.

Phylogenetic reconstruction of gene trees

We generate gene trees to validate the identity of genes involved in putative examples of

paralog substitution and of *Fox* and *Csrp* transcriptional regulators that are identified as

enriched in contractile cells. For this, we first gather protein sequences from potential

homologs using the eggnog version 5.0 orthology database (Huerta-Cepas et al., 2019).

For the *Fox* and *Csrp* phylogenies, we include all Fox clade I (Larroux et al., 2008) and

Csrp/Crip homologs, respectively, from the seven species included in our study.

Alignment of protein sequences is performed with Clustal Omega version 1.2.4 using

default settings as implemented on the EMBL EBI web services platform (Madeira et al.,

2019). Maximum likelihood tree reconstruction is performed using IQ-TREE version

1.6.12 (Nguyen et al., 2015) with the ModelFinder Plus option (Kalyaanamoorthy et al.,

2017). For the *Csrp* tree, we perform 1,000 nonparametric bootstrap replicates to assess

node support. For *Fox*, we utilize the ultrafast bootstrap support option with 1,000

replicates. For each gene tree we choose the model that minimizes the Bayesian

Information Criterion (BIC) score in ModelFinder. This results in selection of the following

models: DCMut+R4 (*Csrp*) and VT+F+R5 (*Fox*). The final consensus trees are visualized

and rendered using the ete3 v3.1.1 python toolkit (Huerta-Cepas et al., 2016) and the

Interactive Tree of Life v4 (Letunic & Bork, 2019).

KOG functional annotation and enrichment analysis

Using the eggnog mapper, KOG functional annotations are transferred to individual

transcripts from their assigned orthology group. For enrichment analysis, all genes

enriched in the set of cell type pairs of interest are lumped to form the target set for each

species. For example, the target set for *Spongilla* archaeocytes used in **Figure 5C** is

composed of all genes enriched between *Spongilla* archaeocytes and other invertebrate

stem cells. Note that this set includes genes from other species that are linked by SAMap

to the *Spongilla* archeocyte genes. We include genes from other species in the target set

to account for differences in KOG functional annotation coverage between species. As

such, the annotated transcripts from all 7 species are combined to form the background

set. We used a hypergeometric statistical test (Eden et al., 2009) to measure the

enrichment of the KOG terms in the target genes compared to the background genes.

Mapping zebrafish and xenopus atlases using existing methods

For benchmarking, we used vertebrate orthologs as determined by Eggnog as input to

Harmony (Korsunsky et al., 2019), Liger (Welch et al., 2019), Seurat (Stuart et al., 2019),

Scanorama (Hie et al., 2019), BBKNN (Polański et al., 2019), which are all run with default

parameters. One-to-one orthologs were selected from one-to-many and many-to-many

orthologs by using the bipartite maximum weight matching algorithm implemented in

networkx. When using the one-to-one orthologs as input for SAMap, we ran for only one

iteration. The resulting integrated lower-dimensional coordinates (PCs for Seurat,

Harmony, and Scanorama and non-negative matrix factorization coordinates for Liger)

and stitched graph (BBKNN and SAMap) were all projected into 2D with UMAP (**Figure**

2 – figure supplement 2A). The integrated coordinates are used to generate a nearest

neighbor graph using the correlation distance metric, which is then used to compute the

alignment scores in **Figure 2 – figure supplement 2B**. The alignment scores for SAMap

and BBKNN are directly computed from their combined graphs.

In-situ hybridization in schistosomes

S. mansoni (strain: NMRI) juveniles are retrieved from infected female Swiss Webster

mice (NR-21963) at ~3 weeks post-infection by hepatic portal vein perfusion using 37°C

DMEM supplemented with 5 % heat inactivated FBS. The infected mice are provided by

the NIAID Schistosomiasis Resource Center for distribution through BEI Resources, NIH-

NIAID Contract HHSN272201000005I. In adherence to the Animal Welfare Act and the

Public Health Service Policy on Humane Care and Use of Laboratory Animals, all

experiments with and care of mice are performed in accordance with protocols approved

by the Institutional Animal Care and Use Committees (IACUC) of Stanford University

(protocol approval number 30366). *In situ* hybridization experiments are performed as

described previously (Tarashansky et al., 2019), using riboprobes synthesized from gene

fragments cloned with the listed primers: collagen (Smp_170340):

GGTGAAGAAGGCTGTTGTGG, ACGATCCCCTTTCACTCCTG; tropomyosin

(Smp_031770): AAGCTGAAGTCGCCTCACTA, CATATGCCTCTTCACGCTGG;

troponin (Smp_018250): CGTAAACCTGGTCAGAAGCG,

ATCCTTTTCCTCCAGAGCGT; myosin regulatory light chain (Smp_132670):

GAGACAGCGAGTAGTGGAGG, TGCCTTCTTTGATTGGAGCT; wnt (Smp_156540):

TGTGGTGATGAAGATGGCAG, CCACGGCCACAACACATATT; frizzled

(Smp_174350): CGAACAGGCGCATGACAATA, TGCTAGTCCTGTTGTCGTGT.

Acknowledgments

We thank D. Wagner and C. Juliano for sharing the data and essential discussions. We

also thank S. Granick, L. Luo, and J. Kebschull for their critical reading of the manuscript.

AJT is a Bio-X Stanford Interdisciplinary Graduate Fellow. This work is supported by a

Beckman Young Investigator Award and an NIH grant (1R35GM138061) to BW.

References

Alié, A., Hayashi, T., Sugimura, I., Manuel, M., Sugano, W., Mano, A., Satoh, N., Agata, K., & Funayama, N. (2015). The ancestral gene repertoire of animal stem cells. *Proceedings of the National Academy of Sciences*, *112*(51), E7093–E7100. https://doi.org/10.1073/pnas.1514789112

Arendt, D., Bertucci, P. Y., Achim, K., & Musser, J. M. (2019). Evolution of neuronal types and families. *Current Opinion in Neurobiology*, *56*, 144–152. https://doi.org/10.1016/j.conb.2019.01.022

Arendt, D., Musser, J. M., Baker, C. V. H., Bergman, A., Cepko, C., Erwin, D. H., Pavlicev, M., Schlosser, G., Widder, S., Laubichler, M. D., & Wagner, G. P. (2016). The origin and evolution of cell types. *Nature Reviews Genetics*, *17*(12), 744–757. https://doi.org/10.1038/nrg.2016.127

Barkas, N., Petukhov, V., Nikolaeva, D., Lozinsky, Y., Demharter, S., Khodosevich, K., & Kharchenko, P. V. (2019). Joint analysis of heterogeneous single-cell RNA-seq dataset collections. *Nature Methods*, *16*(8), 695–698. https://doi.org/10.1038/s41592-019-0466-z

Baron, M., Veres, A., Wolock, S. L., Faust, A. L., Gaujoux, R., Vetere, A., Ryu, J. H., Wagner, B. K., Shen-Orr, S. S., Klein, A. M., Melton, D. A., & Yanai, I. (2016). A single-cell transcriptomic map of the human and mouse pancreas reveals inter- and intra-cell population structure. *Cell Systems*, *3*(4), 346-360.e4. https://doi.org/10.1016/j.cels.2016.08.011

Briggs, J. A., Weinreb, C., Wagner, D. E., Megason, S., Peshkin, L., Kirschner, M. W., & Klein, A. M. (2018). The dynamics of gene expression in vertebrate embryogenesis at single-cell resolution. *Science*, *360*(6392), eaar5780. https://doi.org/10.1126/science.aar5780

Brunet, T., Fischer, A. H., Steinmetz, P. R., Lauri, A., Bertucci, P., & Arendt, D. (2016). The evolutionary origin of bilaterian smooth and striated myocytes. *eLife*, *5*, e19607. https://doi.org/10.7554/eLife.19607

Buzgariu, W., Al Haddad, S., Tomczyk, S., Wenger, Y., & Galliot, B. (2015). Multi-functionality and plasticity characterize epithelial cells in *Hydra*. *Tissue Barriers*, *3*(4), e1068908. https://doi.org/10.1080/21688370.2015.1068908

Cao, C., Lemaire, L. A., Wang, W., Yoon, P. H., Choi, Y. A., Parsons, L. R., Matese, J. C., Wang, W., Levine, M., & Chen, K. (2019). Comprehensive single-cell transcriptome lineages of a proto-vertebrate. *Nature*, *571*(7765), 349–354. https://doi.org/10.1038/s41586-019-1385-y

Dubaissi, E., Rousseau, K., Lea, R., Soto, X., Nardeosingh, S., Schweickert, A., Amaya, E., Thornton, D. J., & Papalopulu, N. (2014). A secretory cell type develops alongside multiciliated cells, ionocytes and goblet cells, and provides a protective, anti-infective function in the frog embryonic mucociliary epidermis. *Development*, *141*(7), 1514–1525. https://doi.org/10.1242/dev.102426

Eddy, S. R. (2008). A probabilistic model of local sequence alignment that simplifies statistical significance estimation. *PLoS Computational Biology*, *4*(5), e1000069. https://doi.org/10.1371/journal.pcbi.1000069

Eden, E., Navon, R., Steinfeld, I., Lipson, D., & Yakhini, Z. (2009). GOrilla: A tool for discovery and visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics*, *10*(1), 48. https://doi.org/10.1186/1471-2105-10-48

El-Brolosy, M. A., Kontarakis, Z., Rossi, A., Kuenne, C., Günther, S., Fukuda, N., Kikhi, K., Boezio, G. L. M., Takacs, C. M., Lai, S.-L., Fukuda, R., Gerri, C., Giraldez, A. J., & Stainier, D. Y. R. (2019). Genetic compensation triggered by mutant mRNA degradation. *Nature*, *568*(7751), 193–197. https://doi.org/10.1038/s41586-019-1064-z

Fincher, C. T., Wurtzel, O., de Hoog, T., Kravarik, K. M., & Reddien, P. W. (2018). Cell type
transcriptome atlas for the planarian *Schmidtea mediterranea*. *Science*, *360*(6391),
eaaq1736. https://doi.org/10.1126/science.aaq1736
Gabaldón, T., & Koonin, E. V. (2013). Functional and evolutionary implications of gene
orthology. *Nature Reviews Genetics*, *14*(5), 360–366. https://doi.org/10.1038/nrg3456
Geirsdottir, L., David, E., Keren-Shaul, H., Weiner, A., Bohlen, S. C., Neuber, J., Balic, A.,
Giladi, A., Sheban, F., Dutertre, C.-A., Pfeifle, C., Peri, F., Raffo-Romero, A., Vizioli, J.,
Matiasek, K., Scheiwe, C., Meckel, S., Mätz-Rensing, K., van der Meer, F., … Prinz, M.
(2019). Cross-Species Single-Cell Analysis Reveals Divergence of the Primate Microglia
Program. *Cell*, *179*(7), 1609-1622.e16. https://doi.org/10.1016/j.cell.2019.11.010
Haghverdi, L., Lun, A. T. L., Morgan, M. D., & Marioni, J. C. (2018). Batch effects in single-cell
RNA-sequencing data are corrected by matching mutual nearest neighbors. *Nature
Biotechnology*, *36*(5), 421–427. https://doi.org/10.1038/nbt.4091
Hie, B., Bryson, B., & Berger, B. (2019). Efficient integration of heterogeneous single-cell
transcriptomes using Scanorama. *Nature Biotechnology*, *37*(6), 685–691.
https://doi.org/10.1038/s41587-019-0113-3
Hu, M., Zheng, X., Fan, C.-M., & Zheng, Y. (2020). Lineage dynamics of the endosymbiotic cell
type in the soft coral Xenia. *Nature*, *582*(7813), 534–538. https://doi.org/10.1038/s41586-
020-2385-7
Huerta-Cepas, J., Serra, F., & Bork, P. (2016). ETE 3: Reconstruction, Analysis, and
Visualization of Phylogenomic Data. *Molecular Biology and Evolution*, *33*(6), 1635–1638.
https://doi.org/10.1093/molbev/msw046
Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K., Cook, H.,
Mende, D. R., Letunic, I., Rattei, T., Jensen, L. J., von Mering, C., & Bork, P. (2019).
eggNOG 5.0: A hierarchical, functionally and phylogenetically annotated orthology resource
based on 5090 organisms and 2502 viruses. *Nucleic Acids Research*, *47*(D1), D309–D314.
https://doi.org/10.1093/nar/gky1085
Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., & Jermiin, L. S. (2017).
ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods*,
14(6), 587–589. https://doi.org/10.1038/nmeth.4285
Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner,
M., Loh, P., & Raychaudhuri, S. (2019). Fast, sensitive and accurate integration of single-
cell data with Harmony. *Nature Methods*, *16*(12), 1289–1296.
https://doi.org/10.1038/s41592-019-0619-0
Larroux, C., Luke, G. N., Koopman, P., Rokhsar, D. S., Shimeld, S. M., & Degnan, B. M. (2008).
Genesis and Expansion of Metazoan Transcription Factor Gene Classes. *Molecular Biology
and Evolution*, *25*(5), 980–996. https://doi.org/10.1093/molbev/msn047
Laumer, C. E., Hejnol, A., & Giribet, G. (2015). Nuclear genomic signals of the
'microturbellarian' roots of platyhelminth evolutionary innovation. *eLife*, *4*, e05503.
https://doi.org/10.7554/eLife.05503
Letunic, I., & Bork, P. (2019). Interactive Tree Of Life (iTOL) v4: Recent updates and new
developments. *Nucleic Acids Research*, *47*(W1), W256–W259.
https://doi.org/10.1093/nar/gkz239
Li, P., Sarfati, D. N., Xue, Y., Yu, X., Tarashansky, A. J., Quake, S. R., & Wang, B. (2020).
Single-cell analysis of Schistosoma mansoni *reveals a conserved genetic program
controlling germline stem cell fate* [Preprint]. https://doi.org/10.1101/2020.07.06.190033
Littlewood, D. T. J., & Waeschenbach, A. (2015). Evolution: A Turn Up for the Worms. *Current
Biology*, *25*(11), R457–R460. https://doi.org/10.1016/j.cub.2015.04.012
Madeira, F., Park, Y. mi, Lee, J., Buso, N., Gur, T., Madhusoodanan, N., Basutkar, P., Tivey, A.
R. N., Potter, S. C., Finn, R. D., & Lopez, R. (2019). The EMBL-EBI search and sequence

759  analysis tools APIs in 2019. *Nucleic Acids Research*, *47*(W1), W636–W641.
760  https://doi.org/10.1093/nar/gkz268
761 Malkov, Y. A., & Yashunin, D. A. (2020). Efficient and robust approximate nearest neighbor
762  search using hierarchical navigable small world graphs. *IEEE Transactions on Pattern*
763  *Analysis and Machine Intelligence*, *42*(4), 824–836.
764  https://doi.org/10.1109/TPAMI.2018.2889473
765 Miles, L. B., Darido, C., Kaslin, J., Heath, J. K., Jane, S. M., & Dworkin, S. (2017). Mis-
766  expression of grainyhead-like transcription factors in zebrafish leads to defects in
767  enveloping layer (EVL) integrity, cellular morphogenesis and axial extension. *Scientific*
768  *Reports*, *7*(1), 17607. https://doi.org/10.1038/s41598-017-17898-7
769 Musser, J. M., Schippers, K. J., Nickel, M., Mizzon, G., Kohn, A. B., Pape, C., Hammel, J. U.,
770  Wolf, F., Liang, C., Hernández-Plaza, A., Achim, K., Schieber, N. L., Francis, W. R., Vargas
771  R., S., Kling, S., Renkert, M., Feuda, R., Gaspar, I., Burkhardt, P., … Arendt, D. (2019).
772  *Profiling cellular diversity in sponges informs animal cell type and nervous system evolution*
773  [Preprint]. https://doi.org/10.1101/758276
774 Nehrt, N. L., Clark, W. T., Radivojac, P., & Hahn, M. W. (2011). Testing the ortholog conjecture
775  with comparative functional genomic data from mammals. *PLoS Computational Biology*,
776  *7*(6), e1002073. https://doi.org/10.1371/journal.pcbi.1002073
777 Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: A fast and
778  effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular*
779  *Biology and Evolution*, *32*(1), 268–274. https://doi.org/10.1093/molbev/msu300
780 Pijuan-Sala, B., Griffiths, J. A., Guibentif, C., Hiscock, T. W., Jawaid, W., Calero-Nieto, F. J.,
781  Mulas, C., Ibarra-Soria, X., Tyser, R. C. V., Ho, D. L. L., Reik, W., Srinivas, S., Simons, B.
782  D., Nichols, J., Marioni, J. C., & Göttgens, B. (2019). A single-cell molecular map of mouse
783  gastrulation and early organogenesis. *Nature*, *566*(7745), 490–495.
784  https://doi.org/10.1038/s41586-019-0933-9
785 Plass, M., Solana, J., Wolf, F. A., Ayoub, S., Misios, A., Glažar, P., Obermayer, B., Theis, F. J.,
786  Kocks, C., & Rajewsky, N. (2018). Cell type atlas and lineage tree of a whole complex
787  animal by single-cell transcriptomics. *Science*, *360*(6391), eaaq1723.
788  https://doi.org/10.1126/science.aaq1723
789 Polański, K., Young, M. D., Miao, Z., Meyer, K. B., Teichmann, S. A., & Park, J.-E. (2019).
790  BBKNN: Fast batch alignment of single cell transcriptomes. *Bioinformatics*, btz625.
791  https://doi.org/10.1093/bioinformatics/btz625
792 Prince, V. E., & Pickett, F. B. (2002). Splitting pairs: The diverging fates of duplicated genes.
793  *Nature Reviews Genetics*, *3*(11), 827–837. https://doi.org/10.1038/nrg928
794 Reddien, P. W. (2018). The Cellular and Molecular Basis for Planarian Regeneration. *Cell*,
795  *175*(2), 327–345. https://doi.org/10.1016/j.cell.2018.09.021
796 Regev, A., Teichmann, S. A., Lander, E. S., Amit, I., Benoist, C., Birney, E., Bodenmiller, B.,
797  Campbell, P., Carninci, P., Clatworthy, M., Clevers, H., Deplancke, B., Dunham, I.,
798  Eberwine, J., Eils, R., Enard, W., Farmer, A., Fugger, L., Göttgens, B., … Human Cell Atlas
799  Meeting Participants. (2017). The Human Cell Atlas. *eLife*, *6*, e27041.
800  https://doi.org/10.7554/eLife.27041
801 Scimone, M. L., Cote, L. E., & Reddien, P. W. (2017). Orthogonal muscle fibres have different
802  instructive roles in planarian regeneration. *Nature*, *551*(7682), 623–628.
803  https://doi.org/10.1038/nature24660
804 Sebé-Pedrós, A., Chomsky, E., Pang, K., Lara-Astiaso, D., Gaiti, F., Mukamel, Z., Amit, I.,
805  Hejnol, A., Degnan, B. M., & Tanay, A. (2018). Early metazoan cell type diversity and the
806  evolution of multicellular gene regulation. *Nature Ecology & Evolution*, *2*(7), 1176–1188.
807  https://doi.org/10.1038/s41559-018-0575-6
808 Shafer, M. E. R. (2019). Cross-Species Analysis of Single-Cell Transcriptomic Data. *Frontiers in*
809  *Cell and Developmental Biology*, *7*, 175. https://doi.org/10.3389/fcell.2019.00175

Siebert, S., Farrell, J. A., Cazet, J. F., Abeykoon, Y., Primack, A. S., Schnitzler, C. E., & Juliano,
C. E. (2019). Stem cell differentiation trajectories in *Hydra* resolved at single-cell resolution.
Science, *365*(6451), eaav9314. https://doi.org/10.1126/science.aav9314
Stamboulian, M., Guerrero, R. F., Hahn, M. W., & Radivojac, P. (2020). The ortholog conjecture
revisited: The value of orthologs and paralogs in function prediction. *Bioinformatics*,
36(Supplement_1), i219–i226. https://doi.org/10.1093/bioinformatics/btaa468
Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., Hao, Y.,
Stoeckius, M., Smibert, P., & Satija, R. (2019). Comprehensive Integration of Single-Cell
Data. *Cell*, *177*(7), 1888-1902.e21. https://doi.org/10.1016/j.cell.2019.05.031
Studer, R. A., & Robinson-Rechavi, M. (2009). How confident can we be that orthologs are
similar, but paralogs differ? *Trends in Genetics*, *25*(5), 210–216.
https://doi.org/10.1016/j.tig.2009.03.004
Tarashansky, A. J., Xue, Y., Li, P., Quake, S. R., & Wang, B. (2019). Self-assembling manifolds
in single-cell RNA sequencing data. *eLife*, *8*, e48994. https://doi.org/10.7554/eLife.48994
Tatusov, R. L., Fedorova, N. D., Jackson, J. D., Jacobs, A. R., Kiryutin, B., Koonin, E. V.,
Krylov, D. M., Mazumder, R., Mekhedov, S. L., Nikolskaya, A. N., Rao, B. S., Smirnov, S.,
Sverdlov, A. V., Vasudevan, S., Wolf, Y. I., Yin, J. J., & Natale, D. A. (2003). The COG
database: An updated version includes eukaryotes. *BMC Bioinformatics*, 14.
Tosches, M. A., & Arendt, D. (2013). The bilaterian forebrain: An evolutionary chimaera. *Current
Opinion in Neurobiology*, *23*(6), 1080–1089. https://doi.org/10.1016/j.conb.2013.09.005
Tosches, M. A., Yamawaki, T. M., Naumann, R. K., Jacobi, A. A., Tushev, G., & Laurent, G.
(2018). Evolution of pallium, hippocampus, and cortical cell types revealed by single-cell
transcriptomics in reptiles. *Science*, *360*(6391), 881–888.
https://doi.org/10.1126/science.aar4237
Traag, V. A., Waltman, L., & van Eck, N. J. (2019). From Louvain to Leiden: Guaranteeing well-
connected communities. *Scientific Reports*, *9*(1), 5233. https://doi.org/10.1038/s41598-019-
41695-z
Wagner, D. E., Weinreb, C., Collins, Z. M., Briggs, J. A., Megason, S., & Klein, A. M. (2018).
Single-cell mapping of gene expression landscapes and lineage in the zebrafish embryo.
Science, *360*(6392), 981–987.
Wang, B., Lee, J., Li, P., Saberi, A., Yang, H., Liu, C., Zhao, M., & Newmark, P. A. (2018). Stem
cell heterogeneity drives the parasitic life cycle of Schistosoma mansoni. *eLife*, 7, e35449.
https://doi.org/10.7554/eLife.35449
Weir, K., Dupre, C., van Giesen, L., Lee, A. S.-Y., & Bellono, N. W. (2020). A molecular filter for
the cnidarian stinging response. *eLife*, *9*, e57578. https://doi.org/10.7554/eLife.57578
Welch, J. D., Kozareva, V., Ferreira, A., Vanderburg, C., Martin, C., & Macosko, E. Z. (2019).
Single-Cell Multi-omic Integration Compares and Contrasts Features of Brain Cell Identity.
Cell, *177*(7), 1873-1887.e17. https://doi.org/10.1016/j.cell.2019.05.006
Wendt, G. R., & Collins, J. J. (2016). Schistosomiasis as a disease of stem cells. *Current
Opinion in Genetics & Development*, *40*, 95–102. https://doi.org/10.1016/j.gde.2016.06.010
Wendt, G. R., Collins, J. N., Pei, J., Pearson, M. S., Bennett, H. M., Loukas, A., Berriman, M.,
Grishin, N. V., & Collins, J. J. (2018). Flatworm-specific transcriptional regulators promote
the specification of tegumental progenitors in Schistosoma mansoni. *eLife*, *7*, e33221.
https://doi.org/10.7554/eLife.33221
Wolf, F. A., Angerer, P., & Theis, F. J. (2018). SCANPY: Large-scale single-cell gene
expression data analysis. *Genome Biology*, *19*(1), 15. https://doi.org/10.1186/s13059-017-
1382-0
Wong, E., Mölter, J., Anggono, V., Degnan, S. M., & Degnan, B. M. (2019). Co-expression of
synaptic genes in the sponge Amphimedon queenslandica uncovers ancient neural
submodules. *Scientific Reports*, *9*(1), 15781. https://doi.org/10.1038/s41598-019-51282-x

Yan, K.-K., Wang, D., Rozowsky, J., Zheng, H., Cheng, C., & Gerstein, M. (2014). OrthoClust:
An orthology-based network framework for clustering data across multiple species. *Genome*
Biology, *15*(8), R100. https://doi.org/10.1186/gb-2014-15-8-r100
Zeng, A., Li, H., Guo, L., Gao, X., McKinney, S., Wang, Y., Yu, Z., Park, J., Semerad, C., Ross,
E., Cheng, L.-C., Davies, E., Lei, K., Wang, W., Perera, A., Hall, K., Peak, A., Box, A., &
Sánchez Alvarado, A. (2018). Prospectively Isolated Tetraspanin+ Neoblasts Are Adult
Pluripotent Stem Cells Underlying Planaria Regeneration. *Cell*, *173*(7), 1593-1608.e20.
https://doi.org/10.1016/j.cell.2018.05.006

Figures

Figure 1: SAMap addresses challenges in mapping cell atlases of distantly related

species. (A) Schematic showing the phylogenetic relationships among 7 species

analyzed. (B) Challenges in mapping single-cell transcriptomes. Gene duplications cause

large numbers of homologs per gene, determined by reciprocal BLAST (cut-off: e-value

$< 10^{-6}$), and frequent gene losses and the acquisition of new genes results in large

fractions of transcriptomes lacking homology, which limits the amount of information

comparable across species. (C) SAMap workflow. Homologous gene pairs initially

weighted by protein sequence similarity are used to align the manifolds, low dimensional

representations of the cell atlases. Gene-gene correlations calculated from the aligned

manifolds are used to update the edge weights in the bipartite graph, which are then used

to improve manifold alignment. (D) Mutual nearest neighborhoods improve the detection

of cross-species mutual nearest neighbors by connecting cells that target one other's

within-species neighborhoods. (E) Convergence of SAMap is evaluated by the root mean

square error (RMSE) of the alignment scores between mapped clusters in adjacent

iterations for all 21 pairwise comparisons of the 7 species.

Figure 2: SAMap successfully maps *D. rerio* and *X. tropicalis* atlases and reveals prevalent paralog substitutions. (A) UMAP projection of the combined zebrafish (yellow) and *Xenopus* (blue) manifolds, with example cell types circled. (B) Sankey plot summarizing the cell type mappings. Edges with alignment score < 0.1 are omitted. Edges that connect developmentally distinct secretory cell types are highlighted in black. (C)

Heatmaps of alignment scores between developmental time points for ionocyte,

forebrain/midbrain, placodal, and neural crest lineages. X-axis: zebrafish. Y-axis:

Xenopus. (D) SAMap alignment scores compared to those of benchmarking methods

using one-to-one vertebrate orthologs as input. Each dot represents a cell type pair

supported by ontogeny annotations. (E) Expression of orthologous (left) and paralogous

(right) gene pairs overlaid on the combined UMAP projection. Expressing cells are color-

coded by species, with those that are connected across species colored cyan. Cells with

no expression are shown in gray. More examples are provided in **Figure 2 – figure**

supplement 3.

Figure 3: SAMap transfers cell type information from a well-annotated organism (planarian *S. mediterranea*) to its less-studied cousin (schistosome *S. mansoni*) and identifies parallel stem cell compartments. (A) UMAP projection of the combined manifolds. Tissue type annotations are adopted from the *S. mediterranea* atlas (Fincher et al., 2018). The schistosome atlas was collected from juvenile worms, which we found to contain neoblasts with an abundance comparable to that of planarian neoblasts (Li et al., 2020). (B) Overlapping expressions of selected tissue-specific TFs with expressing cell types circled. (C) UMAP projection of the aligned manifolds showing planarian and schistosome neoblasts, with homologous subpopulations circled. Planarian neoblast data

912   is from (Zeng et al., 2018), and cNeoblasts correspond to the Nb2 population, which are

913   pluripotent cells that can rescue neoblast-depleted planarians in transplantation

914   experiments. (D) Distributions of conserved TF expressions in each neoblast

915   subpopulation. Expression values are *k*-nearest-neighbor averaged and standardized,

916   with negative values set to zero. Blue: planarian; yellow: schistosome.

Figure 4: Mapping evolutionarily distant species identifies densely connected cell type groups. (A) Violin plots showing the number of enriched gene pairs in cell type mappings from all 21 pairwise mappings between the 7 species. 87% of cell type mappings have greater than 40 enriched gene pairs (dotted line). Species acronyms are

the same as in **Figure 1A**. (B) Schematic illustrating edge (left) and node (right)

transitivities, defined as the fraction of triads (set of three connected nodes) in closed

triangles. (C) The percentage of cell type pairs that are topologically equivalent to the

green edge in each illustrated motif. (D) Network graphs showing highly connected cell

type families. Each node represents a cell type, color-coded by species (detailed

annotations are provided in **Supplementary Table 5**). Mapped cell types are connected

with an edge. (E) Boxplot showing the median and interquartile ranges of node

transitivities for highly connected cell type groups. For all box plots, the whiskers denote

the maximum and minimum observations. The average node transitivity per group is

compared to a bootstrapped null transitivity distribution, generated by repeatedly

sampling subsets of nodes in the cell type graph and calculating their transitivities. *p <

5×10^{-3}, ** p < 5×10^{-5}, ***p < 5×10^{-7}. (F) Boxplot showing the median and interquartile

ranges of the number of enriched gene pairs in highly connected cell type groups. All cell

type connections in these groups have at least 40 enriched gene pairs (dotted line).

Figure 5: SAMap identifies muscle and stem cell transcriptional signatures conserved across species. (A) Enrichment of KOG functional annotations calculated for genes shared in contractile cell types. For each species, genes enriched in individual contractile cell types are combined. (B) Expression and enrichment of conserved muscle genes in contractile cell types. Color: mean standardized expression. Symbol size: the fraction of cells each gene is expressed in per cell type. Homologs are grouped based on overlapping eukaryotic Eggnog orthology groups. If multiple genes from a species are contained within an orthology group, the gene with highest standardized expression is shown. Genes in blue: core transcriptional program of bilaterian muscles; red: transcription factors conserved throughout Metazoa. (C) Enrichment of KOG functional annotations for genes shared by stem cell types. (D) Boxplot showing the median and

949 interquartile ranges of the mean standardized expressions of genes in hydra and

950 planarian stem cells/progenitors that are conserved across all invertebrate species in this

951 study. Planarian progenitors: *piwi+* cells that cluster with differentiated tissues in Fincher

952 et al. (Fincher et al., 2018). Neoblasts: cluster 0 in Fincher et al. (Fincher et al., 2018) that

953 does not express any tissue-specific markers.

Figure 2 – figure supplement 1: Expression of selected genes enriched in *D. rerio* and *X. tropicalis* secretory cell types. Expressions of orthologous gene pairs linked by SAMap are overlaid on the combined UMAP projection. Expressing cells are color-coded by species, with those connected across species colored cyan. Cells with no expression are shown in gray. The mapped secretory cell types are highlighted with circles.

Figure 2 – figure supplement 2: Existing methods failed to map *D. rerio* and *X. tropicalis* atlases. (A) UMAP projections of the integration results from SAMap using the

965　full homology graph, compared to Liger, BBKNN, Scanorama, Seurat, Harmony, and

966　SAMap using 1-1 orthologs. For fair comparisons, all methods were run on the *D. rerio*

967　and *X. torpicalis* atlases subsampled to approximately 15,000 cells to satisfy

968　computational constraints of Seurat and Liger. (B) Distribution of alignment scores

969　between individual cells.

Figure 2 – figure supplement 3: Representative examples of paralog substitution events in _D. rerio_ and _X. tropicalis_ atlases. Expressions of orthologous and paralogous gene pairs are overlaid on the combined UMAP projection. Expressing cells are color-coded by species, with those that are connected across species colored in cyan. Cells with no expression are shown in gray.

Figure 3 – figure supplement 1: **SAMap-linked gene pairs that are enriched in cell type pairs between *S. mediterranea* and *S. mansoni*.** (A) Rows: linked cell types. Schistosome cell types correspond to leiden clusters. Columns: genes linked by SAMap with overlapping eukaryotic Eggnog orthology groups. We calculate the average standardized expression of each gene in an orthology group for its corresponding cell type in a particular pair and report the highest expression. A selected set of orthology groups corresponding to transcriptional regulators are labeled. (B) Fluorescence *in situ* hybridization shows the co-expression of *wnt* (Smp_156540) and a panel of muscle markers (*collagen*, *troponin*, *myosin* and *tropomyosin*) in *S. mansoni* juveniles. The body wall muscles are expected to be located close to the parasite surface (dashed outline).

The images are maximum intensity projections constructed from ~10 confocal slices with

optimal axial spacing recommended by the Zen software collected on a Zeiss LSM 800

confocal microscope using a 40× (N.A. = 1.1, working distance = 0.62 mm) water-

immersion objective (LD C-Apochromat Corr M27). (C) Whole mount *in situ* hybridization

images showing that the expression of *wnt* and *frizzled* (Smp_174350) are concentrated

in the parasite tail (arrows) with decreasing gradients extending anteriorly. In planarian

muscles, Wnt genes provide the positional cues for setting up the body plan during

regeneration (Scimone et al., 2017; Reddien, 2018). The presence of an anterior-

posterior expression gradient of *wnt* and *frizzled* in muscles of schistosome juveniles

suggests that they may have similar functional roles in patterning during development.

1000

Figure 3 – figure supplement 2: Schistosome neoblasts express canonical muscle markers in muscle progenitors. UMAP projections of schistosome neoblasts with gene expressions overlaid. μ and μ' cells are circled. Colormap: expression in units of $log_2(D + 1)$. For visualization, expression was smoothed via nearest-neighbor averaging using SAM. Note that *myod1* and *cabp* are expressed in both presumptive muscle progenitor populations, whereas all other markers are enriched in μ' cells. All genes displayed are also expressed in fully differentiated muscle tissues.

1009

Figure 4 – figure supplement 1: **Alignment scores are mostly independent of edge transitivity.** Top left: alignment scores and edge transitivity for all cell type pairs in the connectivity graph including the 7 species. Dotted line: the linear best fit, with the Pearson

1013   correlation coefficient reported at the top. Alignment scores and edge transitivity for

1014   individual species pairs are shown in the remaining subplots.

1015

1016

Figure 4 – figure supplement 2: **Number of enriched gene pairs are mostly independent of edge transitivity.** Top left: The edge transitivity is plotted against the number of enriched gene pairs for all cell type pairs in the connectivity graph. Dotted line: the linear best fit, with the Pearson correlation coefficient reported at the top. Top right: magnified view of the mapped cell type pairs supported by small numbers of gene pairs

1022   (<40) to show those edges have low transitivity scores (<0.4). The sublots below show

1023   the number of enriched gene pairs and edge transitivity for individual species pairs.

1024

Figure 5 – figure supplement 1: Phylogenetic reconstruction of animal contractile

cell transcriptional regulators. Trees depict *Csrp/Crip* (A) and Fox group I (B) gene

families. Genes labelled red are enriched in at least one contractile gene pair identified

via SAMap. Support values indicate bootstrap support from 1,000 nonparametric (*Csrp*)

or ultrafast (*Fox*) bootstrap replicates. Besides these two transcriptional regulators,

contractile cells in all seven species were found to be also enriched for transcription

factors from the C2H2 Zinc Finger, Lim Homeobox, and Paired Homeobox families,

although in different cell types we found enrichment of a number of distinct orthologs.

Whether this reflects an ancestral role for these transcription factor families in regulating

contractility or their independent evolution will require additional taxonomic sampling and

broader coverage of muscle cell diversity to resolve.

Supplementary table captions

1037

1038

Supplementary Table 1: Cell atlas metadata and cell annotations. Metadata includes

the number of cells, number of transcripts in the transcriptome, median number of

transcripts detected per cell, the reference transcriptome used in this study, database

through which the transcriptomes are provided, technology used for constructing the cell

atlases, atlas data accessions, processing notes, and references. Leiden clusters and

cell type annotations are reported for cells in each atlas. The Zebrafish and *Xenopus*

tables include both the original cell type annotations and those used in this study. *D. rerio*,

X. tropicalis, and mouse annotations include developmental stages.

1047

1048 **Supplementary Table 2: Cell type annotations for the zebrafish-*Xenopus* mapping.**

1049 Correspondence between the cell type annotations provided in the original study (Briggs

1050 et al., 2018; Wagner et al., 2018) and corresponding annotations used in this study is

1051 provided for both *D. rerio* and *X. tropicalis* atlases.

1052

Supplementary Table 3: Identified paralogs with greater expression similarity than

orthologs in the zebrafish-*Xenopus* mapping. Each row contains a pair of vertebrate-

orthologous genes and a corresponding pair of eukaryotic paralogs with higher correlation

in expression compared to the orthologs, the expression correlations for ortholog and

paralog pairs, the difference between their correlations, and whether the paralogs are

considered as a paralog substitution (defined as when the substituted ortholog is either

absent or lowly-expressed with no cell-type specificity). Highlighted rows are shown in

Figure 2E and **Figure 2 – figure supplement 3**.

1061

Supplementary Table 4: Genes enriched in contractile cell types and invertebrate

stem cells highlighted in Figure 4D. The IDs of the genes enriched in the contractile

and invertebrate stem cell types are provided along with the IDs of the Eggnog orthology

groups to which they belong. In cases where multiple genes from a species belonging to

the same orthology group are enriched, the most differentially expressed gene is shown.

The descriptions in the stem cell table are orthology annotations associated with the

Spongilla genes provided in the original study (Musser et al., 2019).

1069

Supplementary Table 5: Cell types in the cell type families shown in Figure 4D. For

the schistosome cell types, we annotated two neural clusters, both of which express the

neural marker *complexin* (Li et al., 2020). One of the clusters expresses the antigen

SmKK7, so we label the clusters "Neural" and "Neural_KK7", respectively. The "Muscle"

population contains non-neoblast cells expressing *troponin*. The "Tegument_prog" and

"Tegument" populations consist of cells expressing tegument progenitor and

differentiated marker genes, respectively, as reported in a previous study (Wendt et al.,

2018).