Adult spiny mice (*Acomys*) exhibit endogenous cardiac recovery in response to myocardial infarction

Hsuan Peng[1*], Kazuhiro Shindo[1*], Renée R. Donahue[1], Erhe Gao[2], Brooke M. Ahern[3], Bryana M. Levitan[3,4], Himi Tripathi[1], David Powell[4], Ahmed Noor[4], Jonathan Satin[3#], Ashley W. Seifert[5#], Ahmed Abdel-Latif[1,4,6#]

[1] *Saha Cardiovascular Research Center, College of Medicine, University of Kentucky*

[2] *The Center for Translational Medicine, Lewis Katz School of Medicine, Temple University, Philadelphia, PA, USA*

[3] *Department of Physiology, College of Medicine, University of Kentucky*

[4] *Gill Heart and Vascular Institute and Division of Cardiovascular Medicine, University of Kentucky*

[5] *Department of Biology, University of Kentucky*

[6] *The Lexington VA Medical Center*

*These authors contributed equally to the study

[#]To whom correspondence should be addressed: Jonathan Satin: Jonathan.Satin@uky.edu;

Ashley W. Seifert: awseifert@uky.edu and Ahmed Abdel-Latif: abdel-latif@uky.edu

Abstract

Complex tissue regeneration is extremely rare among adult mammals. An exception, however,

is the superior tissue healing of multiple organs in spiny mice (*Acomys*). While *Acomys* species

exhibit the remarkable ability to heal complex tissue with minimal scarring, little is known

about their cardiac structure and response to cardiac injury. In this study, we first examined

baseline *Acomys* cardiac anatomy and function in comparison with the commonly used

laboratory *Mus* strains (C57BL6 and SWR). Our results demonstrated comparable cardiac

anatomy and function between *Acomys* and *Mus*, but *Acomys* exhibited a higher percentage of

cardiomyocytes exhibiting immature characteristics. In response to myocardial infarction, all

animals experienced a comparable level of initial cardiac damage. However, *Acomys*

demonstrated superior ischemic tolerance and cytoprotection in response to injury as evidenced

by cardiac functional stabilization, higher survival rate and smaller scar size 50 days after injury

compared to the inbred and outbred mouse strains. Overall, these findings demonstrate

augmented myocardial preservation in spiny mice post-MI and establish *Acomys* as a new adult

mammalian model for cardiac research.

Introduction

Significant clinical advances for heart revascularization and next generation medical therapies have improved the mortality rate after myocardial infarction (MI).[1] The initial ischemic insult, even with timely revascularization, is followed by microvascular damage and infarct expansion leading to exacerbated damage.[2] Unfortunately, no therapies exist to limit myocardial damage from MI and the ensuing infarct expansion where millions of recovering patients progress to develop heart failure (HF). This is because the normal healing response to tissue injury in adult mammals is fibrotic repair; an effective short-term strategy, but one that leads to compromised tissue function. Fibrotic repair is the default strategy in the adult heart where it is the leading cause of the clinical HF epidemic.[3] Multiple therapies developed for limiting infarct expansion or cardiac repair based on animal models have achieved only modest clinical success in humans. A failure to translate basic research findings into successful clinical treatments can be traced, in part, to a lack of adult mammalian models possessing the ability of myocardial preservation.

In contrast to humans[4-6], the ability to regenerate injured organs is widespread among vertebrates. Fishes, newts, and salamanders have extensive regenerative ability, and can functionally replace heart tissue after amputation or severe injury by expediting revascularization[7] and by mobilizing a highly proliferative cardiomyocyte pool[8,9]. Specifically, zebrafish are a well-characterized model for adult cardiac regeneration with the documented ability to recover from a plethora of heart injuries including apical resection,[10] cryo-injury,[11] and coronary artery ligation. Neonatal mice are also capable of limiting the initial damage and repairing the myocardium after excision and ischemic injury during the first few days after birth.[12] In contrast, most adult mammals generally exhibit poor regenerative capacity, especially as it pertains to recovering from heart damage. Spiny mice (*Acomys spp.*) are murid rodents found throughout Africa, the Middle East and Western Asia. These rodents exhibit a number of special traits;[13] tantamount among them is the ability to regenerate skin, complex

tissue[14-19] and nephric tissue.[20] At present, how spiny mice respond to heart injury and the

extent to which they can recover remains unknown, although a recent review on the topic raised

the intriguing possibility that they might exhibit enhanced cardiac healing.[21]

To rigorously examine how spiny mice respond to heart damage, we first detail

Acomys heart structure and cardiac function in comparison to the most widely used inbred

laboratory *Mus* model of cardiac injury (C57BL6). For comparative purposes we also used the

Swiss Webster outbred strain (SWR) to account for greater genetic diversity within our spiny

mouse population. This cross-species characterization indicates similar cardiac structure and

comparable baseline function among spiny mice and the examined mouse strains. We next

conducted a comparative analysis of these animals in response to permanent coronary artery

ligation (MI). In contrast to the mouse strains studied, *Acomys* showed significantly improved

survival and enhanced myocardial preservation following ischemic injury. The ability of

Acomys to recover from ischemic injury is coincident with greatly improved angiogenesis

rarely seen in adult mammals and unique cardiomyocyte characteristics (higher percentage of

small, mono-nucleated and diploid CMs, T-type calcium channels). These findings in *Acomys*

support a rapid angiogenic response into fibrotic tissue which limits scar spread and enhances

ischemic tolerance in response to MI.

Methods

Detailed methods are described in the Supplementary Methods

Animal care

Male *Acomys cahirinus* (6-8 months old, sexually mature animals) were obtained from our in-house breeding colony. Male *Mus musculus* (8-12 weeks, sexually mature animals) C57BL6 were obtained from the Jackson Laboratory, Bar Harbor, ME and outbred SWR were obtained from Charles River. Both strains were housed at the University of Kentucky, Lexington, KY. All animal procedures were approved by the University of Kentucky Institutional Animal Care and Use Committee (IACUC #: 2019-3254, 2013-1155 and 2011-0889).

Murine model of myocardial infarction

MI surgery was performed using a minimally invasive permanent left anterior descending artery (LAD) ligation model as previously described.[22]

Cardiomyocyte isolation and measurements

Ventricular cardiomyocytes were isolated using the Langendorff method as previously described[23] and are further detailed in the supplementary methods. Images used to analyze cardiomyocyte surface area and nucleation were acquired using an Olympus IX-71 and Olympus BX53 microscopes (Olympus, Tokyo, Japan) at 10x magnification. Olympus cellSens software was used to determine cell surface area and number of nuclei. Nuclear volume and ploidy were determined by Imaris reconstruction of z-stack images acquired using a Nikon Ti, A1 confocal microscope (Nikon, Japan) with a step size of 3 μm and 40x oil immersion objective.

Echocardiography

Echocardiography was acquired using a Vevo 3100 (VisualSonics, Toronto, Canada)

equipped with a MX550D 25-55 MHz linear array transducer. Cardiac function was assessed

as previously described.[22] Echocardiographic imaging and analyses were performed by a

blinded investigator.

Cardiac Magnetic Resonance Imaging

Cardiac magnetic resonance imaging (CMR) was performed on a 7-Tesla ClinScan

system (Bruker, Ettlingen, Germany, http://www.bruker.com) equipped with a 4-element

phased-array cardiac coil and a gradient system with a maximum strength of 450 mT/m and a

maximum slew rate of 4,500 mT/m/s. Studies were performed within 72 hours of MI and at 14

days following injury. For late gadolinium-enhanced magnetic resonance imaging, a 0.6

mmol/kg bolus of gadolinium-diethylene triamine pentaacetic acid (Gd-DTPA; Gadavist,

Bayer Health Care, Whippany, NJ) was injected using the intraperitoneal route.

Electrophysiological recordings and calcium transients

$I_{Ca,L}$ was recorded from freshly isolated cardiomyocytes using the whole-cell

configuration of the patch clamp technique, and cytosolic Ca^{2+} transients were collected using

fura2-AM as previously described.[23]

Tissue collection and Processing

Tissue from all species was collected and weighed immediately at sacrifice. The dry

lung weight was collected after incubation at 65° C for three days. Hearts were perfused with

PBS (VWR International) followed by 4% PFA (VWR International) fixation via cannulation

of the ascending aorta. Hearts were post-fixed overnight at 4°C. Hearts were then cut in half

along the long axis at the level of the ligation and transferred to 70% ethanol until embedding.

Tissues were then paraffin embedded and sectioned at 5um for immunofluorescence and at

8um for Masson's Trichrome and Picrosirius Red staining.

Histology

Paraffin-embedded heart sections were stained with Masson's Trichrome or

Picrosirius Red. Morphometric analyses and image acquisition were performed using a

Olympus BX53 microscope (Olympus, Tokyo, Japan) and NIH ImageJ 1.46R software.

Quantification of percent fibrosis and scar thickness are detailed in the supplementary methods.

Scar collagen fiber quality was examined under 40x magnification. Cellular density in scar

tissue was determined by quantifying DAPI+ nuclei in the infarct region at 60x magnification.

Infarct size was also quantified three days post-MI from Masson's Trichrome stained slides as

the percentage of left ventricle affected. These images were acquired on Zeiss Axioscan slide

scanner at 10x magnification. Images were quantified using Nis-Elements software (Nikon).

Immunofluorescence

Immunofluorescence assessments were carried out on deparaffinized and rehydrated

sections as previously described.[22] Images were taken with a 40x Oil Immersion lens on a

Nikon A1 Confocal Microscope in the University of Kentucky Confocal Microscopy facility.

Isolectin staining was performed on tissues from day 50 post-MI and was quantified at both

the peri-infarct border and the center of the scar and is presented as total capillary density per

mm^2. Baseline isolectin measurements were taken at comparable location to the peri-infarct

area. A complete list of antibodies is included in the supplemental data.

Statistics

Values are expressed as mean ± standard error of mean (SEM). Data were analyzed

using a Student's t-test, one-way ANOVA, repeated-measures ANOVA or mixed-effects

ANOVA with Tukey correction to compare data across species as appropriate. Animal numbers

are presented as a range based on the number of animals included in each analysis. These

numbers are included in the figure legends. All statistical analyses were performed using the

Prism 9 software package (GraphPad, La Jolla, CA).

Results

Heart and coronary tree anatomy are similar among *Acomys* and *Mus* (C57BL6 and SWR)

To assess whether interpretable comparisons could be made across species, we first characterized hearts from *Acomys* and the most commonly used laboratory mouse (*Mus*) strain: C57BL6. To control for lifespan differences between species, we used 6 month old, sexually mature *Acomys* and 8-12 week old *Mus* for these studies. In an attempt to control for increased genetic diversity in *Acomys* as well as their larger size, we also included an outbred Swiss Webster strain (SWR – Charles River). As these outbred mice have not been evaluated for myocardial ischemia research, this strain also allowed us to assess whether they shared similarities with the SWR/J strain that is known to exhibit a high percentage of mononuclear cardiomyocytes (CMs) that contribute to enhanced cardiac recovery after ischemic injury.[24] First, we set out to determine whether baseline differences in cardiac gravimetrics exist between *Acomys* and *Mus* (C57BL6 and SWR). We assessed heart weight (HW) and normalized it to body size (body weight [BW] and tibia length [TL]) to account for overall size differences between species (Fig. 1A-D, and Supplementary Table 1). Although *Acomys* demonstrated a higher mean HW compared to C57BL6 (Fig. 1B), normalized to BW, *Acomys* was not significantly different compared to SWR and exhibited only a slighly lower HW/BW compared to C57BL6 (Fig. 1C). SWR mice exhibited a slightly higher HW compared to *Acomys* and C57BL6 when HW was normalized to TL (Fig. 1D). *Acomys* demonstrated heavier wet lung weight compared to *Mus* (C57BL6 and SWR) (Supplementary Fig. 1A), but lower wet and dry lung weight after normalization by BW (Supplementary Fig. 1B, E). Importantly, dry lung weight before normalization by TL and dry and wet lung weight after normalization by TL were not significantly different across species (Supplementary Fig. 1D-F). The complete data is summarized in Supplementary Table 1.

Next, we compared cardiac functional parameters between the three strains and

observed comparable baseline cardiac function under physiological conditions across species.

Importantly, the global cardiac function as presented by left ventricular ejection fraction and

fractional shortening were similar between all groups (Fig. 1E-F). Measures of left ventricular

cavity diameter were also comparable between species except for SWR-*Mus* showing slightly

larger left ventricular end-diastolic diameter and anterior wall thickness compared to *Acomys*

(Fig. 1G-J and Supplementary Table 1). We also assessed gross cardiac structure and coronary

tree among *Acomys* and *Mus* (C57BL6 and SWR) and found that cardiac structure was not

grossly different between the species upon visual comparison of whole hearts along the short-

or long-axis (Supplementary Fig. 1G). Using corrosion casting and intra-aortic infusion of

Batson's 17 polymer mixture, we delineated the left coronary anatomy in all three strains.

Relevant to permanent coronary ligation, all three animals demonstrated similar course of the

left anterior descending artery (LAD) supplying the anterior wall, and of the left circumflex

artery supplying the lateral aspect of the left ventricle (Fig. 1K). Coronary blood vessels adopt

a hierarchal structure and vascular density leads to enhanced blood supply, angiogenesis and

ischemic resistance.[25,26] Thus, we analyzed the bifurcation points of the coronary tree as it

supplies the anterior wall of the left ventricle. These analyses showed a similar density of

branching points per major artery across species (Fig. 1L). Finally, we assessed capillary

density at the mid left ventricular level between species and observed slightly higher capillary

density in *Acomys* and SWR-*Mus* compared to C57BL6-*Mus* (Supplementary Fig. 2). These

data demonstrate that cardiac anatomy and physiology are comparable between spiny mice and

the two inbred laboratory mouse strains C57BL6 and SWR.

Adult *Acomys* exhibit a high percentage of cardiomyocytes possessing fetal characteristics

Ventricular cardiomyocyte (CM) phenotype varies across species and chronological

age where it can influence the cardiac response to injury.[12,24] In mammals, the presence of

mononuclear diploid CMs are indicative of a 'young' heart phenotype that is rarely seen in

adult mice.[27, 28] To examine CM phenotype between species, we isolated ventricular CMs from

Mus and *Acomys* and examined their size, number of nuclei/CM, nuclear characteristics and

ploidy. Similar to a prior report,[29] *Acomys* CMs were significantly smaller than C57BL6 and

were approximately four times more likely to be mononucleated (*Acomys*: 25.9±5.1% vs.

C57BL6: 6.3±2.8%, $P < 0.001$; Fig. 2A-C). Similarly, CM nuclei in *Acomys* were smaller in

size (Fig. 2D).[30] We then assessed the ploidy of ventricular CMs across species using 3-

dimensional confocal microscopy[31] and found a significantly higher percentage of diploid CMs

in *Acomys* with a corresponding lower percentage of >4N CMs (Fig. 2E). Interestingly, SWR-

Mus CMs were intermediate between *Acomys* and C57BL6 with smaller size and higher

percentage of mononuclear CMs as shown for the inbred strain SWR/J in a previous study.[24]

The relatively small cell size and higher mononucleation we observed in *Acomys* CMs

suggested a higher percentage of CMs exhibiting immature or fetal characteristics. T-tubule

organization is highly regular in mature mouse CMs.[32] To assess T-tubule organization we

stained CMs with di-8-ANEPPS (Fig. 3A) and evaluated transverse and axial density along

with T-tubule spacing. *Acomys* showed significantly reduced transverse and long element

densities (Fig. 3B and 3C) without a change in resting T-tubule spacing (Fig. 3D). To further

explore the physiological implications of these findings, we conducted electrophysiological

studies on isolated ventricular CMs from C57BL6-*Mus* and *Acomys*. In mammalian hearts, the

expression of low-voltage activated (T-type calcium channels) is an index of the fetal gene

program;[33] therefore, we recorded voltage-dependent I_{Ca} from *Acomys* and *Mus* ventricular

myocytes. From a V_{hold} -80mV, a V_{test} step to -25mV elicited little if any discernable current in

Mus (C57BL6) whereas, at V_{test} -25mV, *Acomys* cardiomyocytes exhibited a transient 'T-type'

calcium current ($I_{Ca,T}$) (Fig. 3E, upper panel). Longer lasting calcium current ($I_{Ca,L}$) was

prominent with larger depolarizations in *Mus* (C57BL6) and *Acomys* (Fig. 3E, lower panel).

The current-voltage relationship showed a prominent $I_{Ca,T}$ component with a peak current

between V_{test} -20 and -25mV in *Acomys*, but not in *Mus* (C57BL6) (Fig. 3E).

To further dissect $I_{Ca,T}$ from $I_{Ca,L}$, we performed current-voltage curve protocols from

V_{hold} -50mV to voltage-inactivate the $I_{Ca,T}$ component (Fig. 3F). Current-voltage relationships

for $I_{Ca,T}$-expressing *Acomys* cells showed the prominent $I_{Ca,T}$ appearing as a low-voltage

activated current for V_{hold} -80mV but not from V_{hold} –50 mV (Fig. 3F, left). 33% of *Acomys*

ventricular cardiomyocytes showed no $I_{Ca,T}$ (Fig 3F, center) but there was no observed

correlation of cell size to presence of $I_{Ca,T}$ (Fig. 3F, right). Cell capacitance was significantly

lower in *Acomys* compared to *Mus* (C57BL6) (Fig. 3G), consistent with our morphometric

analysis (Fig. 2B). Additionally, maximal conductance density trended greater in *Mus*

(C57BL6) than *Acomys* (Fig. 3H). Voltage-dependent activation and inactivation of $I_{Ca,L}$ was

significantly shifted positive for *Acomys* compared to *Mus* (C57BL6) (Fig. 3I-J).

We next explored cytosolic calcium handling. Cytosolic Ca^{2+} transients (CaT) had a

larger amplitude (Fig. 3K, left), faster upstroke (Fig. 3K, middle), and more rapid decay (Fig.

3K, right) in *Mus* (C57BL6) compared to that observed in *Acomys*. As with CM morphometrics

(Fig. 3A-D) and the prevalent expression of a T-type calcium current (Fig. 3E-F) the smaller,

slower CaT segregated with a less mature CM phenotype. Immature ventricular CMs also tend

to show reduced β-adrenergic receptor (β-AR) acute responsiveness.[34] Therefore, we tested the

effect of isoproterenol challenge (ISO). To assess acute ISO responsiveness, we compared

within cell before – after ISO. *Mus* (C57BL6) showed increased amplitude and more rapid

kinetics (Fig. 3L). By contrast, *Acomys* CaT amplitude and upstroke velocity were not

significantly different ($P>0.05$). Ca^{2+} re-uptake was accelerated by ISO in *Mus* but not in

Acomys (Fig. 3L, right). Taken together, these data are consistent with an *Acomys*

cardiomyocyte phenotype distinct from that of C57BL6-*Mus*. Moreover, these data

demonstrate that a high percentage of *Acomys* CMs exhibit characteristics usually associated

with *Mus* fetal CMs.

***Acomys* hearts demonstrate enhanced cardiac preservation after ischemic injury**

Acomys can regenerate complex tissue wounds, but the response to myocardial

ischemic (MI) injury remains untested. We next sought to compare the cardiac injury response

between *Acomys* and the two strains of *Mus* using the permanent LAD-ligation model of

myocardial infarction.[35-37] After injury, all groups showed a comparable drop in cardiac

function, reflecting similar injury response as assessed by serial echocardiography (Fig. 4,

Supplementary Fig. 3 and Supplementary Table 1). Indeed, there was no significant difference

in most of cardiac fucntional parameters at 48 hours after MI (Fig. 4B-F, *P*>0.05). When we

assessed the extent of cardiac tissue injury in a subset of animals three days after MI using

Mason Trichrome staining, we observed comparable percent infarct between species

(Supplementary Fig. 3A). Furthermore, our cardiac MRI imaging showed comparable injury

and involvement of the distal anterior wall and apex three days after MI beween species as

demonstrated by similar area of late gadolinium enhancement (Supplementary Fig. 4).

Interestingly, we observed stabilization in cardiac function and remodeling pararmeters in

Acomys compared to the progressive deterioration seen in *Mus* strains (C57BL6 and SWR) up

to 50 days after injury (Fig. 4B-F). In these analyses, we utilized the delta change from baseline

to account for any differences in baseline function between species which allowed us to isolate

the analyses to the changes occurring after MI. We observed consistent changes in left

ventricular global ejection fraction, as well as measures of left ventricular adverse remodeling

such as end-systolic and end-diastolic diameters and volumes (Fig. 4A-F and Supplementary

Fig. 3B-F). The enhanced cardiac preservation and reduced adverse cardiac remodeling in

Acomys was reflected by lower heart/body weight ratio throughout the follow-up period

extending to 50 days after MI (Fig. 4G), as well as the aforementioned echocardiographic

evidence of smaller ventricular diameters. We also observed significantly lower wet-dry lung

weight in *Acomys* compared to *Mus* which is a measure of pulmonary edema and heart failure

(Fig. 4H). Importantly, enhanced myocardial preservation in *Acomys* compared to *Mus* was

associated with a significant survival advantage (Fig. 4I). As seen in prior reports, the majority

of mortality across all species was observed in the first week after MI. However, of those

animals that died, while we observed cardiac rupture in 17 C57BL6, we observed none in

Acomys as visualized in post-mortem necropsies.

CMR studies demonstrated reduced scar progression between day 3 (D3) and D14 in

Acomys compared to the continued infarct expansion seen in *Mus* (Fig. 4J). Our long-term

follow-up studies reflected significantly smaller infarct size in *Acomys* compared to *Mus*

(C57BL6 and SWR) when assessed using Masson's Trichrome staining 50 days after MI (Fig.

5A). We also saw similar results in our studies quantifying the degree of fibrosis using

Picrosirius red staining at 50 days after MI (Supplementary Fig. 3G). Furthermore, the resultant

scar in *Acomys* appeared thicker with higher cellular density (Fig. 5C), a finding that could

potentially explain the lower rupture and mortality rates. To further explore the fibrous tissue

alignment of the thicker scar in *Acomys*, we stained cardiac tissue using Picrosirius Red. While

the fiber alignment in the *Mus* strains was typical of a scar with highly parallel and compressed

collagen bundles, fiber organization in the center of the *Acomys* scar was more porous with less

compression (Fig. 5B). Furthermore, our serial scar analysis at days 7, 17 and 50 after MI

demonstrated a unique evolution in infarct size between *Mus* and *Acomys* which corroborated

the CMR studies (Fig. 6A-D). In contrast to previous reports,[24] our studies showed limited

recovery of cardiac functional parameters after MI in SWR-*Mus* (Fig. 4B-F). Taken together,

our results demonstrate that *Acomys* exhibit ischemic tolerance and enhanced myocardial

preservation after ischemic injury compared to the inbred C57BL6- and outbred SWR-*Mus*

strains, respectively.

***Acomys* exhibit rapid and enhanced vascularization of the infarct region after ischemic**

injury.

Blood vessel formation following ischemic injury is essential for functional stabilization. Following ischemic injury, vascular damage ensues in the area surrounding the original infarct region and leads to infarct expansion, development of adverse cardiac remodeling and heart failure.[2,38-40] Several studies have shown successful reduction of infarct expansion through effective angiogenesis and vascular maturation.[41,42] We assessed the vascular density in the peri-infarct region in *Acomys* and *Mus* strains and observed an increased density of capillaries (isolectin+ cells) in *Acomys* compared to C57BL6-*Mus* (Fig. 7A). As commonly observed in adult mammals following MI, both *Mus* strains exhibited reduced vascular density in the scar center; by contrast, *Acomys* showed enhanced vascular density extending into the scar center (Fig. 7A). Additionally, enhanced angiogenesis was linked to more proliferating endothelial cells (isolectin+/EdU+) in the peri-infarct region and scar center in *Acomys* compared to both *Mus* strains (Fig. 7B). In support of enhanced angiogenesis, we observed a significantly higher density of mature blood vessels (α-SMA+) in the peri-infarct region in *Acomys* while the presence of these mature vessels was negligible in *Mus* strains (Fig. 7C). Taken together, in response to MI our data support rapid and enhanced angiogenesis followed by vascular maturation in *Acomys* compared to *Mus*. This can explain, at least in part, the ischemic tolerance, reduced infarct expansion and the thicker, cellularly dense scar in *Acomys* compared to *Mus*.

Discussion

Spiny mice (*Acomys*) represent a novel mammalian model to explore endogenous cardiac repair consistent with their enhanced regenerative ability for a number of tissues and organs. In this study, we established the anatomical, functional, and cellular characteristics pre- and post-MI in *Acomys* compared to the inbred laboratory mouse strain C57BL6 and the outbred SWR strain. Our results demonstrated comparable cardiac structure, coronary anatomy and functional parameters between the three animals allowing us to appropriately use these animal models for comparative heart injury studies. Interestingly, when comparing cardiomyocytes across species we found that adult *Acomys* possessed a distinct CM phenotype that included features usually observed in immature or fetal *Mus* CMs such as small size, mononucleation and reduced t-tubule density and organization. After ischemic injury, *Acomys* exhibited enhanced ischemic tolerance and significant myocardial preservation, resulting in reduced adverse cardiac remodeling, smaller scar size and, importantly, better survival. Overall, our data supports that *Acomys* show resistance to the type of cardiac damage that compromises other adult mammals, including humans and supports future cardiac studies using this model.

Similar to other highly regenerative vertebrates, *Acomys* has emerged as a bona fide mammalian regeneration model for their ability to naturally regrow complex tissues following full-thickness skin injury[14-19] and excision of musculoskeletal tissue from the ear pinna.[16,18,43] In each type of injury, adult *Acomys* were capable of restoring functional tissues, a phenotype that has not been observed in other adult rodents. Our gross anatomical studies established a similar coronary anatomy across species [44] and we found that *Acomys* hearts were similar structurally and anatomically to *Mus*. Importantly, *in vivo* cardiac functional parameters were comparable between the three groups. These findings provided the appropriate foundation for the current study and lay the groundwork for follow-up studies that can inform new therapeutic strategies for ischemic heart disease.

Increased mortality and incidence of heart failure following myocardial infarction

remains prevalent in modern medicine. While advanced revascularization therapies and

mechanical support have slightly improved survival in patients with myocardial infarction,

millions develop progressive heart failure every year. Remarkably, *Acomys* exhibited

significantly lower mortality after MI compared to both *Mus* strains and this reduction in

mortality was related to a reduced incidence of cardiac rupture among *Acomys*. In fact, our

histological assessment combined with cardiac MRI studies indicated significant differences in

infarct expansion and scar evolution between species. While *Mus* demonstrated progressive

infarct expansion, associated with thinning of the anterior wall, *Acomys* showed minimal

expansion despite starting with comparable area of injury. These differences in infarct

expansion led to favorable cardiac remodeling and functional preservation demonstrated in our

echocardiography studies.

This unique phenomenon was associated with a rapid angiogenic response that

eventually led to superior ischemic tolerance. *Acomys* scars appeared to be more cellularized

and had higher capillary and mature vessel density compared to *Mus*. This could explain the

dramatically lower cardiac rupture rate and enhanced survival we observed in *Acomys*

including the stabilization of cardiac function in *Acomys* during long-term follow-up.

Preclinical and human MI studies have linked enhanced angiogenesis and vessel maturation

with lower rates of infarct expansion.[42,45] Our studies demonstrated superior endothelial cell

proliferation and vessel maturation in *Acomys* compared to *Mus* strains which likely accounted

for some of the protection seen in our studies.

The mammalian heart loses its reparative capability after the first week of life where

heart damage results in permanent loss of myocardium combined with fibrotic scarring and

adverse remodeling.[12] The transition to fibrotic repair is closely linked with the loss of CM

proliferation as CMs enter cell cycle arrest and become mononuclear polyploid or

multinucleated.[12,46] However, in other vertebrate species where CMs maintain the lifelong

ability to proliferate (e.g., zebrafish and newts), cardiac regeneration is still possible.[46, 47] One

of the hallmarks of regenerative hearts is the relatively high frequency of mononucleated

diploid CMs.[46-48] This evidence suggests that higher proportions of mononuclear diploid

CMs are associated with higher regenerative potential following ischemic injury.[24]

Interestingly, our results showed that outbred SWR-*Mus* possessed an intermediate CM

phenotype between *Acomys* and C57BL6 with respect to CM size and nuclei number. Despite

possessing this CM phenotype, outbred SWR had nearly the same mortality as C57BL6

suggesting that rapid angiogenesis may be a key component to facilitate the protective features

conferred by having a larger population of smaller, mononucleate CMs. In fact, cardiac tissue

requires vascularization for supporting the high metabolic activity of CMs,[49] and our current

findings support rapid angiogenic invasion into new fibrotic tissue as a mechanism to resist

cardiac damage. Future studies are necessary to explore CM behavior across species and how

this may relate to enhanced angiogenesis.

There are two types of Ca^{2+} channels in cardiomyocytes, L-type and T-type. L-type

Ca^{2+} channels are highly expressed in the adult heart and are important therapeutic targets for

the management of various cardiovascular diseases. In contrast, T-type Ca^{2+} channels are rarely

found in adults and are present in fetal and early postnatal mice.[50] In our cell electrophysiology

studies, *Acomys* exhibited a higher percentage of T-type Ca^{2+} channels. This feature is

consistent with a higher prevalence of phenotypically 'young' CMs in *Acomys* compared to

Mus (C57BL6). Furthermore, immature ventricular cardiomyocytes in *Mus* showed reduced

Ca^{2+} transient amplitude, slower kinetics,[51] and attenuated β-adrenergic receptor

responsiveness.[52] While this unique phenomena in *Acomys* was associated with superior

myocardial preservation after ischemic injury, because we did not examine Ca^{2+} channels in

outbred SWR mice, we do not know if this feature is associated with the prevalence of small,

diploid, mononucleated CMs. However, it is interesting to note that although our CM

characterization studies showed that outbred SWR possessed a higher prevalence of small,

diploid, mononucleated CMs, this feature alone did not confer the type of enhanced myocardial

observed in previous work using inbred SWR/J mice.[24] This would suggest a synergy between

CM phenotype and enhanced angiogenesis with respect to cardiac preservation following MI.

Future studies are necessary to examine the proliferative potential of adult CMs in *Acomys* and

Mus as well as their potential to adequately respond to myocardial injury.

In conclusion, our study presents *Acomys* as a novel model for cardiac research with

comparable cardiac structure, function and anatomy to C57BL6- and SWR-*Mus*. *Acomys*

exhibited a distinct cardiomyocyte phenotype and angiogenic response resulting in enhanced

recovery and ischemic tolerance after ischemic injury. It is important to note that while we do

not see clear evidence of cardiac regeneration, the enhanced myocardial preservation seen in

Acomys could uncover important therapeutic targets for millions of patients who develop

ischemic cardiomyopathy after myocardial infarction. Future studies will focus on the

mechanism and potential for these phenotypic differences to foster myocardial preservation

and repair instead of scarring and heart failure in models of cardiac disease.

Acknowledgments: The authors would like to thank Josh Sarli for his help with *Acomys* husbandry and Thomas Wilkop Ph.D at the microscopy core at the University of Kentucky for his assistance with imaging. We thank the Markey Cancer Center and the COBRE histology core for their assistance with the tissue specimen preparation. We thank Binoy Joseph for his assistance in imaging and the Spinal Cord and Brain Injury Research Center for use of their imaging equipment. We thank Beverly Meacham and the University of Kentucky MRISC center for the assistance with cardiac MRI studies.

JS, AWS, AAL, KS, HP, RRD designed the project and experiments. KS, HP, RRD, EG, BMA, BML, DP, HT, and AN performed all the experiments.

KS, HP, RRD, BMA, BML, JS, AAL analyzed the data and KS, HP, JS, AWS, AAL wrote the manuscript. All authors commented on and edited the final version.

Funding: Dr. Abdel-Latif is supported by NIH Grant R01 HL124266. Work in Dr. Seifert's lab is supported by NIH R01 AR070313. The content in this article is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Markey Cancer Center Core: This research was supported by the Biospecimen Procurement and Translational Pathology Shared Resource Facility of the University of Kentucky Markey Cancer Center (P30CA177558)

COBRE Core: Research reported in this publication was supported by an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health under grant number P30 GM127211.

Competing Interests: The authors declare that there are no competing interests.

456   **Data Availability**:   The authors declare that all the data supporting the findings of this study

457   are available within the paper and are contained within supplementary information.

References.

1. Benjamin, E.J.*, et al.* Heart Disease and Stroke Statistics-2017 Update: A Report From the American Heart Association. *Circulation* **135**, e146-e603 (2017).

2. Iwakura, K. Modulation of individual susceptibility to the no-reflow phenomenon after acute myocardial infarction. *Curr Pharm Des* **19**, 4519-4528 (2013).

3. McAloon, C.J.*, et al.* The changing face of cardiovascular disease 2000-2012: An analysis of the world health organisation global health estimates data. *Int J Cardiol* **224**, 256-264 (2016).

4. Pfeffer, M.A. & Braunwald, E. Ventricular remodeling after myocardial infarction. Experimental observations and clinical implications. *Circulation* **81**, 1161-1172 (1990).

5. Jugdutt, B.I. Ventricular remodeling after infarction and the extracellular collagen matrix: when is enough enough? *Circulation* **108**, 1395-1403 (2003).

6. Struthers, A.D. Pathophysiology of heart failure following myocardial infarction. *Heart* **91 Suppl 2**, ii14-16; discussion ii31, ii43-18 (2005).

7. Marin-Juez, R.*, et al.* Coronary Revascularization During Heart Regeneration Is Regulated by Epicardial and Endocardial Cues and Forms a Scaffold for Cardiomyocyte Repopulation. *Dev Cell* **51**, 503-515 e504 (2019).

8. Bader, D. & Oberpriller, J.O. Repair and reorganization of minced cardiac muscle in the adult newt (Notophthalmus viridescens). *J Morphol* **155**, 349-357 (1978).

9. Flink, I.L. Cell cycle reentry of ventricular and atrial cardiomyocytes and cells within the epicardium following amputation of the ventricular apex in the axolotl, Amblystoma mexicanum: confocal microscopic immunofluorescent image analysis of bromodeoxyuridine-labeled nuclei. *Anat Embryol (Berl)* **205**, 235-244 (2002).

10. Poss, K.D., Wilson, L.G. & Keating, M.T. Heart regeneration in zebrafish. *Science* **298**, 2188-2190 (2002).

11. Honkoop, H.*, et al.* Single-cell analysis uncovers that metabolic reprogramming by ErbB2 signaling is essential for cardiomyocyte proliferation in the regenerating heart. *Elife* **8**(2019).

12. Porrello, E.R.*, et al.* Transient regenerative potential of the neonatal mouse heart. *Science* **331**, 1078-1080 (2011).

13. Haughton, C.L., Gawriluk, T.R. & Seifert, A.W. The Biology and Husbandry of the African Spiny Mouse (Acomys cahirinus) and the Research Uses of a Laboratory Colony. *J Am Assoc Lab Anim Sci* **55**, 9-17 (2016).

14. Harn, H.I.*, et al.* Symmetry breaking of tissue mechanics in wound induced hair follicle regeneration of laboratory and spiny mice. *Nat Commun* **12**, 2595 (2021).

15. Brant, J.O., Yoon, J.H., Polvadore, T., Barbazuk, W.B. & Maden, M. Cellular events during scar-free skin regeneration in the spiny mouse, Acomys. *Wound Repair Regen* **24**, 75-88 (2016).

16. Gawriluk, T.R.*, et al.* Comparative analysis of ear-hole closure identifies epimorphic
regeneration as a discrete trait in mammals. *Nat Commun* **7**, 11164 (2016).
17. Jiang, T.X., Harn, H.I., Ou, K.L., Lei, M. & Chuong, C.M. Comparative regenerative
biology of spiny (Acomys cahirinus) and laboratory (Mus musculus) mouse skin. *Exp*
Dermatol **28**, 442-449 (2019).
18. Matias Santos, D.*, et al.* Ear wound regeneration in the African spiny mouse Acomys
cahirinus. *Regeneration (Oxf)* **3**, 52-61 (2016).
19. Seifert, A.W.*, et al.* Skin shedding and tissue regeneration in African spiny mice (Acomys).
Nature **489**, 561-565 (2012).
20. Okamura, D.M.*, et al.* Scarless repair of acute and chronic kidney injury in African Spiny
mice (Acomys cahirinus). *bioRxiv* **315069.**(2018).
21. Maden, M. & Varholick, J.A. Model systems for regeneration: the spiny mouse, Acomys
cahirinus. *Development* **147**(2020).
22. Al-Darraji, A.*, et al.* Azithromycin therapy reduces cardiac inflammation and mitigates
adverse cardiac remodeling after myocardial infarction: Potential therapeutic targets in
ischemic heart disease. *PLoS One* **13**, e0200474 (2018).
23. Ahern, B.M.*, et al.* Myocardial-restricted ablation of the GTPase RAD results in a pro-
adaptive heart response in mice. *J Biol Chem* **294**, 10913-10927 (2019).
24. Patterson, M.*, et al.* Frequency of mononuclear diploid cardiomyocytes underlies natural
variation in heart regeneration. *Nat Genet* **49**, 1346-1353 (2017).
25. Carmeliet, P. & Jain, R.K. Molecular mechanisms and clinical applications of angiogenesis.
Nature **473**, 298-307 (2011).
26. Potente, M., Gerhardt, H. & Carmeliet, P. Basic and therapeutic aspects of angiogenesis.
Cell **146**, 873-887 (2011).
27. Alkass, K.*, et al.* No Evidence for Cardiomyocyte Number Expansion in Preadolescent
Mice. *Cell* **163**, 1026-1036 (2015).
28. Soonpaa, M.H., Kim, K.K., Pajak, L., Franklin, M. & Field, L.J. Cardiomyocyte DNA
synthesis and binucleation during murine development. *Am J Physiol* **271**, H2183-2189
(1996).
29. Hirose, K.*, et al.* Evidence for hormonal control of heart regenerative capacity during
endothermy acquisition. *Science* **364**, 184-188 (2019).
30. Lozoya, O.A., Gilchrist, C.L. & Guilak, F. Universally Conserved Relationships between
Nuclear Shape and Cytoplasmic Mechanical Properties in Human Stem Cells. *Sci Rep* **6**,
23047 (2016).
31. Broughton, K.M.*, et al.* Cardiac interstitial tetraploid cells can escape replicative
senescence in rodents but not large mammals. *Commun Biol* **2**, 205 (2019).
32. Sipido, K.R., Carmeliet, E. & Van de Werf, F. T-type Ca2+ current as a trigger for Ca2+
release from the sarcoplasmic reticulum in guinea-pig ventricular myocytes. *J Physiol* **508**

(Pt 2), 439-451 (1998).

33. Nuss, H.B. & Houser, S.R. T-type Ca2+ current is expressed in hypertrophied adult feline left ventricular myocytes. *Circ Res* **73**, 777-782 (1993).

34. Schroder, E., Byse, M. & Satin, J. L-type calcium channel C terminus autoregulates transcription. *Circ Res* **104**, 1373-1381 (2009).

35. Dick, S.A.*, et al.* Self-renewing resident cardiac macrophages limit adverse remodeling following myocardial infarction. *Nat Immunol* **20**, 29-39 (2019).

36. Epelman, S., Liu, P.P. & Mann, D.L. Role of innate and adaptive immune mechanisms in cardiac injury and repair. *Nat Rev Immunol* **15**, 117-129 (2015).

37. Lindsey, M.L.*, et al.* Guidelines for experimental models of myocardial ischemia and infarction. *Am J Physiol Heart Circ Physiol* **314**, H812-H838 (2018).

38. Heusch, G. The Coronary Circulation as a Target of Cardioprotection. *Circ Res* **118**, 1643-1658 (2016).

39. Wang, J., Toan, S. & Zhou, H. Mitochondrial quality control in cardiac microvascular ischemia-reperfusion injury: New insights into the mechanisms and therapeutic potentials. *Pharmacol Res* **156**, 104771 (2020).

40. Kloner, R.A. No-reflow phenomenon: maintaining vascular integrity. *J Cardiovasc Pharmacol Ther* **16**, 244-250 (2011).

41. Nazir, S.A.*, et al.* Strategies to attenuate micro-vascular obstruction during P-PCI: the randomized reperfusion facilitated by local adjunctive therapy in ST-elevation myocardial infarction trial. *Eur Heart J* **37**, 1910-1919 (2016).

42. Zhang, S.*, et al.* Comparison of various niches for endothelial progenitor cell therapy on ischemic myocardial repair: coexistence of host collateralization and Akt-mediated angiogenesis produces a superior microenvironment. *Arterioscler Thromb Vasc Biol* **32**, 910-923 (2012).

43. Simkin, J., Gawriluk, T.R., Gensel, J.C. & Seifert, A.W. Macrophages are necessary for epimorphic regeneration in African spiny mice. *Elife* **6**(2017).

44. Ostadal, B., Schiebler, T.H. & Rychter, Z. Relations between development of the capillary wall and myoarchitecture of the rat heart. *Adv Exp Med Biol* **53**, 375-388 (1975).

45. Fitzpatrick, J.R., 3rd*, et al.* Tissue-engineered pro-angiogenic fibroblast scaffold improves myocardial perfusion and function and limits ventricular remodeling after infarction. *J Thorac Cardiovasc Surg* **140**, 667-676 (2010).

46. Kikuchi, K.*, et al.* Primary contribution to zebrafish heart regeneration by gata4(+) cardiomyocytes. *Nature* **464**, 601-605 (2010).

47. Oberpriller, J.O. & Oberpriller, J.C. Response of the adult newt ventricle to injury. *J Exp Zool* **187**, 249-253 (1974).

48. Jopling, C.*, et al.* Zebrafish heart regeneration occurs by cardiomyocyte dedifferentiation and proliferation. *Nature* **464**, 606-609 (2010).

49. Chiu, L.L., Radisic, M. & Vunjak-Novakovic, G. Bioactive scaffolds for engineering
vascularized cardiac tissues. *Macromol Biosci* **10**, 1286-1301 (2010).
50. Cribbs, L.L.*, et al.* Identification of the t-type calcium channel (Ca(v)3.1d) in developing
mouse heart. *Circ Res* **88**, 403-407 (2001).
51. Balaguru, D.*, et al.* Role of the sarcoplasmic reticulum in contraction and relaxation of
immature rabbit ventricular myocytes. *J Mol Cell Cardiol* **29**, 2747-2757 (1997).
52. Kuznetsov, V., Pak, E., Robinson, R.B. & Steinberg, S.F. Beta 2-adrenergic receptor actions
in neonatal and adult rat ventricular myocytes. *Circ Res* **76**, 40-52 (1995).
582

Figure legends.

Figure 1. Characteristics of *Acomys* heart physiology and anatomy at baseline.

(A-J) Measures of body weight (BW) and heart weight in *Mus*-C57BL6 (N = 10), *Mus*-SWR

(N = 19) and *Acomys* (N = 22). Analyses demonstrate comparable heart weight across species

when normalized by body weight (BW) and tibia length (TL) (values are means ± S.E.M, *P

< 0.05, **P<0.01, and ***P<0.001 by one-way ANOVA and Dunnett correction with *Acomys*

as control). Ejection fraction (F=4.1, *P<0.05) (E), fractional shortening (F=0.96, P>0.05) (F),

left ventricular end-diastolic diameter (LVEDD) (F=14.8, **P<0.01) (G), left ventricular end-

systolic diameter (LVESD) (F=4.6) (H), left ventricular anterior wall (LVAW) (F=14.9,

*P<0.05) (I), left ventricular posterior wall (LVPW) (F=6.4) (J) (N = 30 C57BL6-*Mus*, 10

SWR-*Mus* and 10 *Acomys*). (K) Representative images showing comparable left anterior

descending artery coronary anatomy across species using intra-aortic infusion of Batson's 17

polymer mixture and corrosion casting (N = 5 each group, scale bar =1 mm). (L) Quantification

of coronary branching points of the arterial supply of the left ventricle across species. This

analysis shows comparable number of branching points across species with numerically higher

(but not statistically significant) branching points in C57BL6-*Mus* and *Acomys* compared to

SWR-*Mus*. Detailed statistical output for Figure 1 can be found in the Supplementary

Information.

Figure 2. *Acomys* cardiomyocytes exhibit features associated with *Mus* immature

cardiomyocytes.

(A) Representative images of single-cell ventricular suspension stained with DAPI (blue),

identifying small, mononuclear and binucleated cardiomyocytes in *Acomys* and larger

binucleated cardiomyocytes in *Mus*-C57BL6 with SWR cardiomyocytes being intermediate in

size between C57BL6 and *Acomys* (scale bar = 50 μm) with mononuclear and binucleated

cardiomyocytes. (B) Quantitative analyses of cell surface area of *Mus*-C57BL6 and *Acomys* (n

= 80 to 120 randomly selected CMs/animal and N = 4 each for *Mus*-C57BL6, *Mus*-SWR and

Acomys, ***P < 0.001 by one-way ANOVA; values are means ± S.E.M). (C) Percentage of

mono-, bi- and poly-nuclear cardiomyocytes in C57BL6-, SWR-*Mus* and *Acomys* (n = 80 to

120 randomly selected CMs/animal, N = 4 mice for each group, values are means ± S.E.M,

**P<0.01, ***P < 0.001 by One-way ANOVA and Dunnet post-hoc analysis, compared to

C57BL6-*Mus*). (D) Nuclear volume among isolated cardiomyocytes showing small nuclear

volume in SWR-*Mus* and *Acomys* compared to C57BL6-*Mus* (n = 80 to 120 randomly selected

CMs/animal, N = 4 mice per group, values are means ± S.E.M, ***P < 0.001 by One-way

ANOVA and Dunnet's post-hoc analysis, compared to C57BL6-*Mus*). (E) Violin plots, on left,

demonstrating the distribution of DAPI fluorescence intensity in cardiomyocyte nuclei as

assessed using Imaris on 3D (Z-stack) microscopy images. The right panel shows the

percentage of CMs within each ploidy category from the total pool of CMs from each species

(2N, 4N, and > 4N) (n = 25 to 35 randomly selected CMs/animal, N = 3 mice for each group).

Figure 3. Physiological characteristics of *Acomys* and *Mus* cardiomyocytes.

(A) Representative images of isolated cardiomyocytes from C57BL6-*Mus* and *Acomys* stained

with di-8-ANEPPS to visualize T-tubules.   (B-D) T-tubule analysis shows significantly

greater T-tubule density in transverse (B) and axial directions (C) along with wider T-tubule

spacing (D) in *Acomys* compared to *Mus* (C57BL6; t=6.2, 12, and 4.4; F=39, 141, and 19, for

panels B, C, and D, respectively; *Mus* N=5, n=67; *Acomys* N=3, n=23. (E) Current traces (left)

current-voltage relationship (right) elicited by V_{test} -25mV (left, upper) & V_{test} +15mV (left,

lower) from V_{hold} -80mV. (F, left). Current-voltage curves from V_{hold} -80mV and -50 mV

superimposed for *Acomys* CM exhibiting $I_{Ca,T}$ (A) or no $I_{Ca,T}$. (F, center) $I_{Ca,T}$ expression was

heterogeneous but more *Acomys* ventricular CMs showed $I_{Ca,T}$ compared to rare occurrences

in *Mus*-C57BL6.   (F, right) Cell capacitance was not different for *Acomys* CM with or without

$I_{Ca,T}$. (G) Cell capacitance was greater in *Mus*-C57BL6 compared to *Acomys* (t=2.88, F=7.9,

**P<0.01). (H) Maximal conductance density trended greater in *Mus*-C57BL6 compared to

Acomys (t=2.0, F=7.7, P=0.06). (I-J) Voltage-dependent activation and inactivation of $I_{Ca,L}$ was

significantly shifted positive for *Acomys* compared to *Mus*-C57BL6 (t=5.8 and 5.2, F=3.7 and

4.5, for panels I and J, respectively, ****P<0.001). (K) Representative calcium transients from

isolated ventricular cardiomyocytes loaded with fura2-AM, *Mus*-C57BL6 (top, red) and

Acomys (bottom, blue) paced at 1 Hz. Scale bar: 2 seconds. (K, right) Amplitude of the

transients (t=3.7, F=1.4, ***P = 0.0004). Velocity at which calcium enters the cytosol (upstroke

of the transient: t=2.1, F=1.5, *P=0.04). (τ) Calcium transient decay (t=3.6, F=1.5,

***P=0.0005). (L) Representative calcium transients treated with 100 nM isoproterenol (ISO),

Mus-C57BL6 (top, red) and *Acomys* (bottom, blue), paced at 1 Hz. Scale bar: 2 seconds. (Left)

Before and after ISO Amplitude of the transients (t=2.8 for Mus, P=0.53, t=0.7 for Acomys,

**P= 0.007). (Middle) Before and after ISO Velocity at which calcium enters the cytosol (t=2.5,

P = 0.02). (Right) Calcium transient decay (t=3.4 for *Mus*, **P = 0.002; N=7 animals, n=37

cells for C57BL6-*Mus*; and N=4 animals, n=31 cells for *Acomys)*.

Figure 4. Myocardial ischemic injury response in Acomys and Mus.

(A) Representative M-mode echocardiography images of *Mus*-C57BL6) (left), *Mus*-SWR

(middle) and *Acomys* (right) at baseline (upper images) and day 50 after MI (lower images).

Delta (Δ) ejection fraction (EF) (B), Δ left ventricular end-diastolic diameter (LVEDD) (C),

Δ left ventricular end-systolic diameter (LVESD) (D), Δ left ventricular end-diastolic volume

(LVEDV) (E), and Δ left ventricular end-systolic volume (LVESV) (F) after injury, calculated

as difference between each time point and baseline value in the same animal (N = 16-45 *Mus*-

C57BL6, 10-20 *Mus*-SWR and 8-11 *Acomys*) (Values are means ± S.E.M, *P < 0.05 by Mixed

-effects ANOVA, compared to *Mus*-C57BL6 or [#]*Mus*-SWR). (G) Quantitation of heart weight

(HW) normalized by body weight (BW) for *Mus*-C57BL6 (N = 2-26 animals), *Mus*-SWR (N

= 3-11 animals) and *Acomys* (N = 2-13 animals) at baseline and various timepoints after MI

661 suggesting maintenance of HW/BW ratio in *Acomys* and progressive increase in *Mus* (Values

662 are means ± S.E.M, $P < 0.05$ by two-way ANOVA, compared to *Mus-C57BL6 or [#]*Mus*-SWR).

663 (H) Quantitation of wet-dry lung weight normalized by body weight of *Mus*-C57BL6 (2-26

664 animals), *Mus*-SWR (3-11 animals) and *Acomys* (1-13 animals) at various timepoints after MI

665 suggesting maintenance of wet-dry lung weight/body weight ratio in *Acomys* and progressive

666 increase in *Mus* (Values are means ± S.E.M, *$P < 0.05$ by two-way ANOVA, compared to

667 *Mus-C57BL6 or [#]*Mus*-SWR). (I) Kaplan-Meier analysis depicting mortality after MI and

668 showing significantly lower mortality in *Acomys* compared with *Mus* (Gehan-Bareslow-

669 Willcoxon test, $P<0.05$). (J) Representative cardiac magnetic resonance (CMR) images

670 showing comparably medium-sized infarct area in both species at three days post MI (arrows

671 show the boundaries of late gadolinium enhancement/injury). The progression of infarct area

672 was significantly slower in *Acomys* compared to *Mus* at 14 days after MI (N= 9-10 animals per

673 group, t=3.28, *$P< 0.05$ by paired T-test).

675 **Figure 5. *Acomys* exhibit cardiac tissue preservation in response to myocardial infarction.**

676 (A) Representative images of Masson trichrome stained long-axis left ventricular cavity

677 sections of *Mus* (C57BL6 and SWR) and *Acomys* 50 days post-MI (scale bar = 500 μm).

678 Quantification of the infarct size, corresponding to the ratio between infarcted length and left

679 ventricular length showing significantly smaller scar in *Acomys* compared to *Mus* strains (N=

680 17 *Mus*-C57BL6, 11 *Mus*-SWR, and 5 *Acomys,* values are means ± S.E.M, *$P< 0.05$ by one-

681 way ANOVA and Tukey multiple comparison test). (B) Representative images of picrosirius

682 red stained scar center 50 days after MI across species showing alignment of collagen fibers in

683 *Acomys* compared to *Mus* species. Both *Mus* strains show highly compressed, parallel collagen

684 fibers, while *Acomys* fibers show more porosity between fibers and a wavy organization.

685 Quantification of minimum and maximum scar thickness showing that *Acomys* has on average

686 a comparatively thicker scar compared to *Mus* strains (N= 3 animal/group, values are means ±

S.E.M, **P< 0.01 by one-way ANOVA and Tukey multiple comparison test). There was no

statistically significant difference in maximum scar thickness. (C) DAPI images were obtained

in the center of the scar and demonstrate higher cellularity in *Acomys* compared to *Mus* strains

(scale bar = 100 μm in upper panel and 10 μm in lower panel, yellow boxes indicating insets).

Quantitative analysis showed higher density of nuclei and scar percentage cellularity in *Acomys*

vs. *Mus* strains (N= 3 animal/group, values are means ± S.E.M, **P< 0.01 by one-way ANOVA

and Tukey multiple comparison test).

Figure 6. Cardiac scar tissue exhibits limited expansion in *Acomys*

(A) Representative images of Masson trichrome stained long-axis left ventricular cavity

sections of *Mus* (C57BL6 and SWR) and *Acomys* in uninjured and MI animals at 7-, 17- and

50-days post-MI showing the progression of scar across species (scale bar = 500 μm). The

lower panels represent the boxed areas and show the composition of the scar center at each

stage. (B and C) Quantification of the infarct size, corresponding to the ratio between infarcted

length and left ventricular length showing significantly smaller scar in *Acomys* compared to

Mus strains (N= 17 *Mus*-C57BL6, 11 *Mus*-SWR, and 5 *Acomys,* values are means ± S.E.M,

*P< 0.05 by one-way ANOVA and Tukey multiple comparison test). (D) Delta change in scar

size among species showing the reduced delta in *Acomys* compared to *Mus*.

Figure 7. Enhanced angiogenesis and vessel maturation in the infarct region following MI

in *Acomys* compared to *Mus* strains.

(A) Representative images of isolectin staining (red) defining endothelial cells in the peri-

infarct region and the center of the scar at the mid left ventricular cavity level of *Mus* (C57BL6

and SWR) and *Acomys* 50 days post-MI (scale bar = 25 μm). Quantification of capillary density

showing significantly higher capillary density in *Acomys* compared to *Mus*-C57BL6 in the peri-

infarct region and to both *Mus* strains in the center of the scar (N= 4 *Mus*-C57BL6, 4 *Mus*-

SWR, and 5 *Acomys,* values are means ± S.E.M, *P< 0.05 by one-way ANOVA and Tukey

multiple comparison test). (B) Representative images of isolectin (red) and EdU staining

(white) marking proliferating endothelial cells in the peri-infarct region and the center of the

scar at the mid left ventricular cavity level of *Mus* (C57BL6 and SWR) and *Acomys* 50 days

post-MI (scale bar = 25 μm). Quantification of isolectin+/EdU+ cells showing significantly

higher endothelial cell proliferation in *Acomys* compared to the two *Mus* strains in both peri-

infarct and scar center areas (N= 4 *Mus*-C57BL6, 4 *Mus*-SWR, and 5 *Acomys,* values are means

± S.E.M, $F = 28.4$, *P< 0.05 by one-way ANOVA and Tukey multiple comparison test). (C)

Representative images of smooth muscle actin staining (red) in the scar center region showing

higher prevalence of medium sized blood vessels in *Acomys* compared to *Mus* strains.

Autofluorescence (green) was used to identify the infarct boundary (scale bar = 25 μm).

Figure 1

Figure 2

Figure 3

Figure 5

Figure 6

A

Uninjured | Day 7 | Day 17 | Day 50

C57BL6

SWR

Acomys

☐ C57BL6
☐ SWR
☐ *Acomys*

B Day 7

Infarct Percentage

* ** ***

C Day 17

Infarct Percentage

* ** ***

D Chronological scar size

Infarct percentage

Days post-MI

Figure 7

C