**Running head:** Rainfall control on detritivore activity

**Title:** Synergistic interactions between detritivores disappear under reduced rainfall

**Author names and affiliations:** François-Xavier Joly[*,a], Euan McAvoy[a], Jens-Arne Subke[a]

[a] Biological and Environmental Sciences, School of Natural Sciences, University of Stirling,

Stirling, Scotland, UK

**Corresponding author:** François-Xavier Joly; francois-xavier.joly [at] stir.ac.uk

**Abstract**

Understanding the consequences of altered rainfall patterns on litter decomposition is critical

to predicting the feedback effect of climate change on atmospheric $CO_2$ concentrations.

While their effect on microbial decomposition received considerable attention, their effect on

litter fragmentation by detritivores, the other dominant decomposition pathway, remains

largely unexplored. Particularly, it remains unclear how different detritivore species and their

interactions responds to changes in rainfall quantity and frequency. To fill this knowledge

gap, we determined the contribution to litter decomposition of two detritivore species

(millipede and isopod), separately and in combination, under contrasting rainfall quantity and

frequency in a temperate forest. Although halving rainfall quantity and frequency decreased

top-soil moisture by 7.8 and 13.1%, respectively, neither millipede- nor isopod-driven

decomposition were affected by these changes. In contrast, decomposition driven by both

detritivore species in combination was 65.5% higher than expected based on monospecific

treatments under high rainfall quantity, but unchanged or even lower under low rainfall

quantity. This indicate that while detritivore activity is relatively insensitive to changes in

rainfall patterns, large synergistic interactions between detritivore species may disappear

under future rainfall patterns. Incorporating interspecific interactions between decomposers

thus seems critical to evaluate the sensitivity of decomposition to altered rainfall patterns.

**Keywords**

Carbon cycling – Climate change – Isopod – Litter decomposition – Millipede – Precipitation

regime – Rainfall pattern – Soil moisture

**Introduction**

Understanding the consequences of climate change for processes underpinning the

global carbon cycle is essential to predicting its feedback effect on atmospheric $CO_2$

concentrations. This critical challenge prompted major research efforts evaluating the

response of plant litter decomposition – the largest flux of $CO_2$ to the atmosphere

(Schlesinger 2005) – to expected changes in climate change. In particular, predictions of

altered rainfall patterns with larger but less frequent rainfall events leading to enhanced

probabilities of drought and heavy rains (IPCC 2013) led numerous studies to evaluate how

decomposition responds to various rainfall manipulation. To date, the vast majority of these

studies has focused on the response of microbial decomposition (Sanaullah et al. 2012,

Walter et al. 2013, Joly et al. 2017), and reported strong negative responses of microbial

decomposition to reduced rainfall quantity and frequency. In contrast, the response of

detritivorous soil animals to altered rainfall patterns remains poorly documented.

Litter-feeding soil invertebrates (detritivores hereafter) play a major role in organic

matter turnover by consuming large amounts of decomposing litter (García-Palacios et al.

2013) and returning most of it to the soil as feces (David 2014), thereby facilitating further

microbial decomposition (Joly et al. 2018). Despite the importance of this pathway, its

response to expected changes in rainfall patterns is difficult to predict. Among the few studies

that evaluated this response, Joly et al. (2019) found that the feeding activity of one isopod

species (*Armadillidium vulgare*) was not affected by large changes in rainfall quantity, but

controlled by rainfall frequency with higher activity at low rainfall frequency. In contrast,

Coulis et al. (2013) reported a limited decrease in the feeding activity of one millipede

species (*Glomeris marginata*) in response to reduced rainfall quantity and frequency. These

contrasted findings suggest that the influence of rainfall pattern on detritivore-driven

decomposition may largely depend on detritivore species, but species-specific sensitivities to

rainfall patterns are unclear. Additionally, because both studies were performed under

laboratory conditions with detritivores collected from populations commonly exposed to

drought (Mediterranean shrubland and dryland, respectively), there are so far no data on the

importance of this rainfall pattern control under field conditions in mesic ecosystems.

In addition to interspecific differences in the response of feeding activity to rainfall

pattern, the contribution of detritivores to decomposition also depends on the nature of

interspecific interactions between cooccurring detritivores (Gessner et al. 2010). Several

studies reported that when different soil animal species cooccur, their contribution to

decomposition can be larger than predicted based on monospecific treatments (Heemsbergen

et al. 2004, Hedde et al. 2010, DeOliveira et al. 2010). These synergistic effects of detritivore

diversity may contribute substantially to decomposition, but the response of these interactions

to climatic change is largely unknown. Theory predicts that as environmental stress increases,

interspecific interactions switch from competition to complementarity and facilitation

(Bertness and Callaway 1994). This stress-gradient hypothesis implies that an increase in

environmental stress could drive a switch from antagonistic or non-additive diversity effects

on ecosystem functioning to synergistic effects. The hypothesis is relatively well supported

for plant-plant interactions and associated primary productivity (He et al. 2013), but remains

poorly considers for interactions between soil organisms and associated decomposition

processes, with the rare studies focusing on interactions between detritivore diversity and

rainfall pattern reporting contrasted results (Collison et al. 2013, Coulis et al. 2015).

Examining the nature of interactions between detritivores and their response to altered

rainfall patterns thus appears critical for accurate predictions of decomposition under future

climate change scenarios.

We evaluated the consequences of halving rainfall quantity and frequency on the

feeding activity of two common detritivore species from distant phylogenetic groups

(millipedes and isopods), and on their interactions. We hypothesized that (H1) both

detritivores species are more strongly affected by reductions in rainfall frequency than

quantity, and that (H2) synergistic interactions between detritivore species increase with

reduced rainfall quantity and frequency. We tested these hypotheses by measuring the

contribution of millipedes (*Glomeris marginata*) and isopods (*Armadillidium vulgare*),

separately and in combination, to leaf litter decomposition during a six-week field experiment

in a Scottish temperate forest during which we fully manipulated rainfall patterns.

**Methods**

We conducted this experiment in a mixed woodland of the University of Stirling

(56°08'34''N, 3°55'12''W; 40 m a.s.l), located within the Scottish Lowlands, UK. The climate

is characterized by mean annual precipitation of 1019 mm and a mean annual temperature of

8.5 °C (MetOffice, 1981-2010). The months of August and September, during which we ran

this experiment, have mean monthly precipitation of 72.4 and 90.6 mm, respectively and

have 11.4 and 11.9 days with rainfall > 1 mm, respectively. The vegetation at the site consists

in a mixture of broadleaf and coniferous tree species, including *Acer pseudoplatanus* L.,

*Fagus sylvatica* L., *Aesculus hippocastanum* L., *Picea sitchensis* (Bong.) Carr, and *Larix*

*decidua* Mill. In May 2018, we collected two detritivore species from nearby woodlands,

including the millipede *Glomeris marginata* (Villers, 1789) in a woodland near Peebles,

Scotland, UK (55°38'46''N, 3°07'55''W), and the isopod *Armadillidium vulgare* (Latreille,

1804) in a coastal woodland near Dunfermline, Scotland, UK (56°01'35''N 3°23'14''W). We

chose these species as they are widespread across Europe, occurring in diverse ecosystems

from Mediterranean to temperate. Detritivores were kept in plastic boxes and fed with moist

litter until use. In July 2018, we collected *Aesculus hippocastanum* leaf litter from the study

site. We used decomposing leaf litter rather than freshly senesced litter because detritivores

prefer feeding on partially decomposed litter (David and Gillon, 2002). We air-dried the

litter, cleaned it of debris (twigs, non-targeted litter species), and stored it in paper bags until

the start of the experiment. This litter consisted in 0.465 g C g$^{-1}$ litter and 0.0089 g N g-1

litter, for a C:N ratio of 52.0.

We manipulated rainfall quantity (18 vs. 36 mm/month), rainfall frequency (one vs.

two pulses a week) and detritivore community composition (control without detritivore;

millipedes only; isopods only; millipedes and isopods together) in a full-factorial design of

field mesocosms during a six-week period in August and September 2018. Each treatment

combination was replicated five times for a total of 80 mesocosms (2 rainfall quantities × 2

rainfall frequencies × 4 detritivore treatments × 5 replicates). To do so, we set up five

replicate full-rainout shelters (6.25 m$^2$ surface area), consisting of 2.5 × 2.5 m wooden frames

covered by a transparent polythene sheet (0.127 mm thick, 508 g/m$^2$), elevated at 0.75 m

above the soil surface with wooden legs in corners to allow air passage. To prevent rainwater

from accumulating in the middle of the polythene sheet, we placed a 1 m stake in the center

of each shelter to elevate the sheet and allow the rainwater to runoff. Under each shelter, we

set up four subplots (0.5 m$^2$ surface area); one per rainfall treatment. These subplots consisted

in 0.7 × 0.7 m wooden frames placed on the soil surface and inserted 5 cm into the soil. We

placed these frames such that they were 40 cm away from the shelter edges to avoid moisture

from shelter run-off, and 20 cm from one another. Within subplots, we removed the existing

litter layer and inserted four mesocosms (one per detritivore treatment) made of PVC pipe (9

cm long; 16 cm diameter) 7 cm into the soil. Because detritivores shelter during dry

conditions to avoid desiccation, we added small tunnels made of PVC pipe (35 mm long; 35

mm diameter) buried halfway, lengthwise, in the middle of each mesocosm. We then added

$6.0 \pm 0.01$ g of air-dried leaf litter to each mesocosm. We converted air-dried litter mass into

60°C dry mass by weighing air-dried litter subsamples, drying them at 60 °C for 48 h, and

reweighing them to obtain dry mass. In each subplot, we randomly attributed a detritivore

treatment to mesocosms and added either (1) no detritivore for the control treatment without

detritivore, or $0.6 \pm 0.01$ g of (2) millipede, (3) woodlice, or (4) millipede and woodlice.

Because of variation in body size, the number of detritivores per microcosm varied between 5

and 6. To prevent detritivores from escaping while allowing evaporation, we covered

mesocosms with a $2 \times 2$ mm nylon mesh secured with rubber bands. Within each subplot, we

also added leaf litter between mesocosms, in a similar density as within mesocosms to ensure

homogeneous surface conditions (including desiccation) throughout the subplot. We

randomly attributed a rainfall treatment to each subplot for each rainout shelter. Rainfall

quantity and frequency treatments consisted in 18 and 36 mm/month, with the equivalent

weekly amounts added as large pulses once a week or smaller pulses twice a week. These

quantities approximately reflected the actual throughfall (ca. 42 mm/month) during the

incubation period and a 50% reduction (ca. 21 mm/month), respectively. This throughfall was

estimated by correcting the mean monthly rainfall by an estimated 50% rainfall losses due to

canopy interception and stemflow (Johnson 1990). Similarly, the frequencies approximately

reflected the actual number of days with $> 1$ mm of throughfall and a 50% reduction,

respectively. We added the water pulses for both frequency treatments on Monday mornings

and those for the high frequency treatment additionally on Thursday afternoons. These pulses

were applied to mesocosms by watering subplots with a Knapsack pressure sprayer (PS16,

Kingfisher). We monitored the moisture of the topsoil (top 1 cm) throughout the incubation

by collecting soil samples from all subplots on Mondays and Thursdays, before and after

each watering events. Soil samples were weighed, dried at 60°C for 48 h and weighed again

to determine their water content. After six weeks of incubation, detritivores were collected,

counted and weighed, and remaining leaf litter was collected carefully, dried at 60°C for 48

h, and weighed. All decomposed litter samples and five samples of initial litter were ground

with a disc mill and analyzed for C concentration with a flash CHN Elemental Analyser

(Flash Smart, ThermoScientific).

The percentage of C lost after the incubation was calculated as

$[(M_i \times C_i - M_f \times C_f) / (M_i \times C_i)] \times 100$        (Equation 1)

where $M_i$ and $M_f$ are the initial and final 60°C dry masses, respectively, and $C_i$ and $C_f$ are the

initial and final C concentrations, respectively. We used litter C loss rather than total litter

mass loss to correct for inorganic contamination of leaf litter retrieved from mesocosms

where they were in direct contact with soil. To isolate detritivore from microorganism effects

on total C loss, we subtracted the C loss without detritivore from the C loss with detritivores

(Bocock's formula; David 1998). This detritivore-driven litter C loss was then corrected for

slight differences in detritivore mass among mesocosms by dividing it by the detritivore mass

for each mesocosm (average throughout the incubation) and multiplying it by the mean

detritivore mass across all mesocosms with detritivores (0.596 g). For mesocosms including

both millipede and isopods, the relative mixing effect was calculated as

[(observed value - expected value)/expected value] $\times 100$   (Equation 2)

for each rainfall treatment separately, where the expected value is the mean of litter C loss

driven by millipede and isopods separately, in monospecific treatments. Fourteen mesocosms

for which we noted anomalies upon harvest (e.g. presence of snails or lower numbers of

detritivores than initially added) were not used for response variable calculations and further

data analyses.

We used two-way ANOVAs to evaluate the effect of rainfall quantity and frequency

and their interaction on (1) soil moisture, on the contribution of (2) millipedes, (3) isopods,

(4) millipedes and isopods to litter C loss, and on (5) the relative effect of mixing millipede

and isopods. To determine differences in detritivore-driven decomposition among detritivore

treatments across all rainfall treatments, we evaluated the effect of detritivore community

composition on detritivore-driven decomposition with a one-way ANOVA. For all tests,

Tukey HSD post hoc mean comparisons identified difference among treatments. Student's *t*

tests were used to determine if the relative mixing effect for each rainfall treatment was

significantly different from zero. All statistical analyses were performed using R version

3.6.1.

**Results**

The exclusion of natural rainfall and application of simulated rainfall according to

four treatments (current rainfall vs. 50% reduction, each delivered at current frequency vs.

50% reduction) led to variable mean soil moistures throughout the six-week incubation,

ranging from 0.25 g $H_2O$ g soil$^{-1}$ for the low rainfall quantity treatment delivered at low

frequency, to 0.31 g $H_2O$ g soil$^{-1}$ for the high rainfall quantity treatment delivered at high

frequency (Fig. 1). Halving rainfall quantity and frequency led to significant relative

decreases in mean soil moisture by 7.8% (P < 0.05) and 13.1% (P < 0.001; Fig. 1).

After six weeks of incubation in the field, millipedes contributed significantly more to

decomposition than isopods (P < 0.001), with an average detritivore-driven litter C loss of

26.2% for millipedes compared to 10.0% for isopods (Fig. 2), across rainfall treatments. In

these monospecific treatments, neither millipede- nor isopod-driven litter C loss were

significantly affected by halving rainfall quantity (P = 0.08 and P = 0.62 for millipede- and

isopod-driven litter C loss, respectively) or frequency (P = 0.79 and P = 0.76 for millipede-

and isopod-driven litter C loss, respectively; Fig. 2). When millipede and isopods cooccurred,

their combined contribution to litter C loss reached 22.1%, i.e. a level similar to the

contribution of millipedes to litter C loss, but significantly higher than that of isopods ($P <$

$0.001$; Fig. 2). In contrast to monospecific treatments, detritivore-driven litter C loss was

negatively affected by halving rainfall quantity ($P < 0.05$) by not frequency ($P = 0.57$), and

ranged from 15.3% for the low rainfall quantity treatment delivered at low frequency to

27.4% for the high rainfall quantity treatment delivered at low frequency (Fig. 2).

The relative effect of mixing isopods and millipedes, i.e. the relative difference

between observed values of detritivore-driven litter C loss when millipede and isopods

cooccurred and predicted values based on the respective monospecific detritivore treatments,

ranged from -15.4% for the low rainfall quantity treatment delivered at low frequency to

+70.1% for the high rainfall quantity treatment delivered at high frequency (Fig. 3). The

mixing effect was negatively affected by halving rainfall quantity ($P < 0.001$) but not

significantly affected by halving rainfall frequency ($P = 0.16$). Except for the low rainfall

quantity and high frequency treatment, relative mixing effects were significantly different

from zero (Fig. 3).

**Discussion**

The lack of response of either millipede- or isopod-driven decomposition to halving

rainfall quantity supports the view that detritivore feeding activities are largely insensitive to

changes in rainfall quantity. This concords with previous studies focusing on our two studied

species, separately, that reported no response of detritivore-driven decomposition for the

millipede species (Coulis et al. 2013) and isopod species (Joly et al. 2019) to altered rainfall

patterns. Our results indicate that this pattern observed in these studies performed under

controlled laboratory conditions, also applies under field conditions. Furthermore, while these

previous studies used detritivores collected from drought-prone ecosystems (Mediterranean

shrublands and drylands, respectively), the use of detritivores from temperate ecosystems

which are not commonly exposed to drought (Scottish Lowlands) in our study suggests that

this lack of sensitivity to altered rainfall patterns does not result from local adaptations and

reflects species rather than population adaptations. Possibly, the capacity to coil into a sphere

shared by both the millipede and isopod species used in this study (a feature known as

'volvation' that gave them the vernacular name of common pill millipede and common pill

woodlice, respectively) may underpin their resistance to desiccation. It is important to note

that despite halving rainfall quantity, we only reported a limited yet significant reduction of

topsoil moisture (-7.8%), which may explain the lack of detritivore response to reduced

rainfall. On the other hand, halving rainfall frequency induced a larger reduction in topsoil

moisture (-13.1%) but also had no effect on either millipede- or isopod-driven

decomposition, unlike the expectation in our first hypothesis. It also contrasts with the

previous findings that isopod-driven decomposition increases with decreasing rainfall

frequency (Joly et al. 2019). In turn, strong differences in detritivore-driven decomposition

between millipede and isopod treatments suggest that detritivore community composition

rather than rainfall patterns controls detritivore-driven decomposition. Such differences may

be due to distinct consumption rates between detritivore species, and the higher detritivore-

driven decomposition for the millipede compared to the isopod species is in line with

previously reported differences in consumption rate for these two species (Coulis et al. 2015).

While altered rainfall patterns may have limited effects on detritivore-driven decomposition,

effects on detritivore community composition (Zimmer 2005) may have important

consequences for decomposition. Furthermore, there is a strong need to consider multiple

global change factors (Rillig et al. 2019) such as combined changes in temperature and

rainfall that may have interactive effects on the detritivore-driven decomposition (Thakur et

al. 2018).

A key result of our study is that interactions between detritivore species markedly

varied depending on rainfall pattern with the effect of mixing millipedes and isopods on

decomposition ranging from antagonistic (-15.4%) at low levels of rainfall quantity and

frequency to synergistic (+70.1%) at high levels of rainfall quantity and frequency (Fig. 3),

contrary to our second hypothesis. The magnitude of these effects indicates that important

facilitative interactions between detritivores may be at play and that detritivore diversity may

be a strong factor driving plant litter decomposition. While such interactions were previously

reported (Heemsbergen et al. 2004, DeOliveira et al. 2010), this aspect of diversity on

decomposition remains poorly investigated compared with litter diversity (Gessner et al.

2010, Handa et al. 2014). Importantly, our study suggests that the nature and magnitude of

such interactions are strongly dependent on environmental conditions, as evidenced by the

change from large positive interactions to non-additive or even negative interactions

associated with the reduction in rainfall quantity and frequency. This finding from our field

experiment contrasts with laboratory studies that reported no response of detritivore

interactions to changes in moisture treatments (Collison et al. 2013, Coulis et al. 2015). Such

discrepancies may be due to the consideration of a reduction in rainfall frequency only

(Coulis et al. 2015), or due to a lower study duration (two weeks in Collison et al. (2013)).

More generally, our results do not support the stress-gradient hypothesis (Bertness and

Callaway 1994, He et al. 2013) and suggests that rather than mitigating predicted changes in

climate change ("insurance hypothesis",  Yachi and Loreau 1999) positive interspecific

interactions on ecosystem functioning may be vulnerable to predicted changes in climate

change. Understanding the mechanisms underpinning these detritivore interactions and their

sensitivity to rainfall, at larger temporal, spatial and phylogenetic scales thus represents an

urgent research avenue.

275            Collectively, our results indicate that the feeding activities of the detritivore species

*Armadillidium vulgare* and *Glomeris marginata* are relatively insensitive to changes of in

rainfall patterns, but that their interspecific interactions may switch from synergistic to

antagonist under reduced rainfall quantity and frequency. Although these species are

common across Europe and cooccur in a wide range of ecosystems, our findings may not

extend to other detritivore species and their interactions. Evaluating the response of diverse

detritivore species with different phylogenetic origins and distribution, and their interactions

will be necessary to identify general patterns in the climatic sensitivity of detritivores and

their interactions, thereby better predicting the consequences of climate change on

decomposition. To this end, research efforts aiming to identify morphological traits

underpinning such sensitivities appear particularly promising (e.g. Dias et al. (2013) for

isopods). Last, considering interspecific interactions between decomposers appears critical to

evaluating the sensitivity of decomposition to altered rainfall patterns.

**Acknowledgments**

We thank Ian Washbourne for laboratory assistance. This work was funded by a British

Ecological Society Small Research grant (SR18/1215) and by a UK Natural Environment

Research Council grant (NE/P011098/1).

**References**

Bertness, M. D., and R. Callaway. 1994. Positive interactions in communities. Trends in

296            Ecology and Evolution 9:191–193.

Collison, E. J., T. Riutta, and E. M. Slade. 2013. Macrofauna assemblage composition and

soil moisture interact to affect soil ecosystem functions. Acta Oecologica 47:30–36.

Coulis, M., N. Fromin, J.-F. David, J. Gavinet, A. Clet, S. Devidal, J. Roy, and S.

Hättenschwiler. 2015. Functional dissimilarity across trophic levels as a driver of soil

processes in a Mediterranean decomposer system exposed to two moisture levels. Oikos

124:1304–1316.

Coulis, M., S. Hättenschwiler, N. Fromin, and J.-F. David. 2013. Macroarthropod-

microorganism interactions during the decomposition of Mediterranean shrub litter at

different moisture levels. Soil Biology & Biochemistry 64:114–121.

David, J. F. 1998. How to calculate leaf litter consumption by saprophagous macrofauna?

European Journal of Soil Biology 34:111–115.

David, J. F. 2014. The role of litter-feeding macroarthropods in decomposition processes: A

reappraisal of common views. Soil Biology and Biochemistry 76:109–118.

DeOliveira, T., S. Hättenschwiler, and I. T. Handa. 2010. Snail and millipede

complementarity in decomposing Mediterranean forest leaf litter mixtures. Functional

Ecology 24:937–946.

Dias, A. T. C., E. J. Krab, J. Mariën, M. Zimmer, J. H. C. Cornelissen, J. Ellers, D. A.

Wardle, and M. P. Berg. 2013. Traits underpinning desiccation resistance explain

distribution patterns of terrestrial isopods. Oecologia 172:667–677.

García-Palacios, P., F. T. Maestre, J. Kattge, and D. H. Wall. 2013. Climate and litter quality

differently modulate the effects of soil fauna on litter decomposition across biomes.

Ecology Letters 16:1045–1053.

Gessner, M. O., C. M. Swan, C. K. Dang, B. G. McKie, R. D. Bardgett, D. H. Wall, and S.

Hättenschwiler. 2010. Diversity meets decomposition. Trends in Ecology and Evolution

25:372–380.

Handa, I. T., R. Aerts, F. Berendse, M. P. Berg, A. Bruder, O. Butenschoen, E. Chauvet, M.

O. Gessner, J. Jabiol, M. Makkonen, B. G. McKie, B. Malmqvist, E. T. H. M. Peeters, S.

Scheu, B. Schmid, J. Van Ruijven, V. C. A. Vos, and S. Hättenschwiler. 2014.

Consequences of biodiversity loss for litter decomposition across biomes. Nature

509:218–221.

He, Q., M. D. Bertness, and A. H. Altieri. 2013. Global shifts towards positive species

interactions with increasing environmental stress. Ecology Letters 16:695–706.

Hedde, M., F. Bureau, M. Chauvat, and T. Decaëns. 2010. Patterns and mechanisms

responsible for the relationship between the diversity of litter macro-invertebrates and

leaf degradation. Basic and Applied Ecology 11:35–44.

Heemsbergen, D. A., M. P. Berg, M. Loreau, J. R. Van Hal, J. H. Faber, and H. A. Verhoef.

2004. Biodiversity effects on soil processes explained by interspecific functional

dissimilarity. Science 306:1019–1020.

IPCC. 2013. Climate Change 2013: The Physical Science Basis. Contribution of Working

Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate

Change. Page (T. F. Stocker, D. Qin, G. K. Plattner, M. Tignor, S. K. Allen, J.

Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley, Eds.). Cambridge Univ.

Press, Cambridge, UK, and New York, p. 1535.

Johnson, R. C. 1990. The interception, throughfall and stemflow in a forest in highland

Scotland and the comparison with other upland forests in the UK. Journal of Hydrology

118:281–287.

Joly, F.-X., S. Coq, M. Coulis, J. Nahmani, and S. Hättenschwiler. 2018. Litter conversion

into detritivore faeces reshuffles the quality control over C and N dynamics during

decomposition. Functional Ecology 32:2605–2614.

Joly, F.-X., K. L. Kurupas, and H. L. Throop. 2017. Pulse frequency and soil-litter mixing

alter the control of cumulative precipitation over litter decomposition. Ecology 98:2255–

2260.

Joly, F.-X., A. K. Weibel, M. Coulis, and H. L. Throop. 2019. Rainfall frequency, not

350            quantity, controls isopod effect on litter decomposition. Soil Biology and Biochemistry

351            135:154–162.

352      Rillig, M. C., M. Ryo, A. Lehmann, C. A. Aguilar-Trigueros, S. Buchert, A. Wulf, A.

353            Iwasaki, J. Roy, and G. Yang. 2019. The role of multiple global change factors in

354            driving soil functions and microbial biodiversity. Science 366:886–890.

355      Sanaullah, M., C. Rumpel, X. Charrier, and A. Chabbi. 2012. How does drought stress

356            influence the decomposition of plant litter with contrasting quality in a grassland

357            ecosystem? Plant and Soil 352:277–288.

358      Schlesinger, W. H. 2005. Biogeochemistry. Page Treatise on Geochemistry. Elsevier, New

359            York.

360      Thakur, M. P., P. B. Reich, S. E. Hobbie, A. Stefanski, R. Rich, K. E. Rice, W. C. Eddy, and

361            N. Eisenhauer. 2018. Reduced feeding activity of soil detritivores under warmer and

362            drier conditions. Nature Climate Change 8:75–78.

363      Walter, J., R. Hein, C. Beierkuhnlein, V. Hammerl, A. Jentsch, M. Schädler, J. Schuerings,

364            and J. Kreyling. 2013. Combined effects of multifactor climate change and land-use on

365            decomposition in temperate grassland. Soil Biology and Biochemistry 60:10–18.

366      Yachi, S., and M. Loreau. 1999. Biodiversity and ecosystem productivity in a fluctuating

367            environment: the insurance hypothesis. Proceedings of the National Academy of

368            Sciences 96:1463–1468.

369      Zimmer, M. 2005. Effects of temperature and precipitation on a flood plain isopod

370            community: A field study. European Journal of Soil Biology 40:139–146.

 **Figure legends**

Figure 1: Mean moisture of the topsoil (top 1 cm) throughout the six-week incubation, under

two different rainfall quantities (18 and 36 mm/month) and two different rainfall frequencies

(one or two pulses per week). Lowercase letters indicate, for each detritivore treatment,

significant differences among rainfall treatments (Tukey HSD tests).

Figure 2: Percentage of litter carbon (C) loss driven by isopods, millipedes, as well as isopods

and millipedes, after a six-week incubation under two different rainfall quantities (18 and 36

mm/month) and two different rainfall frequencies (one or two pulses per week). Uppercase

letters indicate significant differences among detritivore treatments, while lowercase letters

indicate, for each detritivore treatment, significant differences among rainfall treatments

(Tukey HSD tests).

Figure 3: Relative difference between the observed detritivore-driven litter C loss for the

millipede-isopod mixture and the expected value based on monospecific treatments after a

six-week incubation under two different rainfall quantities (18 and 36 mm/month) and two

different rainfall frequencies (one or two pulses per week). Letters indicate differences

among rainfall treatments (Tukey HSD tests). Asterisks indicate a relative mixing effect

significantly different from 0 (*: P < 0.05; **: P < 0.01).