9   **Sexually divergent cortical control of affective-autonomic integration**

16   Tyler Wallace[1], Derek Schaeuble[1], Sebastian A. Pace, Morgan K. Schackmuth, Shane T.
17   Hentges, Adam J. Chicco, Brent Myers
18
19   [1]Equal contribution
20
21   *Biomedical Sciences, Colorado State University, Fort Collins, CO*

31   Abstract: 150 words
32   Introduction: 562 words
33   Discussion: 1702 words
34   Figures: 6
35   Supplement: 4 figures,1 table

Address for correspondence
Brent Myers, Ph.D.
Department of Biomedical Sciences
Colorado State University
1617 Campus Delivery
Fort Collins, CO 80523
brent.myers@colostate.edu

Abstract

Depression and cardiovascular disease reduce quality of life and increase mortality risk.

These conditions commonly co-occur with sex-based differences in incidence and severity.

However, the biological mechanisms linking the disorders are poorly understood. In the current

study, we hypothesized that the infralimbic (IL) prefrontal cortex integrates mood-related

behaviors with the cardiovascular burden of chronic stress. In a rodent model, we utilized

optogenetics during behavior and *in vivo* physiological monitoring to examine how the IL

regulates affect, social motivation, neuroendocrine-autonomic stress reactivity, and the cardiac

consequences of chronic stress. Our results indicate that IL glutamate neurons increase socio-

motivational behaviors specifically in males. IL activation also reduced endocrine and

cardiovascular stress responses in males, while increasing reactivity in females. Moreover, prior

IL stimulation protected males from subsequent chronic stress-induced sympatho-vagal

imbalance and cardiac hypertrophy. Our findings suggest that cortical regulation of behavior,

physiological stress responses, and cardiovascular outcomes fundamentally differ between

sexes.

17
18 **Keywords:** infralimbic cortex, social motivation, corticosterone, chronic stress, ventricular
19 hypertrophy

1. Introduction

Major depressive disorder (MDD) and cardio-metabolic conditions including

hypertension, glucose intolerance, and heart failure significantly contribute to global disease

burden. Epidemiological evidence implicates life stressors as a risk factor for both MDD and

cardiovascular disease (CVD) (Binder and Nemeroff, 2010; Grippo and Johnson, 2009; Myers

et al., 2014b; Sgoifo et al., 2015). Furthermore, sex differences in the incidence of MDD, CVD,

and MDD-CVD co-morbidity suggest that sex-specific factors contribute to outcomes (Goldstein

et al., 2019). However, the biological basis for stress effects on health, particularly the

integration of affective and physiological systems, is poorly understood. Human brain imaging

studies indicate that ventral medial prefrontal cortex (vmPFC) activity associates with sadness

and blood pressure reactivity, suggesting that top-down cortical control may integrate diverse

aspects of mood and systemic physiology.

The vmPFC is involved in numerous cognitive and emotional processes (McKlveen et

al., 2015; Myers-Schulz and Koenigs, 2012; Wood and Grafman, 2003). A subregion of the

vmPFC, the subgenual cingulate cortex (BA25), is activated by sadness-provoking stimuli,

responds to social isolation, and has reduced volume in MDD patients (Beckmann et al., 2009;

Liotti et al., 2000; Vijayakumar et al., 2017). BA25 is also targeted for deep brain stimulation in

patients with treatment-resistant depression, where larger volumes predict better treatment

outcomes (Mayberg et al., 2005; Sankar et al., 2019). Although, broader investigation of BA25

activity in mood disorders has yielded mixed results with reports of both hyper- (Hamani et al.,

2011; Mayberg et al., 2005) and hypo-activity (Drevets et al., 2008, 1997). Subgenual regions of

vmPFC have also been identified as components of a central autonomic network monitoring

visceral functions (Beissner et al., 2013; Gianaros and Sheu, 2009; Gianaros and Wager, 2015;

Myers, 2016; Shoemaker et al., 2016). Furthermore, recent pharmacological studies employing

glutamate uptake inhibitors in non-human primates have implicated BA25 activity in reduced

reward motivation (Alexander et al., 2019) and enhanced threat-related autonomic responses

(Alexander et al., 2020). The rodent putative anatomical homolog of BA25, the infralimbic cortex

(IL) (Öngür et al., 2003; Roberts and Clarke, 2019; Uylings et al., 2003; Vertes, 2004),

innervates limbic and stress-regulatory nuclei including the amygdala, thalamus, and

hypothalamus (Gabbott et al., 2005; Myers et al., 2016, 2014a; Wood et al., 2018). Additionally,

IL stimulation reduces passive coping and increases pyramidal neuron spine density in male

rodents (Fuchikami et al., 2015). Moreover, knockdown of IL glutamatergic output exacerbates

chronic stress effects on hypothalamic-pituitary-adrenal (HPA) axis reactivity and vascular

function (Myers et al., 2017; Schaeuble et al., 2019). However, the potential sex-specific roles of

IL activity to integrate socio-motivational behaviors with physiological stress reactivity and the

cardiac outcomes of chronic stress remain to be determined.

To identify how the male and female vmPFC coordinates mood-related behaviors and

cardiovascular outcomes, genetically-identified vmPFC projection neurons received temporally-

specific stimulation in an *in vivo* rodent model. Specifically, channelrhodopsin-2 (ChR2) was

expressed under the calcium/calmodulin-dependent protein kinase type II α (CaMKIIα) promoter

in the IL to permit optogenetic activation of IL pyramidal neurons in both male and female rats

(Wood et al., 2018). This approach was combined with measures of place preference and social

behavior to examine affective valence and sociability. Behavioral assessment was followed by

measures of physiological stress reactivity, including radiotelemetry and echocardiography over

the course of chronic variable stress (CVS). Ultimately, these findings identify the IL as an

affective-autonomic integrator that links motivation and stress responding divergently in males

and females.

2. Methods

2.1 Animals

Age-matched adult male and female Sprague-Dawley rats were obtained from Envigo

(Denver, CO) with male rat weight ranging from 250-300 g and female from 150-200 g. After

stereotaxic surgery, rats were housed individually in shoebox cages with cardboard tubes for

enrichment in a temperature- and humidity-controlled room with a 12-hour light-dark cycle (lights

on at 07:00h, off at 19:00h) and food and water *ad libitum*. In accordance with ARRIVE

guidelines, all treatments were randomized and experimenters blinded. All procedures and

protocols were approved by the Colorado State University Institutional Animal Care and Use

Committee (protocol: 16-6871A) and complied with the National Institutes of Health Guidelines

for the Care and Use of Laboratory Animals. Signs of poor health and/or weight loss ≥ 20% of

pre-surgical weight were *a priori* exclusion criteria. These criteria were not met by any animals

in the current experiments; however, animals were removed from experimentation if fiber optic

or radiotelemetry devices failed.

2.2 Microinjections

Rats were anesthetized with isoflurane (1-5%) and administered analgesic (0.6 mg/kg

buprenorphine-SR, subcutaneous). Rats received bilateral microinjections (Males 1.5 - 2 µL,

Females 0.75 – 1.25 µL) of adeno-associated virus (AAV) into the IL (males: 2.7 mm anterior to

bregma, 0.6 mm lateral to midline, and 4.2 mm ventral from dura, females: 2.3 mm anterior to

bregma, 0.5 mm lateral to midline, and 4 mm ventral from dura). These volumes correspond

with prior studies utilizing viral vector transduction in rat vmPFC (Ferenczi et al., 2016; Ji and

Neugebauer, 2012; Wood et al., 2018). AAV5-packaged constructs (University of North Carolina

Vector Core, Chapel Hill, NC) either expressed yellow fluorescent protein (YFP) or ChR2

conjugated to YFP under the CaMKIIα promoter to achieve pyramidal cell-predominant

expression (Wood et al., 2018). All microinjections were carried out with a 25-gauge, 2-µL

microsyringe (Hamilton, Reno, NV) using a microinjection unit (Kopf, Tujunga, CA) at a rate of 5

minutes/µL. The needle was left in place for 5 minutes before and after injections to reduce

tissue damage and allow diffusion. Skin was closed with wound clips that were removed 2

weeks after injections and animals were allowed at least 6 weeks for recovery and ChR2

expression.

2.3 Electrophysiology

Adult male rats (n = 8) were injected with AAV constructs as described above and, after

8-12 weeks, exposed to 5% isoflurane prior to decapitation and brain removal. As previously

described (Rau and Hentges, 2017), brains and sections were collected in ice-cold artificial CSF

(aCSF) consisting of the following (in mM): 126 NaCl, 2.5 KCl, 1.2 $MgCl_2$ Λ $6H_2O$, 2.4 $CaCl_2$ Λ

$2H_2O$, 1.2 NaH_2PO_4, 11.1 glucose, and 21.4 $NaHCO_3$, bubbled with 95% O_2 and 5% CO_2.

Coronal slices containing the IL were cut at a thickness of 240 µm using a model VT1200S

vibratome (Leica Microsystems, Buffalo Grove, IL). After resting 1 hr at 37°C in aCSF, slices

were transferred to the recording chamber and perfused with oxygenated 37° C aCSF at a 2

ml/min flow rate. For whole-cell recordings, the internal recording solution contained the

following (in mM): KCL 57.5, K-methyl sulfate 57.5, NaCl 20, $MgCl_2$ 1.5, HEPES 5; EGTA 0.1;

ATP 2; GTP 0.5, and phosphocreatine 10. The pH was adjusted to 7.3. Recording electrodes

had a resistance of 2 – 4 MΩ when filled with this solution. IL pyramidal neurons were identified

for recording based on the expression of ChR2-YFP under the control of CaMKIIα. Whole-cell

patch-clamp recordings were acquired in voltage-clamp at a holding potential of -60 mV using

an Axopatch 200B Amplifier (Molecular Devices, San Jose, CA). Current-clamp recordings were

acquired while holding current at 0 pA. Electrophysiological data were collected and analyzed

using Axograph X software on a Mac OS X operating system (Apple, Cupertino, CA). Light

activation of IL neurons expressing ChR2 occurred via 473 nm LED (Thorlabs, Newton, NJ) 1.1

mW light pulse driven by a LEDD1B driver (Thorlabs, Newton, NJ) triggered through the TTL

output on an ITC-18 computer interface board (HEKA Instruments, Holliston, MA). Current-

clamp experiments utilized 5, 10, and 20 Hz stimulation frequencies for 5 min bouts. Recordings

were excluded if access resistance exceeded 10 Ω during recording.

2.4 Radiotelemetry implantation

A subset of rats was instrumented with ECG-enabled radiotelemetry transmitters (HD-

S11 F0, Data Sciences International, St. Paul, MN) as previously described (Flak et al., 2011;

Goodson et al., 2017; Schaeuble et al., 2019). Briefly, rats were anesthetized with inhaled

isoflurane anesthesia (1-5%) and given a subcutaneous injection of analgesic (0.6 mg/kg

Buprenorphine-SR) and an intramuscular injection of antibiotic (5 mg/kg gentamicin). The

descending aorta was exposed via an abdominal incision, allowing implantation of a catheter

extending from the transmitter. The catheter was secured with tissue adhesive (Vetbond; 3M

Animal Care Products, St. Paul, MN) applied over a cellulose patch. ECG leads were passed

through abdominal musculature and sutured subcutaneously above the rib cage and pectoral

muscles. The transmitter body was then sutured to the abdominal musculature, followed by

closure of the abdominal and skin with suture and wound clips, respectively. Rats then

recovered for 2 weeks before wound clips were removed.

2.5 Fiber optic cannulas

Rats were anesthetized with isoflurane (1-5%) followed by analgesic (0.6 mg/kg

buprenorphine-SR, subcutaneous) and antibiotic (5 mg/kg gentamicin, intramuscular)

administration. Bilateral fiber-optic cannulas (flat tip 400/430 μm, NA = 0.66, 1.1 mm pitch with

4.5 mm protrusion for males and 4.2 mm protrusion for females; Doric Lenses, Québec,

Canada) were aligned with the IL injection sites and lowered to the ventral PL/dorsal IL

approximately 1 mm dorsal to the injection to enable optic stimulation of the IL. Cannulas were

secured to the skull with metal screws (Plastics One) and dental cement (Stoelting, Wood Dale,

IL). Skin was sutured and, following 1 week of recovery, rats were handled daily and acclimated

to the stimulation procedure for another week before experiments began. Rat handling and

cannula habituation continued daily throughout experiments.

2.6 Optogenetic stimulation

Light pulses (1.0 - 1.3 mW, 5 ms pulses, 10 or 20 Hz) were delivered through a

fiber-optic patch cord (240 μm core diameter, NA = 0.63; Doric Lenses) connected to a

473 nm LED driver (Doric Lenses). Optic power was measured with a photodiode sensor

(PM160, Thorlabs Inc, Newton, NJ) at the cannula fiber tip. Initial male experiments used

20 Hz stimulation but, following the results of slice electrophysiology experiments, all

subsequent studies used 10 Hz stimulation. Rats received a single 20-minute session of

stimulation (1 min on/1 min off) in their homecage the week prior to stress exposure. In

total, rats received self-determined stimulation in the stimulation zone of RTPP day 2,

throughout the social behavior assay, and during acute stress exposure (restraint and

novel environment), all prior to CVS. Immediately before and after CVS, ultrasound

measures were conducted with optics both off and on to isolate long-term structural

changes and acute functional changes. Further optic details are elaborated in the

specified sections below.

2.7 Estrous cycle cytology

All female rats went through experiment 3 simultaneously, housed in the same room and

randomly cycling. Immediately following acute measures of behavior and physiology, vaginal

cytology was examined to approximate the estrous cycle stage. A damp (deionized water)

cotton swab was used to collect cells from the vaginal canal and place them onto a glass slide.

When dried, slides were viewed under a 10x objective light microscope by a minimum of two

blind observers and were categorized as proestrus, estrus, metestrus, or diestrus (Cora et al.,

2015; Smith et al.; Solomon et al., 2015).

2.8 Real-time place preference

The real-time place preference (RTPP) assay was used to assess the valence of IL

stimulation (Bimpisidis et al., 2020; Stamatakis and Stuber, 2012). Cannulas were connected

via patch cords to LEDs for light delivery and rats placed in a custom-made fiberglass arena

with two chambers connected by a corridor (chambers: 15 x 15'', corridor: 8 x 6'', 15'' deep).

Rats explored the arena for 10 minutes on two consecutive days. The first day was a

habituation day and no stimulation was delivered on either side. On the second day, rats

received LED-generated 470 nm light pulses upon entry and throughout the time spent in the

assigned stimulation side. Stimulation stopped when rats exited the assigned stimulation side

but re-commenced upon re-entry. Thus, rats determined the amount of stimulation received

through time spent in the stimulation side. Trials were recorded by a camera mounted above the

arena and animal position was tracked by Ethovision software (Noldus Information

Technologies) for automated optic hardware control. Stimulation side assignment was

counterbalanced and animal testing was randomized. The time rats spent in the stimulation side

was divided by the total time and multiplied by 100 to generate a percentage of time spent in the

stimulation side.

2.9 Social behavior

A modified version of the 3-chambered social behavior assay was used to accommodate

optic patch cords (Felix-Ortiz and Tye, 2014; Moy et al., 2004). To examine social interaction,

each rat was connected to a patch cord and placed in a black rectangular fiberglass arena (36 x

23'', 15.8'' deep). Initially, the arena was empty and experimental rats were allowed to explore

for 5 minutes without optic stimulation. The experimental rat was then returned to their home

cage while an empty enclosure (ventilated with small round openings) was placed on one side

of the arena, defined as the object, and an identical enclosure containing an age- and sex-

matched conspecific was placed on the other side of the arena, defined as the social cage. The

experimental rat was then placed in the middle of the arena and allowed to explore freely for 10

minutes with 5 ms pulsatile stimulation delivered throughout to quantify social motivation. The

experimental rat was then placed again into its homecage while the empty enclosure was

replaced with a new enclosure containing a novel age- and sex-matched conspecific. The

experimental rat freely explored for 10 minutes while receiving optic stimulation to assess social

novelty preference. Behavior was recorded with an overhead camera and interactions were

defined as nose pokes onto cages and scored by a treatment-blinded observer. The duration of

interactions was divided by the total time of each interaction period and multiplied by 100 to give

a percent interaction value. Sides for object cage, social cage, and novel cage were

counterbalanced and animal order randomized.

2.10 Novel environment

In rats with radiotelemetry implants, baseline cardiovascular measurements were

collected the weekend prior to novel environment stress. Rats were then exposed to a 30-

minute novel environment stressor to assess acute cardiovascular reactivity. During novel

environment, rats were connected to fiber-optic patch cords for stimulation and placed into a

brightly-lit semi-transparent arena. Radiotelemetry receiver pads (Data Sciences International,

St. Paul, MN) were arrayed under the arena with 10 Hz optic stimulation during the stressor to

record hemodynamics and activity in 1-minute bins. Heart rate (HR), mean arterial pressure

(MAP), systolic arterial pressure (SAP), and diastolic arterial pressure (DAP) were collected and

analyzed with Ponemah software (Version:6.4x Data Sciences International).

2.11 Restraint stress

Restraint was used to examine neuroendocrine responses to acute stress. Rats were

placed in plastic film decapicones (Braintree Scientific, Braintree, MA) and connected to fiber-

optic patch cords for optic stimulation throughout the 30-minute restraint. Blood samples

(approximately 250 μL) were collected by tail clip at the initiation of restraint with additional

samples taken 15 and 30 min after (Vahl et al., 2005). At the conclusion of restraint, patch cords

were disconnected and rats returned to their homecage with recovery blood samples collected

at 60 and 90 min after the initiation of restraint. Blood glucose was determined with Contour

Next EZ glucometers (Bayer, Parsippany, NJ) and 2 independent readings for each time point

were averaged. Blood samples were centrifuged at 3000 X g for 15 minutes at 4° C and plasma

was stored at –20° C until radioimmunoassay (RIA). Plasma corticosterone levels were

measured with an [125]I RIA kit (MP Biomedicals, Orangeburg, NY) as previously described

(Myers et al., 2017). All samples for each sex were run in duplicate and all time points were run

in the same assay. RIA intra-assay coefficient of variation was 8.6% and interassay 13.6%.

2.12 Chronic variable stress

Chronic variable stress was comprised of twice daily (AM and PM) repeated and

unpredictable stressors presented in a randomized manner including: exposure to a brightly-lit

open field (28.5 x 18'' 13'' deep, 1 hour), cold room (4° C, 1 hour), cage tilt (45°, 1 hour), forced

swim (23° to 27° C, 10 minutes), predator odor (fox or coyote urine, 1 hour), restraint (1 hour),

and shaker stress (100 rpm, 1 hour). Additionally, overnight stressors were variably included,

comprised of damp bedding (400 mL water) and overnight light. During the 2 weeks of CVS,

rats were weighed every 3-4 days to monitor body weight.

2.13 Spectral analysis of heart rate variability

Throughout CVS, resting homecage recordings of heart rate variability (HRV) were

collected and analyzed using Ponemah software (Version:6.4x Data Sciences International, St.

Paul, MN). Using guidelines for heart rate variance (Electrophysiology, 1996) in the frequency

domain, low (LF) and high frequency (HF) components were collected during two-hour periods

in the AM and PM and averaged within treatment groups. Recordings were sampled at least

one hour after stressors and did not follow overnight stressors, swim, or restraint. Spectral

analysis is used to measure autonomic balance as LF predominantly represents sympathetic

and HF predominantly parasympathetic contributions to HRV (Electrophysiology, 1996; Sgoifo

et al., 2015). Accordingly, LF/HF represents net cardiac sympathetic drive.

2.14 Echocardiography

Left ventricle structure and function were assessed before and after CVS via

echocardiography. In preparation, rats were anesthetized with inhaled isoflurane (5% induction,

2% maintenance), connected to fiber optic patch cord, and shaved over the ventral thorax. A 12-

mHz pediatric transducer connected to a Phillips XD11 ultrasound system was used to image

the heart in transverse (parasternal short axis) and 4-chamber angles. M-mode

echocardiograms were utilized to measure left ventricular end-systolic and end-diastolic

chamber dimensions and anterior/posterior wall thickness (Chicco et al., 2008; Le et al., 2014).

Once the measurements were taken for a rat, optics were turned on (1.3 mW, 5 ms pulses, 10

Hz) and a second exam was conducted to investigate acute stimulation-induced changes. Total

exam time for each trial was approximately 5 minutes with approximately 2 minutes of optic

stimulation.

2.15 Tissue collection

At the conclusion of experiments, rats were given an overdose of sodium pentobarbital

and perfused transcardially with 0.9% saline followed by 4.0% paraformaldehyde in 0.1 M PBS.

Brains were removed and post-fixed in 4.0% paraformaldehyde for 24 h at room temperature,

followed by storage in 30% sucrose in PBS at 4 °C. Coronal sections were made on a freezing

microtome at 30 μm thickness and then stored in cryoprotectant solution at −20 °C until

processing. A subset of rats received optogenetic stimulation prior to tissue collection. Rats

were tethered to a fiber optic patch cord and received 5 minutes of optic stimulation (1 mW, 5

ms pulses, 10 Hz) followed by 90 minutes of recovery for immediate-early gene (c-Fos)

expression prior to euthanasia, as described above.

2.16 Immunohistochemistry and microscopy

For fluorescent labeling of c-Fos, coronal brain sections were removed from

cryoprotectant and rinsed in PBS (5 x 5 min) at room temperature. Sections were then placed in

blocking solution (PBS, 0.1% bovine serum albumin, and 0.2% Triton X-100) for 1 hour. Next,

sections were incubated overnight in rabbit anti-c-Fos primary antibody (1:200 in blocking

solution, Cell Signaling Technologies, Ab #2250). Following overnight incubation in primary

antibody, sections were rinsed in PBS (5 x 5 min) and incubated in donkey anti-rabbit Cy5

secondary (1:1000 in PBS, Jackson ImmunoResearch, AB_2340607) for 1 hour. The tissue was

then washed (5 x 5 min in PBS), mounted with polyvinyl medium, and cover slipped for imaging.

For chromogen labeling of c-Fos positive nuclei, sections were first rinsed in PBS (5 x 5 min) at

room temperature. Sections were then incubated in 1% hydrogen peroxide in PBS for 10

minutes followed by a second PBS (5 x 5 min) wash. Sections were then placed in blocking

solution for 1 hour. Following blocking, sections were incubated overnight in rabbit anti-c-Fos

primary antibody (1:2000 in blocking solution, Abcam, ab190289). Sections were then rinsed in

PBS (5 x 5 min) followed by a 1 hour incubation in biotinylated goat anti-rabbit secondary (1:500

1 in PBS, Vector Laboratories, BA-1000). Sections were rinsed in PBS (5 x 5 min) followed by a 1

2 hour incubation in Vectastain ABC Solution (1:1,000; Vector Laboratories). Sections were rinsed

3 again prior to incubation in diaminobenzidine and hydrogen peroxide (0.02%

4 diaminobenzidine/0.09% hydrogen peroxide in KPBS) for 10 min. Following incubation, slices

5 were rinsed, slide mounted, dehydrated in graded ethanol, and cover slipped.

6   To determine injection placement, YFP was imaged with a Zeiss Axio Imager Z2

7 microscope using the 10x objective, while YFP and c-Fos dual fluorescence were acquired as

8 tiled 20x objective images. For quantification of c-Fos positive cells, slides were imaged in

9 brightfield with the 10x objective.

11 *2.17* Quantification of c-Fos

12   To quantify the number of c-Fos positive cells, 3-4 IL micrographs adjacent to visible

13 cannula tracts were imaged and positive cells within a half-width 10x frame were hand-counted

14 by a treatment-blind observer.

16 *2.18 Experimental design*

17   Timelines for experiments are outlined in figure 1. Experiment 1 was comprised of 2

18 cohorts of male rats (n = 13-14 each) to yield 10 YFP and 17 ChR2. For this experiment, real-

19 time place preference and social behavior were assessed prior to restraint stress with 20 Hz

20 used for all stimulation. A parallel group of injected male rats (n = 8) were used for slice

21 electrophysiology. Experiment 2 consisted of 3 cohorts of male rats (n = 7-8 each) for a total of

22 12 YFP and 10 ChR2. A subset of these rats was equipped with radiotelemetry transmitters

23 (YFP n = 7, ChR2 n = 8) and all underwent real-time place preference and social behavior prior

24 to novel environment with 10 Hz stimulation. Afterward, rats in experiment 2 underwent

25 echocardiography before and after CVS. Experiment 3 was designed as a single cohort of

26 female rats (n = 29) with 12 YFP and 17 ChR2. A subset of these rats was equipped with

radiotelemeters (YFP n = 7, ChR2 n = 8). Real-time place preference and social motivation

were followed by novel environment and restraint with 10 Hz stimulation for all measures.

Female rats also underwent echocardiography before and after CVS. A subset of rats in

experiments 2 and 3 (n = 6-7/group/sex) received optic stimulation prior to tissue collection to

verify neuronal activation.

2.19 Data analysis

Data are expressed as mean ± standard error of the mean. Data were analyzed using

Prism 8 (GraphPad, San Diego, CA), with statistical significance set at $p < 0.05$ for rejection of

null hypotheses. Stimulation induced c-Fos was analyzed with Welch's unpaired t-test

comparing virus groups. RTPP stimulation preference was assessed via repeated measure two-

way analysis of variance (ANOVA) with virus and day (repeated) as factors. In the case of

significant main or interaction effects, Sidak's multiple comparisons were run to determine group

differences. Social motivation and novelty preference were assessed with unpaired t-tests to

compare virus groups within social, object, novel rat, or familiar rat interactions. Total distance

traveled during RTPP and social interaction was assessed with unpaired t-tests comparing virus

groups. Stress responses over time (corticosterone, glucose, HR, MAP, SAP, DAP, and HRV

spectra) were analyzed using mixed-effects analysis with virus and time (repeated) as factors,

followed by Fisher's post-hoc if significant main or interaction effects were present. Baseline HR

and MAP, as well as mean activity during novel environment, were assessed with Mann-

Whitney U non-parametric test comparing virus groups. Fractional shortening and ventricular

structure were assessed with paired Wilcoxon signed-rank tests comparing pre- and post-CVS

or optic status within virus groups.

3. Results

3.1 Validation and design

AAV viral vectors were targeted to the IL for expression of membrane-targeted ChR2-

YFP or cytosolic YFP under the pyramidal neuron promoter, CaMKIIα. Fiber optic cannulas

implanted in the IL permitted selective stimulation of IL glutamatergic neurons (**Fig. S1**), as

previously reported (Fuchikami et al., 2015; Wood et al., 2018). In female rats, representative c-

Fos labeling demonstrated *in vivo* neural activation in response to light-stimulation (1 mW, 10

Hz, 5 ms, 5 min) (**Fig. 1A**). c-Fos quantification in both male and female rats (**Fig. S2**) further

validated stimulation efficacy. Whole-cell patch-clamp recordings in slice demonstrated light-

evoked depolarizing current in male IL neurons expressing ChR2 (**Fig. 1B**). Further, 10 Hz

stimulation led to high-fidelity action potential generation. 20-Hz light pulses induced action

potential firing, but also increased resting membrane potential (**Fig. 1C**). These results,

combined with our previous studies quantifying increased immediate-early gene expression

after 20 Hz stimulation (Wood et al., 2018), suggest that both 10 Hz and 20 Hz stimulation

activate IL CaMKIIα-positive neurons. Although, 10 Hz stimulation is more in line with the

reported 4-10 Hz intrinsic firing rate of IL pyramidal neurons (Homayoun and Moghaddam,

2007; Ji and Neugebauer, 2012). The experimental design (**Fig. 1D**) is detailed in the methods.

Male and female rats received optogenetic stimulation during behavioral and physiological

measures in separate experiments. At the conclusion of all experiments, the placement of viral

injections and fiber optics was determined. Only animals with cannula placement verified to be

within or immediately (within 0.5 mm) dorsal to the IL were included in analyses (**Fig. 1E**).

Figure 1: Approach validation and experimental design. (**A**) Injection of AAV-packaged
constructs led to expression of cytosolic YFP or membrane-targeted ChR2-YFP under the
CaMKIIα promoter. Blue light stimulation in ChR2 females (1 mW, 10 Hz, 5 ms pulses for 5 min)
led to robust expression of the immediate-early gene marker c-Fos. Bregma +2.8mm. White
arrows indicate representative c-Fos-positive nuclei. Dashed white lines indicate fiber tip. Scale
bar: 200 μm and 40 μm for combined. (**B**) Voltage-clamp recordings from male IL-containing
slices illustrated light-evoked (1.1 mW, 100 ms pulse) depolarizing current. (**C**) Current-clamp
recordings found stimulation-locked spiking with 10 and 20 Hz stimulation (1.1 mW, 5 ms pulse).
(**D**) Experimental timelines. RTPP: real-time place preference, SI: social interaction, RS:
restraint stress, NE: novel environment, CVS: chronic variable stress. (**E**) Optic fiber placements
(YFP: yellow, ChR2: blue) were mapped within or immediately dorsal to the IL (red outline).
Coronal sections adapted from Swanson Rat Brain Atlas (3rd edition).

3.2 Real-time place preference

RTPP was used to examine the affective valence of stimulating IL glutamatergic

neurons. Group mean heat maps illustrate time spent in the stimulation versus non-stimulation

chambers (**Fig. 2A**). Male rats showed a preference for the stimulation chamber (**Fig. 2B**) with

20 Hz (repeated-measures 2-way ANOVA: stimulation x ChR2 $F_{(1, 23)}$ = 15.04, $p < 0.001$) and 10

Hz stimulation (repeated-measures 2-way ANOVA: stimulation $F_{(1, 20)}$ = 6.03, $p < 0.05$, ChR2 $F_{(1,}$

$_{20)}$ = 9.04, $p < 0.01$, stimulation x ChR2 $F_{(1, 20)}$ = 6.43, $p < 0.05$). Specifically, Sidak's *post hoc*

indicated a preference for the 20 Hz and 10 Hz stimulation chambers compared to habituation

day in the ChR2 groups (n = 10-15, $p < 0.01$). ChR2 groups also preferred the stimulation

chamber relative to YFP animals on the stimulation day (n = 10-12, $p < 0.01$). In contrast,

female rats demonstrated no preference or aversion for the chamber paired with IL stimulation

(n = 11-13/group, repeated-measures 2-way ANOVA: stimulation $F_{(1, 22)}$ = 5.79, $p < 0.05,$

stimulation x ChR2 $F_{(1, 22)}$ = 0.005, $p = 0.94$). Additionally, IL stimulation did not affect general

locomotor activity in either sex (**Fig. 2C**). Together, these findings indicate that the activity of

male IL glutamatergic neurons has a positive affective valance as animals were motivated to

seek out the stimulation. However, this cell group does not appear to modify affective state in

females.

3.3 Social behavior

The three-chamber social interaction assay was used to determine the influence of IL

glutamatergic neurons on sociability. During the social motivation test (**Fig. 2D**), there was no

difference in social interactions with 20 Hz stimulation in males (n = 9-10/group, unpaired t-test:

ChR2 vs YFP $t_{(17)}$ = 0.77, $p = 0.45$). However, 10 Hz stimulation in males increased social

interactions (**Fig. 2E**; n = 10-12/group, unpaired t-test: ChR2 vs YFP $t_{(18)}$ = 3.00, $p < 0.01$). In

contrast, 10 Hz stimulation in females did not impact social interaction (n = 12-16/group,

unpaired t-tests: ChR2 vs YFP $t_{(24)}$ = 1.01, $p = 0.32$). Furthermore, neither male nor female IL

stimulation affected preference for novel vs. familiar interactors (**Table S1**). Overall, these

results indicate that IL glutamatergic activity increases male social motivation in a frequency-

dependent manner; however, this cell population does not alter female social motivation.

3.4 Endocrine reactivity

To determine the effect of stimulating IL pyramidal neurons on neuroendocrine

responses, blood glucose and plasma corticosterone were monitored during restraint stress. In

males, optic stimulation decreased corticosterone (n = 10-17/group, mixed-effects: time $F_{(4,81)}$ =

139.4, p < 0.0001, ChR2 $F_{(1,25)}$ = 8.93, $p < 0.01$) at the 30-minute timepoint (**Fig. 3A**; $p < 0.01$).

Additionally, corticosterone was decreased post-stimulation in the ChR2 group during stress

recovery (90 min; $p < 0.05$). Stimulation also decreased blood glucose (mixed-effects: time $F_{(4,90)}$

= 62.86, $p < 0.0001$, time x ChR2 $F_{(4,90)}$ = 3.43, $p < 0.05$) in male rats during restraint (**Fig. 3B**;

15 min, $p < 0.01$; 30 min, $p < 0.05$). In contrast, stimulation did not alter plasma corticosterone in

female rats (**Fig. 3C**; n = 10-17/group, mixed-effects: time $F_{(4,93)}$ = 62.94, $p < 0.0001$) with

effects of time limited to within treatment. Additionally, stimulation increased glucose responses

to stress in female rats (n = 10-17/group, mixed effects: time $F_{(4,90)}$ = 24.96, $p < 0.0001$)

specifically at the 15-min timepoint (**Fig. 3D**; $p < 0.05$). Collectively, these results suggest that IL

activity in males reduces HPA axis activation as well as glucose mobilization. In females, IL

activity does not appear to alter HPA axis response to stress. However, IL stimulation in females

increases glucose, possibly through sympathetic mobilization of epinephrine and/or glucagon.

Figure 2: IL pyramidal neuron activity was preferred and increased social motivation in males but not females. A) Heat maps illustrate mean animal position in RTPP arena on stimulation day (n = 10-15/group). (**B**) Male ChR2 rats preferred the chamber paired with 20 Hz or 10 Hz stimulation (blue shading) relative to habituation day and YFP controls. Females showed no preference or aversion for IL stimulation. ** p < 0.01, *** p < 0.001 vs. YFP within stimulation. ## p < 0.01 vs. habituation within ChR2. Inset: portion of rats in each estrous cycle phase. D: diestrus, P: proestrus, M: metestrus, E: estrous. (**C**) Total distance traveled during

1   stimulation found no treatment-based differences in locomotion. (**D**) Representative movement
2   traces during the social interaction test (n = 7-16/group). (**E**) Interaction, defined as nose-poke
3   onto social cage, was increased by 10 Hz stimulation in males but not 20 Hz in males or 10 Hz
4   in females. ** $p < 0.01$ vs. YFP. Inset: portion of rats in each estrous cycle phase.

Figure 3: IL activation during acute restraint stress attenuated endocrine responses in
males but increased female stress reactivity. Stimulation (blue shading) during restraint
lowered plasma corticosterone **(A)** and blood glucose **(B)** in ChR2 males. **(C)** Stimulation in
females did not alter corticosterone responses. Inset: portion of rats in each estrous cycle
phase. D: diestrus, P: proestrus, M: metestrus, E: estrous. **(D)** IL stimulation during restraint
increased glucose mobilization in ChR2 females. n = 10-17/group, * $p < 0.05$, ** $p < 0.01$ vs.
YFP.

3.5 Cardiovascular reactivity

Baseline hemodynamic measures were recorded in the homecage prior to handling (**Fig.**

S4). Animals were then connected to fiber optic patch cords and exposed to a brightly lit novel

environment as a psychogenic stimulus to examine IL glutamatergic effects on cardiovascular

reactivity. Over the course of the stressor, there were no differences in activity between male

YFP (n = 7) and ChR2 (n = 8) rats (**Fig. 4A**; Mann-Whitney: U = 407, p = 0.53). In terms of

hemodynamic responses, stimulation decreased HR reactivity in male ChR2 rats (mixed-effects:

time $F_{(30,280)}$ = 19.61, $p < 0.0001$, time x ChR2 $F_{(30,280)}$ = 2.06, $p < 0.01$), an interaction present

at the 8-minute timepoint (**Fig. 4B**; $p < 0.05$). After the first minute of stimulation, arterial

pressures including mean (mixed-effects: time $F_{(30,172)}$ = 25.99, $p < 0.0001$, time x ChR2

$F(30,172)$ = 1.86, $p < 0.01$), systolic (mixed-effects: time $F_{(30,213)}$ = 27.61, $p < 0.0001$), and

diastolic (mixed-effects: time $F_{(30,189)}$ = 20.56, $p < 0.0001$, ChR2 $F_{(1,12)}$ = 8.05, $p < 0.05$, time x

ChR2 $F_{(30,189)}$ = 2.38, $p < 0.01$) were decreased in male ChR2 rats. Post-hoc analysis indicated

these effects were present across numerous timepoints (**Fig. 4C-E**; 7-28 minutes; $p < 0.05$). In

contrast, stimulation increased MAP and DAP in minute 1 ($p < 0.05$). For females, there was no

effect of ChR2 on activity in the novel environment (**Fig. 4F**; n = 7-8, Mann-Whitney: ChR2 vs

YFP U = 352, p = 0.15). However, IL activation increased female HR responses (mixed effects:

time $F_{(30,390)}$ = 25.83, $p < 0.0001$) early in the stressor (**Fig. 4G**; 1-9 min; $p < 0.05$) as well as at

25 min ($p < 0.05$). Mixed effects analysis of arterial pressure found main effects of time for MAP

(**Fig. 4H**; time $F_{(30,390)}$ = 31.96, $p < 0.0001$) and SAP (**Fig. 4I**; time $F_{(30,390)}$ = 35.01, $p < 0.0001$)

without virus-specific effects. Taken together, these results indicate activation of male IL

glutamate neurons decreases HR and arterial pressure responses to psychological stress. In

contrast, activation in females increases HR.

Figure 4: Activation of glutamatergic IL neurons during novel environment exposure reduced male cardiovascular stress responses but increased female HR reactivity. (A) Male IL activation did not affect mean activity in the novel environment. Despite an increase in MAP and DAP during the first minute of stimulation (blue shading) and stress, all recorded hemodynamic responses were decreased in ChR2 males **(B-E)**. **(F)** Stimulation did not alter activity in females. **(G)** Female IL stimulation elevated HR reactivity compared to YFP controls but did not affect arterial pressures **(H-J)**. Inset: portion of rats in each estrous cycle phase. D: diestrus, P: proestrus, M: metestrus, E: estrous. n = 7-8/group, * p < 0.05 vs. YFP.

3.6 Effects of chronic stress on cardiac function

Resting homecage LF/HF of HRV was recorded via radiotelemetry to assess how prior

optic stimulation affected circadian autonomic balance and net cardiac sympathetic drive during

CVS. In the absence of ongoing stimulation, ChR2 (n = 8) males had lower LF/HF than YFP (n

= 7) rats (**Fig. 5A**; mixed-effects: time $F_{(5,62)}$ = 4.45, $p < 0.01$, ChR2 $F_{(1,13)}$ = 5.33, $p < 0.05$). In

the AM the day prior to CVS, ChR2 males had lower baseline LF/HF ($p < 0.05$) leading to

decreased sympathetic drive the morning of CVS day 14 ($p < 0.05$). In females, there were

circadian effects on LF/HF (**Fig. 5B**; n = 5-6/group, mixed-effects: time $F_{(5,45)}$ = 5.98, p < 0.01)

but no differences due to virus (mixed-effects: ChR2 $F_{(1,9)}$ = 1.28, p > 0.05).

Left ventricle echocardiography was performed before and after CVS to examine the *in*

vivo cardiac consequences of chronic stress and the effects of IL stimulation on cardiac

contractility. Fractional shortening (FS) was measured as the portion of diastolic dimension lost

in systole. In male rats treated with YFP (n = 8), CVS increased FS (**Fig. 5C**; Wilcoxon: Post vs

Pre W = 30.0, $p < 0.05$). In contrast, there was no effect of CVS on FS in ChR2 (n = 8) males

(Wilcoxon: Post vs Pre W = -4.0, $p = 0.84$). To determine whether post-CVS FS was affected by

acute stimulation, measurements were taken with optics off and then on (**Fig. 5D**). Here, a

cohort of males showed no effect of acute optic status on FS (n = 3-4/group, Wilcoxon: YFP On

vs Off W = 4.0, $p = 0.13$; ChR2 On vs Off W = 2.0, $p = 0.75$). In females there was no effect of

CVS on FS (**Fig. 5E**; n = 10, YFP Post vs Pre W = -21.0, $p = 0.25$; n = 17, ChR2 Post vs Pre W

= 53.0, $p = 0.07$). However, optic stimulation increased post-CVS FS in ChR2 animals (**Fig. 5F**;

Wilcoxon: On vs Off W = 87.0, $p < 0.01$). In aggregate, these findings indicate that a history of IL

stimulation limits net cardiac sympathetic drive during chronic stress in males, an effect that

may protect against CVS-increased FS. Conversely, neither CVS nor prior IL stimulation

affected autonomic balance or cardiac function in females. However, acute IL stimulation after

CVS increased female cardiac contractility.

Figure 5: The effects of chronic stress and IL activity on cardiac function. (A) The ratio of
LF to HF components of HRV were reduced in ChR2 males at baseline and the end of CVS. **(B)**
Females had no treatment-based effects on net sympathetic drive. L: light phase, D: dark
phase, LF: low frequency, HF: high frequency, HRV: heart rate variability. **(C)** CVS increased
FS in YFP but not ChR2 males. **(D)** Acute optic status (On vs. Off) did not affect FS in males.
(E) Females showed no changes in FS after CVS. **(F)** However, acute optogenetic activation
(blue shading) increased female FS after CVS. * p < 0.05 vs YFP or Pre CVS YFP, ** p < 0.01
vs. ChR2 Off.

3.7 Chronic stress effects on cardiac structure

Left ventricular morphological analysis was carried out to examine potential structural

contributions to altered function. In YFP males (**Fig. 6A**; n = 8), CVS increased wall thickness of

the posterior wall in systole (**Fig 6B**; Wilcoxon: Post vs Pre W = 28, $p < 0.05$) and the anterior

wall in both systole (Wilcoxon: Post vs Pre W = 36, $p < 0.01$) and diastole (Wilcoxon: Post vs

Pre W = 28, $p < 0.05$). Furthermore, increased wall thickness was sufficient to decrease

ventricle size in systole (**Fig. 6C**; Wilcoxon: Post vs Pre W = -30, $p < 0.05$). Critically, male rats

that had previously received IL stimulation (ChR2, n = 8) were protected from the effects of CVS

on cardiac hypertrophy (**Fig 6D-E**; Wilcoxon: Post vs Pre W = -9 to 20, $p > 0.05$). In contrast to

males, female rats (**Fig. 6F**) showed minimal effects of CVS on cardiac remodeling as YFP rats

(n = 10) had decreased posterior wall thickness in diastole (**Fig. 6G**; Wilcoxon: Post vs Pre W =

-37.0, $p < 0.05$) with no other structural changes (**Fig. 6H**; $p > 0.05$). Additionally, there were no

significant changes in cardiac structure in the female ChR2 group (**Fig. 6I-J**; n = 16, Wilcoxon:

Post vs Pre W = -45 to 25, $p > 0.05$). Overall, CVS-induced inward hypertrophic remodeling was

prevented by prior IL activation in males. In contrast, female rats were generally resistant to the

hypertrophic effects of chronic stress.

Figure 6: Effects of chronic stress and prior IL stimulation on cardiac structure.
(A) Representative echocardiographic images of the left ventricle in bisected m-mode view
before and after CVS in YFP males. V: ventricle, A: anterior wall, P: posterior wall. **(B)** CVS
increased wall thickness in diastole and systole in YFP males, **(C)** reducing ventricle size. **(D, E)**
ChR2 males that received prior stimulation were protected from the cardiac consequences of
CVS. **(F)** Representative m-mode view of the left ventricle in YFP females before and after CVS.
(G, H) Females did not exhibit ventricular hypertrophy after CVS as posterior wall thickness in
diastole was decreased. **(I, J)** No myocardial structural changes were evident in ChR2 females.
n = 8-16/group, * p < 0.05, ** p < 0.01 vs. Pre CVS YFP

4. Discussion

In the current study, optogenetic stimulation of glutamatergic IL pyramidal neurons was

combined with behavioral, endocrine, and cardiovascular assessments. Our results show that,

in males, IL pyramidal neuron activity was preferred, increased social motivation, and reduced

acute physiological stress reactivity. Intriguingly, prior IL activation lowered net cardiac

sympathetic drive and protected against subsequent myocardial remodeling after chronic stress.

However, IL activity had fundamentally different regulatory effects in females. Stimulation did

not have motivational valence or alter social behavior but increased acute physiological stress

reactivity and cardiac contractility following chronic stress. Collectively, these findings identify

sexual divergence in the cortical integration of affective and physiological systems, suggesting

that vmPFC output signaling may differentially impact health outcomes in males and females.

The comorbidity of CVD and MDD shows sexual divergence with females at twice the

risk (Goldstein et al., 2019; Möller-Leimkühler, 2007; Naqvi et al., 2005). Given the interactions

between stress, mood disorders, and CVD, stress-reactive neural populations are well

positioned to regulate affective and cardiovascular outcomes. Importantly, chronic stress

exposure in male rats reduces IL pyramidal neuron dendritic arborization and increases local

GABAergic signaling, suggesting reduced glutamatergic output (McKlveen et al., 2019, 2016;

Radley et al., 2008). Further, long-term reduction of male IL glutamatergic output increases HPA

axis activity and impairs vascular function (Myers et al., 2017; Schaeuble et al., 2019).

Collectively, IL output neurons represent a target for modulating behavioral and physiological

responses to stress. Thus, we sought to test this hypothesis through real-time *in vivo*

assessments of behavior, physiology, and organ function. Altogether, we found that the

neurobiology of emotional behaviors, endocrine-autonomic integration, and cardiovascular

outcomes differed substantially between sexes.

The RTPP test was used to assess the affective state of experimental animals by

measuring preference or aversion for IL neural activity. Here, male rats preferred the chamber

paired with either 10 Hz or 20 Hz stimulation, but the valence of female IL activation was neither

positive nor negative. Previous work indicates that male mPFC stimulation reduces social

avoidance after social defeat (Covington et al., 2010), yet relatively few studies have examined

IL function in females. Although social interaction induces less IL immediate early-gene

expression in females than males (Mikosz et al., 2015; Stack et al., 2010), the current study is,

to our knowledge, the first examination of IL regulation of female social behavior. IL activity did

not alter social behavior in females, indicating a significant sexual divergence in the neural

regulation of sociability. However, males exhibited frequency-dependent increases in social

motivation. These results suggest that stimulation near the intrinsic pyramidal neuron firing rate

(4-10 Hz) (Homayoun and Moghaddam, 2007; Ji and Neugebauer, 2012) is required to increase

male social motivation.

Activation of IL glutamatergic neurons also reduced male glucose and corticosterone

responses to stress. Conversely, female IL glutamatergic stimulation increased glucose

mobilization without affecting corticosterone, suggesting a role in sympatho-excitation. Female

IL activity also increased tachycardic responses to stress, while male IL stimulation reduced

cardiovascular stress responses including HR and arterial pressures. These experiments

indicate that IL activity has opposing effects on endocrine-autonomic integration in males and

females whereby male glutamatergic IL neurons cause widespread inhibition of the stress

response and female IL neurons facilitate sympathetic responses.

Chronic stress exposure induced ventricular hypertrophy and increased endocardial FS

in males, suggesting that increased wall thickness likely accounted for the increase in FS. This

effect was prevented by a history of IL stimulation in males, likely arising from a reduction in

symaptho-vagal imbalance throughout CVS. In fact, previous IL stimulation limited resting net

sympathetic drive, a major risk for CVD (Thayer et al., 2012; Wulsin et al., 2015). Given that rats

received no optic stimulation during CVS and that acute optic status did not impact male FS,

this effect was likely driven by stimulation-induced IL plasticity. IL stimulation has been shown to

induce persistent morphological changes in males including increased excitatory synapses onto

pyramidal neurons, suggesting a prolonged state of enhanced excitability (Fuchikami et al.,

2015; Moda-Sava et al., 2019). Thus, lower resting net sympathetic tone and/or reduced

sympathetic activity during stressors may have prevented the consequences of chronic stress to

induce inward hypertrophic remodeling of the myocardium. These effects were not present in

females as hypertrophic remodeling was not evident and acute stimulation increased FS.

Collectively, these results indicate that prior male IL glutamatergic activity is sufficient to restrain

responses to chronic stress while female IL activity increases cardiac contractility independent

of structural changes.

Physiological responses to stress are critical for survival, tightly defended, and limited in

maximum capacity; accordingly, our manipulations only caused moderate changes in response

magnitude to acute challenges. However, our findings suggest that the interaction of altered

stress responding with chronic stressors is sufficient to account for differences in cardiac

structure and function. It is also worth noting that prior studies on the role of vmPFC in stress

responding have yielded equivocal results. Specifically, lesions studies in male rats have

observed both increased (Diorio et al., 1993; Figueiredo et al., 2003) and decreased (Radley et

al., 2006) neuroendocrine stress responding. Similarly, pharmacological studies found that male

vmPFC both reduces (Müller-Ribeiro et al., 2012) and enhances (Tavares et al., 2009)

cardiovascular stress responses. These discrepancies are not limited to rodents as

pharmacological studies in non-human primates suggest that vmPFC increases cardiovascular

stress responding (Alexander et al., 2020), while electrical stimulation in humans causes

hypotension (Lacuey et al., 2018). Although it is possible that variations in acute stressors,

selectivity of cortical subregion targeting, and/or species may account for these divergent

findings, all of the described studies used non-specific blockade or stimulation. Given the

complexity of cortical circuitry and the numerous local inhibitory mechanisms (McKlveen et al.,

2019), cell-type specific targeting has the ability to isolate the precise contribution of vmPFC

principal cells independent of afferents, interneurons, and glia. To address this issue, our prior

studies developed anatomical and cell-type specific approaches to reduce glutamate outflow

from IL pyramidal neurons. Genetic knockdown of IL glutamate release in males increased

neuroendocrine and cardiovascular responses to both acute and chronic stress (Myers et al.,

2017; Schaeuble et al., 2019), demonstrating the necessity of IL output for inhibiting

physiological stress responses. The current cell-type specific optogenetic targeting adds further

temporal specificity and identifies the sufficiency of male IL glutamate neurons for stress

inhibition. Albeit with opposing effects in females.

The IL does not directly innervate the neurons that govern endocrine and autonomic

stress responses. Accordingly, downstream glutamate signaling from IL synaptic terminals

requires intermediary synapses. The exact circuits engaged by IL pyramidal cells to bring about

the observed effects remain to be determined. However, anterograde mapping studies indicate

that IL projections widely innervate the forebrain and brainstem (Vertes, 2004; Wood et al.,

2018). We previously found that, in males, stress-activated IL neurons innervate local inhibitory

GABAergic neurons in the posterior hypothalamus (PH) (Myers et al., 2016). Furthermore,

blocking GABAergic tone in the PH reduces social behavior and increases HPA axis reactivity

suggesting inhibition of the PH may be important for limiting behavioral and physiological stress

responses (Myers et al., 2016). Additionally, IL inputs to the amygdala are critical for fear

extinction and reducing anxiety-like behavior (Adhikari et al., 2015; Sierra-Mercado et al., 2011).

Interestingly, amygdala-projecting IL neurons are both resistant to stress-induced dendritic

retraction as well as sensitive to estrogen (Shansky et al., 2010, 2009). Thus, the IL-amygdala

circuit could play a key role in sex differences in behavioral regulation. Although the

downstream mechanisms of IL cardiovascular regulation are unknown, male IL projections

target pre-autonomic cell groups in the brainstem and give rise to multi-synaptic pathways that

innervate the adrenal medulla (Dum et al., 2019; Gabbott et al., 2005). Further sex-specific

analysis of IL synaptic signaling in forebrain and brainstem nuclei is necessary to determine the

basis of divergent behavioral and physiological integration.

While ovarian hormones have far-reaching effects on behavior and physiological

systems, we did not control for estrous cycle phase. Instead, we used randomly cycling females.

Cycle phase was reported for each treatment, but statistical power was insufficient to examine

phase as a factor. It remains to be determined how gonadal hormones might contribute to the

sexually divergent effects observed. Estrogen receptor (ER) α, β, and the g-protein coupled ER

are expressed in pyramidal and non-pyramidal PFC neurons in both sexes (Almey et al., 2014;

Montague et al., 2008). Further, ER localized to PFC dendritic spines regulates synaptic

morphology and ionotropic glutamate receptor ubiquitination/degradation following repeated

stress (Hao et al., 2006; Wei et al., 2014; Yuen et al., 2016). These protective effects are

dependent on PFC ERα and estradiol-synthesizing aromatase (Wei et al., 2014), indicating a

role for extra-ovarian estrogen synthesis. In addition, ER expressed in axons and axonal

terminals rapidly alters pre-synaptic transmission in pyramidal cells (Almey et al., 2014).

Although studies have described multiple interactions between ER and cortical glutamate

signaling (Galvin and Ninan, 2014; Hara et al., 2018), much less is known about PFC

progesterone signaling. Progesterone receptors are expressed in frontal cortex (Guerra-Araiza

et al., 2003) but progesterone derivatives also signal through GABA$_A$ receptors and regulate

GABA subunit expression (Andrade et al., 2012). Moreover, sex differences in IL regulation of

behavior and physiology could arise from the actions of androgens. Androgen receptors (AR)

are expressed in PFC neurons and glia (Finley and Kritzer, 1999). AR is also enriched in VTA-

projecting PFC neurons that influence extracellular dopamine in PFC through downstream VTA

glutamate signaling (Aubele and Kritzer, 2012). It is particularly interesting that androgens and

estrogens act in opposition to modify PFC dopamine, norepinephrine, and serotonin metabolism

during novel environment stress (Handa et al., 1997). Thus, gonadal steroids and locally

synthesized modulators affect PFC cellular processes and projection activity, ultimately

engaging multiple neurotransmitter systems. Accordingly, the specific cellular and synaptic

processes contributing to sex-dependent stress reactivity and resilience are promising avenues

for identifying therapeutic targets.

The current study identified a cortical node integrating mood-related behaviors with

cardiovascular outcomes. Moreover, activity in this vmPFC cell population produced sex-

specific effects on behavior and physiology. In addition to highlighting the necessity of sex-

based investigation, these data point to a neurochemical basis for sex differences in stress-

related health determinants. Ultimately, further investigation of brain-body interactions in the

face of prolonged stress may provide a better understating of disease risk and resilience factors.

Funding

This work was supported by NIH grants R00 HL122454 and R01 HL150559 to Brent

Myers. Data generated by the University of Cincinnati Mouse Metabolic Phenotyping Core were

supported by NIH grant U2C DK059630.

Author contributions

Tyler Wallace: Formal analysis, Investigation, Validation, Visualization, Writing - original

draft, Writing - review & editing. **Derek Schaeuble:** Formal analysis, Investigation, Validation,

Visualization, Writing - original draft, Writing - review & editing. **Sebastian A. Pace:** Formal

analysis, Investigation, Validation. **Morgan K. Schackmuth:** Formal analysis, Validation.

Shane T. Hentges: Methodology, Resources, Supervision, Writing - review & editing. **Adam J.**

Chicco: Formal analysis, Investigation, Methodology, Resources, Writing - review & editing.

Brent Myers: Conceptualization, Funding acquisition, Investigation, Methodology, Resources,

Supervision, Writing - original draft, Writing - review & editing.

Disclosures

The authors have no conflicts of interest to disclose.

Acknowledgments

References

Adhikari, A., Lerner, T.N., Finkelstein, J., Pak, S., Jennings, J.H., Davidson, T.J., Ferenczi, E., Gunaydin, L.A., Mirzabekov, J.J., Ye, L., Kim, S.Y., Lei, A., Deisseroth, K., 2015. Basomedial amygdala mediates top-down control of anxiety and fear. Nature 527, 179–185. https://doi.org/10.1038/nature15698

Alexander, L., Gaskin, P.L.R., Sawiak, S.J., Fryer, T.D., Hong, Y.T., Cockcroft, G.J., Clarke, H.F., Roberts, A.C., 2019. Fractionating Blunted Reward Processing Characteristic of Anhedonia by Over-Activating Primate Subgenual Anterior Cingulate Cortex. Neuron 101, 307-320.e6. https://doi.org/10.1016/j.neuron.2018.11.021

Alexander, L., Wood, C.M., Gaskin, P.L.R., Sawiak, S.J., Fryer, T.D., Hong, Y.T., McIver, L., Clarke, H.F., Roberts, A.C., 2020. Over-activation of primate subgenual cingulate cortex enhances the cardiovascular, behavioral and neural responses to threat. Nat. Commun. 11, 1–14. https://doi.org/10.1038/s41467-020-19167-0

Almey, A., Cannell, E., Bertram, K., Filardo, E., Milner, T.A., Brake, W.G., 2014. Medial prefrontal cortical estradiol rapidly alters memory system bias in female rats: Ultrastructural analysis reveals membrane-associated estrogen receptors as potential mediators. Endocrinology 155, 4422–4432. https://doi.org/10.1210/en.2014-1463

Andrade, S., Arbo, B.D., Batista, B.A.M., Neves, A.M., Branchini, G., Brum, I.S., Barros, H.M.T., Gomez, R., Ribeiro, M.F.M., 2012. Effect of progesterone on the expression of GABA $_A$ receptor subunits in the prefrontal cortex of rats: implications of sex differences and brain hemisphere. Cell Biochem. Funct. 30, 696–700. https://doi.org/10.1002/cbf.2854

Aubele, T., Kritzer, M.F., 2012. Androgen influence on prefrontal dopamine systems in adult male rats: Localization of cognate intracellular receptors in medial prefrontal projections to the ventral tegmental area and effects of gonadectomy and hormone replacement on glutamate-stimulated extracellular dopamine level. Cereb. Cortex 22, 1799–1812. https://doi.org/10.1093/cercor/bhr258

Beckmann, M., Johansen-Berg, H., Rushworth, M.F.S., 2009. Connectivity-based parcellation of human cingulate cortex and its relation to functional specialization. J. Neurosci. 29, 1175–1190. https://doi.org/10.1523/JNEUROSCI.3328-08.2009

Beissner, F., Meissner, K., Bär, K.J., Napadow, V., 2013. The autonomic brain: An activation likelihood estimation meta-analysis for central processing of autonomic function. J. Neurosci. 33, 10503–10511. https://doi.org/10.1523/JNEUROSCI.1103-13.2013

Bimpisidis, Z., König, N., Wallén-Mackenzie, Å., 2020. Two different real-time place preference paradigms using optogenetics within the ventral tegmental area of the mouse. J. Vis. Exp. 2020, 60867. https://doi.org/10.3791/60867

Binder, E.B., Nemeroff, C.B., 2010. The CRF system, stress, depression and anxietyinsights from human genetic studies. Mol. Psychiatry. https://doi.org/10.1038/mp.2009.141

Chicco, A.J., McCune, S.A., Emter, C.A., Sparagna, G.C., Rees, M.L., Bolden, D.A., Marshall, K.D., Murphy, R.C., Moore, R.L., 2008. Low-intensity exercise training delays heart failure and improves survival in female hypertensive heart failure rats. Hypertension 51, 1096–1102. https://doi.org/10.1161/HYPERTENSIONAHA.107.107078

Cora, M.C., Kooistra, L., Travlos, G., 2015. Vaginal Cytology of the Laboratory Rat and Mouse:Review and Criteria for the Staging of the Estrous Cycle Using Stained Vaginal Smears. Toxicol. Pathol. 43, 776–793. https://doi.org/10.1177/0192623315570339

Covington, H.E., Lobo, M.K., Maze, I., Vialou, V., Hyman, J.M., Zaman, S., LaPlant, Q., Mouzon, E., Ghose, S., Tamminga, C.A., Neve, R.L., Deisseroth, K., Nestler, E.J., 2010. Antidepressant effect of optogenetic stimulation of the medial prefrontal cortex. J. Neurosci. 30, 16082–16090. https://doi.org/10.1523/JNEUROSCI.1731-10.2010

Diorio, D., Viau, V., Meaney, M.J., 1993. The Role of the Medial Prefrontal Cortex (Cingulate Gyrus) in the Regulation of Hypothalamic-Pituitary-Adrenal Responses to Stress, The

Journal of Neuroscience.

Drevets, W.C., Price, J.L., Simpson, J.R., Todd, R.D., Reich, T., Vannier, M., Raichle, M.E., 1997. Subgenual prefrontal cortex abnormalities in mood disorders. Nature 386, 824–827. https://doi.org/10.1038/386824a0

Drevets, W.C., Savitz, J., Trimble, M., 2008. The subgenual anterior cingulate cortex in mood disorders. CNS Spectr. 13, 663–681. https://doi.org/10.1017/S1092852900013754

Dum, R.P., Levinthal, D.J., Strick, P.L., 2019. The mind-body problem: Circuits that link the cerebral cortex to the adrenal medulla. Proc. Natl. Acad. Sci. U. S. A. 116, 26321–26328. https://doi.org/10.1073/pnas.1902297116

Electrophysiology, T.F. of the E.S., 1996. Heart Rate Variability. Circulation 93, 1043–1065. https://doi.org/10.1161/01.CIR.93.5.1043

Felix-Ortiz, A.C., Tye, K.M., 2014. Amygdala inputs to the ventral hippocampus bidirectionally modulate social behavior. J. Neurosci. 34, 586–595. https://doi.org/10.1523/JNEUROSCI.4257-13.2014

Ferenczi, E.A., Zalocusky, K.A., Liston, C., Grosenick, L., Warden, M.R., Amatya, D., Katovich, K., Mehta, H., Patenaude, B., Ramakrishnan, C., Kalanithi, P., Etkin, A., Knutson, B., Glover, G.H., Deisseroth, K., 2016. Prefrontal cortical regulation of brainwide circuit dynamics and reward-related behavior. Science (80-.). 351. https://doi.org/10.1126/science.aac9698

Figueiredo, H.F., Bruestle, A., Bodie, B., Dolgas, C.M., Herman, J.P., 2003. The medial prefrontal cortex differentially regulates stress-induced c-fos expression in the forebrain depending on type of stressor. Eur. J. Neurosci. 18, 2357–2364. https://doi.org/10.1046/j.1460-9568.2003.02932.x

Finley, S.K., Kritzer, M.F., 1999. Immunoreactivity for intracellular androgen receptors in identified subpopulations of neurons, astrocytes and oligodendrocytes in primate prefrontal cortex. J. Neurobiol. 40, 446–457. https://doi.org/10.1002/(SICI)1097-4695(19990915)40:4<446::AID-NEU3>3.0.CO;2-J

Flak, J.N., Jankord, R., Solomon, M.B., Krause, E.G., Herman, J.P., 2011. Opposing effects of chronic stress and weight restriction on cardiovascular, neuroendocrine and metabolic function. Physiol. Behav. 104, 228–234. https://doi.org/10.1016/j.physbeh.2011.03.002

Fuchikami, M., Thomas, A., Liu, R., Wohleb, E.S., Land, B.B., DiLeone, R.J., Aghajanian, G.K., Duman, R.S., 2015. Optogenetic stimulation of infralimbic PFC reproduces ketamine's rapid and sustained antidepressant actions. Proc. Natl. Acad. Sci. U. S. A. 112, 8106–8111. https://doi.org/10.1073/pnas.1414728112

Gabbott, P.L.A., Warner, T.A., Jays, P.R.L., Salway, P., Busby, S.J., 2005. Prefrontal cortex in the rat: Projections to subcortical autonomic, motor, and limbic centers. J. Comp. Neurol. 492, 145–177. https://doi.org/10.1002/cne.20738

Galvin, C., Ninan, I., 2014. Regulation of the mouse medial prefrontal cortical synapses by endogenous estradiol. Neuropsychopharmacology 39, 2086–2094. https://doi.org/10.1038/npp.2014.56

Gianaros, P.J., Sheu, L.K., 2009. A review of neuroimaging studies of stressor-evoked blood pressure reactivity: Emerging evidence for a brain-body pathway to coronary heart disease risk. Neuroimage. https://doi.org/10.1016/j.neuroimage.2009.04.073

Gianaros, P.J., Wager, T.D., 2015. Brain-Body Pathways Linking Psychological Stress and Physical Health. Curr. Dir. Psychol. Sci. 24, 313–321. https://doi.org/10.1177/0963721415581476

Goldstein, J.M., Hale, T., Foster, S.L., Tobet, S.A., Handa, R.J., 2019. Sex differences in major depression and comorbidity of cardiometabolic disorders: impact of prenatal stress and immune exposures. Neuropsychopharmacology. https://doi.org/10.1038/s41386-018-0146-1

Goodson, M.L., Packard, A.E.B., Buesing, D.R., Maney, M., Myers, B., Fang, Y., Basford, J.E.,

Hui, D.Y., Ulrich-Lai, Y.M., Herman, J.P., Ryan, K.K., 2017. Chronic stress and Rosiglitazone increase indices of vascular stiffness in male rats. Physiol. Behav. 172, 16–23. https://doi.org/10.1016/j.physbeh.2016.03.031

Grippo, A.J., Johnson, A.K., 2009. Stress, depression and cardiovascular dysregulation: A review of neurobiological mechanisms and the integration of research from preclinical disease models. Stress. https://doi.org/10.1080/10253890802046281

Guerra-Araiza, C., Villamar-Cruz, O., González-Arenas, A., Chavira, R., Camacho-Arroyo, I., 2003. Changes in progestrone receptor isoforms content in the rat brain during the oestrous cycle and after oestradiol and progesterone treatments. J. Neuroendocrinol. 15, 984–990. https://doi.org/10.1046/j.1365-2826.2003.01088.x

Hamani, C., Mayberg, H., Stone, S., Laxton, A., Haber, S., Lozano, A.M., 2011. The Subcallosal Cingulate Gyrus in the Context of Major Depression. BPS 69, 301–308. https://doi.org/10.1016/j.biopsych.2010.09.034

Handa, R.J., Hejna, G.M., Lorens, S.A., 1997. Androgen inhibits neurotransmitter turnover in the medial prefrontal cortex of the rat following exposure to a novel environment. Brain Res. 751, 131–138. https://doi.org/10.1016/S0006-8993(96)01394-7

Hao, J., Rapp, P.R., Leffler, A.E., Leffler, S.R., Janssen, W.G.M., Lou, W., McKay, H., Roberts, J.A., Wearne, S.L., Hof, P.R., Morrison, J.H., 2006. Estrogen alters spine number and morphology in prefrontal cortex of aged female rhesus monkeys. J. Neurosci. 26, 2571–2578. https://doi.org/10.1523/JNEUROSCI.3440-05.2006

Hara, Y., Crimins, J.L., Puri, R., Wang, A.C.J., Motley, S.E., Yuk, F., Ramos, T.M., Janssen, W.G.M., Rapp, P.R., Morrison, J.H., 2018. Estrogen Alters the Synaptic Distribution of Phospho-GluN2B in the Dorsolateral Prefrontal Cortex While Promoting Working Memory in Aged Rhesus Monkeys. Neuroscience 394, 303–315. https://doi.org/10.1016/j.neuroscience.2018.09.021

Homayoun, H., Moghaddam, B., 2007. NMDA receptor hypofunction produces opposite effects on prefrontal cortex interneurons and pyramidal neurons. J. Neurosci. 27, 11496–11500. https://doi.org/10.1523/JNEUROSCI.2213-07.2007

Ji, G., Neugebauer, V., 2012. Modulation of medial prefrontal cortical activity using in vivo recordings and optogenetics. Mol. Brain 5, 36. https://doi.org/10.1186/1756-6606-5-36

Lacuey, N., Hampson, J.P., Theeranaew, W., Zonjy, B., Vithala, A., Hupp, N.J., Loparo, K.A., Miller, J.P., Lhatoo, S.D., 2018. Cortical structures associated with human blood pressure control. JAMA Neurol. 75, 194–202. https://doi.org/10.1001/jamaneurol.2017.3344

Le, C.H., Mulligan, C.M., Routh, M.A., Bouma, G.J., Frye, M.A., Jeckel, K.M., Sparagna, G.C., Lynch, J.M., Moore, R.L., McCune, S.A., Bristow, M., Zarini, S., Murphy, R.C., Chicco, A.J., 2014. Delta-6-desaturase links polyunsaturated fatty acid metabolism with phospholipid remodeling and disease progression in heart failure. Circ. Hear. Fail. 7, 172–183. https://doi.org/10.1161/CIRCHEARTFAILURE.113.000744

Liotti, M., Mayberg, H.S., Brannan, S.K., McGinnis, S., Jerabek, P., Fox, P.T., 2000. Differential limbic-cortical correlates of sadness and anxiety in healthy subjects: Implications for affective disorders. Biol. Psychiatry 48, 30–42. https://doi.org/10.1016/S0006-3223(00)00874-X

Mayberg, H.S., Lozano, A.M., Voon, V., McNeely, H.E., Seminowicz, D., Hamani, C., Schwalb, J.M., Kennedy, S.H., 2005. Deep brain stimulation for treatment-resistant depression. Neuron 45, 651–660. https://doi.org/10.1016/j.neuron.2005.02.014

McKlveen, J.M., Moloney, R.D., Scheimann, J.R., Myers, B., Herman, J.P., 2019. "Braking" the Prefrontal Cortex: The Role of Glucocorticoids and Interneurons in Stress Adaptation and Pathology. Biol. Psychiatry. https://doi.org/10.1016/j.biopsych.2019.04.032

McKlveen, J.M., Morano, R.L., Fitzgerald, M., Zoubovsky, S., Cassella, S.N., Scheimann, J.R., Ghosal, S., Mahbod, P., Packard, B.A., Myers, B., Baccei, M.L., Herman, J.P., 2016. Chronic Stress Increases Prefrontal Inhibition: A Mechanism for Stress-Induced Prefrontal

Dysfunction. Biol. Psychiatry 80, 754–764. https://doi.org/10.1016/j.biopsych.2016.03.2101

McKlveen, J.M., Myers, B., Herman, J.P., 2015. The Medial Prefrontal Cortex: Coordinator of Autonomic, Neuroendocrine and Behavioural Responses to Stress. J. Neuroendocrinol. https://doi.org/10.1111/jne.12272

Mikosz, M., Nowak, A., Werka, T., Knapska, E., 2015. Sex differences in social modulation of learning in rats. Sci. Rep. 5. https://doi.org/10.1038/srep18114

Moda-Sava, R.N., Murdock, M.H., Parekh, P.K., Fetcho, R.N., Huang, B.S., Huynh, T.N., Witztum, J., Shaver, D.C., Rosenthal, D.L., Alway, E.J., Lopez, K., Meng, Y., Nellissen, L., Grosenick, L., Milner, T.A., Deisseroth, K., Bito, H., Kasai, H., Liston, C., 2019. Sustained rescue of prefrontal circuit dysfunction by antidepressant-induced spine formation. Science (80-.). 364. https://doi.org/10.1126/science.aat8078

Möller-Leimkühler, A.M., 2007. Gender differences in cardiovascular disease and comorbid depression. Dialogues Clin. Neurosci. 9, 71–83.

Montague, D., Weickert, C.S., Tomaskovic-Crook, E., Rothmond, D.A., Kleinman, J.E., Rubinow, D.R., 2008. Oestrogen receptor α localisation in the prefrontal cortex of three mammalian species. J. Neuroendocrinol. 20, 893–903. https://doi.org/10.1111/j.1365-2826.2008.01743.x

Moy, S.S., Nadler, J.J., Perez, A., Barbaro, R.P., Johns, J.M., Magnuson, T.R., Piven, J., Crawley, J.N., 2004. Sociability and preference for social novelty in five inbred strains: An approach to assess autistic-like behavior in mice. Genes, Brain Behav. 3, 287–302. https://doi.org/10.1111/j.1601-1848.2004.00076.x

Müller-Ribeiro, F.C. de F., Zaretsky, D. V., Zaretskaia, M. V., Santos, R.A.S., DiMicco, J.A., Fontes, M.A.P., 2012. Contribution of infralimbic cortex in the cardiovascular response to acute stress. Am. J. Physiol. Integr. Comp. Physiol. 303, R639–R650. https://doi.org/10.1152/ajpregu.00573.2011

Myers-Schulz, B., Koenigs, M., 2012. Functional anatomy of ventromedial prefrontal cortex: Implications for mood and anxiety disorders. Mol. Psychiatry. https://doi.org/10.1038/mp.2011.88

Myers, B., 2016. Corticolimbic regulation of cardiovascular responses to stress. Physiol. Behav. 172, 49–59. https://doi.org/10.1016/j.physbeh.2016.10.015

Myers, B., Carvalho-Netto, E., Wick-Carlson, D., Wu, C., Naser, S., Solomon, M.B., Ulrich-Lai, Y.M., Herman, J.P., 2016. GABAergic Signaling within a Limbic-Hypothalamic Circuit Integrates Social and Anxiety-Like Behavior with Stress Reactivity. Neuropsychopharmacology 41, 1530–9. https://doi.org/10.1038/npp.2015.311

Myers, B., Mark Dolgas, C., Kasckow, J., Cullinan, W.E., Herman, J.P., 2014a. Central stress-integrative circuits: Forebrain glutamatergic and GABAergic projections to the dorsomedial hypothalamus, medial preoptic area, and bed nucleus of the stria terminalis. Brain Struct. Funct. 219. https://doi.org/10.1007/s00429-013-0566-y

Myers, B., McKlveen, J.M., Herman, J.P., 2014b. Glucocorticoid actions on synapses, circuits, and behavior: Implications for the energetics of stress. Front. Neuroendocrinol. https://doi.org/10.1016/j.yfrne.2013.12.003

Myers, B., McKlveen, J.M., Morano, R., Ulrich-Lai, Y.M., Solomon, M.B., Wilson, S.P., Herman, J.P., 2017. Vesicular Glutamate Transporter 1 Knockdown in Infralimbic Prefrontal Cortex Augments Neuroendocrine Responses to Chronic Stress in Male Rats. Endocrinology 158, 3579–3591. https://doi.org/10.1210/en.2017-00426

Naqvi, T.Z., Naqvi, S.S.A., Merz, C.N.B., 2005. Gender differences in the link between depression and cardiovascular disease. Psychosom. Med. https://doi.org/10.1097/01.psy.0000164013.55453.05

Öngür, D., Ferry, A.T., Price, J.L., 2003. Architectonic subdivision of the human orbital and medial prefrontal cortex. J. Comp. Neurol. 460, 425–449. https://doi.org/10.1002/cne.10609

Pratt, W.E., Mizumori, S.J.., 2001. Neurons in rat medial prefrontal cortex show anticipatory rate

changes to predictable differential rewards in a spatial memory task. Behav. Brain Res. 123, 165–183. https://doi.org/10.1016/S0166-4328(01)00204-2

Radley, J.J., Arias, C.M., Sawchenko, P.E., 2006. Regional differentiation of the medial prefrontal cortex in regulating adaptive responses to acute emotional stress. J. Neurosci. 26, 12967–12976. https://doi.org/10.1523/JNEUROSCI.4297-06.2006

Radley, J.J., Rocher, A.B., Rodriguez, A., Ehlenberger, D.B., Dammann, M., Mcewen, B.S., Morrison, J.H., Wearne, S.L., Hof, P.R., 2008. Repeated stress alters dendritic spine morphology in the rat medial prefrontal cortex. J. Comp. Neurol. 507, 1141–1150. https://doi.org/10.1002/cne.21588

Rau, A.R., Hentges, S.T., 2017. The relevance of AgRP neuron-derived GABA inputs to POMC neurons differs for spontaneous and evoked release. J. Neurosci. 37, 7362–7372. https://doi.org/10.1523/JNEUROSCI.0647-17.2017

Roberts, A.C., Clarke, H.F., 2019. Why we need nonhuman primates to study the role of ventromedial prefrontal cortex in the regulation of threat- And reward-elicited responses. Proc. Natl. Acad. Sci. U. S. A. 116, 26297–26304. https://doi.org/10.1073/pnas.1902288116

Sankar, T., Chakravarty, M.M., Jawa, N., Li, S.X., Giacobbe, P., Kennedy, S.H., Rizvi, S.J., Mayberg, H.S., Hamani, C., Lozano, A.M., 2019. Neuroanatomical predictors of response to subcallosal cingulate deep brain stimulation for treatment-resistant depression. J. Psychiatry Neurosci. 44, 1–10. https://doi.org/10.1503/jpn.180207

Schaeuble, D., Packard, A.E.B., McKlveen, J.M., Morano, R., Fourman, S., Smith, B.L., Scheimann, J.R., Packard, B.A., Wilson, S.P., James, J., Hui, D.Y., Ulrich-Lai, Y.M., Herman, J.P., Myers, B., 2019. Prefrontal Cortex Regulates Chronic Stress-Induced Cardiovascular Susceptibility. J. Am. Heart Assoc. 8, e014451. https://doi.org/10.1161/JAHA.119.014451

Sgoifo, A., Carnevali, L., Pico Alfonso, M.D.L.A., Amore, M., 2015. Autonomic dysfunction and heart rate variability in depression. Stress. https://doi.org/10.3109/10253890.2015.1045868

Shansky, R.M., Hamo, C., Hof, P.R., Lou, W., McEwen, B.S., Morrison, J.H., 2010. Estrogen promotes stress sensitivity in a prefrontal cortex-amygdala pathway. Cereb. Cortex 20, 2560–2567. https://doi.org/10.1093/cercor/bhq003

Shansky, R.M., Hamo, C., Hof, P.R., McEwen, B.S., Morrison, J.H., 2009. Stress-induced dendritic remodeling in the prefrontal cortex is circuit specific. Cereb. Cortex 19, 2479–2484. https://doi.org/10.1093/cercor/bhp003

Shoemaker, J.K., Badrov, M.B., Al-Khazraji, B.K., Jackson, D.N., 2016. Neural control of vascular function in skeletal muscle. Compr. Physiol. 6, 303–329. https://doi.org/10.1002/cphy.c150004

Sierra-Mercado, D., Padilla-Coreano, N., Quirk, G.J., 2011. Dissociable roles of prelimbic and infralimbic cortices, ventral hippocampus, and basolateral amygdala in the expression and extinction of conditioned fear. Neuropsychopharmacology 36, 529–538. https://doi.org/10.1038/npp.2010.184

Smith, B.L., Morano, R.L., Ulrich-Lai, Y.M., Myers, B., Solomon, M.B., Herman, J.P.,. Adolescent environmental enrichment prevents behavioral and physiological sequelae of adolescent chronic stress in female (but not male) rats. Stress 21. https://doi.org/10.1080/10253890.2017.1402883

Solomon, M.B., Loftspring, M., De Kloet, A.D., Ghosal, S., Jankord, R., Flak, J.N., Wulsin, A.C., Krause, E.G., Zhang, R., Rice, T., McKlveen, J., Myers, B., Tasker, J.G., Herman, J.P., 2015. Neuroendocrine function after hypothalamic depletion of glucocorticoid receptors in male and female mice. Endocrinology 156, 2843–2853. https://doi.org/10.1210/en.2015-1276

Stack, A., Carrier, N., Dietz, D., Hollis, F., Sorenson, J., Kabbaj, M., 2010. Sex differences in social interaction in rats: Role of the immediate-early gene zif268.

Neuropsychopharmacology 35, 570–580. https://doi.org/10.1038/npp.2009.163
Stamatakis, A.M., Stuber, G.D., 2012. Activation of lateral habenula inputs to the ventral
midbrain promotes behavioral avoidance. Nat. Neurosci. 15, 1105–1107.
https://doi.org/10.1038/nn.3145
Tavares, R.F., Corrêa, F.M.A., Resstel, L.B.M., 2009. Opposite role of infralimbic and prelimbic
cortex in the tachycardiac response evoked by acute restraint stress in rats. J. Neurosci.
Res. 87, 2601–2607. https://doi.org/10.1002/jnr.22070
Thayer, J.F., Åhs, F., Fredrikson, M., Sollers, J.J., Wager, T.D., 2012. A meta-analysis of heart
rate variability and neuroimaging studies: Implications for heart rate variability as a marker
of stress and health. Neurosci. Biobehav. Rev.
https://doi.org/10.1016/j.neubiorev.2011.11.009
Uylings, H.B.M., Groenewegen, H.J., Kolb, B., 2003. Do rats have a prefrontal cortex? Behav.
Brain Res. https://doi.org/10.1016/j.bbr.2003.09.028
Vahl, T.P., Ulrich-Lai, Y.M., Ostrander, M.M., Dolgas, C.M., Elfers, E.E., Seeley, R.J., D'Alessio,
D.A., Herman, J.P., 2005. Comparative analysis of ACTH and corticosterone sampling
methods in rats. Am. J. Physiol. Endocrinol. Metab. 289, E823-8.
https://doi.org/10.1152/ajpendo.00122.2005
Vertes, R.P., 2004. Differential Projections of the Infralimbic and Prelimbic Cortex in the Rat.
Synapse 51, 32–58. https://doi.org/10.1002/syn.10279
Vijayakumar, N., Cheng, T.W., Pfeifer, J.H., 2017. Neural correlates of social exclusion across
ages: A coordinate-based meta-analysis of functional MRI studies. Neuroimage.
https://doi.org/10.1016/j.neuroimage.2017.02.050
Wei, J., Yuen, E.Y., Liu, W., Li, X., Zhong, P., Karatsoreos, I.N., McEwen, B.S., Yan, Z., 2014.
Estrogen protects against the detrimental effects of repeated stress on glutamatergic
transmission and cognition. Mol. Psychiatry 19, 588–598.
https://doi.org/10.1038/mp.2013.83
Wood, J.N., Grafman, J., 2003. Human prefrontal cortex: Processing and representational
perspectives. Nat. Rev. Neurosci. 4, 139–147. https://doi.org/10.1038/nrn1033
Wood, M., Adil, O., Wallace, T., Fourman, S., Wilson, S.P., Herman, J.P., Myers, B., 2018.
Infralimbic prefrontal cortex structural and functional connectivity with the limbic forebrain: a
combined viral genetic and optogenetic analysis. Brain Struct. Funct. 1–25.
https://doi.org/10.1007/s00429-018-1762-6
Wulsin, L.R., Horn, P.S., Perry, J.L., Massaro, J.M., D'Agostino, R.B., 2015. Autonomic
Imbalance as a Predictor of Metabolic Risks, Cardiovascular Disease, Diabetes, and
Mortality. J. Clin. Endocrinol. Metab. 100, 2443–2448. https://doi.org/10.1210/jc.2015-1748
Yuen, E.Y., Wei, J., Yan, Z., 2016. Estrogen in prefrontal cortex blocks stress-induced cognitive
impairments in female rats. J. Steroid Biochem. Mol. Biol.
https://doi.org/10.1016/j.jsbmb.2015.08.028
39
40

1   ***Supplemental Data***
2
3
4
5   **Sexually divergent cortical control of affective-autonomic integration**
6
7
8
9   Tyler Wallace[1], Derek Schaeuble[1], Sebastian A. Pace, Morgan K. Schackmuth, Shane T.
10   Hentges, Adam J. Chicco, Brent Myers
11
12
13
14   **Injection and cannula placement**

15   Viral vectors were injected in the IL. Fiber optic cannulas were targeted to the dorsal

16   boundary of the IL, allowing cone-shaped illumination ventral to the fiber tip to stimulate IL

17   CaMKIIα-positive neurons (**Fig. S1**).

19
Figure S1: Photomicrograph of IL cannulation site. 10x image of cannula tract targeting the
IL with AAV-delivered YFP fluorescence. White arrow indicates location of fiber tip. Dashed
white lines indicate region borders. ACB - nucleus accumbens, aco – anterior commissure,
CCfa – corpus callosum anterior forceps, IL – infralimbic, PL- prelimbic.

Stimulation-induced c-Fos expression

To determine the impact of light stimulation on neural activity, we quantified c-Fos

positive nuclei labeled through chromogen immunohistochemistry (**Fig. S2**). Stimulation

increased c-Fos positive nuclei in males (n = 6-7/group, Welch's unpaired t-test: ChR2 vs YFP

$t_{(6.416)}$ =3.268, $p < 0.05$) and females (n = 6-7/group, unpaired t-test: ChR2 vs YFP $t_{(11)}$ = 4.475,

$p < 0.005$). These results indicate, that at the termination of experimentation, 10 Hz, 1 mW

pulsatile stimulation significantly increased cellular activation in the IL of both sexes. While the

mean value of c-Fos positive cells appears higher in males, there was no significant (p = 0.08)

sex difference between the ChR2 groups.

Figure S2: Optogenetic stimulation increased c-Fos positive cell count in males and
females. At the conclusion of experiments 2 and 3**,** rats received 5 minutes of optic stimulation
(10 Hz, 5 ms pulses, 1 mW) followed by 90 minutes of recovery prior to tissue collection.
Chromogen immunohistochemistry was used to identify and quantify c-Fos positive cells in the
IL. * p < 0.05, *** p < 0.005 vs YFP within sex.

1   **Social novelty preference**

2   Time spent interacting with novel and familiar rats. Stimulation of IL pyramidal

3   neurons did not affect preference for social novelty in males or females.

Experiment	Virus	Novel interaction (% time)	Novel interaction (SEM)	Familiar interaction (% time)	Familiar interaction (SEM)	n
Male 20 Hz	YFP	16.63	1.27	13.06	2.81	7
	ChR2	12.84	1.72	9.73	0.94	10
Male 10 Hz	YFP	16.43	2.07	13.13	2.17	12
	ChR2	18.19	1.81	16.32	2.47	10
Female 10 Hz	YFP	24.84	2.61	16.56	1.62	12
	ChR2	21.58	1.55	16.96	1.51	16

5
6
7   **Table S1: Social novelty preference.**
8

Homecage stimulation

Following behavioral experiments, male rats from experiment 2 and female rats from

experiment 3 went through a homecage stimulation protocol the week prior to novel

environment. Each rat was removed from the homecage, tethered to a fiber optic patch cord,

and then returned to the cage. After 2 minutes, optics were turned on (0.5 mW, 5 ms pulses, 5

Hz) for one minute then turned off for one minute. This procedure repeated for 10 Hz and 20 Hz

stimulation before power was increased to 1.3 mW and the protocol repeated. In total, each rat

was tethered for 14 minutes and received stimulation for 6 minutes. For male rats, heart rate

(HR) and mean arterial pressure (MAP) were recorded. Photostimulation of the IL did not affect

HR or MAP (**Fig. S3**). Specifically, the 10 Hz stimulation used for cardiovascular reactivity

studies had no effect at 0.5 or 1.3 mW on HR (n = 7-8, Mann-Whitney: ChR2 vs YFP 10 Hz 0.5

mW U = 25, p = 0.78; ChR2 vs YFP 10 Hz 1.3 mW U = 24, p = 0.69) or MAP (n = 7-8, Mann-

Whitney: ChR2 vs YFP 10 Hz 0.5 mW U = 27, p = 0.96; ChR2 vs YFP 10 Hz 1.3 mW U = 21, p

= 0.46). Although, there was an overall trend for decreased HR and MAP as a function of time

post-handling. Hemodynamic parameters were not recorded in females.

Figure S3: Optogenetic stimulation did not affect homecage HR or MAP in non-stressed
male rats. HR and MAP were recorded over 2-minute periods of 10 Hz stimulation (0.5 and 1.3
mW).

Baseline hemodynamic measures

The weekend prior to novel environment, baseline cardiovascular measurements were

recorded at approximately the same time of day that experiments were conducted. Data were

collected while rats were untethered in their homecages without optic stimulation (**Fig. S4**). In

males, HR (n = 7-8, Mann-Whitney: ChR2 vs YFP U = 19, p = 0.34) and MAP (n = 7-8, Mann-

Whitney: ChR2 vs YFP U = 23, p = 0.59) were not different. Likewise, female HR (n = 7-8,

Mann-Whitney: ChR2 vs YFP U = 25, p = 0.78) and MAP (n = 7-8, Mann-Whitney: ChR2 vs

YFP U = 28, p > 0.99) had no differences.

Figure S4: Baseline HR and MAP in male and female rats. Male rats **(A,B)** had no
differences in baseline HR or MAP (p > 0.05). In female rats **(C,D)**, there were also no
differences in HR or MAP (p > 0.05).

14