A human monoclonal antibody targeting a conserved pocket in the SARS-CoV-2
receptor-binding domain core

Juliette Fedry[1=], Daniel L. Hurdiss[1,2=], Chunyan Wang[2=], Wentao Li[2=], Gonzalo Obal[3], Ieva
Drulyte[4], Stuart C. Howes[1], Frank J.M. van Kuppeveld[2], Friedrich Förster[1#] and Berend-Jan
Bosch[2#]

[1]Cryo-Electron Microscopy, Bijvoet Centre for Biomolecular Research, Department of Chemistry, Faculty of Science, Utrecht
University, Padualaan 8, 3584 CH Utrecht, The Netherlands
[2]Virology Section, Infectious Diseases and Immunology Division, Department of Biomolecular Health Sciences, Faculty of
Veterinary Medicine, Utrecht University, 3584CL Utrecht, The Netherlands.
[3]Crystal and Structural Chemistry, Bijvoet Centre for Biomolecular Research, Department of Chemistry, Faculty of Science,
Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands
[4]Materials and Structural Analysis, Thermo Fisher Scientific, Eindhoven, 5651 GG, The Netherlands.

= Contributed equally to this work.
Address correspondence to Friedrich Förster (f.g.forster@uu.nl) and Berend-Jan Bosch
(b.j.bosch@uu.nl).

Abstract

SARS-CoV-2 has caused a global outbreak of severe respiratory disease (COVID-19), leading
to an unprecedented public health crisis. To date, there has been over 33 million diagnosed
infections, and over one million deaths. No vaccine or targeted therapeutics are currently
available. We previously identified a human monoclonal antibody, 47D11, capable of cross-
neutralising SARS-CoV-2 and the related 2002/2003 SARS-CoV. Here we present the
structural basis of its neutralization mechanism. We describe cryo-EM structures of trimeric
SARS-CoV and SARS-CoV-2 spike ectodomains in complex with the 47D11 Fab. These data
reveal that 47D11 binds specifically to the closed conformation of the receptor binding domain,
distal to the ACE2 binding site. The CDRL3 stabilises the N343 glycan in an upright
conformation, exposing a conserved and mutationally constrained hydrophobic pocket, into
which the CDRH3 loop inserts two aromatic residues. Interestingly, 47D11 preferentially
selects for the partially open conformation of the SARS-CoV-2 spike, suggesting that it could
be used effectively in combination with other antibodies that target the exposed receptor-
binding motif. Taken together, these results expose a cryptic site of vulnerability on the SARS-
CoV-2 RBD and provide a structural roadmap for the development of 47D11 as a prophylactic
or post-exposure therapy for COVID-19.

Introduction

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged from a zoonotic event in China, late 2019(*1*). To date, the resulting coronavirus induced disease 19 (COVID-19) pandemic has been responsible for over 33 million infections and over a million deaths, as of September 29, 2020(https://covid19.who.int/). SARS-CoV-2 and SARS-CoV, another highly lethal respiratory pathogen which emerged in 2002/2003(*2*), belong to the *Sarbecovirus* subgenus (genus *Betacoronavirus*, family *Coronaviridae*)(*3*). At present, no targeted therapeutics have been approved for COVID-19, meaning there is an urgent clinical need for potent antiviral therapies to halt the spread of SARS-CoV-2 and to pre-empt future outbreaks caused by SARS-like viruses. Antibodies are a promising class of drugs for combatting infectious diseases and have shown therapeutic efficacy for a number of viruses(*4, 5*), including in the treatment of SARS and COVID-19(*6, 7*). Such antibodies function by targeting vulnerable sites on viral surface proteins.

The coronavirus trimeric spike (S) glycoprotein, located on the viral envelope, is the key mediator of viral entry into host cells. The spike protein consists of two main parts: S1 is involved in receptor binding and S2 is the membrane fusion domain. The S1 domain itself is further subdivided into an N-terminal domain (NTD, or S1A) and a receptor binding domain (RBD, or S1B)(*8, 9*). The spike proteins of SARS-CoV-2 (SARS2-S; 1273 residues, strain Wuhan-Hu-1) and SARS-CoV (SARS-S, 1255 residues, strain Urbani) exhibit 77.5% identity in their primary amino acid sequence and are structurally conserved. The spike trimer exists in equilibrium between a closed conformation, where all three RBD are lying flat, and a partially open conformation, where one RBD stands upright and is exposed for receptor engagement(*10-12*). Both viruses use the human angiotensin converting enzyme 2 (ACE2) protein as a host receptor, with binding mediated through interactions with the receptor-binding motif (RBM) located on the RBD, and the N-terminal helix of ACE2(*13*). The spike-mediated fusion of viral and cellular membranes is tightly regulated and triggered by a cascade of preceding events. The first step involves the attachment of SARS-CoV-2 to the target cell surface via the interaction between spike and ACE2(*13, 14*). In the second step, the spike protein needs to be primed for membrane fusion by host proteases (e.g. cellular transmembrane serine protease 2) which cleave the spike at multiple sites(*15*), enabling shedding of S1. Finally, the free S2 catalyses the fusion of the viral and the host membranes(*16, 17*), causing the release of the viral genome into the host cell cytoplasm.

The S glycoprotein is the primary target for neutralising antibodies, making it the main target for vaccine development(*18*). Indeed, a number of SARS-CoV-2 neutralising antibodies have

now been described(*19-33*). However, comparatively few cross-neutralising antibodies have
been reported(*20, 25, 30, 34*), of which only a handful have been structurally characterised(*29,*
35-37). The most commonly identified antibodies neutralize coronaviruses by binding to the
receptor interaction site in S1, blocking receptor interactions and/or promoting premature
conformational change of spike to the post-fusion state. However, a smaller number of
antibodies have been reported to bind sites which are distal to the ACE2 binding site. Such
antibodies target the RBD-core(*27, 35, 37, 38*), or even the NTD(*39*). Several promising
vaccines are currently being developed but it is estimated that at least a year will be needed
before they can be introduced on the market. Hence there is an urgent need for characterized
antibodies to form cocktails for the treatment by passive immunization of COVID-19 patients.
Combined structural and functional studies are thus required to determine the epitopes and
investigate the molecular mechanisms of SARS-CoV-2 neutralizing antibodies. Moreover,
such studies may identify cryptic sites of vulnerability which can guide vaccine and antiviral
development(*40*).
90
91 We recently reported the first human monoclonal antibody, 47D11, capable of cross-
92 neutralising SARS-CoV and SARS-CoV-2 at 1.3 and 3.8 nM, respectively, but its mode of
93 engagement with the spike protein remained unclear. We thus employed structural and
94 functional studies to decipher the molecular basis for 47D11-mediated neutralisation.
95
Results
47D11 specifically binds to the closed receptor binding domain
To understand how 47D11 binds to the SARS-CoV and SARS-CoV-2 spike proteins, we used
cryo-electron microscopy (cryo-EM) to determine structures of prefusion stabilised
ectodomain trimers in complex with the 47D11 Fab fragment. The resulting cryo-EM maps
have global resolutions of 3.8 Å and 4.0 Å resolution for SARS-S and SARS2-S, respectively
(Supplementary Figure1A-F). For previously reported apo S trimers, both the open and closed
conformation are observed, with the latter being predominant (56% for SARS and 67% for
SARS2(*11, 41*)). Upon incubation with 47D11, only the closed conformation of the SARS spike
was observed, with stochiometric binding of 47D11 to each RBD (Figure 1A). Interestingly, for
SARS-CoV-2, only the partially open conformation of spike was observed, with one Fab bound
to each of the closed RBDs, and the remaining open RBD unoccupied and, in principle,
accessible to ACE2 binding (Figure 1B). The sub stochiometric binding observed for SARS2-
S may partially explain our previous observations that 47D11 binds to the SARS-S with higher
affinity than SARS2-S (equilibrium dissociation constant [K_D] of 0.745 nM and 10.8 nM,
respectively)(*42*). To understand why 47D11 favours different spike conformations for SARS-
CoV and SARS-CoV-2, we first superposed the Fab-bound structures with their apo

counterparts. Compared to the apo partially open SARS2-S structure, the RBDs are less
compact when 47D11 is bound (Figure 2A). The apo conformation of the closed RBD would
preclude binding of 47D11 through steric hindrance. To accommodate the bound Fab, this
RBD shifts outwards by ~7 Å (Figure 2B). Unlike S309 and H014(35, 38), two other RBD-core
targeting SARS neutralizing antibodies, there was no indication from our cryo-EM data that
47D11 can bind to the open conformation of the SARS2-S RBD. In line with this,
superimposition of open and closed SARS2-S RBDs revealed that 47D11 would clash with
the adjacent N-terminal domain (NTD) and the N331 glycan in the latter conformation (Figure
2C). Similar to SARS2-S, the RBDs of the 47D11 bound SARS-S are also less compact than
the reported apo fully closed structure (Figure 2D). However, in contrast to SARS2-S, there is
a potential stabilising salt bridge between D463, located on the receptor binding ridge (RBR),
and R18 on the 47D11 light chain (Figure 2E). Indeed, the RBR exhibits the most prominent
structural differences between SARS2-S and SARS-S(13). This epitope distal loop, located
within the ACE2-binding region, contains an essential disulfide bridge in both viruses, but is
more compact in SARS-S. In order to test whether the epitope distal RBR impacts binding of
47D11 to the SARS-S and SARS2-S, we swapped loop residues 470-490 (SARS2-S
numbering) and produced chimeric ectodomains. In support of our hypothesis, the SARS2-S,
containing the SARS-S RBR loop, exhibited increased binding to 47D11. However, we did not
observe an equivalent loss of binding for the chimeric SARS-S, suggesting that other
differences in protein sequence or quaternary structure may be involved (Figure 2F). Taken
together, our data shows that 47D11 binding to the closed RBDs of the trimeric spike protein
has differing outcomes for SARS-S and SARS2-S, trapping them in the fully closed and the
partially open conformation, respectively (Figure S2).
136

47D11 targets a conserved hydrophobic pocket in the RBD

The 47D11 epitope is distinct from the ACE2 binding site (Figure 3A), consistent with our
recently reported functional data(42). The protein/glycan epitope is located on the core domain
of the SARS-S and SARS2-S RBD. As expected, the mode of binding is highly similar for
SARS-S and SARS2-S (Figure S3A), with the aligned 47D11:RBD complexes having an
RMSD value of 1.4 Å. The paratope is composed of CDRL3 and CDRH3 loops, that form a
primarily hydrophobic interaction with the RBD surface of ~830 Å^2 and ~800 Å^2 for SARS-S
and SARS2-S, respectively. The side chain of 47D11 CDRL3 tryptophan W94 stacks against
the N330/N343 (SARS/SARS2) glycan tree, contributing to its stabilization in an upright
conformation (Figure 3B). This reveals a hydrophobic pocket into which the CDRH3 loop
projects, allowing Fab residues W102 and F103 to interact with RBD core residues F338,
F342, Y365, V367, L368, F374 and W436 (F325, F329, Y352, V354, L355, F361 and W423
in SARS-S) – figure 3B and S3B. Interestingly, this pocket is normally shielded by the N343

glycan in previously reported SARS2-S structures (Figure 3C-D)(*11, 12*). In order to
accommodate the CDRH3 loop residues, the helix encompassing residues 365-370 is
displaced outwards by 2 Å, creating 55 Å^3 of solvent accessible volume which is not present
in the apo RBD (Figure 3C-D and Figure S3C-D). Of note, the region directly below this
hydrophobic pocket was recently shown bind to linoleic acid, which stabilises the closed
conformation of spike by spanning two adjacent RBDs(*43*). However, the distance between
the 47D11-bound RBDs is too great to be bridged by linoleic acid. Consistent with this, no
density consistent with linoleic acid was present in any of our reconstructions.
158
To verify the 47D11 epitope, we introduced alanine mutations at each of the identified contact
residues in the context of full-length spike protein. In addition, a spike mutant with the naturally
occurring V367F minority variant was generated(*44*). Binding of 47D11 to surface expressed
wildtype and mutant spike proteins was assessed by flow cytometry. Soluble Fc-tagged ACE2
and the RBD core binding mAb CR3022 were taken along as controls. The V367F substitution
and the alanine substitute at this position only had a minor effect on 47D11 antibody binding
(Figure 3E), consistent with data showing that this polymorphism had no effect on
neutralisation of SARS2-S pseudo type virus (Figure 3I). Collectively, this indicates that 47D11
would be effective against this SARS-CoV-2 variant. In contrast, all other amino acid
substitutions in the hydrophobic core not only reduced cell-surface binding by 47D11 (Figure
3E), but also prevented binding of ACE2 and the core targeting antibody CR3022, despite
being distal to their respective interaction sites (Figure 3F-G and Figure S4A-B). Total cellular
expression of mutants was comparable to wildtype spike protein as demonstrated by an
antibody targeting the C-terminal appended Flag-tag on the spike proteins (Figure 3H),
suggesting that mutations in the RBD hydrophobic core have a detrimental effect on protein
folding, compromising the tertiary structure of the RBD. A recent study reported deep
mutational scanning of SARS2-S RBD residues, revealing how mutation of each of the RBD
residues affects expression of folded protein and its affinity for ACE2(*45*). When the mean
mutation effect on expression was mapped on the 47D11 bound RBD, we observed that the
hydrophobic pocket, targeted by 47D11, is highly mutationally constrained (Figure 3J).
Another SARS-CoV and SARS-CoV-2 neutralising antibody, S309, targets a similar region to
47D11, but here the orientation of the N343 glycan prohibits access to the hydrophobic pocket,
similarly to apo structures (Figure 3K)(*35*). The 47D11 epitope is distinct from other reported
RBD-core targeting antibodies/nanobodies, such as CR3022, H014 and VHH-72 (Figure
S4)(*27, 38, 46*).
184
Comparative sequence analysis revealed that the 47D11 epitope is highly conserved across
circulating SARS-like viruses (Figure 4A and S5). This is in contrast to the ACE2 binding

region which exhibits the greatest sequence variability. In order to assess whether 47D11 has
broad reactivity, we recombinantly expressed the RBD from WIV16, HKU3-3 and HKU9-3 and
analysed 47D11 binding to these related sarbecoviruses. The results demonstrated that
47D11 can bind to the WIV16 RBD with similar affinity to SARS-S and SARS2-S (Figure 4B).
Closer inspections of the aligned RBD amino acid sequences revealed that the N343
glycosylation site, as well as the hydrophobic pocket, are strictly conserved between these
three SARS-like viruses (Figure 4C). Therefore, the lack of binding observed for HKU3-3 and
HKU9-3 may be due to epitope adjacent sequence differences which preclude binding of the
CDRH3 loop. Nevertheless, the potent binding observed for WIV16 underscores the potential
of 47D11 as a treatment for future outbreaks caused by SARS-like viruses.
197
In conclusion, our structural and functional analyses demonstrate that 47D11 is able to
unmask a conserved and mutationally constrained epitope on the SARS-CoV-2 RBD by
stabilising the N343 glycan in an upright conformation. Once this site is exposed, the CDRH3
loop is able to insert two aromatic residues into the hydrophobic core of the RBD, inducing
conformational changes which lead to the formation of a 55 Å^3 cavity. This cryptic site offers
an attractive target for design of vaccines and targeted therapeutics, such as small molecule
inhibitors. The 47D11 epitope is distal to the ACE2 receptor binding motif, rationalising its
ability to cross-neutralize SARS-CoV and SARS-CoV-2 independently of receptor-binding
inhibition. Our structural analysis also shows that 47D11 exhibits differing conformational
selectivity for the SARS-S and SARS2-S, providing a possible explanation for the differences
in observed binding affinity. Given that 47D11 selects for the partially open conformation of
the SARS-CoV-2 spike protein, we speculate that 47D11 may render the spike more
susceptible to other monoclonal antibodies which target the exposed receptor-binding
subdomain, making it a prime candidate for combination treatment. Antibody combinations
targeting non-overlapping epitopes may act synergistically permitting a lower dosage and an
increased barrier to immune escape(*47*). Genetic diversity for SARS2 is currently limited, as
the virus has gone through a genetic bottleneck during the singular animal-to-human spill-over
event. However, the genetic/antigenic variation will increase in time, as observed for the
endemic human coronavirus HCoV-229E, which exhibits cumulative sequence variation in the
RBD loops which engage its cellular receptor(48). The limited mutational space of the 47D11
epitope may confer the antibody sustainable applicability in neutralizing a wide range of future-
emerging virus variants.
220
221   **Methods**

Expression and purification of coronavirus spike proteins

To express the prefusion spike ectodomain, gene encoding residues 1−1200 of SARS2 S (GenBank: QHD43416.1) with proline substitutions at residues 986 and 987, a "AAARS" substitution at the furin cleavage site (residues 682–685) and residues 1-1160 of SARS S (GenBank: AAP13567.1) with proline substitutions at residues 956 and 957, a C-terminal T4 fibritin trimerization motif, a StrepTag was synthesized and cloned into the mammalian expression vector pCAGGS. Similarly, pCAGGS expression vectors encoding S1 or its subdomain $S1_B$ of SARS (S1, residues 1-676; $S1_B$, residues, 325-533), and SARS2 (S1, residues 1-682; $S1_B$, residues 333-527) C-terminally tagged with Fc domain of human or mouse IgG or Strep-tag were generated as described before(*42*). Recombinant proteins and antibody 47D11 were expressed transiently in FreeStyle™ 293-F Cells (Thermo Fisher Scientific) and affinity purified from the culture supernatant by protein-A sepharose beads (GE Healthcare) or streptactin beads (IBA) purification. Purity and integrity of all purified recombinant proteins was checked by coomassie stained SDS-PAGE.

Pseudotyped virus neutralization assay

Neutralization with SARS2-S VSV pseudotyped viruses was performed as described previously (24). HEK-293T cells were transfected with pCAGGS expression vectors encoding SARS2-S carrying a 18-a.a. cytoplasmic tail truncation, respectively. One day post transfection, cells were infected with the VSV-G pseudotyped VSVΔG expressing the firefly (*Photinus pyralis*) luciferase. Twenty-four hours later, cell supernatants containing SARS2-S pseudotyped VSV particles were harvested and titrated on African green monkey kidney VeroE6 (ATCC#CRL-1586) cells. In the virus neutralization assay, mAbs were threefold serially diluted and mixed with an equal volume of pseudotyped VSV particles and incubated for 1 hour at room temperature (RT). The virus/antibody mix was subsequently added to confluent VeroE6 monolayers in 96-well plate, and incubated at 37°C. After 24 hours, cells were washed and lysis buffer (Promega) was added. Luciferase activity was measured on a Berthold Centro LB 960 plate luminometer using D-luciferin as a substrate (Promega). The percentage of infectivity was calculated as ratio of luciferase readout in the presence of mAbs normalized to luciferase readout in the absence of mAb. The half maximal inhibitory concentration (IC_{50}) was determined using 4-parameter logistic regression (GraphPad Prism version 8).

ELISA analysis of antibody binding to CoV spike antigens

ELISA was performed as described previously (24). Briefly, NUNC Maxisorp plates (Thermo Scientific) coated with equimolar antigen amounts were blocked with 3% bovine serum albumin (Bio-Connect) in PBS containing 0.1% Tween-20 at RT for 2 hours. Fourfold serial

dilutions of mAbs starting at 10 µg/ml (diluted in blocking buffer) were added and plates were

incubated for 1 hour at RT. Plates were washed three times and incubated with horseradish

peroxidase (HRP)-conjugated goat anti-human secondary antibody (ITK Southern Biotech)

diluted 1:2000 in blocking buffer for 1 hour at RT. An HRP-conjugated anti-StrepMAb (IBA)

antibody was used to corroborate equimolar coating of the Strep-tagged spike antigens. HRP

activity was measured at 450 nanometer using tetramethylbenzidine substrate (BioFX) using

an ELISA plate reader (EL-808, Biotek). Half-maximum effective concentration (EC_{50}) binding

values were calculated by non-linear regression analysis on the binding curves using

GraphPad Prism (version 8).

Preparation of Fab-47D11 from IgG

47D11 Fab was digested from IgG with papain using a Pierce Fab Preparation Kit (Thermo

Fisher Scientific), following the manufacturer's standard protocol.

Cryo-EM sample preparation and data collection

3 µL of SARS2-S or SARS-S at 1.6 mg/mL was mixed with 0.85 µL of Fab 47D11 at 4 mg/mL

and incubated for 50 s at RT. The sample was applied onto a freshly glow discharged R1.2/1.3

Quantifoil grid in a Vitrobot Mark IV (Thermo Fisher Scientific) chamber pre-equilibrated at

4°C and 100% humidity. The grid was immediately blotted at force 0 for 5 s and plunged into

liquid ethane. Data was acquired on a 200 kV Talos Arctica (Thermo Fisher Scientific)

equipped with a Gatan K2 Summit direct detector and Gatan Quantum energy filter operated

in zero-loss mode with a 20 eV slit width. To account for the preferred orientation exhibited by

the spike ectodomains, automated data collection at tilts 0°, 20°, and 30° was carried out using

EPU 2 software (Thermo Fisher Scientific), and data at tilt 40° using SerialEM(49). A nominal

magnification of 130,000x, corresponding to an effective pixel size of 1.08 Å, was used. Movies

were acquired in counting mode with a total dose of 40e/$Å^2$ distributed over 50 frames. 4,231

movies were acquired for SARS2 and 3,247 movies for SARS-S, with defocus ranging

between 0.5 µm and 3 µm.

Cryo-EM data processing

Single-particle analysis was performed in Relion version 3.1.(*50*). The data was processed in

four separate batches, corresponding to the stage tilt angle used for the acquisition. Drift and

gain correction were performed with MotionCor2(*51*), CTF parameters were estimated using

CTFFind4(*52*) and particles were picked using the Laplacian picker in Relion(*50*). One round

of 2D classification was performed on each batch of data and particles belonging to well

defined classes were retained. Subsequently, 3D classification was performed, using a 50 Å

low-pass filtered partially open conformation as an initial model (EMD-21457,(*12*)), without

imposing symmetry. All particles belonging to Fab bound class were then selected for 3D auto-
refinement. Before merging the different batches, iterative rounds of per particle CTF
refinement, 3D auto-refinement and post-processing were used to account for the stage tilt
used during data collection. The refined particle star files from each batch were then combined
and subjected to a final round of 3D auto refinement, per particle defocus estimation, 3D auto-
refinement and post processing, both with and without imposed C3 symmetry. Overviews of
the single-particle image processing pipelines are shown in supplementary figure 6 and 7.
303

Model building and refinement
UCSF Chimera (version 1.12.0) and Coot (version, 1.0.) were used for model building and
analysis(*53*, *54*). The SARS2-S model, in the partially open conformation (one RBD up, pdb
6VYB)(*12*), was used for the spike and fitted into our density using the UCSF Chimera 'Fit in
map' tool(*53*). For SARS a closed protomer of the pdb 6NB6 was used as starting model(*55*).
To build a model for the Fab the sequence of the variable regions of the HC and the LC were
separately blasted against the pdb. For the HC variable region, the corresponding region of
the pdb 6IEB (human monoclonal antibody R15 against RVFV Gn) was used(*56*). The LC
variable region was modelled using the pdb 6FG1 as template (Fab Natalizumab)(*57*). For
both chains, the query sequence of 47D11 was aligned to the template sequence. Sequence
identity was particularly high (87% and 97% for the HC and LC, respectively). Phenix sculptor
was used to create an initial model for the Fab chains(*58*), removing the non-aligning regions
(notably the CDRH3). This model was fitted into our density and the missing regions were built
manually in the density map using Coot(*54*). Models were refined against the respective EM
density maps using Phenix Real Space Refinement and Isolde(*59*)(*60*), and validated with
MolProbity and Privateer (glycans)(*59*, *61-63*).
320

Analysis and visualisation
PDBePISA was used to identify spike residues interacting with 47D11(*64*). Surface colouring
of the SARS-CoV-2 RBD using the Kyte-Doolittle hydrophobicity scale was performed in
UCSF chimera(*53*). Volume measurements were performed using CASTp 3.0, using a probe
radius of 1.2 Å(*65*). In order to colour the 47D11 bound RBD surface according to each
residues mean mutational effect on expression, the pdb file was populated with the mean
mutation effect on expression values described by Starr et al(*45*). The UCSF Chimera
'MatchMaker' tool was used to obtain RMSD values, using default settings. Figures were
generated using UCSF Chimera(*53*) and UCSF ChimeraX(*66*).
330

331 **Figure Legends**

Figure 1: **47D11 has differing conformational selectivity for the SARS-CoV and SARS-CoV-2 spike.** A) Surface rendering of the fully closed SARS spike bound to three 47D11 antibody Fab fragments, shown as two orthogonal views. (B) Surface rendering of the partially open SARS2 spike in complex with two 47D11 antibody Fab fragments, shown as two orthogonal views. The spike protomers are coloured pink, blue and grey, and the 47D11 heavy and light chain are coloured yellow and purple, respectively. For clarity, only the Fab variable region is shown.

Figure 2: **47D11 binds specifically to the closed RBD and prevents their full compaction.** A) Top view of the 47D11 bound SARS2 spike. The spike protomers are coloured pink, blue and grey, and the 47D11 heavy and light chain are coloured yellow and purple, respectively. Glycans, and the N-terminal domain, are omitted for clarity and only the Fab variable region is shown. The superposed structure of the partially open apo SARS2 spike (PDB ID: 6ZGG) is shown as a silhouette. B) Zoomed in view of the boxed region in panel A. C) Zoomed in view of the SARS2 open RBD and adjacent NTD. The overlaid 47D11 Fab is shown semi-transparent and the N343 glycan is shown in ball-and-stick representation and coloured tan. D) Top view of the 47D11 bound SARS2 spike coloured as shown in panel A. The superposed structure of the closed apo SARS spike (PDB ID: 5XLR) is shown as a silhouette. E) Zoomed in view of the boxed region in panel D, showing a putative salt bridge between the 47D11 variable light chain and the RBD loop. F) ELISA-binding curves of 47D11 binding to wildtype and loop swapped spike ectodomains.

Figure 3: **The 47D11 epitope comprises a mutationally constrained hydrophobic pocket which is normally shielded by glycan N343.** A) Ribbon diagram of the SARS2-S receptor-binding domain (RBD) in complex with the 47D11 antibody Fab fragment. For comparison, residues 1-84 of the RBD bound ACE2 (PDB ID: 6M0J) are shown as a silhouette. B) Close-up view of the 47D11 epitope with the hydrophobic pocket residues shown as sticks and coloured dark blue. The N343 glycan is shown in ball-and-stick representation and coloured tan. For clarity, only the core pentasaccharide is shown. C) Slice through the surface rendered 47D11 bound SARS2-S RBD. D) Equivalent view as shown in panel C for the apo RBD (PDB ID:6VYB). E) Relative surface binding of 47D11, (F) ACE2, (G) CR3022 and (H) an anti-FLAG antibody to full-length SARS2 spike epitope mutants, determined by fluorescence-activated cell sorting. I) Antibody-mediated neutralization of infection of luciferase-encoding VSV particles pseudotyped with wild-type, V367A or V367F SARS2-S. J) Surface representation of the 47D11 bound SARS2-S RBD coloured according to mean mutation effect on expression (red indicates more constrained)(*45*). The Fab is shown as a ribbon diagram. K) As shown in E for the S309 bound SARS2 RBD.

Figure 4: The 47D11 epitope is conserved in SARS-like viruses. A) Surface representation

of the 47D11 bound RBD coloured according to sequence conservation across SARS-1,

SARS2 and 11 SARS-like viruses (Supplementary Figure 5). The 47D11 Fab variable chains

are shown as a ribbon diagram and coloured grey. Heavy chain residues W102 and F103 are

shown as sticks. For comparison, residues 1-84 of the RBD bound ACE2 (PDB ID: 6M0J) are

shown as a silhouette. B) ELISA-binding curves of 47D11 to the S1B domain of SARS,

SARS2, WIV16, HKU3-3 and HKU9-3. The average ± SD from two independent experiments

with technical duplicates is shown. C) Aligned RBD sequences of SARS2, SARS-1, WIV16,

HKU3-3 and HKU9-3. Key residues in the 47D11 epitope are indicated by red arrows.

 Figure 1

Closed SARS-CoV S trimer

Partially open SARS-CoV-2 S trimer

383   **Figure 2**

Figure 3

387 **Figure 4**

Data availability

Coordinates for the 47D11-bound SARS-CoV and SARS-CoV-2 spike proteins are deposited in the Protein Data Bank under accession codes XXXX and XXXX, respectively. The corresponding EM density maps have been deposited to the Electron Microscopy Data Bank under the accessions EMD-XXXX and EMD-XXXX. All reagents and relevant data are available from the authors upon request.

Acknowledgements

This work was supported by the European Research Council under the European Union's Horizon2020 Programme (ERC Consolidator Grant Agreement 724425 - BENDER). J.F. and D.L.H. are funded from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement (No 792575 and 842333). J.F. and D.L.H. also hold EMBO non-stipendiary long-term Fellowships (ALTF 948-2017 and ALTF 1172-2018). Research reported in this publication was supported by a China Scholarship Council grant to C.W. (CSC201708620178). We thank Dr Mihajlo Vanevic for IT support.

Author contributions

J.F., D.L.H., C.W., W.L., F.J.M.K., F.F. and B.J.B. conceived, designed, and coordinated the study. J.F., D.L.H., C.W., W.L., I.D., G.O., and S.H. conducted the experiments. F.J.M.K, F.F. and B.J.B. supervised part of the experiments. All authors contributed to the interpretations and conclusions presented. J.F. D.L.H. and B.J.B. wrote the manuscript, J.F., D.L.H, C.W., W.L. F.F. and B.J.B participated in editing the manuscript.

Competing interests

A patent application has been filed on 12 March 2020 on monoclonal antibodies targeting SARS-CoV-2 (United Kingdom patent application no. 2003632.3; patent applicants: Utrecht University, Erasmus Medical Center and Harbour BioMed). I.D. is an employee of Thermo Fisher Scientific. The other authors declare no competing interests.

Corresponding authors

Correspondence to Friedrich Förster (f.g.forster@uu.nl) and Berend-Jan Bosch (b.j.bosch@uu.nl).

References

1. P. Zhou *et al.*, A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. **579**, 270–273 (2020).

2. T. G. Ksiazek *et al.*, A novel coronavirus associated with severe acute respiratory syndrome. *N. Engl. J. Med.* **348**, 1953–1966 (2003).

3. Coronaviridae Study Group of the International Committee on Taxonomy of Viruses, (Nature Publishing Group, 2020), vol. 5, pp. 536–544.

4. P. Prabakaran *et al.*, Potent human monoclonal antibodies against SARS CoV, Nipah and Hendra viruses. *Expert Opin Biol Ther*. **9**, 355–368 (2009).

5. E. O. Saphire, S. L. Schendel, B. M. Gunn, J. C. Milligan, G. Alter, Antibody-mediated protection against Ebola virus. *Nat. Immunol.* **19**, 1169–1178 (2018).

6. A. Casadevall, L.-A. Pirofski, The convalescent sera option for containing COVID-19. *J. Clin. Invest.* **130**, 1545–1548 (2020).

7. A. C. Cunningham, H. P. Goh, D. Koh, Treatment of COVID-19: old tricks for new challenges. *Crit Care*. **24**, 91–2 (2020).

8. F. Li, Structure, Function, and Evolution of Coronavirus Spike Proteins. *http://dx.doi.org/10.1146/annurev-virology-110615-042301*. **3**, 237–261 (2016).

9. B.-J. Bosch, R. van der Zee, C. A. M. de Haan, P. J. M. Rottier, The coronavirus spike protein is a class I virus fusion protein: structural and functional characterization of the fusion core complex. *Journal of Virology*. **77**, 8801–8811 (2003).

10.   M. Gui *et al.*, Cryo-electron microscopy structures of the SARS-CoV spike glycoprotein
reveal a prerequisite conformational state for receptor binding. *Cell Res.* **27**, 119–129
(2017).

11.   D. Wrapp *et al.*, Cryo-EM structure of the 2019-nCoV spike in the prefusion
conformation. *Science*. **367**, 1260–1263 (2020).

12.   A. C. Walls *et al.*, Structure, Function, and Antigenicity of the SARS-CoV-2 Spike
Glycoprotein. *Cell*. **181**, 281–292.e6 (2020).

13.   J. Lan *et al.*, Structure of the SARS-CoV-2 spike receptor-binding domain bound to the
ACE2 receptor. *Nature*. **579**, 1–6 (2020).

14.   R. Yan *et al.*, Structural basis for the recognition of SARS-CoV-2 by full-length human
ACE2. *Science*. **367**, 1444–1448 (2020).

15.   M. Hoffmann *et al.*, SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is
Blocked by a Clinically Proven Protease Inhibitor. *Cell*. **181**, 271–280.e8 (2020).

16.   A. C. Walls *et al.*, Tectonic conformational changes of a coronavirus spike glycoprotein
promote membrane fusion. *Proc. Natl. Acad. Sci. U.S.A.* **114**, 11157–11162 (2017).

17.   Y. Cai *et al.*, Distinct conformational states of SARS-CoV-2 spike protein. *Science*,
eabd4251 (2020).

18.   D. van Riel, E. de Wit, Next-generation vaccine platforms for COVID-19. *Nat. Mater.*
19, 810–812 (2020).

19.   J. Hansen *et al.*, Studies in humanized mice and convalescent humans yield a SARS-
CoV-2 antibody cocktail. *Science*. **369**, 1010–1014 (2020).

20.   P. J. M. Brouwer *et al.*, Potent neutralizing antibodies from COVID-19 patients define
multiple targets of vulnerability. *Science*. **369**, 643–650 (2020).

21.   R. Shi *et al.*, A human neutralizing antibody targets the receptor-binding site of SARS-
CoV-2. *Nature*. **584**, 120–124 (2020).

22.   S. J. Zost *et al.*, Potently neutralizing and protective human antibodies against SARS-
CoV-2. *Nature*. **584**, 443–449 (2020).

23.   S. J. Zost *et al.*, Rapid isolation and profiling of a diverse panel of human monoclonal
antibodies targeting the SARS-CoV-2 spike protein. *Nat. Med.* **26**, 1422–1427 (2020).

24.   D. F. Robbiani *et al.*, Convergent antibody responses to SARS-CoV-2 in convalescent
individuals. *Nature*. **584**, 437–442 (2020).

25.   T. F. Rogers *et al.*, Isolation of potent SARS-CoV-2 neutralizing antibodies and
protection from disease in a small animal model. *Science*. **369**, 956–963 (2020).

26.   Y. Cao *et al.*, Potent neutralizing antibodies against SARS-CoV-2 identified by high-
throughput single-cell sequencing of convalescent patients' B cells. *Cell* (2020),
doi:10.1016/j.cell.2020.05.025.

27.   M. Yuan *et al.*, A highly conserved cryptic epitope in the receptor binding domains of
SARS-CoV-2 and SARS-CoV. *Science*. **368**, 630–633 (2020).

28. W. Li *et al.*, Potent neutralization of SARS-CoV-2 in vitro and in an animal model by a
human monoclonal antibody. *bioRxiv*. **525**, 129 (2020).

29. A. Z. Wec *et al.*, Broad neutralization of SARS-related viruses by human monoclonal
antibodies. *Science*. **369**, 731–736 (2020).

30. B. Ju *et al.*, Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature*.
584, 115–119 (2020).

31. Y. Wu *et al.*, A noncompeting pair of human neutralizing antibodies block COVID-19
virus binding to its receptor ACE2. *Science*. **368**, 1274–1278 (2020).

32. E. Seydoux *et al.*, Analysis of a SARS-CoV-2-Infected Individual Reveals Development
of Potent Neutralizing Antibodies with Limited Somatic Mutation. *Immunity*. **53**, 98–
105.e5 (2020).

33. C. O. Barnes *et al.*, Structures of Human Antibodies Bound to SARS-CoV-2 Spike
Reveal Common Epitopes and Recurrent Features of Antibodies. *Cell*. **182**, 828–
842.e16 (2020).

34. H. Lv *et al.*, Cross-reactive Antibody Response between SARS-CoV-2 and SARS-CoV
Infections. *Cell Rep*. **31**, 107725 (2020).

35. D. Pinto *et al.*, Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-
CoV antibody. *Nature*. **583**, 1–10 (2020).

36. D. Zhou *et al.*, Structural basis for the neutralization of SARS-CoV-2 by an antibody
from a convalescent patient. *Nat. Struct. Mol. Biol.* **20**, 773–9 (2020).

37. H. Liu *et al.*, Cross-neutralization of a SARS-CoV-2 antibody to a functionally conserved
site is mediated by avidity. *bioRxiv* (2020), doi:10.1101/2020.08.02.233536.

38. Z. Lv *et al.*, Structural basis for neutralization of SARS-CoV-2 and SARS-CoV by a
potent therapeutic antibody. *Science*. **369**, 1505–1509 (2020).

39. X. Chi *et al.*, A neutralizing human antibody binds to the N-terminal domain of the Spike
protein of SARS-CoV-2. *Science*. **369**, 650–655 (2020).

40. B. R. West *et al.*, Structural Basis of Pan-Ebolavirus Neutralization by a Human
Antibody against a Conserved, yet Cryptic Epitope. *MBio*. **9**, 365 (2018).

41. Y. Yuan *et al.*, Cryo-EM structures of MERS-CoV and SARS-CoV spike glycoproteins
reveal the dynamic receptor binding domains. *Nat Commun*. **8**, 15092–9 (2017).

42. C. Wang *et al.*, A human monoclonal antibody blocking SARS-CoV-2 infection. *Nat
Commun*. **11**, 2251–6 (2020).

43. C. Toelzer *et al.*, Free fatty acid binding pocket in the locked structure of SARS-CoV-2
spike protein. *Science*, eabd3255 (2020).

44. T. Koyama, D. Platt, L. Parida, Variant analysis of SARS-CoV-2 genomes. *Bull. World
Health Organ.* **98**, 495–504 (2020).

45. T. N. Starr *et al.*, Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain
Reveals Constraints on Folding and ACE2 Binding. *Cell* (2020),
doi:10.1016/j.cell.2020.08.012.

46. D. Wrapp *et al.*, Structural Basis for Potent Neutralization of Betacoronaviruses by
Single-Domain Camelid Antibodies. *Cell*. **181**, 1004–1015.e15 (2020).

47. M. A. Tortorici *et al.*, Ultrapotent human antibodies protect against SARS-CoV-2
challenge via multiple mechanisms. *Science*. **4**, eabe3354 (2020).

48. A. H. M. Wong *et al.*, Receptor-binding loops in alphacoronavirus adaptation and
evolution. *Nat Commun*. **8**, 1735–10 (2017).

49. M. Schorb, I. Haberbosch, W. J. H. Hagen, Y. Schwab, D. N. Mastronarde, Software
tools for automated transmission electron microscopy. *Nat. Methods*. **16**, 471–477
(2019).

50. J. Zivanov, T. Nakane, S. H. W. Scheres, A Bayesian approach to beam-induced
motion correction in cryo-EM single-particle analysis. *IUCrJ*. **6**, 5–17 (2019).

51. S. Q. Zheng *et al.*, MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat. Methods*. **14**, 331–332 (2017).

52. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron
micrographs. *Journal of Structural Biology*. **192**, 216–221 (2015).

53. E. F. Pettersen *et al.*, UCSF Chimera--a visualization system for exploratory research
and analysis. *J Comput Chem*. **25**, 1605–1612 (2004).

54. P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta
Crystallogr. D Biol. Crystallogr.* **60**, 2126–2132 (2004).

55. A. C. Walls *et al.*, Unexpected Receptor Functional Mimicry Elucidates Activation of
Coronavirus Fusion. *Cell*. **176**, 1026–1039.e15 (2019).

56. Q. Wang *et al.*, Neutralization mechanism of human monoclonal antibodies against Rift
Valley fever virus. *Nat Microbiol*. **4**, 1231–1241 (2019).

57. A. Cassotta *et al.*, A single T cell epitope drives the neutralizing anti-drug antibody
response to natalizumab in multiple sclerosis patients. *Nat. Med.* **25**, 1402–1407
(2019).

58. G. Bunkóczi, R. J. Read, Improvement of molecular-replacement models with Sculptor.
Acta Crystallogr. D Biol. Crystallogr. **67**, 303–312 (2011).

59. J. J. Headd *et al.*, Use of knowledge-based restraints in phenix.refineto improve
macromolecular refinement at low resolution. *Acta Crystallogr. D Biol. Crystallogr.* **68**,
381–390 (2012).

60. T. I. Croll, ISOLDE: a physically realistic environment for model building into low-
resolution electron-density maps. *Acta Crystallogr D Struct Biol*. **74**, 519–530 (2018).

61. V. B. Chen *et al.*, MolProbity: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 12–21 (2010).

62. J. Agirre *et al.*, Privateer: software for the conformational validation of carbohydrate
structures. *Nat. Struct. Mol. Biol.* **22**, 833–834 (2015).

63. J. Agirre, G. Davies, K. Wilson, K. Cowtan, Carbohydrate anomalies in the PDB. *Nat.
Chem. Biol.* **11**, 303–303 (2015).

64.   E. Krissinel, K. Henrick, Inference of macromolecular assemblies from crystalline state.
J. Mol. Biol. **372**, 774–797 (2007).

65.   W. Tian, C. Chen, X. Lei, J. Zhao, J. Liang, CASTp 3.0: computed atlas of surface
topography of proteins. *Nucleic Acids Res.* **46**, W363–W367 (2018).

66.   T. D. Goddard *et al.*, UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. *Protein Sci.* **27**, 14–25 (2018).

562