Stabilization demands of walking modulate the vestibular contributions to gait

Rina M. Magnani[1], Sjoerd M. Bruijn[2,3], Jaap H. van Dieën[2], Patrick A. Forbes[4] *

[1]Department of Physiotherapy, School of Physical Education and Physical Therapy, State University

of Goiás, Goiânia/GO, Brazil

[2]Department of Human Movement Sciences, Faculty of Behavioral and Movement Sciences, Vrije

Universiteit Amsterdam, Amsterdam Movement Sciences, Amsterdam, The Netherlands

[3]Institute Brain and Behavior Amsterdam, Amsterdam, The Netherlands

[4]Department of Neuroscience, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The

Netherlands

rinamagnani@gmail.com

s.m.bruijn@gmail.com

j.van.dieen@vu.nl

p.forbes@erasmusmc.nl

*Corresponding author:

Patrick Forbes
Department of Neuroscience
Erasmus MC
University Medical Center Rotterdam
P.O. Box 2040
3000 CA, Rotterdam
The Netherlands

Abstract *[200 words]*

Stable walking relies critically on motor responses to signals of head motion provided by the vestibular system, which are phase-dependent and modulated differently within each muscle. It is unclear, however, whether these vestibular contributions also vary according to the stability of the walking task. Here we investigate how vestibular signals influence muscles relevant for gait stability (medial gastrocnemius, gluteus medius and erector spinae) – as well as their net effect on ground reaction forces – while humans walked normally, with mediolateral stabilization, wide and narrow steps. We estimated local dynamic stability of trunk kinematics together with coherence of electrical vestibular stimulation (EVS) with muscle activity and mediolateral ground reaction forces. Walking with external stabilization increased local dynamic stability and decreased coherence between EVS and all muscles/forces compared to normal walking. Wide-base walking also decreased vestibulomotor coherence, though local dynamic stability did not differ. Conversely, narrow-base walking increased local dynamic stability, but produced muscle-specific increases and decreases in coherence that resulted in a net increase in vestibulomotor coherence with ground reaction forces. Overall, our results show that while vestibular contributions may vary with gait stability, they more critically depend on the stabilization demands (i.e. control effort) needed to maintain a stable walking pattern.

Introduction

The stability of walking is commonly assessed as the ability to maintain upright locomotion in the presence of self-generated and/or external perturbations [1-3]. To ensure stable walking, the nervous system relies on the integration and modulation of sensory signals from visual, somatosensory and vestibular sources to generate ongoing and/or corrective postural responses throughout the different phases of the gait cycle [4-7]. The vestibular system, for instance, which encodes signals of head movement in space [8], is assumed to contribute to gait stability because impaired gait is often observed in vestibulopathic patients [9,10]. Recent studies have further revealed that vestibular contributions to locomotion undergo phase- and muscle-specific responses that appear to align with each muscle's functional role in gait stability [11,12]. Specifically, the vestibular contributions to mediolateral stability during walking may be related to mediolateral foot placement produced by muscles around the hip [11,13,14] and to ankle torque at push-off driven by muscles of the ankle [15]. In the sagittal plane, however, corrective responses to a vestibular disturbance are almost entirely absent throughout the gait cycle [16]. Because stability in the sagittal plane is maintained largely passively [17] – due to passive dynamics of the legs [18] – it may be possible that the control of whole-body stability during locomotion requires less feedback-driven control as compared to the frontal plane [16]. Thus, the question arises whether changes in the stability of walking in the mediolateral plane also influences the vestibular contribution to stability during locomotion. Therefore, our aim is to identify how vestibular contributions to balance are modulated across varying stabilization demands of walking by characterizing changes in coupling of vestibular input to motor outputs during externally-imposed and natural variations in stability.

Experimentally, gait stability can be improved by adding external lateral stabilization to the body, thereby removing the need for the nervous system to control upright balance in the mediolateral direction [19-21]. Under these stabilized conditions, step width variability, trunk and pelvis motion and premotor cortical involvement all decrease compared to normal walking [22-25]. These observations indicate a reduced need to actively control gait stability, and as a result, may also diminish the necessity for vestibular sensory feedback control. This may be similar to observations during upright standing, where compensatory responses to mediolaterally-directed vestibular disturbances are absent when participants are stabilized in the mediolateral direction, despite having to maintain balance in an anterior-posterior direction [26]. Therefore, we first hypothesized that external lateral stabilization of the body would diminish the muscle and whole-body responses to a mediolateral vestibular disturbance during locomotion.

Natural changes in gait are also thought to influence stability. For instance, adopting a wider step width has been described as a response to decreased lateral stability during locomotion [22,27], and seems to be an effective approach to increase margins of stability (i.e. base of support) in older adults [28,29]. If the vestibular system influence on gait is tightly coupled to stability, then similar to external stabilization, we expected that the evoked muscle and whole-body responses to a vestibular

disturbance should diminish when walking with wider steps. On the other hand, narrow-base walking requires increased effort to control gait stability in the frontal plane in young and old adults [30,31], since the margins of stability, or tolerance for errors, are substantially reduced [32,33]. These observations, however, seem to contrast with measures of local dynamic stability, which quantify the likelihood of departing from a steady-state gait pattern in the absence of external disturbances, and instead indicate increased gait stability during narrow-base walking and decreased gait stability during wide-base walking [31,34]. We aim to test a potential explanation for this conflict by examining whether the increased local dynamic stability observed when walking with narrow step width is (partially) subserved by increased vestibular sensory feedback. Indeed, support for this is seen by the additional contribution of vestibular signals (via the lateral vestibular nuclei) to limb muscle activity in mice when walking on a narrow beam that is absent when walking on level ground [35]. Here, we tested these hypotheses by comparing muscle and whole-body responses evoked by a vestibular disturbance – together with estimates of stability – across normal, mediolaterally stabilized, wide-base and narrow-base walking.

Methods

Participants

We measured 23 healthy young adults between 24 and 33 years recruited from the university campus. Twelve participants were excluded during data analysis due to technical problems in the collection of electromyography (n=3) or kinematic data (n=9) in any of the trials recorded. Here we present results from eleven young adults (four females, 28.5±2.9 years old, 71.6±8.6 kg and 1.77±0.10 m, body mass index 22.2±3.1 kg/m^2). Exclusion criteria included self-reported history of injury and/or disfunction of the nervous, musculoskeletal or vestibular systems, or the use of medications that can cause dizziness. Participants were also instructed not to participate in intense physical exercise on the day of the experiment. The participants agreed to participate in the study by signing the informed consent form, and the study was approved by the VU Amsterdam Research Ethics Committee (VCWE-2017-158).

Electrical vestibular stimulation

A continuous electrical vestibular stimulus (EVS) was used to deliver an isolated vestibular disturbance to participants during all walking trials. Coupling of the electrical stimulus with muscle activity and ground reaction forces was quantified over the gait cycle to determine the magnitude and timing of the vestibular contribution to ongoing muscle and whole-body responses. The electrical stimulus modulates the afferent firing rate of both semicircular canal and otolith afferents [36-38], and when delivered in a binaural-bipolar configuration, EVS evokes a sensation of head roll rotational velocity [39] about an axis directed posteriorly and superiorly by 18° relative to the Reid plane [40-42]. When the head is facing forward, this stimulus configuration results in a postural response in the frontal plane to compensate for the induced roll error signal [16,26,43-46].

The electrical stimulus was applied to participants using flexible carbon rubber electrodes (9 cm^2). The electrodes were coated with Spectra 360 electrode gel (Parker La, USA) and fixed to participants' mastoid processes using adhesive tape and an elastic head band. The stimulus was delivered as an analog signal via a data acquisition board (National Instruments Corp., Austin, TX, USA) to an isolated constant current stimulator (STMISOLA, Biopac, Goleta, CA, USA). All participants were exposed to the same stochastic EVS designed with a limited bandwidth of 0 to 25 Hz [47], zero-mean low-pass filtered white noise, 25 Hz cutoff, zero lag, fourth-order Butterworth, peak amplitude of 5.0 mA, root mean square of ~ 1.2 mA, lasting 8 minutes and created with Matlab software (MathWorks, Natick, MA, USA). Because this binaural-bipolar stimulus oscillates around a zero-mean, the imposed sensations of roll motion, and the accompanying compensatory responses, occur in both a left and right direction.

Protocol

Participants walked on a dual-belt treadmill at a belt speed of 0.8 m/s in four different conditions: normal walking, stabilized walking, wide-base walking, and narrow-base walking. During normal walking and stabilized walking, participants were instructed to walk with their naturally preferred step width. Stabilized walking was achieved using a custom-made spring-loaded mediolateral pelvic stabilization frame. The stabilization frame was attached through springs to two carts that allowed for movement in the anterior-posterior direction. The springs were pre-tensioned to provide a stabilizing stiffness of 1260 N/m [48] and the height of the carts was aligned with the height of the pelvis for each subject [49]. During wide-base walking, participants were instructed to increase their step width beyond the approximate width of their hips. During narrow-base walking, participants were instructed to adopt a step width smaller than both the width of their hips and their usual step width. Throughout wide- and narrow-base trials, participants received repeated verbal instruction to maintain wider and narrower step widths, respectively, compared to normal walking. Participants walked in each condition for 8 minutes while being exposed to continuous EVS and were guided by the beat of a metronome at 78 steps/min to control for effects of varying cadence on vestibular contributions during locomotion [11,12]. The walking speed of 0.8 m/s and cadence 78 steps/min were chosen to replicate the conditions of Dakin et al. (2013) [11]. These walking parameters also ensure that vestibular-evoked balance responses, which are known to decrease as velocity and cadence increase [11,12], could be measured throughout the gait cycle.

Prior to starting the experiments, participants were allowed to walk for 3-4 minutes to familiarize themselves with walking on the treadmill at the specific cadence and with the electrical stimulus. In addition, participants walked for 2 minutes in each condition before the electrical stimulus was applied, which together with the familiarization period was considered a sufficient exposure period to remove acclimatization effects of walking on a treadmill [50,51]. Trial order for the different walking conditions was also randomized for each subject and subjects were given a short 5 minute break between trials to limit the influence of any long-term habituation to the electrical stimulus

178 throughout the walking trials [52]. Finally, participants maintained their head in a slightly extended
179 position with the Reid's plane pitched ~18° up from horizontal [40,41] by keeping a headgear-mounted
180 laser on a target located 3 m in front of them. This head position was chosen to maximize the
181 amplitude of vestibulomotor balance responses in the mediolateral direction [41,53,54].

Instrumentation

Kinematic data were recorded using a 3D motion capture system (Optotrak, Northern Digital
Inc., Waterloo, Ontario, Canada) sampling at 100 samples/s. Clusters of three light emitting diodes
(LED) were positioned at the occipital lobe, the spinous process of the sixth thoracic vertebra (T6),
the posterior superior iliac spine and at the calcaneus bilaterally. Ground reaction forces (GRF) were
measured from each belt by force plates embedded in the treadmill (Motekforce Link, The
Netherlands) at a sampling rate of 200 samples/s. In addition to analysis of the coupling between
EVS and mediolateral forces as described below, these signals were used for the identification of
toe-off and heel-strike and gait events.

Surface electromyography (EMG) (TMSI Porti system, TMSI Enschede, the Netherlands)
was collected at 2000 samples/s bilaterally from the medial gastrocnemius, gluteus medius and
erector spinae muscles, using pairs of disposable self-adhesive Ag/AgCl surface electrodes (Ambu,
Balerrup, Denmark; model Blue sensor; diameter 30x22mm) for each muscle. Electrodes were
placed over the recorded muscles according to SENIAM electrodes placement recommendations [55]
after abrading and cleaning the skin with alcohol. A reference electrode was placed over the medial
bony part of the left wrist (styloid process). The three muscles measured were chosen based on their
supposed roles in different stabilizing strategies that may be employed throughout the gait cycle and
across walking conditions. The medial gastrocnemius muscle (and other ankle plantarflexors) act as
the foot`s prime movers during normal walking [56] and are most sensitive to vestibular input in the
late stance phase [11,12,15] when modulating push-off force. The gluteus medius muscle contributes to
foot placement strategies; its activity is correlated to the next foot placement [57,58] and is most
sensitive to imposed vestibular errors just prior to heel strike [11]. Finally, trunk muscles serve to
directly influence trunk motion relative to the pelvis and are primarily activated to stabilize the trunk
during weight transfer around heel strike [59,60]. Therefore, erector spinae muscle may be especially
suited to contribute to angular momentum control of the torso during narrow base walking [61] when
push-off modulation and foot-placement are rendered ineffective.

Data analysis

Force-plate data were used first to calculate center of pressure positions, which were in turn
used to identify heel contacts [62]. From these estimates, stride time was calculated as the duration
between two consecutive heel strikes of the same foot and the step width was determined as the
mediolateral distance between the centroids of the feet cluster markers at heel strike. These
measures of limb kinematics (stride time and step width) were compared across walking conditions
to determine whether participants adhered to our instructions (that is, walk at the metronome-guided

cadence and with a modified step width). We compared muscle activity across each walking condition after rectifying and low-pass filtering (20 Hz cutoff, zero lag, sixth order Butterworth) the EMG signals and then time-normalizing the data per stride and averaging the data across the 256 strides. Prior to averaging, each rectified and low-pass filtered EMG signal was normalized to the maximum amplitude across all conditions.

To assess changes in local dynamic stability during the different walking conditions, we estimated the local divergence exponent (LDE). Measures of local dynamic stability for walking indicate the ability for a participant to return to the steady-state periodic motion after infinitesimally small perturbations [63], which occur, for example, through natural variability in the walking surface or the neuromuscular system. These measures have been shown to be particularly useful for detecting patients at risk of falling [3]. The LDE measures the exponential rate of divergence of neighboring trajectories of a state space constructed from kinematic data of gait [64], whereby an increasing LDE indicates reduced stability. We calculated the LDE using Rosenstein's algorithm [65] and as input the velocity of a maker placed over the T6 vertebrae, which was estimated using a three-point differentiation of the position trace [66]. Velocity time series were first resampled so that each time series of 256 strides (the minimum number of strides collected from every participant) contained 25600 samples. The LDE was then calculated as the slope of the mean divergence curve, whose horizontal axis was normalized by stride time from 0-0.5 stride [67,68].

To examine the vestibulomotor coupling between the input stimulus (i.e. EVS) and motor output (EMG and ground reaction forces) across our conditions, we computed the time-frequency coherence and gain assuming a linear stimulus-response relationship [69]. Prior to estimating coherence and gain, EVS, EMG and ground reaction forces were cut into segments synchronized to heel strikes. Based on the symmetry of our walking conditions, we chose to align the data on each limb's heel strike in order to pool responses from muscles in the left and right limbs. Symmetry of the evoked muscle responses was confirmed prior to pooling the data (see Statistical analysis). This approach, however, was not possible for coherence and gain between EVS and GRFs since in our narrow-base walking condition because participants regularly made contact of one limb with the opposing belt such that forces from each limb could not be measured on the separate force plates. Therefore, forces from both plates were first summed before estimating the coherence. To avoid distortion in the coherence estimates at the beginning and end of the signal, each stride was padded with data from the neighboring strides (50%). EVS and rectified EMG were low-pass filtered (100 Hz cutoff, zero lag, sixth-order Butterworth) and down-sampled to 200 samples/s. To account for stride-to-stride variation, stride duration was normalized by resampling the data according to the average stride duration of all trials. This normalization was performed on the auto-spectra of the EVS, EMG and force signals, as well as on their cross-spectra (see below).

Our analysis of coherence and gain was performed based on continuous Morlet wavelet decomposition [15,70] using equations (1) and (2):

$$252 \qquad C(\tau,f) = \frac{|P_{xy}(\tau,f)|^2}{P_{xx}(\tau,f)P_{yy}(\tau,f)} \qquad (1)$$

$$253 \qquad G(\tau,f) = \left|\frac{P_{xy}(\tau,f)}{P_{xx}(\tau,f)}\right| \qquad (2)$$

where $Pxy(\tau,f)$ (τ and f denote the stride time and frequency, respectively) is the time-
dependent cross-spectrum between the EVS and rectified EMG or GRF, and $Pxx(\tau,f)$ and $Pyy(\tau,f)$
are the time-dependent auto-spectra of the EVS and rectified EMG or GRF, respectively. Coherence
ranges from zero to one, and provides a measure of the linear relationship between two signals [71]
as well as an estimate of their coupling (i.e. shared variance) at each frequency [72]. Because
coherence is normalized to the auto-spectra of the input and output signals, it can be sensitive to
variation in non-vestibular contributions to the measured output signal (i.e. magnitude changes of
EMG or ground reaction forces). Gain on the other hand indicates the magnitude of the output
relative to the input and is not normalized by the output signal power spectrum; as a result, it is not
expected to change even if non-vestibular input leads to changes in output signal magnitude. A
preliminary evaluation of our results indicated that similar to previous studies [12,15], both coherence
and gain followed parallel changes in magnitude and timing across the four walking conditions. This
suggests that the modulation in coherence was not simply dependent upon the magnitude of the
measured output signals, and as a result, we present only the coherence to describe the modulation
of vestibular contributions to mediolateral stability. Finally, we evaluated the coherence between
EVS and both EMG and GRFs to estimate muscle-specific vestibular contributions and to provide a
net estimate of the vestibular input to ongoing locomotor behavior, respectively.

Statistical analysis

We compared all gait parameters (stride time, step width and local dynamic stability) between
the four conditions using one-way repeated measures ANOVAs. Subsequently, we performed
planned pairwise comparisons (t-test) between normal walking condition and each modified walking
condition (i.e. normal vs. stabilized; normal vs. narrow and normal vs. wide-base walking). EVS-EMG
coherence and EVS-GRF coherence were used to identify the phase-dependent coupling between
vestibular stimulation and motor responses throughout the walking cycle. For each participant,
coherence was defined as significant for those points in the gait cycle where it exceeded 0.018,
corresponding to $p < 0.01$ (for 256 strides) in view of the bi-dimensional nature of the correlations [15].
To determine whether our walking manipulations modified the EVS-EMG coherence when compared
to normal walking, we performed cluster-based permutation tests (paired t-tests, 5000 permutations)
[73] between conditions aimed to identify whether the time-frequency-coherence spectra significantly
differed from the normal condition. In doing so, we did not disregard non-significant coherence
values. Since our initial analysis showed no significant between leg differences in coherence, we
averaged coherence values over legs.

Results

Effects of condition on gait parameters, muscle activity and local dynamic stability

To characterize changes in gait across walking trials, we first evaluated the gait parameters, muscle activity and stability measures. During all trials, participants were able to maintain stable upright locomotion while exposed to the stochastic electrical stimulation. Despite walking to the beat of a metronome in all conditions, we found a significant main effect of condition on stride time ($F_{(3,30)}=4.14$; $p=0.014$; $\eta^2=0.127$). Pairwise comparisons revealed that stride time increased by approximately 6.12±15.65% (0.09±0.02 s; $p=0.018$) during stabilized walking when compared to normal walking (Figure 1a), but it did not change significantly during either wide-base ($p=0.928$) or narrow-base walking ($p=0.155$). As intended, walking condition significantly affected step width ($F_{(3,30)}=73.4$; $p<0.001$; $\eta^2=0.786$, see Figure 1b). Consistent with previous results [20,23-25], pairwise analysis revealed that participants reduced their step width by 65.31±40% (0.18±0.02 m; $p<0.001$) during stabilized walking, in spite of no explicit instruction to do so. We also found that participants adhered to the step width instructions during the other two conditions, increasing step width by 30.25±20% (0.08±0.02 m; $p<0.001$) during wide-base walking and reducing step width by 38.74±5% (0.11±0.01 m; $p<0.001$) during narrow-base walking (Figure 1b).

Figure 1: Spatiotemporal parameters of gait during normal walking (blue), walking with external lateral stabilization (green), wide-base walking (black) and narrow-base walking (red). Mean values (bars), standard deviation (error bar) and participants individual changes across conditions (lines) for the stride time (a) and stride width (b) for the four conditions.

As expected, when compared to normal walking, significant differences in mediolateral ground reaction forces [30,31,74] during all other walking conditions were observed primarily during single-support phases of the gait cycle (see Figure 2a). Forces decreased when subjects were externally supported or walked with a narrow-base, and increased when subjects walked with a wide-base (see Supplementary Figure S1). In the medial gastrocnemius and gluteus medius muscles, peak activity in each muscle occurred during the double stance phase (~50% of the stride cycle) and after heel strike (~20% of stride cycle), respectively. EMG responses for these two muscles were for the most part overlapping in all conditions, with only limited significant differences observed during the stance phase for some of the conditions relative to normal walking (see Figure 2b/c and

Supplementary Figure S1). Muscle activity in the erector spinae demonstrated more complex phasic activity throughout the gait cycle with two peaks occurring just after heel strike of each limb. Limited significant differences were observed during stabilized walking and narrow-base walking when compared to normal walking (see Figure 2d and Supplementary Figure S1).

Figure 2: Ground reaction force and electromyographic amplitudes throughout the gait cycle during normal walking (blue line), walking with external lateral stabilization (green line), wide-base walking (black line) and narrow-base walking (red line). Mean values (lines) and standard deviation (shadow area) of the ground reaction force (a) and the EMG envelopes of the medial gastrocnemius (b), gluteus medius (c) and erector spinae (d) muscles for all conditions. For each muscle, EMG signals were normalized to the maximum amplitude across all conditions.

Our manipulations also had a significant effect on mediolateral stability, as expressed by the
local divergence exponent ($F_{(3,30)}$=129; $p<0.001$; η^2=0.712). Pairwise comparisons showed that
participants had a higher stability (i.e., lower LDE values) during the stabilized ($p<0.001$) and narrow-
base walking conditions (p=0.019) compared to normal walking. Although the mean local divergence
exponent was highest (i.e. lowest stability) during wide-base walking, this was not significantly
different from normal walking (p=0.265) (Figure 3).

Figure 3: Local divergence exponents during normal walking (blue), walking with external
stabilization (green), wide-base walking (black) and narrow-base walking (red). Mean values (bars),
standard deviation (error bar) and participants individual changes across conditions (lines) for the
evaluated conditions.

EVS-mediolateral GRF and EVS-EMG coherence in normal walking

We next characterized the coupling of the electrical stimulus with both the ground reaction
forces and muscle activity in normal walking as a baseline for comparison to our manipulated
conditions (Figure 4 – first column). During normal walking, significant EVS-GRF coherence was
seen in all participants over the entire gait cycle, with phase-dependent group mean responses that
peaked during single stance (Figure 4 top row). Significant phase-dependent EVS-EMG coupling
was also prominent in the mean responses during normal walking, but muscle-specific variations
were observed: coherence peaked in mid stance in the medial gastrocnemius, just before heel strike
in gluteus medius, and at heel strike and at mid stride (though at a lower magnitude) for the erector
spinae. Consistent with previous reports [11,12,15], peak coherences did not align with peak EMG for
any of the muscles, further confirming that vestibular contributions do not depend purely on the
excitation of the motoneuron pool. In addition, as commonly observed in vestibular-evoked muscles
responses during standing [75], the bandwidth of significant EVS-EMG coherence spanned ~0-25 Hz
while significant EVS-GRF coherence was observed from ~0-10 Hz (Figure 4).

Figure 4: Coherence plots of EVS-GRF (first row) and EVS-EMG for medial gastrocnemius (second row), gluteus medius (third row) and erector spinae (fourth row) for normal walking (first column), walking with external stabilization (second column), wide-base walking (third column) and narrow-base walking (fourth column). Coherence magnitude is indicated by the color bars.

Stabilization demands but not dynamic stability modulate EVS-GRF and EMG-EVS coupling

To establish the effects of stabilization demands on vestibulomotor coupling, we examined the difference in coherence between normal walking and all other conditions (Figure 5). During externally stabilized walking, both EVS-GRF and EVS-EMG coherences decreased significantly relative to normal walking (see Figure 4 and 5). More specifically, the reduced coupling during the stabilized condition was primarily observed during the periods of peak coherence in normal walking (i.e. single stance for GRF and medial gastrocnemius, and before/at heel strike for gluteus medius and erector spinae). Although the increased stride time (i.e. decreased cadence) during stabilized walking (see Figure 1) may have acted as a confounding factor to these changes [11], this effect commonly increases vestibulomotor responses in contrast to the observed decrease in coherence seen here. During wide-base walking, EVS-GRF and EVS-EMG coherences were also significantly decreased compared to normal walking (Figure 5). The EVS-GRF coherence decreased over the majority of the gait cycle with the most prominent changes observed during the periods of peak

coherence seen in the normal condition. Further, EVS-EMG coherence was reduced for all three
muscles in wide-base walking, again with the greatest differences observed at instants of peak
coherence in normal walking (Figure 5). Taken together, the results of stabilized and wide-base
walking show a reduction in vestibular input to the net muscle activity of the body (i.e. GRFs), which
is driven at least in part by the three muscles measured when stabilization demands (but not dynamic
stability) are decreased.

During narrow-base walking, we observed more complex changes in coupling between EVS
and GRF and between EVS and muscle activity. Figures 4 and 5 together show that EVS-GRF
coherence during narrow-base walking significantly increased compared to normal walking over the
entire gait cycle. While this was matched by an increase in EVS-EMG coherence in the erector
spinae muscle, we found unchanged EVS-EMG coupling in the gluteus medius muscle and a
decrease in coherence in the medial gastrocnemius muscle. These more complex changes in
vestibular-evoked motor responses suggest that while the net output of the vestibular-evoked muscle
activity (i.e. EVS-GRF coherence) increases with increased stabilization demands, as well as the
dynamic stability (see Figure 3), this trend is not reflected in EVS-EMG coupling of all muscles.

Figure 5: Differences in time-frequency coherence between the normal walking condition and
walking with external lateral stabilization (first column), wide-base walking (second column) and

narrow-base walking (third column). Color coding refers to the difference in coherence between two

walking conditions (e.g. normal minus stabilized) for the EVS-GRF coherence (first row) and EVS-

EMG coherence of medial gastrocnemius (second row), gluteus medius (third row) and erector

spinae (forth row) muscles. For illustrative purposes, differences that were not significant were

plotted slightly opaque.

Discussion

We characterized how coupling of vestibular input with muscle activity and ground reaction

forces modulate as a function of the stabilization demands during locomotion. We found that as

participants walked with decreased stabilization demands through either external stabilization or

wider step widths, coherence between electrical vestibular stimulation and both muscle activity and

ground reaction forces decreased compared to normal walking. These overall reductions in

vestibulomotor coupling were accompanied by an increase or no change in the stability of the gait

pattern – measured as a decreased or constant local divergence exponent – during stabilized and

wide-base walking, respectively. In contrast, the increased stabilization demands of walking with

narrow steps invoked complex changes in vestibulo-muscular coupling that increased or decreased

specific to each muscle's involvement in correcting for the imposed vestibular error. Nevertheless,

these changes in vestibulo-muscular coupling increased the collective contribution of vestibular

signals to the ground reaction forces and occurred together with a decrease in the local divergence

exponent (i.e. increased gait stability). This suggests that participants maintained a more stable gait

pattern during narrow walking that was at least partially subserved through increased use of

vestibular feedback. Ultimately, these results indicate that vestibular contributions to gait stability

may be modulated with frontal plane stability, but that they more specifically depend on the

stabilization demands (i.e. control effort) required to maintain a stable gait pattern and not the stability

of the gait pattern itself.

When participants walked with external stabilization, the stability of the gait pattern increased

(i.e. decreasing LDE) while vestibular-evoked muscle and force responses decreased as compared

to normal walking. Both of these results are not entirely surprising since the control of mediolateral

motion is aided by the forces generated by the springs [19,20,25]. As a result, there is a reduced reliance

on vestibular signals to maintain upright locomotion during stabilized walking. This is similar to the

task dependent reductions in vestibular input observed during standing [44,45,76-78] when participants

are externally supported; stimulus-evoked responses are suppressed since the vestibular feedback

is no longer relevant to balancing the body. Our results reveal that these task dependent changes in

the vestibular control of standing also apply during the more dynamic task of walking. In addition,

they also support the proposal that anteroposterior control of whole-body stability during locomotion

is controlled passively [79]. By making the body passively stable in the mediolateral direction, we saw

a reduction in vestibular-evoked response that matched the near absence of vestibular contributions

when the vestibular error is directed in the anterior-posterior direction [16].

When participants walked with a wide base, vestibular-evoked muscle and force responses also decreased in a manner that parallels the effects of wide stance during standing [44,80]. Walking (and standing) with a wide foot placement increases the base of support, and the passive stiffness in the frontal plane. The current results show that the corrective contribution of vestibular signals during walking with a wide base decrease in a manner similar to the effects seen during external stabilization. Our measure of dynamic stability, however, did not follow the same trend. Instead, we observed a slight (albeit non-significant) increase in the local divergence exponent compared to normal walking. This aligns with previous estimates of a constant or decreased dynamic gait stability during wide-base walking [30,34]. A key difference between stabilized and wide-base walking is that in the former, increased gait stability and upright balance is an inevitable result of the external support. Wide-base walking, on the other hand, despite the increased base of support, still demands active stabilization, and the manner in which this is achieved differs from normal walking. For example, push-off modulation and step-by-step foot placement precision both decrease when walking with wide steps [30]. In addition, the increased moment arm of the ground reaction forces about the center of mass generates greater fluctuations in angular plane momentum [81], which in and of themselves are destabilizing. These changes may be possible because the margins for error of balance control are inherently increased, making wide-base walking more robust to external disturbances. The current results therefore suggest that vestibular contributions may also decrease under walking conditions with reduced control effort to maintain mediolateral stability. This is in line with recent findings in patients with vestibular hypofunction who walk slower, with increased cadence and wider steps [82-86]. Our results suggest that they may adopt these changes in walking behavior to be less dependent on vestibular input.

The possibility that vestibular contributions are modulated with the control effort (i.e. stabilization demand) of gait is further supported by our narrow-base walking results. Both the dynamic stability of gait and vestibular-evoked ground reaction forces during narrow-base walking increased relative to normal walking. The increase in dynamic stability (i.e. decreasing LDE) in particular is thought to be required to constrain torso motion to margins of error within which walking with narrow steps can be maintained [34]. Indeed, foot placement during narrow-base walking is more tightly coupled to center-of-mass motion [30,31] and participants maintain smaller step-to-step oscillations [87]. Our results show that under these highly regulated conditions, this increase in control effort may be driven, at least in part, by a net increase in vestibular input (as reflected in the ground reaction forces). This overall increase in vestibular contribution to mediolateral stability, however, does not seem to originate from a general upregulation of vestibular responses in all muscles. Instead, we found decreasing and unchanging vestibular responses in medial gastrocnemius and gluteus medius muscles, respectively, while vestibular responses in erector spinae muscles increased. These muscle specific changes align with the scaling of evoked responses according to the muscle's involvement in correcting for imposed errors [11,26]. For example, moments generated by the ankle muscles, such as the medial gastrocnemius, strongly contribute to balance corrections

through push-off modulation [88]. During narrow-base walking, however, push-off modulation is not a viable option since the push off force has only a minimal moment arm. Similarly, regulation of mediolateral foot placement as provided by the gluteus medius [11,57] is rendered less effective, since narrow-base walking constrains the foot to a restricted range. Instead, humans more commonly rely on the modulation of the torso's angular moment to maintain upright balance [61,87]. Although the muscles around the ankles and hips also contribute to frontal plane angular momentum during normal walking [81], the direct influence of spinal muscles on trunk motion may make them more suited to contribute to balance during narrow walking by producing or responding to rapid trunk tilts. A limitation to this interpretation, however, is that we cannot rule out the influence of arm movements, which are known to influence the stability of human gait [89-91], since we did not measure arm movements or provide specific instructions to participants to control their movement. Nevertheless, our erector spinae results indicate that spinal muscles provide a key, and perhaps primary, contribution to the net vestibular output to ground reaction forces to maintain upright balance during narrow-base walking.

Significant coherence between the electrical stimulus and erector spinae muscle activity in all conditions also demonstrates that the phasic contribution of this particular axial muscle to whole-body mediolateral stability can be flexibly modulated to address varying stabilizing demands. This is similar to this muscle's differential response to vestibular disturbances in standing and sitting [92]. Others have reported, however, that erector spinae muscles, unlike lower limb muscles, maintain a fixed response sensitivity to vestibular input throughout all phases of walking [93]. This follows a similar argument made in neck muscles where vestibulocollic reflexes are maintained regardless of the requirement to maintain head on trunk balance [94,95]. By delivering square wave EVS pulses at and slightly after (15% of the gait cycle) the heel strike, Guillaud et al. (2020) argued that the invariant response of the muscle to the electrical stimulation at these two time points indicate a fixed vestibular sensitivity throughout the entire gait cycle [93]. The improved resolution of the time-varying techniques used here, however, reveals that this muscle does indeed modulate its response to vestibular input throughout locomotion: EVS-EMG coherence peaked at heel strike and dropped to near-zero at approximately 40% of the stride cycle. The restricted time-window of the two stimulation occurrences considered (i.e. at and slightly after heel strike) in the study from Guillaud et al.(2020), however, may have masked any changes in EVS-evoked responses. In addition, the modulation of this muscle's vestibular sensitivity extends across walking conditions, which nearly doubled during narrow-base walking as compared to normal walking.

A limitation of our study is that while vestibular-evoked responses were quantified throughout the gait cycle, the local divergence exponent remains a mean measure of stability for the entire gait cycle. It may therefore be possible that time-varying changes in gait stability contribute to the phase-dependent vestibular responses seen here. While early studies using phase dependent metrics of gait stability suggested that these measures may hold promise [96,97], a recent study from our lab doubted their use, as we found only limited correlation between phase dependent gait stability

measures and the probability of falling in a simple dynamic walking model [98]. However, if sufficient phase dependent gait stability measures become available, the relationship between phase dependent gait stability and phase-dependent vestibular responses may be an interesting study topic.

In conclusion, we have shown that the muscle and whole-body responses evoked by a vestibular stimulation differ according to the gait stabilization demands. When stability is increased by external support, the muscle and whole-body responses to the vestibular stimulus are substantially reduced. During wide-base walking vestibular-evoked muscle and force responses also decrease, though these changes in vestibular contribution are not accompanied by increased dynamic gait stability. Conversely, narrow-base walking produced complex muscle-specific responses that resulted in an increase in the net vestibular contribution to ground reaction forces and increased stability of gait. Overall, our results show that although the vestibular control of gait stability may vary with frontal plane stability, they critically depend on the stabilization demands (i.e. control effort) needed to maintain stable walking patterns.

Acknowledgements:

RMM was funded by CAPES (PDSE 19/2016). SMB was funded by a VIDI grant (016.Vidi.178.014) from the Netherlands Organization for Scientific Research (NWO). PAF received funding from the Netherlands Organization for Scientific Research (NWO #016.Veni.188.049).

Competing Interests Statement:

The authors have no conflict of interest to declare.

Author contribution statement:

JVD and PAF contributed to the conception or design of the work. RMM, SMB and PAF contributed to the acquisition, analysis, or interpretation of data. SMB and PAF made the figures. All authors wrote the main manuscript text, reviewed the manuscript and have approved the submitted version.

References

1. Bruijn, S. M., Meijer, O. G., Beek, P. J. & van Dieen, J. H. Assessing the stability of human locomotion: a review of current measures. *J R Soc Interface* **10**, 20120999, doi:10.1098/rsif.2012.0999 (2013).
2. van Emmerik, R. E. A., Ducharme, S. W., Amado, A. C. & Hamill, J. Comparing dynamical systems concepts and techniques for biomechanical analysis. *Journal of Sport and Health Science* **5**, 3-13, doi:10.1016/j.jshs.2016.01.013 (2016).
3. Mehdizadeh, S. The largest Lyapunov exponent of gait in young and elderly individuals: A systematic review. *Gait Posture* **60**, 241-250, doi:10.1016/j.gaitpost.2017.12.016 (2018).
4. Misiaszek, J. Neural control of walking balance: if falling then react else continue. *Exercise and Sport Sciences Review* **34**, 128-134 (2006).

5   Chien, J. H., Mukherjee, M. & Stergiou, N. Mastoid Vibration Affects Dynamic Postural
Control During Gait. *Ann Biomed Eng* **44**, 2774-2784, doi:10.1007/s10439-016-1556-z
(2016).
6   Stergiou, N. & Decker, L. M. Human movement variability, nonlinear dynamics, and
pathology: is there a connection? *Hum Mov Sci* **30**, 869-888,
doi:10.1016/j.humov.2011.06.002 (2011).
7   Rossignol, S., Dubuc, R. & Gossard, J. P. Dynamic sensorimotor interactions in locomotion.
Physiol. Rev. **86**, 89-154, doi:10.1152/physrev.00028.2005 (2006).
8   Cullen, K. E. The vestibular system: multimodal integration and encoding of self-motion for
motor control. *Trends Neurosci* **35**, 185-196, doi:10.1016/j.tins.2011.12.001 (2012).
9   Brandt, T. Vestibulopathic gait: you're better off running than walking. *Curr Opin Neurol* **13**,
3-5 (2000).
10   Borel, L. *et al.* Walking performance of vestibular-defective patients before and after
unilateral vestibular neurotomy. *Behave Brain Res* **150**, 191-200 (2004).
11   Dakin, C. J., Inglis, J. T., Chua, R. & Blouin, J. S. Muscle-specific modulation of vestibular
reflexes with increased locomotor velocity and cadence. *Journal of neurophysiology* **110**,
86-94, doi:10.1152/jn.00843.2012 (2013).
12   Forbes, P. A. *et al.* Rapid limb-specific modulation of vestibular contributions to ankle
muscle activity during locomotion. *J Physiol* **595**, 2175-2195, doi:10.1113/JP272614 (2017).
13   Bent, L. R., Inglis, J. T., bradford, J. & McFadyen, B. J. When is Vestibular Information
Important During Walking? *J Neurophysiol* **92**, 1269-1275 (2004).
14   Iles, J. F., Baderin, R., Tanner, R. & Simon, A. Human standing and walking: comparison of
the effects of stimulation of the vestibular system. *Exp. Brain Res.* **178**, 151-166,
doi:10.1007/s00221-006-0721-2 (2007).
15   Blouin, J. S. *et al.* Extracting phase-dependent human vestibular reflexes during locomotion
using both time and frequency correlation approaches. *J Appl Physiol (1985)* **111**, 1484-
1490, doi:10.1152/japplphysiol.00621.2011 (2011).
16   Tisserand, R. *et al.* Down regulation of vestibular balance stabilising mechanisms to enable
transition between motor states. *eLife* **7**, doi:https://doi.org/10.7554/eLife.36123
(2018).
17   Bauby, C. E. & Kuo, A. D. Active control of lateral balance in human walking *Journal of
Biomechanics* **33**, 1433-1440 (2000).
18   McGeer, T. Passive Dynamic Walking. *The International Journal of Robotics Research T* **9**,
62-82, doi:https://doi.org/10.1177/027836499000900206 (1990).
19   Mahaki, M., Bruijn, S. M. & van Dieen, J. H. The effect of external lateral stabilization on the
use of foot placement to control mediolateral stability in walking and running. *PeerJ* **7**,
e7939, doi:10.7717/peerj.7939 (2019).
20   Donelan, J. M., Shipman, D. W., Kram, R. & Kuo, A. D. Mechanical and metabolic
requirements for active lateral stabilization in human walking. *Journal of biomechanics* **37**,
827-835, doi:10.1016/j.jbiomech.2003.06.002 (2004).
21   Ortega, J. D., Fehlman, L. A. & Farley, C. T. Effects of aging and arm swing on the metabolic
cost of stability in human walking. *J Biomech* **41**, 3303-3308,
doi:10.1016/j.jbiomech.2008.06.039 (2008).
22   Dean, J. C., Alexander, N. B. & Kuo, A. D. The effect of lateral stabilization on walking in
young and old adults. *IEEE Trans Biomed Eng* **54**, 1919-1926 (2007).
23   Ijmker, T., Houdijk, H., Lamoth, C. J., Beek, P. J. & van der Woude, L. H. Energy cost of
balance control during walking decreases with external stabilizer stiffness independent of
walking speed. *Journal of biomechanics* **46**, 2109-2114,
doi:10.1016/j.jbiomech.2013.07.005 (2013).

24 Ijmker, T. *et al.* Can external lateral stabilization reduce the energy cost of walking in
persons with a lower limb amputation? *Gait & posture* **40**, 616-621,
doi:10.1016/j.gaitpost.2014.07.013 (2014).
25 Bruijn, S. M., Van Dieen, J. H. & Daffertshofer, A. Beta activity in the premotor cortex is
increased during stabilized as compared to normal walking. *Front Hum Neurosci* **9**, 593,
doi:10.3389/fnhum.2015.00593 (2015).
26 Forbes, P. A., Luu, B. L. & Blouin, J. S. Transformation of vestibular signals for the control of
standing in humans. *The Journal of Neuroscience* **36**, 11510-11520,
doi:10.1523/JNEUROSCI.1902-16.2016 (2016).
27 Kubinski, S. N., McQueen, C. A., Sittloh, K. A. & Dean, J. C. Walking with wider steps
increases stance phase gluteus medius activity. *Gait & Posture* **41**, 130-135,
doi:http://dx.doi.org/10.1016/j.gaitpost.2014.09.013 (2015).
28 Aboutorabi, A., Arazpour, M., Bahramizadeh, M. & W., H. S. The effect of aging on gait
parameters in able-bodied older subjects: a literature review. . 2015; doi:. *Aging Clin Exp*
Res **28**, 393–405, doi:s40520-015-0420-6 (2016).
29 Schrager, M. A., Kelly, V. E., Price, R., Ferrucci, L. & Shumway-Cook, A. The effects of age on
medio-lateral stability during normal and narrow base walking. *Gait Posture* **28**, 466-471,
doi:10.1016/j.gaitpost.2008.02.009 (2008).
30 Perry, J. A. & Srinivasan, M. Walking with wider steps changes foot placement control,
increases kinematic variability and does not improve linear stability. *R Soc Open Sci* **4**,
160627, doi:10.1098/rsos.160627 (2017).
31 Arvin, M. *et al.* Effects of narrow base gait on mediolateral balance control in young and
older adults. *Journal of biomechanics* **49**, 1264-1267, doi:10.1016/j.jbiomech.2016.03.011
(2016).
32 Hak, L. *et al.* Speeding up or slowing down?: Gait adaptations to preserve gait stability in
response to balance perturbations. *Gait Posture* **36**, 260-264,
doi:10.1016/j.gaitpost.2012.03.005 (2012).
33 Young, P. M. M. & Dingwell, J. B. Voluntary changes in step width and step length during
human walking affect dynamic margins of stability. *Gait & Posture* **36**, 219-224,
doi:10.1016/j.gaitpost.2012.02.020 (2012).
34 Young, P. M. M. & Dingwell, J. B. Voluntarily changing step length or step width affects
dynamic stability of human walking. *Gait Posture* **35**, 472-477,
doi:10.1016/j.gaitpost.2011.11.010 (2012).
35 Murray, A. J., Croce, K., Belton, T., Akay, T. & Jessell, T. M. Balance control mediated by
vestibular circuits directing lim extension or antagonist muscle co-activation. *Cell Reports*
22, 1325-1338, doi:https://doi.org/10.1016/j.celrep.2018.01.009 (2018).
36 Goldberg, J. M., Fernandez, C. & Smith, C. E. Responses of vestibular-nerve afferents in the
squirrel monkey to externally applied galvanic currents. *Brain Research* **252**, 156-160
(1982).
37 Kim, J. & Curthoys, I. S. Responses of primary vestibular neurons to galvanic vestibular
stimulation (GVS) in the anaesthetised guinea pig. *Brain Research* **64**, 265-271,
doi:https://doi.org/10.1016/j.brainresbull.2004.07.008 (2004).
38 Kwan, A., Forbes, P. A., Mitchell, D. E., Blouin, J. S. & Cullen, K. E. Neural substrates,
dynamics and thresholds of galvanic vestibular stimulation in the behaving primate. *Nat*
Commun **10**, 1904, doi:10.1038/s41467-019-09738-1 (2019).
39 Peters, R. M., Rasman, B. G., Inglis, J. T. & Blouin, J. S. Gain and phase of perceived virtual
rotation evoked by electrical vestibular stimuli. *J Neurophysiol* **114**, 264-273,
doi:10.1152/jn.00114.2015 (2015).

40 Khosravi-Hashemi, N., Forbes, P. A., Dakin, C. J. & Blouin, J. S. Virtual signals of head
rotation induce gravity-dependent inferences of linear acceleration. *J. Physiol.* **597**, 5231-
5246, doi:10.1113/jp278642 (2019).

41 Fitzpatrick, R. C. & Day, B. L. Probing the human vestibular system with galvanic
stimulation. *J Appl Physiol (1985)* **96**, 2301-2316 (2004).

42 St George, R. J. & Fitzpatrick, R. C. The sense of self-motion, orientation and balance
explored by vestibular stimulation. *The Journal of Physiology* **589**, 807-813,
doi:10.1113/jphysiol.2010.197665 (2011).

43 Nashner, L. M. & Wolfson, P. Influence of head position and proprioceptive cues on short
latency postural reflexes evoked by galvanic stimulation of the human labyrinth. *Brain Res*
67, 255-268, doi:10.1016/0006-8993(74)90276-5 (1974).

44 Mian, O. S. & Day, B. L. Violation of the craniocentricity principle for vestibularly evoked
balance responses under conditions of anisotropic stability. *J Neurosci* **34**, 7696-7703,
doi:10.1523/JNEUROSCI.0733-14.2014 (2014).

45 Britton, T. C. *et al.* Postural electromyographic responses in the arm and leg following
galvanic vestibular stimulation in man. *Exp Brain Res* **94**, 143-151, doi:10.1007/BF00230477
(1993).

46 Lund, S. & Broberg, C. Effects of different head positions on postural sway in man induced
by a reproducible vestibular error signal. *Acta Physiol Scand* **117**, 307-309,
doi:10.1111/j.1748-1716.1983.tb07212.x (1983).

47 Dakin, C. J., Luu, B. L., van den Doel, K., Inglis, J. T. & Blouin, J. S. Frequency-specific
modulation of vestibular-evoked sway responses in humans. *J Neurophysiol* **103**, 1048-
1056, doi:10.1152/jn.00881.2009 (2010).

48 Mahaki, M., Bruijn, S. M. & van Dieën, J. H. The effect of external lateral stabilization on the
control of mediolateral stability in walking and running. . *PeerJ Preprints* **6:e27244v1**
doi:https://doi.org/10.7287/peerj.preprints.27244v1 (2018).

49 Mahaki, M., IJmker, T., Houdijk, H. & Bruijn, S. M. How does external lateral stabilization
constrain normal gait, apart from improving medio-lateral gait stability? *bioRxiv* (2020).

50 Matsas, A., Taylor, N. & McBurney, H. Knee joint kinematics from familiarised treadmill
walking can be generalised to overground walking in young unimpaired subjects. . *Gait &*
Posture **11**, 46-53, doi:10.1016/s0966-6362(99)00048-x (2000).

51 Meyer, C. *et al.* Familiarization with treadmill walking: How much is enough? *Sci Rep* **9**,
5232, doi:https://doi.org/10.1038/s41598-019-41721-0 (2019).

52 Hannan, K. B., Todd, M. K., Pearson, N. J., Forbes, P. A. & Dakin, C. J. Vestibular attenuation
to random - waveform galvanic vestibular stimulation during standing and treadmill
walking. *Scientific Reports* **11**, doi:https://doi.org/10.1038/s41598-021-87485-4 (2021
).

53 Cathers, I., Day, B. L. & Fitzpatrick, R. C. Otolith and canal reflexes in human standing.
Journal of Physiology **563**, 229-234, doi:10.1113/jphysiol.2004.079525 (2005).

54 Fitzpatrick, R. C., Butler, J. E. & Day, B. L. Resolving head rotation for human bipedalism.
Curr Biol **16**, 1509-1514, doi:10.1016/j.cub.2006.05.063 (2006).

55 Hermens, H. J. *et al. SENIAM 8: European recommendations for surface electromyography.*,
Vol. 1 (Roessingh Researchand Development, 1999).

56 Liu, M. Q., Anderson, F. C., Schwartz, M. H. & Delp, S. L. Muscle contributions to support
and progression over a range of walking speeds. *Journal of biomechanics* **41**, 3243-3252,
doi:10.1016/j.jbiomech.2008.07.031 (2008).

57 Rankin, B. L., Buffo, S. K. & Dean, J. C. A neuromechanical strategy for mediolateral foot
placement in walking humans. *Journal of neurophysiology* **112**, 374-383,
doi:10.1152/jn.00138.2014 (2014).

58   van Leeuwen, A. M., van Dieën, J. H., Daffertshofer, A. & Bruijn, S. M. Step width and
frequency to modulate: Active foot placement control ensures stable gait. *bioRxiv* (2020).

59   Waters, R. L. & Morris, J. M. Electrical activity of muscles of the trunk during walking. *J*
Anat **111**, 191-199 (1972).

60   Thorstensson, A., Carlson, H., Zomlefer, M. R. & Nilsson, J. Lumbar back muscle activity in
relation to trunk movements during locomotion in man. *Acta Physiol Scand* **116**, 13-20,
doi:10.1111/j.1748-1716.1982.tb10593.x (1982).

61   Best, A. N. & Wu, A. R. Upper body and ankle strategies compensate for reduced lateral
stability at very slow walking speeds. *bioRxiv* (2020).

62   Roerdink, M. & al, e. Online gait event detection using a large force platform embedded in
a treadmill. . *Journal of biomechanics* **41**, 2628-2632 (2008).

63   Dingwell, J. B. & Cusumano, J. P. Nonlinear time series analysis of normal and pathological
human walking. *Chaos* **10**, 848-863, doi:10.1063/1.1324008 (2000).

64   Dingwell, J. B. & Marin, L. C. Kinematic variability and local dynamic stability of upper body
motions when walking at different speeds. *J Biomech* **39**, 444-452,
doi:10.1016/j.jbiomech.2004.12.014 (2006).

65   Rosenstein, M. T., Coliins, J. J. & De Luca, C. J. A practical method for calculating largest
Lyapunov exponents from small data sets. *Physica* **65**, 117-134 (1993).

66   Kang, H. G. & Dingwell, J. B. Dynamic stability of superior vs. inferior segments during
walking in young and older adults. *Gait & Posture* **30**, 260-263 (2009).

67   Stenum, J., Bruijn, S. M. & Jensen, B. R. The effect of walking speed on local dynamic
stability is sensitive to calculation methods. *Journal of biomechanics* **47**, 3776-3779,
doi:10.1016/j.jbiomech.2014.09.020 (2014).

68   Local Dynamic Stability: first release v. 1 (Zenodo, Zenodo, 2017).

69   Forbes, P. A. *et al.* Electrical vestibular stimuli to enhance vestibulo-motor output and
improve subject comfort. *PLoS One* **9**, e84385, doi:10.1371/journal.pone.0084385 (2014).

70   Zhan, Y., Halliday, D. W., Jiang, P., Liu, X. & Feng, J. Detecting time- dependent coherence
between non-stationary electrophysiological signals. A combined statistical and time-
frequency approach. *J Neurosci Methods* **156**, 322-332 (2006).

71   Pintelon, R. & Schoukens, J. *System identification: a frequency domain approach*. 2nd
edition edn, (Hoboken, N.J.: John Wiley & Sons Inc., 2012).

72   Warner, R. M. *Spectral Analysis of Time-Series Data (Methodology in the Social Sciences)*. 1
edition edn, (The Guilford Press, 1998).

73   Maris, E., Schoffelen, J. M. & Fries, P. Nonparametric statistical testing of coherence
differences. *Journal of neuroscience methods* **163**, 161-175,
doi:10.1016/j.jneumeth.2007.02.011 (2007).

74   Hof, A. L., van Bockel, R. M., Schoppen, T. & Postema, K. Control of lateral balance in
walking. Experimental findings in normal subjects and above-knee amputees. *Gait &*
Posture **25**, 250-258 (2007).

75   Dakin, C. J., Son, G. M., Inglis, J. T. & Blouin, J. S. Frequency response of human vestibular
reflexes characterized by stochastic stimuli. *J Physiol* **583**, 1117-1127,
doi:10.1113/jphysiol.2007.133264 (2007).

76   Fitzpatrick, R. C. & McCloskey, D. I. Proprioceptive, visual and vestibular thresholds for the
perception ofsway during standing in humans. *Journal of Physiology* **478**, 173-177 (1994).

77   Forbes, P. A., Siegmund , G. P., Schouten, A. C. & Blouin, J.-S. b. Task, muscle and
frequency dependent vestibular control of posture. *Frontiers in Integrative Neuroscience* **8**,
doi:10.3389/fnint.2014.00094 (2015).

78   Luu, B. L. *et al.* Human standing is modified by an unconscious integration of congruent
sensory and motor signals. *Journal of physiology* **590**, 5783–5794, doi:10.
1113/jphysiol.2012.230334 (2012).
79   Bauby, C. E. & Kuo, A. D. Active control of lateral balance in human walking. *Journal of*
biomechanics **33**, 1433-1440 (2000).
80   Day, B. L., Cauquil, A. S., Bartolomei, M., Pastor, M. A. & I.N., L. Human body-
segmenttiltsinducedbygalvanicstimulation:
avestibularlydrivenbalanceprotectionmechanism. *Journal of Physiology* **500**, 661-672
(1997).
81   **Neptune, R. R.** & **McGowan, C. P.** Muscle Contributions to Frontal Plane Angular
Momentum during Walking. *J Biomechanics* **49**, 2975-2981,
doi:doi:10.1016/j.jbiomech.2016.07.016
(2016).
82   Liu, P. *et al.* Characterizing Patients with Unilateral Vestibular Hypofunction Using
Kinematic Variability and Local Dynamic Stability during Treadmill Walking. *Behav Neurol*
2017, 4820428, doi:10.1155/2017/4820428 (2017).
83   Schniepp, R. *et al.* Locomotion speed determines gait variability in cerebellar ataxia and
vestibular failure. *Movement Disorders* **27**, 125-131, doi:10.1002/mds.23978 (2012).
84   Gimmon, Y., Millar, J., Pak, R., Liu, E. & Schubert, M. C. Central not peripheral vestibular
processing impairs gait coordination. *Exp Brain Res* **235**, 3345-3355, doi:10.1007/s00221-
017-5061-x (2017).
85   Coelho, A. R., Fontes, R. C., Moraes, R., Barros, C. G. C. & Abreu, D. C. C. Effects of the use
of anchor systems in the rehabilitation of dynamic balance and gait in individuals with
chronic dizziness of peripheral vestibular origin: a single-blinded, randomized, controlled
clinical trial. *Archives of Physical Medicine and Rehabilitation* **101**, 249-257,
doi:https://doi.org/10.1016/j.apmr.2019.07.012
(2020).
86   Herssens, N. *et al.* An exploratory investigation on spatiotemporal parameters, margins of
stability, and their interaction in bilateral vestibulopathy. *Scientific Reports* **11**, 6427
(2021).
87   Reimann, H., Fettrow, T. & Jeka, J. J. Strategies for the control of balance during
locomotion. *Kinesiology Review* **7**, 18-25, doi:https://doi.org/10.1123/kr.2017-0053 (2018).
88   Kim, M. & Collins, S. H. Stabilization of a three-dimensional limit cycle walking model
through step-to-step ankle control. *IEEE Int Conference on Rehabilitation Robotics* **24**
(2013).
89   Bruijn, S. M., Meijer, O. G., Beek, P. J. & van Dieen, J. H. The effects of arm swing on human
gait stability. *J Exp Biol* **213**, 3945-3952, doi:10.1242/jeb.045112 (2010).
90   Collins, S. H., Adamczyk, P. G. & Kuo, A. D. Dynamic arm swinging in human walking.
Proceedings Royal Society **276**, 3679-3688 (2009).
91   Pijnappels, M., Kingma, I., Wezenberg, D., Reurink, G. & van Dieën, J. H. Armed against
falls: the contribution of arm movements to balance recovery after tripping. *Experimental*
Brain Research **201**, 689-699, doi:10.1007/s00221-009-2088-7 (2010).
92   Ali, A. S., Rowen, K. A. & Iles, J. F. Vestibular actions on back and lower limb muscles during
postural tasks in man   *Journal Physiology* **546.2**, 615-624,
doi:10.1113/jphysiol.2002.030031 (2003).
93   Guillaud, E. *et al.* Ancestral persistence of vestibulospinal reflexes in axial muscles in
humans. *Journal of neurophysiology* **123**, 2010-2023,
doi:https://doi.org/10.1152/jn.00421.2019 (2020).

94 Forbes, P. A., Siegmund, G. P., Happee, R., Schouten, A. C. & Blouin, J. S. Vestibulocollic
reflexes in the absence of head postural control. *J Neurophysiol* **112**, 1692-1702,
doi:10.1152/jn.00343.2014 (2014).
95 Forbes, P. A., Fice, J. B., Siegmund, G. P. & Blouin, J. S. Electrical Vestibular Stimuli Evoke
Robust Muscle Activity in Deep and Superficial Neck Muscles in Humans. **9**,
doi:10.3389/fneur.2018.00535 (2018).
96 Ihlen, E. A. *et al.* Phase-dependent changes in local dynamic stability of human gait. *J
Biomech* **45**, 2208-2214, doi:10.1016/j.jbiomech.2012.06.022 (2012).
97 Mahmoudian, A. *et al.* Phase-dependent changes in local dynamic stability during walking
in elderly with and without knee osteoarthritis. *J Biomech* **49**, 80-86,
doi:10.1016/j.jbiomech.2015.11.018 (2016).
98 Jin, J., Kistemaker, D., van Dieën, J. H., Daffertshofer, A. & Bruijn, S. M. The validation of
new phase-dependent gait stability measures: a modeling approach. *bioRxiv*,
doi:https://doi.org/10.1101/2020.06.18.156950 (2020).