Title: GNS561, a clinical-stage PPT1 inhibitor, has powerful antitumor activity against hepatocellular carcinoma via modulation of lysosomal functions

Sonia Brun[1, #, *, 15], Eric Raymond[1, 2, #], Firas Bassissi[1], Zuzana Macek Jilkova[3, 4, 5], Soraya Mezouar[1], Madani Rachid[1], Marie Novello[1], Jennifer Tracz[1], Ahmed Hamaï[6, 7], Gilles Lalmanach[8, 9], Lise Vanderlynden[8, 9], Eloïne Bestion[1, 10], Raphael Legouffe[11], Jonathan Stauber[12], Thomas Schubert[13], Maximilian G. Plach[13], Jérôme Courcambeck[1], Cyrille Drouot[1], Guillaume Jacquemot[1], Cindy Serdjebi[1], Gael Roth[3, 4, 5], Jean-Pierre Baudoin[14], Christelle Ansaldi[1], Thomas Decaens[3, 4, 5, #] and Philippe Halfon[1, #, *]

[1]Genoscience Pharma, Marseille, France

[2]Medical Oncology, Paris Saint-Joseph Hospital, Paris, France

[3]Institute for Advanced Biosciences, Research Center UGA / Inserm U 1209 / CNRS 5309, La Tronche, France

[4]University Grenoble Alpes, Faculté de médecine, France

[5]Clinique Universitaire d'Hépato-gastroentérologie, Pôle Digidune, CHU Grenoble, France

[6]Institut Necker-Enfants Malades, Inserm U1151-CNRS UMR 8253, Paris, France

[7]University Paris Descartes-Sorbonne Paris Cité, Paris, France

[8]INSERM, UMR1100, Centre d'Etude des Pathologies Respiratoires, Equipe "Mécanismes Protéolytiques dans l'Inflammation", Tours, France

[9]Tours University, Tours, France

[10]CNRS, IRD, MEPHI, IHU Méditerranée Infection, Aix Marseille University, Marseille, France

[11]ImaBiotech, Loos, France

[12]ImaBiotech, Billerica, USA

[13]2Bind GmbH, Regensburg, Germany

[14]Aix Marseille University, IRD, APHM, MEPHI, IHU Méditerranée Infection, Marseille, France

[15]Lead contact

[#]These authors contributed equally

*Correspondence: brun.sonia@hotmail.fr and p.halfon@genosciencepharma.com

SUMMARY

Hepatocellular carcinoma (HCC) is the most frequent primary liver cancer. Autophagy

inhibitors have been extensively studied in cancer but, to date, none has reached efficacy in

clinical trials. In this study, we demonstrated that GNS561, a new autophagy inhibitor,

whose anticancer activity was previously linked to lysosomal cell death, displayed high liver

tropism and potent antitumor activity against a panel of human cancer cell lines and in two

HCC in vivo models. We showed that GNS561, which is an effective lysosomotropic agent,

can reach and inhibit its enzyme target, palmitoyl-protein thioesterase 1, resulting in

lysosomal unbound Zn^{2+} accumulation, impairment of cathepsin activity, blockage of

autophagic flux, altered location of mTOR, lysosomal membrane permeabilization, caspase

activation and cell death. Accordingly, GNS561, currently tested in a global Phase 1b/2a

clinical trial against primary liver cancer, represents a promising new drug candidate and a

hopeful therapeutic strategy in cancer treatment.

Keywords: GNS561, liver cancer, lysosome, PPT1, autophagy, lysosomotropism, antitumor,

zinc, mTOR

INTRODUCTION

With an estimated 782,000 deaths in 2018, hepatocellular carcinoma (HCC) stands as the most common primary liver cancer and constitutes the fourth leading cause of cancer-related death worldwide (Bray et al. 2018). The rising incidence of HCC, the high worldwide mortality rate, and limited therapeutic options at advanced stages, make HCC a significant unmet medical need.

Autophagy-related lysosomal cell death, either alone or in connection with several other cell death pathways, has been recognized as a major target for cancer therapy (Aits and Jaattela 2013). Dysregulated autophagic-lysosomal activity and mTOR signaling were shown to allow cancer cells to become resistant to the cellular stress induced by chemotherapy and targeted therapy (Klempner et al. 2013). Recently, several lysosome-specific inhibitors were shown to target palmitoyl-protein thioesterase 1 (PPT1), resulting in the modulation of protein palmitoylation and antitumor activity in melanoma and colon cancer models (Rebecca et al. 2017, Rebecca et al. 2019). PPT1 palmitoylates proteins, enabling their degradation and intracellular trafficking of membrane-bound proteins. This process was shown to play a central role in the control of cellular autophagy. PPT1 was reported to be highly expressed in several cancer cell lines as well as in advanced stage cancers in patients (Rebecca et al. 2019).

Chloroquine (CQ) and hydroxychloroquine (HCQ) have been used for more than 50 years to prevent and treat malarial infections and autoimmune diseases. Based on the lysosomotropic properties and the capacity for autophagy inhibition, these molecules have been proposed as active drugs in cancer (Dolgin 2019, Pérez-Hernández et al. 2019) and have been extensively investigated in recent years (Kimura et al. 2013, Manic et al. 2014, Zhang et al. 2015, Shi et al. 2017, Verbaanderd et al. 2017, Xu et al. 2018). Over 40 clinical trials have been reported to evaluate the activity of both CQ or HCQ as single agent or in combination with chemotherapy in several tumor types (Manic et al. 2014, Shi et al. 2017, Verbaanderd et al. 2017). However, the required drug concentrations to inhibit autophagy were not achieved in humans, leading to inconsistent results in cancer clinical trials (Pascolo 2016, Rebecca et al. 2017, Plantone and Koudriavtseva 2018). This prompted research to identify novel compounds with potent inhibitory properties against autophagy for cancer therapy.

We previously reported that GNS561 was efficient in intrahepatic cholangiocarcinoma (iCCA)

by inhibiting late-stage autophagy and inducing a dose-dependent build-up of enlarged

lysosomes (Brun et al. 2019). In this study, we investigated the lysosomotropism of GNS561

and then the disruption of related lysosomal functions such as autophagy and lysosomal

enzymatic activity. We also identified lysosomal PPT1 as a target GNS561. Exposure to

GNS561 was shown to induce lysosomal unbound zinc ion (Zn^{2+}) accumulation, inhibition of

PPT1 and cathepsin activity, blockage of autophagic flux and mTOR displacement.

Interestingly, these effects resulted in lysosomal membrane permeabilization (LMP) and

caspase activation that led to cancer cell death. This mechanism was associated with dose-

dependent inhibition of cancer cell proliferation and tumor growth inhibition in several HCC

in vivo models. These data establish PPT1 and lysosomes as major targets for cancer cells

and led to the development of a clinical program investigating the effects of GNS561 in

patients with advanced HCC.

 RESULTS

GNS561 displays activity against human cancer cell lines and patient-derived cells

The effects of GNS561 on cell viability were investigated in a panel of human cancer cell

lines, including HCC, iCCA and colon, renal cell, breast, prostate, lung, and ovarian carcinoma

as well as acute myeloid leukemia, glioblastoma, and melanoma. As shown in Table 1,

GNS561 showed potent antitumor activity ranging from 0.22 ± 0.06 μM for the most

sensitive cell line (LN-18, a glioblastoma cell line) to 7.27 ± 1.71 μM for the least sensitive cell

line (NIH:OVCAR3, an ovarian cancer cell line). GNS561 was at least 10-fold more effective

than HCQ in cultured cancer cells. GNS561 also displayed activity in primary HCC patient-

derived cells and was on average 3-fold more potent than sorafenib, a reference drug in HCC

treatment (mean IC_{50} 3.37 ± 2.40 μM for GNS561 vs 10.43 ± 4.09 μM for sorafenib).

| Cancer type | Cell lines | Mean IC_{50} ± SD (μM) | | Primary HCC patient-derived cells | IC_{50} (μM) | |
		GNS561	HCQ		GNS561	sorafenib
Colon Carcinoma	HCT-116	1.22 ± 0.15	14.41 ± 1.5	LI0050	3.54	9.12
	HT-29	1.35 ± 0.04	24.18 ± 5.14	LI0574	2.41	8.65
Renal Cell Carcinoma	786-O	1.72 ± 0.17	21.65 ± 3.15	LI0612	6.93	17.94
	CAKI-1	1.10 ± 0.19	17.69 ± 1.29	LI0752	0.49	6.34
Ovarian Cancer	NIH:OVCAR3	7.27 ± 1.71	98.01 ± 12.75	LI0801	2.07	5.7
Melanoma	A375	1.2 ± 0.13	12.27 ± 2.8	LI1005	3.16	14.49
	SK-MEL-28	1.81 ± 0.5	22.78 ± 2.65	LI1098	6.95	10.85
Breast Cancer	MDA-MB-231	2.17 ± 0.14	14.13 ± 3.06	LI1646	1.44	10.33
Prostate Cancer	DU-145	1.09 ± 0.18	45.74 ± 0.55	Mean	3.37 ± 2.40	10.43 ± 4.09
	PC-3	2.56 ± 0.23	43.43 ± 6.04			
Lung Cancer	A549	1.69 ± 0.34	14.33 ± 1.59			
	NCI-H358	2.54 ± 0.34	54.07 ± 14.19			
HCC	HepG2	0.47 ± 0.15	11.55 ± 1.52			
	Huh7	0.88 ± 0.31	13.62 ± 0.71			
Glioblastoma	LN-229	0.60 ± 0.24	10.87 ± 1.23			
	LN-18	0.22 ± 0.06	5.27 ± 0.74			
Acute Myeloid Leukemia	KG-1	5.86 ± 1.64	43.92 ± 2.76			
	Mean	1.99 ± 1.86	27.52 ± 23.28			

Table 1. In vitro activity of GNS561 and HCQ in human cancer cell lines (left, $IC_{50} \pm SD$, μM)

and in vitro activity of GNS561 and sorafenib in primary hepatocellular carcinoma (HCC)

patient-derived cells (right, IC_{50}, μM).

GNS561 has antitumor properties in HCC in vivo models

The whole-body tissue distribution of GNS561 was investigated in rats after repeated oral

administration of GNS561 at a dose of 40 mg/kg/day for 28 days. Seven hours after the last

administration, the GNS561 level was measured by mass spectrometry imaging in the liver,

lung, stomach, brain, eye, salivary gland, kidney, heart, fat, muscle, testis, and skin (Figure 1).

GNS561 mainly accumulated in the liver, stomach and lung as shown by the calculated

organ/blood ratio (Figure 1B). Lower concentrations of GNS561 were also detected in eyes,

skin, brain and testis, indicating that GNS561 crosses the blood/brain barrier and the

blood/testis barrier to a limited extent (brain to blood and testis to blood ratios were 0.21

and 0.40, respectively).

Figure 1. Whole body tissue distribution of GNS561. (A) Mass spectrometry imaging of a

control rat (top) and a rat treated with GNS561 at a dose of 40 mg/kg/day for 28 days

(bottom). (B) Normalized intensity of GNS561 relative to blood signal in several organs of

GNS561-treated rats (Mean + SEM, n=2 except for eye, n=1). Of note, the GNS561 liver-to-

blood ratio is underestimated due to liver signal saturation.

Based on the high concentrations of GNS561 in the liver and potent in vitro activity against

HCC cells, the effects of GNS561 were investigated in vivo using two liver cancer models,

including the human HCC orthotopic patient-derived LI0752 xenograft mouse model and the

diethylnitrosanime (DEN)-induced immunocompetent rat HCC model.

In the HCC patient-derived LI0752 xenograft BALB/c nude mouse model, tumor volume and

weight were reduced by 37.1% and 34.4%, respectively, in mice treated with GNS561 at 50

mg/kg compared to the control (Figure S1A and B). Consistently, GNS561 treatment induced

a decrease in serum AFP levels in a dose-dependent manner and was significantly different

from the control at days 21 and 28 after treatment (Figure S1C-G).

Since HCC often develops in cirrhotic livers in humans, we further characterized the

antitumor effects of GNS561 in a DEN-induced cirrhotic rat model of HCC. Rats with already

developed HCC were either treated with sorafenib at 10 mg/kg, GNS561 at 15 mg/kg, or the

combination of both drugs (Figure S2). In this model, tumor progression was significantly

reduced by sorafenib (33.0%) and GNS561 (33.0%) compared to an untreated control group,

and the greatest decrease in tumor progression was observed by the combination (68%) that

displayed an additive effect (Figure 2A). Magnetic resonance imaging analyses further

showed a significant increase in the mean tumor size of 9.97 ± 0.97 mm in control rats

compared to 6.45 ± 0.35 mm with sorafenib, 5.48 ± 1.00 mm in GNS561 and 3.83 ± 0.52 mm

in the combination group (Figure 2B). Following liver resection, the macroscopic counting of

tumor nodules revealed significantly lower numbers in all treated groups compared to the

control group (Figure 2C). Immunohistochemical analyses of liver tumors showed a

significantly lower Cyclin D1-positive nuclear staining in the tumors of rats treated with

GNS561 or by the combination of GNS561 with sorafenib compared to the control group

(Figure 2D). GNS561 and combination treatments also significantly reduced Ki67 staining

compared to the control group (Figure 2E). The effects on Cyclin D1 and Ki67 were primarily

related to GNS561 exposure, as sorafenib alone showed no statistically significant

differences in Cyclin D1 or Ki67 staining compared to the control group. Our results further

showed that GNS561 and the combination treatment did not interfere with lipid or glucose

metabolism or kidney function but slightly affected some liver functions (Table S1).

Figure 2. GNS561 activity in a diethylnitrosanime-induced cirrhotic rat model of hepatocellular carcinoma. (A) Tumor progression assessment by comparison of tumor size obtained by magnetic resonance imaging (MRI) 1 and MRI 3 in the control, sorafenib at 10 mg/kg, GNS561 at 15 mg/kg and combination (GNS561 + sorafenib) groups. Macroscopic examination of livers with assessments of (B) tumor size and (C) tumor number at the surface of livers. (D) Representative images of nuclear Cyclin D1 staining and quantification of Cyclin D1-positive staining per high-power field (HPF). (E) Representative images of nuclear Ki67-positive staining and quantification of Ki67 staining per HPF. For all studies, mice n ≥ 6 per group. Data represent the mean + SEM. Comparison of means was performed by one-way ANOVA with Dunnett's post hoc analysis. * represents significant difference, at least $p < 0.05$.

GNS561 activates the caspase-dependent apoptosis pathway

We further wanted to characterize the antitumor effect of GNS561 and to determine whether GNS561 could trigger apoptotic cell death. To this end, annexin V/propidium iodide (PI) analysis was performed by flow cytometry after 48 h of GNS561 exposure in HepG2 cells. Early (Annexin V+/PI- staining) and late (Annexin V+/PI+ staining) apoptosis increased in a dose-dependent manner after GNS561 exposure (Figure 3A). The induction of apoptosis was confirmed by immunodetection of poly-ADP-ribose polymerase cleavage in GNS561-treated cells (Figure 3B). We further examined whether GNS561-induced apoptosis was related to

caspase activation. After 6 h of exposure, GNS561 had no effect on caspase 8 and caspase

3/7 activity in HepG2 cells (Figure 3C). In contrast, activation of caspase 8 and caspase 3/7

was observed after 24 h of treatment with GNS561, and this effect was sustained at 30 h. A

decrease in cell viability was concomitant with caspase activation (Figure 3C). The induction

of caspase activation was confirmed by flow cytometry (Figure 3D) and by detection of

cleavage of caspase 3 using immunoblot analysis (Figure 3E). Moreover, to confirm that

GNS561-induced cell death is caspase-dependent apoptosis, pretreatment (1 h) with the

cell-permeable pan-caspase inhibitor Z-VAD-FMK (5 µM) was performed. Cell viability was

restored in the presence of Z-VAD-FMK (Figure 3F), further confirming that GNS561 induced

a caspase-dependent apoptotic cell death.

Figure 3. GNS561 induces apoptotic cell death in HepG2 cells in a dose and time-dependent

manner through caspase activation. (A) Annexin V (A)/propidium iodide (PI) analysis by flow

cytometry after 48 h of GNS561 treatment. (B) Representative immunoblotting of the
cleaved and non-cleaved forms of poly-ADP-ribose polymer (PARP) after 24 h of GNS561
treatment. (C) Cell viability and activation of caspase 3/7 and 8 after 6, 24 and 30 h of
treatment with GNS561. (D) Representative immunoblotting of cleaved caspase 3 levels
after 24 h of treatment with GNS561. (E) Caspase-glow analysis by flow cytometry after 48 h
of treatment with GNS561. (F) Viable cell (A-/PI-) analysis by flow cytometry after
pretreatment with Z-VAD-FMK (ZVAD) at 5 µM for 1 h and then treatment with ZVAD at 5
µM and GNS561 for 48 h. For all blots, GAPDH was used as a loading control. For all studies,
n ≥ 3 biological replicates. Data represent the mean + SEM. For comparison, Student t-test
was used. * represents significant difference, at least $p < 0.05$.

GNS561 is a lysosomotropic agent

The intracellular localization of GNS561 in HepG2 cells was visualized using GNS561D, the
photoactivable analog of GNS561 containing a diazide moiety (Figure 4A). GNS561D showed
a punctuate fluorescent signal that colocalized with the intracellular vesicle-like structure
stained by LAMP1 (Figure 4B), demonstrating that GN561 accumulated in lysosomes and is a
lysosomotropic agent. Pretreatment with NH_4Cl, a weak base that rapidly increases
lysosomal pH, was further used to validate the lysosomotropic character of GNS561. As
shown in Figure 4B, NH_4Cl pretreatment strongly prevented lysosomal accumulation of
GNS561D. Then, we investigated whether GNS561 lysosomotropism was related to induced
cell death. For this purpose, HepG2 cells were pretreated for 2 h with NH_4Cl and then
treated with GNS561 for 24 h. Although a concentration of 20 mM NH_4Cl alone slightly
decreased viability (Figure 4C), it significantly attenuated the larger decrease in viability
induced by GNS561. These results were confirmed by pretreatment with bafilomycin A1 (Baf
A1), an inhibitor of the vacuolar H+-ATPase (Figure S3). Therefore, disrupting GNS561
lysosomal localization protected against GNS561-mediated cell death. These results
suggested that GNS561 antitumor activity in HepG2 cells is caused by its lysosomotropism.

Figure 4. The lysosomotropic agent GNS561 modulates lysosomal functions in the HepG2 cell line. (A) Chemical labeling of GNS561D in cells. (B) Lysosomal localization of GNS561D after NH₄Cl pretreatment (20 mM) for 30 min and then treatment with GNS561D (10 µM) and NH₄Cl (20 mM) for 90 min. (C) Cell viability after 24 h of GNS561 exposure in the

presence or absence of NH$_4$Cl (20 mM). (D) Staining of lysosomes (LysoTracker) and unbound

Zn^{2+} (Fluozin) after GNS561 treatment (1 h, 10 μM). Quantification of LysoTracker

fluorescence in arbitrary units (a.u.) (middle) and lysosomal unbound Zn^{2+} accumulation by

Pearson correlation coefficient between LysoTracker and Fluozin (right). (E) Fold change of

peptidase activity of cysteine cathepsins (including both cathepsins B and L) (CTSB/L),

cathepsin B (CTSB) and cathepsin D (CTSD) after GNS561 treatment (6 h and 24 h) calculated

in comparison with the control condition. (F) Representative immunoblotting of

procathepsin B (precursor form) and mature CTSB (left), procathepsin L (precursor form) and

mature CTSL (middle) and procathepsin D, intermediate and mature CTSD (right) after

GNS561 treatment for 16 h. For all blots, GAPDH was used as a loading control. For all

studies, n ≥ 3 biological replicates. Data represent the mean + SEM. For comparison, Student

t-test was used for (C), (B) and (D), and one-way ANOVA with Dunnett's post hoc analysis

was performed for (E). * represents significant difference, at least $p < 0.05$.

GNS561 modulates lysosomal functions

The GNS561 lysosomotropism-dependent cell death prompted us to examine GNS561

capacity to modulate lysosomal characteristics and functions.

Following continuous exposure to GNS561, staining of LysoTracker, which is a reagent

allowing the identification of the lysosomal compartment, increased in HepG2 cells (Figure

4D), suggesting that GNS561 prompted a dose-dependent build-up of enlarged lysosomes.

We therefore examined the enzymatic activity of three prominent lysosomal proteinases,

two cysteine cathepsins B (CTSB) and L (CTSL), and aspartic cathepsin D (CTSD). After 6 and

24 h of treatment, GNS561 significantly impaired, in a dose-dependent manner, the

enzymatic activity of cathepsins (Figure 4E). However, this decreased activity did not relate

to a direct GNS561-dependent inhibition of cathepsin activities (Figure S4). Based on the

literature, depressed proteolytic activity of cathepsins may result from an increased Zn^{2+}

lysosomal concentration and/or altered maturation of cathepsin precursors. Indeed, it has

been described that Zn^{2+} may downregulate the proteolytic activity of CSTB and CTSL

(Lockwood 2010, Lockwood 2013, Lockwood 2019). We investigated whether GNS561

modified unbound Zn^{2+} localization in HepG2 cells. As shown in Figure 4D, GNS561 induced a

strong accumulation of unbound Zn^{2+} in lysosomes, as evidenced by colocalization of the

fluorescent signals of Fluozin and LysoTracker in the merged images. This increase in
lysosomal unbound Zn^{2+} could explain the decreased proteolytic activity of CTSL and CTSB.
Cathepsins are synthesized as inactive zymogens, which are converted to their mature active
forms by other proteases or by autocatalytic processing (Turk et al. 2012). As depicted in
Figure 4F, GNS561 did not impact CTSB maturation, while it impaired the maturation of both
CTSL and CSTD (increase of precursor forms) and decreased their catalytic activity
accordingly.

As GNS561 induced lysosomal dysfunction, the effect of GNS561 on the autophagic process
was investigated. Herein, we showed that the GNS561-induced accumulation of light chain 3
phosphatidylethanolamine conjugate was not enhanced in the presence of BafA1 (Figure
S5), suggesting that GNS561 blocked autophagic flux.

PPT1 is a target of GNS561
Since PPT1 is critical for lysosomal function and is described to be the molecular target of
chloroquine derivatives (Rebecca et al. 2017, Rebecca et al. 2019), we investigated whether
PPT1 could be a molecular target of GNS561. First, the binding of GNS561 to recombinant
PPT1 was analyzed in vitro by nano differential scanning fluorimetry using HCQ as a positive
control (Rebecca et al. 2019). In the presence of GNS561 and HCQ, we observed a significant
dose-dependent decrease in PPT1 melting temperature (Figure 5A). Additionally, inhibition
of PPT1 enzymatic activity was observed in HepG2 cells treated with GNS561 (Figure 5B).
Moreover, the chemical mimetic N-*tert*-butylhydroxylamine (NtBuHA) attenuated autophagy
inhibition associated with GNS561 (Figure 5C), indicating that inhibition of PPT1 function by
GNS561 induced the observed anti-autophagy effect.
To determine whether inhibition of PPT1 function was responsible for the antitumoral
activity of GNS561, HepG2 cells were treated with GNS561 with or without NtBuHA
pretreatment. As shown in Figure 5D, NtBuHA partially prevented the antitumor activity of
GNS561, as evidenced by the increased viability of cells pretreated with NtBuHA. The same
rescue effect of NtBuHA pretreatment was observed for HCQ used as a positive control
(Figure S7). After demonstrating that NtBuHA had no impact on GNS561 lysosomal
localization (Figure S6), we validated that the impact of NtBuHA pretreatment was due to its

PPT1 mimetism, suggesting that inhibition of PPT1 function by GNS561 was partially liable

for its antitumoral activity.

The results of Rebecca et al. suggested that PPT1 inhibition could result in mTOR inhibition

through the displacement of mTOR from the lysosomal membrane (Rebecca et al. 2017,

Rebecca et al. 2019). Thus, we investigated the localization of mTOR after GNS561 treatment

using immunofluorescence microscopy. HCQ and EAD1 were used as positive controls (Sironi

et al. 2019). As shown in Figure 5E, GNS561 treatment, as well HCQ and EAD1 treatments,

significantly impaired mTOR localization to the lysosomal surface. Therefore, GNS561-

induced PPT1 inhibition resulted in displacement of mTOR from the lysosomal membrane

and consequently likely inhibited the mTOR signaling pathway.

Figure 5. GNS561 targets PPT1. (A) Nano differential scanning fluorimetry assays comparing

GNS561 + PPT1 and HCQ + PPT1 against the apo-PPT1 ligand. Data represent the mean (solid

lines) ± SEM (shaded areas) of two experiments. (B) PPT1 enzymatic activity of HepG2 cells

treated with GNS561 for 3 h. HCQ and HDSF were used as positive controls. The results were

compared to the diluent of GNS561 (control condition). (C) Representative immunoblotting

of LC3-II in HepG2 cells treated with GNS561 for 16 h in the presence or absence of NtBuHA
(8 mM). GAPDH was used as a loading control. Fold changes of normalized LC3-II level were
calculated against the control condition (diluent of GNS561 + diluent of NtBuHA). (D) Cell
viability percent against the control condition (diluent of GNS561 + diluent of NtBuHA) after
24 h of treatment with GNS561 in the presence or absence of NtBuHA (8 mM). (E) Staining of
lysosomes (LAMP2, green), mTOR (red) and nucleus (DAPI, blue) after treatment with
GNS561 and two positive controls, EAD1 and HCQ, for 16 h. Pearson correlation coefficient
between mTOR and LAMP2 was represented using box and whisker representation (min to
max). Scale bars represent 20 µm. In (B), (C) and (D), data represent the mean + SEM. For
comparison, Student t-test was used for (C) and (D) and one-way ANOVA with Dunnett's
post hoc analysis was performed for (B) and (E). For all studies except (A), n ≥ 3 biological
replicates. * represents significant difference, at least $p < 0.05$.
315

GNS561 induces LMP and cathepsin-dependent cell death
To characterize GNS561-induced changes in lysosomes, we analyzed LMP. To this end, we
took advantage of the steady endocytic capacity of cells to load fluorescent dextran into
lysosomes and the translocation of lysosomal localized dextran into the cytosol after LMP-
inducing insult. Fluorescent dextran in healthy cells appears in dense punctate structures
representing intact lysosomes, whereas after LMP, a diffuse staining pattern throughout the
cytoplasm is seen. After GNS561 treatment, such diffuse dextran staining was observed
(Figure 6A), suggesting an induction of LMP. As seen in Figure 6B, the loss of membrane
integrity, which is the hallmark of LMP, was observed by transmission electron microscopy of
HepG2 cells treated with 3 µM GNS561 for 24 h. To confirm this effect, cathepsin localization
was studied after GNS561 treatment. After 48 h of treatment, GNS561 decreased cathepsin
staining (Figure 6C), indicating that cathepsins were released into the cytosol, thus validating
LMP.
As cathepsin release into the cytosol after LMP may trigger cytosolic cellular death signaling
(Oberle et al. 2010), we evaluated the role of cathepsins in GNS56-induced cell death. To this
end, HepG2 cells were pretreated with an inhibitor of CTSD, pepstatin A, or an inhibitor of
CTSB, CA-074-Me. Under these conditions, cell viability was partially rescued (Figure 6D and
E), suggesting that the GNS561-induced apoptotic pathway is at least partially cathepsin-
dependent.

Figure 6. GNS561 induces LMP and cathepsin-dependent cell death in HepG2 cells. (A)
Localization of FITC-dextran after GNS561 treatment for the indicated times. (B) Electron
microscopy imaging of lysosomal membrane permeabilization (arrows) after GNS561
treatment (3 µM) for 24 h. (C) Localization of cathepsin B (CTSB), cathepsin D (CTSD) and
cathepsin L (CTSL) after GNS561 treatment for 48 h. (D) Viable cell (A-/PI-) analysis by flow
cytometry of cells pretreated or not with pepstatin A (Pep A) (5 µM) for 1 h and then treated
with Pep A (5 µM) and GNS561 or with GNS561 alone for 48 h. (E) Viable cell (A-/PI-) analysis
by flow cytometry of cells pretreated or not with CA-074-Me (20 µM) for 1 h and then
treated with CA-074-Me (20 µM) and GNS561 or with GNS561 alone for 48 h. Scale bars in
(A) and (C) represent 10 µm. For all studies, n ≥ 3 biological replicates. Data represent the
mean + SEM. For comparison, Student t-test was used. * represents significant difference, at
least $p < 0.05$.
349
350

DISCUSSION

Rapidly dividing and invasive cancer cells are strongly dependent on effective lysosomal
functions. Lysosomes are acidic and catabolic organelles found in nucleated human cells that
are responsible for the disposal and recycling of used and damaged macromolecules and
organelles, as well as the assimilation of extracellular materials incorporated into the cell by
endocytosis, autophagy, and phagocytosis. Increased autophagic flux and changes in
lysosomal compartments in cancer cells have been shown to promote invasion, proliferation,
tumor growth, angiogenesis, and drug resistance. Consistently, lysosomal changes are
expected to sensitize cells to lysosome-targeting anticancer drugs (Kallunki et al. 2013).
Many steps in the autophagy pathway represent potentially druggable targets and several
clinical trials have aimed to inhibit autophagy by inhibiting lysosomal functions using CQ and
HCQ. Unfortunately, CQ and HCQ failed to demonstrate consistent antitumor effects
possibly due to subeffective anticancer concentrations in humans, even with high doses.
Drug screening led us to identify GNS561 as a lead compound that displays lysosomotropism
and significantly higher antiproliferative effects in human cancer cells compared to HCQ.
We previously reported that GNS561 yielded antiproliferative activity in iCCA, inhibited late-
stage autophagy, and induced a dose-dependent enlargement of lysosomes (Brun et al.
2019). Based on these preliminary results, we further investigated the cellular mechanisms
by which GNS561 may lead to lysosomal changes and death in cancer cells. In this study, we
confirmed that GNS561 antitumor properties are strongly dependent on its lysosomotropic
properties. In accordance with the hypothesis proposed in our previous study (Brun et al.
2019), we showed here that GNS561 induced a dose-dependent increase in the number of
enlarged lysosomes, LMP leading to cytosolic cathepsin release, caspase activation, and
apoptotic cell death. These observations confirm prior reports that highlight the capability of
lysosomotropic agents to cause lysosomal stress and lysosomal enlargement (Wang et al.
2018). Moreover, studies demonstrated that lysosomal unbound Zn^{2+} buildup led to
lysosomal swelling, LMP, release of lysosomal enzymes, and cell death (Hwang et al. 2008,
Chung et al. 2009, Hwang et al. 2010, Yu et al. 2010). Further investigations are needed to
identify the upstream signals that initiate LMP in GNS561-treated cells.
PPT1, an enzyme involved in the removal of thioester-linked fatty acyl groups in proteins and
thus subsequently enabling the degradation and intracellular trafficking of membrane-bound

proteins, plays a central role in the control of cellular autophagy. PPT1 is highly expressed in
several cancer cell lines as well as in advanced stage cancers in patients (Rebecca et al.
2019). Recent data have shown that lysosome-specific inhibitors targeting PPT1 can
modulate protein palmitoylation and display antitumor activity in melanoma and colon
cancer models (Rebecca et al. 2017). Our data showed that PPT1 acts as a molecular target
of GNS561. GNS561 bound to PPT1 and inhibited its activity in cells. Cells treated with the
chemical mimetic NtBuHA were partially resistant to GNS561-mediated cytotoxicity and
attenuated GNS561-associated autophagic flux inhibition, suggesting that inhibition of the
thioesterase activity of PPT1 is essential for the anti-autophagic and antitumoral effects of
GNS561.

In our study, we observed that GNS561 modified the intracellular distribution and
localization of mTOR. This is in accordance with previous studies showing that inhibition of
PPT1 may displace the mTOR protein from the lysosomal membrane as a result of the
inhibition of vATPase/Ragulator/Rag GTPase interactions (Sancak et al. 2010, Korolchuk et al.
2011, Carroll et al. 2016, Rebecca et al. 2017, Rabanal-Ruiz and Korolchuk 2018, Rebecca et
al. 2019, Sironi et al. 2019). It was also described that lysosomal mTORClocalization brings it
in close vicinity to its main regulator, Rheb, and that as a result, the mTOR/Rheb interaction
can activate mTOR kinase activity leading to the phosphorylation of downstream effectors
(Carroll et al. 2016). Consistently, we hypothesized that GNS561-induced PPT1 inhibition led
to mTOR signaling pathway inhibition.

As previously observed in iCCA (Brun et al. 2019), we showed here that GNS561 induced a
significant decrease in the enzymatic activity of cathepsins. This decreased activity is unlikely
due to a direct inhibition of CTSL, CTSB and CTSD by GNS561 but rather could be the
consequence of both impairment of CTSL and CTSD maturation and lysosomal unbound Zn^{2+}
accumulation. As cathepsin activity is optimal in acidic pH (Gieselmann et al. 1985, Turk et al.
1999), we could also speculate that GNS561 may negatively influence the proteolytic activity
of cathepsins by inducing an increase in lysosomal pH via PPT1 inhibition. In fact, other
authors have shown that PPT1 deficiency in Cln1-/- mice disrupted the delivery of the v-
ATPase subunit V0a1 to the lysosomal membrane, leading to a dysregulation of lysosomal
acidification (Bagh et al. 2016). The authors suggested that S-palmitoylation by PPT1 may
play a critical role in the trafficking of the V0a1 subunit of v-ATPase to the lysosomal
membrane and in lysosomal pH regulation.

Based on prior studies, GNS561 was neither a zinc ionophore nor a zinc chelator (data not

shown), unlike CQ (Xue et al. 2014). However, our hypothesis that GNS561-induced PPT1

inhibition could lead to lysosomal deacidification could also explain the observed lysosomal

unbound Zn^{2+} accumulation after GNS561 treatment. In fact, as lysosomal pH is mainly

regulated by cation/anion movement across the lysosomal membrane, it was suggested that

a proton motive force was required to mediate unbound Zn^{2+} efflux (Lockwood 2013, Bin et

al. 2019).

In summary, GNS561-induced PPT1 inhibition may lead to two main mechanisms inducing

cancer cell death. One is related to lysosomal deacidification, which induces lysosomal

unbound Zn^{2+} accumulation, a decrease in the enzymatic activity of cathepsins, inhibition of

autophagic flux, lysosomal swelling, LMP, cathepsin release, and caspase-dependent

apoptosis. The other is linked to prevention of the interaction between v-ATPase and the

Ragulator complex, blockage of mTOR lysosomal recruitment, impairment of mTOR–Rheb

interaction and finally the inhibition of mTOR signaling pathway. Thus, by targeting PPT1,

GNS561 acts as a regulator of autophagy and mTOR, two major processes that drive cancer

aggressiveness. Finally, as lysosomes and autophagy are associated with adaptive

mechanisms of resistance to mTOR inhibition (Xie et al. 2013), GNS561 can disable mTOR

function and downregulate adaptive mechanisms of resistance.

An extensive preclinical program has been conducted to evaluate the antitumor activity,

pharmacological properties and toxicology of GNS561. Our data showed that GNS561

displays antiproliferative effects in several human cancer cells (cell lines and primary patient-

derived cells) and that GNS561 was more potent than HCQ. Analysis of the whole-body

tissue distribution of GNS561 in rats after repeated oral dosing of GNS561 showed that

GNS561 was mainly concentrated in the liver, stomach and lung. The data are consistent

with the basic lipophilic nature of GNS561 and with studies showing that basic lipophilic

drugs show high lysosomal tropism and high uptake in lysosome-profuse tissues, such as the

liver and the lung (Daniel and Wójcikowski 1997). As GNS561 had a high liver tropism, the

effect of GNS561 on tumor growth in vivo was evaluated using two liver cancer models: one

orthotopic human liver cancer xenograft mouse model (with an HCC patient-derived cell line,

LI0752) and one DEN-induced cirrhotic rat model with HCC. These studies showed that

GNS561 administered by oral gavage was well tolerated up to the doses of 50 mg/kg/day for

6 days in mice and up to 15 mg/kg/day for 6 weeks in rats and induced significant antitumor

growth activity that was either comparable to or higher than sorafenib. In addition, ina DEN-
induced cirrhotic rat model with HCC, the combination of GNS561 with sorafenib exerted an
additive effect in controlling tumor progression and cell proliferation. Furthermore, instead
of that observed with CQ and HCQ (Harder et al. 2018), the distribution of GNS561 into the
central nervous system was limited. Inactivating PPT1 mutations have long been known to
induce infantile neuronal cerebral lipofuscinosis and induce retinopathy during childhood
(Metelitsina et al. 2016). Germline PPT1 mutations were shown to selectively affect the
central nervous system, with no effects in other tissues. Prior clinical experience using CQ
and HCQ showed that retinopathy was one of the major toxicities in patients (Marmor et al.
2011). Authors have suggested that novel PPT1 inhibitors may take advantage of not
crossing the blood-brain barrier to avoid retinal toxicity (Rebecca et al. 2019). Interestingly,
our data shown that the disposition of GNS561 displays limited penetration into the brain in
rats, consistent with the lack of neurological and retinal toxicity observed in the current
Phase 1b/2a clinical trial of GNS561 (ClinicalTrials.gov).

In brief, our findings strengthen the importance of PPT1 and lysosomes as cancer targets.
Recently, it was shown that PPT1 inhibition by CQ derivatives or genetic Ppt1 inhibition
increases the antitumor activity of anti-PD-1 antibody in melanoma by M2 to M1 phenotype
switching in macrophages and a reduction in myeloid-derived suppressor cells in the tumor
(Sharma et al. 2020). As such, GNS561 represents a promising new candidate for drug
development in HCC either alone or in combination with other drugs, such as anti-PD-1
antibody.

METHODS

Details of the materials and methods are provided in the Supplementary Methods.

Cell culture

All cell lines were cultured in the presence of 5% CO_2 and 95% air in a humidified incubator and were maintained in medium containing 1% penicillin-streptomycin (Dutscher, #P06-07100) and 10% fetal bovine serum (HyClone, #SV30160.03C), except NIH:OVCAR3 and KG-1 cell lines, which were cultured in medium supplemented with 20% fetal bovine serum.

Animal models

The animals were checked daily for clinical signs, effects of tumor growth and any other abnormal effects. For experiments involving the mouse model (performed in CrownBio facilities), the protocol and any amendment(s) or procedures involving the care and use of animals were reviewed and approved by the Institutional Animal Care and Use Committee of CrownBio prior to experimentation, and during the study, the care and use of animals was conducted in accordance with the regulations of the Association for Assessment and Accreditation of Laboratory Animal Care. For the rat model, all animals received humane care in accordance with the Guidelines on the Humane Treatment of Laboratory Animals (Directive 2010/63/EU), and experiments were approved by the animal Ethics Committee: GIN Ethics Committee No.004.

Statistical analysis

Statistical analyses were performed using Prism 8.4.3 software (GraphPad Software Inc., CA, USA). For datasets with normal distribution, multiple comparisons were performed using one-way ANOVA with Dunnett's post hoc analysis. The parametric Student t-test was used to compare two groups of data with normal distribution. Data are presented as the mean values ± standard error mean (SEM) unless stated otherwise. Statistical significance was defined as a p-value < 0.05 and has been indicated by an asterisk in all figures.

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online.

ACKNOWLEDGMENTS

The authors are very grateful to Dr. Sebastian Müller and Dr. Raphaël Rodriguez from Curie

Institute for mechanistic analysis, Pr. Thierry Levade and Dr. Nathalie Andrieu from CRCT for

the PPT1 enzymatic assay, Keerthi Kurma and Seyedeh Tayebeh Ahmad Pour for the Institute

for Advanced Biosciences for technical support during animal experiments and Dr. François

Autelitano, Dr. Marie Guillemot and Philippe Fabre for Zn^{2+} localization analysis.

AUTHOR CONTRIBUTIONS

Conceptualization, S.B, E.R, F.B and P.H.; Methodology, S.B., F.B., T.D, P.H; Validation, S.B.,

Z.M.J, G.L, J.S, T.S, T.D; Formal Analysis, S.B., Z.M.J, M.N, J.T, A.H, G.L, R.L, T.S, M.G.P, J.P.B;

Investigation, Z.M.J, S.M., M.N, J.T, A.H, L.V, E.B, R.L, M.G.P, G.R; Resources, J.C, C.A;

Data Curation, C.D, G.J; Visualization, S.B, Z.M.J; Supervision, S.B, F.B, P.H; Project

Administration, S.B; Writing – original draft preparation, S.B.; Writing – review and editing,

S.B., E.R, F.B., Z.M.J, M.R, G.L, C.S, and P.H.; Funding Acquisition, P.H.

REFERENCES

Aits, S. and Jaattela, M. (2013). Lysosomal cell death at a glance. Journal of Cell Science
126(9): 1905-1912.

Bagh, M.B., Peng, S., Chandra, G., Zhang, Z. and Mukherjee, A.B. (2016). Dysregulation of V-
ATPase Function Impairs Lysosomal Acidification in a Mouse Model of a Rare Lysosomal
Storage Disease, INCL. The FASEB Journal 30(1_supplement): 879.871-879.871.

Bin, B.-H., Lee, S.-H., Bhin, J., Irié, T., Kim, S., Seo, J., Mishima, K., Lee, T.R., Hwang, D.,
Fukada, T., et al. (2019). The epithelial zinc transporter ZIP10 epigenetically regulates human
epidermal homeostasis by modulating histone acetyltransferase activity. The British Journal
of Dermatology 180(4): 869-880.

Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R.L., Torre, L.A. and Jemal, A. (2018). Global
cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36
cancers in 185 countries. CA: a cancer journal for clinicians 68(6): 394-424.

Brun, S., Bassissi, F., Serdjebi, C., Novello, M., Tracz, J., Autelitano, F., Guillemot, M., Fabre,
P., Courcambeck, J., Ansaldi, C., et al. (2019). GNS561, a new lysosomotropic small molecule,
for the treatment of intrahepatic cholangiocarcinoma. Investigational New Drugs 37(6):
1135-1145.

Carroll, B., Maetzel, D., Maddocks, O.D., Otten, G., Ratcliff, M., Smith, G.R., Dunlop, E.A.,
Passos, J.F., Davies, O.R., Jaenisch, R., et al. (2016). Control of TSC2-Rheb signaling axis by
arginine regulates mTORC1 activity. eLife 5.

Chung, H., Yoon, Y.H., Hwang, J.J., Cho, K.S., Koh, J.Y. and Kim, J.-G. (2009). Ethambutol-
induced toxicity is mediated by zinc and lysosomal membrane permeabilization in cultured
retinal cells. Toxicology and Applied Pharmacology 235(2): 163-170.

ClinicalTrials.gov. " Study of GNS561 in Patients With Liver Cancer." ClinicalTrials.gov.

Daniel, W.A. and Wójcikowski, J. (1997). Contribution of lysosomal trapping to the total
tissue uptake of psychotropic drugs. Pharmacology & Toxicology 80(2): 62-68.

Dolgin, E. (2019). Anticancer autophagy inhibitors attract 'resurgent' interest. Nature
Reviews. Drug Discovery 18(6): 408-410.

Gieselmann, V., Hasilik, A. and von Figura, K. (1985). Processing of human cathepsin D in
lysosomes in vitro. The Journal of Biological Chemistry 260(5): 3215-3220.

Harder, B.G., Blomquist, M.R., Wang, J., Kim, A.J., Woodworth, G.F., Winkles, J.A., Loftus, J.C.
and Tran, N.L. (2018). Developments in Blood-Brain Barrier Penetrance and Drug
Repurposing for Improved Treatment of Glioblastoma. Frontiers in Oncology 8.

Hwang, J.J., Kim, H.N., Kim, J., Cho, D.-H., Kim, M.J., Kim, Y.-S., Kim, Y., Park, S.-J. and Koh, J.-
Y. (2010). Zinc(II) ion mediates tamoxifen-induced autophagy and cell death in MCF-7 breast

cancer cell line. Biometals: An International Journal on the Role of Metal Ions in Biology,
Biochemistry, and Medicine 23(6): 997-1013.

Hwang, J.J., Lee, S.-J., Kim, T.-Y., Cho, J.-H. and Koh, J.-Y. (2008). Zinc and 4-hydroxy-2-
nonenal mediate lysosomal membrane permeabilization induced by H2O2 in cultured
hippocampal neurons. The Journal of Neuroscience: The Official Journal of the Society for
Neuroscience 28(12): 3114-3122.

Kallunki, T., Olsen, O.D. and Jäättelä, M. (2013). Cancer-associated lysosomal changes:
friends or foes? Oncogene 32(16): 1995-2004.

Kimura, T., Takabatake, Y., Takahashi, A. and Isaka, Y. (2013). Chloroquine in cancer therapy:
a double-edged sword of autophagy. Cancer Research 73(1): 3-7.

Klempner, S.J., Myers, A.P. and Cantley, L.C. (2013). What a tangled web we weave:
emerging resistance mechanisms to inhibition of the phosphoinositide 3-kinase pathway.
Cancer Discovery 3(12): 1345-1354.

Korolchuk, V.I., Saiki, S., Lichtenberg, M., Siddiqi, F.H., Roberts, E.A., Imarisio, S., Jahreiss, L.,
Sarkar, S., Futter, M., Menzies, F.M., et al. (2011). Lysosomal positioning coordinates cellular
nutrient responses. Nature Cell Biology 13(4): 453-460.

Lockwood, T.D. (2010). The lysosome among targets of metformin: new anti-inflammatory
uses for an old drug? Expert Opinion on Therapeutic Targets 14(5): 467-478.

Lockwood, T.D. (2013). Lysosomal metal, redox and proton cycles influencing the CysHis
cathepsin reaction. Metallomics: Integrated Biometal Science 5(2): 110-124.

Lockwood, T.D. (2019). Biguanide is a modifiable pharmacophore for recruitment of
endogenous Zn2+ to inhibit cysteinyl cathepsins: review and implications. Biometals: An
International Journal on the Role of Metal Ions in Biology, Biochemistry, and Medicine 32(4):
575-593.

Manic, G., Obrist, F., Kroemer, G., Vitale, I. and Galluzzi, L. (2014). Chloroquine and
hydroxychloroquine for cancer therapy. Molecular & Cellular Oncology 1(1): e29911.

Marmor, M.F., Kellner, U., Lai, T.Y.Y., Lyons, J.S., Mieler, W.F. and Ophthalmology, A.A.o.
(2011). Revised recommendations on screening for chloroquine and hydroxychloroquine
retinopathy. Ophthalmology 118(2): 415-422.

Metelitsina, T.I., Waggoner, D.J. and Grassi, M.A. (2016). BATTEN DISEASE CAUSED BY A
NOVEL MUTATION IN THE PPT1 GENE:. Retinal Cases & Brief Reports 10(3): 211-213.

Oberle, C., Huai, J., Reinheckel, T., Tacke, M., Rassner, M., Ekert, P.G., Buellesbach, J. and
Borner, C. (2010). Lysosomal membrane permeabilization and cathepsin release is a
Bax/Bak-dependent, amplifying event of apoptosis in fibroblasts and monocytes. Cell Death
and Differentiation 17(7): 1167-1178.

Pascolo, S. (2016). Time to use a dose of Chloroquine as an adjuvant to anti-cancer
chemotherapies. European Journal of Pharmacology 771: 139-144.

Pérez-Hernández, M., Arias, A., Martínez-García, D., Pérez-Tomás, R., Quesada, R. and Soto-
Cerrato, V. (2019). Targeting Autophagy for Cancer Treatment and Tumor
Chemosensitization. Cancers 11(10): 1599.

Plantone, D. and Koudriavtseva, T. (2018). Current and Future Use of Chloroquine and
Hydroxychloroquine in Infectious, Immune, Neoplastic, and Neurological Diseases: A Mini-
Review. Clinical Drug Investigation 38(8): 653-671.

Rabanal-Ruiz, Y. and Korolchuk, V.I. (2018). mTORC1 and Nutrient Homeostasis: The Central
Role of the Lysosome. International Journal of Molecular Sciences 19(3).

Rebecca, V.W., Nicastri, M.C., Fennelly, C., Chude, C.I., Barber-Rotenberg, J.S., Ronghe, A.,
McAfee, Q., McLaughlin, N.P., Zhang, G., Goldman, A.R., et al. (2019). PPT1 Promotes Tumor
Growth and Is the Molecular Target of Chloroquine Derivatives in Cancer. Cancer Discovery
9(2): 220-229.

Rebecca, V.W., Nicastri, M.C., McLaughlin, N., Fennelly, C., McAfee, Q., Ronghe, A., Nofal,
M., Lim, C.-Y., Witze, E., Chude, C.I., et al. (2017). A Unified Approach to Targeting the
Lysosome's Degradative and Growth Signaling Roles. Cancer Discovery 7(11): 1266-1283.

Sancak, Y., Bar-Peled, L., Zoncu, R., Markhard, A.L., Nada, S. and Sabatini, D.M. (2010).
Ragulator-Rag complex targets mTORC1 to the lysosomal surface and is necessary for its
activation by amino acids. Cell 141(2): 290-303.

Sharma, G., Ojha, R., Noguera-Ortega, E., Rebecca, V.W., Attanasio, J., Liu, S., Piao, S., Lee,
J.J., Nicastri, M.C., Harper, S.L., et al. (2020). PPT1 inhibition enhances the antitumor activity
of anti-PD-1 antibody in melanoma. JCI insight 5(17).

Shi, T.-T., Yu, X.-X., Yan, L.-J. and Xiao, H.-T. (2017). Research progress of hydroxychloroquine
and autophagy inhibitors on cancer. Cancer Chemotherapy and Pharmacology 79(2): 287-
294.

Sironi, J., Aranda, E., Nordstrøm, L.U. and Schwartz, E.L. (2019). Lysosome Membrane
Permeabilization and Disruption of the Molecular Target of Rapamycin (mTOR)-Lysosome
Interaction Are Associated with the Inhibition of Lung Cancer Cell Proliferation by a
Chloroquinoline Analog. Molecular Pharmacology 95(1): 127-138.

Turk, B., Dolenc, I., Lenarcic, B., Krizaj, I., Turk, V., Bieth, J.G. and Björk, I. (1999). Acidic pH as
a physiological regulator of human cathepsin L activity. European Journal of Biochemistry
259(3): 926-932.

Turk, V., Stoka, V., Vasiljeva, O., Renko, M., Sun, T., Turk, B. and Turk, D. (2012). Cysteine
cathepsins: From structure, function and regulation to new frontiers. Biochimica et
Biophysica Acta (BBA) - Proteins and Proteomics 1824(1): 68-88.
664
665 Verbaanderd, C., Maes, H., Schaaf, M.B., Sukhatme, V.P., Pantziarka, P., Sukhatme, V.,
666 Agostinis, P. and Bouche, G. (2017). Repurposing Drugs in Oncology (ReDO)-chloroquine and
667 hydroxychloroquine as anti-cancer agents. Ecancermedicalscience 11: 781.
668
669 Wang, F., Gómez-Sintes, R. and Boya, P. (2018). Lysosomal membrane permeabilization and
670 cell death. Traffic 19(12): 918-931.
671
672 Xie, X., White, E.P. and Mehnert, J.M. (2013). Coordinate Autophagy and mTOR Pathway
673 Inhibition Enhances Cell Death in Melanoma. PLOS ONE 8(1): e55096.
674
675 Xu, R., Ji, Z., Xu, C. and Zhu, J. (2018). The clinical value of using chloroquine or
676 hydroxychloroquine as autophagy inhibitors in the treatment of cancers: A systematic review
677 and meta-analysis. Medicine 97(46): e12912.
678
679 Xue, J., Moyer, A., Peng, B., Wu, J., Hannafon, B.N. and Ding, W.-Q. (2014). Chloroquine Is a
680 Zinc Ionophore. PLoS ONE 9(10): e109180.
681
682 Yu, L., McPhee, C.K., Zheng, L., Mardones, G.A., Rong, Y., Peng, J., Mi, N., Zhao, Y., Liu, Z.,
683 Wan, F., et al. (2010). Termination of autophagy and reformation of lysosomes regulated by
684 mTOR. Nature 465(7300): 942-946.
685
686 Zhang, Y., Liao, Z., Zhang, L.-j. and Xiao, H.-t. (2015). The utility of chloroquine in cancer
687 therapy. Current Medical Research and Opinion 31(5): 1009-1013.
688