<u>Estimating the extrinsic incubation period of malaria using a mechanistic model of</u>

<u>sporogony</u>

Isaac J. Stopard, Thomas S. Churcher & Ben Lambert

MRC Centre for Global Infectious Disease Analysis, Imperial College London

Keywords: epidemiology, systems biology, thermal ecology, multiscale, temperature,

climate, *Plasmodium falciparum*, mosquito, *Anopheles*, development rate, within-host

Running title: estimating the extrinsic incubation period of malaria

Abstract

During sporogony, malaria-causing parasites infect a mosquito, reproduce and migrate to

the mosquito salivary glands where they can be transmitted the next time blood-feeding

occurs. The time required for sporogony, or extrinsic incubation period (EIP), is a crucial

determinant of malaria transmission intensity. The EIP is typically estimated as the time for

a given percentile of infected mosquitoes to have salivary gland sporozoites (the infectious

parasite life stage). Many mechanisms, however, affect the observed sporozoite prevalence

including the human-to-mosquito transmission probability and possibly differences in

mosquito mortality according to infection status. To account for these various mechanisms,

we present a mechanistic mathematical model ("mSOS"), which explicitly models key

processes at the parasite, mosquito and observational scales. Fitting this model to

experimental data, we find greater variation in EIP than previously thought: we estimated

the range between two percentiles of the distribution, EIP_{10}–EIP_{90} (at 27°C), as 4.5 days,

compared to 0.9 days using existing methods. This pattern holds over the range of study

temperatures included in the dataset. Increasing temperature from 21°C to 34°C decreased

the EIP_{50} from 16.1 to 8.8 days and the human-to-mosquito transmission probability from

84% to 42%. Our work highlights the importance of mechanistic modelling of sporogony to

(1) improve estimates of malaria transmission under different environmental conditions or

disease control programs and (2) evaluate novel interventions that target the mosquito life

stages of the parasite.

Author summary

Anopheles mosquitoes become infected with malaria-causing parasites when blood feeding

on an infectious human host. The parasites then process through a number of life stages,

which begin in the mosquito gut and end in the salivary glands, where the newly formed

infectious parasites can be transmitted to another host the next time a mosquito blood-

feeds. The large variability in parasite numbers and development times that exists between

mosquitoes, environments and parasites, mean that understanding parasite population

dynamics from individual mosquito dissections is difficult. Here, we introduce a

mathematical model of the mosquito life stages of parasites that mimics key characteristics

of the biology. We show that the model's parameters can be chosen so that its predictions

correspond with experimental observations. In doing so, we estimate key system

characteristics that are crucial determinants of malaria transmission intensity. Our work is a

step towards a realistic model of within-mosquito parasite dynamics, which is increasingly

important given that many recently proposed disease interventions specifically target

mosquito life stages of the parasite.

1   Introduction

Malaria remains a leading cause of morbidity and mortality worldwide, with an extremely

inequitable distribution: over 400,000 people, primarily children under the age of five in

sub-Saharan Africa, die annually due to malaria (1). The widespread use of vector control

tools that kill adult *Anopheles* mosquitoes is largely responsible for a historical decline in

malaria incidence (2); a result foretold by early mathematical models, which predicted the

sensitivity of malaria transmission to adult mosquito survival (3,4). For a newly infected

mosquito to become infectious, it must survive the extrinsic incubation period (EIP). Since

the EIP is long relative to mosquito life expectancy, only older mosquitoes can pass on

infection meaning malaria transmission responds acutely to changes in survival (3,5).

The EIP is defined as the duration of sporogony: the obligate reproduction of

malaria-causing *Plasmodium* parasites (henceforth parasites) within the mosquito (6). First,

female mosquitoes feed on an infectious host. A proportion of these mosquitoes, as

determined by the human-to-mosquito transmission probability (5), ingest male and female

Plasmodium gametocytes within the red blood cells (RBCs) of the blood-meal (7). The

change of parasite host (from human to mosquito) involves certain environmental changes,

including a decrease in temperature, which collectively trigger gametogenesis and the

parasites to emerge from the RBCs (8,9). Fertilisation occurs within the mosquito midgut,

where gametes fuse into a single zygote, which differentiates into a motile ookinete (10).

Within a few days, ookinetes migrate across the mosquito midgut epithelial wall, and the

parasites that survive the mosquito innate immune response (11) go on to form immobile

oocysts beneath the midgut basal lamina (12). The number of oocysts remains relatively

constant as they grow in size and their genome mitotically replicates (12). Oocysts then

burst with each releasing hundreds of infectious sporozoites, which migrate to the mosquito

salivary glands, completing the EIP (6,13).

The EIP and human-to-mosquito transmission probability are typically estimated in

the laboratory using experimentally introduced infections. Laboratory reared mosquitoes

are fed on infectious blood through a membrane feeder, and the parasites are allowed to

develop before the mosquitoes are dissected to determine the presence and number of

oocysts or sporozoites (13,14). Since a mosquito can be dissected only once, it is not

possible to observe parasites dynamics within a single mosquito. Numerous dissections are

therefore used to reconstruct the temporal dynamics of sporogony in the population at

large (6).

Historically, the EIP has been recorded as the time from blood-feeding until the first

mosquito is observed with any salivary gland sporozoites (15,16). Parasite development

rates are, however, highly heterogenous due to differences in mosquito nutrition (17,18)

and environmental temperature (14,19), and considerable variation remains even after

accounting for these factors (14,17). Mosquito and parasite genetic differences may also

contribute to variation (15), and even in genetically identical cells reaction rates are noisy,

as genes are regulated by small fluctuations in regulatory molecules (20). Given this

variation, any single measure – especially one based on a single mosquito, which has

historically been the case – does not adequately represent the EIP. Instead, more recent

studies attempt to characterise the variability in the EIP, which is reported as the time for

given percentiles of infected mosquitoes to display sporozoites (15). A rigorous framework

for characterising the distribution of the EIP is yet to be defined.

Not all mosquitoes fed infectious blood will develop observable oocysts or

sporozoites. The human-to-mosquito transmission probability depends on many factors

including differences in blood meal size, gametocyte density (7,21,22), the mosquito

immune response (11,23) and midgut microbiota (24,25) amongst others. To estimate the

EIP percentiles, it is necessary to determine the number of infected mosquitoes within the

sample as a proportion of all infected individuals. To do this, it is assumed that the

maximum observed oocyst- or sporozoite- prevalence is the actual proportion of

mosquitoes with viable infection (15). Focussing on raw temporal changes in observed

prevalence without an underlying mechanistic model may, however, overlook key

processes. In the laboratory, malaria infections may alter mosquito survival in infected

mosquitoes (26), meaning the observed maximum prevalence may not represent the true

proportion of infected mosquitoes. What's more, certain variables can impact multiple

mechanisms simultaneously (27), requiring a finer-scale understanding of the constituent

processes.

Temperature, for example, can modulate parasite development rate (14,19), survival

of laboratory mosquitoes (14) and mosquito immune response (28), causing differences in

vector competence (29). To date, the relationship between temperature and the EIP has

been modelled using the degree-day model, which parameterises the total amount of heat

required to complete sporogony (30). Polynomial (quadratic or Brière) functions have also

been fit to the relationship between the EIP and temperature (31,32). Parameterisations of

both these methods rely on a point estimate of the EIP and do not model the constituent

processes that produce these observations. Mechanistic models that simulate parasite

population dynamics during sporogony may provide a more informative framework (33,34).

Here, we introduce a mathematical model, the "multiscale Stochastic model Of

Sporogony (mSOS)", to capture the temporal dynamics of *Plasmodium* sporogony. By

explicitly modelling the underlying biology, at the parasite, mosquito and experimental

scales, we aim to provide an accurate representation of parasite development (35). We fit

mSOS to laboratory membrane feeding assay data and present new EIP estimates which

contrast with the existing literature. We also demonstrate how mSOS can be used to

estimate the impact of temperature on *Plasmodium falciparum* sporogony within *Anopheles*

mosquitoes. All code and data needed to recapitulate our results is available at

https://github.com/IsaacStopard/mSOS.

2 Models and methods

2.1 Data

We are unaware of a single study with sufficient data to fit the model. To capture the full population dynamics of oocysts and sporozoites, we combined data from four published studies (Table 1). Most data came from Standard Membrane Feeding Assays (SMFA) from the same laboratory, conducted with the same parasite-vector combination (14,18,36). These studies collected data over a range of days and temperatures using a laboratory strain of *P. falciparum* (NF54) and laboratory-reared *Anopheles stephensi* mosquitoes. To examine oocyst prevalence prior to day five, we also included data from a Direct Membrane Feeding Assay (DMFA) (37), which used wild *P. falciparum* parasites to infect laboratory-reared *Anopheles gambiae sensu stricto* mosquitoes. It is assumed there is no difference between the two parasite-vector combinations. All studies replicated the following experimental protocol: mosquitoes were raised from larvae under laboratory conditions; adult female mosquitoes were then fed on *P. falciparum*-infected blood via a membrane feeder; at various days post infection, blood-fed mosquitoes were collected, dissected and the presence of oocysts or salivary-gland sporozoites identified by microscopy. The oocyst load within each mosquito was estimated by counting observable oocysts. Sporozoites were quantified by presence or absence. Only mosquito larvae kept on a high food diet were used in these analyses, and, in all cases, mosquitoes were housed at a constant temperature and humidity (see Table 1 for the range of different temperatures explored).

147 **Table 1 Summary of different studies, parasite-vector combinations and experimental proceedures used to parameterise the model.**

Malaria species (strain)	Vector species (strain)	Days post blood-feeding when oocyst presence or load was determined by dissection	Days post blood-feeding when sporozoite presence was determined by dissection	Dead mosquitoes counted?	Adult mosquito housing temperature (°C)	Reference
P. falciparum (NF54)	*A. stephensi* (laboratory)	7 (oocyst load)	15	Yes, infectious blood-fed & uninfected blood-fed (control) mosquitoes	27, 30 & 33	Murdock et al. (36)
P. falciparum (NF54)	*A. stephensi* (laboratory)	Not recorded	Varied with temperature: 10 to 23 & 25 (21°C), 9 to 23 & 25 (24°C), 8 to 18, 20, 22, 24 & 25 (27°C), 6 to 17 & 19 (30°C), 5 to 15, 17 & 19 (32°C) and 5 to 14 & 16 (34°C).	Yes, infectious blood-fed mosquitoes only	21, 24, 27, 30, 32 & 34	Shapiro et al. (14)
P. falciparum (NF54)	*A. stephensi* (laboratory)	5, 6, 7, 8, 9 & 10 (oocyst load)	9, 10, 11, 12, 13, 14, 15 & 16	Yes, infectious blood-fed mosquitoes only	27	Shapiro et al. (18)
P. falciparum (wild)	*A. gambiae s.s.* (laboratory)	3, 4, 5, 6, 7 & 8 (oocyst presence used)	Not used	No	27 - 28	Bompard et al. (37)

149   2.2   Model overview
150
mSOS models the oocyst and salivary gland sporozoite (henceforth sporozoite) life stages, as

oocysts are the most widely recorded outcome of membrane-feeding assays, and

sporozoites are the most epidemiologically important life stage. We explicitly model the

counts of oocyst- and sporozoite-positive mosquitoes to determine how the prevalence (i.e.

the percent of positive mosquitoes among the sample) of each life stage varies over time.

We also model oocyst intensity (mean oocyst load among the sample), as it provides more

information on the underlying parasite dynamics and is considered a more reliable estimate

of human-to-mosquito transmission (38,39).

Sporogony is modelled at multiple scales, and all parameters are fit simultaneously

in a single model. At the parasite scale, we model the sequence of three life stages:

inoculant → oocyst → sporozoite (Figure 1A). We do not model the pre-oocyst life stages

since they are harder to count and are rarely recorded. Rather, these are represented in a

notional "inoculant" stage ingested during blood-feeding, each of which develops into a

single oocyst. mSOS captures both the time it takes for each oocyst to appear, and the time

for each oocyst to develop into sporozoites. The time taken for each of these transitions to

occur is assumed independent of the other, and we model these times stochastically.

At the mosquito scale, mosquitoes are initially infected with different numbers of

inoculant stage parasites, with each load determined stochastically. We also allow infectious

blood-fed mosquitoes to avoid infection (via a human-to-mosquito transmission probability)

(Figure 1B). Potential differences in survival of the mosquito population due to differences

in malaria infection (26) are explicitly modelled.

At the observation level, to reconstruct the temporal dynamics of sporogony at a

population level, we recreate populations of parasites nested within a population of

mosquito vectors, which then form samples of dissected mosquitoes (Figure 1C). We allow

the composition of each sample of dissected mosquitoes to vary according to the

proportions of infected and uninfected mosquitoes alive at the time of dissection.

To incorporate temperature-dependency, we allow the time for the appearance of

oocysts (40,41), and the human-to-mosquito transmission probability, to be influenced by

temperature. The development time from oocyst to sporozoites and the mean parasite load

among infected mosquitoes are assumed to be independent of temperature.

In what follows, we describe the model in detail. To provide greater clarity, a

Mathematica file that steps through the derivations is available in the supplementary

information.

Figure 1 Structure of mSOS: a multiscale model of the population dynamics of *Plasmodium falciparum* during sporogony. (a) Multiple malaria parasites are found within a single mosquito; we separately model development time from inoculation at blood-feeding (G) to oocyst (O) and from oocyst to salivary-gland sporozoites (S). If dissected, a mosquito is "positive" for a particular parasite life stage if at least a single parasite has developed. We do not model the decline in observed oocyst numbers due to oocyst bursting, since we do not have sufficient later oocyst observations. (b) Mosquitoes are infected via a membrane feeder; parasite load-varies in each due to differences in the number of parasites ingested and variation in

mosquito immune response. (c) Temporal dynamics of sporozoite prevalence within a mosquito population:

following the infectious blood-feed, a proportion of the population is infected with malaria parasites. The

parasites develop into sporozoites causing the mosquitoes to become infectious. Throughout the experiments,

mosquito mortality (in the laboratory) may be greater in infected mosquitoes, resulting in an eventual decline

in observed sporozoite prevalence.

2.2.1   Parasite scale

We model the time taken, T_{GO} , for each undeveloped oocyst at the time of blood-feeding

(the notional inoculant stage denoted 'G') to develop into an observable oocyst ('O'). We

then model the time taken, T_{OS} , for each oocyst to develop, burst, and for the sporozoites

to migrate to the salivary glands (we term sporozoites in the salivary glands 'S'). So, these

transitions follow:

206
$$G \xrightarrow{T_{GO}} O \xrightarrow{T_{OS}} S$$

These development times are modelled stochastically as: $T_{GO} \overset{i.i.d.}{\sim} \Gamma(\alpha_{GO}, \beta_{GO})$ and

$T_{OS} \overset{i.i.d.}{\sim} \Gamma(\alpha_{OS}, \beta_{OS})$; that is, we assume that the time it takes a subsequent oocyst to burst

into sporozoites is independent of the initial time taken for the oocyst to develop. This

assumption also implies the transitions of each parasite occur independently, meaning

density dependent population dynamics do not occur. We assume a parameterisation of the

gamma distribution such that its mean is $E(T_i) = \alpha_i/\beta_i$, where $i \in (GO, OS)$, α_i is the

shape parameter and β_i is the rate parameter.

The time required for G to develop into S is given by the sum: $T_{GS} = T_{GO} + T_{OS}$. An

analytic form for the cumulative density function (CDF) of the sum of two gamma

distributed random variables with different rate (β) parameters is currently not known. This

aggregate distribution can, however, be approximated by a gamma distribution where the

mean, λ, and variance, σ^2, match that of the true distribution (42),

219
$$T_{GS} \sim \Gamma(\alpha_{GS}, \beta_{GS}), \tag{2.01}$$

where,

$\alpha_{GS} = \dfrac{\lambda^2}{\sigma^2}, \qquad \beta_{GS} = \dfrac{\lambda}{\sigma^2}, \qquad \lambda = \dfrac{\alpha_{os}\beta_{GO} + \alpha_{GO}\beta_{os}}{\beta_{GO}\beta_{os}}, \qquad \sigma^2 = \dfrac{\alpha_{os}\beta_{GO}^2 + \alpha_{GO}\beta_{os}^2}{\beta_{GO}^2\beta_{os}^2}. \tag{2.02}$

2.2.2   Mosquito scale

Mosquitoes are treated as self-contained populations of parasites. When feeding on

infectious blood, mosquitoes receive a heterogeneous load of gametocytes, with some

receiving none at all (7). Additionally, not all parasites develop into the observed oocyst

stage due to the innate immune response of some mosquitoes (13,38). Evidence indicates

that, assuming the mosquito is still alive, the majority of oocysts will produce sporozoites

(43). Hence, we term a mosquito "successfully infected" if at least one viable oocyst could

develop from the initial infective load given sufficient time and define this probability of

successful infection (i.e. the human-to-mosquito transmission probability) as δ. Among

successfully infected mosquitoes, the initial load of G-stage parasites, n, is modelled by a

zero-truncated negative binomial distribution,

$$n \sim NB(\mu, k), \qquad n \in [1, \infty), \tag{2.03}$$

where, μ is the mean and k is the overdispersion parameter. Evidence indicates μ and

δ may be jointly influenced by gametocyte density, which is modulated by temperature (29).

We treated these parameters as independent, as there was an insufficient combination of

gametocyte density and temperature treatments in the data.

To model the observed oocyst or sporozoite presence, we assume a mosquito is

measured as "positive" if at least a single parasite has developed to the given life stage. That

is, we assume that dissection always uncovers some parasites of a given life stage should

any exist.

Let the time taken for parasite j to develop into a subsequent stage be T_j for $j =$

$1, 2, \dots, n$ parasites within a specific mosquito. Whether a mosquito is positive for a

particular stage at time t then depends on whether $t > T_{1:n}^{min} = min(T_1, T_2, \dots, T_n)$. Since,

we are concerned with the minimum of a series of random variables, we are in the realm of

order statistics – see (44). Specifically, $T_{1:n}^{min}$ is the 1^{st} order statistic of the sample of n

development times. Considering a single parasite, j, the probability that it has developed by

time t is given by $Pr(t \geq T_j) = Q(\alpha, 0, \beta t)$, where $Q(\alpha, 0, \beta t)$ is the CDF of the gamma

distribution governing that particular life stage transition. Specifically, Q is the generalized

regularized incomplete gamma function $Q(a, z_0, z_1) = \int_{z_0}^{z_1} t^{a-1} e^{-t} dt / \int_0^\infty t^{a-1} e^{-t} dt$, and

α, β are the shape and rate parameters of the underlying gamma distribution (these

parameters will, in general, be different for T_{GO} and T_{GS}). Considering n parasites within a

single mosquito, the probability that at least one of them has developed by time t is:

255
$$\Pr\left(t \geq T_{1:n}^{min}\right) = 1 - \prod_{j=1}^{n} \Pr\left(t < T_j\right) = 1 - \left(1 - Q(\alpha, 0, \beta t)\right)^n. \qquad (2.04)$$

256   To model the number of oocysts at a given time within an individual infected

257   mosquito, $Y(t)$, we track the number of G-stage parasites, from the initial load n, that have

258   developed by a given time, which follows a binomial distribution,

259
$$Y(t) \sim B\left(n, Q(\alpha_{GO}, 0, \beta_{GO} t)\right), \qquad (2.05)$$

where $Q(\alpha_{GO}, 0, \beta_{GO} t)$ is the cumulative probability that an individual parasite has

developed into an oocyst by time t.

Mosquitoes die throughout the course of the experiments, which we explicitly model

via a Cox regression model. We estimate the probability a mosquito is alive immediately

prior to time t, $A(t)$, by fitting this model to the mosquito survival data (Table 1) and allow

for differences in the survival due to infection by comparing groups of mosquitoes exposed

to uninfected and infected blood and allowing the hazard to vary with infection status.

Mosquitoes exposed to infected blood may not actually become infected, so this group

consists of a mix of infected and uninfected individuals. By fitting the complete model, we

account for this to estimate infection-specific differences in mortality, see S7.1.1 for full

details. A previous analysis of a subset of the data determined that the Gompertz

distribution, in which the mosquito mortality rate increases with age (senescence) (14), was

the best fit, and we use this distribution here.

2.2.3   Observation level

During the experiments, samples of mosquitoes are dissected at particular time intervals.

From these samples, two possible measurements are recorded: (1) the aggregate count of

parasite-positive mosquitoes (either oocysts or sporozoites) and (2) the number of oocysts

counted in each of the dissected mosquitoes.

We model the aggregate count of positive mosquito data thus. At each time, a

sample of mosquitoes, $D(t)$, are dissected, likely comprising a mix of infected and

uninfected individuals. The number of infected mosquitoes within the sample, $I(t)$, is

modelled as a binomial random variable,

$$I(t) \sim B\big(D(t), \delta R(t)\big), \tag{2.06}$$

284 where δ is the probability successful infection occurs during the infectious blood-feeding,

285 and $R(t)$ is the fraction of infected mosquitoes alive. $R(t)$ is the ratio of survival

286 probabilities for infected (E) and uninfected (U) mosquitoes:

$$R(t) = \frac{A_E(t)}{A_U(t)}. \tag{2.07}$$

Note, we assume that once a mosquito is infected it will remain so for the duration of its

lifespan (45).

A mosquito may be successfully infected but no parasites observed during dissection, if

insufficient time has passed since blood-feeding. If n was known for each dissected

mosquito, the probability dissection would detect parasites of a given life stage, at time t, is

given by eq. (2.04). In reality, n is not known. We model this uncertainty using eq. (2.03) and

incorporate it into the probability any infected mosquito has observable parasites by

marginalising n out of the joint distribution,

296
$$\Pr(t \geq T^{min}) = \sum_n \Pr(t \geq T_{1:n}^{min}|n)\Pr(n) \qquad (2.08)$$

297
$$= E\big(\Pr(t \geq T_{1:n}^{min}|n)\big)$$

298
$$= E\big(1 - \big(1 - Q(\alpha, 0, \beta t)\big)^n\big)$$

299
$$= \frac{1 - k^k(k + \mu Q(a, 0, \beta t))^{-k}}{1 - (\frac{k}{k + \mu})^k}$$

300   where T^{min} is the time when first parasite of the given life stage appears (O or S), and $E_n(.)$

301   denotes the expectation with respect to the distribution of n (eq. (2.03)).

302   The count of infected mosquitoes in which the parasite life stage of interest (O or S)

303   is observed, $X(t)$, given a sample of infected mosquitoes, $I(t)$, is then also binomially

304   distributed,

305
$$X(t) \sim B\big(I(t), \Pr(t \geq T^{min})\big). \qquad (2.09)$$

306   Within a given sample of mosquitoes, the number of successfully infected mosquitoes, $I(t)$,

307   is not known – we only observe the count of observed infected individuals, $X(t)$, and the

308   total number dissected, $D(t)$. We incorporate this uncertainty in $I(t)$ by marginalising this

309   quantity out of the joint distribution defined in eqs. (2.06) and (2.09), resulting in the

310   following binomial distribution describing the observed count,

311
$$X(t) \sim B(D(t), \delta R(t)\Pr(t \geq T^{min})). \qquad (2.10)$$

We next detail how our model describes oocyst counts, $Y(t)$. A particular mosquito

may yield a zero count when dissected either if it was not successfully infected or if

insufficient time has elapsed for any oocysts to develop. The probability a sampled

individual is successfully infected is $\delta R(t)$. The count of oocysts to have developed by a

given time within a successfully infected mosquito is described by eq. (2.05): this probability

distribution depends on n (the number of G-stage parasites) – an unknown quantity. To

derive the sampling distribution of oocyst counts, we marginalise n out of the joint

distribution assuming its uncertainty is described by eq. (2.03). Combining this with our

underlying uncertainty regarding the mosquito's infection status results in the following

expression describing the probability of counting Y oocysts at time t, in a dissected

mosquito,

$Pr(Y(t)|Q_{GO}(t), \delta, R(t), t, \mu, k)$

$$
= \begin{cases}
(1 - \delta R(t)) + \delta R(t)\left(\dfrac{\left(\frac{k}{k+\mu}\right)^k - \left(\frac{k}{k+Q_{GO}(t)\mu}\right)^k}{-1 + \left(\frac{k}{k+\mu}\right)^k}\right) & , if\ Y = 0 \\[2em]
\delta R(t) \dfrac{\left(k^k (Q_{GO}(t)\mu)^Y (k + Q_{GO}(t)\mu)^{(-Y-k)} \binom{-1+x+k}{-1+x}\right)}{1 - \left(\frac{k}{k+\mu}\right)^k} & , if\ Y \geq 1.
\end{cases}
$$

(2.11)

where, for notational convenience, we term $Q_{GO}(t) = Q(\alpha_{GO}, 0, \beta_{GO} t)$. From eq. (2.11), the

mean oocyst intensity within the population is given by $\delta R(t) \frac{Q_{GO}(t)\mu}{1-(\frac{k}{k+\mu})^k}$. Mosquitoes are

typically dissected for oocysts before any burst, so we do not model how bursting would

lead to a decline in oocyst prevalence or intensity.

2.3   Incorporating temperature-dependence

First, we fit the model to data collected under standard insectary conditions (27 °C), which

included the complete range of oocyst intensity, oocyst prevalence, sporozoite prevalence

and survival (grouped by infectious blood-fed and control) data. Next, to investigate the

impact of temperature on the EIP, we fit the model to the data collected at each other

temperature, which included sporozoite prevalence and survival data only. Due to the lack

of oocyst data, development times from day of blood-feeding to sporozoite were estimated

as a single gamma distribution with shape, α_{GS}, and rate, β_{GS}. Data collected at 33°C was

excluded as mosquitoes were only sampled on a single day. We refer to these individual

temperature model fits collectively as the "single temperature models".

To estimate a functional relationship between temperature and EIP, we fit the model

to all the data simultaneously (the "all temperature model"). To do so, we first plotted the

α_{GS} and β_{GS} parameters of the single temperature models against temperature (Figure S1).

This indicated an approximate linear relationship between temperature and the

development rate, β_{GS}. Laboratory studies have found the early part of sporogony of

laboratory strains of *P. falciparum*, in *A. stephensi*, is most temperature sensitive (40,41), so

we assumed the temperature variation affected the rate at which parasites develop from *G*

to *O*. The development rate was then modelled as a linear function of temperature, *C*,

347
$$\beta_{GO} = m_\beta C + c_\beta. \tag{2.12}$$

348   The infection of mosquitoes has been found to decline at high temperatures, in the

349   laboratory, due to factors such as the impact of temperature on vector immunity (27) and

350   the establishment of gametocytes (29). To account for this, we included a relationship

351   between temperature and the human-to-mosquito transmission probability, δ, modelled as

352   a logit-linear function,

353
$$\delta = \frac{1}{1 + e^{-(m_\delta C + c_\delta + \rho_i)}}. \tag{2.13}$$

354   To capture differences between membrane feeding assays (38), we included a hierarchical

355   term, ρ_i, in eq. (2.13) where,

356
$$\rho_i \sim N(0, \sigma_\delta), \tag{2.14}$$

and i refers to an individual experiment, defined as any unique combination of study and

temperature treatments (i.e. the combination of reference and adult mosquito housing

temperature values in Table 1).

Kaplan-Meier plots of mosquito survivorship and existing studies indicated that

laboratory *A. stephensi* survival depended on temperature (Figure S2). To account for this, a

temperature-dependent mortality term was incorporated into the Cox model hazard,

$$h_0(t)e^{(\beta_C C + \beta_E E + \varepsilon_i)}, \tag{2.15}$$

364 where C is temperature, E is the infection status ($E = 1$ indicates successful infection; and

365 $E = 0$ for uninfected individuals), and $h_0(t)$ is the baseline hazard modelled by a Gompertz

366 distribution. To account for experimental heterogeneity in mosquito survival between

367 experiments, a hierarchical error term, ε_i, was added to eq. (2.15) where,

$$\varepsilon_i \sim N(0, \sigma_{survival}). \tag{2.16}$$

To facilitate model fitting, we scaled the temperature values such that temperature

was centred around 0 with a standard deviation of 1.

2.4   Model fitting

The models were fit under a Bayesian framework, using the probabilistic programming

language Stan (46), which implements the No-U-turn Markov chain Monte Carlo (MCMC)

sampler (47). We specified weakly informative priors following a literature search (Table S1).

For each model fit, four chains were run, with 1500 warmup iterations and 4500 iterations

in total. Convergence was determined by $\hat{R} < 1.01$ for all parameters, and bulk and tail

effective sample sizes (ESS; an estimate of sampling efficiency) greater than 400 (see Table

S2 for these values). We visually examined the trace plots (Figure S3) and illustrate the

difference between prior and posterior distributions in Figure S4.

2.5 Model investigation

To compare our estimates with the predominant approach, we estimated the EIP and

human-to-mosquito transmission probability using a logistic model (14,15,18,27,48), which

we fit using a non-linear least squares approach (see supplementary information 7.1.3).

Given a decline in the observed sporozoite prevalence at later time points, the data was

subset to include only data points before the time of peak observed sporozoite prevalence,

as has been done previously (14).

Data did not exist to investigate the impact of parasite load experimentally. We

therefore used our fitted model to simulate the temporal dynamics of sporogony with

different mean parasite loads (among infected mosquitoes), holding all other parameters at

their mean posterior values for 27°C (sensitivity analysis).

3 Results

3.1 Temporal dynamics of sporogony

We first considered the observed temporal dynamics of sporogony at a single temperature

(27°C; standard insectary conditions). All values we report are the posterior predictive

means and the credible intervals are the 95% central posterior estimates; times reported

are the number of days post infectious blood-feed. From the model fit, we estimate that

oocyst prevalence was 10% at 2.2 days (CI: 2.1–2.3 days) and peaked at 69% (CI: 68–71%)

after approximately 5.9 days (CI: 5.8–6.0 days) (Figure 2a). The modelled oocyst intensity

peaked at 2.0 (CI: 1.9–2.1) oocysts per mosquito, shortly after oocyst prevalence peaked

(Figure 2b). In the raw data, sporozoite-positive mosquitoes were first observed on day 10

when the sporozoite prevalence was 1.4%; our model estimates are slightly earlier: the

modelled sporozoite prevalence reached 10% at 9.5 days (CI: 9.3–9.7 days) (Figure 2c). The

time required for sporozoites to appear varied between mosquitoes, and, only after 15.9

days (CI: 14.2–16.0 days) did the sporozoite prevalence peak at 63% (CI: 61–65%).

At the individual parasite level, we consider the time taken for each modelled

parasite life stage transition to occur: the mean time taken was 3.4 days (CI: 3.3–3.6 days)

for G to O, 9.2 days (CI: 8.9–9.5 days) for O to S and 12.6 days (CI: 12.3–12.9 days) for G to S

(Figure S5a). The majority of individual parasites (99.5%) had undergone each of these

transitions at 6.1 days (CI: 5.6–6.6 days) for G to O; 14.7 days (CI: 13.9–15.5 days) for O to S;

and 18.5 days (CI: 17.8–19.3 days) for G to S (Figure S5b).

3.2   Impact of temperature

We next considered the impact of temperature on sporogony. The *single temperature*

models fitted the sporozoite prevalence data well across the entire range of temperatures

(Figure S6). *The all temperature model* fitted the sporozoite prevalence data well at lower

temperatures (between 21°C and 30°C); at higher temperatures, the model tended to

overstate the EIP (Figure 3). The survival of both infectious blood-fed and control

mosquitoes was only available at 27°C, 30°C and 33°C, and consequently the *single*

temperature model survival parameters had greater freedom to vary. Indeed, the *all*

temperature model (Figure S7) fits to the survival data were less variable and more

predictable across different temperatures than the *single temperature models* (Figure S8).

Consequently, here we provide the *all temperature model* (Figure 3, Figure S9) results due

to their greater generalisability.

For the *all temperature model*, infection resulted in a higher risk of mosquito death,

with a hazard odds ratio of 1.65 (CI: 1.47–1.86). Increases in temperature elevated mosquito

mortality with a hazard odds ratio of 1.57 (CI: 1.33-1.86) per 3.5°C change. At 10 days post-

infection, the probability of an infected mosquito being alive, *A(t)*, was 0.91 (CI: 0.87–0.94)

at 21°C versus 0.67 (CI: 0.58–0.75) at 34°C, with similar differences for uninfected

individuals.

At higher temperatures, fewer mosquitoes develop sporozoites, but those that do,

do so faster. We estimate that, between 21°C and 34°C, increases in temperature reduced

the EIP: EIP_{10} fell from 12.9 days (CI: 12.5–13.3 days) to 6.9 days (CI: 6.7–7.1 days), EIP_{50}

declined from 16.1 days (CI: 15.5–16.8) to 8.8 days (CI: 8.6–8.9 days) and EIP_{90} fell from 20.4

days (CI: 19.4–21.8 days) to 11.3 days (CI: 11.0–11.7 days) (Figure 4a). Table S3 provides the

EIP percentiles at all temperatures. Increases in temperature reduced the human-to-

mosquito transmission probability, which fell from 84% (CI: 76–89%) at 21°C to 42% (CI: 34–

50%) at 34°C (Figure 4).

3.3   Key differences in transmission parameters estimated using mSOS

Next, we compared our estimates of two important malaria transmission parameters – the

EIP and human-to-mosquito transmission probability – to those obtained using the

predominant literature approach based on logistic regression (logistic model fits are shown

Figure S10). In our framework, we can disentangle the development time of parasites from

other underlying processes, specifically mosquito mortality induced by malaria infection,

which can influence the observed sporozoite prevalence. In doing so, we quantify the EIP

distribution as the time taken for a given percentile of the infected mosquitoes to display

sporozoites, in the absence of other mechanisms that determine the observed sporozoite

prevalence. This is given by the time at which the cumulative probability an infected

mosquito develops any salivary gland sporozoites, $\Pr(T_1 \leq t)$, reaches a given value. Using

this approach, across all temperatures, the variation in the EIP distribution is greater than

the equivalent logistic model estimates (Figure 4a). At 27°C, for example, our *single*

temperature model estimated an EIP_{10}–EIP_{90} range of 9.2–13.7 days compared to 10.3–11.2

days estimated by the logistic approach.

The mSOS estimates of the human-to-mosquito transmission probability were

consistently higher than the equivalent values estimated by the logistic model (Figure 4b).

The 27°C *single temperature* model estimate, for example, was 70% (CI: 68–72%) as

opposed to the logistic growth model estimate of 60%. This is because, in the laboratory

data we analysed, infected mosquitoes died at an elevated rate, meaning that the

proportion of infected mosquitoes declined with time and the observed peak of sporozoite

prevalence is not representative of the initial proportion of infected mosquitoes. Not

accounting for these mechanisms results in an underestimate of the human-to-mosquito

transmission probability.

3.4   Sensitivity analysis of the mean parasite load

Within our model, the time-taken for each simulated parasite to develop is both

independent and stochastic. At each time point, whether or not that parasite has yet

developed can be viewed as a toss of a coin: the more coins are tossed, the more likely it is

that one will land on heads, or equivalently a parasite will have developed, by chance.

Higher parasite loads then lead to earlier rises in parasite prevalence. Figure 5A shows the

simulated sporozoite prevalence over time across populations with different mean parasite

load parameter values, which indicates within the model greater parasite numbers cause

the prevalence to peak earlier. Figure 5B shows the resultant EIP quantiles as a function of

parasite load: increasing the mean parasite load from 1.7 to 25.0 reduced the modelled

EIP_{50} from 10.9 to 8.0 days.

Figure 2 Single temperature 27°C model fit to the oocyst and sporozoite data. The panels show our model fit

to the 27°C dataset: panel A to the oocyst prevalence, panel B to the oocyst intensity data and panel C to the

sporozoite prevalence. (A & C) Points: parasite prevalence of the laboratory mosquito data (95% confidence

intervals are given by the point range). The grey shaded area represents the 95% credible interval of the model

posterior predictive means, the median posterior predictive mean is shown by the black line. (B) The points

show the mean parasite load among all blood fed mosquitoes (intensity), the point range indicates the 2.5%–

97.5% quantiles of the raw data. The shaded area represents the 2.5%–97.5%-quantiles of the negative

binomial distribution; where the location and overdispersion parameters at set to their posterior means.

Figure 3 Model fits to sporozoite prevalence data across all temperatures. These fits were generated by

fitting a single model to all temperatures simultaneously ("all temperature model"), with the functional form

of temperature as described in eqs. (2.11) and (2.12). Black points: parasite prevalence of the laboratory

mosquito data (95% confidence intervals are given by the vertical black lines). The grey shaded area represents

the 95% quantiles of the posterior predictive means; the black lines represent the median posterior predictive

means.

Figure 4 Effect of temperature on malaria transmission parameters. Panel A shows the model impact of temperature on the EIP quantiles (as indicated in legend); Panel B shows its impact on the human-to-mosquito transmission probability. In both panels, the lines show impact as estimated by the *all temperature* mSOS model with 95% posterior intervals indicated by shading; the discrete round points show independent estimates of the *single temperature* mSOS model at each temperature, 95% posterior intervals are shown by the vertical lines. The number of iterations used to calculate the EIP plot were thinned to every 5^{th} iteration for efficiency. The discrete triangle points show the logistic growth model parameter estimates at each temperature.

Figure 5 Modelled impact of parasite load on the extrinsic incubation period. Panel A shows the impact of varying the mean parasite load of infected mosquitoes on the temporal dynamics of sporozoite prevalence in a sensitivity analysis; panel B summarises how the EIP is affected by the same parameter in the sensitivity analysis. All other parameters were held constant at their mean posterior values.

4   Discussion

Membrane feedings assays form the bulk of experiments used to determine key

parasitological parameters of malaria transmission, but despite the seeming simplicity of

these experiments, numerous hidden processes contribute to the observed data. Here, we

applied a systems biology approach to develop a multiscale model of the temporal dynamics

of sporogony (mSOS) that explicitly account for these underlying processes. To illustrate its

use, we fitted mSOS to experimental data and demonstrated its predictive accuracy across a

range of experimental protocols. In doing so, we estimated two important determinants of

malaria transmission intensity – the EIP and human-to-mosquito transmission probability.

By adopting a mechanistic approach, our EIP estimates are defined in terms of the

underlying biology rather than characteristics of the raw experimental data. Our estimates

indicate greater variation in the EIP than previously thought and highlight the importance of

accounting for this variation when making epidemiological predictions (see also (15)). This

variation could be included by embedding mSOS within transmission dynamics models of

malaria.

The influence of parasite-induced mosquito mortality is still a source of debate, and

likely to vary depending on the parasite-vector system (26,36). Nonetheless, accounting for

potential differences in mortality is essential to avoid bias in estimates of both EIP and the

human-to-mosquito transmission probability. By modelling mosquito infection and survival

in a single framework we could estimate malaria infection induced differences in mosquito

survival, whilst accounting for mosquitoes that do not develop oocysts within the infectious

blood fed group. At higher temperatures, we found that these differences only partly

explained the sporozoite prevalence data. Another study observed no difference in the

survivorship of wild caught *A. gambiae* mosquitoes due to malaria infection, yet

P. falciparum detection did decline over time (49). So, it is possible that mosquito

survivorship was not behind these observed declines and alternative explanations, such as

sporozoite loss during sugar-feeding (50,51) or sporozoite mortality could require further

investigation.

Temperature is an important predictor of *P. falciparum* prevalence among humans,

its impact is, however, often non-linear and location-specific (52–54). Differences in the EIP

due to temperature, parameterised using laboratory data, are commonly used in the

prediction of the spatiotemporal limits and endemicity of malaria transmission (32,55,56).

Over the temperature range investigated, our estimates qualitatively match those of (15),

which determined that the EIP falls with increases in temperature, and that the incremental

change is greatest at lower temperatures. In a semi-field setting, increases in temperature,

due to the interaction between deforestation and altitude, similarly decreased the EIP of

wild *P. falciparum* (19), but very little field data exists. To more accurately simulate the EIP

in the field will require consideration of the impact of the time of biting (57), diurnal

temperature fluctuations (58) and mosquito resting location on the temperatures parasites

are exposed to during different life stages. The degree to which the EIP of locally adapted

strains differ to the laboratory strains must also be considered (59). Current estimates

indicate climate change has the potential to change the number of people at risk of malaria

(56,60,61), though there is uncertainty due to the complexity of how the parasite and

mosquito will interact with the local changing environment in the long term. Characterising

spatiotemporal heterogeneity in the EIP in the field is nonetheless critical to assess our

current understanding of malaria and its control as well as the changing risk of malaria

resulting from the interactions between climatic, land-use (62) and socio-economic factors

(63).

Within our model, heterogeneity in the EIP emerges from differences in the

development times of individual parasites. Density-dependent processes (64,65) and

intraspecific variation in individual mosquito characteristics, such as body size (66), may also

impact the observed sporozoite prevalence. To include such processes in the model in a

data-driven way requires higher resolution data to differentiate the impact of different

hypotheses. To parameterise mSOS, we collated data from four previously published

experimental infections. We therefore recommend that future studies dissect mosquitoes

for both oocysts and sporozoites across a greater range of times post infection. It would also

be useful if mosquito mortality were recorded, ideally determining the past infection status

of carcases through molecular methods.

The work shows how higher parasite loads within a mosquito might decrease the

EIP. To our knowledge, this possible relationship between parasite load and development

time has not been considered elsewhere but could have potentially important

epidemiological consequences as it indicates that onwards transmission may be more

efficient from more infectious people. Further work is needed to explore this hypothesis

given that our model assumes the EIP is determined on a per-oocyst basis and that

developmental times are independent within an individual mosquito which needs to be

verified experimentally. What's more, other processes may be operating: emerging

experimental evidence hints that a decrease in resource availability per parasite (at high

parasite loads), as determined by the parasite load and the number of times the mosquito

blood-feeds, may decrease the parasite development rate (48,67). Since (a) transmission is

highly sensitive to changes in EIP, (b) transmission blocking interventions cause a decline in

oocyst intensity (38), and (c) parasite load in the field may be higher than previously thought

(37), the influence of parasite load on EIP merits further investigation.

It is now well over a century since *Plasmodium* parasites were first uncovered in

dissected anopheline mosquitoes. Today, we remain dependent on dissection for

understanding the parasite lifecycle in mosquitoes, and there remains much to learn. The

model we introduce here provides a new way to parse dissection data to probe the

underlying biology. Our model can be extended to incorporate more fine scale

characteristics of parasite ecology, but to do so in a principled manner requires more fine

scale data. As such, we foresee a great necessity and opportunity for closer collaboration

between experimentalists and modellers in the future.

5   Acknowledgements.

We would like to thank Professor Matthew Thomas for useful insight and the Thomas lab at

Penn State University for producing and publishing the high-quality data upon which our

study is based.

IJS would like to thank the Natural Environment Research Council (NERC) for PhD

funding administered through the Quantitative Methods in Ecology and Evolution (QMEE)

Centre for Doctoral Training (grant reference: NE/P012345/1). The authors would like to

acknowledge the MRC Centre for Global Infectious Disease Analysis (grant reference:

MR/R015600/1), this award is jointly funded by the UK Medical Research Council (MRC) and

the UK Department for International Development (DFID) under the MRC/DFID Concordat

agreement and is also part of the EDCTP2 programme supported by the European Union.

6   References

1.   World Health Organization. World malaria report 2019 [Internet]. Geneva; 2019.

2.   Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, et al. The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. Nature. 2015 Sep 16;526:207.

3.   Macdonald G. Epidemiological basis of malaria control. Bull World Health Organ. 1956;15(3–5):613.

4.   Macdonald G. The analysis of the sporozoite rate. Trop Dis Bull. 1952;49(6).

5.   Smith DL, Ellis McKenzie F. Statics and dynamics of malaria infection in Anopheles mosquitoes. Malar J. 2004;3(1):13.

6.   Vaughan JA. Population dynamics of Plasmodium sporogony. Trends Parasitol. 2007;23(2):63–70.

7.   Pichon G, Awono-Ambene HP, Robert V. High heterogeneity in the number of Plasmodium falciparum gametocytes in the bloodmeal of mosquitoes fed on the same host. Parasitology. 2000;121(2):115–20.

8.   Arai M, Billker O, Morris HR, Panico M, Delcroix M, Dixon D, et al. Both mosquito-derived xanthurenic acid and a host blood-derived factor regulate gametogenesis of Plasmodium in the midgut of the mosquito. Mol Biochem Parasitol. 2001;116(1):17–24.

9.   Billker O, Lindo V, Panico M, Etienne AE, Paxton T, Dell A, et al. Identification of xanthurenic acid as the putative inducer of malaria development in the mosquito. Nature. 1998;392(6673):289–92.

10.   Kuehn A, Pradel G. The coming-out of malaria gametocytes. J Biomed Biotechnol. 2010/01/05. 2010;2010:976827.

11.   Clayton AM, Dong Y, Dimopoulos G. The Anopheles innate immune system in the defense against malaria infection. J Innate Immun. 2013/08/28. 2014;6(2):169–81.

12.   Gouagna LC, Mulder B, Noubissi E, Tchuinkam T, Boudin C, Boudin C. The early sporogonic cycle of Plasmodium falciparum in laboratory-infected Anopheles gambiae: an estimation of parasite efficacy. Trop Med Int Heal. 1998 Jan 1;3(1):21–8.

13.   Vaughan JA, Noden BH, Beier JC. Population dynamics of Plasmodium falciparum sporogony in laboratory-infected Anopheles gambiae. J Parasitol. 1992;78(4):716–24.

14.   Shapiro LLM, Whitehead SA, Thomas MB. Quantifying the effects of temperature on mosquito and parasite traits that determine the transmission potential of human malaria. PLOS Biol. 2017 Oct 16;15(10):e2003489.

15.   Ohm JR, Baldini F, Barreaux P, Lefevre T, Lynch PA, Suh E, et al. Rethinking the extrinsic incubation period of malaria parasites. Parasit Vectors. 2018;11(1):178.

16.   Nikolaev BP. On the influence of temperature on the development of malaria plasmodia in the mosquito. Leningr Pasteur Inst Epidemiol Bacteriol. 1935;2:108–9.

17.   Hien DFDS, Dabiré KR, Roche B, Diabaté A, Yerbanga RS, Cohuet A, et al. Plant-Mediated Effects on Mosquito Capacity to Transmit Human Malaria. PLoS Pathog. 2016 Aug 4;12(8):e1005773–e1005773.

18.   Shapiro LLM, Murdock CC, Jacobs GR, Thomas RJ, Thomas MB. Larval food quantity affects the capacity of adult mosquitoes to transmit human malaria. Proc R Soc B Biol Sci. 2016 Jul 13;283(1834):20160298.

19.   Afrane YA, Little TJ, Lawson BW, Githeko AK, Yan G. Deforestation and vectorial

capacity of Anopheles gambiae Giles mosquitoes in malaria transmission, Kenya. Emerg Infect Dis. 2008 Oct;14(10):1533–8.

20. Kærn M, Elston TC, Blake WJ, Collins JJ. Stochasticity in gene expression: from theories to phenotypes. Nat Rev Genet. 2005;6(6):451–64.

21. Paul REL, Bonnet S, Boudin C, Tchuinkam T, Robert V. Aggregation in malaria parasites places limits on mosquito infection rates. Infect Genet Evol. 2007;7(5):577–86.

22. Da DF, Churcher TS, Yerbanga RS, Yaméogo B, Sangaré I, Ouedraogo JB, et al. Experimental study of the relationship between Plasmodium gametocyte density and infection success in mosquitoes; implications for the evaluation of malaria transmission-reducing interventions. Exp Parasitol. 2015;149:74–83.

23. Yassine H, Osta MA. Anopheles gambiae innate immunity. Cell Microbiol. 2010 Jan 1;12(1):1–9.

24. Dong Y, Manfredini F, Dimopoulos G. Implication of the mosquito midgut microbiota in the defense against malaria parasites. PLoS Pathog. 2009/05/08. 2009 May;5(5):e1000423–e1000423.

25. Blanford S, Chan BHK, Jenkins N, Sim D, Turner RJ, Read AF, et al. Fungal Pathogen Reduces Potential for Malaria Transmission. Science (80-). 2005 Jun 10;308(5728):1638 LP – 1641.

26. Ferguson HM, Read AF. Why is the effect of malaria parasites on mosquito survival still unresolved? Trends Parasitol. 2002;18(6):256–61.

27. Paaijmans KP, Blanford S, Chan BHK, Thomas MB. Warmer temperatures reduce the vectorial capacity of malaria mosquitoes. Biol Lett. 2011/12/21. 2012 Jun 23;8(3):465–8.

28. Murdock CC, Paaijmans KP, Bell AS, King JG, Hillyer JF, Read AF, et al. Complex effects of temperature on mosquito immune function. Proceedings Biol Sci. 2012/05/16. 2012 Aug 22;279(1741):3357–66.

29. Pathak AK, Shiau JC, Thomas MB, Murdock CC. Field Relevant Variation in Ambient Temperature Modifies Density-Dependent Establishment of Plasmodium falciparum Gametocytes in Mosquitoes [Internet]. Vol. 10, Frontiers in Microbiology . 2019. p. 2651.

30. Detinova TS, Bertram DS, Organization WH. Age-grouping methods in Diptera of medical importance, with special reference to some vectors of malaria. World Health Organization; 1962.

31. Paaijmans KP, Read AF, Thomas MB. Understanding the link between malaria risk and climate. Proc Natl Acad Sci. 2009 Aug 18;106(33):13844 LP – 13849.

32. Mordecai EA, Paaijmans KP, Johnson LR, Balzer C, Ben-Horin T, Moor E, et al. Optimal temperature for malaria transmission is dramatically lower than previously predicted. Ecol Lett. 2012 Oct 11;16(1):22–30.

33. Teboh-Ewungkem MI, Yuster T. A within-vector mathematical model of Plasmodium falciparum and implications of incomplete fertilization on optimal gametocyte sex ratio. J Theor Biol. 2010;264(2):273–86.

34. Childs LM, Prosper OF. Simulating within-vector generation of the malaria parasite diversity. PLoS One. 2017 May 22;12(5):e0177941–e0177941.

35. Gog JR, Pellis L, Wood JLN, McLean AR, Arinaminpathy N, Lloyd-Smith JO. Seven challenges in modeling pathogen dynamics within-host and across scales. Epidemics. 2015;10:45–8.

36. Murdock CC, Sternberg ED, Thomas MB. Malaria transmission potential could be

reduced with current and future climate change. Sci Rep. 2016 Jun 21;6:27771.

37. Bompard A, Da DF, Yerbanga SR, Morlais I, Awono-Ambéné PH, Dabiré RK, et al. High Plasmodium infection intensity in naturally infected malaria vectors in Africa. Int J Parasitol. 2020;

38. Churcher TS, Blagborough AM, Delves M, Ramakrishnan C, Kapulu MC, Williams AR, et al. Measuring the blockade of malaria transmission – An analysis of the Standard Membrane Feeding Assay. Int J Parasitol. 2012;42(11):1037–44.

39. Miura K, Swihart BJ, Deng B, Zhou L, Pham TP, Diouf A, et al. Transmission-blocking activity is determined by transmission-reducing activity and number of control oocysts in Plasmodium falciparum standard membrane-feeding assay. Vaccine. 2016/06/29. 2016 Jul 29;34(35):4145–51.

40. Noden BH, Kent MD, Beier JC. The impact of variations in temperature on early Plasmodium falciparum development in Anopheles stephensi. Parasitology. 2009/04/06. 1995;111(5):539–45.

41. Eling W, Hooghof J, Van De Vegte-Bolmer M, Sauerwein R, Van Gemert GJ. Tropical temperatures can inhibit development of the human malaria parasite Plasmodium falciparum in the mosquito. In: Proceedings of the Section Experimental and Applied Entomology-Netherlands Entomological Society. 2001. p. 151–6.

42. Stewart T, Strijbosch LWG, Moors H, Batenburg P van. A simple approximation to the convolution of gamma distributions. 2007;

43. Stone WJR, Eldering M, van Gemert G-J, Lanke KHW, Grignard L, van de Vegte-Bolmer MG, et al. The relevance and applicability of oocyst prevalence as a read-out for mosquito feeding assays. Sci Rep. 2013 Dec 4;3:3418.

44. David HA, Nagaraja HN. Order statistics. Encycl Stat Sci. 2004;

45. Beier JC. Malaria parasite development in mosquitoes. Annu Rev Entomol. 1998;43(1):519–43.

46. Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, et al. Stan: A probabilistic programming language. J Stat Softw. 2017;76(1).

47. Hoffman MD, Gelman A. The No-U-Turn sampler: adaptively setting path lengths in Hamiltonian Monte Carlo. J Mach Learn Res. 2014;15(1):1593–623.

48. Shaw WR, Holmdahl IE, Itoe MA, Werling K, Marquette M, Paton DG, et al. Multiple blood feeding in mosquitoes shortens the Plasmodium falciparum incubation period and increases malaria transmission potential. bioRxiv. 2020 Jan 1;2020.03.24.991356.

49. Chege GMM, Beier JC. Effect of Plasmodium falciparum on the Survival of Naturally Infected Afrotropical Anopheles (Diptera: Culicidae). J Med Entomol. 1990 Jul 1;27(4):454–8.

50. Brugman VA, Kristan M, Gibbins MP, Angrisano F, Sala KA, Dessens JT, et al. Detection of malaria sporozoites expelled during mosquito sugar feeding. Sci Rep. 2018;8(1):7545.

51. Billingsley PF, Hodivala KJ, Winger LA, Sinden RE. Detection of mature malaria infections in live mosquitoes. Trans R Soc Trop Med Hyg. 1991;85(4):450–3.

52. Snow RW, Gouws E, Omumbo J, Rapuoda B, Craig MH, Tanser FC, et al. Models to predict the intensity of Plasmodium falciparum transmission: applications to the burden of disease in Kenya. Trans R Soc Trop Med Hyg. 1998 Nov 1;92(6):601–6.

53. Gosoniu L, Vounatsou P, Sogoba N, Maire N, Smith T. Mapping malaria risk in West Africa using a Bayesian nonparametric non-stationary model. Comput Stat Data Anal.

2009;53(9):3358–71.

54. Reiner RC, Geary M, Atkinson PM, Smith DL, Gething PW. Seasonality of Plasmodium falciparum transmission: a systematic review. Malar J. 2015;14(1):343.

55. Gething PW, Van Boeckel TP, Smith DL, Guerra CA, Patil AP, Snow RW, et al. Modelling the global constraints of temperature on transmission of Plasmodium falciparum and P. vivax. Parasit Vectors. 2011;4(1):92.

56. Ryan SJ, Lippi CA, Zermoglio F. Shifting transmission risk for malaria in Africa with climate change: a framework for planning and intervention. Malar J. 2020;19(1):170.

57. Suh E, Grossman MK, Waite JL, Dennington NL, Sherrard-Smith E, Churcher TS, et al. The influence of feeding behaviour and temperature on the capacity of mosquitoes to transmit malaria. Nat Ecol Evol. 2020;

58. Paaijmans KP, Blanford S, Bell AS, Blanford JI, Read AF, Thomas MB. Influence of climate on malaria transmission depends on daily temperature variation. Proc Natl Acad Sci. 2010 Aug 24;107(34):15135 LP – 15139.

59. Mordecai EA, Caldwell JM, Grossman MK, Lippi CA, Johnson LR, Neira M, et al. Thermal biology of mosquito-borne disease. Ecol Lett. 2019 Oct 1;22(10):1690–708.

60. Tanser FC, Sharp B, le Sueur D. Potential effect of climate change on malaria transmission in Africa. Lancet. 2003;362(9398):1792–8.

61. Siraj AS, Santos-Vega M, Bouma MJ, Yadeta D, Carrascal DR, Pascual M. Altitudinal Changes in Malaria Incidence in Highlands of Ethiopia and Colombia. Science (80-). 2014 Mar 7;343(6175):1154 LP – 1158.

62. Franklinos LH V, Jones KE, Redding DW, Abubakar I. The effect of global change on mosquito-borne disease. Lancet Infect Dis. 2019;19(9):e302–12.

63. Gething PW, Smith DL, Patil AP, Tatem AJ, Snow RW, Hay SI. Climate change and the global malaria recession. Nature. 2010;465(7296):342–5.

64. Sinden RE, Dawes EJ, Alavi Y, Waldock J, Finney O, Mendoza J, et al. Progression of Plasmodium berghei through Anopheles stephensi is density-dependent. PLoS Pathog. 2007 Dec 28;3(12):e195–e195.

65. Mendes AM, Awono-Ambene PH, Nsango SE, Cohuet A, Fontenille D, Kafatos FC, et al. Infection intensity-dependent responses of Anopheles gambiae to the African malaria parasite Plasmodium falciparum. Infect Immun. 2011;79(11):4708–15.

66. Lyimo EO, Koella JC. Relationship between body size of adult Anopheles gambiae sl and infection with the malaria parasite Plasmodium falciparum. Parasitology. 1992;104(2):233–7.

67. Habtewold T, Sharma AA, Wyer CAS, Masters EKG, Windbichler N, Christophides GK. Plasmodium oocysts respond with dormancy to crowding and nutritional stress. bioRxiv. 2020 Jan 1;2020.03.07.981951.

68. Pringle G. A quantitative study of naturally-acquired malaria infections in Anopheles gambiae and Anopheles funestus in a highly malarious area of East Africa. Trans R Soc Trop Med Hyg. 1966;60(5):626–32.

69. Rosenberg R, Andre RG, Somchit L. Highly efficient dry season transmission of malaria in Thailand. Trans R Soc Trop Med Hyg. 1990 Jan 1;84(1):22–8.

70. Taylor LH. Infection rates in, and the number of Plasmodium falciparum genotypes carried by Anopheles mosquitoes in Tanzania. Ann Trop Med Parasitol. 1999 Sep 1;93(6):659–62.

71. Billingsley PF, Medley GF, Charlwood D, Sinden RE. Relationship between prevalence and intensity of Plasmodium falciparum infection in natural populations of Anopheles

mosquitoes. Am J Trop Med Hyg. 1994;51(3):260–70.

7 Supplementary Information

7.1 Supplementary methods

7.1.1 Mosquito survival (A(t))

To estimate the difference in mosquito survival due to infection status, E, we model the mosquito survival data (see Table 1) using the Cox proportional hazards model as follows. The Kaplan-Meier survival curves of the mosquito survival data (Figure S2) show that the proportion of mosquitoes alive does not decline exponentially indicating that the mosquito mortality rate is not constant. Indeed, a previous analysis of a subset of the data found the best fitting survival distribution to be the Gompertz, in which the mosquito mortality rate increases with age (i.e. there is senescence) (14). The baseline hazard, $h_0(t)$, was modelled as Gompertzian:

$$h_0(t) = ae^{bt}. \tag{1.01}$$

The hazard, $h(t)$, is modelled accounting for differences in mosquito infection status, E (= 1, if infected; = 0, if uninfected), and calculated as,

$$h(t) = h_0(t)e^{\beta_E E}, \tag{1.02}$$

where β_E is a constant. The probability of being alive at time t, $A(t)$, is calculated by integrating the hazard to time t, as follows,

$$A(t) = e^{-\int_0^t h(t)dt} = e^{\frac{a}{b}(1-e^{bt})e^{\beta_E E}}. \tag{1.03}$$

As mosquitoes were removed for dissection or still alive at the end of the study some observations were right censored, that is we do not know the time when death would have occurred. For these observations, the likelihood is given by, $A(t)$. The probability density function, $f(t)$, describing the event that death occurs at time t, for uncensored observations, is given by,

$$f(t) = h(t)A(t). \tag{1.04}$$

We suppose the infectious blood-fed group is a mix of infected and uninfected mosquitoes. The true infection status of these mosquitoes was not known. The probability of an observable infection in this group is given by δ, but we cannot distinguish between mosquitoes that never received any parasites and those that cleared the infection. There may be a potential cost of immunity and tissue damage caused by the ookinete life stage, but given that the impact of malaria infection is still unclear (26), we assume that if the

mosquito clears the malaria infection prior to the oocyst life stage, there is no additional mortality hazard due to infection. We account for uncertainty in the true infection status of mosquitoes by marginalising infection status out of the joint distribution as follows,

$$f(t) = \delta h_0(t) e^{(\beta_E)} e^{\left(\frac{a}{b}\right)\left(1-e^{bt}\right)e^{(\beta_E)}} + (1-\delta)h_0(t)e^{\left(\frac{a}{b}\right)\left(1-e^{bt}\right)}. \qquad (1.05)$$

7.1.2 Model fitting

Figure S1 Single temperature model estimates of parasite transition parameters. The gamma distribution posterior values of the development time between inoculation and observed sporozoites for the single temperature models. The MCMC median, 5th and 95th posterior quantiles are represented by the points and vertical lines respectively.

Figure S2 Kaplan Meier mosquito survival curves used in the model fitting.

Table S1 Model priors.

Parameter	Description	Prior parameters (subject to limits in next column)	Parameter limits	Justification of parameterisation (reference)
Temperature-independent parameters				
α_{OS}	Shape parameter for the gamma distribution governing oocyst to sporozoite development time	$\alpha_{OS} \sim N(15.0, 2.5)$	Lower: 0	Yields a mean sporozoite development time of 8 days
β_{OS}	Rate parameter for the gamma distribution governing oocyst to sporozoite development time	$\beta_{OS} \sim N(1.875, 2.5)$		
μ	Mean number of G-stage parasites inoculating an infected mosquito	$\mu \sim N(3.0, 2.5$	Lower: 0 Upper: 50	The median and mean number of oocysts per mosquito has been found to be approximately 3 (68–71)
k	Overdispersion in the number of G-stage parasites inoculating an infected mosquito	$k \sim N(0.1, 2.5)$	Lower: 0 Upper: 20	The overdispersion of oocysts has been found to be below slightly below 0.1 (70,71)
a	Gompertz distribution parameter governing the baseline hazard	$a \sim N(0.05, 2.5)$	Lower: 0 Upper: 5	Assuming mosquito survival is solely determined by the baseline hazard, the mean prior values result in a median survival time of 10.3 days (0.5 – 30.9 days 95% survival quantiles)
b	Gompertz distribution parameter governing the baseline hazard	$b \sim N(0.05, 2.5)$		
β_E	Cox proportional hazards model coefficient of infection status	$\beta_E \sim N(0.5, 1.0)$	Lower: -5 Upper: 5	The hazard odds ratio of infection is 1.65, given the mean prior value
σ_δ	Standard deviation of the delta error term	$\sigma_\delta \sim N(0, 0.25)$	Lower: 0	There are no differences between experiments, given the mean prior values

$\sigma_{survival}$	Standard deviation of the Cox model error term	$\sigma_{survival} \sim N(0, 0.25)$		
Temperature-dependent parameters				
$\alpha_{GO,}$	Shape parameter for the gamma distribution governing G-stage parasite to oocyst development time	$\alpha_{IO} \sim N(15.0, 2.5)$	Lower: 0	At the prior mean value of m_b (0) temperature has no effect on the oocyst development rate. The shape and rate priors give a mean oocyst development time of 5 days
m_β	Temperature-coefficient in eq (2.12)	$m_\beta \sim N(0, 2.5)$	m_b has no limits	
c_β	Intercept in eq (2.12)	$c_\beta \sim N(3.0, 2.5)$	Lower: 0	
m_δ	Temperature-coefficient in eq (2.13)	$m_\delta \sim N(0, 2.5)$		At the prior mean values of temperature has no effect on the transmission probability. Overall these result in a prior mean transmission probability of 0.69
c_δ	Intercept in eq (2.13)	$c_\delta \sim N(0.8, 2.5)$		
β_C	Temperature coefficient in Cox proportional hazards model (eq (2.15))	$\beta_C \sim N(0.5, 1.0)$	Lower: -5 Upper: 5	Mosquitoes have a hazard odds ratio of 1.15 for each 1°C increase in temperature

Figure S3 Trace plots of the MCMC parameter sampling for the all temperature model. The horizontal and vertical axis give the iteration and parameter values respectively. Iterations during the warmup are not included.

Figure S4 Posterior and prior probability densities. The posterior distributions are estimated by kernel density estimation with a gaussian smoothing kernel.

Table S2 All temperature model posterior values.

Parameter	Mean value	SE of the mean	Standard deviation	2.5% quantile	25% quantile	50% quantile	75% quantile	97.5% quantile	ESS	$\hat{R}$
α_{GO}	13.67	0.02	1.8	10.36	12.42	13.59	14.84	17.3	5808	1
m_β	1.44	0	0.19	1.09	1.31	1.43	1.57	1.83	6182	1
c_β	4.19	0.01	0.6	3.1	3.78	4.16	4.59	5.42	5847	1
α_{OS}	17.87	0.02	1.69	14.69	16.71	17.82	18.99	21.29	10275	1
β_{OS}	2.19	0	0.24	1.75	2.03	2.18	2.35	2.68	10101	1
μ	2.34	0	0.1	2.14	2.27	2.34	2.4	2.53	11620	1
k	3	0	0.53	2.1	2.63	2.95	3.32	4.2	14393	1
m_δ	-0.53	0	0.12	-0.76	-0.61	-0.53	-0.46	-0.3	6435	1
c_δ	0.59	0	0.13	0.34	0.51	0.6	0.68	0.85	4900	1
a	0.01	0	0	0.01	0.01	0.01	0.01	0.01	4772	1
b	0.12	0	0	0.11	0.11	0.12	0.12	0.12	16156	1
β_E	0.5	0	0.06	0.39	0.46	0.5	0.54	0.62	10200	1
β_C	0.45	0	0.08	0.29	0.4	0.45	0.5	0.62	4727	1
σ_δ	0.31	0	0.08	0.17	0.25	0.3	0.36	0.48	7127	1
$\sigma_{survival}$	0.27	0	0.09	0.15	0.21	0.26	0.31	0.51	5358	1

7.1.3 Logistic growth model

The cumulative sporozoite prevalence at time t, $g(t)$, was estimated under the logistic growth model as,

$$g(t) = \frac{g_{max}}{1 + e^{-c(t-t_m)}},$$

where g_{max} is the upper asymptote equivalent to the maximum possible sporozoite prevalence (used as an estimate of the human-to-mosquito transmission probability), t_m and c are constants that determine the growth rate (27). The binary logistic model was fit to the all dissected mosquitoes (binary coding; 1: sporozoites observed, 0: no sporozoites observed) simultaneously, using non-linear least squares.

7.2   Supplementary results

Figure S5 Modelled times required for individual malaria parasites to transition between life stages. Panels A and B show the probability density and the cumulative probability density that an individual malaria parasite within a mosquito (maintained under standard insectary conditions - 27 °C) will transition from a given life stage to the next life stage at a given time post blood-feed. The grey shaded area represents 5%-95% posterior quantiles of estimated distributions.

Figure S6 Single temperature model fits of the temporal variation in sporozoite prevalence. Black points: parasite prevalence of the laboratory mosquito data (95% confidence intervals are given by the vertical black lines). The grey shaded area represents the 95% uncertainty intervals of the mean prevalence (posterior predictive means). The black line represents the median of the poster predictive means.

Figure S7 Kaplan-Meier survival curves and all temperature Cox proportional-hazards survival model fit. Kaplan Meier curves are stepped. The shaded area shows the 95% uncertainty intervals of the posterior predictive mean survival probability (A(t)) modelled by the Cox proportional hazards model.

Figure S8 Kaplan Meier survival curves and the single temperature Cox proportional-hazards survival model fits. Kaplan Meier curves are stepped. The shaded area shows the 95% uncertainty intervals of the posterior predictive mean survival probability (A(t)) modelled by the Cox proportional hazards model.

Figure S9 All temperature model fits of the temporal variation in oocyst prevalence and oocyst intensity.
Oocyst prevalence plot: black points represent the prevalence of the laboratory mosquito data and the lines represent the 95% confidence intervals.: The grey shaded area represents the 95% uncertainty interval of the posterior predictive mean prevalence. Oocyst intensity plots: black points represent the mean oocyst intensity among all blood-fed mosquitoes and the grey shaded area represents the uncertainty in the oocyst count; the 5% and 95% negative binomial quantiles given the mean posterior predictive mean and mean posterior overdispersion parameter are shown.

Table S3 EIP estimates by temperature. The posterior predictive median, 5% and 95% quantiles of the "all temperature model" are given to one decimal place.

Temperature / °C	EIP_{10}	EIP_{50}	EIP_{90}
21	12.9 (12.5 - 13.3)	16.1 (15.5 - 16.8)	20.4 (19.4 - 21.8)
24	9.5 (9.3 - 9.7)	11.8 (11.6 - 12)	14.9 (14.5 - 15.2)
27	8.2 (8 - 8.3)	10.2 (10 - 10.4)	13 (12.7 - 13.3)
30	7.5 (7.3 - 7.6)	9.4 (9.2 - 9.6)	12 (11.7 - 12.4)
32	7.2 (7 - 7.3)	9 (8.9 - 9.2)	11.6 (11.3 - 12)
33	7 (6.9 - 7.2)	8.9 (8.7 - 9.1)	11.5 (11.1 - 11.8)
34	6.9 (6.7 - 7.1)	8.8 (8.6 - 8.9)	11.3 (11 - 11.7)

Figure S10 Logistic growth model fits. A logistic growth model (black line) was fit a subset of the sporozoite prevalence data (as determined by the time of peak sporozoite prevalence; black points were included and grey points were excluded).