Multiple pathways to homothallism in closely related yeast lineages in the Basidiomycota

Alexandra Cabrita[1], Márcia David-Palma[1*], Patrícia H. Brito[1], Joseph Heitman[2], Marco A. Coelho[1*] and Paula Gonçalves[1#]

[#] Corresponding author: pmz@fct.unl.pt

1. Applied Molecular Biosciences Unit-UCIBIO, Departamento de Ciências da Vida, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal

2. Department of Molecular Genetics and Microbiology, Duke University, Duke University Medical Center, Durham, NC 27710, USA.

Running title: Homothallism in *Cystofilobasidium*

Abstract

Sexual reproduction in fungi relies on proteins with well-known functions encoded by the mating-type (*MAT*) loci. In the Basidiomycota, *MAT* loci are often bipartite, with the *P/R* locus encoding pheromone precursors and pheromone receptors and the *HD* locus encoding heterodimerizing homeodomain transcription factors (Hd1/Hd2). The interplay between different alleles of these genes within a single species usually generates at least two compatible mating types. However, a minority of species are homothallic, reproducing sexually without an obligate need for a compatible partner. Here we examine the organization and function of the *MAT* loci of *Cystofilobasidium capitatum,* a species in the order Cystofilobasidiales, which is unusually rich in homothallic species. We determined *MAT* gene content and organization in *C. capitatum* and found that it resembles a mating type of the closely related heterothallic species *Cystofilobasidium ferigula.* To explain the homothallic sexual reproduction observed in *C. capitatum* we examined HD-protein interactions in the two *Cystofilobasidium* species and determined *C. capitatum MAT* gene expression both in a natural setting and upon heterologous expression in *Phaffia rhodozyma,* a homothallic species belonging to a clade sister to *Cystofilobasidium.* We conclude that the molecular basis for homothallism in *C. capitatum* appears to be distinct from that previously established for *P. rhodozyma.* Unlike the latter species, homothallism in *C. capitatum* may involve constitutive activation or dispensability of the pheromone receptor and the functional replacement of the usual Hd1/Hd2 heterodimer by an Hd2 homodimer. Overall, our results suggest that homothallism evolved multiple times within the Cystofilobasidiales.

Importance

Sexual reproduction is important for the biology of eukaryotes because it strongly impacts the dynamics of genetic variation. In fungi, although sexual reproduction is usually associated with the fusion between

---

[*] Present address: Department of Molecular Genetics and Microbiology, Duke University, Duke University Medical Center, Durham, NC 27710, USA.

cells belonging to different individuals (heterothallism), sometimes a single individual is capable of completing the sexual cycle alone (homothallism). Homothallic species are unusually common in a fungal lineage named Cystofilobasidiales. Here we studied the genetic bases of homothallism in one species in this lineage, *Cystofilobasidium capitatum*, and found it to be different in several aspects when compared to another homothallic species, *Phaffia rhodozyma*, belonging to the genus most closely related to *Cystofilobasidium*. Our results strongly suggest that homothallism evolved independently in *Phaffia* and *Cystofilobasidium*, lending support to the idea that transitions between heterothallism and homothallism are not as infrequent as previously thought. Our work also helps to establish the Cystofilobasidiales as a model lineage in which to study these transitions.

Introduction

In Fungi, as in all eukaryotes, sexual reproduction is widespread, and some of the underlying mechanisms and molecular components are conserved even among distant lineages. The specific molecular pathways involved and the recognition systems responsible for triggering sexual reproduction may nonetheless vary greatly (1). Generally, sexual reproduction occurs through mating of two haploid individuals of the same species, possessing distinct mating types (2, 3). Mating types are defined by specific regions of the genome, the mating type (*MAT*) loci, which encode proteins responsible for triggering the major pathways leading to sexual development. Distinct mating types differ in the genetic content of the *MAT* loci. In basidiomycetes, mating-type determination relies on two *MAT* loci (named *P/R* and *HD*) that encode two different classes of proteins (4–6). The *HD* locus contains two divergently transcribed genes encoding homeodomain transcription factors and the *P/R* locus comprises pheromone receptor and pheromone precursor encoding genes - *STE3* and *MFA,* respectively (4, 5, 7). In the Ascomycota, only one *MAT* locus encoding transcription factors is required to determine mating-type identity. Hence, the participation of receptors and pheromones in the determination of mating type is found only in the Basidiomycota.

The *HD* and *P/R MAT* loci in Basidiomycota can be either genetically linked or unlinked in the genome of a given species. If these loci are unlinked, they may segregate independently during meiosis, leading to the generation of four distinct mating types among the haploid progeny, which is the hallmark of the tetrapolar breeding system (4, 5, 7). The bipolar breeding system results from linkage of the two *MAT* loci and yields only two mating types in the haploid progeny (4, 5, 7). Bipolar mating also takes place if one of the two (unlinked) *MAT* loci loses its function in determining mating-type identity, which has been reported occasionally for the *P/R* locus (4, 5, 7).

Although heterothallism involving the fusion between compatible mating types as a requirement for sexual reproduction is common, in some fungal species individuals are universally compatible, meaning that they can undergo sexual reproduction with any other individual, or even with or by itself, a pattern termed homothallism (3, 7, 8). In basidiomycetes few homothallic organisms have been found and even fewer have had the genetic basis of their homothallism elucidated (1, 3, 7, 8). Two relevant examples of basidiomycete yeasts with distinct molecular strategies that result in a homothallic sexual mode of reproduction are the human pathogenic yeast *Cryptococcus deneoformans* (9, 10) and *Phaffia*

rhodozyma (11). *Cryptococcus deneoformans* exhibits a particular form of homothallism named
unisexual reproduction, where cells of the same mating type can either fuse or undergo endoreplication
of the entire genome resulting in a diploid nucleus that subsequently develops into hyphae, basidia, and
gives rise to four haploid spores that are products of a meiotic event. Interestingly, the key genes for
heterothallic reproduction like *MFA*, *STE3* and even *HD1* (*SXI1*) and *HD2* (*SXI2*) appear to be
dispensable for unisexual reproduction of some *C. deneoformans* strains (10, 12, 13).

More recently, the genetic basis of homothallism was dissected in *Phaffia rhodozyma* in our
laboratory (11). This astaxanthin-producing basidiomycetous yeast (14) belongs to the order
Cystofilobasidiales, a lineage with an unusually high proportion of homothallic species (15–20), but
comprising also heterothallic species and others for which sexual reproduction has not yet been
observed. In *P. rhodozyma*, deletion mutants were used to show that the two pairs of *STE3* and *MFA*
genes, and the single *HD1/HD2* gene pair present in the genome, are all required for robust sexual
reproduction (11). The two pairs of pheromone and pheromone receptors turned out to have reciprocal
compatibility, a single compatible *STE3* and *MFA* pair being sufficient for sexual reproduction. This is
what might be expected if the *P/R* loci of two putative mating types were present in the same genome,
in accord with the definition of primary homothallism. The only pair of Hd1 and Hd2 proteins encoded in
the genome is also essential for sexual development, but the mode of action is not fully understood (11).
This is because Hd1 and Hd2 are usually expected to heterodimerize to form functional transcription
factors, but proteins encoded in the same *HD* locus do not normally form dimers. This imposes
heterozygosity at the *HD* locus, with dimerization occurring only between proteins encoded by different
gene pairs, as a condition for sexual development. In line with this, the Hd1 and Hd2 proteins of *P.*
rhodozyma that are encoded in the same gene pair, do not appear to interact strongly, leading to the
tentative conclusion that a weak interaction between the two proteins might suffice for function (11).
Therefore, the *HD* locus configuration in *P. rhodozyma* is not fully aligned with the concept of primary
homothallism, where the presence of two distinct pairs of *HD* genes supporting strong cross dimerization
between Hd1 and Hd2 proteins would be expected.

Here, we examine in detail the content and function of the *MAT* loci of *Cystofilobasidium*
capitatum, a homothallic species closely related to *P. rhodozyma*, in order to understand if there are
common features between the molecular basis of homothallism in both species. We aim to shed some
light on the diversity of molecular mechanisms through which homothallism can occur in the phylum
Basidiomycota and improve the understanding of the evolution of mating patterns in the
Cystofilobasidiales.

109
110
111

RESULTS
MAT* loci in *Cystofilobasidium capitatum* and *Cystofilobasidium ferigula
Cystofilobasidium capitatum and *Cystofilobasidium ferigula* belong to the order
Cystofilobasidiales that also contains the genus *Phaffia*, for which two new species were recently
described (21), in addition to six other genera (22). The phylogenetic relationships within the order were

only recently clarified by a comprehensive genome-based phylogeny (21). This order contains a
strikingly large number of homothallic species (approximately one third of those described so far; **Table**
S1 (10.6084/m9.figshare.13176422)), a sexual mode uncommon in the Basidiomycota (3). In particular,
Phaffia is composed entirely of homothallic species (21) while *Cystofilobasidium* comprises species with
a variety of sexual properties, with *C. capitatum* and *C. intermedium* (23) representing the only two fully
homothallic species in the genus. The remaining species of *Cystofilobasidium* are heterothallic except
for *C. macerans*, which comprises strains exhibiting diverse sexual patterns (heterothallic, homothallic,
and asexual) and *C. alribaticum* for which no sexual reproduction has been observed under the
conditions tested (23). The genome-based phylogeny published for the Cystofilobasidiales (21) robustly
supported *Phaffia* and *Cystofilobasidium* as sister genera, a relationship that is recapitulated in the
phylogeny shown in **Fig. 1,** where a more limited number of species were included.

Fig. 1. Maximum likelihood phylogeny reconstructed from the concatenated protein alignments of 1147
single-copy genes shared across the studied taxa and the outgroup represented by *Cryptococcus*
deneoformans. Branch support was assessed by 1000 replicates of ultrafast bootstrap approximation
(UFBoot) with NNI optimization, and branch lengths are given in number of substitutions per site.
133

The availability of draft genomes for several *Cystofilobasidium* species allowed us to examine
the *MAT* loci of *C. capitatum* as well as that of the heterothallic species *C. ferigula* (**Fig. 2** and **Fig. S1**)
and to compare these with *MAT* loci in the homothallic *P. rhodozyma* concerning gene content and
organization (24). In the genome of *C. capitatum* PYCC 4530 a single pair of divergently transcribed
homeodomain transcription factors (*HD1* and *HD2*) was found, as well as a single pheromone receptor
gene (*STE3*) flanked by two identical pheromone precursor genes (*MFA1a* and *MFA1b*; **Fig 2**). These
two sets of genes are located on different scaffolds and encode proteins with similar lengths to their
counterparts in *P. rhodozyma* and the receptor is predicted to possess seven transmembrane domains
as expected (**Fig. 2B and Fig. S2**). Both predicted Hd proteins have homeobox domains and nuclear
localization signals (NLS; **Fig.S2**). However, the *MFA* genes in *C. capitatum* encode a 41 amino acid
pheromone precursor protein where no site for N-terminal processing could be identified (**Fig. 2C**), which
may compromise the formation of a mature pheromone (25). Furthermore, analysis of gene synteny
conservation between *Phaffia* and *C. capitatum* indicates that the *P/R* locus of *C. capitatum* may be
restricted to the region containing the pheromones/receptors (highlighted in yellow in **Fig. 2B**) and the

HD locus most likely includes only the *HD1* and *HD2* genes, as observed in many other basidiomycetes
(11, 26, 27).

Fig. 2. Gene content and organization of the (A) *HD* and (B) *P/R* mating-type loci in *C. capitatum* and
C. ferigula (PYCC 4410)*,* compared to *P. rhodozyma*. Genes are depicted as arrows denoting the
direction of transcription. The genomic regions corresponding to the putative *MAT* loci are highlighted
in yellow. Vertical bars connect orthologs that are in the same (blue) or inverted (pink) orientation. Non-
syntenic genes are shown in white, and genes in grey in *C. capitatum* and *C. ferigula* are found scattered
in the corresponding *P/R*- or *HD*-containing scaffold in *P. rhodozyma*, suggesting low level of synteny
conservation beyond the core *MAT* regions. Gene names are based on top BLASTp hits in the *P.
rhodozyma* genome and "hyp" denotes hypothetical proteins. The *P/R* locus of *C. ferigula* was omitted
because the available genome assemblies are too fragmented in these regions to allow for a
comparison. (C) Sequence alignment of pheromone precursors with amino acid positions colored in a
blue gradient according to conservation. Sequences proposed as the putative mature pheromones are
outlined by a red box, while those resembling the CAAX motif required for farnesylation are underlined.
In *C. capitatum* the absence of a conserved position for N-terminal processing (the two charged amino
acids indicated by a red arrowhead; ref 25), precludes the prediction of a mature pheromone sequence.
164

For *C. ferigula,* the genomes of two compatible mating types were obtained (PYCC 4410 and
PYCC 5628). Findings concerning *MAT* gene content were similar to *C. capitatum* (**Fig. 2** and **Fig. S1**),

and in line with the genetic composition typically found in haploid mating types of basidiomycetes. In *C.*
ferigula, two genes encoding pheromone precursors were also found in each of the mating types, but in
strain PYCC 5628 the two genes seem to encode slightly different mature pheromones. Because these
genes are found in very small contigs in the current genome assemblies, it was not possible to determine
their position in relation to the *STE3* gene nor the exact number of copies in the genome. It is conceivable
that additional pheromone genes may become apparent when more complete assemblies of *C.*
capitatum and *C. ferigula* genomes are available. As in *C. capitatum*, the *HD1/HD2* and the *STE3* genes
were also found on different scaffolds in both *C. ferigula* strains, but the higher level of fragmentation of
the current assemblies precludes a precise determination of the length and configuration of the *P/R* and
HD MAT loci.

While heterothallic species are expected to harbour at least two different mating types with
distinct alleles of *MAT* genes, this does not necessarily apply to homothallic species because there is
no requirement for an operational self/nonself-recognition system in this case. To assess allele diversity
of *MAT* genes in the *Cystofilobasidium* species under study, we obtained *MAT* gene sequences for as
many strains as possible for both species. For *C. ferigula* two clearly distinct *STE3* alleles (sharing ~51%
amino acid sequence identity) could be recognized among the five strains examined, as expected for a
heterothallic species (**Fig. 3A**). For *C. capitatum,* although several different alleles could be identified in
the 11 strains examined, they were much less divergent (sharing ~98% amino acid sequence identity)
than *STE3* alleles known to encode proteins with different specificities, like those of *P. rhodozyma* (~50%
sequence identity; (11)) and *C. ferigula* (**Fig. 3A**)*.*

188
Fig. 3. Sequence diversity of *STE3* and *HD* mating-type genes in *C. capitatum* and *C. ferigula*. (A)
Maximum likelihood phylogeny of pheromone receptors (Ste3) obtained from various strains of *C.*
capitatum (homothallic) and *C. ferigula* (heterothallic), along with the previously characterized Ste3
sequences of *P. rhodozyma* (homothallic) and *C. deneoformans* (heterothallic). The tree was inferred
with the LG+F+G4 model of amino acid substitution and was rooted in the midpoint. Branch support
values separated by a slash were assessed by 10,000 replicates of both the Shimodaira-Hasegawa
approximate likelihood ratio test (SH-aLRT) and the ultrafast bootstrap approximation (UFBoot).
Compared to the other species, the low sequence divergence of Ste3 sequences among *C. capitatum*
strains suggest the absence of functionally distinct, mating-type specific, alleles in this species. (B) and
(C) Sequence alignments of the *HD1* and *HD2* gene products of *C. capitatum* and *C. ferigula*. Sequence
identity between each pair of Hd1 and Hd2 proteins is given for the variable (N-termini) and conserved

(homeodomain and C-termini) regions with amino acid positions colored in a blue gradient according to
conservation. The comparison between Hd2 proteins of *C. ferigula* was not performed because the *HD2*
gene of PYCC 5628 is fragmented in the current genome assembly. Values in brackets for the N-
terminal regions are the average identity as calculated from different allele products (see **Fig. S3C** for
details on number of alleles). In *C. capitatum*, variable amino acid positions are evenly distributed
throughout the length of the Hd1 and Hd2 proteins. In contrast, the N-terminal region of Hd1 in *C. ferigula*
is comparatively more variable, as commonly observed in Hd1 and Hd2 proteins of other heterothallic
basidiomycetes (5, 7, 12, 26).
208

The comparison of *C. capitatum* Hd proteins suggest the existence of three main Hd variants in

the species with a degree of divergence between them that is lower than observed for functionally

different alleles of heterothallic species (**Fig. S3B and S3C**). Moreover, for the two proteins that could

be examined over their entire length, the differences in the sequences are distributed homogeneously

throughout the protein (**Fig. 3B**), like observed for *P. rhodozyma* (11)*. This is unlike the divergence

observed in Hd1 sequences of *C. ferigula* and of other heterothallic basidiomycetes, which is more

extensive and concentrated in the N-terminal region responsible for self/nonself-recognition (**Fig. 3C**).

C. ferigula was reported to be bipolar, as only two different mating types had been identified so

far (16). This would mean that each *STE3* allele is expected to be linked to a single *HD* allele, defining

two mating types. However, as shown in **Fig. S3C**, our analysis uncovered three *HD* alleles, instead of

the expected two, and the *HD.B1* allele appears associated with the two receptors in different strains,

whereas the *STE3.A1* allele is associated with *HD.B1* and *HD.B3*. From these observations it seems

more likely that *C. ferigula* may have a tetrapolar mating system, in contrast to previous assumptions

(16).

Involvement of the Hd proteins of *C. capitatum* in the homothallic sexual cycle

In heterothallic basidiomycetes, the Hd1 and Hd2 proteins encoded in the *HD* locus control the

later stages of sexual reproduction through heterodimerization of non-allelic Hd1 and Hd2 proteins

brought together by cell fusion (4, 12, 26). Previous studies in our laboratory concerning the molecular

mechanisms of sexual reproduction of *P. rhodozyma* revealed that both the Hd1 and Hd2 proteins are

required for normal sporulation (which for this species consists of basidia formation), and likely act

through heterodimerization despite the weak nature of their interaction (11).

To understand if *C. capitatum* resembled *P. rhodozyma* in this respect, the yeast two-hybrid

assay was employed to assess the ability of the Hd1 and Hd2 proteins of *C. capitatum* to interact with

each other. *HD1* and *HD2* cDNAs were isolated from *C. capitatum* strain PYCC 5626 and utilized for

the construction of the Gal4 fusion genes for this assay. The results of the assay, presented in **Fig. 4**,

are consistent with a complete absence of interaction between the Hd1 and Hd2 proteins of *C.*

capitatum, unlike the results for *P. rhodozyma* (11). Notably, a strong homodimerization was detected

for the Hd2 protein from this species. (**Fig. 4**). Weak homodimerizations of Hd proteins were also

previously observed for *P. rhodozyma* (11). For *C. ferigula* a strong heterodimerization of the Hd1 and

Hd2 proteins derived from strains of different mating types (PYCC 5628 and PYCC 4410, respectively)

was noted (**Fig. 4**), in line with observations in other heterothallic basidiomycete species (12, 26).

Consistently, β-galactosidase activity resulting from activation of the *lacZ* reporter gene was similar to

the positive control both for heterodimerization of Hd proteins from *C. ferigula* and for homodimerization

of Hd2 in *C. capitatum* (**Fig. 4B**)*.* Homodimerization of Hd1 in *C. capitatum* could not be tested, because

the construct of the fusion protein between Hd1 and the Gal4 DNA binding domain could not be stably

expressed in the pertinent *S. cerevisiae* strain.

Fig. 4. Results of the Yeast Two-Hybrid assay for Hd proteins of *C. capitatum* and *C. ferigula*. (A)
Qualitative results concerning the dimerization of Hd1 and Hd2 of *C. capitatum* and of *C. ferigula*, as
well as homodimerization of Hd2 of *C. capitatum*, through the activation of the reporter genes *MEL1,*
ADE2, and *HIS3*. Activity of the three reporter genes was tested separately in appropriate media, namely
containing X-α-Gal and all required supplements *for MEL1,* or lacking supplementation with adenine
(*ADE2)* or histidine *(HIS3)*. Activity of the two latter reporter genes is denoted by the ability of the strains
to grow on these media, while *MEL1* expression results in the formation of blue colonies. In *C. ferigula*
the Hd1 protein sequence is derived from strain PYCC 5628 and the Hd2 protein is derived from strain
PYCC 4410. (B) Quantitative results of the β-galactosidase assay of interactions shown in panel A. Each
datapoint is the average of the activity measured in two replicate assays; the three datapoints shown
for each strain represent β-galactosidase activities measured in reactions stopped after 2, 6 and 24
hours (see also Table S8 (10.6084/m9.figshare.13176422)). Blue bars denote the mean and vertical
bars the standard error of the means. The interaction between a fusion protein containing the Gal4
activation domain fused to the SV40 large T-antigen and a fusion between Gal4 binding domain and
p53 was used as positive control, while the negative control employed the same Gal4 activation domain
fusion in combination with a fusion between Gal4 binding domain and lamin. Plasmids encoding the
positive and negative control proteins were provided with the Matchmaker Gold Yeast Two-Hybrid
System, by Takara Bio USA. AD, activation domain of Gal4; BD, DNA binding domain of Gal4.
Significant differences between means were calculated using the Tukey's HSD test (*p<0.05, **p<0.01).

The *HD* locus of *C. capitatum* partially complements a *P. rhodozyma HD* deletion mutant

The results obtained in the yeast two-hybrid assay suggest that homodimerization of Hd2 may

play a role in homothallic sexual development in *C. capitatum,* possibly functionally replacing the usual

Hd1/Hd2 heterodimer. To investigate this, we set out to assess function of the *C. capitatum HD* locus

by heterologous expression in a *P. rhodozyma HDΔ* mutant*.*

Integration of the complete *HD* locus of *C. capitatum* strain PYCC 4530 in the rDNA locus of the

P. rhodozyma HDΔ mutant (construct *HDΔ+HD1/HD2-PYCC4530*) resulted in a very weak but

consistent recovery of sporulation (**Table 1, Table S2 and Table S3 (10.6084/m9.figshare.13176422)**)*.*

Because no interaction between the Hd1 and Hd2 proteins could be detected in the yeast two-hybrid

assay but instead strong homodimerization of the Hd2 protein was observed, we subsequently decided

to assess whether expression of *HD2* alone was sufficient to restore sexual development of the *P.*

rhodozyma HDΔ mutant. To this end, a construct containing in addition to the intergenic region only a

residual (344 bp) 5´portion of the *HD1* gene (excluding the homeodomain; *HDΔ+HD2-PYCC4530*) was

used to transform the *P. rhodozyma HDΔ* mutant. In this mutant that expresses only the Hd2 protein,

sporulation levels like those observed upon transformation of the complete *C. capitatum HD* locus were
observed (**Table 1, Table S2 and Table S3 (10.6084/m9.figshare.13176422)**), suggesting that only the
Hd2 protein is required for the observed complementation.
287
288

289 **Table 1.** Complementation of *P. rhodozyma MAT* loci mutant strains with *C. capitatum* genes.
290 Sporulation (basidia formation) patterns were scored qualitatively using the criteria explained as given
291 in the key.
292

Strain	*P. rhodozyma* WT	*HDΔ*	*HDΔ+HD2-PYCC4530*	*HDΔ+HD1/HD2-PYCC4530*	*P/RΔ*	*P/RΔ+STE3/MFA-PYCC4530*
Sporulation* phenotype	+++*	–	+	+	–	–

293
294 *Basidia counted on the surface of 30 colonies per assay; results were scored from two to four essays per strain;
295 (–): < 5 basidia; (+): 5-30 basidia; (+++) > 1000 basidia.
296
297

The *HD1 and HD2* genes of *C. capitatum* are differently expressed during growth in sporulation
conditions and in *P. rhodozyma* mutants
To substantiate the hypothesis that Hd2 might be the sole important component in the *HD* locus
of *C. capitatum*, we compared the expression levels of *HD1* and *HD2* in *C. capitatum* PYCC 5626, in
sporulation conditions, and in the *P. rhodozyma* mutant containing the complete *HD* locus of *C.*
capitatum PYCC 4530 (construct *HDΔ+HD1/HD2-PYCC4530*). The results, depicted in **Fig. 5**, show
that of the *HD* gene pair, only the *HD2* gene seems to be substantially expressed in the *C. capitatum*
strain. Heterologous expression of *HD2* in the *P. rhodozyma* mutant, although much lower than in *C.*
capitatum, can be easily detected, while heterologous expression of *HD1* seems to be only vestigial.
307

Fig. 5. Real time quantitative PCR results of *MAT* gene expression in *C. capitatum* under sporulation conditions and in *P. rhodozyma* complemented mutants. Expression of *MAT* genes is given as $\log_2$ fold difference relative to the expression of actin in each strain. These results are derived from two biological replicates, each assayed in triplicate, resulting in six datapoints for each measurement. For *HD1* expression assays less than six dots are plotted because in some reactions *HD1* expression was undetectable. Significant differences between the expression of different genes were calculated with the Mann-Whitney test (*$p<0.05$, **$p<0.01$).

_P/R_ locus of _C. capitatum_ does not restore sporulation in a _P. rhodozyma_ cognate deletion mutant

Because Hd function could be assessed using heterologous expression, a *P. rhodozyma* deletion mutant of both *P/R* clusters (*P/RΔ*) was similarly used as host for the *P/R* locus of *C. capitatum* strain PYCC 4530, encompassing the *STE3* and *MFA1*a genes and the respective native promotor regions (*P/RΔ+STE3/MFA-PYCC4530*). However, integration of the *P/R* locus of *C. capitatum* in the rDNA of the *P. rhodozyma P/RΔ* mutant failed to restore sporulation (**Table 1 and Table S2 (10.6084/m9.figshare.13176422)**), although the *C. capitatum STE3* gene was expressed in *P. rhodozyma* (**Fig. 5**). Interestingly, *STE3* is the most expressed among *MAT* genes of *C. capitatum* PYCC 5626 (**Fig. 5**), suggesting it may have a role in sexual reproduction despite our inability to observe it in the heterologous setting.

DISCUSSION

The main aim of this study was to ascertain to which extent common features could be found between the molecular bases of homothallism in different species of the Cystofilobasidiales, a lineage in the Basidiomycota particularly rich in species exhibiting this uncommon sexual behaviour. The molecular basis of homothallism was previously dissected in the genetically tractable Cystofilobasidiales

species *P. rhodozyma* (11). Here, we characterized the *MAT* loci of a second homothallic species belonging to a sister genus, *C. capitatum,* by examining the structure of the loci in the available genome of strain PYCC 4530 and by comparing *MAT* gene sequences in a total of 11 *C. capitatum* strains. The most striking difference between the *MAT* loci in *P. rhodozyma* and *C. capitatum* is the presence in the latter species of a single pheromone receptor gene, instead of the two distinct and functionally complementary sets of pheromone receptor and pheromone precursor genes found in *P. rhodozyma*. Therefore, the *C. capitatum MAT* gene content resembles a haploid mating type of a heterothallic species (4), while that of *P. rhodozyma* is reminiscent of a fusion between two mating types (11). However, like in *P. rhodozyma*, no evidence for functionally distinct variants (alleles) of *MAT* genes that might form different mating types were found in *C. capitatum,* consistent with its homothallic behaviour. We subsequently devised various experimental approaches to try to bring to light functional features of *MAT* genes in *C. capitatum.* In these experimental approaches two *C. capitatum* strains were used: in addition to sequenced strain PYCC 4530, strain PYCC 5626 was employed for the isolation of cDNAs because unlike other strains, it readily sporulates in different experimental conditions.

 P. rhodozyma can be described as a primary homothallic basidiomycete, with reciprocal compatibility between the pheromones and receptors of the two *P/R* clusters and with a weak heterodimerization between the only pair of Hd1 and Hd2 proteins being responsible for triggering of its sexual cycle, with possible hints at homodimerization events (11). Therefore, in *P. rhodozyma*, the *P/R* system seems to work similarly to what might be expected for a heterothallic mating type, while the *HD* locus is apparently operating through weak dimerization between adjacently encoded Hd1 and Hd2. This dimerization normally occurs only between proteins encoded by different *HD* alleles. Hence, the weak interaction in *P. rhodozyma* may have evolved to support sexual development in the absence of a second *HD* allele. Because the *MAT* loci in *C. capitatum* have the same gene content as haploid mating types of heterothallic species, in this case both the *P/R* and the *HD* loci probably underwent changes in their mode of operation to permit homothallism. For the *HD* locus, the complete absence of interaction between the Hd1 and Hd2 proteins, the formation of Hd2 homodimers, complementation results of the *P. rhodozyma HDΔ* mutant and the very low expression of the *HD1* gene suggest that the *HD* locus is relevant for sporulation but also that regulatory functions normally fulfilled by the Hd1/Hd2 heterodimer may have been taken over by the Hd2 homodimer. Hd1 homodimerization was not tested due to technical difficulties. Although it would have been very interesting to be able to test also Hd1 homodimerization, this lost importance when gene expression experiments showed that *HD1* expression was barely detectable. Interestingly, the α2 transcription factor, which is the Hd1 *S. cerevisiae* homologue, is capable of forming both a homodimer that represses transcription of **a**-specific genes in haploid α-cells, and a heterodimer with the a1 transcription factor (Hd2 homologue) that represses haploid-specific genes in diploid cells (28). This suggests that these proteins can operate both as homodimers and as heterodimers. However, to our knowledge, functional Hd1 or Hd2 homodimers have not been reported so far in basidiomycetes, although there was some evidence for homodimerization of Hd proteins in *P. rhodozyma* (11). Our hypothesis for the mode of action of the *HD* locus predicts that Hd1 might be dispensable. In some heterothallic basidiomycetes, it has been shown that the Hd1 and Hd2 proteins hold different functional domains that are essential for function of

the transcription regulation complex (29–31). One such species is the mushroom *Coprinopsis cinerea*,
where it has been shown that Hd1 contributes with the NLS region, allowing the heterodimer to be
transported into the nucleus, while the homeodomain of Hd2 is required for binding of the complex to
DNA (29, 32). Hence, in this case, formation of a heterodimer is required for function. In *C. capitatum*,
Hd2 possesses both an NLS and a homeodomain so in that perspective it seems possible that Hd2 may
indeed play a role in the homothallic life cycle of *C. capitatum* that is sufficient and independent of Hd1.
It is possible that this hypothesized function of Hd2 evolved recently in this species, which could explain
the fact that the *HD1* is not pseudogenized. Alternatively, the Hd1 may have acquired a function
unrelated to sexual reproduction, which might entail its expression under different conditions, and justify
its maintenance in the genome. Despite multiple attempts, the methods used to transform *P. rhodozyma*
failed to yield transformants of *C. capitatum,* which precluded the possibility of testing the hypothesis
regarding the role of Hd2 in this species by deletion of the *HD2* gene.
Introduction of the *P/R* locus into the *P. rhodozyma P/RΔ* mutant failed to restore sporulation of
the cognate mutant, even though expression of the *STE3* gene in the heterologous setting could be
demonstrated. The explanations for this are presently unclear, since the ability of Ste3-like receptors to
activate heterologously a sexual development pathway over a much larger phylogenetic distance was
previously demonstrated (33). As to the mode of action of the receptor in its normal setting, the possibility
that it could be constitutively active should be considered, in line with the fact that the predicted
pheromone precursor genes lack the N-terminal processing signals, likely precluding the formation of
an active pheromone. Mutations within the pheromone receptor that can lead to constitutively active
receptors have been previously reported, resulting in a bypass of the pheromone for sexual reproduction
(25, 33–36). Mutations in the pheromone that allow it to activate the receptor encoded in the same *P/R*
cluster are also known (25, 37). In fact, mutations that lead to self-activation or self-compatibility have
been proposed to form the basis for the transition from a tetrapolar to a bipolar mating system, or even
from a heterothallic to a homothallic mating behaviour (3, 5) for some species. If the pheromone system
does not operate normally in *C. capitatum*, this may be a reason for the lack of complementation of the
P. rhodozyma P/RΔ mutant by the *C. capitatum P/R* locus.
Hence, taken together, our results suggest that *P. rhodozyma* and *C. capitatum* attained
homothallic breeding systems through different mechanisms, which is consistent with the hypothesis
that the *P. rhodozyma* mating system arose at the origin of the genus, possibly by fusion of *P/R* loci of
compatible mating types and the adaptation of the HD dimerization system. *C. capitatum*, on the other
hand, has an extant makeup of its *MAT* loci that suggests it may have evolved from a heterothallic
mating type, in which the Hd2 homodimer acquired a prominent role. If a mature pheromone is formed
despite the absence of an N-terminal processing signal, it might be that the pheromone-receptor pair in
C. capitatum have become self-activating. On the other hand, if no mature pheromone is formed, the
pheromone receptor gene might have bypassed the pheromone requirement and is now constitutively
active. Whatever the case may be, the *P/R* system still likely has a role in some aspects of the sexual
development, which in agreement with the fact that *STE3* gene seems to be highly expressed. Although
we cannot discard the hypothesis that the receptor is fulfilling a role unrelated to sexual reproduction,

and while such receptors have been reported (38), they were not associated with pheromone precursor
genes.

The different particularities of homothallism in the two Cystofilobasidiales species studied so far
are suggestive of remarkable levels of plasticity in the evolution of sexual reproduction in this order. It
will be interesting to conduct similar studies in other homothallic species of this order, which would allow
us to get a more complete insight in the array of mechanisms involved as well as possible genomic
rearrangement that may have been involved in the transitions between heterothallic and homothallic
species. Having uncovered *P. rhodozyma* as a viable host for heterologous expression, opens the
possibility of assessing the functionality of other *MAT* proteins from uncharacterised species in this
order.

MATERIALS AND METHODS
Strains and culture conditions
P. rhodozyma, *C. capitatum* and *C. ferigula* strains (**Table S4**
(10.6084/m9.figshare.13176422)) were routinely grown in YPD medium (2% Peptone, 1% Yeast
Extract, 2% Glucose, 2% Agar) at 17 to 20ºC. For the preparation of electrocompetent cells, *P.*
rhodozyma strains were grown in YPD liquid medium at 20ºC, and transformants were incubated at
17ºC in selective medium, consisting of YPD plates supplemented with the appropriate antifungal drugs
(100 µg/ml of geneticin and/or 100 µg/ml of zeocin and/or 100 µg/ml of hygromycin B).
Escherichia coli strain DH5α (Gibco-BRL, Carlsbad, CA, USA) was used as a cloning host and
was grown at 37ºC in LB medium (1% NaCl, 1% Tryptone, 0.5% Yeast Extract and 2% Agar for solid
medium) supplemented with ampicillin at 100 µg/ml when appropriate.

DNA extraction, genome sequencing and assembly
Genomic DNA of *C. ferigula* PYCC 5628 was extracted from single cell-derived cultures using
the ZR Fungal/Bacterial DNA MiniPrep kit (ZYMO Research). DNA samples were quantified using Qubit
2.0. Genome sequencing was carried out by commercial providers, at the Genomics Unit of Instituto
Gulbenkian de Ciência, and at the Sequencing and Genomic Technologies Core Facility of the Duke
Center for Genomic and Computational Biology. Two short insert-size libraries (~500 bp) were prepared
with the Nextera Kit and subsequently sequenced using the Illumina MiSeq and HiSeq2500 systems to
generate paired 300- and 151-nt reads, respectively. After adaptor clipping using Trimmomatic (v0.36)
the two sets of reads were assembled with SPAdes (v3.11.1) (39) (with parameters: "--careful" to reduce
the number of mismatches and short indels in the final assembly, and the k-mer sizes: 21, 33, 55, 77,
99, 127, automatically selected based on read length). Genome assembly quality was assessed by the
QUAST (v.5.0.2) (40), and gene models were predicted ab initio using Augustus (41) trained on
Cryptococcus neoformans. Genome sequencing data generated, and final genome assembly statistics
are given in **Table S5 (10.6084/m9.figshare.13176422)**.

Identification of mating-type genes and synteny analyses

Scaffolds containing *MAT* genes, namely the homeodomain transcription factors (*HD1*/*HD2*)
and the mating pheromones (*MFA*) and receptors (*STE3*), were identified by BLASTP or TBLASTN in
the genome assemblies of *C. capitatum* PYCC 4530 (=CBS 7420; BioProject: PRJNA371774) and *C.*
ferigula PYCC 4410 (BioProject: PRJNA371786), and in the newly obtained assembly of *C. ferigula*
PYCC 5628 (BioProject: PRJNA371793). Well-annotated *P. rhodozyma MAT* proteins (11) were used
as search query. The retrieved *MAT* genes were manually reannotated if required and analysed further:
(i) the transmembrane regions in the pheromone receptor protein were predicted by HMMTOP software
(42); (ii) the Homeodomain regions in Hd1 and Hd2 proteins were predicted by InterPro server (43) and
compared to the previously characterized homeodomain proteins in Pfam database; (iii) nuclear
localization signals (NLS) and coiled-coil motifs were identified in the complete Hd1 and Hd2 sequences
using, respectively, the SeqNLS (with a 0.8 cut-off) (44) and Jpred4 (45) (see **Figs. S1B** and **S1C** and
Fig. S2). Synteny between *MAT* regions of different strains and species was based on bidirectional
BLAST analyses of the corresponding predicted proteins. The short pheromone precursor genes in the
genomes of *C. ferigula* and *C. capitatum* were identified manually as they usually fail automatic
detection.
468
Species and MAT gene phylogenies
A phylogenetic analysis representing major lineages within Cystofilobasidiales was inferred on
a concatenated protein dataset of single copy core genes of four *Cystofilobasidium* species, *Phaffia*
rhodozyma CBS 6938, *Mrakia frigida* PYCC 3896, *Krasilnikovozyma curviuscula* PYCC 5836, and the
outgroup *Cryptococcus deneoformans* JEC21 (**Table S**4 (10.6084/m9.figshare.13176422)).
Orthologous clusters were inferred with all-against-all BLASTP (NCBI Blast-2.2) searches and the
Markov cluster algorithm (OrthoMCL v1.4; (46)) with inflation factor (F) of 1.5, and minimum pairwise
sequence alignment coverage of 50% implemented in GET_HOMOLOGUES package (47). Clusters
present in single copy in all analyzed genomes were retained, aligned with MAFFT v7.407 using the G-
INS-I method and default parameter values (48), trimmed with BMGE v1.12 using the amino acid option
(49), and finally concatenated into a single data set. The species phylogeny was inferred with IQ-TREE
v1.6.12 (50) using maximum-likelihood (ML) inference under a LG+F+I+G4 model of sequence
evolution. ModelFinder (51) was used to determine best-fit model according to Bayesian Information
Criterion (BIC) and branch support was estimated using ultrafast bootstrap approximation (UFBoot) with
NNI optimization (52), both implemented in IQ-TREE package.
To analyse the *MAT* gene content across strains of *C. capitatum* and *C. ferigula*, protein
sequences of the *HD1*, *HD2* and *STE3* genes were retrieved from the genome assemblies and aligned
separately. Conserved regions were used to design primers to amplify the corresponding genomic
regions across the available strains of each species (**Table S6 (10.6084/m9.figshare.13176422)**).
These regions include a ~ 870-bp region of the *STE3* gene, and a ~1.5-kb fragment encompassing the
5′ end and intergenic regions of the *HD1* and *HD2* genes (**Table S6 (10.6084/m9.figshare.13176422)**
and Fig. S3). Genomic DNA was extracted through a standard Phenol-Chloroform method (53) and the
regions of interest were PCR-amplified, purified using Illustra GFX PCR DNA and Gel Band Purification
Kit (GE Healthcare Life Sciences), and then sequenced by Sanger Sequencing, at STABVida (Portugal).

For phylogenetic analysis of *MAT* genes, amino acid or nucleotide sequences were individually aligned
with MAFFT v7.310 (48) using the L-INS-i strategy (--localpair --maxiterate 1000) and poorly aligned
regions were trimmed with TrimAl (--gappyout) (54). The resulting alignments were input to IQ-TREE
v.1.6.5 (50) ML phylogenies using best-fit models automatically determined by ModelFinder (51)
(parameter: -m MFP). The exact model employed in tree reconstruction is given in the respective figure
legends. Branch support values were obtained from 10,000 replicates of both UFBoot (52) and the
nonparametric variant of the approximate likelihood ratio test (SH-aLRT) (55). In addition, the option "-
bnni" was employed to minimize the risk of overestimating branch supports with UFBoot when in
presence of severe model violations. The resulting phylogenies were midpoint rooted and graphically
visualized with iTOL v5.5.1 (56).

Yeast Two-Hybrid assay
To assess the interaction between the Hd1 and Hd2 proteins, the Matchmaker Gold Yeast Two-
Hybrid System kit (Takara) was used. In this system, fusion proteins containing the Gal4 DNA binding
domain (BD) are expressed from plasmid pGBKT7 in *MAT***a** haploid strain Y2HGold, while fusion
proteins containing the Gal4 activation domain (AD) are expressed from plasmid pGADT7 in *MATα*
haploid strain Y187. Diploid strains expressing both AD and BD fusion proteins are used to test
interactions and are obtained by mating the haploid strains expressing each of the fusion proteins of
interest. Hence, coding DNA sequences of the pertinent *HD* genes were cloned into pGADT7 and
pGBKT7, so as to yield plasmids expressing the desired fusion proteins.
Synthetic genes designed using the coding DNA sequences of the *HD1* and *HD2* genes of *C.*
ferigula and were synthesised at Eurofins Genomics (Germany). The *HD1* and *HD2* gene sequences of
strains PYCC 4410 and PYCC 5628 from *C. ferigula* respectively were adapted to the *S. cerevisiae*
codon usage (**Fig. S4**). The synthetic genes were obtained as inserts of pEX-A258 plasmids.
cDNAs of *HD1* and *HD2* from *C. capitatum* were obtained from total RNA isolated from strain PYCC
5626, briefly as follows. Strain PYCC 5626 was cultivated in GSA medium (0.2% Glucose, 0.2%
Soytone) in 10% of the flask volume, for 8 days at 20ºC and 90 rpm (Sartorius Certomat IS incubator),
until inspection under the microscope revealed the presence of teliospores (16). RNA extraction was
performed using the *ZR Fungal/Bacterial RNA MiniPrep kit* (by ZYMO Research), with a single step of
in-column DNase I Digestion to free the RNA samples of genomic DNA. cDNA was synthesized from
total mRNA using Maxima H Minus Reverse Transcriptase (by Thermo Scientific) and oligo (dT)20 as
primer, and synthesis of the second DNA strand was performed using specific primers for the complete
HD1 and *HD2* genes (**Table S6 (10.6084/m9.figshare.13176422)**). The fragments corresponding to the
HD1 and *HD2* cDNA sequences of strain PYCC 5626 were sequenced by Sanger Sequencing, at
STABVida (Portugal). The protein sequences of Hd1 and Hd2 of strain PYCC 5626 were aligned with
those of strain PYCC 4530 (**Fig. 3.B**) using the software MUSCLE (implemented in the software Unipro
UGENE v1.30.0 (57)) and the level of intraspecific variability was calculated.
The *HD1* and *HD2* complete synthetic genes from *C. ferigula* and cDNA fragments from *C.*
capitatum were amplified using primers that contained 40 bp tails at their 5' ends that correspond to the

flanking regions of the Multiple Cloning Sites present in pGADT7 and in pGBKT7 (**Table S6**
(10.6084/m9.figshare.13176422))). Plasmid pGADT7 was then linearized at the Multiple Cloning Site
by digestion with Cla I (Thermo Scientific), while pGBKT7 was linearized by digestion with Pst I (Thermo
Scientific). *S. cerevisiae* strains Y187 and Y2HGold were transformed with inserts and linearized vectors
using the transformation method described in the Yeastmaker Yeast Transformation System 2 protocol.
Transformants were selected in appropriate media (Yeast Nitrogen Base without amino acids, with
appropriate supplements (**Table S7 (10.6084/m9.figshare.13176422)**)) at 30ºC. Transformantions with
linearized pGBKT7 and the *C.capitatum* Hd1 insert resulted in numbers of transformants similar to those
observed in other transformations. However, unlike in other transformations, all transformants
investigated harboured plasmids without insert. This was observed only for this combination of vector
and insert and persisted in different transformation attempts.

Mating of haploid *S. cerevisiae* strains was performed by incubating a single colony from each
of the two haploid transformants to be mated in 200 µl of YPD medium at 30ºC, for 24 hours, at 250
rpm. After incubation, the cells were recovered and thoroughly washed with distilled sterile water and
plated on appropriate selective media (**Table S7 (10.6084/m9.figshare.13176422)**).

In the Matchmaker Gold Yeast Two-Hybrid System the *MEL1, ADE2 and HIS3* reporter genes
are used for qualitative assessment of interactions using plate assays, while the *LacZ* reporter gene is
used only to quantify the strength of the interaction, through quantification of β-galactosidase activity in
cell extracts. To test the activation of the *ADE2* and *HIS3* reporter genes, diploid strains were plated on
appropriate selective media without adenine or histidine, respectively, while to test the activation of the
yeast *MEL1* reporter gene encoding a secreted α–galactosidase, haploid transformants (to check for
autoactivation; Fig. S5) and diploid derivatives were plated on appropriate selective medium
supplemented with X-α-Gal (Takara Bio) at a final concentration of 40 µg/ml.

To quantify activation of the *LacZ* reporter gene, a β-galactosidase activity assay was performed, using
o-nitrophenyl β-D-galactopyranoside (ONPG) as a substrate. The assay was performed as described in
the Yeast Protocols Handbook (Takara Bio). All reactions were performed in triplicate, so that they could
be stopped at 3 different points in time (after 2 hours, 6 hours, and 24 hours), by the addition of 0.4 mL
of 1M Na_2CO_3 to each suspension. Raw data concerning these assays is shown in **Table S8**
(10.6084/m9.figshare.13176422).

Construction of recombinant plasmids and gene deletion cassettes

For the construction of *P. rhodozyma* mutants, recombinant plasmids and gene deletion
cassettes were constructed, as follows. Primer sequences (**Table S6 (10.6084/m9.figshare.13176422)**)
were based on available genome sequences of *P. rhodozyma* strain CBS 6938 (NCBI project
PRJEB6925 (58)) and *C. capitatum* strain PYCC 4530 (Bioproject PRJNA371774). Plasmids used for
the constructions were pJET1.2/blunt (Thermo Scientific), pPR2TN containing a geneticin resistance
cassette (59) and pBS-HYG (60) containing a hygromycin resistance cassette. All fragments used for
cloning and deletion cassettes were amplified by PCR using Phusion High-Fidelity DNA polymerase
(Thermo Scientific) and the amplified products were purified using either *Illustra GFX PCR DNA and Gel*

Band Purification Kit (GE Healthcare) or *GeneJET Gel Extraction Kit* (Thermo Scientific). Constructions were performed using standard molecular cloning methods (61) and *E. coli* strain DH5α as host.

Transformation of *P. rhodozyma*

 P. rhodozyma strains (CBS 6938 and mutants) (**Table S4 (10.6084/m9.figshare.13176422)**) were transformed by electroporation, with the linearized recombinant plasmids or deletion cassettes, as previously described by Visser *et al*, 2005 (62), reducing the amount of DNA used to 2 μg for the transformations of the complemented *P. rhodozyma* mutants. The electroporation conditions (Gene Pulser II Electroporation System, Bio-Rad) consisted of an internal resistance of 1000 Ω, an electric pulse of 0.8 kV, a capacitance of 25 μF, resulting in a pulse ranging from 18 to 20 ms (63, 64). The cells recovered subsequently in YPD liquid medium, for at least 2.5 hours at 17 ºC before being plated on YPD medium supplemented with the appropriate antifungal drug and incubated at 17ºC. The genotypes of the transformants were determined as previously described (11, 65).

***P. rhodozyma* sporulation (basidia formation) assays**

 To determine the ability of the *P. rhodozyma* mutants to sporulate, basidia formation assays were performed, where the strains were incubated in DWR solid medium with 0.5% of ribitol (0.5% Ribitol and 2.5% Agar), as previously described (11). Each basidia formation assay was conducted on 3 plates containing DWR+0.5% ribitol. On each plate, 10 colonies of each strain to be tested were spotted. Different strains were employed in each assay as indicated in **Table S2 (10.6084/m9.figshare.13176422)**, but in all assays the *P. rhodozyma* wild type was used as a positive control. Colonies were examined under the microscope using 100x magnification after 10 and 20 days of incubation at 18ºC, and basidia formation patterns were scored qualitatively. The numbers of basidia counted in experiment E8 for the complemented mutants concerning the study of the *HD* locus of *C. capitatum* after 20 days of incubation are listed in **Table S3 (10.6084/m9.figshare.13176422)**. In all cases the entire colony was submitted to microscopic observation.

Real-time quantitative PCR to assess expression of the *MAT* genes

 Total RNA was extracted from a sporulating culture of *C. capitatum* strain PYCC 5626. Sporulation (teliospore formation in the case of this species) was induced by incubation in GSA liquid medium (0.2% Glucose, 0.2% Soytone) in 10% the volume of the flask, at 17ºC, without agitation, until microscopic inspection revealed hyphae and teliospores, the latter being thick-walled resting spores from which basidia arise (16). Total RNA extraction was performed using the *ZR Fungal/Bacterial RNA MiniPrep kit* (by ZYMO Research). *P. rhodozyma* complemented mutants were grown in YPD liquid medium to an OD_{600nm} of 1.0. The cultures were then collected and frozen at -80ºC for 1 hour before proceeding with total RNA extraction through a standard Trizol method. In-column DNase I digestion to free the RNA samples of genomic DNA using the RNA Clean & Concentrator kit (by ZYMO Research) was used for all samples, and absence of gDNA was verified by PCR. cDNA was synthesized from total mRNA using Maxima H Minus Reverse Transcriptase (Thermo Scientific) and oligo (dT)20 as primer. Real-Time PCR was performed using the SensiFAST SYBR No-ROX kit (by Bioline, London), with 20µl

reactions, in a Rotor-Gene 6000 Corbett apparatus. The reaction parameters consisted of an initial denaturation step at 95ºC, for 2 min, followed by 40 cycles of 95ºC for 5 seconds, 57ºC for 10 seconds and 72ºC for 20 seconds. Two biological replicates were performed, with triplicates performed for each reaction (**Table S9 (10.6084/m9.figshare.13176422)**). Relative expression of the *MAT* genes was calculated using the $2^{-\Delta Ct}$ method, where $\Delta Ct = Ct_{test} - Ct_{reference}$, as described by Livak and Schmittgen (66), and expression values were represented as the $\log_2 (2^{-\Delta Ct})$. Mann-Whitney tests were performed to determine if the differences in expression of genes within each strain were statistically different.

Data availability

Nucleotide sequences been deposited in NCBI/EMBL (GenBank) under accession numbers: MT561333-MT561342 (*C. capitatum STE3*); MT561330-MT561332 (*C. ferigula STE3*); MT592882-MT592890 and MT592891-MT592894 (respectively for *C. capitatum* and *C. ferigula* 5'-end and intergenic region of *HD1* and *HD2*). Sequencing reads for *C. ferigula* PYCC 5628 (BioProject PRJNA371793) are available in the NCBI SRA database. The *C. ferigula* PYCC 5628 draft genome assembly has been deposited at DDBJ/ENA/GenBank under accession numbers MVAN00000000.

ACKNOWLEDGEMENTS
The authors are very thankful to Alex Idnurm, Brenda Wingfield, Timothy James, and Paul Dyer for their insightful and thorough reviews of this manuscript. This work was supported by UCIBIO-Unidade de Ciências Biomoleculares Aplicadas, which is financed by Portuguese funds from Fundação para a Ciência e Tecnologia, Ministério da Ciência, Tecnologia e Ensino Superior (FCT/MCTES; https://www.fct. pt/) grant UID/Multi/04378/2019 and FCT/MCTES grants PTDC/BIA-GEN/112799/2009 (to P.G.), SFRH/BPD/79198/2011 (M.C.) and SFRH/BD/81895/2011 (M.D.P.). This work also benefited from the support of the INCD computing infrastructure funded by FCT and FEDER under the project 01/SAICT/2016 nº 022153. This work was also supported by NIH/NIAID R37 Award AI39115-23 and R01 award AI50113-16. J.H. is co-director and fellow for CIFAR program Fungal Kingdom: Threats & Opportunities.

References

1.   Lee SC, Ni M, Li W, Shertz C, Heitman J. 2010. The evolution of sex: a perspective from
        the fungal kingdom. Microbiol Mol Biol Rev 74:298–340.

2.   Stanton BC, Hull CM. 2007. Mating-Type Locus Control of Cell Identity, p. 59–73. *In*
        Heitman, J, Kronstad, JW, Taylor, JW, Casselton, LA (eds.), Sex in fungi: molecular
        determination and evolutionary implications. ASM Press, Washington, DC.

3.   Lin X, Heitman J. 2007. Mechanisms of homothallism in fungi and transitions between

heterothallism and homothallism, p. 35–57. *In* Heitman, J, Kronstad, JW, Taylor, JW,

Casselton, LA (eds.), Sex in fungi: molecular determination and evolutionary

implications. ASM Press, Washington, D.C.

4.   Coelho MA, Bakkeren G, Sun S, Hood ME, Giraud T. 2017. Fungal sex: the

Basidiomycota. Microbiol Spectr 5:FUNK-0046-2016.

5.   Fraser JA, Hsueh Y-P, Findley KM, Heitman J. 2007. evolution of the mating-type locus:

the Basidiomycetes, p. 19–34. *In* Heitman, J, Kronstad, JW, Taylor, JW, Casselton, LA

(eds.), Sex in Fungi: Molecular Determination and Evolutionary Implications. ASM

Press, Washington, DC.

6.   Raudaskoski M, Kothe E. 2010. Basidiomycete mating type genes and pheromone

signaling. Eukaryot Cell 9:847–859.

7.   Kües U, James TY, Heitman J. 2011. Mating type in basidiomycetes: unipolar, bipolar,

and tetrapolar patterns of sexuality, p. 97–160. *In* Pöggeler, S, Wöstemeyer, J (eds.),

The Mycota, Evolution of Fungi and Fungal-Like Organisms, Volume XIV. Springer,

Berlin Heidelberg.

8.   Wilson AM, Wilken PM, van der Nest MA, Steenkamp ET, Wingfield MJ, Wingfield BD.

2015. Homothallism: an umbrella term for describing diverse sexual behaviours. IMA

Fungus 6:207–214.

9.   Lin X, Hull CM, Heitman J. 2005. Sexual reproduction between partners of the same

mating type in *Cryptococcus Neoformans*. Nature 434:1017–1021.

10. Wang L, Lin X. 2011. Mechanisms of unisexual mating in *Cryptococcus neoformans*.

Fungal Genet Biol 48:651-660.

11. David-Palma M, Sampaio JP, Gonçalves P. 2016. Genetic dissection of sexual

reproduction in a primary homothallic basidiomycete. PLoS Genet 12: e1006110.

12. Hull CM, Boily M, Heitman J. 2005. Sex-specific homeodomain proteins sxi1 α and sxi2

a coordinately regulate sexual development in *Cryptococcus neoformans*. Eukaryot

Cell 4:526–535.

13. Gyawali R, Zhao Y, Lin J, Fan Y, Xu X, Upadhyay S, Lin X. 2017. Pheromone independent

unisexual development in *Cryptococcus neoformans*. PLOS Genet 13:e1006772.

14. Barredo J, García-Estrada C, Kosalkova K, Barreiro C. 2017. Biosynthesis of astaxanthin

as a main carotenoid in the heterobasidiomycetous yeast *Xanthophyllomyces*

dendrorhous. J Fungi 3:44.

15.   Kurtzman CP, Fell JW, Boekhout T. 2011. The Yeasts, a Taxonomic Study, 5th ed.

Elsevier.

16.   Sampaio JP. 2011. *Cystofilobasidium* Oberwinkler & Bandoni (1983), p. 1423–1432. *In*

Kurtzman, CP, Fell, JW, Boekhout, T (eds.), The Yeasts, a Taxonomic Study, 5th ed.

Elsevier.

17.   Fell JW, Johnson EA, Scorzetti G. 2011. *Xanthophyllomyces* Golubev (1995), p. 1595–

1599. *In* Kurtzman, CP, Fell, JW, Boekhout, T (eds.), The Yeasts, a Taxonomic Study,

5th ed. Elsevier.

18.   Fell JW. 2011. *Mrakia* Y. Yamada & Komagata (1987), p. 1503–1510. *In* Kurtzman, CP,

Fell, JW, Boekhout, T (eds.), The Yeasts, a Taxonomic Study, 5th ed. Elsevier.

19.   Sampaio JP. 2011. *Tausonia* Bab'eva (1998), p. 1999–2001. *In* Kurtzman, CP, Fell, JW,

Boekhout, T (eds.), The Yeasts, a Taxonomic Study, 5th ed. Elsevier.

20.   Takashima M, Nakase T. 2011. *Udeniomyces* Nakase & Takematsu (1992), p. 2063–

2068. *In* Kurtzman, CP, Fell, JW, Boekhout, T (eds.), The Yeasts, a Taxonomic Study,

5th ed. Elsevier.

21.   David-Palma M, Libkind D, Brito PH, Silva M, Bellora N, Coelho MA, Heitman J,

Gonçalves P, Sampaio JP. 2020. The untapped australasian diversity of astaxanthin-

producing yeasts with biotechnological potential - *Phaffia australis* sp. nov. and

Phaffia tasmanica sp. nov. Microorganisms 8, 1651.

22.   Kachalkin A V, Turchetti B, Inácio J, Carvalho C, Mašínová T, Pontes A, Röhl O,

Glushakova AM, Akulov A, Baldrian P, Begerow D, Buzzini P, Sampaio JP, Yurkov AM.

2019. rare and undersampled dimorphic basidiomycetes. Mycol Prog 18:945–971.

23.   Pontes A, Röhl O, Carvalho C, Maldonado C, Yurkov AM, Paulo Sampaio J. 2016.

Cystofilobasidium intermedium sp. nov. and *Cystofilobasidium alribaticum* f.a. sp.

nov., isolated from mediterranean forest soils. Int J Syst Evol Microbiol 66:1058–1062.

24.   Bellora N, Moliné M, David-Palma M, Coelho MA, Hittinger CT, Sampaio JP, Gonçalves

P, Libkind D. 2016. Comparative genomics provides new insights into the diversity,

physiology, and sexuality of the only industrially exploited tremellomycete: *Phaffia*

rhodozyma. BMC Genomics 17:901.

25.   Kües U. 2015. From two to many: multiple mating types in basidiomycetes. Fungal Biol

Rev 29:126–166.

26.   Kämper J, Reichmann M, Romeis T, Bölker M, Kahmann R. 1995. multiallelic
714   recognition: nonself-dependent dimerization of the bE and bW homeodomain
715   proteins in *Ustilago maydis*. Cell 81:73–83.

716   27.   Maia TM, Lopes ST, Almeida JMGCF, Rosa LH, Sampaio JP, Gonçalves P, Coelho MA.
717   2015. evolution of mating systems in basidiomycetes and the genetic architecture
718   underlying mating-type determination in the yeast *Leucosporidium scottii*. Genetics
719   201:75–89.

720   28.   Dranginis AM. 1990. Binding of yeast a1 and α2 as a heterodimer to the operator dna
721   of a haploid-specific gene. Nature 347:682–685.

722   29.   Asante-Owusu RN, Banham AH, Böhnert HU, Mellor EJC, Casselton LA. 1996.
723   heterodimerization between two classes of homeodomain proteins in the mushroom
724   *Coprinus cinereus* brings together potential dna-binding and activation domains. Gene
725   172:25–31.

726   30.   Schlesinger R, Kahmann R, Kämper J. 1997. The homeodomains of the heterodimeric
727   Be And Bw proteins of *Ustilago maydis* are both critical for function. Mol Gen Genet
728   254:514–519.

729   31.   Van Diepen LTA, Olson A, Ihrmark K, Stenlid J, James TY. 2013. Extensive trans-specific
730   polymorphism at the mating type locus of the root decay fungus *Heterobasidion*. Mol
731   Biol Evol 30:2286–2301.

732   32.   Kües U, Göttgens B, Stratmann R, Richardson W V, O'Shea SF, Casselton L a. 1994. A
733   chimeric homeodomain protein causes self-compatibility and constitutive sexual
734   development in the mushroom *Coprinus cinereus*. EMBO J 13:4054–4059.

735   33.   Hsueh Y-P, Xue C, Heitman J. 2009. A constitutively active GPCR governs morphogenic
736   transitions in *Cryptococcus neoformans*. EMBO J 28:1220–1233.

737   34.   Olesnicky NS, Brown AJ, Dowell SJ, Casselton LA. 1999. A constitutively active G-
738   protein-coupled receptor causes mating self-compatibility in the mushroom *Coprinus*.
739   EMBO J 18:2756–2763.

740   35.   Boone C, Davis NG, Sprague GF. 1993. Mutations that alter the third cytoplasmic loop
741   of the a-factor receptor lead to a constitutive and hypersensitive phenotype. Proc Natl
742   Acad Sci 90:9921–9925.

743   36.   Fowler TJ, Mitton MF, Vaillancourt LJ, Raper CA. 2001. changes in mate recognition
744   through alterations of pheromones and receptors in the multisexual mushroom

fungus *Schizophyllum commune*. Genetics 158:1491–1503.

37.   Riquelme M, Challen MP, Casselton LA, Brown AJ. 2005. The origin of multiple *b*

mating specificities in *Coprinus cinereus*. Genetics 170:1105–1119.

38.   Díaz-Valderrama JR, Aime MC. 2016. The cacao pathogen *Moniliophthora roreri*

(Marasmiaceae) possesses biallelic *A* and *B* mating loci but reproduces clonally.

Heredity 116:491–501.

39.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, Pyshkin A V, Sirotkin A V, Vyahhi N, Tesler G,

Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its

applications to single-cell sequencing. J Comput Biol 19:455–477.

40.   Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. 2018. Versatile genome

assembly evaluation with QUAST-LG. Bioinformatics 34:i142–i150.

41.   Stanke M, Diekhans M, Baertsch R, Haussler D. 2008. Using native and syntenically

mapped cDNA alignments to improve *de novo* gene finding. Bioinformatics 24:637–

644.

42.   Tusnády GE, Simon I. 2001. The HMMTOP transmembrane topology prediction server.

Bioinformatics 17:849–850.

43.   Mitchell A, Chang H, Daugherty L, Fraser M, Hunter S, Lopez R, McAnulla C,

McMenamin C, Nuka G, Pesseat S, Sangrador-Vegas A, Scheremetjew M, Rato C, Yong

S, Bateman A, Punta M, Attwood T, Sigrist C, Redaschi N, Rivoire C, Xenarios I, Kahn D,

Guyot D, Bork P, Letunic I, Gough J, Oates M, Haft D, Huang H, Natale D, Wu C, Orengo

C, Sillitoe I, Mi H, Thomas P, Finn R. 2015. The InterPro protein families database: the

classification resource after 15 years. Nucleic Acids Res 43:D213-221.

44.   Lin J rong, Hu J. 2013. SeqNLS: nuclear localization signal prediction based on frequent

pattern mining and linear motif scoring. PLoS One 8:e76864.

45.   Drozdetskiy A, Cole C, Procter J, Barton GJ. 2015. JPred4: a protein secondary

structure prediction server. Nucleic Acids Res 43:W389–W394.

46.   Li L, Stoeckert Jr CJ, Roos DS. 2003. OrthoMCL: identification of ortholog groups for

eukaryotic genomes. Genome Res 13:2178–2189.

47.   Contreras-Moreira B, Vinuesa P. 2013. GET_HOMOLOGUES, a versatile software

package for scalable and robust microbial pangenome analysis. Appl Environ Microbiol

79:7696–7701.

48.   Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. Mol Biol Evol 30:772–780.

49.   Criscuolo A, Gribaldo S. 2010. BMGE (Block Mapping and Gathering with Entropy): a new software for selection of phylogenetic informative regions from multiple sequence alignments. BMC Evol Biol 10:210.

50.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol 32:268–274.

51.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods 14:587–589.

52.   Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the Ultrafast Bootstrap Approximation. Mol Biol Evol 35:518–522.

53.   Gonçalves P, Valério E, Correia C, de Almeida JMGCF, Sampaio JP. 2011. Evidence for divergent evolution of growth temperature preference in sympatric *Saccharomyces* species. PLoS One 6:e20739.

54.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–1973.

55.   Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. Syst Biol 59:307–321.

56.   Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new developments. Nucleic Acids Res 47:W256–W259.

57.   Okonechnikov K, Golosova O, Fursov M, Varlamov A, Vaskin Y, Efremov I, German Grehov OG, Kandrov D, Rasputin K, Syabro M, Tleukenov T. 2012. Unipro UGENE: A unified bioinformatics toolkit. Bioinformatics 28:1166–1167.

58.   Sharma R, Gassel S, Steiger S, Xia X, Bauer R, Sandmann G, Thines M. 2015. The genome of the basal agaricomycete *Xanthophyllomyces dendrorhous* provides insights into the organization of its acetyl-CoA derived pathways and the evolution of Agaricomycotina. BMC Genomics 16:233.

59.   Verdoes JC, Krubasik P, Sandmann G, Van Ooyen AJJ. 1999. Isolation and functional

characterisation of a novel type of carotenoid biosynthetic gene from

Xanthophyllomyces dendrorhous. Mol Gen Genet 262:453-461.

60. Niklitschek M, Alcaíno J, Barahona S, Sepúlveda D, Lozano C, Carmona M, Marcoleta

A, Martínez C, Lodato P, Baeza M, Cifuentes V. 2008. Genomic organization of the

structural genes controlling the astaxanthin biosynthesis pathway of

Xanthophyllomyces dendrorhous. Biol Res 41:93-108.

61. Sambrook J, Russell D. 2001. Molecular Cloning: A Laboratory Manual, 3rd ed. Cold

Spring Harbor Laboratory Press, New York.

62. Visser H, Sandmann G, Verdoes JC. 2005. Xanthophylls in Fungi. Metabolic engineering

of the astaxanthin biosynthetic pathway in *Xanthophyllomyces dendrorhous.*

Microbial Processes and Products.pp 257-272.

63. Rodríguez-Sáiz M, Godio RP, Álvarez V, De La Fuente JL, Martín JF, Barredo JL. 2009.

The NADP-dependent glutamate dehydrogenase gene from the astaxanthin producer

Xanthophyllomyces dendrorhous: Use of its promoter for controlled gene expression.

Mol Biotechnol 41:165-172.

64. Wery J, Verdoes JC, Van Ooyen AJJ. 1998. Efficient transformation of the astaxanthin-

producing yeast *Phaffia rhodozyma*. Biotechnol Tech 12:399-405.

65. Lin G, Bultman J, Johnson EA, Fell JW. 2012. Genetic manipulation of

Xanthophyllomyces dendrorhous and *Phaffia rhodozyma*. Methods Mol Biol 898:235–

249.

66. Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-

time quantitative PCR and the 2-ΔΔCT method. Methods 25:402–408.

843 **Supplementary Figures**

845
846
Fig. S1. Analysis of the *HD* locus of *C. ferigula*. (A) Genomic organization of the *HD1* and *HD2* genes
in *C. ferigula* PYCC 4410 and PYCC 5628. The *HD2* gene in strain PYCC 5628 is fragmented and
localizes, as depicted, at the ends of two different scaffolds. However, comparative analysis indicates
this region is syntenic in both *C. ferigula* strains. (B) Hd1 amino acid sequences of strains PYCC 4410
and PYCC 5628 shown as a sequence alignment (on the top) and as a schematic representation (on
the bottom). (C) Hd2 amino acid sequence of *C. ferigula* PYCC 4410. In panels B and C, typical protein
secondary structure features are highlighted according to the key on the top right.

Fig. S2. Sequence alignment of the (A) *HD1* and (B) *HD2* gene products from *C. capitatum* PYCC 4530 and PYCC 5626. In both panels, the amino acid sequence alignment is shown on the top and a schematic representation is shown below. Typical protein secondary structure features are highlighted according to the key on the bottom.(C) Sequence alignment of the Ste3 receptors from *C. capitatum* and *C. neoformans,* and the *C. deneoformans* pheromone receptor-like Cpr2, highlighting the seven transmembrane domains. The constitutive activity of Cpr2 in *C. deneoformans* results from an

864 unconventional residue, Leu^{222}, in place of a conserved proline in transmembrane domain six (red box;
865 Hsueh Y-P, Xue C, and Heitman J, EMBO J 28:1220–1233, 2009).

874
875

876
Fig. S3.
Fig. S3. Determination of different alleles of the *HD* loci in *C. capitatum* and *C. ferigula*. (A) A ~1.5-kb-
long genomic region spanning the homeodomain, the 5' end region, and the common intergenic region
of the *HD1* and *HD2* genes was PCR-amplified and sequenced from available 10 strains of *C.*
capitatum and 5 strains of *C. ferigula*. Primers locations are shown as yellow arrowheads below the
genes. (B) Maximum likelihood phylogenies inferred from the 5' end of the *HD1* and *HD2* genes
(regions underlined in panel A) of *C. capitatum* using the best-fit substitution models HKY+F and K2P,
respectively. (C) Maximum likelihood phylogeny inferred from the 5' end of the *HD1* gene of *C. ferigula*
using the best-fit model K2P. Note that the branch length in the two *HD1* gene trees is quite different
(scale bars in nucleotide substitutions per site), which implies a much higher divergence of the *C.*
ferigula HD1 alleles compared to the *HD1* alleles of *C. capitatum*. The trees are rooted in the midpoint
and branch support values separated by a slash were assessed by 10,000 replicates of both
Shimodaira-Hasegawa approximate likelihood ratio test (SH-aLRT) and the ultrafast bootstrap
approximation (UFBoot). The molecular mating type assigned for each of the analyzed strains of *C.*
ferigula provides evidence that this species has a biallelic *P/R* locus and a multiallelic *HD* locus, which
are genetically unlinked (the *B1* allele appears associated with both *A1* and *A2* alleles).
893

C. ferigula PYCC5628 *HD1* synthetic gene sequence

```
   1 ATGGCAACCCAAACTATTCTCAAAAACCGACTTCAGGACGGGCTGCAGAATTTCTTCAGTGTACTAAAAAGCGGTGACAACCAGTCATGTCGAAGTTTCC
 101 AGGCACAACTTCTTTCACTAGTCCAGGCTGTTGGACAAGCAGCACAGCAGGGGCGGTTGGATGAGGAAACCAGAACGGTATGTGGTGCTGTAGCAGCCAT
 201 ACTTGAATCGAGCGCCACTGTCCTTCTTCAGGTTAAGTCTGACGCCTGTGTTATGGAACAGCAGCTGCATTTTGATGTTAGCAAAGATCTAGGCTCGCAT
 301 GCCGTGATAAGGTCGTCCGTGGAAGAACACACGGCGCCATTGCCCTACCCTTGCCACTCCTCGCCTGACTCGTTTAGTTTAAGCCCACCGCCCCCGCCAT
 401 TCTCCTCTAGCATCACACCAACGGCGGCACCCGCTCCGCCTCCACCATCTGCTTTTCGCCACATCAGAAACTGGTTTTTGGAGCATCTGGAACACCCGTT
 501 CCCTACACCTGACGAAAAAGCCGCGTTAGCCAAAGCAAAAGGTTTAACTGTTCCACAAGTTAACGGTTGGCTAACAAATATGAGGCGCAGGTCGGGTTGG
 601 ACTCAAACAGAAAGGGATTTCGCCAACGCAGATCATGAACAAATGAGAGCGCTCGTGCTAGCCTTTCGGAGAGGCCCTTCCGGTGGCGTAGCCAGAGATG
 701 TCGTGTTGGCCCTAGAAAAAATAGAAGACTATGTGGATGAAAGGGGACAAGAAAGAGTCAGAGATTGGATGGTAGGTGTGATCGAGCAAACCGCTAAAGA
 801 CGTACAAGAAGGAAAGGACGTTCTGGGTGAGGATTCCGAGGATGAAGAAGGAGCATTTGATGAGGATGACGGAGAAGCGCCCAGTACGAGTGCGGCCGTC
 901 AGCGAAGCTGTGGGGAAGAAACGTAAAACAAGTTCTTTGGCTGCACCATCCGCCAGCTCAGTCAAACAACCTAAATCTAAACCAGCGCCAAAGACTGAAG
1001 ATCGTATTCTGAAGCCGTTGCCGAAGTTGAGGCTCGTAAGGCCTTTGTCATCAACTAGTTACTCTGCAACTGTTCCCGCCAGTCCGCCACCATTGCCCCC
1101 AAATGCGAGGACCATTACCCCGCCCTACAGTCGTCACCCAAGTGAGCCTAGAAGTACTGTTGGCTTGGGGTACCCATCTGTTTCAACTAATCCTCATCAA
1201 CATAGAACCTCTTCTGTCAGAAATTCATCAGGAACATCATCACTAGAAACCGCCGTTGGACCGGGAACTCCGGAAGAATTCGAAGGCGACTTCGAGCCAA
1301 ATGCTGTTTTTCAACAACAACATGAGTTAGGCGTTTTTTCAGCTCCCTTACCGCCTCCTTTGTCCCAATACCCCTCTCACGCCCCCCCCCCCCATCAAGC
1401 ATATTACCCTTCCCACTCACCATTGCCTCCCCCCCACTACTTTGCTCCTTTGGCTCCCGCATTACCACCTTCCCATTCCGATGTTTTTCCTCACCCATAC
1501 GCATATGATCCACCTTATTCCCACGCTAATTTTCCACCACCTCCTCCACCAACTAATGGCCGTTTAGCACACCCTCGTACATATTCTTGGTCAGCTGGTG
1601 AAGCTCATCCTCCACCACAACCACCAGCTAAGAAGCATAAGAGAAGAGCTACATCTGATCAACATCCTTCTACATTATCATCTGCACATGCTTATTATCA
1701 TCCAGCACCACAAGAACCTTATCCATATGCTCCACCTCATGCTTTTTCTCCTCCTCATCCATCTTATTTACATCCATTACCTCCACCTGCTCATGCTGCA
1801 CCACCACCACCACAAACAGATGATGATAGACATGCAGCTTTATTAGCTAGAAAGGAAAGAATGAGAGAAGAGCAAGAACAAATTAGAAGAGAGGAGGAAG
1901 CTTTAGCTTCTGGTGGTGGTGGTGGTGGTTCTGGTGGTGGTCAATAA
```

C. ferigula PYCC4410 *HD2* synthetic gene sequence

```
   1 ATGCCGGATGATACTGCTCTTCTGCAGAATGTTTTGAAGCTTGCACAGGAGTTAGGTCAGTCTGCTGGAGTCCCGCTCAGTACAGGAAGACCACTTACCT
 101 TACCGGAGTACCTCCCGCTGGATCTCCCACCTCCCCCCACCTTGTCGAGATATCTAGAGTCGATTGGGATTTCACCGGCTTTGCAAGCAGCTCTGCAACT
 201 TGCATTCGAAAAAAGAATCCGAACATTTGCCATATCAACTCAACAATTGTATCAGCGATCGTCTGGGGGAGTGAGTGCACCATTGGGTGTCTTGGAGGAG
 301 GTCAATGGGTCTAGGGGCTACGCGGCTAGATTGTTCCGTGACTTCCGTGCTCAAGTGATAGCCCACTTTGTAACCGAAGTTGCCAAATTGCAAGCCATGG
 401 TTTTAAGCAAGGCAAGGGACCATGTACTGAAGGCTGGATCCACTAAACCGCCTCCCCAGTTCACTAAAGCTCACCCATTGAGCAGTCATCCTCCCACTAG
 501 CGCTATAGAAACACTGCGGAAGCTGTACTTGTTAAATACTTACCCTAACCCTGCAGAAATACGTTTGATAGCTAAAAAAATCCAGATGGATTACAAACAG
 601 GTGAGGGTCTGGTTTCAAAACCGTAGGAACAGAAAAGCTGTCTCCGCACGCGGACTAAAGGAAGAAGTAAGGATCGCACAGGCTTCCGGCATCAAATTAC
 701 CGTCCACGCACAAGCGTGAGTTGAGAAAACTTGAGACAGCCGCTGCAAAGCAAACTAGGAAGAAGGATGGCGAGAGAAGACGCAGAGGAGGCGTAGTCGT
 801 CACAGCTCAGGATGTACAAGATGAAGACAAAGAAGGCATGCATGTGGACGATGCCGGCATCGCTCCTCCGACAGCCAAACATAGCGCTCTAGACACGGTA
 901 TCCTTCAATTATGGGGATGTGTCAATTCTTGGTTATGGCAGACCACCTCCCGTTTATTTCGGAGCGGGAAGTTCAGAGACATCGCCAGCACCCCACAGTG
1001 CTAGTGCTGCTAGCTCCTCTCTATCGTTGTCTGGTCAACCCCCTCCCCCATCATCCTTAGTGCCCAGTGGTTCTTGGAGCTATGATGCTTCCCCCTCCCA
1101 ATGTTCTCAATCGATGACTATTCCAGATGAACAATGGAATCTTGTTCCAGCTAGACCAGACTACTATCAACAAAGACATTCTGACGGTTCTGTTTTAGTT
1201 CCACCTCCAGCAACGGCACAACCTCATAGAACGACCTCATTCGGTAGCAGCTCAAGTTCATCAGGTCTTTCGGATCTAGATTCCTTGTGGTCAATTGCAA
1301 GTACCGATCCAGCGGCGACTTACTCCTCTTTCACATCAGCTCCTTGTGTTCCTACTGCTTTAGACTTTACCCCTCCTTTGCCTTCTGATGAAAGTACTTT
1401 GGCGGCTGCACATGAAAACCCTCTATTCGAACCACCAGAATATTTATTACAATCTTCAATTCCTGAATTCAGTTTTTCTCCATCTCCTCCACCAGCCCTA
1501 AATCCAGCATGTGCCTCTAATCTAGGTTTTTCTGCAGATGATTTTAGAGCCGTTTTAAACGCCGATCCTTTTTCAGACTCTTCTGCCGACGCGTTAGCGC
1601 AAGCTGACTCTAACGGTTTTTTTTGGAATTGCTCAGGTGGTGCGGGTGCAGACGTTTTAGAATTATCAGCCGAAATTGATCAAGTTTTAAGAGCATTTAC
1701 ACCAGAATTTTTAGGTGCTCCAGGTGCCGAAGGTGAAGCATTTGATCCATATGATTGGGGTATTCCAGCCCCAGCAGATACAACCTATCCAGAACCAGTT
1801 TGGTAA
```

Fig. S4. DNA sequences of the designed synthetic *HD1* and *HD2* genes of *C. ferigula* PYCC 5628 and PYCC 4410, respectively. Sequences correspond to the coding DNA sequences of the *HD1* and *HD2* genes of PYCC 5628 and PYCC 4410, respectively, optimized for the codon usage of *S. cerevisiae*.

Fig. S5 Activation of the reporter genes in the Yeast Two-Hybrid Assay by the individual fusion proteins. This control consists in assessing activation of reporter genes in haploid transformants carrying each of the fusion proteins to be tested. Absence of this so-called autoactivation shows that activation of reporter genes in diploid strains requires the presence of both interacting partners. Controls are the same as in Figure 4. AD, activation domain of Gal4; BD, DNA binding domain of Gal4.