Positive but variable effects of crop diversification on biodiversity and ecosystem services

D. Beillouin [1,2*], T. Ben-Ari [3,4], E. Malézieux [1,2], V. Seufert [5] and D. Makowski [6]

[1] CIRAD, UPR HORTSYS, F-34398 Montpellier, France
[2] HortSys, Univ Montpellier, CIRAD, Montpellier, France
[3] UMR 211, INRAE, AgroParisTech, Université Paris-Saclay, F-78850 Thiverval-Grignon, France
[4] Institut d'écologie et des sciences de l'environnement de Paris (IEES), Campus Pierre et Marie Curie, France
[5] Institute for Environmental Studies (IVM), Vrije Universiteit (VU) Amsterdam, The Netherlands
[6] UMR MIA 518, INRAE, AgroParisTech, Univsersité Paris-Saclay, Paris, France

* *damien.beillouin@cirad.fr*

Abstract

Ecological theory suggests that biodiversity has a positive and stabilizing effect on the delivery of ecosystem services. Yet, the impacts of increasing the diversity of cultivated crop species or varieties in agroecosystems are still under scrutiny. The empirical evidence available is scattered in scope, agronomic and geographical contexts and impacts on ecosystem services may depend on the type of diversification strategy used. To robustly assess the effects of crop diversification in agroecosystems, we compiled the results of 95 meta-analyses integrating 5,156 experiments conducted over 84 experimental years and representing more than 54,500 paired observations on 120 crop species in 85 countries. Overall, our synthesis of experimental data from across the globe shows that crop diversification enhances not only crop production (median effect +14%), but also the associated biodiversity (+24%, i.e. the biodiversity of non-cultivated plants and animals), and several supporting and regulating ecosystem services including water quality (+51%), pest and disease control (+63%) and soil quality (+11%). However, there was substantial variability in the results for each individual ecosystem service between different diversification strategies like agroforestry, intercropping, cover-crops, crop rotation or variety mixtures. Agroforestry is particularly effective in delivering multiple ecosystem services, i.e. water use and quality, pest and diseases regulation, associated biodiversity, long-term soil productivity and quality. Variety mixtures, instead, provide the lowest benefits, while the other strategies show intermediate results.

Our results highlight that while increasing the diversity of cultivated crop species or varieties in

agroecosystems represents a very promising strategy for more sustainable land management,

contributing to enhanced yields, enhanced biodiversity and ecosystem services, some crop

diversification strategies are more effective than others in supporting key ecosystem services.

Keywords: meta-synthesis, agroforestry, crop rotation, intercropping, rotation,

agrobiodiversity, ecosystem services, research synthesis

Introduction

Driven by productivity-oriented policies, agricultural practices throughout the 20[th] century have

converged towards a strong intensification of agroecosystems[1], which was associated with a

large decrease in the number of cultivated plant species[2,3]. These intensive agricultural practices

have not only resulted in a homogenization of agricultural systems, they have also resulted in

agriculture becoming the major contributor of the crossing of several planetary boundaries[4]. A

variety of different already existing agricultural systems have been proposed as more

sustainable alternatives to current intensive systems, to ensure adequate food availability while

minimizing the environmental impacts of food production[5], including, for example, organic

agriculture, agroecological farming, ecological or sustainable intensification[3,6,7]. Most of these

alternative systems share the implicit or explicit goal of increasing agrobiodiversity, i.e. the

diversity of cultivated crop species on-farm and in agricultural landscapes.

Ecological theory suggests that more diverse systems can provide a wider range of different

ecosystem services in higher quantity and at increased stability over time compared to less

diverse systems[8–10]. The mechanisms behind the proposed relationship between biodiversity,

ecosystem function and the delivery of ecosystem services are manifold, complex and debated

but have been summarized in four (not necessarily mutually exclusive) hypotheses: systems

with higher biodiversity i) offer more different niches and thus provide more different functions

(*species complementarity*); ii) have built-in redundancy (i.e. there are more species than there

are functions, *species redundancy);* iii*)* increase resilience, due to their redundancy, over time

in the face of environmental fluctuations (*diversity-stability*); iv) provide extremely variable

relationships between a system's biodiversity and function depending on the type and strength

of species interactions (*idiosyncrasy*) [9,11,12]. In agroecosystems, these mechanisms are further

complicated, since many ecosystem functions are substituted by human labour or by

agricultural inputs[8].

The growing popularity and promise of crop diversification as a tool for more sustainable land

management is evidenced in a fastly growing empirical literature (*Fig. 1a*), as well as numerous

qualitative[13–19] and quantitative[20] synthesis. But when synthesizing this vast body of literature

to provide guidance to decision-makers about the promises and shortfalls of these practices, it

is crucial to (1) start with a clear definition and conceptualization of what agricultural

diversification means, (2) provide a systematic and global synthesis of the current evidence

base, while (3) addressing and exploring bias and heterogeneity in the data analysed, as well as

(4) exploring the quality and robustness of the results.

Our objective is hence to provide a rigorous global synthesis of the effects of various crop

diversification practices at field scale on biodiversity and ecosystem services. We define crop

diversification as any type of agricultural diversification practice that involves increasing the

diversity of cultivated annual or perennial crops (including non-harvested crops like cover crops

or catch crops) at genetic or species level. To allow for additional insights into the impacts of

different types of crop diversification practices we group crop diversification strategies into 5

categories: *crop rotations* (i.e. temporal succession of a set of selected crops grown on a field

each season or each year), *cover crops* (i.e. plant sown in addition to the main crop for

agronomic or environmental purposes), *agroforestry* (i.e. inclusion of at least one woody

perennial species), *intercropping* (i.e. the simultaneous cultivation in the same field of two or

more different crop species for all or part of their growth cycle) and *variety mixtures* (i.e. the

simultaneous cultivation in the same field of multiple varieties of the same crop species; *Fig. 1*;

for a more detailed definition of the crop diversification practices see Supplementary

Materials). To provide a synthesis of the empirical evidence we compile an exhaustive database

of published meta-analyses on the subject and perform a meta-analysis of published meta-

analyses, i.e. a second-order meta-analysis. This approach has been scarcely applied in

agronomy so far, but is already common practice in the medical sciences[21]. We also focused on

assessing the quality of selected meta-analyses and their redundancy (i.e., how many common

primary studies are shared between distinct meta-analyses). The overall summary effect sizes

computed in our analysis are weighted according to these criteria (i.e., we perform a weighted

second-order meta-analysis, see methods).

Figure 1. The five crop diversification strategies included in the analysis, the temporal
evolution of the number of meta-analyses studying this practice (A), as well as illustrative
examples of each practice (B-F). The colors of the border of each photo correspond to the
color of the crop diversification strategies in panel A. The crop diversification strategies are:
(B) crop rotation (example: wheat and canola rotation), (C) cover-crops (here, a mix of oats,
buckwheat, turnip, radish, mustard, millet, soybean, and hairy vetch in wheat residues, USA),
(D) agroforestry (here, peanuts and acacia, Senegal), (E) intercropping (here, organic cotton
cultivation combined with pineapple and banana crops, Paraguay), and (F) cultivar mixture
(here, wheat mixtures, France). Note that one meta-analysis could study one or multiple crop
diversification strategies. For a detailed definition of each crop diversification practice see the
Supplementary Materials. Credits of photo are in supplementary.

Results and Discussion

We identified 95 meta-analyses synthesizing the effects of increasing crop diversity in agrosystems at the field scale. Our dataset includes the results of 5,156 primary studies equivalent to 54,554 experiments published since 1936, spanning more than 120 crops and 85 countries in Africa, America, Asia, Europe, and Oceania. Different crop diversification practices vary in the degree they have been studied through experimental trials and ensuing meta-analyses: cover-crops (35 meta-analyses; 23,000 experiments), crop rotation (20 meta-analyses; 3400 experiments), intercropping (20 meta-analyses; 13,000 experiments), agroforestry (19 meta-analyses, 6,900 experiments), variety mixtures (5 meta-analyses; 7,800 experiments), (*Fig 1a*). These experiments and meta-analyses examined a broad suite of different agronomic and environmental outcomes, which we summarize into the following biodiversity and ecosystem services categories (for further details about how each category was defined see Supplementary Materials): agricultural production (40 meta-analyses; 23,000 experiments), soil quality (32 meta-analyses; 13,000 experiments), associated biodiversity (14 meta-analyses; 6,500 experiments), pest and disease control (12 meta-analyses; 3,100 experiments), water quality (11 meta-analyses; 1,450 experiments), water regulation (7 meta-analyses; 1,400 experiments), greenhouse gas (GHG) emissions (7 meta-analyses; 1,300 experiments), input use efficiency (5 meta-analyses; 2,000 experiments), product quality (4 meta-analyses; 1,800 experiments), profitability (3 meta-analyses; 100 experiments), yield stability (1 meta-analysis; 390 experiments).

Crop diversification improves biodiversity and ecosystem service delivery

Considered together, the various crop diversification strategies examined improved associated biodiversity (median effect estimate equal to +24%; 95% confidence interval CI: 15%, 33%), as well as the majority of ecosystem services analyzed (*Fig. 2*), for example resulting in

improved water quality (+51%; 3%, 123%), more efficient pest and disease control (+63%;

CI: 31%, 103%), better soil quality (+11%; CI: 7%, 16%) and higher production levels

(+14%; CI: 8%, 20%). Our results are thus in line with the positive effects found for highly

diverse plant communities in natural ecosystems[22–25] and a previous second order-meta-analysis

for diversified agricultural systems[20].

The mechanisms behind the positive effect of crop diversification on associated biodiversity,

regulating and supporting, as well as provisioning ecosystem services are manifold. Crop

diversification can, for example, provide access to additional resources, or improve the diversity

and connectivity of habitats for wildlife[13]. Regulatory services like pest and disease control,

water or soil quality are, in turn, supported by the higher levels of associated biodiversity, as

well as by changes in habitats and resource competition in more diversified systems. Globally,

increasing crop diversity has, for example, a strong positive effect on all sub-categories of pest

and disease control examined, i.e. weed reduction (60, CI:13%, 116%), pests reduction (33, CI:

2%, 74%), disease control (41, CI: 15%, 73%) and reduction of crop damages (62, CI: 37%,

93%). Increasing crop diversity could alter weed dynamics by limiting available resources

(light, water, minerals) for weed species[26]. The increased canopy complexity of diversified

systems also provides refuge to natural enemies[27,28], reducing the spread of pests and diseases

through microclimatic changes or physical barriers[29]. Similarly, crop diversification also

improved all measured indicators of soil quality: soil physics, soil chemistry and soil carbon

content (see supplementary Materials). The latter indicator is of particular interest to assess the

potential for diversified systems to contribute to soil carbon sequestration and climate

mitigation (although whether and to what degree agricultural practices can result in long-term

carbon accumulation in soils is still debated[30,31].

The higher levels of provisioning services observed in diversified systems could largely be

driven by these positive effects on regulatory ecosystem services[32–35]. Garibaldi et al., 2018

showed that synergistic effects (positive interactions) between yield and regulatory services

generally outnumber negative ones, suggesting that positive impacts of crop diversification on

regulating services like soil quality and pest control lead to the positive impacts on provisioning

services observed[36].

The results of our analysis are, however, inconclusive for the regulating and supporting services

of water regulation (+19%; CI: -1%, 44%) and GHG emissions (+25%; CI: 0%, 44%) despite

thousands of experiments for these services (1400 and 1300 experiments, respectively). It is

also important to note that GHG emissions is the only outcome examined that shows a negative

trend - crop diversification practices thus might actually increase GHG emissions, but with

results being non-significant due to high uncertainty and/or few syntheses. In addition, our

literature synthesis, although exhaustive, shows that there is currently insufficient aggregated

data to precisely evaluate the impacts of crop diversification on input use efficiency, yield

stability, product quality and profitability - even though the tendency for these outcomes are

also positive (i.e., improved input-use efficiency, and product quality, increased profitability).

Our results are robust to the type of statistical model used and to publication bias (see

Supplementary Materials).

a. Biodiversity

b. Supporting & regulating ecosystem services

c. Provisioning ecosystem services

Effect size of crop diversification (lnRR)

182
Figure 2. **Impacts of crop diversification (all strategies) on associated biodiversity, supporting and**
regulating ecosystem services and provisioning ecosystem services. Points, thin lines and thick lines
represent the estimated median summary effect, 95% confidence intervals (CI), and 95% prediction
intervals (PI- see methods) of the logRatios estimated from a weighted mixed effect model. CIs indicate
the range of values that likely contains the mean estimate of the data analysed, while PIs indicate the
range of values that would likely contain the mean estimate if an additional new meta-analysis would
be added. The number of meta-analyses and experiments are displayed between parentheses below each
category. Colored shaded areas represent the distribution probability of the effect sizes of the meta-
analyses.

Different crop diversification strategies vary in the magnitude of their positive impacts

Despite these often-clear positive outcomes across all diversification practices, we observe high

heterogeneity among the results of different meta-analyses (I2 statistics higher than 90% for

most ecosystem services- Supplementary Materials), which implies that the effects and their

magnitude depend on the crop diversification practice and/or bioclimatic conditions. Further

analysis reveals that a substantial part of the variability of the mean effect sizes is due to

significant differences in performances between the five crop diversification strategies.

The global positive effect of crop diversification on **associated biodiversity** strongly depends

on the type of strategy considered (p-value <0.001 – *Fig. 3; Supplementary Materials*). The

biodiversity benefits are the largest for agroforestry (+61%, CI: 26, 105%), crop rotation

(+37%; CI:16,62%) and cover crops (+21%, CI: 17, 25%)- which are all strategies that are often

associated with heterogeneous non-crop habitats. Note that we also find substantial differences

between different types of agroforestry practices (i.e., +9% of associated biodiversity for

silvopasture; +45% for alley cropping; +62% for parklands, +86% for shaded perennial

systems, see *Fig. 3*; for illustrative examples of these different agroforestry practices and for

detailed definitions see Supplementary Materials). Intercropping and variety mixtures, on the

other hand, have smaller or non-significant effects on associated biodiversity (+7%; CI: 3,12%

and CI: 2; -12,18%, respectively).

The variability of **agricultural production** observed for aggregated results is, also, largely

explained by the different crop diversification strategies (67% of the variability of the effect

sizes - p< 0.02 – *Fig. 3, Supplementary Materials*). The two diversification strategies relying

on the simultaneous cultivation of different plant species within one field (i.e., agroforestry and

intercropping) stand out as improving agricultural crop production most strongly (+35%;

CI: 12%, 62%; +22%, CI: 14%, 32%, respectively; note that we do not consider tree production

for agroforestry). Including an additional crop in a rotation leads to a 16% median increase

(CI: 11%, 21%) in production levels of the following crops. Cover crops (+6%; CI: 1%, 12%)

and variety mixtures (+2%; CI:1%, 3%) promote modest but still significant yield

improvements compared to more simplified systems. The remaining variability of the effect

sizes within each crop diversification strategy (reflected by the confidence intervals for each of

the five strategies) could be caused by the diversity of conditions covered (e.g., a variety of

agricultural practices, climates and soil characteristics, crop species or taxa, and initial

diversification levels of the control plot). For agroforestry, for example, the different types of

systems considered differed in their impact on agricultural production (i.e., -19% production

for perennial shaded systems; +3% for hedgerows; +31% for alley cropping, +47% for

parklands, see *Fig. 3*). For the other provisioning ecosystem services (i.e. product quality, input

use efficiency, yield stability and profitability), the numbers of meta-analyses by crop

diversification strategies are rather low, and do not allow a detailed comparison between

strategies.

Most **supporting and regulating ecosystem services** also depend on the type of crop

diversification strategy implemented. Pest and disease control is, for example, improved by all

crop diversification strategies, but with different magnitudes of effects (*Fig. 4*). Our results

suggest that the benefits are highest when using cover crops (+125%; CI: 83%, 178%), while

intercropping (+66%; CI: 40%, 98%) and agroforestry (+59%; CI: 38%, 82%) still have high

but slightly smaller positive impacts. For agroforestry, available evidence shows strong positive

effects on pest and disease control for hedgerows (median effect = 84%) and shaded perennial

systems (median effect = 40%; *Fig. 4*). The benefits of crop diversification for soil quality is

also variable across strategies (p-value < 0.08- *Fig. 4*). The benefits are particularly important

for agroforestry and intercropping (+19%; CI: 16%, 23% and +11%; CI: 5%, 18%

respectively). Crop rotation promotes modest soil quality improvements (5%; CI: 2%, 8%), and

variety mixtures have non-significant effects (-5, CI: -25%, 18%). For agroforestry, the median

effect is -12% (non-significant) for silvopasture, +13% for hedgerows, +17% for alley cropping

and +21% for parklands (*Fig. 4*). Other supporting and regulating ecosystem services such as

water regulation (which includes variables like run-off reduction or soil water infiltration,

Supplementary Material) also vary between strategies: Agroforestry leads to significant

improvements (45%, CI: 13, 87%) while crop rotation and cover crops effects are non-

significant (18%, CI: -5, 48% and 10, CI: -10, 34%, Supplementary Materials). On the contrary,

no differences are observed among strategies for water quality, partly because of the very large

estimated confidence intervals (89%, CI: 20, 200% for agroforestry; 87%, CI: 36, 156% for

intercropping and 61, CI: 12, 132% for cover crops).

255
Figure 3. **Impacts of the five crop diversification strategies on biodiversity (A.) and agricultural**
production (B.). Results are presented globally over all strategies (a.), for each of the five crop
diversification strategy separately (b.), and for selected agroforestry practices (c.). The numbers of meta-
analyses and experiments are displayed between parentheses below each category (NA indicates the
absence of reported information the meta-analyses) on the left. Points, thin lines and thick lines represent
the estimated median summary effect, 95% confidence intervals (CI), and 95% prediction intervals (PI)
of the logRatios estimated from a weighted mixed effect model. The numbers between parentheses in
sub-plots b. represent the p-values of statistical tests of differences between crop diversification
strategies.
265

266
267
Figure 4. **Impacts of the five crop diversification strategies on soil quality (A.) and pest**
and disease control (B.). Results are presented globally over all strategies (a.), for each of
the five crop diversification strategy separately (b.), and selected agroforestry practices (c.).
The numbers of meta-analyses and experiments are displayed between parentheses below
each category (NA indicates the absence of reported information the meta-analyses) on the
left. Points, thin lines and thick lines represent the estimated median summary effect, 95%
confidence intervals (CI), and 95% prediction intervals (PI) of the logRatios estimated from a
weighted mixed effect model. The numbers between parentheses in sub-plots b. represent the
p-values of statistical tests of differences between crop diversification strategies.
277

Climate impacts of crop diversification are variable and context-dependent

Given the importance of agriculture as a contributor to but also a potential solution to

anthropogenic climate change[37] , any sustainable agricultural practice needs to be scrutinized

for its climate mitigation potential. We find that agroforestry (+19%; CI: 14%, 24%),

intercropping (+13%; CI: 6%, 10%) and cover crops (+13%; CI: 10%, 15%) contribute

significantly to increased soil carbon content (see Supplementary Materials). On the other hand,

both crop rotations and crop variety mixtures have a small and non-significant impact (+3%;

CI: 0%, 4% and -5%; CI: -24%, 4% respectively). These different effects of different crop

diversification strategies on soil carbon content may be linked with the fact that some strategies

such as agroforestry, intercropping and cover crops tend to produce more above- and below-

ground litter, which favors the formation and accumulation of soil organic carbon[38] (see

Supplementary Materials). This is in line with other recent studies that have highlighted the

promises of agroforestry or cover crops for sequestering carbon in agricultural soils[39,40]. To

contribute to soil carbon sequestration and climate mitigation, such practices need, not only

provide additional biomass and carbon uptake but also (i) not result in a trade-off with yields

(which would result in displacement effects), (ii) not result in the increase of other GHGs like

nitrous oxide (N2O) or methane or increased energy use for example for fertilizer production,

and (iii) persist over long periods[41,42]. While all diversification practices improve yields (see

above), we find based on 6 meta-analyses that cover crops actually increase GHG emissions

(+29%; CI: +49%, +1%). Experimental evidence suggests that the use of leguminous cover

crops but also incorporation of cover crops into the soil, as well as the higher availability of

mineralizable carbon under cover crop management can result in an increase of direct N2O

emissions compared to non-cover crop management[43,44]. Yet, a literature synthesis assessing

the net GHG balance of cover crops suggests that improved soil carbon stocks, as well as

decreased indirect N2O emissions due to reduced nitrate leaching could outweigh the

sometimes increased direct N2O emissions[44]. Another shortcoming of the use of cover crops

for carbon sequestration is that cover crop management can be easily reversed and carbon stored

released again in following years[41]. The net benefits of cover crops for climate mitigation are

thus complex and appear to vary depending on factors like cover crop species, climate and

tillage practices[41].

More diversified agroforestry systems, instead, do not only improve yields (*Fig. 3*), result in

more below-ground soil carbon (Supplementary materials), as well as more above-ground

biomass (*Fig. 3*), but - being perennial systems - they are also likely to persist longer[45] and

some analyses suggest that they reduce other soil-based GHG emissions like N2O[46] (which we

were not able to assess due to a lack of agroforestry meta-analyses on GHG emissions). These

important differences in the performances of different crop diversification practices for soil

carbon and climate mitigation highlight the need to unpack such differences instead of treating

crop diversification as a monolithic and always sustainable practice.

Knowledge gaps and future research agenda

The transition to more sustainable diversified systems is a long and constantly evolving process

requiring up-to-date and comprehensive multidisciplinary evaluation. The data gaps that we

identified on some important outcomes like yield stability or farm profitability could partly

explain the actual low rate of adoption of more diversified strategies in practice[47,48]. Most of

the scientific literature on crop diversification has, to date, focused on ecological outcomes and

production. Other agronomic outcomes are, instead, rarely studied. For example, few results

are available on the economic performance of crop diversification strategies (*Fig. 2*), certainly

a major factor limiting the rate of adoption of any new agricultural practice[47]. Restoring

ecosystem services or substituting ecosystem functions provided by petrochemical energy will

only likely be realized by farmers and land users if there is a direct economic or other benefit

to be derived from it[8]. Similarly, evidence on the stability of the production of diversified crop

management is scarce (only one meta-analysis, *Fig. 2*), despite the abundant theoretical

literature that suggests d that diversified agro-ecosystems might be better able to cope with

perturbations (e.g. [5,48]). More scientific evidence on the impacts of crop diversity on product

quality (e.g. micronutrient content) and input use efficiency - also two variables of interests to

farmers and consumers - could also favor a wider adoption.

Beyond the field scale that constitutes the basis of this study, it is also important to maintain

ecosystem services at larger scales, such as farm or landscape, and to promote a higher diversity

of land-uses (e.g.[49]). Most of the ecological processes are strongly dependent on the complex

interactions between land use, biophysical context and human activity at the landscape scale,

such as water quantity and quality, pollination, pest regulation or carbon storage[50,51].

Conclusion

Our results, based on the largest assessment examining effects of increasing the diversity of

cultivated crop species or varieties in agroecosystems, show that these effects are positive but

variable and depend on the type of crop diversification strategy considered. Our analysis is

based on hundreds or even thousands of field-scale experiments where the effects of increased

crop diversity have been compared to less diverse agrosystems. Our synthesis allows the

identification of strategies that effectively and consistently reduce air, soil and water pollution

and increase biodiversity, pest and diseases control and crop production. Agroforestry strikes

out as a particularly promising strategy that is able to substantially increase all the ecosystem

services considered in our analysis (and for which sufficient data was available to allow drawing

robust conclusions), i.e. associated biodiversity, production, water regulation, water quality,

pest and diseases control and soil quality. The other diversification strategies also have

important effects on specific ecosystem services, e.g. cover crops and intercropping on water

quality and pests and disease control or crop rotations on associated biodiversity and

production. Variety mixtures, instead, provide the lowest benefits for most of the ecosystem

services considered.

Despite a sometimes-uneven geographic repartition of available studies[19], our exhaustive and

quantitative review of the literature, by precisely weighing the evidence based on its quality,

redundancy, and precisely analyzing its heterogeneity, provides a solid basis for policy makers

and scientists alike to build future decision-making and future research upon the existing

knowledge base. It can also contribute to help the Food agricultural Organization (FAO)

monitoring and orienting policies towards Sustainable Development Goals or provide support

for the new European "Green Deal" as a starting point for regional debates on the effectiveness

of the various strategies analyzed globally[52]. These broad conclusions should, however, not

hide the fact that heterogeneity exists and that the results of a single meta-analysis could

sometimes diverge from our mean estimates, as indicated by the wide prediction intervals found

in our analysis. Agro-ecological transitions must therefore be designed to take into account

local contexts and constraints[53]

Methods

Data collection and extraction

We identified gray and published meta-analyses on the impacts of crop diversity on biodiversity and ecosystem services through a systematic search using Web of Science, CAB abstract, Greenfile, Environment Complete Database, Agricola and Google Scholar. The search was performed in May 2020 and updated in November 2020 with the equation: *(meta-analysis OR meta analysis) AND (cropping system OR crop* OR agriculture) AND ((rotation OR Diversification OR intercrop* OR cover crop OR mixture) OR (organic AND (system OR agriculture)) OR (conservation AND (system OR agriculture)) OR no till* OR agroforestry OR agroecology).* The query was carried out on the article title, abstract and keywords, with no restriction on the date of publication. We also screened the references cited in each selected meta-analysis. This resulted in a total of 984 publications (844 once duplicates removed – see Supplementary Materials). Only publications satisfying the following criteria were retained for the analysis (i) the publication reports the results of a quantitative analysis based on several primary studies, i.e. a meta-analysis, (ii) The meta-analysis focus on at least one crop diversification strategies defined in table S1, (iii) The adjacent control plot used in the meta-analysis is less diversified as the treatment (we exclude comparisons to diversified natural ecosystems), (iv) the meta-analysis present indicators of variability of the effect-sizes. Studies dealing with pure forestry or wood production were excluded. Based on these criteria, 95 meta-analyses were selected.

All quantitative measures of the effects of crop diversity compared to the less diversified reference, i.e effect-sizes, were extracted from text, table or figure using Plot Digitizer (http://plotdigitizer.sourceforge.net/). The effect-sizes of each meta-analysis were precisely documented (unit, name of the variable, soil, climate, management conditions, etc.). We also retrieved the number of primary studies that were used to calculate the effect-sizes, the number of experiments (one primary study can provide several independent estimates), and the confidence intervals or other indicators of dispersion, when available. Effect sizes expressed as standardized mean difference or percent changes were then converted into ratios, to ensure the comparability of the results between the meta-analyses[54].

Different meta-analyses could be partly based on the same pool of individual experiments. To handle

this pseudo replicated data for our second-order meta-analysis, we retrieved all the references of primary

studies of each meta-analysis (see Supplementary Materials). We then calculated a redundancy index

between each pair of meta-analyses, and included it in our meta-analytical model (see *DataAnalysis*

section).

The meta-analyses differ in the quality of the methodology used to retrieve the primary study and

analyze the data[19,55] which can result in biased and misleading results[56]. To lower the importance of

low-quality studies in our second-order meta-analysis, we calculated a quality score, based on 20 criteria

analyzing the literature review, statistical analyses and analysis of potential bias of each meta-analysis

(see Supplementary Materials). We then integrate this score as a proxy of the internal quality of the

meta-analyses in our meta-analytical model (see *Data Analysis section*).

All retrieved data are freely available on internet web platform:

https://cropdiversification.shinyapps.io/Crop_diversification_2020

Data analysis

We categorized the retrieved response variables reported in the meta-analyses into main- and sub-

categories of biodiversity, supporting and regulating ecosystem services and provisioning ecosystem

services. The 12 main categories are associated biodiversity, pest and disease control, water quality,

water regulation, soil quality, greenhouse gas (GHG) emissions, agricultural production, product

quality, yield stability, profitability and input use efficiency. Sub-categories are shown in Supplementary

Materials.

We quantified the impacts of crop diversity on each main and sub-categories of biodiversity, ecosystem

services. We then detailed the impacts for each crop diversification strategy (e.g. agroforestry,

intercropping). The global form of our models is:

$$Y_{ij} = \mu + b_i + \varphi_{ij} + \varepsilon_{ij} \qquad (Eq.1)$$

$$\text{with } b_i \sim N(0, \tau^2), \varphi_{ij} N \sim (0, \upsilon^2) \text{ and } \varepsilon_{ij} \sim N(0, {\sigma_{ij}}^2)$$

Where Y_{ij} are the j[th] effect-size of the i[th] a meta-analysis, μ is the mean effect, b_i is the random meta-

analysis effect, φ_{ij} is the random effect-size effect within the ith meta-analysis, and ε_{ij} is the random

estimation errors associated with the j[th] effect size of the i[th] meta-analysis. We refer to τ^2 as the between-

meta-analysis variance, υ^2 as the between effect-size variance, and $\sigma_{ij}{}^2$ as the variance of the j[th]

estimated effect-size of the i[th] meta-analysis.

We weighted each meta-analysis by the inverse of the variance of the effect-size, as recommended by

[57]. We then reduced the weight of the studies of lower quality following doi et al., 2008 methodology[58].

We considered the non-independence between the effect sizes of different meta-analyses by calculating

variance-covariance matrix based on a pseudo correlation between meta-analyses[59]. The proxy of the

correlation between each pair of meta-analyses was estimated as, where m is the number of common

primary studies between each pair of meta-analyses, and n_1 and n_2 are the total number of primary studies

in the two meta-analyses, respectively.

We analyzed the heterogeneity of the estimated effect-sizes by calculating i) the 95% confidence-

intervals, i.e. the range of values that likely contains the mean estimates, ii) the 95% prediction

intervals, i.e. the range that likely contains the mean value of a single new observation, a new

meta-analysis in our case, iii) the I2 statistics, i.e. the percentage of variation across studies that

is due to heterogeneity rather than chance[60]. We also precised the percentage of variability of

each biodiversity and ecosystem services explained by the crop diversification strategy. The

added-value of adding crop diversification strategy the models was assessed by a likelihood

ratio test between model including crop diversification as moderators and a null model.

The potential publication biases were assessed with funnel plots and egger tests[61]. The funnel

plots assume that studies with high precision will be plotted near the average mean effect, and

studies with low precision will be spread evenly on both sides. The Egger value tests for the

asymmetry of the funnel plot.

We also test the robustness of our results to the type of model used (with or without random

effect), the type of random effects included in the model (ID meta-analysis and scenario vs.

only ID meta-analyses), and the potential impact of quality of the results (model considering

quality of the meta-analyses or not). The sensitivity of the results against publication bias was

tested with Rosenthal fail-safe number, i.e. the number of additional studies with a mean null

result necessary to reduce the nullify the global estimated effects (see Supplementary Materials).

We also provide an estimation of the mean effect considering the missing studies, based on

trim-and-fill methodology[62].

The parameters of the model were estimated by restricted maximum likelihood. We used log-

transformed ratios for performing the statistical analyses. All statistical analyses were

conducted with R[45] (version 3.0.2), dplyr[63] for data management, metafor for model analysis[64]

and ggplot2 for data visualization[65].

Contribution:

DM, TBA and DB participated in the design and the coordination of the study. DB performs

the data collection, with the help of DM and TBA. DB performed the statistical analyses with

the help of DM, and designed the figures. DB and TBA wrote the draft, all authors provided

critical feedback and contributed significantly to the writing.

Acknowledgments

This work was produced within the framework of the European project 'Diversification through

Rotation, Intercropping, Multiple Cropping, Promoted with Actors and value-Chains towards

Sustainability' (DiverIMPACTS), funded by the European Commission under Grant Agreement

number 727482. It was also supported by the INRAE-CIRAD metaprogram GloFoods and by

the Institute of Convergence CLAND (16-CONV-0003). We are grateful to Mathilde Duvallet

for her contribution to the database.

Competing interests

The authors declare no competing interests.

Data availability statement

The data that support the findings of this study are openly available on an internet web platform:

https://cropdiversification.shinyapps.io/Crop_diversification_2020

482
483
484
485

Literature

1. Hazell, P. & Wood, S. Drivers of change in global agriculture. *Philos. Trans. R. Soc. B Biol. Sci.* **363**, 495–515 (2008).

2. Kleijn, D. *et al.* On the relationship between farmland biodiversity and land-use intensity in Europe. *Proc. R. Soc. B Biol. Sci.* **276**, 903–909 (2009).

3. Pretty, J. & Bharucha, Z. P. Sustainable intensification in agricultural systems. *Ann. Bot.* **114**, 1571–1596 (2014).

4. Campbell, B. M. *et al.* Agriculture production as a major driver of the Earth system exceeding planetary boundaries. *Ecol. Soc.* **22**, art8 (2017).

5. Tamburino, L., Bravo, G., Clough, Y. & Nicholas, K. A. From population to production: 50 years of scientific literature on how to feed the world. *Glob. Food Secur.* **24**, 100346 (2020).

6. Cassman, K. G. & Grassini, P. A global perspective on sustainable intensification research. *Nat. Sustain.* **3**, 262–268 (2020).

7. Potts, S. G., Imperatriz-Fonseca, V. L. & Ngo, H. T. IPBES The assessment report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on pollinators, pollination and food production. (2017).

8. Swift, M. J., Izac, A.-M. N. & van Noordwijk, M. Biodiversity and ecosystem services in agricultural landscapes—are we asking the right questions? *Agric. Ecosyst. Environ.* **104**, 113–134 (2004).

9. Tscharntke, T., Klein, A. M., Kruess, A., Steffan-Dewenter, I. & Thies, C. Landscape perspectives on agricultural intensification and biodiversity – ecosystem service management. *Ecol. Lett.* **8**, 857–874 (2005).

10. Lefcheck, J. S. *et al.* Biodiversity enhances ecosystem multifunctionality across trophic levels and habitats. *Nat. Commun.* **6**, 1–7 (2015).

11. Johnson, K. H., Vogt, K. A., Clark, H. J., Schmitz, O. J. & Vogt, D. J. Biodiversity and the productivity and stability of ecosystems. *Trends Ecol. Evol.* **11**, 372–377 (1996).

12. Vandermeer, J., van Noordwijk, M., Anderson, J., Ong, C. & Perfecto, I. Global change and multi-
species agroecosystems: concepts and issues. *Agric. Ecosyst. Environ.* **67**, 1–22 (1998).

13. Landis, D. A. Designing agricultural landscapes for biodiversity-based ecosystem services. *Basic*
Appl. Ecol. **18**, 1–12 (2017).

14. Tscharntke, T. *et al.* Global food security, biodiversity conservation and the future of agricultural
intensification. *Biol. Conserv.* **151**, 53–59 (2012).

15. Rosa-Schleich, J., Loos, J., Mußhoff, O. & Tscharntke, T. Ecological-economic trade-offs of
Diversified Farming Systems – A review. *Ecol. Econ.* **160**, 251–263 (2019).

16. Kremen, C. & Miles, A. Ecosystem Services in Biologically Diversified versus Conventional
Farming Systems: Benefits, Externalities, and Trade-Offs. *Ecol. Soc.* **17**, art40 (2012).

17. Isbell, F. *et al.* Benefits of increasing plant diversity in sustainable agroecosystems. *J. Ecol.* **105**,
871–879 (2017).

18. Malézieux, E. *et al.* Mixing Plant Species in Cropping Systems: Concepts, Tools and Models: A
Review. in *Sustainable Agriculture* (eds. Lichtfouse, E., Navarrete, M., Debaeke, P., Véronique, S.
& Alberola, C.) 329–353 (Springer Netherlands, 2009). doi:10.1007/978-90-481-2666-8_22.

19. Beillouin, D., Ben-Ari, T. & Makowski, D. Evidence map of crop diversification strategies at the
global scale. *Environ. Res. Lett.* **14**, 123001 (2019).

20. Tamburini, G. *et al.* Agricultural diversification promotes multiple ecosystem services without
compromising yield. *Sci. Adv.* **6**, eaba1715 (2020).

21. Schmidt, F. L. & Oh, I.-S. Methods for second order meta-analysis and illustrative applications.
Organ. Behav. Hum. Decis. Process. **121**, 204–218 (2013).

22. Duffy, J. E., Godwin, C. M. & Cardinale, B. J. Biodiversity effects in the wild are common and as
strong as key drivers of productivity. *Nature* **549**, 261–264 (2017).

23. Loreau, M. & Hector, A. Partitioning selection and complementarity in biodiversity experiments.
Nature **412**, 72–76 (2001).

24. Naeem, S., Bunker, D. E., Hector, A., Loreau, M. & Perrings, C. *Biodiversity, ecosystem*
functioning, and human wellbeing: an ecological and economic perspective. (Oxford University
Press, 2009).

25. Tilman, D., Isbell, F. & Cowles, J. M. Biodiversity and ecosystem functioning. *Annu. Rev. Ecol.*

Evol. Syst. **45**, 471–493 (2014).

26. Weisberger, D., Nichols, V. & Liebman, M. Does diversifying crop rotations suppress weeds? A

meta-analysis. *PloS One* **14**, e0219847 (2019).

27. Gurr, G. M., Wratten, S. D., Landis, D. A. & You, M. Habitat Management to Suppress Pest

Populations: Progress and Prospects. *Annu. Rev. Entomol.* **62**, 91–109 (2017).

28. Langellotto, G. A. & Denno, R. F. Responses of invertebrate natural enemies to complex-structured

habitats: a meta-analytical synthesis. *Oecologia* **139**, 1–10 (2004).

29. Finckh, M. R. *et al.* Cereal variety and species mixtures in practice, with emphasis on disease

resistance. *Agronomie* **20**, 813–837 (2000).

30. Poulton, P., Johnston, J., Macdonald, A., White, R. & Powlson, D. Major limitations to achieving

"4 per 1000" increases in soil organic carbon stock in temperate regions: Evidence from long-term

experiments at Rothamsted Research, United Kingdom. *Glob. Change Biol.* **24**, 2563–2584 (2018).

31. Amundson, R. & Biardeau, L. Opinion: Soil carbon sequestration is an elusive climate mitigation

tool. *Proc. Natl. Acad. Sci.* **115**, 11652–11656 (2018).

32. Dainese, M. *et al.* A global synthesis reveals biodiversity-mediated benefits for crop production.

Sci. Adv. **5**, eaax0121 (2019).

33. Lundin, O., Smith, H. G., Rundlöf, M. & Bommarco, R. When ecosystem services interact: crop

pollination benefits depend on the level of pest control. *Proc. R. Soc. B Biol. Sci.* **280**, 20122243

(2013).

34. Sutter, L. & Albrecht, M. Synergistic interactions of ecosystem services: florivorous pest control

boosts crop yield increase through insect pollination. *Proc. R. Soc. B Biol. Sci.* **283**, 20152529

(2016).

35. Gagic, V., Marcora, A. & Howie, L. Additive and interactive effects of pollination and biological

pest control on crop yield. *J. Appl. Ecol.* **56**, 2528–2535 (2019).

36. Garibaldi, L. A. *et al.* Complementarity and synergisms among ecosystem services supporting crop

yield. *Glob. Food Secur.* **17**, 38–47 (2018).

37. Shukla, P. *et al.* IPCC, 2019: Climate Change and Land: an IPCC special report on climate change,

desertification, land degradation, sustainable land management, food security, and greenhouse gas

fluxes in terrestrial ecosystems. (2019).

38. Wang, Y. *et al.* Below ground residues were more conducive to soil organic carbon accumulation

than above ground ones. *Appl. Soil Ecol.* **148**, 103509 (2020).

39. Chapman, M. *et al.* Large climate mitigation potential from adding trees to agricultural lands. *Glob.*

Change Biol. **26**, 4357–4365 (2020).

40. Bolinder, M. A. *et al.* The effect of crop residues, cover crops, manures and nitrogen fertilization

on soil organic carbon changes in agroecosystems: a synthesis of reviews. *Mitig. Adapt. Strateg.*

Glob. Change **25**, 929–952 (2020).

41. Powlson, D. S., Whitmore, A. P. & Goulding, K. W. Soil carbon sequestration to mitigate climate

change: a critical re-examination to identify the true and the false. *Eur. J. Soil Sci.* **62**, 42–55 (2011).

42. Guenet, B. *et al.* Can N2O emissions offset the benefits from soil organic carbon storage? *Glob.*

Change Biol. **27**, 237–256 (2021).

43. Mitchell, D. C., Castellano, M. J., Sawyer, J. E. & Pantoja, J. Cover crop effects on nitrous oxide

emissions: role of mineralizable carbon. *Soil Sci. Soc. Am. J.* **77**, 1765–1773 (2013).

44. Abdalla, M. *et al.* A critical review of the impacts of cover crops on nitrogen leaching, net

greenhouse gas balance and crop productivity. *Glob. Change Biol.* **25**, 2530–2543 (2019).

45. Searchinger, T. *et al. Creating a sustainable food future: A menu of solutions to feed nearly 10*

billion people by 2050. Final report. (WRI, 2019).

46. Mutuo, P. K., Cadisch, G., Albrecht, A., Palm, C. & Verchot, L. Potential of agroforestry for carbon

sequestration and mitigation of greenhouse gas emissions from soils in the tropics. *Nutr. Cycl.*

Agroecosystems **71**, 43–54 (2005).

47. Marra, M., Pannell, D. J. & Ghadim, A. A. The economics of risk, uncertainty and learning in the

adoption of new agricultural technologies: where are we on the learning curve? *Agric. Syst.* **75**, 215–

234 (2003).

48. Newbold, T. *et al.* Global effects of land use on local terrestrial biodiversity. *Nature* **520**, 45–50

(2015).

49. Redlich, S., Martin, E. A. & Steffan-Dewenter, I. Landscape-level crop diversity benefits biological

pest control. *J. Appl. Ecol.* **55**, 2419–2428 (2018).

50. Swinton, S. M., Lupi, F., Robertson, G. P. & Hamilton, S. K. *Ecosystem services and agriculture:*

cultivating agricultural ecosystems for diverse benefits. (Elsevier, 2007).

51. Li, C. *et al.* Crop diversity for yield increase. *PLoS One* **4**, e8049 (2009).

52. European Commision. *The European Green Deal. COM 2019(640)FInal* (2019).

53. Barrios, E. *et al.* The 10 Elements of Agroecology: enabling transitions towards sustainable

agriculture and food systems through visual narratives. *Ecosyst. People* **16**, 230–247 (2020).

54. Borenstein, M., Hedges, L. V., Higgins, J. P. T. & Rothstein, H. R. *Introduction to Meta-Analysis.*

(John Wiley & Sons, Ltd, 2009). doi:10.1002/9780470743386.

55. Philibert, A., Loyce, C. & Makowski, D. Assessment of the quality of meta-analysis in agronomy.

Agric. Ecosyst. Environ. **148**, 72–82 (2012).

56. Nakagawa, S., Noble, D. W., Senior, A. M. & Lagisz, M. Meta-evaluation of meta-analysis: ten

appraisal questions for biologists. *BMC Biol.* **15**, 1–14 (2017).

57. Marín-Martínez, F. & Sánchez-Meca, J. Weighting by inverse variance or by sample size in random-

effects meta-analysis. *Educ. Psychol. Meas.* **70**, 56–73 (2010).

58. Doi, S. A. R., Barendregt, J. J., Khan, S., Thalib, L. & Williams, G. M. Advances in the meta-

analysis of heterogeneous clinical trials II: The quality effects model. *Contemp. Clin. Trials* **45**,

123–129 (2015).

59. Lajeunesse, M. J. On the meta-analysis of response ratios for studies with correlated and multi-

group designs. *Ecology* **92**, 2049–2055 (2011).

60. Higgins, J. P. & Thompson, S. G. Quantifying heterogeneity in a meta-analysis. *Stat. Med.* **21**,

1539–1558 (2002).

61. Egger, M., Smith, G. D., Schneider, M. & Minder, C. Bias in meta-analysis detected by a simple,

graphical test. *Bmj* **315**, 629–634 (1997).

62. Duval, S. & Tweedie, R. A nonparametric "trim and fill" method of accounting for publication bias

in meta-analysis. *J. Am. Stat. Assoc.* **95**, 89–98 (2000).

63. Hadley Wickham, R. F., Henry, L. & Müller, K. *dplyr: A Grammar of Data Manipulation. R*

package version 0.7. 4. (2017).

64. Viechtbauer, W. Conducting Meta-Analyses in *R* with the **metafor** Package. *J. Stat. Softw.* **36**,

(2010).

65. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York, 2016).

Supplementary materials

Identification and selection of the studies

Figure S1. Prisma flowchart of literature search and screening process. C1: Selection criterion 1 (several individual studies are analyzed); C2: Selection criterion 2 (assessment of the impact of at least one strategy of crop diversification). C3: Selection criterion 3 (control plots are present next to treatment plots)

Definition of the crop diversification considered and illustrative examples

Table S1. Definition of the crop diversification considered

Figure S2. Illustrative example of the six types of agroforestry practices considered in our study. A. Parkland (Faidherbia albida, Pennisetum glaucum and Acacia albida , Senegal); B. Improved fallows (Gmelina_arborea); C. Alley cropping/silvoarable (Lactuca sativa under J*uglans regia* L., France), D. Hedgerows (England), E. shaded perennial crops (Coffea arabica, India), F. Silvopasture (dehesa, Spain).

Text1. Photos credits.

Further details on the effects-sizes, and their variability

Figure S3. Percent of intra-(blue) and the sum of intra-and inter- (orange) variability explained by diversification strategies for different biodiversity and ecosystem services. The numbers at the top of the plot represent the p-values of the test of difference between diversification strategies.

Table S2. Global description of available data presented as ratio and their characteristics. Rosenthal's Fail-Safe Number represent the number of additional 'negative' studies (studies in which the intervention effect was zero) that would be needed to increase the p value for the meta-analysis to above 0.05 (*Rosenthal 1979)*. The heterogeneity (I2) represent the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error. The proportion of total variance that can be attributable to within-cluster heterogeneity is indicated in the last column.

Table S3. Detailed effect-sizes for soil carbon (sub-category of soil quality). The effects-sizes are presented in logRatio for main and sub-categories of crop diversification.
All effect-sizes are available in the web app: https://cropdiversification.shinyapps.io/Crop_diversification_2020

Figure S4. Description of individual variables considered within each Main categories of biodiversity and ecosystem services,

Quality of the included meta-analyses

Table S5. Description of the 20 individual quality items

684 **Figure S5. Boxplot of the quality score for each crop diversification strategies.** The
685 quality score represents the proportion of individual quality criterions met. Precise
686 description of quality items are available in Table S3.
687
688 **Figure S6. Percentage of meta-analyses meeting the 20 individual quality criteria.**
689 Quality criteria are organized in three main groups: review criteria (grey bars), statistical
690 analyses (yellow bars) and bias (blue bars). Precise description of quality items are
691 available in Table S3.
692
693 ## Redundancy of the included meta-analyses
694
695 **Figure S7. Apparent redundancy of primary studies among meta-analyses.** Results
696 on the x-axis are expressed as fractions of common primary studies in two meta-
697 analyses.
698
Analysis of the robustness of our conclusion
Figure S8. Analysis of the impact of potential publication bias. The mean effect-
sizes calculated with a random effect model (ID and scenario as random effects) are
plotted against the same effect-sizes calculated with the trim and fill method (see *Duval*
et al., 2000 for further details). Blue points indicate significant egger tests (p-value
<0.005). Results are presented for the five strategies of crop diversification on the main
ecosystem services. For details on the few categories presenting significant egger test,
see Table S2
707
Figure S9. Analysis of the potential impact of the quality of the meta-analyses. The
mean effect-sizes calculated with a random effect model (ID and scenario as random
effects) are plotted against the same effect-sizes calculated considering the quality of
each meta-analyses in the random effect model (see *Doi et al., 2005* for further details).
Blue points indicate significant egger tests (p-value <0.005). Results are presented for
the five strategies of crop diversification on the main ecosystem services.

719 **Figure S1.**

Strategies of crop diversification	Definition and details
Agroforestry	Agroforestry satisfies three conditions: i) at least two crops interact biologically, ii) at least one of the plant species is a woody perennial, and iii) at least one of the plant species is managed for forage, annual or perennial crop production. We further classified agroforestry practices as follows, based on table 1 of *Cardinael et al.,2018* and from *Nair, 2009: See illustrations in Figure S2* - **Alley cropping**: Fast-growing, usually leguminous, woody species (mainly shrubs) grown in crop fields, usually at high densities. The woody species are regularly pruned and the prunings are applied as mulch into the alleys as a source of organic matter and nutrients. Usually found in tropical regions. Sometimes referred as 'intercropping systems' -Improved **Fallows**: Only sequential agroforestry system considered here. Include both improved and natural fallows - **Hedgerows:** They consist of linear plantation around the fields. They include also shelterbelts, windbreaks and live fences. - **Multistrata systems**: Multistorey combinations of a large number of various trees at high density, and perennial and annual crops. They include home gardens and agroforests. -**Parklands**: Intercropping of agricultural crops or grazing land under low density mature scattered trees. Typical of dry areas like Sahel (e.g. *Faidherbia albida*) - **Shaded perennial-crop systems**: Growing shade-tolerant species such as cacao and coffee under, or in between, overstorey shade trees that can be used for timber or other commercial tree products. - **Silvoarable systems:** Woody species planted in parallel tree rows to allow mechanization and intercropped with an annual crop; usually used for timber (e.g. *Juglans* spp), but also for fuel (e.g. *Populus* spp). Usually low tree density per hectare. Usually found in temperate regions, but not exclusively. - **Silvopastures**: Woody species planted on permanent grasslands, often grazed.
Cover crops	Plant sown in addition to the main crop for agronomic or environmental purposes (e.g., to manage soil erosion, soil fertility, soil quality, weeds, pests, diseases, biodiversity or nitrate leaching). The associated plant could be harvested or not, permanent or not. We further classified associated plant practices as follows: -**Cover crops:** plants that are planted within fields to cover the soil rather than for the purpose of being harvested. Cover crops may be an off-season crop planted after harvesting the cash crop. -**Grass-strips:** an agroecological infrastructure formed by a linear plant cover, often placed on or near the edge of a field, perpendicular to the slope. Also named buffer strip.
Cultivar mixture	The simultaneous cultivation in the same field of multiple cultivars of the same species. All cultivars are harvested. The cultivars could differ for many characters including disease resistance or physical characteristics, but have sufficient similarity to be grown together. For further details, see *Wolfe (1985).*

Intercropping	The simultaneous cultivation in the same field of two or more crops (different species) for all or part of their growth cycle. In this category, we do not include the cultivation with non-woody crops (see Agroforestry category). We further classified intercropping practices as follows (see Brookers et al., 2015 for further details) -**Mixed intercropping:** the plants are grown without any distinct row arrangement, but in an orderly manner. -**Relay intercropping:** the planting of the second crop occur after the first crop has flowered or completed its development or just before harvesting. -**Strip intercropping:** the different plants are grown in strips, allowing crop interactions and independent cultivation. This category also include: -**in flowers/push/pull plants:** Plants or flowers are sown within fields or at field margin for a behavioral manipulation of insect pests and their natural enemies via the integration of stimuli that act to make the protected resource unattractive or unsuitable to the pests (push) while luring them toward an attractive source (pull) from where the pests are subsequently removed (see cook et al., 2006 for further details) - **Inter-row grass cover:** A particular type of cover-cropping, for perennial plantation. The inter-row of the main crops (e.g. vineyard, orchards) are sown with non-harvesting purposes plants.
Rotation	Recurrent succession of a set of selected crops grown on a particular agricultural land each season or each year according to a definite plan. Here, we do not consider as rotation, a system with temporal overlap of two or more crops.

729 **Figure S2**

736   Text 1.
737
738
Figure 1
B. Canola fields: Maria Eklind, Svedala, Sweden, under Attribution-ShareAlike 2.0 Generic
(CC BY-SA 2.0) licence. Flickr. Wheat field. the aucitron. unkown location. under Attribution-
ShareAlike 2.0 Generic (CC BY-SA 2.0) licence. Flickr.
C. Cover crops.   USDA NRCS Montana. USA. Public Domain Mark 1.0. flick
D. Agroforestry. Caroline Dangleant, Senegal. CIRAD Copiright.
E. Intercropping. Pierre Sylvie, Paraguay. CIRAD Copiright.
F. Wheat variety mixture. Mathilde Chen. Copiright.
747
748   Figure S2.:
749   A. Caroline Dangleant. Senegal. CIRAD Copiright.
750   B. Gmelina arborea, Wikipedia, (CC-BY-SA3.0)
751   C. Pierre Sylvie. CIRAD Copiright.
752   D. Unkown location. under Attribution-ShareAlike 2.0 Generic (CC BY-SA 2.0)
753   licence. Flickr.
754   E. Verena Seufert. Copiright
755   F. Verena Seufert. Copiright.
756
757
758
759

Table S2.

Ecosystem services	number of primary meta-analyses	number of effect_sizes	number of paired -data	Rosenthal's Fail-Safe Number	I2
Yield stability	1	4	366	NA	NA
Profitability	3	7	97	456	88,20
Product quality	4	21	1824	36	99,84
Input use efficiency	5	10	2029	2393	98,58
Greenhouse gas emissions	7	13	1299	7257	99,39
Water regulation	7	22	1399	3423	98,76
Water quality	11	33	1453	6722	98,5
Associated biodiversity	14	100	6508	63928	99,37
Pest and disease control	12	59	3081	17866	94,87
Soil quality	32	151	13107	347734	99,92
Production	40	122	23049	101676	99,92

767 **Table S3**

Crop diversification	Sub-category	ES	lower	higher
ALL together	ALL together	0,11137234	0,08777598	0,1349687
Associated plants	ALL together	0,11894124	0,0969786	0,14090389
Agroforestry	ALL together	0,17557429	0,13287615	0,21827243
Crop rotation	ALL together	0,02551628	0,00230497	0,0487276
Intercropping	ALL together	0,12014069	0,05573356	0,18454782
Variety mixture	ALL together	-0,0564958	-0,2842574	0,1712657
Associated plants	grass strips	0,19021408	0,12715615	0,25327201
Agroforestry	silvopastures alley	0,01074449	-0,0516865	0,07317547
Agroforestry	cropping/silvoarable	0,15245917	0,07003978	0,23487856
Associated plants	cover crops	0,10630035	0,08434918	0,12825152
Agroforestry	fallows	0,28912678	0,20529286	0,3729607
Agroforestry	miscellaneous	0,19000209	0,15673624	0,22326795
Agroforestry	parklands	0,28575668	0,19639826	0,3751151
Crop rotation	incl, Legumes	0,00450161	-0,0296758	0,03867906
Crop rotation	miscellaneous	0,0371161	0,00829405	0,06593815
Agroforestry	hedgerows	0,10177642	0,07427732	0,12927553
Agroforestry	multistrata systems	0,27711799	0,19137031	0,36286568
Associated plants	inter-row grass cover	0,13251734	0,12698772	0,13804696
Intercropping	miscellaneous	0,12014069	0,05573356	0,18454782
Variety mixture	miscellaneous	-0,0564958	-0,2842574	0,1712657

   Table S4

Ecosystem services	Details effect-sizes	Number of effect-sizes
associated biodiversity	Animalia	49
associated biodiversity	Bacteria	60
associated biodiversity	Fungi	13
associated biodiversity	Fungi/bacteria	3
associated biodiversity	Miscellaneous	9
associated biodiversity	Plantae	5
greenhouse gas emissions	CH4	1
greenhouse gas emissions	CO2/CO2e	23
greenhouse gas emissions	Energy	1
greenhouse gas emissions	Miscellaneous	2
greenhouse gas emissions	N2O	14
input use efficiency	Fertilizer nitrogen equivalent ratio	5
input use efficiency	N fixed	2
input use efficiency	N use	1
input use efficiency	N use efficiency	2
pest and disease control	Damage	9
pest and disease control	Diseases	4
pest and disease control	Miscellaneous	6
pest and disease control	Pests	25

pest and disease control	Weeds	43
product quality	Fiber	14
product quality	N content	7
production	Grain yield	179
production	Grass production	9
production	Land equivalent ratio	10
profitability	Benefit/margin	5
profitability	Income/revenue	12
profitability	Price/cost	3
soil quality	Carbon stock	98
soil quality	Other soil characterictics	2
soil quality	Soil chemistry	79
soil quality	Soil physics	12
water quality	Erosion/sediment loss reduction	9
water quality	Nutrient leaching reduction	25
water regulation	Infiltration/Drainage	6
water regulation	Miscellaneous	4
water regulation	Runoff reduction	2
water regulation	Water Content	12
yield stability	Stability	4

   Table S5.

Item	description
A protocol is published before the meta-analysis	An a priori review protocol exists, and can be accessed (e.g., Web address), and, if available, provide registration information including registration number.
The aim of the study is clearly mentioned	The list of criteria used to defined the scope of the study are clearly mentioned PICO : Population, variable response, treatment and control.
Literature databases are mentioned	The study describes the databases in the search.
Additional literature search is performed	Additional research (grey literature?)
Search strings are presented	Are the search string clearly defined? All search terms, so that the exact search is repeatable by a third party
Inclusion and exclusion criteria are mentioned	Description of the criteria used to select articles from the original result list (to ensure repeatability). If additional exclusion motives - these are also detailed (e.g., years considered, language, publication status, date last searched)
The number of studies is given at each selection steps	Detailed summary of studies screened, assessed for eligibility, and included in the review, ideally with a flow diagram. Diagram PRISMA
The list of included studies is provided	The study includes the final list of the references of the selected individual studies used in the meta-analysis
The list of excluded studies is provided	The study includes the list of the excluded studies
Method of data extraction is described	Do the authors explicate how data were extracted from the figures?
The dataset is available	Availability of the original dataset? (In results?)
All tools used for the review are mentioned	Name of the program/package used for statistical analysis
Statistical models are described	Statistical analysis of the database: description (model, effect(s), equation...)
Studies are weighted according to their accuracy	Are individual studies weighted according to their accuracy or quality?
Effect sizes of individual studies are presented	Provide information of the size effect for each individual study includes in the meta-analysis? (table, plot)

Confidence intervals of individual studies are mentioned	Provide information of the confidence interval for each individual study included in the meta-analysis? (table, plot)
Quantitative results of the meta-analyses are fully described	Results: Quantitative information on summary effects (effect estimates and confidence intervals, ideally with a forest plot).
Heterogeneity of results is analysed	Is the heterogeneity analyzed using sensitivity analysis or meta-regression?
Publication bias is analysed	Does the synthesis consider possible publication bias?
Funding sources are mentioned	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.

Figure S4.

784   **Figure S5.**

785
786

787 **Figure S6.**

790   **Figure S7**

Figure S8

798 **Figure S9**