Soil microbial diversity impacts plant microbiota more than herbivory

Antonino Malacrinò[1]*, Alison J. Karley[2], Leonardo Schena[3], Alison E. Bennett[1]

Affiliations:

[1] Department of Evolution, Ecology and Organismal Biology, The Ohio State University, Columbus (OH) 43210 USA

[2] Ecological Sciences Department, The James Hutton Institute, Invergowrie, Dundee, UK, DD2 5DA

[3] Dipartimento di Agraria, Università degli Studi "Mediterranea" di Reggio Calabria, Reggio Calabria, Italy

* Corresponding author: A. Malacrinò; malacrino.1@osu.edu and antonino.malacrino@gmail.com

Abstract

Interactions between plants and microbiomes play a key role in ecosystem functioning and are of

broad interest due to their influence on nutrient cycling and plant protection. However, we do not yet

have a complete understanding of how plant microbiomes are assembled. Here, we tested and

quantified the effect of different factors driving the diversity and composition of plant-associated

microbial communities. We manipulated soil microbial diversity (high or low diversity), plant species

(*Solanum tuberosum* or *Solanum vernei*), and herbivory (presence or absence of a phloem-feeding

insect *Macrosiphum euphorbiae*), and found that soil microbial diversity influenced the herbivore-

associated microbiome composition, but also plant species and herbivory influenced the soil

microbiome composition. We quantified the relative strength of these effects and demonstrated that

the initial soil microbiome diversity explained the most variation in plant- and herbivore-associated

microbial communities. Our findings strongly suggest that soil microbial community diversity is a

driver of the composition of multiple associated microbiomes (plant and insect), and this has

implications for the importance of management of soil microbiomes in multiple systems.

Keywords

29   rhizosphere; phyllosphere; bacteria; fungi; metabarcoding; *Solanum*; *potato aphid*

Introduction

Microbiomes can be considered an extension of the plant genome (Berg *et al.*, 2014; Schlaeppi and Bulgarelli, 2015; Rosenberg and Zilber-Rosenberg, 2016; Levy *et al.*, 2018). While their functional importance has been widely dissected in the last two decades of microbiome research, how plant microbial communities assemble, respond to environmental stimuli, and interact with their host remains to be determined (Cordovez *et al.*, 2019; Saikkonen *et al.*, 2020). In addition, research has rarely examined or compared multiple drivers and, to the best of our knowledge, no study has tested the relative strength of different drivers of microbial community composition.

There have been a number of studies identifying individual factors that drive the microbiome composition of plants and their associated organisms and environments. It is well established that plant microbiota structures mainly by plant compartment (e.g. different plant organs, and diverse between endosphere and ectosphere) (Trivedi *et al.*, 2020). Plant genotype and developmental stage have been shown to influence the composition of both plant and soil microbiomes (Wagner *et al.*, 2016). Soil microbiome composition has also been shown to shape plant microbiome composition (Cordovez *et al.*, 2019), and plant pathogens and herbivory produce compositional shifts in plant-associated microbial communities (Lareen *et al.*, 2016). However, we still know little about the relative strength of these factors in shaping plant microbiomes.

Soil provides microbial inoculum and sets the conditions for both plant and microbial growth (Schlaeppi *et al.*, 2014). While different plant tissues can develop distinct microbiomes, soil provides an important reservoir of microbial inoculum for both the phyllosphere and rhizosphere (Bai *et al.*, 2015). The overlap of soil and plant microbiota has been found in a variety of plants, for example *Saccharum officinarum* (de Souza *et al.*, 2016), *Boechera stricta* (Wagner *et al.*, 2016), *Vitis vinifera*

(Zarraonaindia *et al.*, 2015; Mezzasalma *et al.*, 2018) and in the biofuel crops *Panicum virgatum* and

Miscanthus x giganteus (Grady *et al.*, 2019). This suggests that soil microbial communities can

represent a major factor shaping plant microbiomes in both above and belowground plant

compartments. Furthermore, it has been shown that soil microbial community can influence the

feeding behaviour of insect herbivores (Badri *et al.*, 2013). However, only one study to our knowledge

has reported that soil microbial community can directly influence both the aboveground microbiota

of plants (*Taraxacum officinale*) and the microbiota of an insect herbivore (*Mamestra brassicae*),

showing overlap between the microbial communities of the insect and soil (Hannula *et al.*, 2019).

However, it was unclear whether the influence of soil microbiome on the caterpillar's microbiome

was due to passive transfer (e.g. microbe dispersal when watering) or an active colonisation

mechanism (Hannula *et al.*, 2019). Thus, there is potential for the influence of soil microbial

communities on plant-associated microbial communities to extend beyond their plant host.

Plant species and genotype also contribute to the composition of multiple plant-associated

microbiota (Turner *et al.*, 2013). Plant microbial communities assemble differently in different organs,

and their composition varies according to plant phylogeny (Dastogeer *et al.*, 2020). Plant traits (tissue

morphology and physicochemical properties) and resources might drive these plant-genotype effects

on plant microbiota (Fitzpatrick *et al.*, 2018; Dastogeer *et al.*, 2020). For example, an analysis of 30

species of angiosperms revealed differences in the diversity and composition of root microbiomes

across plant species (Fitzpatrick *et al.*, 2018). Similarly, the rhizosphere microbiota of wheat, maize,

tomato and cucumber each had unique microbial communities (Ofek *et al.*, 2014), and leaf and root

microbial communities of *Agave* species clustered according to the host plant species (Coleman-Derr

et al., 2016). Furthermore, several studies reported that host plant identity is an important factor in

the assembly of insect herbivore-associated microbial communities (Colman *et al.*, 2012; Malacrinò,

2018). For example, host plant species influenced the composition of the microbiome associated with

the herbivores *Ceratitis capitata* (Malacrinò *et al.*, 2018) and *Thaumetopoea pytiocampa* (Strano *et*

al., 2018). Therefore, plant species identity has an effect on the communities of microorganisms in

the rhizosphere, living in the different plant organs, and even within plant herbivores.

Only a few studies have tested the effects of herbivory on plant microbiomes. For example,

whitefly infestation of pepper plants led to an increased proportion of Gram-positive bacteria in the

rhizosphere (Yang *et al.*, 2011), and aphid herbivory on pepper plants increased the abundance of

Bacillus subtilis and decreased that of the pathogen *Ralstonia solanacearum* in roots (Lee *et al.*, 2012).

While whitefly herbivory shifted the rhizosphere microbiome composition in pepper plants (Kong *et*

al., 2016), there was no effect of aphid herbivory on the rhizosphere microbiota of *Brassica oleracea*

var. *capitata* (O'Brien *et al.*, 2018). Herbivore attack may alter plant root exudates which, in turn,

promote the assembly of beneficial rhizosphere microbiota (Hu *et al.*, 2018; Rolfe *et al.*, 2019). Thus,

herbivory can alter plant and rhizosphere microbiomes, but the relative impact of herbivory versus

plant species or initial soil diversity on plant-associated microbiomes has not been investigated.

To date most studies have focused on one or two drivers of microbiome composition (e.g., soil

microbiota, plant species or herbivory) and, to the best of our knowledge, no study has tested the

comparative strength of multiple biotic drivers expected to shape above- and belowground

microbiomes *in vivo*. This represents a major gap in our understanding of the relative importance of

factors determining microbiome assembly. Here we ask how plant microbiome composition is shaped

by three different major drivers of plant-associated microbial communities — soil microbial diversity,

plant species, and herbivory — and whether they have equal impact on plant-associated

microbiomes. By manipulating insect herbivory (presence/absence), plant species identity and soil

microbial diversity in a microcosm system, and by quantifying their effects on both bacterial and

fungal plant-associated microbiome composition, we tested the relative strength of these three biotic

factors in shaping rhizosphere, plant (root and shoot), and herbivore microbiomes. As outlined above,

there is evidence of overlaps between soil and plant microbiota, thus we hypothesize that soil

microbial diversity is the major driver structuring plant microbiomes at different compartments.

Previous studies reported that, within the same plant species, geographical location greatly

contributes to assemble the plant microbiota (Christian *et al.*, 2016; Lin *et al.*, 2020). Similarly, there is

evidence that soil microbial community can influence herbivores (Badri *et al.*, 2013) and their

microbiota (Hannula *et al.*, 2019). Thus, we hypothesize that plant species and herbivory will impact

plant-associated microbial communities, but their magnitude would be lower than soil microbial

diversity.

Methods

Experimental design

In this study, we used a 2 × 2 × 3 factorial design to test our hypothesis. We grew two *Solanum*

species (*Solanum tuberosum* and *Solanum vernei*) in soil with different microbial diversities: high

diversity and low diversity (see below). To evaluate the effects of herbivory on plant and rhizosphere

microbiota, we infested plants (within each 'soil' × 'plant species' combination) with two clonal lines

of the polyphagous aphid species *Macrosiphum euphorbiae* (potato aphid); uninfested plants served

as a control. Each treatment combination of plant species (*n*=2), soil microbial diversity (*n*=2), and

aphid clonal line and presence/absence (*n*=3) was replicated five times, involving 60 plants in total.

Study System

Solanum tuberosum (genotype TBR-5642) and Solanum vernei (genotype VRN-7630) seeds were obtained from the Commonwealth Potato Collection at The James Hutton Institute (Dundee, Scotland, UK). Seeds were germinated in steam-sterilized coir, and then transplanted to the experimental pots after 3 weeks.

We used two aphid clones of *M. euphorbiae* (AK13/08 and AK13/18) previously collected in the field (James Hutton Institute, Dundee, UK — 56.457 N, 3.065 W) and reared for several generations on excised leaves of *Solanum tuberosum* (cv. Desirée) in ventilated cups at 20 °C and 16:8 h (light:dark).

All inoculum was prepared from soil collected from an uncultivated field at the James Hutton Institute (56.457 N, 3.065 W) (Bennett *et al.*, 2016; Karley *et al.*, 2017), sieved to 3 cm to remove rocks and large debris, and homogenized. The high diversity inoculum consisted of whole soil, and half of the high diversity inoculum was steam sterilized by autoclaving at 121 °C for 3 h, allowing it to cool for 24 h and then autoclaving it again at 121 °C for a further 3 h. The low diversity inoculum was prepared by blending 50 ml of high diversity inoculum with twice the volume of water, filtering the solution through a 38 μm sieve, and vacuum filtering the collected solution through a Whatman filter paper no. 1 (Bennett *et al.*, 2011). Half of the filtrate was autoclaved at 121°C for 20 min. When preparing the experimental pots, those assigned to the high-diversity treatment were inoculated with high-diversity inoculum (whole soil) and autoclaved low-diversity inoculum (filtrate), while those assigned to the low-diversity treatment were inoculated with low-diversity inoculum and autoclaved high-diversity inoculum. This process allowed us to homogenize the abiotic inputs across our pots while varying the microbial community. For further details please see the section below and Figure S2 (Supplementary material).

The filtration process eliminated larger soil microbes, such as arbuscular mycorrhizal (AM)

fungi, from the low diversity inoculum. We found significant differences in microbial phylogenetic

diversity (see below) between high and low diversity inoculum (Faith's phylogenetic diversity index

38.31±3.66 for high diversity soil, 8.76±5.94 for low diversity soil; $F_{1, 10}$ = 22.05, P < 0.001) which also

translated into different microbial diversities in the rhizosphere (see Results below).

Microcosm setup

Experimental pots (1 L) were assembled as depicted in Fig. S2 (Supplementary material). We

added 100 ml of sterile background soil to the bottom and top of each pot to reduce the risk of

microbial contamination between pots when watering. Sterile background soil was prepared by

mixing Sterilized Loam (Keith Singleton, Cumbria, UK) and sand (ratio 1:1), autoclaving this mixture at

121 °C for 3 h, allowing it to cool for 24 h and then autoclaving it again at 121 °C for a further 3 h.

Between the layers of sterile background soil, we added a mix of 100 ml live or sterile high diversity

inoculum (10% of the pot volume) and 700 ml of sterile background soil. Pots assigned to the high

diversity treatment were filled with live high diversity inoculum and received 1 ml of sterile low

diversity inoculum, while pots assigned to the low diversity treatment were filled with sterile high

diversity inoculum and received 1 ml of live low diversity inoculum. In this way, we controlled for

physical and chemical differences between pots, which only differed in terms of their microbial

community. One potato seedling was transplanted into each pot, pots were randomized into two

blocks, and left to grow in an insect-screened greenhouse with an average temperature of 25 °C and

16:8 h (light:dark) photoperiod.

Five weeks after transplanting, two apterous adult aphids of *M. euphorbiae* clone AK13/08

were added to 20 plants, two apterous adult aphids of clone AK13/18 were added to 20 plants, and

20 plants were left uninfested. All plants were screened with a microperforated plastic bag (Sealed

Air, UK) that allowed transpiration while preventing aphid escape. Three weeks following infestation,

we collected from each pot five aphids, leaves, roots and rhizosphere soil (≈500 mg each) and stored

them at -80 °C. The five aphids were randomly collected from five different leaves on each plant. Leaf

samples were also randomly collected from each plant, being careful to not sample from leaves

infested by aphids. Rhizosphere soil was sampled from soil still adhering to the roots after removing

loose soil, released by vigorously shaking roots and then sampled. Roots were carefully washed with

tap water before collecting a random sample of root tissue. Aphid infestation was scored using a 0–5

scale of severity (scale 0 = no aphids, scale 1 = between 1 and 250 aphids, scale 2 = between 251 and

500 aphids, scale 3 = between 501 and 750 aphids, scale 4 = between 751 and 1,000 aphids, scale 5 =

more than 1,000 aphids).

DNA extraction, Illumina Miseq libraries preparation and sequencing

Samples were crushed in an extraction buffer (10 mM Tris, 100 mM NaCl, 10 mM EDTA, 0.5%

SDS) using three 1 mm Ø stainless steel beads per tube, with the aid of a bead mill homogenizer set at

30 Hz for 5 min (TissueLyzer II, Qiagen, UK). Total DNA was extracted using phenol/chloroform, and it

was subsequently checked for quantity and quality with a Nanodrop 2000 (Thermo Fisher Scientific

Inc., USA). We conducted a metabarcoding analysis for both bacterial and fungal communities of

leaves, roots and rhizosphere soil, and bacterial communities of aphids. Bacterial communities were

characterized by targeting the 16S rRNA gene with primers 515f/806rB (Caporaso *et al.*, 2012). Fungal

communities were analysed by amplifying the fungal ITS2 region of the rRNA with primers ITS3-

KYO/ITS4 (Toju *et al.*, 2012). Amplifications were also carried out on DNA extracted from soil

inoculum, and non-template controls where the sample was replaced with nuclease-free water in

order to account for possible contamination of instruments, reagents and consumables used for DNA

extraction (see Supplementary material).

PCR reactions were performed in a total volume of 25 µl, containing about 50 ng of DNA, 0.5

µM of each primer, 1X KAPA HiFi HotStart ReadyMix (KAPA Biosystems, USA) and nuclease-free water.

Amplifications were performed in a Mastercycler Ep Gradient S (Eppendorf, Germany) set at 95 °C for

3 minutes, 98 °C for 30 s, 55 °C for 30 s and 72 °C for 30 s, repeated 35 times, and ended with 10

minutes of extension at 72 °C. Reactions were carried out in technical triplicate, in order to reduce the

stochastic variability during amplification (Schmidt *et al.*, 2013), and a no-template control in which

nuclease-free water replaced target DNA was utilized in all PCR reactions (see Supplementary

material). We also PCR-tested all root samples for the presence of AM fungi using specific primers

(Lee *et al.*, 2008), finding presence of AM fungi just in plants grown on high-diversity treatment soil.

Libraries were checked on agarose gel for successful amplification and purified with Agencourt

AMPure XP kit (Beckman and Coulter, CA, USA) using the supplier's instructions. A second short-run

PCR was performed in order to ligate the Illumina i7 and i5 barcodes and adaptors following the

supplier's protocol, and amplicons were purified again with Agencourt AMPure XP kit. Libraries were

then quantified through Qubit spectrophotometer (Thermo Fisher Scientific Inc., USA), normalized

using nuclease-free water, pooled together and sequenced on an Illumina MiSeq platform using the

MiSeq Reagent Kit v3 300PE chemistry following the supplier's protocol.

Raw reads processing

De-multiplexed forward and reverse reads were merged using PEAR 0.9.1 algorithm using

default parameters (Zhang *et al.*, 2014). Data handling was carried out using QIIME 1.9 (Caporaso *et*

al., 2012), quality-filtering reads using default parameters, binning OTUs with a 97% cut-off and

discarding chimeric sequences using VSEARCH (Rognes *et al.*, 2016). Singletons and OTUs coming from

amplification of chloroplast DNA were discarded from the downstream analyses. Within the ITS2

dataset, all non-fungal OTUs were discarded using ITSx (Bengtsson-Palme *et al.*, 2013). Taxonomy was

assigned to each OTU through the BLAST method by querying the SILVA database (v. 132) for 16S

(Quast *et al.*, 2012), and UNITE database (v. 8.0) for ITS2 (Nilsson *et al.*, 2019).

Data analysis

Data analysis was performed using R statistical software 3.5 (R Core Team, 2013) with the

packages *phyloseq* (McMurdie and Holmes, 2013), *vegan* (Dixon, 2003) and *picante* (Kembel *et al.*,

2010).

Core microbiota. The core microbiota was identified separately for each compartment using

the package *ampvis2* (Andersen *et al.*, 2018), considering an OTU as a member of the core microbiota

if it was retrieved at a relative abundance of >0.1% in more than 50% of samples. We also tested if the

number of core OTUs shared by pairs of compartments was greater/less than an overlap generated by

random chance. To do so, for each compartment we draw a group of random OTUs from all those

identified in that compartment, in a quantity equal to the number of observed OTUs for that

compartment (i.e., if for a compartment we observed 10 core OTUs, we drew 10 random OTUs from

the pool of OTUs identified in that compartment). Then, we calculated the number of overlapping

random OTUs between all pairs of compartments. We did this separately for bacteria and fungi, and

we repeated it 10,000 times. Then, for each pair of compartments, we selected the highest number of

random overlapping OTUs among the 10,000 permutations and we tested it against the number of

observed overlapping OTUs for that pair of compartments using a χ^2 test.

Phylogenetic diversity. We selected Faith's phylogenetic diversity index (Faith, 1992) to

estimate the diversity of microbiotas in our system because, in contrast to other indices, it takes into

account the phylogenetic relationship between taxa within the community. Comparison of diversity

indices among groups was performed by fitting a linear mixed-effects model, separately for bacterial

and fungal community, specifying *compartment* (i.e., rhizosphere soil, root, leaf, aphid), *soil*

treatment, *plant species*, *herbivory* (and their interactions) as fixed factors and *aphid clonal line* and

block as a random effect (Table 1). We also ran separate analyses for plant (rhizosphere soil, root,

leaf) and aphids obtaining comparable results (Tab. S1 and S2). The use of *aphid clonal line* as a

random variable in the mixed-effects model allowed for the control of differences in the performance

of aphid clonal lines. Models were fitted using the *lmer*() function under the *lme4* package (Bates *et*

al., 2015) and the package *emmeans* was used to infer pairwise contrasts (corrected using False

Discovery Rate, FDR).

Community structure. We analyzed the effects of treatment factors (*compartment*, *soil*

treatment, *plant species*, *herbivory* and their interactions) on the structure of the microbial

communities using a multivariate approach. Distances between pairs of samples, in terms of

community composition, were calculated using a unweighted Unifrac matrix, and then visualized

using a Canonical Analysis of Principal Coordinates (CAP) procedure (Anderson and Willis, 2003).

Differences between sample groups were inferred through PERMANOVA multivariate analysis (999

permutations stratified at the level of *block* and *aphid clonal line*). The use of *aphid clonal line* for

stratification in PERMANOVA allowed for the control of differences in the performance of aphid clonal

lines. We also ran separate analyses for plant (rhizosphere soil, root, leaf) and aphids obtaining

comparable results (Tab. S1 and S2).

Soil diversity vs. plant species vs. herbivory driven effects. Which is strongest? We assessed the

impact of soil treatment, plant species and herbivory for each OTU using the R package *DESeq2* (Love

et al., 2014). Using this package, we calculated the effect of each factor (herbivory, plant species, and

soil microbial diversity) to the abundance of OTU (expressed as abs $\log_2$ Fold Changes) in each plant

compartment. We first built a model using *compartment* (leaves, roots and rhizosphere), *soil*

treatment, *herbivory* and *plant species* as factors. Then, we extracted the appropriate contrasts (*Low*

diversity/High diversity for soil treatment, *S. vernei/S. tuberosum* for plant species and *Herbivore/No*

herbivore for herbivore treatment) for each compartment (leaves, roots and rhizosphere). From each

contrast, we used the absolute log2 Fold Change values (*ashr* shrinked (Stephens, 2017)) for each OTU

to quantify the impact of soil, plant and herbivore treatments on the microbiota in each

compartment. Comparisons of absolute log2 Fold Change values were performed by fitting a linear

mixed-effects model, specifying *compartment*, *treatment* (herbivory, plant or soil) and their

interaction as fixed factors and *OTU identity* as a random effect, and using the package *emmeans* to

infer contrasts (FDR corrected).

Aphid infestation. We tested whether the aphid infestation levels were influenced by soil

microbial diversity by fitting a cumulative link mixed model using the *ordinal* R package (Christensen,

2015), specifying *soil treatment*, *plant species*, and their interaction as fixed factors and *block* and

aphid clonal line as a random effect.

Results

Dataset summary and community composition

Overall, we identified 43,879 bacterial and 4,713 fungal OTUs. The analysis of the core

microbiota resulted in identifying 150 core bacterial OTUs and 26 fungal OTUs (Fig. 1). We identified

81 bacterial and 19 fungal OTUs as the core microbiota of rhizosphere soil, where the bacterial

community was largely dominated by uncultured taxa (50.6%), *Ramlibacter* (10.1%), *Pseudomonas*

(9.1%), *Massilia* (8.6%) and *Chitinophaga* (6.1%), while the fungal community was mostly represented

by *Peziza* (44.9%), unidentified fungi (18.1%), *Humicola* (9.1%), *Mortierella* (8.8%), *Penicillium* (8.6%),

Mucor (5.5%) and *Trichoderma* (4.9%). The core microbiota of roots was represented by 74 bacterial

and 16 fungal OTUs. Root tissues were mainly associated with *Flavobacterium* (39.4%) and uncultured

bacterial taxa (23.7%), while *Peziza* (22.9%), uncultured taxa (18.1%) and *Fusarium* (11.1%) dominated

the fungal community. Leaf core microbiota was represented by 20 bacterial and 6 fungal OTUs, with

a higher abundance of *Stenotrophomonas* (24.1%), *Candidatus* Hamiltonella (20.9%), *Flavobacterium*

(10.5%) and *Pedobacter* (9.4%), while fungi were mainly represented by *Cladosporium* (42.3%),

Penicillum (34.3%) and *Peziza* (13.6%). Aphids were mainly associated with *Buchnera* (64.1%) and

Candidatus *Hamiltonella* (35.8%). We identified 5 bacterial OTUs (*Flavobacterium* − 2 OTUs,

Citrobacter, *Pedobacter*, *Terrimonas*) shared between all compartments, and 4 fungal OTUs

(*Cladosporium*, *Humicola*, *Penicillium* and *Peziza*) shared between leaves, roots and rhizosphere. For a

more detailed analysis of the core microbiota, please see the Supplementary material.

To gain understanding on the overlap of core OTUs between compartments, we tested the

number of observed core OTUs overlapping between two compartments against a the number of

OTUs overlapping between two compartments due to random sampling. We found that the overlap of

bacterial core OTUs between pairs of plant compartments was always greater than random chance (P

< 0.05). Aphids also showed a higher number of shared bacterial core OTUs with leaves than random

chance (8 OTUs, χ^2 = 5.4; P = 0.01), but not with roots (7 OTUs, χ^2 = 2.7; P = 0.09) and rhizosphere soil

(5 OTUs, χ^2 = 2; P = 0.15). In fungi, we saw that the OTUs shared between leaves and roots (χ^2 = 1.3; P

= 0.25) or rhizosphere soil (χ^2 = 0.5; P = 0.47) were not different than those shared by chance, while

OTUs shared between roots and rhizosphere soil were greater than those shared by chance (χ^2 = 6.2;

P = 0.01).

Phylogenetic diversity

For bacterial communities, we found a significant compartment × soil treatment × plant

species interaction (Tab. 1). In all plant compartments (leaves, roots, rhizosphere) we found a higher

phylogenetic diversity in *S. vernei* than in *S. tuberosum* when plants were grown on low-diversity soil

treatment, and no differences between the two plant species were found when plants were grown on

high-diversity soil treatment (Supplementary material Tab. S3). Plant species did not influence aphid

bacterial diversity when they were exposed to plants grown on high- or low-diversity soil treatment

(Supplementary material Tab. S3). In fungal communities we found a significant effect of the factor

"plant species", reporting a higher diversity in *S. vernei* than *S. tuberosum* plants (P = 0.001, Tab. 1),

although we did not find any significant interaction with other factors.

We found a significant compartment × soil treatment × herbivory interaction in both bacterial

and fungal communities. Post-hoc contrasts show a higher leaf bacterial diversity in aphid-infested

plants compared with uninfested control plants when they were grown on low-diversity soil

treatment (Supplementary material, Tab. S4). Root bacterial and fungal communities, in both soil

treatments, had higher diversity values in infested plants compared with uninfested control plants

(Supplementary material, Tab. S4). In the rhizosphere we observed differences between infested and

uninfested plants in both bacterial and fungal community diversity of plants grown on high-diversity

soil treatment, while this difference was found just in the fungal community of plants grown on low-

diversity soil treatment (Supplementary material, Tab. S4).

We found phylogenetic diversity of the aphid microbiota was highest in the low diversity

treatment (Figure S3), which mirrored differences in aphid infestation levels (χ^2=8.19, df=1, *P*=0.004;

mean infestation scores: 3.10±0.23 for high diversity soil, 2.45±0.28 for low diversity soil).

Microbial community composition

The multivariate analysis (i.e., the PERMANOVA) reported a significant compartment × soil

treatment × plant species interaction (Tab. 1). Post-hoc contrasts showed differences between *S.*

vernei and *S. tuberosum* in the structure of leaf, root and rhizosphere bacterial and fungal

communities when plants were grown on low-diversity soil treatment (Fig. 2a-f, Tab. S5). In high-

diversity soil treatment, only root fungal communities differed between *S. vernei* and *S. tuberosum*

(Fig. 2e, Supplementary material Tab. S5). On the other hand, differences between low-diversity and

high-diversity treatments were found in both *S. vernei* and *S. tuberosum* in root and rhizosphere

communities (both bacterial and fungal, Fig. 2b-c and 2e-f, Supplementary material Tab. S6). In leaves,

differences between soil treatments were found just in the bacterial community of *S. vernei* (Fig. 2a,

Supplementary material Tab. S6). We also found a significant compartment × herbivory interaction

(Tab. 1), with herbivory influencing bacterial communities in all compartments, but fungal community

just in leaves and roots (Tab. 2). Our multivariate analysis demonstrated that the strongest driver of

rhizosphere, root and leaf microbial community structure was soil diversity treatment (Tab. 2) for

both bacterial and fungal communities. Indeed, bacterial and fungal communities responded to soil

diversity treatment and plant species across all compartments (roots, rhizosphere and leaves) (Table

2, Fig. 2a-f). Based on the variation explained by each factor included in the model, soil microbial

diversity was the most important factor shaping the microcosm's microbiota in all compartments

(Table 2). The variation explained by soil microbial diversity tended to decrease when moving across

compartments from rhizosphere to leaves and aphids (Table 2).

Soil diversity vs. plant species vs. herbivory driven effects. Which is strongest?

We answered this question in two ways, focusing on the single factors included in our design

(soil treatment, plant species, herbivory). First, as discussed above, using the variation explained by

each of our predictor variables in our PERMANOVA model, we determined that the predictor that

explained the most variation was the initial soil community diversity. Both soil diversity, plant species

and herbivory influenced bacterial and fungal assemblies in our system. Soil treatment explained

≈30% (rhizosphere), ≈20% (root), and ≈7% (leaf) of variation in microbiome community composition

(Table 2). However, the variance in community composition explained by both plant species and

herbivory (3–5%) was always lower than the variance explained by the soil treatment. Furthermore,

soil treatment explained ≈8% of variation in aphid microbiota (Table 2). This suggests that the soil-

driven effect is stronger than the other effects in our system.

Second, to investigate in more detail which factor (soil, plant, herbivore) had a stronger

influence on plant microbiome composition we examined the magnitude of change in abundance for

each OTU (absolute $\log_2$ Fold Changes) in relation to soil treatment, plant species and herbivory. For

both bacterial and fungal communities, and in all compartments, the changes produced by soil

treatment were greater than those produced by herbivory and plant species ($\chi^2_{bacteria}$=23331.3 and

χ^2_{fungi}=1055, df=2, P<0.001, Fig. 3), with the only exception being the leaf fungal community, where

no differences were found between the three factors (Fig. 3b). Also, in all cases, there was no

difference between the changes produced by herbivory and those produced by plant species (Fig. 3).

The analysis of changes in the abundance of OTUs in aphids revealed that soil diversity treatment had

a greater influence than plant species in shaping aphid bacterial communities (χ^2=766.8, df=1,

P<0.001; abs(log$_2$FoldChange)$_{soil}$ = 1.7±0.1 and abs(log$_2$FoldChange)$_{plant}$ = 0.37±0.1). Collectively, these

results demonstrate that the strongest effect on microbial taxa in the rhizosphere, roots, shoots, and

aphid herbivores is driven by the initial soil community diversity.

Discussion

Here we test the influence of multiple drivers (and their interaction) on plant microbiome

diversity and composition, and we show that, of all the drivers tested, soil microbial diversity had the

greatest influence on the microbial community composition of rhizosphere, roots, leaves, and even

aphid herbivores. Thus, we correctly hypothesized that soil microbial diversity drives changes in plant

and herbivore microbiota and that this effect would be much stronger than plant species or

herbivory. This influence of soil microbial diversity correlated with aphid abundance on infested

plants. Furthermore, we showed herbivory and plant species also affect the microbiome community

composition of leaves, roots and rhizosphere, but their effects are weaker than those driven by soil

diversity. We also observed that the response of plant microbiome to herbivory or plant species

differs according to soil treatment.

Plant and insect microbial diversity was influenced by soil microbiota and interactions between

soil microbiota and plant species and herbivory. When we quantified the relative contribution of the

main effects (soil, plant species, and herbivory), soil treatment was the strongest driver of plant

related microbiota composition. Soil community composition is well-known to influence plant

microbial composition (Bulgarelli *et al.*, 2012; Schlaeppi *et al.*, 2014; Bai *et al.*, 2015; Zarraonaindia *et*

al., 2015; de Souza *et al.*, 2016; Wagner *et al.*, 2016; Mezzasalma *et al.*, 2018; Grady *et al.*, 2019), but

here we reveal that soil produces a stronger effect when compared to other factors (i.e. plant species

and herbivory). Soil represents a major reservoir of microbial inoculum for plants (Zarraonaindia *et*

al., 2015; Trivedi *et al.*, 2020), especially belowground. However, we also observed that the effect

driven by soil is a function of plant species, likely because the intrinsic characteristics of each plant

species can modulate specific changes in the diversity and composition of microbial communities at

each compartment (Turner *et al.*, 2013; Dastogeer *et al.*, 2020).

Interestingly, we found that soil microbial diversity influences the phylogenetic diversity and

structure of aphid microbiota. The influence of the soil microbial diversity on aphid bacterial

communities and aphid infestation level could potentially be explained through two, non-mutually

exclusive, mechanisms: (i) translocation of microbes from the rhizosphere through or on the plant and

(ii) changes in plant physiology and/or metabolome. Although leaves are physically separated from

roots, their microbiomes can still interact at interfaces such as the stem, and microbial translocation

could occur due to active and passive mechanisms (Bai *et al.*, 2015; Wagner *et al.*, 2016). A recent

study comparing the microbiota of caterpillars feeding on detached leaves and intact plants found the

microbiota of caterpillars that fed on intact plants had a similar community composition to the soil

microbiota (Hannula *et al.*, 2019) suggesting direct (splashing of soil microbiota on leaves) or indirect

(movement through the plant) microbial translocation. However, our data does not show this pattern,

as the core microbiome belowground is different from shoots and herbivores, few OTUs are common

to all compartments in the system, and the overlap of core OTUs between aphids and belowground

compartments is not greater than random chance. The aphids in our system employ a different

feeding strategy (sap-feeding) compared to the caterpillars (chewing) in the previous study (Hannula

et al., 2019), and chewing herbivores may have an increased likelihood of environmental uptake of

microbes (Paniagua Voirol *et al.*, 2018). We thus find it unlikely that microbes in our system were

translocated through or on the plant to the herbivore.

The second potential mechanism is through changes in plant physiology. Many soil microbes

are able to modulate plant nutrient intake, or prime plant defences (Martinez-Medina *et al.*, 2016).

The composition of belowground microbial communities can alter plant metabolism (Li *et al.*, 2019),

which in turn influences herbivore fitness (Mason *et al.*, 2019). Our low diversity treatment lacked

large microbes including arbuscular mycorrhizal fungi, a group well known to prime plant defences

(Jung *et al.*, 2012; Bennett *et al.*, 2018), although aphids are less susceptible to changes in defenses

primed by arbuscular mycorrhizal fungi (Koricheva *et al.*, 2009). Thus, the changes we observed in the

aphid microbiome could be due to changes in host plant physiology and metabolome, for example

triggering of plant defences which has been shown to decrease the diversity of plant-associated

microbial communities (Kniskern *et al.*, 2007). The higher aphid abundance on plants grown in the

high diversity soil provides indirect evidence for such changes in plant biochemical composition, as

does a previous study showing increased aphid suitability as a host for parasitoids when feeding on

Solanum plants grown with AM fungi from the same site (Bennett *et al.*, 2016). While no previous

study found a relationship between soil microbial diversity and aphid infestation, previous studies

show that soil microbiota can influence insect feeding behaviour through changes in the plant

metabolome (Badri *et al.*, 2013), which can represent the likely mechanism driving our observations.

While a clear consumer-driven effect was observed on the plant microbiome in our study, it

was a weaker effect than soil microbial diversity. Thus, herbivory plays a less significant role in

determining plant microbiome composition. The herbivory-driven effect on the bacterial community

composition of the roots and rhizosphere could be driven by changes in plant physiology (e.g. defence

activation, carbon metabolism) and root exudation due to aphid feeding activity (Züst and Agrawal,

2016; Hoysted *et al.*, 2018). Herbivory has been shown to alter the types of organic compounds

released at the root surface leading to changes in the composition of rhizosphere microbial

communities (Lareen *et al.*, 2016; Hu *et al.*, 2018). Previous research has shown that *Bemisia tabaci* (whitefly) herbivory can alter the rhizosphere microbiome of pepper plants (Kong *et al.*, 2016), and artificial induction of plant defences (Hein *et al.*, 2008), or their deactivation (Kniskern *et al.*, 2007), has been shown to shape rhizosphere microbial communities. In our study we focused on a sap-feeding insect, which is known to trigger specific physiological responses in plants. Thus, further research is needed to test if different herbivores with different feeding strategies are able to drive changes in the microbiota thriving in different plant compartments.

Plant species was also a predictor shaping the microbiome community composition of the rhizosphere, plants, and herbivores in our study. Plant species is known to be a strong driver of root and rhizosphere microbiota because plant species differ in morphology, chemistry and relationship with microorganisms (Trivedi *et al.*, 2020). *Solanum* plants are known to produce toxic glycoalkaloids as chemical defense against herbivory (Altesor *et al.*, 2014), and these compounds could also shape the microbiota associated with the plant. The strength of this effect might be context-dependent due to the interaction with the soil microbial composition (Bulgarelli *et al.*, 2013; Fierer, 2017). This might explain why we observed a greater impact of soil treatment than plant species in the belowground microbiotas in our study. Also, it might explain why we found differences in the bacterial community between plant species just in one soil treatment. It has been previously shown that soil represents a reservoir for leaf microbial communities and that phyllosphere habitat selects for specific members (Grady *et al.*, 2019), which partially explains our observation that the impact of soil microbial diversity was greater than plant species on the phyllosphere bacterial and fungal communities. The differences in the microbiome composition of aphids feeding and reproducing on the two different plant species is not surprising, as it is well known that the identity of the host plant is a major factor in shaping insect-associated microbial community composition (Colman *et al.*, 2012; Malacrinò, 2018). The fact

that seeds were not surface-sterilized might be a caveat of our study, potentially influencing the

structure of plant microbiome differently for each species. However, previous studies show that even

surface-sterilized seeds show differences in their endophytic microbiome according to the plant

genotype (Walitang *et al.*, 2018; Raj *et al.*, 2019; Liu *et al.*, 2020), and our results show a major effect

driven by soil microbial community composition, thus we are confident that our results were not

biased by not surface-sterilizing seeds.

By quantifying and comparing the relative strength of multiple biotic drivers of microbial

community composition, our work contributes to a more comprehensive understanding of the factors

determining the outcome of plant–microbe–insect interactions, and how plant-associated

microbiomes assemble and respond to resource- and consumer-driven effects. Thus, if understood

and managed correctly, these interactions have potential to be applied in natural and managed

systems to improve food security and safety, or the success of ecological restoration efforts.

Declarations

Availability of data and material

Raw data is available at the NCBI SRA database under accession number PRJNA557499.

Competing interests

We declare that none of the authors have competing financial or non-financial interests.

Authors' contributions

All authors designed the experiment. AM and AEB performed the greenhouse experiment. AM prepared library and performed data analysis. AM wrote the first draft of the manuscript and all authors contributed substantially to revisions.

Acknowledgements

We would also like to thank Davide Bulgarelli (University of Dundee, UK), Marta Jarzyna (Ohio State University, USA), Jeff Powell (Hawksbury Institute for the Environment, Western Sydney University, AU), and Philip Smith (Dundee, UK) for helpful comments on the manuscript.

Funding

We would like to acknowledge funding from EU COST Action FA1405 and the strategic research programme and the underpinning capacity programme funded by the Scottish Government's Rural and Environment Science and Analytical Services Division.

References

Altesor, P., García, Á., Font, E., Rodríguez-Haralambides, A., Vilaró, F., Oesterheld, M., et al. (2014) Glycoalkaloids of Wild and Cultivated Solanum: Effects on Specialist and Generalist Insect Herbivores. *J Chem Ecol* **40**: 599–608.

Andersen, K., Kirkegaard, R.H., Karst, S.M., and Albertsen, M. (2018) ampvis2: an R package to analyse and visualise 16S rRNA amplicon data. *bioRxiv* 299537.

Anderson, M.J. and Willis, T.J. (2003) Canonical analysis of principal coordinates: a useful method of constrained ordination for ecology. *Ecology* **84**: 511–525.

Badri, D. V., Zolla, G., Bakker, M.G., Manter, D.K., and Vivanco, J.M. (2013) Potential impact of soil microbiomes on the leaf metabolome and on herbivore feeding behavior. *New Phytol* **198**: 264–273.

Bai, Y., Müller, D.B., Srinivas, G., Garrido-Oter, R., Potthoff, E., Rott, M., et al. (2015) Functional overlap of the *Arabidopsis* leaf and root microbiota. *Nature* **528**: 364–369.

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015) Fitting linear mixed-effects models using lme4. *J Stat Softw* **67**: 1–48.

Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z., Godhe, A., et al. (2013) Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. *Methods Ecol Evol* **4**: 914–919.

Bennett, A.E., Millar, N.S., Gedrovics, E., and Karley, A.J. (2016) Plant and insect microbial symbionts alter the outcome of plant–herbivore–parasitoid interactions: implications for invaded, agricultural and natural systems. *J Ecol* **104**: 1734–1744.

Bennett, A.E., Orrell, P., Malacrino, A., Pozo, M.J., Bennett, A.E., Orrell, P., et al. (2018) Fungal-

Mediated Above-Belowground Interactions: The Community Approach, Stability, Evolution,

Mechanisms, and Applications. In *Aboveground–Belowground Community Ecology*. Ohgushi, T.,

Johnson, S.N., and Wurst, S. (eds). Springer, pp. 85–116.

Bennett, A.E., Thomsen, M., and Strauss, S.Y. (2011) Multiple mechanisms enable invasive species to

suppress native species. *Am J Bot* **98**: 1086–1094.

Berg, G., Grube, M., Schloter, M., and Smalla, K. (2014) Unraveling the plant microbiome: looking back

and future perspectives. *Front Microbiol* **5**: 148.

Bulgarelli, D., Rott, M., Schlaeppi, K., Ver Loren van Themaat, E., Ahmadinejad, N., Assenza, F., et al.

(2012) Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial

microbiota. *Nature* **488**: 91–95.

Bulgarelli, D., Schlaeppi, K., Spaepen, S., van Themaat, E.V.L., and Schulze-Lefert, P. (2013) Structure

and Functions of the Bacterial Microbiota of Plants. *Annu Rev Plant Biol* **64**: 807–838.

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., et al. (2012) Ultra-

high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME*

J **6**: 1621–1624.

Christensen, R.H.B. (2015) ordinal—regression models for ordinal data. R package.

http://www2.uaem.mx/r-mirror/web/packages/ordinal/.

Christian, N., Sullivan, C., Visser, N.D., and Clay, K. (2016) Plant Host and Geographic Location Drive

Endophyte Community Composition in the Face of Perturbation. *Microb Ecol* **72**: 621–632.

Coleman-Derr, D., Desgarennes, D., Fonseca-Garcia, C., Gross, S., Clingenpeel, S., Woyke, T., et al.

(2016) Plant compartment and biogeography affect microbiome composition in cultivated and

native *Agave* species. *New Phytol* **209**: 798–811.

Colman, D.R., Toolson, E.C., and Takacs-Vesbach, C.D. (2012) Do diet and taxonomy influence insect

gut bacterial communities? *Mol Ecol* **21**: 5124–5137.

Cordovez, V., Dini-Andreote, F., Carrión, V.J., and Raaijmakers, J.M. (2019) Ecology and Evolution of

Plant Microbiomes. *Annu Rev Microbiol* **73**: 69–88.

Dastogeer, K.M.G., Tumpa, F.H., Sultana, A., Akter, M.A., and Chakraborty, A. (2020) Plant

microbiome–an account of the factors that shape community composition and diversity. *Curr*

Plant Biol **23**: 100161.

Dixon, P. (2003) VEGAN, a package of R functions for community ecology. *J Veg Sci* **14**: 927–930.

Faith, D.P. (1992) Conservation evaluation and phylogenetic diversity. *Biol Conserv* **61**: 1–10.

Fierer, N. (2017) Embracing the unknown: disentangling the complexities of the soil microbiome. *Nat*

Rev Microbiol **15**: 579–590.

Fitzpatrick, C.R., Copeland, J., Wang, P.W., Guttman, D.S., Kotanen, P.M., and Johnson, M.T.J. (2018)

Assembly and ecological function of the root microbiome across angiosperm plant species. *Proc*

Natl Acad Sci **115**: E1157–E1165.

Grady, K.L., Sorensen, J.W., Stopnisek, N., Guittar, J., and Shade, A. (2019) Assembly and seasonality of

core phyllosphere microbiota on perennial biofuel crops. *Nat Commun* **10**: 4135.

Hannula, S.E., Zhu, F., Heinen, R., and Bezemer, T.M. (2019) Foliar-feeding insects acquire

microbiomes from the soil rather than the host plant. *Nat Commun* **10**: 1254.

Hein, J.W., Wolfe, G. V., and Blee, K.A. (2008) Comparison of rhizosphere bacterial communities in

Arabidopsis thaliana mutants for systemic acquired resistance. *Microb Ecol* **55**: 333–343.

Hoysted, G.A., Bell, C.A., Lilley, C.J., and Urwin, P.E. (2018) Aphid Colonization Affects Potato Root

Exudate Composition and the Hatching of a Soil Borne Pathogen. *Front Plant Sci* **9**:.

Hu, L., Robert, C.A.M., Cadot, S., Zhang, X., Ye, M., Li, B., et al. (2018) Root exudate metabolites drive

plant–soil feedbacks on growth and defense by shaping the rhizosphere microbiota. *Nat*

Commun **9**: 2738.

Jung, S.C., Martinez-Medina, A., Lopez-Raez, J.A., and Pozo, M.J. (2012) Mycorrhiza-Induced

Resistance and Priming of Plant Defenses. *J Chem Ecol* **38**: 651–664.

Karley, A.J., Emslie-Smith, M., and Bennett, A.E. (2017) Potato aphid *Macrosiphum euphorbiae*

performance is determined by aphid genotype and not mycorrhizal fungi or water availability.

Insect Sci **24**: 1015–1024.

Kembel, S.W., Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly, D.D., et al. (2010)

Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* **26**: 1463–1464.

Kniskern, J.M., Traw, M.B., and Bergelson, J. (2007) Salicylic acid and jasmonic acid signaling defense

pathways reduce natural bacterial diversity on *Arabidopsis thaliana*. *Mol Plant-Microbe Interact*

20: 1512–1522.

Kong, H.G., Kim, B.K., Song, G.C., Lee, S., and Ryu, C.-M. (2016) Aboveground whitefly infestation-

mediated reshaping of the root microbiota. *Front Microbiol* **7**: 1314.

Koricheva, J., Gange, A.C., and Jones, T. (2009) Effects of mycorrhizal fungi on insect herbivores: a

meta-analysis. *Ecology* **90**: 2088–2097.

Lareen, A., Burton, F., and Schäfer, P. (2016) Plant root-microbe communication in shaping root

microbiomes. *Plant Mol Biol* **90**: 575–587.

Lee, B., Lee, S., and Ryu, C.-M. (2012) Foliar aphid feeding recruits rhizosphere bacteria and primes

plant immunity against pathogenic and non-pathogenic bacteria in pepper. *Ann Bot* **110**: 281–

290.

Lee, J., Lee, S., and Young, J.P.W. (2008) Improved PCR primers for the detection and identification of

arbuscular mycorrhizal fungi. *FEMS Microbiol Ecol* **65**: 339–349.

Levy, A., Salas Gonzalez, I., Mittelviefhaus, M., Clingenpeel, S., Herrera Paredes, S., Miao, J., et al.

(2018) Genomic features of bacterial adaptation to plants. *Nat Genet* **50**: 138–150.

Li, X., Jousset, A., de Boer, W., Carrión, V.J., Zhang, T., Wang, X., and Kuramae, E.E. (2019) Legacy of

land use history determines reprogramming of plant physiology by soil microbiome. *ISME J* **13**:

738–751.

Lin, M., Xiong, H., Xiang, X., Zhou, Z., Liang, L., and Mei, Z. (2020) The Effect of Plant Geographical

Location and Developmental Stage on Root-Associated Microbiomes of *Gymnadenia conopsea*.

Front Microbiol **11**:.

Liu, Y., Yan, H., Zhang, X., Zhang, R., Li, M., Xu, T., et al. (2020) Investigating the endophytic bacterial

diversity and community structures in seeds of genetically related maize (*Zea mays* L.)

genotypes. *3 Biotech* **10**: 27.

Love, M.I., Huber, W., and Anders, S. (2014) Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol* **15**: 550.

Malacrinò, A. (2018) Meta-Omics Tools in the World of Insect-Microorganism Interactions. *Biology*

(Basel) **7**: 50.

Malacrinò, A., Campolo, O., Medina, R.F., and Palmeri, V. (2018) Instar- and host-associated

differentiation of bacterial communities in the Mediterranean fruit fly *Ceratitis capitata*. *PLoS*

One **13**: e0194131.

Martinez-Medina, A., Flors, V., Heil, M., Mauch-Mani, B., Pieterse, C.M.., Pozo, M.J., et al. (2016)

Recognizing Plant Defense Priming. *Trends Plant Sci* **21**: 818–822.

Mason, C.J., Ray, S., Shikano, I., Peiffer, M., Jones, A.G., Luthe, D.S., et al. (2019) Plant defenses

interact with insect enteric bacteria by initiating a leaky gut syndrome. *Proc Natl Acad Sci* **116**:

15991–15996.

McMurdie, P.J. and Holmes, S. (2013) phyloseq: An R package for reproducible interactive analysis

and graphics of microbiome census data. *PLoS One* **8**: e61217.

Mezzasalma, V., Sandionigi, A., Guzzetti, L., Galimberti, A., Grando, M.S., Tardaguila, J., and Labra, M.

(2018) Geographical and Cultivar Features Differentiate Grape Microbiota in Northern Italy and

Spain Vineyards. *Front Microbiol* **9**:.

Nilsson, R.H., Larsson, K.-H., Taylor, A.F.S., Bengtsson-Palme, J., Jeppesen, T.S., Schigel, D., et al.

(2019) The UNITE database for molecular identification of fungi: handling dark taxa and parallel

taxonomic classifications. *Nucleic Acids Res* **47**: D259–D264.

O'Brien, F.J.M., Dumont, M.G., Webb, J.S., and Poppy, G.M. (2018) Rhizosphere Bacterial

Communities Differ According to Fertilizer Regimes and Cabbage (*Brassica oleracea* var. *capitata*

L.) Harvest Time, but Not Aphid Herbivory. *Front Microbiol* **9**:.

Ofek, M., Voronov-Goldman, M., Hadar, Y., and Minz, D. (2014) Host signature effect on plant root-

associated microbiomes revealed through analyses of resident vs . active communities. *Environ*

Microbiol **16**: 2157–2167.

Paniagua Voirol, L.R., Frago, E., Kaltenpoth, M., Hilker, M., and Fatouros, N.E. (2018) Bacterial

Symbionts in Lepidoptera: Their Diversity, Transmission, and Impact on the Host. *Front Microbiol*

9: 556.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2012) The SILVA ribosomal

RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res*

41: D590–D596.

R Core Team (2013) R: A language and environment for statistical computing, Vienna, Austria: R

Foundation for Statistical Computing.

Raj, G., Shadab, M., Deka, S., Das, M., Baruah, J., Bharali, R., and Talukdar, N.C. (2019) Seed interior

microbiome of rice genotypes indigenous to three agroecosystems of Indo-Burma biodiversity

hotspot. *BMC Genomics* **20**: 924.

Rognes, T., Flouri, T., Nichols, B., Quince, C., and Mahé, F. (2016) VSEARCH: a versatile open source

tool for metagenomics. *PeerJ* **4**: e2584.

Rolfe, S.A., Griffiths, J., and Ton, J. (2019) Crying out for help with root exudates: adaptive

mechanisms by which stressed plants assemble health-promoting soil microbiomes. *Curr Opin*

Microbiol **49**: 73–82.

Rosenberg, E. and Zilber-Rosenberg, I. (2016) Microbes Drive Evolution of Animals and Plants: the

Hologenome Concept. *MBio* **7**:.

Saikkonen, K., Nissinen, R., and Helander, M. (2020) Toward Comprehensive Plant Microbiome

Research. *Front Ecol Evol* **8**:.

Schlaeppi, K. and Bulgarelli, D. (2015) The Plant Microbiome at Work. *Mol Plant-Microbe Interact* **28**:

212–217.

Schlaeppi, K., Dombrowski, N., Oter, R.G., Ver Loren van Themaat, E., and Schulze-Lefert, P. (2014)

Quantitative divergence of the bacterial root microbiota in *Arabidopsis thaliana* relatives. *Proc*

Natl Acad Sci **111**: 585–592.

Schmidt, P.-A., Bálint, M., Greshake, B., Bandow, C., Römbke, J., and Schmitt, I. (2013) Illumina

metabarcoding of a soil fungal community. *Soil Biol Biochem* **65**: 128–132.

de Souza, R.S.C., Okura, V.K., Armanhi, J.S.L., Jorrín, B., Lozano, N., da Silva, M.J., et al. (2016)

Unlocking the bacterial and fungal communities assemblages of sugarcane microbiome. *Sci Rep*

6: 28774.

Stephens, M. (2017) False discovery rates: a new deal. *Biostatistics* **18**: 275–294.

Strano, C.P., Malacrinò, A., Campolo, O., and Palmeri, V. (2018) Influence of Host Plant on

Thaumetopoea pityocampa Gut Bacterial Community. *Microb Ecol* **75**: 487–494.

Toju, H., Tanabe, A.S., Yamamoto, S., and Sato, H. (2012) High-Coverage ITS Primers for the DNA-

Based Identification of Ascomycetes and Basidiomycetes in Environmental Samples. *PLoS One* **7**:

e40863.

Trivedi, P., Leach, J.E., Tringe, S.G., Sa, T., and Singh, B.K. (2020) Plant–microbiome interactions: from

community assembly to plant health. *Nat Rev Microbiol* **18**: 607–621.

Turner, T.R., James, E.K., and Poole, P.S. (2013) The plant microbiome. *Genome Biol* **14**: 209.

Wagner, M.R., Lundberg, D.S., del Rio, T.G., Tringe, S.G., Dangl, J.L., and Mitchell-Olds, T. (2016) Host

genotype and age shape the leaf and root microbiomes of a wild perennial plant. *Nat Commun* **7**:

12151.

Walitang, D.I., Kim, C.-G., Kim, K., Kang, Y., Kim, Y.K., and Sa, T. (2018) The influence of host genotype

and salt stress on the seed endophytic community of salt-sensitive and salt-tolerant rice

cultivars. *BMC Plant Biol* **18**: 51.

Yang, J.W., Yi, H.-S., Kim, H., Lee, B., Lee, S., Ghim, S.-Y., and Ryu, C.-M. (2011) Whitefly infestation of

pepper plants elicits defence responses against bacterial pathogens in leaves and roots and

changes the below-ground microflora. *J Ecol* **99**: 46–56.

Zarraonaindia, I., Owens, S.M., Weisenhorn, P., West, K., Hampton-Marcell, J., Lax, S., et al. (2015) The

Soil Microbiome Influences Grapevine-Associated Microbiota. *MBio* **6**:.

Zhang, J., Kobert, K., Flouri, T., and Stamatakis, A. (2014) PEAR: a fast and accurate Illumina Paired-

End reAd mergeR. *Bioinformatics* **30**: 614–620.

Züst, T. and Agrawal, A.A. (2016) Mechanisms and evolution of plant resistance to aphids. *Nat Plants*

2: 15206.

Tables

Table 1. Models testing the effect of compartment (aphids, leaves, roots, rhizosphere soil), soil treatment (high diversity, low diversity), plant species (*S. tuberosum*, *S. vernei*), herbivory (infested, control) and their interaction on the phylogenetic diversity (linear mixed-effect model) and taxonomical structure (PERMANOVA) of bacterial and fungal communities.

		Bacterial community					Fungal community			
		Phylogenetic diversity		PERMANOVA			Phylogenetic diversity		PERMANOVA	
Factors	df	χ^2	P	F	P	df	χ^2	P	F	P
Compartment (Cp)	3	843.62	**<0.001**	23.39	**0.001**	2	895.91	**<0.001**	12.9894	**0.001**
Soil treatment (S)	1	2.19	0.13	13.26	**0.001**	1	225.62	**<0.001**	14.906	**0.001**
Plant species (P)	1	18.11	**<0.001**	2.37	**0.001**	1	10.67	**<0.01**	1.9625	**0.003**
Herbivory (H)	1	34.66	**<0.001**	2.52	**0.001**	1	47.21	**<0.001**	3.7912	**0.001**
Cp × S	3	37.23	**<0.001**	5.67	**0.001**	2	252.44	**<0.001**	5.7238	**0.001**
Cp × P	3	8.18	**0.04**	1.51	**0.005**	2	1.66	0.43	1.3138	**0.03**
S × P	1	23.6	**<0.001**	1.69	**0.025**	1	3.74	0.05	1.7088	**0.014**
Cp × H	2	16.51	**<0.001**	1.61	**0.007**	2	20.7	**<0.001**	2.0339	**0.001**
S × H	1	0.01	0.9	1.25	0.11	1	5.61	**0.01**	1.2592	0.107
P × H	1	2.26	0.13	1.17	0.18	1	0.16	0.68	1.1442	0.217
Cp × S × P	3	12.57	**<0.01**	1.42	**0.005**	2	0.28	0.86	1.34	**0.029**
Cp × S × H	2	6.86	**0.03**	1.28	0.07	2	12.38	**<0.01**	1.107	0.211
Cp × P × H	2	1.9	0.38	1.04	0.31	2	1.18	0.55	1.0136	0.402
S × P × H	1	2.59	0.1	1.13	0.20	1	1.33	0.24	1.1739	0.152
Cp × S × P × H	2	0.25	0.87	1.05	0.29	2	0.84	0.65	1.1249	0.186

683 **Table 2**. Analysis of the effects of soil treatment (high diversity, low diversity), plant species (*S.*

684 *tuberosum*, *S. vernei*), herbivory (infested, control) on the bacterial and fungal community

685 taxonomical structure for each compartment (aphids, leaves, roots and rhizosphere) performed

686 through PERMANOVA (Unifrac distance matrix). Analyses were run separately for each compartment

687 within each community. Values in bold represent $P<0.05$.

	Aphids		Leaves		Roots		Rhizosphere	
Bacterial community	R^2	*P*	R^2	*P*	R^2	*P*	R^2	*P*
Soil treatment	0.08	**<0.01**	0.07	**<0.01**	0.16	**<0.01**	0.28	**<0.01**
Plant species	0.03	**0.01**	0.03	**<0.01**	0.03	**<0.01**	0.02	**0.04**
Herbivory	-	-	0.03	**<0.01**	0.04	**<0.01**	0.02	**0.04**
	Aphids		Leaves		Roots		Rhizosphere	
Fungal community	R^2	*P*	R^2	*P*	R^2	*P*	R^2	*P*
Soil treatment	-	-	0.06	**<0.01**	0.21	**<0.01**	0.31	**<0.01**
Plant species	-	-	0.02	0.26	0.03	**<0.01**	0.02	0.17
Herbivory	-	-	0.05	**<0.01**	0.03	**0.01**	0.02	0.07

Figure captions

Figure 1. Core bacterial (A) and fungal (B) OTUs shared between compartments. Core OTUs were identified among those with relative abundance of >0.1% in more than 50% of samples.

Figure 2. Canonical Analysis of Principal coordinates (CAP) analysis of bacterial and fungal

communities Unifrac distance matrix for each compartment. We report the response of bacterial (*a-c*)

and fungal (*d-f*) communities to soil microbial diversity, plant species and herbivory in leaves (*a, d*),

roots (*b, e*) and rhizosphere soil (*c, f*). Aphid bacterial community (*g*) responded to both plant species

and soil microbial diversity. For each graph, percentages in parentheses inside each graph along the

axes report the variance explained by the respective axis.

Figure 3. Magnitude of changes in abundance for each OTU (absolute $\log_2$ Fold Changes). For each

compartment (leaves, roots and rhizosphere) we investigated the response of single OTUs to soil

microbial diversity (red), plant species (green) and herbivory (blue), for (*a*) bacterial and (*b*) fungal

communities. Comparisons were tested using a linear mixed-effect model, and contrasts were

extracted using the function *emmeans*. *** *P*<0.001.

Supplementary material

Supplementary material. Supplementary results, tables and figures.