Cryo-electron tomography of the herpesvirus procapsid reveals

interactions of the portal with the scaffold and a shift on maturation

Michael H. C. Buch[a,$], William W. Newcomb[a,$], Dennis C. Winkler[a,b], Alasdair C. Steven[a,#], J.

Bernard Heymann[a,#]

[a] Laboratory of Structural Biology Research, National Institute of Arthritis and Musculoskeletal

and Skin Diseases, National Institutes of Health, Bethesda, MD, US.

[b] Advanced Imaging Core, National Institute on Deafness and Other Communication Disorders,

National Institutes of Health, Bethesda, MD, US.

[$] These authors contributed equally to this study.

[#] Corresponding authors. Contact: heymannb@mail.nih.gov , stevena@mail.nih.gov

Running title: Visualizing the herpes portal in the procapsid

Abstract

Herpes simplex virus type 1 (HSV-1) requires seven proteins to package its genome through a

vertex in its capsid, one of which is the portal protein, pUL6. The portal protein is also thought to

facilitate assembly of the procapsid. While the portal has been visualized in mature capsids, we

aimed to elucidate its role in the assembly and maturation of procapsids using cryo-electron

tomography. We identified the portal vertex in individual procapsids, calculated a subtomogram

average, and compared that with the portal vertex in empty mature capsids (A-capsids). The

resulting maps show the portal on the interior surface with its narrower end facing outwards,

while maintaining close contact with the capsid shell. In the procapsid, the portal is embedded in

the underlying scaffold, suggesting that assembly involves a portal – scaffold complex. During

maturation, the capsid shell angularizes with a corresponding outward movement of the vertices.

We found that in A-capsids, the portal translocates further than the adjacent capsomers and

strengthens its contacts with the capsid shell. Our methodology also allowed us to determine the

number of portal vertices in each capsid, with most having one per capsid, but some none or two,

and rarely three. The predominance of a single portal per capsid supports facilitation of the

assembly of the procapsid.

Importance

Herpes simplex virus type 1 (HSV-1) infects a majority of humans, causing mostly mild disease

but in some cases progressing towards life-threatening encephalitis. Understanding the life cycle

of the virus is important to devise countermeasures. Production of the virion starts with the

assembly of an icosahedral procapsid including DNA packaging proteins at a vertex, one of which

is the dodecameric portal protein. The procapsid then undergoes maturation and DNA packaging

through the portal, driven by a terminase complex. We used cryo-electron tomography to

visualize the portal in procapsids and compare them to mature empty capsids. We found the

portal located inside one vertex interacting with the scaffold protein in the procapsid. On

maturation, the scaffold dissociates, the capsid angularizes, and the portal moves outward,

associating closely with the capsid shell. These transformations may provide a basis for the

development of drugs to prevent HSV-1 infections.

Introduction

Members of the *Herpesviridae* family cause persistent infections in humans, ranging from cold

sores on the lips and genitals to shingles and tumors. While individual members of this family

exhibit differences in pathogenicity and structural details, reflected in their classification into

three subfamilies, their overall architecture is largely conserved. The large genome of HSV-1, the

prototypical representative of *Herpesviridae*, is accommodated in a correspondingly massive and

complex capsid. During their elaborate life cycle (1), nascent capsids assemble in the nucleus as

spherical precursor particles called procapsids (2-4), which conform to T=16 quasi-icosahedral

symmetry and contain a spherical internal scaffold. The scaffold shell consists of the viral proteins

pUL26.5 and pUL26, of which the latter has an N-terminally module, the viral protease (4). Upon

completion of the procapsid, the protease is activated and proceeds to cleave the scaffold,

initiating maturation of the capsid. Maturation involves: (I) angularization of the capsid; (II)

reorganization of the interactions that engage the triplex proteins as well as rearrangement of

capsomers (1-3) from skew to more symmetric orientations; (III) conformational changes (2, 5-8)

that establish interactions between capsomer subunits; and (IV) decoration of the mature capsid

with additional proteins (9, 10). Maturation increases the capsid's stability dramatically,

countering the internal pressure exerted by the tightly packed genome (11). This process yields

three capsid species, two of which remain in the nucleus. The A-capsid has a mature shell that

does not contain any DNA and is essentially empty, while B-capsids retain a shrunken and

somewhat disordered scaffold despite having matured into an angular shape. Only mature, DNA-

filled capsids, called C-capsids, leave the nucleus and do so via the perinuclear space (reviewed

in (12, 13)), to undergo the subsequent steps in the viral life cycle.

DNA packaging occurs at a capsid vertex and requires seven proteins, pUL6, pUL15, pUL17,

pUL25, pUL28, pUL32 and pUL33 (14). pUL6 forms a circular dodecamer similar to the portals of

bacteriophages (15) and may facilitate procapsid assembly (16). Heterotrimers of pUL17 (one

copy) and pUL25 (two copies) are bound to the pentons and peripentonal triplexes (17-19). A

terminase complex composed of pUL15, pUL28 and pUL33 provides the force for packaging and

subsequent cleavage of the DNA (20-26). The final protein, pUL32, has not been found as a

component of capsids (14). These proteins replace a penton presumably at one vertex of the

capsid (27). While the structural disposition and transformations of some of these components

have been elucidated, many details of assembly, maturation and packaging remain to be

described.

Early attempts to locate the portal-bearing vertex in capsids employed cryo-electron tomography

(cryoET) (28, 29). The tomograms were of low resolution (55-60 Å), complicating analysis. This

led to placing the portal either at the level of capsomers (28) or inside the capsid shell (29).

Subsequent analysis employing cryo-electron microscopy (cryoEM) and single particle analysis

(SPA), indicated that the portal is located inside the capsid shell (30, 31). One of the difficulties in

the SPA approach is that the overlap of one vertex with the large mass of the capsid (~100 fold

ratio) makes it hard to distinguish a portal vertex from the penton vertices. Nevertheless, the

extensive averaging of structural information means that correct assignments would dominate

incorrect assignments distributed over the eleven other possibilities and reveal the location of

the portal. The recent technological advances in cryoEM enabled the high resolution

reconstruction of various herpesvirus capsids showing the portal on the inside of one vertex (18,

31-34).

One omission from the existing studies is the predisposition of the portal in the initial assembly

product, the procapsid. It is also not clear that only one vertex bears a portal as assembly can

occur without portal (2). In this study, we returned to cryoET to examine the portal vertices in

procapsids and compared them to A-capsids as a control. In this effort, we developed the

computational capability to find the portal vertices in a more reliable manner, allowing an

estimate of the number of portal vertices in each capsid. From subtomogram averages of the

portal vertices, we found abundant contacts between the portal and the scaffolding layer in the

procapsid. We also found that the portal moves closer to the capsid shell on maturation.

Results

Production of procapsids. Procapsids were produced using the M100 mutant that lacks the viral

protease VP24 required for rapid maturation (35). We followed a gentle purification protocol (5,

7) to minimize damage to the particles. For successful identification of the portal vertex in

procapsids, several precautions were taken: since procapsids are fragile and malleable, we used

an antibody binding to the major capsid protein (2) that stabilized the particles. We performed

cryoET on the isolated procapsids, collecting and analyzing 80 tilt series where the particles were

well separated (Fig. 1A).

Production of capsid samples using a pUL25-deficient mutant. We decided to analyze mature

A-capsids as a control to avoid interference from internal density features such as scaffolding (cf.

B-capsids) or packaged DNA (cf. C-capsids). The established protocol for purifying HSV-1

nucleocapsids from a wild-type infection yields only a minor fraction of A-capsids. To increase

their yield, we used a mutant lacking the vertex protein, pUL25, that is defective in DNA packaging

(36). This virus replicates in a complementary African Green Monkey cell line transfected with

the *UL25* gene, but in wild-type cells cannot produce progeny. In our protocol, centrifugation on

small-scale sucrose gradients yielded bands for A-capsids and B-capsids but no band for the DNA-

filled C-capsids (Fig. 1C). We isolated the UL25-Null A-capsid band (hereafter referred to as A-

capsids) and collected tilt series for cryoET. Individual capsids in the tomograms were well spread

out in the sample, and almost all capsids could be confirmed as A-capsids from their empty

appearance (Fig. 1B).

Fig. 1. Isolated capsids. A. Central slice of a tomogram of procapsids. Some particles contained visibly degraded scaffold or showed imperfections (red arrowheads) and were excluded from further processing. We took care to include only particles that were separated, spherical, and had a visibly intact scaffold (turquoise arrowheads). **B.** Central slice of a tomogram of A-capsids. The individual particles are well defined and separated. Occasionally, a capsid with some internal content – probably some scaffold – were found in the field of view (yellow arrowhead). These particles were not included in further processing. **C.** Small-scale sucrose gradient of a capsid mix obtained during initial purification from a UL25-Null mutant, showing bands of A- and B-capsids. Scale bars, 500 Å.

Identification of the portal vertex in cryo-electron tomograms. Our aim was to locate the portal using cryoET to take advantage of the straightforward identification of unique features in 3D maps. The approach used in an earlier tomographic protocol relied on simple template matching to find the portal vertex (28). Another protocol made use of the comparison of opposite vertices and assessing the density differences between vertices (30). In an effort to improve the detection of portal vertices, we developed a different algorithm aimed at detecting the vertex that is most different from the other vertices. We implemented this algorithm in a new program called bico added to the existing Bsoft package (37). An overview of our workflow is given in Fig. 2. From reconstructed tomograms, subtomogram volumes were extracted and aligned to an icosahedral reference, in our case a reconstruction of a HSV-1 C-capsid (accession number EMDB-7472 (38)).

Next, the fivefold vertices of each capsid were extracted, and an average of all extracted and

correctly oriented vertices was calculated. To focus only on the central part of a vertex where we

expect the biggest difference between penton and portal, we used a soft cylindrical mask in the

comparisons of each vertex with the average (see Materials and Methods for mask details). For

each set of 12 vertices, bico then classified each vertex set into two subsets: the first subset

contained the eleven best-fitting vertices. The second subset contained the vertex with the worst

fit, taken to be the portal vertex. The coordinates of the identified portal vertices were then

transferred back into the extracted and aligned capsids, and a reconstruction of the capsid

featuring the portal vertex was calculated. To reduce noise, this reconstruction was symmetrized

fivefold in the final step and filtered to 30 Å (just beyond the 34 Å resolution estimated by Fourier

shell correlation – see Fig. S1).

Fig. 2. Workflow used to identify and average the portal vertices. Individual particles (volumes

each containing a capsid) extracted from tomograms and aligned with an icosahedral reference

using a frequency space mask to compensate for the missing wedge. This allows the location of

the twelve 5-fold vertices of each capsid, which are then extracted and averaged. Each vertex is

correlated to this average using a missing wedge mask in frequency space and a cylindrical real

space soft mask to isolate the region in a vertex occupied by the penton or portal (cyan dashed

line). The vertex with the lowest (worst) correlation is assumed to contain the portal. This is

followed by a further refinement to find the best 5-fold orientation around the vertex axis. The

coordinates of the portal vertices were then used to transform their corresponding capsids into

orientations that placed the portal vertex on the z-axis at the top of the volume.

The portal-bearing vertex of UL25-Null A-capsids. We extracted a total of 164 subtomogram

volumes of A-capsids from our tomograms. We identified the likely portal-bearing vertex in the

map of each of the A-capsids and calculated an average (Fig. 3), obtaining a resolution of 34 Å

(Fig. S1). In line with recent publications (18, 32, 33), we visualized the portal on the inside of the

capsid, with its wider end facing towards the center of the particle (Fig. 3A). A superposed atomic

model of the HSV-1 portal (PBD ID 6OD7 (32)) fits well into the portal density (Fig. 3B). The portal

has a narrow end called the "clip domain" that engages the triplexes (green arrow in Fig. 3B) (for

the assignment of this density as a triplex, see Huet et al. (31)). The widest part of the portal, the

"wing domain" extends towards the floors of the neighboring capsomers (blue arrow in Fig. 3B).

The overall appearance and placement of this connection is consistent with a set of β-barrels and

spine helices originating from the pUL19 hexons that were identified earlier in a high-resolution

reconstruction (32). Knowing what it looks like, we could identify the portal vertex in individual

particles in our tomograms (white arrows in Fig. 3C).

Fig. 3. The portal vertex in UL25-Null A-capsids. A. Central slice of the C5-symmetrized capsid
with the portal vertex located at the top. Scale bar, 200 Å. B. Magnified view of the vertex. The
portal is located beneath the capsid floor with its wider end facing the inside of the particle. A
high-resolution model (pink, PDB ID 6OD7 (32)) is superimposed on the left half of the portal
density. The portal forms connections to the triplex (green arrow) and the capsid floor (blue
arrow indicates a link). Note that there is a diffuse density above the portal vertex that could be
residual terminase. Scale bar, 100 Å. C. Slices from tomograms of A-capsids that traverse the
portal in each (white arrowheads). Scale bar, 500 Å.

The portal in procapsids interacts with the scaffold. We selected tomograms with well-
separated spherical particles (Fig. 1A) and extracted 368 subvolumes. As with A-capsids, we
identified the portal-bearing vertex, taking care to adjust the mask so that it excluded the
scaffold, which might otherwise outweigh the portal density during vertex sorting. After
identifying the portal vertex in each procapsid, we calculated an average (Fig. 4) with a resolution
of 34 Å (Fig. S1). Consistent with the A-capsids, the portal is located inside the capsid shell (Fig.

4A), and again agrees with an atomic portal model (Fig. 4B). The portal penetrates deeply into

the scaffold layer (yellow arrow in Fig. 4B), almost to the widest part of its wing, a position that

allows the formation of extensive interactions. Interestingly, the nominally spherical scaffold

shows a degree of perturbation caused by the portal, giving the scaffold an overall teardrop

shape (Fig. 4A). We also note that – apart from in the portal region - the scaffold shell shows up

in slices as three concentric shells (arrows in Fig. 4A) consistent with previous studies (5, 7). Only

tenuous contacts (blue arrow in Fig. 4B) are formed between the portal wing domain and the

surrounding hexons, but the clip domain appears to interact with the triplexes (green arrow in

Fig. 4B). The vertex overall has a funnel-like shape that likely accommodates DNA insertion by

the terminase complex. Finally, the procapsid hexons also display faint residual density on their

exterior parts that we attribute to the 6F10 antibody used to stabilize the procapsids during

isolation (Fig. 4A, 5C).

Fig. 4. The portal vertex in procapsids. A. Central slice of the C5-symmetrized procapsid with

the portal vertex at the top, showing distinct rings of scaffold density (yellow arrows). Scale bar,

200 Å. **B.** Magnified view of the portal vertex with a superimposed atomic model (pink, PDB ID

6OD7 (32)). The portal is mainly connected to the capsid through the triplex (green arrow) with

a thin connection to the capsid floor (blue arrow). The bottom of the portal is embedded in the

scaffold (yellow arrow). Scale bar, 100 Å.

Conformational changes and movement of the portal during maturation. While the portal

orientation is similar in procapsids and A-capsids, contrasts in vertex configuration come to light

in a side-by-side comparison (Fig. 5). Upon maturation, the capsid becomes angular, causing all

the vertices (both penton and portal) to translocate outwards. We assessed the movement of

the pentons by shifting the penton vertex of the procapsid outwards until it overlays the A capsid

penton, yielding a translation of 29 Å. Similarly, we measured the translation of the portal to be

54 Å, thus moving closer to the capsid shell. The portal in Fig. 5A appears to be somewhat

separated from the capsid shell but is much closer in the mature capsid (Fig. 5B). During

maturation, the capsid shell stabilizes by closure of the floor and burying interfaces between the

capsomers (8). Our results indicate that the portal also adopts a more stable interaction with the

mature capsid shell.

In addition, we observe a shift in the triplex network surrounding the portal vertex, which

narrows the vertex channel. These changes become even more apparent when the vertices are

viewed from the top (Fig. 5C,D). While triplexes are somewhat weakly defined in the procapsid

portal vertex, they become much more visible in the A-capsid and even extend towards the

fivefold symmetry axis. It is possible that some of this extending density could be ascribed to

components of the tegument, particularly pUL17. We recall that, upon maturation, the hexons

rearrange from a skewed orientation to true hexameric symmetry (8).

We also compared the triplexes of the portal vertex with those of the opposing penton vertex in

both maps. The triplexes in the procapsid portal vertex are poorly defined (Fig. 6A), making it

hard to ascertain whether they are in the same conformation as those at the opposing penton

vertex (Fig. 6B). In contrast, the orientations of the triplexes in the A-capsid are clearly the same

for both vertices (Fig. 6C,D).

Fig. 5. Comparison of portal vertices in procapsid and UL25-Null A-capsid. A. Side view cut

through the procapsid portal vertex, showing the portal (orange) connecting to the capsid shell

(blue) and embedded in the scaffold (yellow). **B.** Side view of the mature A-capsid portal vertex,

with the portal (orange) very closely associated with the capsid shell (blue). The diffuse density

above the vertex (pink) may be residual terminase. **C.** Top view of the procapsid portal vertex

with faint density associated with the tops of the hexons attributed to the 6F10 antibody used

in the preparation of the particles (green arrow). **D.** Top view of the A-capsid portal vertex.

Upon maturation, the triplexes surrounding the vertex shift and constrict the vertex channel,

and the hexons transform from skewed ovals towards hexagonal symmetry. Scale bars, 100 Å.

Fig. 6. The arrangement of triplexes at the vertices. The triplexes (yellow stars) at the

procapsid portal vertex (**A**) are recessed relative to those at the opposite penton vertex (**B**)

(slices at a radius of 307 Å). In contrast, the triplexes at the A-capsid portal vertex (**C**) are closer

in conformation to those at the opposite penton vertex (**D**) (slices at a radius of 325 Å). Note

that the vertices in the procapsid move outwards by ~29 Å on maturation, accounting for the

differences in slices shown here. Scale bars, 100 Å.

The number of portal vertices per capsid. While we implicitly assumed there is exactly one

portal per capsid, that is not necessarily true. To determine the number of portal vertices in

each capsid, we examined the distribution of correlation coefficients for the 12 vertices. Our

aim was to generate two clusters: the few with the worst correlation coefficients indicating

portal vertices, and the rest as penton vertices. To separate the clusters, we took the biggest

difference between successive correlation coefficients in a sorted list as the distinction between

penton and portal vertices. Because we expected only a few portal vertices, only the six with

the lowest coefficients were assessed. However, this strategy assumes there is at least one

portal vertex. We compared the difference between the two worst vertices with the next three

differences and assumed that when it is the lowest, it indicated a capsid without a portal

vertex. The resultant distributions are shown in Fig. 7, giving average portal vertices per capsid

of 1.07 for procapsids and 1.25 for A-capsids, consistent with previous estimations (16, 27). In

our subtomogram averages, we included all of the capsids, but we do not consider the minor

contribution of the capsids without portals to significantly change our conclusions. About 65 %

of the capsids have a single portal vertex, in agreement with the 73 % reported for A-capsids

(28). A still significant number (~20 %) have two portal vertices, while those with three are rare.

If the incorporation of portal vertices follows a random pattern, the expected distribution

would be binomial in nature (blue lines in Fig. 7). The predominance of a singly incorporated

portal indicates that the assembly is more specific than a random process.

Fig. 7. Estimating the number of portal vertices (bars) per procapsid (A) an A-capsid (B). The

blue line in each plot is a binomial distribution calculated from the average number of portal

vertices per capsid as an indication of expected random incorporation.

Discussion

Localization and interactions of the portal in procapsids compared to A-capsids. In a critical step

of the HSV-1 life cycle, newly synthesized DNA is threaded into the capsid which undergoes

extensive conformational changes as it matures (5, 8) although the latter process can occur

without DNA packaging (8) and may precede DNA packaging *in vivo* (39). In both herpesviruses

and tailed bacteriophages, DNA is inserted through a fivefold vertex via the dodecameric portal

protein and driven by terminase complex. While the overall structure of the HSV-1 portal, pUL6,

has been known for some time (15), its orientation and placement in the context of the capsid

shell have been subject to debate. The portal's overall shape makes it difficult to distinguish

visually from capsomer pentons in micrographs, making consistent alignment of 2D projections

a challenge. In an earlier study, we used cryoET, aiming to characterize the portal vertex of HSV-

1 (28). Here, we have refined this approach by developing a new program designed to identify

the portal vertex on individual capsids, thereby allowing coherent averaging. Applying it to HSV-

1 procapsids and A-capsids, we located the portal on the inside of the particle, consistent with

several recent SPA-based studies (18, 30, 32-34). In confirmation, an atomic model of pUL6 (32)

fits well into our portal density (Fig. 8). This density extends beyond the clip domain of the model,

presumably representing some of residues 308-515, which were not modeled (corresponding

residues are also unmodeled in a reconstruction of the Epstein-Barr-virus portal (40)).

Interestingly, a corresponding turret-shaped density was discovered in the portal vertex of

Kaposi's Sarcoma Herpesvirus (KSHV) (34); it was theorized that this helical turret is an extension

of the portal's clip domain and could be a docking site for the terminase complex. If such a

component is indeed an additional protein, it may only be added to mature particles upon the

successful retention of DNA.

Fig 8. **The model of the portal within the context of the procapsid portal vertex. A.** The portal

model (PDB ID 6OD7 (32)) fitted into the portal density (orange) within a lower threshold

envelope (grey) to show the connections to the triplexes around the portal (green) and the

scaffold (yellow). The connections to the capsid shell (composed of the major capsid protein

and other triplexes shown in blue) are not resolved in this map. Scale bar, 100 Å. **B.** Close-up

view of the portal with key tryptophan residues (yellow balls) at the wing (orange arrow: W163,

W241, W262, W532) and at the bottom (blue arrow: W90, W127) that were shown to interact

with the scaffold (41). Scale bar, 40 Å.

Applying our workflow to procapsids led to several novel observations. The procapsid portal is

located inside the vertex forming contacts with the scaffolding shell and deforming the scaffold

proteins' spherical arrangement at the interaction site. Conserved tryptophan residues have

been identified in HSV-1 pUL6 ((41), see also Fig. 2 in (42)) that play crucial roles in interaction

between portal and, most likely, conserved scaffold residues 143-151 of pUL26, particularly

YYPGE (43, 44). Figure 8B shows these critical tryptophan residues arrange in a ring in the wing

domain (orange arrow in Fig. 8B) as well as a ring in the base of the portal (blue arrow in Fig. 8B).

We know the scaffold C-terminus is bound to the capsid (45) which puts it in the proximity of the

portal wing domain. Conserved tryptophan residues are a feature not only of HSV-1 but also of

some bacteriophages ((46), reviewed in (42)). The scaffold probably shares the symmetry of the

surface shell with which it co-assembles but is not highly ordered and its apparent sphericity is

enhanced by averaging. From previous studies we know that the protease at the N-terminal part

of about 10 % of the scaffold proteins is located towards the center of the procapsid (4) and the

C-terminus interacts with the capsid shell (47) and portal (43). This means the protein is in an

extended radial conformation consistent with a side-by-side packing of coiled coil dimers as

inferred by Pelletier et al. (48). In the procapsid we see three peaks in the concentric density

shells (Fig. 4) in agreement with the three domains proposed by Trus et al. (7).

Incorporation of the portal during procapsid assembly. The stage at which the portal is

incorporated into the procapsid is still unknown, with possibilities that it could initiate assembly,

inserted at some intermediate point, or at the end (16). Procapsids can assemble *in vitro* without

portal present (2, 3) and some of the capsids we analyzed appear to lack a portal vertex (Figure

7). We also found an average copy number slightly larger than one and consistent with previous

studies (16, 27). If one assumes a random model of incorporation, it is expected to follow a

binomial distribution. However, Figure 7 shows it clearly deviates from a binomial pattern,

suggesting a predisposition towards a single portal vertex per capsid. This suggests that the portal

may facilitate procapsid assembly and thus ensure that a majority has at least one portal vertex.

However, if it initiates assembly, it does not preclude incorporation of one or two additional

portals.

Morphological changes upon capsid maturation. Cleavage of the C-terminal 25 residues of the

scaffolding protein by the protease (47) triggers maturation of the capsid shell. Our preparation

of procapsids is defective in the protease to allow us to purify it. Nevertheless, the capsids are

still able to mature slowly (8). On cleavage, the scaffold changes from extended in the procapsid

(i.e., large core particles) to somewhat contracted in the B-capsid (i.e., small core particles). At

the same time, the vertices of the capsid shell shift outwards during angularization by ~29 Å, with

the portal itself moving even further by ~54 Å (note that the capsid shells in A- and B-capsids are

very similar (31)). The portal therefore disengages from the scaffold to some extent, while

strengthening interactions with the capsid shell. The portal movement is accompanied by the

establishment of a support network with the newly established capsid floor (termed suspension

network in another study (32)).

The overall shape of the portal vertex differs significantly between procapsids and mature capsids

(Fig. 6). In procapsids, we note that the center of the vertex is essentially open, which would, in

a biological context, be advantageous for the insertion of DNA. Triplex density is present around

the portal vertex, but it is somewhat fainter than around a penton vertex, which could result

from a lack of interaction between the portal and the procapsid itself. In the mature capsid, the

vertex is narrower, and it appears that density originating at the triplexes extends towards the

vertex axis. This extended density may indeed be contributed by components of the tegument,

particularly pUL17. While our A-capsids lack pUL25, it has been shown that procapsids contain

practically no pUL25 (25), and given that the procapsid undergoes substantial conformational

changes during maturation, it is likely that no tegument component protein is present on these

immature particles. The extended, star-shaped density we see on the A-capsid portal vertex

(Fig. 5D) may therefore act as a docking platform for copies of pUL25 that would be added in the

course of a wild type infection and α-helices contributed by pUL36 at a later stage. This platform,

however, is evidently insufficient to retain DNA in the capsid, as the absence of pUL25 alone is

enough to completely prevent the emergence of propagation-competent, DNA-filled C-capsids

(36), underscoring pUL25's functional versatility as a DNA retention protein and a key player in

nuclear egress as well (49).

Conclusion. Our routine for processing cryo-electron tomograms enabled us to visualize the

portal in procapsids in comparison with mature A-capsids. Particularly revealing is the

interactions of scaffold with the portal likely to be involved in procapsid assembly. Quantitation

of the number of portal vertices per capsid further solidifies a role in assembly. We also gained

insight into the intricate maturation-induced changes in the portal vertex, visualizing an outward

movement of the portal exceeding vertex movement and thus stabilizing the interaction with the

capsid shell.

Material and Methods

Cell handling and propagation of virus stocks. Stocks of the UL25 deletion mutant (UL25-null)

(36) were generated by using a complementary African Green Monkey cell line (both gifts from

Prof. Fred L. Homa, University of Pittsburgh, Pittsburgh, PA). At roughly 75% confluency,

monolayers in 175 cm^2 flasks were infected with virus stock at a MOI of 0.1 in 1 mL of PBS without

calcium or magnesium per flask for 40 min at room temperature, followed by 5 min at 37˚C. The

cells were then overlaid with 15 mL virus growth medium (1x MEM Alpha, Corning, 1.5% (*v/v*)

penicillin/streptomycin) and incubated at 37˚C. At 48 h post infection, virus-containing medium

was clarified by centrifugation at 1,500 rpm for 5 min, and crude cell debris was removed from

the supernatant by pelleting at 6,000 rpm for 15 min in a Beckman SW28 rotor. Virus was pelleted

for 45 min at 19,000 rpm in a Beckman SW28 rotor. The virus pellets were resuspended in small

volumes of PBS without calcium or magnesium, and aliquots of 500 μL were frozen at -80˚C. The

virus titer was assessed in complementary cells 48 hr. after infection.

Stocks of the procapsid-producing M100 line (35) were generated similarly using the

complementary F-3 cell line. Stock titers were assessed as described earlier (5).

Production of capsids. Roller bottles of African Green Monkey cells were grown to about 75%

confluency and infected with UL25-null virus stocks at a MOI of 10. 14 hr post infection, cells had

started to round up, but not detach from the surface. Cells were harvested by centrifugation at

1,500 rpm for 5 min, and any virus still attached to the cell surface was removed by incubation

with TNE (20 mM Tris, pH 7.4, 500 mM NaCl, 1 mM EDTA) on ice for 15 min. Cells were again

pelleted for 5 min at 2,000 rpm, resuspended in TNE, and broken up by sonication. 1.5% (*v/v*)

TX-100 was added to solubilize nuclear membranes, followed by pelleting of crude cell debris for

30 min at 6,000 rpm in a Beckman SW28 rotor. The resulting supernatant was supplemented with

a cushion of 30% (*w/v*) sucrose in TNE, and a mix of A- and B-capsids was pelleted for 1 h at

20,000 rpm in a Beckman SW28 rotor. The pellets were resuspended in TNE, supplemented with

complete protease inhibitor cocktail (Roche), and applied to a preparative gradient containing a

range of 20-50% (*w/v*) sucrose in TNE. The gradient was run for 1 h at 22,000 rpm in a Beckman

SW55Ti rotor. Distinct bands containing the different capsid species were isolated, pelleted for

50 min at 22,000 rpm in a Beckman SW55Ti rotor, and resuspended in small amounts of TNE to

reach an appropriate concentration for cryoET.

Procapsids of the M100 mutant in the viral protease were produced as described earlier (5). They

were precipitated with the 6F10 antibody (raised against the triplex protein pUL19) for

purification and capsid stability purposes (4).

Grid preparation and cryo-electron tomography. Purified capsids were mixed with Aurion

Anionic 10 nm BSA gold tracers (EMS) at a 1:1 ratio, and 3 μL were applied to Quantifoil copper

grids covered with 300-mesh holey carbon (EMS), which were previously plasma-cleaned for 12

s with an argon-oxygen mixture (25% oxygen) in a Model 1020 plasma cleaner (Fischione, Export,

PA). Using a Leica EM GP plunger (Leica Microsystems, Buffalo Grove, IL), excess liquid was

blotted off for 2.5 s, and grids were flash-frozen in liquid ethane. For the A-capsids, 21 tilt series

ranging from -57˚ to +57˚ at 3˚ increments were collected at 20,000x magnification and a defocus

target of -2.5 μm on a JEOL JEM-2200FS cryo-electron microscope (JEOL USA, Peabody, MA)

operating at 200 kV and equipped with an in-column energy filter and a K2 Summit detector

(Gatan, Warrendale, PA). For procapsids, 80 tilt series ranging from -57˚ to +57˚ at 3˚ increments

at 81,000x magnification and a -2 µm defocus target were acquired on a 300 kV Titan Krios

(Thermo Fisher Scientific, FEI, Hillsboro, OR) equipped with a post-column Model 967 GIF

Quantum-LS with a K2 Summit detector (Gatan, Warrendale, PA). In both cases, data were

acquired using SerialEM (50).

Tomogram reconstruction and identification of the portal-bearing vertex. Image processing and

tomogram reconstruction were carried out using Bsoft (50, 51). Tomograms were binned 2-fold,

resulting in a final pixel size of 3.66 Å for A-capsids and 3.61 Å for procapsids. For both capsids,

promising particles were extracted in 450 x 450 x 450 pixel boxes (program bpick) and aligned

(program bfind) to a reference of a HSV-1 C-capsid (38) and using an appropriate frequency space

mask to compensate for the missing wedge (created with program bmissing). Given the correct

alignment parameters, the program bpick was used to extract the 12 vertices for each capsid

(150 x 150 x 150 pixel boxes) offset 140 pixels outward from the center of the particle on the

fivefold symmetry axes. Using the new Bsoft program, bico, for each set of 12 vertices was cross-

correlation against an average calculated from all vertices, and the vertex with the lowest

correlation was deemed the portal-bearing vertex. As for the whole capsid alignment, a

frequency space mask was used in the correlation calculation (created with program bmissing).

In addition, a real space mask was used to isolate a cylindrical region occupied by the penton at

the vertex. The mask was created with program beditimg specifying a cylinder centered at

75,75,45 with radius 20 and height 90, and a soft edge with a gaussian width of 3 pixels. The

softness of this mask was tuned to avoid high frequency correlations in the comparison of vertex

volumes. To address the symmetry mismatch between the fivefold vertex and the twelvefold

portal, the best 5-fold orientations of the portal-bearing vertices were found and refined

(program bico). The coordinates corresponding to the portal vertices were then transferred to

the parameter file referencing the full capsid volumes (program bico). Finally, the average of the

whole capsids was calculated with the portal vertex located at the top of the volume. Graphics

were prepared using PyMOL (52) and UCSF Chimera (53).

Data Availability:

Maps for the UL25-Null A-capsid and the procapsid, both showing the portal vertex, were

deposited in the Electron Microscopy Database under accession numbers EMDB-22378 and

EMDB-22379, respectively.

Acknowledgements

We thank Dr. Fred L. Homa (University of Pittsburgh) for providing us with the HSV-1 UL25-Null

mutants and the complementary, UL25-expressing African Green Monkey cell line as well as a

wild type African Green Monkey cell line. We thank Dr. Jay Brown for useful discussions. This

work was supported by the Intramural Research Program of the National Institute for Arthritis,

Musculoskeletal and Skin Diseases, NIH and used the NIH Multi-Institute CryoEM Facility (MICEF).

Competing Interests:

The authors declare no competing interests.

References:

1.   Mettenleiter TC, Klupp BG, Granzow H. 2009. Herpesvirus assembly: an update. Virus
Res 143:222-34.
2.   Newcomb WW, Homa FL, Thomsen DR, Booy FP, Trus BL, Steven AC, Spencer JV, Brown
JC. 1996. Assembly of the herpes simplex virus capsid: characterization of intermediates
observed during cell-free capsid formation. Journal of Molecular Biology 263:432-46.
3.   Newcomb WW, Homa FL, Thomsen DR, Trus BL, Cheng N, Steven A, Booy F, Brown JC.
1999. Assembly of the herpes simplex virus procapsid from purified components and
identification of small complexes containing the major capsid and scaffolding proteins.
Journal of Virology 73:4239-50.
4.   Newcomb WW, Trus BL, Cheng N, Steven AC, Sheaffer AK, Tenney DJ, Weller SK, Brown
JC. 2000. Isolation of herpes simplex virus procapsids from cells infected with a
protease-deficient mutant virus. Journal of Virology 74:1663-73.
5.   Aksyuk AA, Newcomb WW, Cheng N, Winkler DC, Fontana J, Heymann JB, Steven AC.
2015. Subassemblies and asymmetry in assembly of herpes simplex virus procapsid.
mBio 6:e01525-15.
6.   Huet A, Makhov AM, Huffman JB, Vos M, Homa FL, Conway JF. 2016. Extensive subunit
contacts underpin herpesvirus capsid stability and interior-to-exterior allostery. Nat
Struct Mol Biol 23:531-9.
7.   Trus BL, Booy FP, Newcomb WW, Brown JC, Homa FL, Thomsen DR, Steven AC. 1996.
The herpes simplex virus procapsid: structure, conformational changes upon

maturation, and roles of the triplex proteins VP19c and VP23 in assembly. J Mol Biol
263:447-62.
8.   Heymann JB, Cheng N, Newcomb WW, Trus BL, Brown JC, Steven AC. 2003. Dynamics of
herpes simplex virus capsid maturation visualized by time-lapse cryo-electron
microscopy. Nat Struct Biol 10:334-41.
9.   Booy FP, Trus BL, Newcomb WW, Brown JC, Conway JF, Steven AC. 1994. Finding a
needle in a haystack: detection of a small protein (the 12-kDa VP26) in a large complex
(the 200-MDa capsid of herpes simplex virus). Proceedings of the National Academy of
Sciences, USA 91:5652-6.
10. Trus BL, Homa FL, Booy FP, Newcomb WW, Thomsen DR, Cheng N, Brown JC, Steven AC.
1995. Herpes simplex virus capsids assembled in insect cells infected with recombinant
baculoviruses: structural authenticity and localization of VP26. Journal of Virology
69:7362-6.
11. Bauer DW, Huffman JB, Homa FL, Evilevitch A. 2013. Herpes virus genome, the pressure
is on. J Am Chem Soc 135:11216-21.
12. Johnson DC, Baines JD. 2011. Herpesviruses remodel host membranes for virus egress.
Nat Rev Microbiol 9:382-94.
13. Mettenleiter TC, Muller F, Granzow H, Klupp BG. 2013. The way out: what we know and
do not know about herpesvirus nuclear egress. Cell Microbiol 15:170-8.
14. Heming JD, Conway JF, Homa FL. 2017. Herpesvirus Capsid Assembly and DNA
Packaging. Adv Anat Embryol Cell Biol 223:119-142.
15. Trus BL, Cheng N, Newcomb WW, Homa FL, Brown JC, Steven AC. 2004. Structure and
polymorphism of the UL6 portal protein of herpes simplex virus type 1. J Virol 78:12668-
71.
16. Newcomb WW, Thomsen DR, Homa FL, Brown JC. 2003. Assembly of the herpes simplex
virus capsid: identification of soluble scaffold-portal complexes and their role in
formation of portal-containing capsids. J Virol 77:9862-71.
17. Trus BL, Newcomb WW, Cheng N, Cardone G, Marekov L, Homa FL, Brown JC, Steven
AC. 2007. Allosteric signaling and a nuclear exit strategy: binding of UL25/UL17
heterodimers to DNA-Filled HSV-1 capsids. Mol Cell 26:479-89.
18. McElwee M, Vijayakrishnan S, Rixon F, Bhella D. 2018. Structure of the herpes simplex
virus portal-vertex. PLoS Biol 16:e2006191.
19. Liu YT, Jiang J, Bohannon KP, Dai X, Gant Luxton GW, Hui WH, Bi GQ, Smith GA, Zhou ZH.
2017. A pUL25 dimer interfaces the pseudorabies virus capsid and tegument. J Gen Virol
98:2837-2849.
20. Heming JD, Huffman JB, Jones LM, Homa FL. 2014. Isolation and characterization of the
herpes simplex virus 1 terminase complex. J Virol 88:225-36.
21. al-Kobaisi MF, Rixon FJ, McDougall I, Preston VG. 1991. The herpes simplex virus UL33
gene product is required for the assembly of full capsids. Virology 180:380-8.
22. Baines JD, Poon AP, Rovnak J, Roizman B. 1994. The herpes simplex virus 1 UL15 gene
encodes two proteins and is required for cleavage of genomic viral DNA. J Virol 68:8118-
24.

23.   Yu D, Weller SK. 1998. Herpes simplex virus type 1 cleavage and packaging proteins
550   UL15 and UL28 are associated with B but not C capsids during packaging. J Virol
551   72:7428-39.
552   24.   Adelman K, Salmon B, Baines JD. 2001. Herpes simplex virus DNA packaging sequences
553   adopt novel structures that are specifically recognized by a component of the cleavage
554   and packaging machinery. Proc Natl Acad Sci U S A 98:3086-91.
555   25.   Sheaffer AK, Newcomb WW, Gao M, Yu D, Weller SK, Brown JC, Tenney DJ. 2001. Herpes
556   simplex virus DNA cleavage and packaging proteins associate with the procapsid prior to
557   its maturation. J Virol 75:687-98.
558   26.   White CA, Stow ND, Patel AH, Hughes M, Preston VG. 2003. Herpes simplex virus type 1
559   portal protein UL6 interacts with the putative terminase subunits UL15 and UL28. J Virol
560   77:6351-8.
561   27.   Newcomb WW, Juhas RM, Thomsen DR, Homa FL, Burch AD, Weller SK, Brown JC. 2001.
562   The UL6 gene product forms the portal for entry of DNA into the herpes simplex virus
563   capsid. J Virol 75:10923-32.
564   28.   Cardone G, Winkler DC, Trus BL, Cheng N, Heuser JE, Newcomb WW, Brown JC, Steven
565   AC. 2007. Visualization of the herpes simplex virus portal in situ by cryo-electron
566   tomography. Virology 361:426-34.
567   29.   Chang JT, Schmid MF, Rixon FJ, Chiu W. 2007. Electron cryotomography reveals the
568   portal in the herpesvirus capsid. J Virol 81:2065-8.
569   30.   Rochat RH, Liu X, Murata K, Nagayama K, Rixon FJ, Chiu W. 2011. Seeing the portal in
570   herpes simplex virus type 1 B capsids. J Virol 85:1871-4.
571   31.   Huet A, Huffman JB, Conway JF, Homa FL. 2020. The role of the herpes simplex virus
572   CVSC proteins at the capsid portal vertex. J Virol doi:10.1128/JVI.01534-20.
573   32.   Liu YT, Jih J, Dai X, Bi GQ, Zhou ZH. 2019. Cryo-EM structures of herpes simplex virus
574   type 1 portal vertex and packaged genome. Nature 570:257-261.
575   33.   Wang N, Chen W, Zhu L, Zhu D, Feng R, Wang J, Zhu B, Zhang X, Chen X, Liu X, Yan R, Ni
576   D, Zhou GG, Liu H, Rao Z, Wang X. 2020. Structures of the portal vertex reveal essential
577   protein-protein interactions for Herpesvirus assembly and maturation. Protein Cell
578   11:366-373.
579   34.   Gong D, Dai X, Jih J, Liu YT, Bi GQ, Sun R, Zhou ZH. 2019. DNA-Packing Portal and Capsid-
580   Associated Tegument Complexes in the Tumor Herpesvirus KSHV. Cell 178:1329-1343
581   e12.
582   35.   Gao M, Matusick-Kumar L, Hurlburt W, DiTusa SF, Newcomb WW, Brown JC, McCann PJ,
583   3rd, Deckman I, Colonno RJ. 1994. The protease of herpes simplex virus type 1 is
584   essential for functional capsid formation and viral growth. J Virol 68:3702-12.
585   36.   McNab AR, Desai P, Person S, Roof LL, Thomsen DR, Newcomb WW, Brown JC, Homa FL.
586   1998. The product of the herpes simplex virus type 1 UL25 gene is required for
587   encapsidation but not for cleavage of replicated viral DNA. Journal of Virology 72:1060-
588   70.
589   37.   Heymann JB. 2018. Guidelines for using Bsoft for high resolution reconstruction and
590   validation of biomolecular structures from electron micrographs. Protein Sci 27:159-171.
591   38.   Dai X, Zhou ZH. 2018. Structure of the herpes simplex virus 1 capsid with associated
592   tegument protein complexes. Science 360.

39. Church GA, Dasgupta A, Wilson DW. 1998. Herpes simplex virus DNA packaging without
measurable DNA synthesis. J Virol 72:2745-51.
40. Machon C, Fabrega-Ferrer M, Zhou D, Cuervo A, Carrascosa JL, Stuart DI, Coll M. 2019.
Atomic structure of the Epstein-Barr virus portal. Nat Commun 10:3891.
41. Yang K, Baines JD. 2009. Tryptophan residues in the portal protein of herpes simplex
virus 1 critical to the interaction with scaffold proteins and incorporation of the portal
into capsids. J Virol 83:11726-33.
42. Dedeo CL, Cingolani G, Teschke CM. 2019. Portal Protein: The Orchestrator of Capsid
Assembly for the dsDNA Tailed Bacteriophages and Herpesviruses. Annu Rev Virol
6:141-160.
43. Singer GP, Newcomb WW, Thomsen DR, Homa FL, Brown JC. 2005. Identification of a
region in the herpes simplex virus scaffolding protein required for interaction with the
portal. J Virol 79:132-9.
44. Yang K, Baines JD. 2009. Proline and tyrosine residues in scaffold proteins of herpes
simplex virus 1 critical to the interaction with portal protein and its incorporation into
capsids. J Virol 83:8076-81.
45. Hong Z, Beaudet-Miller M, Durkin J, Zhang R, Kwong AD. 1996. Identification of a
minimal hydrophobic domain in the herpes simplex virus type 1 scaffolding protein
which is required for interaction with the major capsid protein. J Virol 70:533-40.
46. Guasch A, Pous J, Ibarra B, Gomis-Ruth FX, Valpuesta JM, Sousa N, Carrascosa JL, Coll M.
2002. Detailed architecture of a DNA translocating machine: the high-resolution
structure of the bacteriophage phi29 connector particle. J Mol Biol 315:663-76.
47. Thomsen DR, Newcomb WW, Brown JC, Homa FL. 1995. Assembly of the herpes simplex
virus capsid: requirement for the carboxyl-terminal twenty-five amino acids of the
proteins encoded by the UL26 and UL26.5 genes. J Virol 69:3690-703.
48. Pelletier A, Do F, Brisebois JJ, Lagace L, Cordingley MG. 1997. Self-association of herpes
simplex virus type 1 ICP35 is via coiled-coil interactions and promotes stable interaction
with the major capsid protein. J Virol 71:5197-208.
49. Draganova EB, Zhang J, Zhou ZH, Heldwein EE. 2020. Structural basis for capsid
recruitment and coat formation during HSV-1 nuclear egress. Elife 9.
50. Mastronarde DN. 2005. Automated electron microscope tomography using robust
prediction of specimen movements. J Struct Biol 152:36-51.
51. Heymann JB. 2021. High resolution electron tomography and segmentation-by-
modeling interpretation in Bsoft. Protein Science 30:44-59.
52. Schrodinger L. 2015. The PyMOL Molecular Graphics System, Version 18:1.
53. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE.
2004. UCSF Chimera--a visualization system for exploratory research and analysis. J
Comput Chem 25:1605-12.
631
632

633 **Supplementary material**

636 **Fig. S1.** Fourier shell correlation of subtomogram half-set averages of procapsids (red) and A-

637 capsids (blue) with the resolution indicated at a threshold of 0.143 (dashed line). A soft mask

638 covering the capsid shell and portal vertex was used in the analysis.