A time-lagged association between the gut microbiome, nestling growth and nestling

survival in wild great tits

Gabrielle L. Davidson[a,b,*✉] , Shane E. Somers[a,*], Niamh Wiley[c,d], Crystal N. Johnson[c,d], Michael

S. Reichert[a,e], R. Paul Ross[c,d], Catherine Stanton[c,d], John L. Quinn[a]

*Joint first authors

✉ Correspondence: gd339@cam.ac.uk

a. School of Biological, Earth and Environmental Sciences, Distillery Fields, North Mall,

University College Cork, Cork, Ireland.

b. Department of Psychology, Downing Street, University of Cambridge, Cambridge,

UK.

c. APC Microbiome Ireland, University College Cork, Cork, Ireland.

d. Teagasc Food Research Centre, Moorepark, Fermoy, Ireland.

e. Department of Integrative Biology, Oklahoma State University, USA

Abstract:

1. Natal body mass is a key predictor of viability and fitness in many animals. While

variation in body mass and therefore viability of juveniles may be explained by

genetic and environmental factors, emerging evidence points to the gut microbiota

as an important factor influencing host health. The gut microbiota is known to

change during development, but it remains unclear whether the microbiome

predicts fitness, and if it does, at which developmental stage it affects fitness traits.

2. We collected data on two traits associated with fitness in wild nestling great tits

(*Parus major*): weight and survival to fledging. We characterised the gut microbiome

using 16S rRNA sequencing from nestling faeces and investigated temporal

associations between the gut microbiome and fitness traits across development at

day 8 (D8) and day 15 (D15) post-hatching. We also explored whether particular

microbial taxa were 'indicator species' that reflected whether nestlings survived or

not.

3. There was no link between mass and microbial diversity on D8 or D15. However, we
detected a time-lagged relationship whereby the microbial diversity at D8 was
negatively associated with weight at D15, while controlling for weight at D8,
therefore reflecting weight gain over the intervening period. Indicator species
analysis revealed that while several taxa were unique to birds that either survived or
did not survive, there were no universal taxa that were consistently found across all
birds within either survival group. This suggests that the presence of particular
bacterial taxa may be sufficient, but not necessary for determining future survival,
perhaps owing to functional overlap in microbiota.

4. We highlight that measuring microbiome-fitness relationships at just one time point
may be misleading, especially early in life. Instead, microbial-host fitness effects
should be investigated longitudinally as there may be critical development windows
in which key microbiota are established and prime host traits associated with
nestling growth. Pinpointing which features of the gut microbial community impact
on host fitness, and when during development this occurs, will shed light on
population level processes and has the potential to support conservation.

Introduction:

Understanding the numerous factors that drive fitness has long been the focus of research
in ecology and evolution (Merila, 1996; Pickett et al., 2013). This has primarily included
behavioural factors, such as foraging (Patrick & Weimerskirch, 2014) and parental care
(Schwagmeyer & Mock, 2008), and ecological factors, for example, food availability (Van
Noordwijk et al., 1995), predation (Götmark, 2002), and more recently climate change
(Blomberg et al., 2014). Proximate factors such as genetics (e.g. Merila (1996), Norris
(1993)), stress (Crino & Breuner, 2015; Weber et al., 2018) and natal environment effects
(Keller & Van Noordwijk, 1994; Pickett et al., 2013) have also been studied intensively. More
recently, a growing body of research is beginning to show the gut microbiome - the
microbial community in the gut and the host environment - as a powerful proximate
mechanism influencing host health and fitness among animals (Kinross et al., 2011; Lloyd-

Price et al., 2016; Rosshart et al., 2017). Emerging evidence has highlighted that the gut

microbiota varies substantially within and between individuals (David et al., 2014; Maurice

et al., 2015), and that microbe-host interactions are likely to be taxonomically widespread

and central to many behavioural, ecological and evolutionary processes (Alberdi et al., 2016;

Davidson et al., 2020; Sherwin et al., 2019). However, the role of the gut microbiome in

determining fitness in natural populations, and whether there are critical developmental

windows whereby it affects future growth and survival, remains poorly understood.

Evidence suggests that the microbiome is associated with many aspects of survival and

fitness. In humans, the maternal vaginal microbiota is associated with pre-term birth

(Fettweis et al., 2019) and sudden infant death syndrome (Fettweis et al., 2019; Highet et

al., 2014), and the gut microbiome is associated with longevity (Bian et al., 2017).

Laboratory-bred mice, which have substantially different gut microbiota than their wild

counterparts, were more likely to survive an influenza infection if they had received a gut

microbiota transplant from wild donors (Rosshart et al., 2017). Notably, wild mice had

significantly higher abundances of Bacteroidetes and Proteobacteria, and lower abundances

of Firmicutes, suggesting these taxa may play a role in regulating host health. The potential

for the gut microbiome to impact animal fitness in natural systems has gained much

attention in recent years (Amato, 2013; Suzuki, 2017; Trevelline et al., 2019; West et al.,

2019). However, the empirical evidence to support such hypotheses is lacking, perhaps

owing to the logistical constraints associated with collecting longitudinal data on both the

gut microbiome and indicators of fitness such as survival in these systems.

Research on laboratory model species has shown that the microbial community in the gut

can confer specific health benefits to the host (Kinross et al., 2011), and the timing of gut

colonization is critical for normal development (Cox et al., 2014; Hansen et al., 2012). In

particular, gut microbiota are important for the development and regulation of gut

immunity by maintaining intestinal barrier function (Hooper et al., 2012; O'Hara &

Shanahan, 2006) and aiding digestive capabilities through the metabolism of dietary

components (Rowland et al., 2018). Gut bacteria can induce expression of host genes that,

for example, code for proteins in the mucosal layer of the intestines (O'Hara & Shanahan,

2006). Moreover, bacterial genomes can produce enzymes that are absent in the host, yet

are necessary for specific digestive functions such as cellulose digestion in termites

(*Nasutitermes spp.)* and giant pandas (*Ailuropoda melanoleuca*) (Warnecke et al., 2007; Zhu

et al., 2011) and nitrogen metabolism in birds (Vispo & Karasov, 1997). Germ-free mice

devoid of any microbes have depressed immune function and lower nutrient assimilation

efficiency, which can only be improved by introducing intestinal microbes from normal mice

donors (Hansen et al., 2012; Olszak et al., 2012). In many cases, the timing of microbial

colonization is critical and microbiome restoration must occur within an early

developmental window in order to affect host phenotypes that are observed later in life

(Cox et al., 2014; Hansen et al., 2012). Therefore, variation in host microbiota can have

important consequences for digestive capability and body condition, and act as a

mechanism underlying developmental plasticity (Alberdi et al., 2016). Importantly, effects

may be dependent on the timing of microbial colonization if microbial effects on host traits

may have immediate effects (i.e. contemporary associations), or delayed effects (i.e. time-

lagged associations).

In many taxa, natal body size (Ringsby et al., 1998), natal body mass (Both et al., 1999;

Monrós et al., 2002), or growth rate (Martin, 2015) can be key predictors of juvenile viability

and recruitment into the breeding population. The association between the gut microbiome

and weight has been studied extensively in clinical studies for application in human obesity

(Bäckhed et al., 2004; Turnbaugh et al., 2006), and in agricultural studies to increase animal

productivity (Coates et al., 1981; Ford & Coates, 1971; Torok et al., 2011). While having a

more diverse microbiome may intuitively suggest a healthier gut microbiome (Lozupone et

al., 2012), the effects of microbial community structure are likely more complex (Zaneveld

et al., 2017). In humans, higher diversity of gut microbiota is associated with high body

mass index and high amounts of visceral fat (Beaumont et al., 2016; Le Chatelier et al.,

2013). Specific taxa, such as *Lactobacillus* spp. can influence fat storage and weight change,

however, the effects of such bacteria are strain-specific (Drissi et al., 2017). In poultry,

beneficial bacteria can be selectively enriched (e.g. Patterson et al. (1997)), and the use of

antibiotics reduce gut bacterial load, decrease microbial diversity, and improve feed

conversion ratios (Banerjee et al., 2018). Weight gain associated with antibiotic

administration and decreased bacterial diversity has also been shown in house sparrows

(*Passer domesticus*) (Kohl et al., 2018), *Daphnia magna* (Motiei et al., 2020) and magellanic

penguins (*Spheniscus magellanicus*) (Potti et al., 2002). By contrast, microbial diversity has

been positively associated with body condition and growth rate in wild nestling birds

(Noguera et al., 2018; Teyssier et al., 2018).

The great tit (*Parus major*) provides a powerful natural system for addressing microbiome

related effects on host phenotypes. It is an altricial species that readily breeds in nest boxes,

so that the developmental period can be tracked easily in the wild. We investigated the

relationship between natural variation in the gut microbiome and direct and indirect

measures of fitness in nestling great tits when they were 8 and 15 days old. The nestling

stage is a critical time for gut microbiome colonization (Koenig et al., 2010) and the

microbial-programming of adult phenotypes (Cox et al., 2014; Hansen et al., 2012; Sudo et

al., 2004). Nestling great tits undergo substantial shifts in gut microbiota between day 8 and

day 15 of development (Teyssier et al., 2018), a period when we predicted the microbiome

would have a detectable effect on body weight and the probability of survival to fledging.

We tested whether diversity of the microbiome, or the abundances of specific microbes,

were associated with nestling weight and survival to fledging. Given the mixed reports of

microbial links with host fitness in the existing literature (see above), we had no a priori

predictions regarding the direction of such potential relationships. Nevertheless, we

proposed that if diversity was positively related to measures of host fitness, then a rich

and/or evenly distributed microbial community would be important for healthy

development. By contrast, if the relationship was negative, then the host may benefit from

the presence or absence of specific microbial taxa, which should be observed in differences

in the relative abundance of these specific microbes associated with host fitness. Moreover,

because early developmental windows have been reported to be important for microbial

influences on host traits (Cox et al., 2014; Hansen et al., 2012), we sampled the gut

microbiome from chicks at two developmental stages to test whether microbiome effects

on host fitness traits were contemporary or time-lagged. In other words, we asked whether

the gut microbiome at D8 or D15 was a better predictor of mass and survival.

Methods:

Field monitoring and microbiome sampling

Birds were sampled from nine nest box populations across Co. Cork, Ireland, five of these

were mixed/deciduous habitats and four were coniferous habitats (see O'Shea et al. (2018)).

We collected 204 faecal samples from 150 nestling great tits from 54 nests (see below for

the   number   of   samples   that   were   successfully   sequenced   and   sample   size   per

developmental stage) for 16S rRNA gene sequencing in order to map the gastrointestinal

microbial communities. During April-June 2016, nest boxes were monitored to determine

lay dates, hatching dates and nestling survival. Nestlings aged 8 days (+/- 1 day) (D8) and 15

days (D15) were placed into sterile holding bags inside a heated holding case. Coffee filters

were   used   to   line   the   bags   in   order   to   soak   up   urea   from   the   faeces,   as   urea   has   the

potential to affect downstream sequencing (Khan et al., 1991). Samples were collected from

nestlings that defaecated naturally within 15-20 minutes of being out of the nest before

being returned to the nest. Faecal sacks were opened using a sterile inoculation loop to

release the faecal matter and place in a microcentrifuge tube containing 500uL of 100%

ethanol. Samples were stored at -20^{o}C within 8 hours of collection until DNA extraction. D8

birds were weighed, and uniquely individually identified by clipping the tip of one of their

nails, avoiding the quick (i.e. blood vessel). D15 nestlings were weighed and ringed with a

unique   identifiable   metal   ring   (British   Trust   for   Ornithology)   and   matched   against   their

unique nail clipping. Nestlings for which we had repeated samples (i.e. for both D8 and D15)

are referred to as 'repeat samples'.

DNA Extractions

Briefly, DNA was extracted from the dried faecal contents of all birds using the Qiagen

QIAamp DNA Stool Kit, following the "Isolation of DNA from Stool for Pathogen Detection"

protocol (June 2012 edition), with modifications described in (Shutt et al., 2020) to

accommodate dried avian faeces. DNA was stored at -20^{o}C. Full extraction methods are

described in Supporting Information.

Illumina MiSeq sequencing

Full library preparation details are described in Supporting Information. Briefly, the V3-V4

variable region of the 16S rRNA gene was amplified from the DNA extracts using the 16S

metagenomic sequencing library protocol (Illumina). The DNA was amplified with primers

specific to the V3-V4 region of the 16S rRNA gene which also incorporates the Illumina

overhang adaptor. Samples were sequenced on the MiSeq sequencing platform (Clinical

Microbiomics, Denmark), using a 2 x 300 cycle kit, following standard Illumina sequencing

protocols.

Bioinformatics and statistical analysis

The DADA2 pipeline was used to process the raw sequencing data (Callahan et al., 2016) in R

version 3.5 (R Core Team, 2019). Sequence quality was visually inspected. Sequences were

trimmed to remove adapters and lower quality reads (median quality scores below 25-30

threshold) at the extremities of the sequence and filtered to remove sequences with higher

than expected errors. Read errors were estimated before dereplication. Forward and

reverse reads were merged to construct 'contig' sequences, these were used to construct a

sequence table of Amplicon Sequence Variants (ASV's), which counts the number of times

each unique sequence is detected. The previous steps were performed for each run

separately. Then the separate sequence tables were merged and chimeras removed using

the 'consensus' method. Taxonomy was assigned to each ASV by RDP's Naive Bayes

Classifier (Wang et al., 2007) against the Silva reference database (version 132) (Quast et

al., 2012). This is at 100% sequence identity in contrast to the lower resolution OTU method

which groups sequences at 97% identity. ASV's allow greater sensitivity and specificity,

better discrimination of ecological patterns than OTU's and are reusable across studies

(Callahan et al., 2017).

The DADA2 outputs were assembled into a single Phyloseq object (McMurdie & Holmes,

2013). Sequences identified as mitochondrial or chloroplast were removed. Further filtering

of samples and ASV's took place in R. Sample completeness curves were plotted using vegan

(Jari Oksanen, 2019) and helped determine the lower cut-off for sample reads at 10,000

reads. Low read samples (<10000 reads, 9 samples) were removed leaving 195 (Day-8=81,

Day-15=114, repeat samples=41) samples for the analysis. Alpha diversity (both Shannon

and Chao1 diversity) was calculated using the '*estimate_richness*' function from the

phyloseq package on the filtered dataset. Shannon diversity (Shannon, 1948) measured

richness weighted by abundance (the evenness of a community) and Chao1 (Chao, 1984)

measuresd richness, specifically estimating taxa abundance and rare taxa missed from

under sampling.

Statistical analysis

Nestling growth: Alpha diversity and phylum-level relative abundance

Linear mixed models (LMMs) and generalised linear mixed models (GLMMs) were used to examine the association between measures of nestling growth and two measures of alpha diversity: Shannon's index and Chao1 index. Models were fit using the lme4 package (Bates et al., 2015) on each data subset. Numeric variables were centered and scaled. Model diagnostics were assessed using the DHARMa package (Hartig, 2019). Raw weight (g) was used as the response variable instead of change in weight (delta-weight) as this allowed us to control for D8 weight in time-lagged models, as well as to differentiate between, for example, very light D8 nestlings achieving normal D15 weight and heavy D8 nestlings achieving heavy D15 weight.

We examined whether nestling body weight was influenced by the gut microbiome in three different ways: (1) whether the microbiome at an early developmental stage predicted future host weight gain (henceforth 'time-lagged weight'), (2) whether the microbiome at the time of sampling was associated with weight at the same developmental stage (henceforth 'contemporary weight') at D8, and (3) contemporary weight at D15. Due to collinearity (alpha diversity) and non-independence (Phylum-level relative abundance), we performed separate analyses for four different measures of microbiome as explanatory variables: alpha diversity (Shannon diversity and chao1 index), and phylum-level abundance (Proteobacteria and Firmicutes). These two phyla were selected for our analyses because they were the only two taxa that were prevalent across all birds (except for three birds).

For the time-lagged weight analyses, we ran a linear mixed model (LMM) using the lmer function in R with D15 weight as the response variable, and alpha diversity at D8 as a fixed variable. We included D8 weight as a covariable to control for variation in starting weight. We also included habitat, lay date of first egg and brood size as fixed factors as these may influence nestling weight. Site and nest ID were used as nested random effects, as well as sequence plate as a separate random effect. Only birds that had repeat samples were included in this analysis (n= 41).

For the D8 contemporary weight analyses (all D8 samples, n= 81), we ran a LMM as
described above, with D8 as the response variable and alpha diversity at D8 as a fixed
factor. We included the same fixed effects and random effects as described above. The D15
contemporary analyses (all D15 samples, n = 114) were the same as described for D8, using
D15 data as opposed to D8 data. D15 weight was squared in the contemporary D15 model
as the model did not converge with raw weight as the response.

Nestling survival: Alpha diversity and phylum-level relative abundance
We tested whether microbiome diversity at D8 predicted nestling survival. Here we define
survival as surviving to fledgling (i.e. the nestling was not found dead in the nest between
D8 and fledging). We ran a non-parametric cox proportional hazards survival model using
the Coxme package (Therneau, 2020), and included weight (at D8), lay date and brood size
(at D8) as covariables. We included three ranks of survival times (i.e. died before D15, died
after D15 or fledged) as the ranks of survival times provide additional temporal information
beyond a binary outcome, i.e. died/survived for the model to estimate mortality risk. The
habitat variable was excluded from the survival models as it caused premature model
convergence due to a lack of variation in the variable (i.e. all coniferous birds survived).
Because all but one bird that survived to D15 survived to fledging, we only performed this
analysis for birds that were sampled for gut microbiome at D8. We repeated this analysis
with Proteobacteria abundance as the predictor variable, and again with Firmicutes
abundances as the predictor variables.

Nestling survival: Beta diversity
We investigated whether microbial community structure differed between birds that did or
did not survive to fledgling, using the adonis function (PERMANOVA) from the vegan
package (Jari Oksanen, 2019). Beta diversity at D8 was measured using Bray-Curtis distance
(weighted by ASV abundance) and Jaccard (presence/absence of ASVs). We ran separate
analyses for each beta diversity measure in which D8 weight and survival (Yes/No) were
fixed factors. Adonis fits each term sequentially, therefore we used the model formula
beta-diversity~D8-weight + survival, allowing us to estimate the variation in microbiome
community composition associated with survival status after accounting for body weight, as
weight was associated with survival in our population (see results, Table 2). Permutations

288 were constrained to within nest with the 'strata' option to control for pseudo-replication of

289 samples from the same nest. Adonis results were very similar with and without this strata

290 option.

292 The beta diversity dataset (D8 samples) was filtered by prevalence prior to distance

293 calculation, where taxa with less than 1 copy in 5% of samples were removed following

294 Knowles et al. (2019). This resulted in a dataset containing 1254 ASV's. The relative

295 abundance of each phyla was calculated per sample using 'total sum scaling' (TSS) by

296 dividing each feature count by the total library size (McKnight et al., 2019). Bray Curtis and

297 Jaccard distances were calculated from our TSS normalized data. An assumption of the

298 adonis test is that groups have homogeneity of variance. The dispersion of the groups was

299 checked for homogeneity of variance using the Betadisper and permutest functions from

300 the vegan package.

302 ***Indicator ASVs for nestling survival***

303 We carried out indicator species analysis to identify which ASVs were representative of birds

304 that were alive at D8 (n=81) and survived to fledgling (n=64), or did not survive to fledgling

305 (n=17). Indicator analysis identified "indicator species" if they reflected the state of the

306 environment (i.e. likelihood the bird will survive, or not survive). We used the multipatt

307 function from the IndicSpecies package (M. De Caceres, Legendre, P., 2009), which assigns a

308 test statistic, or "indicator value" to each ASV for each group, in our case two groups:

309 Survival-Yes, Survival-No. Taxa with an indicator value of > 0.4 and p <= 0.05 were

310 considered indicator species. We also report the "specificity value" and the "fidelity value".

311 The former is the estimate of the probability that an ASV is detected in a particular group.

312 For example, a value of 1.0 would indicate that the ASV is only detected in that group and

313 not another group. The latter is the probability estimate of finding the species in sites (i.e.

314 birds) belonging to the group. For example, a value of 1.0 would indicate that the ASV is

315 recorded in all birds within that group (De Caceres, 2020). We did not adjust for multiple

316 comparisons for three reasons. First, our analysis did not include any overlap, and thus

317 multiple analyses, across different combinations of groups (De Caceres et al., 2009). Second,

318 the purpose of this analysis was exploratory in order to generate, rather than confirm

319 hypotheses (Bender & Lange, 2001; Ranstam, 2019). As such, we report ASVs as potential

indicator species, as opposed to reporting a quantitative, total number of taxa associated

with survival/non-survival (De Cáceres et al., 2010). Third, this analysis served to generate a

posteriori predictions regarding the relationship between indicator taxa and weight gain

(see below), although we note that independent studies will be necessary to confirm these

exploratory analyses (Ranstam et al 2019). We generated a heatmap of indicator species for

survival and non-survival using the heatmap function from the phyloseq package (McMurdie

& Holmes, 2013).

A posteriori analysis of indicator species

Indicator species from our analysis may reflect whether a bird survives or not. We

hypothesized that indicator species may reflect host survival because such indicator species

may also predict weight gain. We performed a posteriori LMMs with D15 weight as the

response variable and D8 relative abundance (of the indicator species) as a fixed variable.

D8 weight, habitat, brood size and lay date were also included as fixed variables. Site and

nest were included as nested random terms (i.e. 1|site/nest). We restricted this a posteriori

analysis to one indicator species that had the highest test statistic from each group

(Survival-Yes: Lactobacillus (ASV46) and Survival-No: Rahnella (ASV38)). Because there were

a high number of zeros in relative abundance of these taxa, and because our data were

proportions and therefore could not be fit using a zero-inflated (count data) model, we

included only birds where the taxa were present and the birds with 0 abundance for that

taxa were excluded (ASV46: n=21/41; ASV38: n=8/41).

All graphs were created using using ggplot2 (Wickham, 2016), unless otherwise indicated.

Results:

Sequencing results

There were 18,537,795 high quality reads, clustered into 49,655 ASV's before removing

control samples, mitochondria and chloroplast sequences. Mean reads per sample was

88,698 and individual samples ranged from 61 to 579,864 reads. Samples with less than

10,000 reads were discarded before further analysis (n=9). There were 15,183,675 total

high-quality reads, clustered into 47,400 ASV's after removing control samples as well as

mitochondria and chloroplast sequences. Samples ranged from 10,220 to 557,336 reads and

mean reads per sample was 77,865.

The most prominent phyla across all samples (D8 and D15) were as follows (mean

percentage relative abundance ±SE): Firmicutes (43.0%±2.5); Proteobacteria (35.2%±2.2);

Tenericutes (9.7%±1.8); Bacteroidetes (4.2%±0.4) and Actinobacteria (2.2%±0.2). Relative

abundance of the major phyla at D8 (at least >=10% relative abundance in at least 1 sample)

per sample are plotted in Figure 1.

Figure 1. Relative abundance of the 8 most prominent phyla (>=10% relative abundance). Each vertical bar corresponds to a nestling at Day 8. "Other" group represents all low abundance phyla as white space.

Nestling growth/weight

The time-lagged analysis indicated that weight at D15 was negatively related to alpha

diversity at D8 (Figure 2a, Table 1), controlling for weight at D8 (Table 1). Neither habitat,

lay date or brood size were significant in this analysis (Table 1). Cross-sectional analyses

indicated no relationship between contemporary weight and diversity at either D8 or D15

(Figure 2b and 2c, Table S1). Brood size was negatively related to weight at D8. Habitat had

a non-significant effect on weight in D15 birds where deciduous birds had lower weight than

coniferous birds. The relative abundances of Proteobacteria and Firmicutes did not

significantly predict weight in any of these analyses.

Figure 2. The effect of Shannon diversity on (a) time-lagged weight (i.e. Day 15 (D15)), (b) contemporary weight Day 8 (D8) and (c) contemporary weight D15. Black dots are individual data points, blue line is regression line with 95% CI (shaded blue).

Table 1. GLMM output from time-lagged (Day-15) weight analyses. The effect of alpha diversity is reported for (a) Shannon diversity and (b) Chao1 diversity. * p < 0.05.

Dependent~Independent variable	Estimate	Std. Error	df	Test statistic	P_estimate
(a) Nestling weight (time-lagged)					
(Intercept)	16.798	0.525	21.052	31.977	<0.001 *
Shannon (Day-8)	-0.593	0.167	23.749	-3.544	0.002 *
Habitat (deciduous)	0.247	0.630	22.081	0.392	0.699

Weight (Day-8)	0.700	0.173	24.672	4.035	<0.001 *
Lay Date (first egg)	0.252	0.296	21.226	0.851	0.404
Brood size (Day-8)	0.168	0.273	21.991	0.616	0.544
(b) Nestling weight (time-lagged)					
(Intercept)	16.961	0.475	20.856	35.681	<0.001 *
Chao1 (Day-8)	-0.525	0.184	34.314	-2.855	0.007 *
Habitat (deciduous)	0.011	0.572	21.903	0.019	0.985
Weight (Day-8)	0.528	0.192	32.990	2.756	0.009 *
Lay Date (first egg)	0.257	0.270	21.399	0.949	0.353
Brood size (Day-8)	0.121	0.250	22.841	0.483	0.634

Nestling Survival

Survival was significantly related to weight at D8; the hazard ratio of 0.67 indicated that for

every unit increase in weight, mortality risk decreased by a factor of 0.67. Survival was not

influenced by lay date or brood size. Nestling survival to fledging was not predicted by

Shannon's diversity or Chao1 index at D8 (Table 2). The relative abundance of

Proteobacteria and Firmicutes at D8 also did not predict survival to fledging.

Tests to verify the assumption of homogeneity of dispersion for survived/not-survived

groups were marginally non-significant for Jaccard distance (Sum Squares= 0.009, F=3.295,

p=0.081) and non-significant for Bray-Curtis (Sum squares=0.006, F=1.973, p=0.164). Beta

diversity did not differ between surviving and non-surviving nestlings for Jaccard (Sum

Squares=0.563, F=1.233, p=0.172) or Bray-Curtis (Sum Squares=0.523, F=1.193, p=0.228),

see Supporting Information Table S3.

Table 2. Determinants of mortality risk from Cox proportional hazards model. Positive

coefficients indicate increased risk of mortality. HR = Hazard ratio, which is the

exponentiated coefficients, and gives the ratio of the total number of observed to expected events. Note: all numeric covariates are centered and scaled. * p < 0.05.

Dependent~Independent variable	Coef	HR	se	Test statistic (z)	P_estimate
Nestling mortality risk					
Shannon	0.021	1.021	0.327	0.07	0.95
Weight D8	-0.665	0.514	0.282	-2.36	0.018*
Lay Date (First egg)	0.304	1.356	0.382	0.80	0.43
Brood Size (Day-8)	0.551	1.735	0.363	1.52	0.13
Nestling survival					
Chao1	0.314	1.368	0.283	1.11	0.27
Weight D8	-0.609	0.544	0.289	-2.11	0.035*
Lay Date (First egg)	0.297	1.346	0.396	0.75	0.45
Brood Size (Day-8)	0.634	1.885	0.386	1.64	0.1

Indicator species analysis

The indicator species with the highest indicator values associated with survival belonged to
ASV46, ASV310 and ASV150 which all belong to the *Lactobacillaceae* family (Figure 3, Table
S4). The taxa with the highest indicator values associated with non-survival belonged to the
ASV38, ASV1034, ASV895 which belong to *Enterobacteriaceae*, *Methylopilaceae* and
Acidothermaceae families, respectively. The taxa indicative of non-survival and survival (p
value <=0.05 and test statistic >=0.4) were plotted on a heatmap (Figure 3). Overall,
specificity values were high (mean = 0.94±0.008 standard error) and fidelity values were low
(0.21±0.008), suggesting that indicator species were typically only detected within a given
group (i.e. survive/non-survive), but that not all birds within a group were recorded to have

a given indicator species. All indicator species and test statistics (specificity values and
indicator values) are reported in Supporting Information Table S4.
402
403   A posteriori analyses showed that a higher relative abundance of ASV46 (a *Lactobacillus*
404   spp., and the taxa most strongly associated with survival) at D8 predicted a higher weight at
405   D15. Relative abundance of ASV38 (a *Rahnella* sp., and the taxa most strongly associated
406   with non-survival) was not associated with weight at D15 (Table 3).
407

Figure 3. Heatmap of relative abundance of indicator species ASV's for survival. Individual birds are columns with separate panels for individuals that either died or survived. ASVs are in rows where taxa in brackets refer to the ASV's family-level taxonomic assignment, or NA if not resolved to family-level. Taxa associated with survival as opposed to non-survival are in bold. Lighter colours indicate greater abundance.

Table 3. A posteriori GLMM output from time-lagged weight analyses. The effect of indicator species on weight at D15 are reported for a) *Lactobacillus* sp. (ASV46) and b) *Rahnella* sp. (ASV38). * p <0.05

Dependent/Independent variable	Estimate	Std. Error	df	Test statistic	P_estimate
(a) D15 Weight (time-lagged)					
(Intercept)	17.673	0.739	4.679	23.906	<0.001 *
Abundance ***Lactobacillus sp.* (ASV46)**	1.042	0.367	8.601	2.839	0.02 *
Habitat (deciduous)	-1.946	1.007	5.126	-1.933	0.11
Weight (Day-8)	0.835	0.340	11.184	2.460	0.031 *
Brood Size (Day-15)	-0.459	0.352	11.573	-1.303	0.218
Lay Date (First egg)	-0.457	0.389	7.245	-1.175	0.277
(b) D15 Weight (time-lagged)					
(Intercept)	16.473	0.656	2.000	25.111	0.002 *
Abundance ***Rahnella sp.* (ASV38)**	0.040	0.388	2.000	0.103	0.927
Habitat (deciduous)	2.908	1.320	2.000	2.203	0.158
Weight (Day-8)	5.398	1.579	2.000	3.419	0.076
Brood Size (Day-15)	1.690	0.660	2.000	2.558	0.125
Lay Date (First Egg)	-1.069	1.113	2.000	-0.960	0.438

409
410
Discussion:

Overall summary

We demonstrate that the gut microbiota predicts nestling fitness in a population of wild

passerine birds. Neither microbial alpha diversity nor phylum-level abundance predicted

contemporary weight, but alpha diversity did predict future weight gain. Moreover, eight-

day old nestlings with a greater relative abundance of specific *Lactobacillus* taxa were more

likely to fledge, and for at least one of these taxa, greater relative abundance was

specifically associated with increased host weight. Notably, although we show clear links
between gut microbiota and and survival, and a proxy for survival (i.e. weight), the direction
of these relationships or the microbial metrics that predict such relationships are variable
within our own study (Figure 3) and contrast with some, but not other studies. A lack of
consensus on what aspects of microbial taxonomic composition predict host traits is
prominent in the microbiome literature (Zaneveld et al., 2017). We discuss below how
critical developmental windows, environmental matching and the Anna Karenina
Hypothesis may explain such differential findings across studies.

We show a negative relationship between alpha diversity and future weight gain.
Experimentally reducing diversity of the microbiome through antibiotic administration has
been shown to improve host growth (Banerjee et al., 2018; Kohl et al., 2018; Lan et al.,
2005; Potti et al., 2002), possibly because antibiotics reduce bacterial load, diverting
resources towards growth, as opposed to immune function, in a classic life-history trade-off
(Rauw, 2012). Our study demonstrates that this negative relationship is present in the
absence of microbiome manipulation, although we can only speculate as to the mechanisms
responsible for limiting growth in nestlings with high gut microbial diversity. For example, a
conventional gut microbiome compared to a germ-free one lowers absorption capacity of
glucose and B-vitamins by increasing the thickness of the gut epithelium (Ford & Coates,
1971), and non-mutualistic microbes may create greater competition for nutrients
(Wasielewski et al., 2016). While the diversity of the microbial community may be important
for regulating host growth, some taxa may have disproportionate effects. At least one ASV,
Lactobacillus sp., with the strongest potential association with survival, was positively
associated with host growth. Several *Lactobacillus* spp. have beneficial, probiotic effects on
the host (Angelakis & Raoult, 2010; Banerjee et al., 2018; Gomes et al., 2012; Vásquez et al.,
2012), though this is certainly not the case for all *Lactobacillus* spp. Future studies should
look to shotgun sequencing to identify specific bacterial strains and genes for functional
profiling and for developing probiotic treatments. This could be followed up by manipulative
studies to identify the causal relationships between ASV and host traits, such as isolating
and culturing ASV46 to test its potential probiotic effects when administered to nestlings
(Davidson et al., 2020; Gatesoupe, 1999; Holzapfel & Schillinger, 2002).

Although a growing number of studies have demonstrated that the microbiome is an
important predictor of host weight, the microbial profiles and the direction of this
relationship vary across species and studies (Kohl et al., 2018; Potti et al., 2002; Teyssier et
al., 2018; Videvall et al., 2019). For example, a negative relationship between alpha diversity
and future weight gain was found in juvenile ostriches (Videvall et al., 2019), whereas a
positive relationship between alpha diversity and contemporary body condition was
reported in a different population of great tit nestlings (Teyssier et al., 2018). The
contrasting results reported in these studies and our own may be due to variations in the
microbiome sampling method and storage solutions (i.e. cloacal versus faecal, see Videvall
et al. (2017); storing solutions see Vargas-Pellicer et al. (2019)). Nevertheless, it is more
likely that differences in local environments (Song et al., 2020) and the temporal resolution
of microbial and host traits dictate when and how microbiota impacts on host phenotypes
in birds (Videvall et al., 2019).

Time-lagged associations between the gut microbiota and host traits are expected to be
especially important during developmental periods when large shifts in the establishing
microbial community coincide with the programming of host biology (Cox et al., 2014;
Hansen et al., 2012; Sudo et al., 2004), and in systems that are likely to experience abrupt
changes in environmental inputs associated with gut microbiome, such as diet. The latter
may be especially relevant in birds as their microbiomes are tightly linked with
environmental variation, in contrast to mammals where the microbiome is primarily linked
to phylogeny (Song et al., 2020). In ostriches, microbiome diversity predicted future weight
gain, but the direction of this relationship switched after week 1 post-hatching, a time when
ostriches no longer rely on an internal yolk as sustenance (Videvall et al., 2019). In the
current study, alpha diversity at D8 predicted future weight at D15, whereas cross-sectional
analyses were not sufficient to detect such a relationship. This time-lagged association may
be owing to a critical developmental period in which the microbiome impacts on future
phenotypes, a phenomenon that is prominent in clinical microbiome literature (Cox et al.,
2014; Hansen et al., 2012; Sudo et al., 2004). The gut microbiome may also act as a
mechanism for adaptive phenotypic change through its sensitivity to environmental
variability, and its connection to host biological processes (Alberdi et al., 2016). Such a
mechanism would also be in line with the 'Environmental Matching Hypothesis', which

posits that conditions during development cause phenotypic changes that serve to match
individuals to their environments (Crino & Breuner, 2015; Weber et al., 2018). For example,
unpredictable food supply during the development stage can influence adult physiological
and behavioral phenotypes (Spencer et al., 2009; Zimmer et al., 2013). Therefore,
microbiome-mediated developmental plasticity may be adaptive in species where offspring
benefit from a gut microbiome that primes the host for dietary shifts. Dietary shifts are
common in tits where the relative frequency of insect taxa can change throughout the
provisioning period (Arnold et al., 2007; García-Navas et al., 2012). Although Belgian great
tit nestlings had the greatest improvement in body condition if they maintained a stable
microbial diversity throughout development (Teyssier et al., 2018), perhaps this may be
because the environmental conditions at the time were also stable. In our study, it is
difficult to differentiate between whether the gut microbiome is responding to dietary shifts
as opposed to priming the host. Pinpointing if and when early life bacterial diversity
programs host digestive and/or metabolic efficiency in wild animals, and the extent to
which environmental variation, such as food availability and diet, impacts on the developing
microbiome will progress our understanding of microbial-mediated developmental
mechanisms.

The Anna Karenina Hypothesis states that changes in an unhealthy or so-called "dysbiotic"
microbiome can be stochastic rather than deterministic, making them difficult to detect
statistically (Zaneveld et al., 2017). In other words, unhealthy individuals can have greater
variation in their microbial community composition than healthy individuals, and rather
than displaying a specific 'unhealthy' state, the microbiomes of diseased or stressed
individuals can all look very different due to a breakdown in regulation. This phenomenon
may therefore explain why nestling survival was not associated with either alpha or beta
diversity. Moreover, although several indicator species were identified to be associated with
birds that did or did not survive, many of these had low prevalence across nestlings. This is
reflected in the high specificity values and low fidelity values, suggesting that although the
presence of an indicator species in a nestling's gut microbiome reflects a high probability of
survival or non-survival, the same indicator species is not present across all non-surviving
nestlings in our sample population. This distinction is important in, for example,
conservation practices whereby microbes are used as markers for population viability

(Trevelline et al., 2019; West et al., 2019). Our analysis suggests that in the case of non-
survival, there are no microbes that are universally indicative of non-survival across birds,
but rather the presence of one of several candidate microbes predicts a higher probability
of mortality, which is consistent with the Anna Karenina Hypothesis. Whether these same
taxa affect survival in other populations and/or other species of birds is unknown; thus
these candidate taxa should be examined in independent confirmatory studies. For
example, ASVs reported for non-survival may be pathogenic to great tits, although we found
no evidence in the literature to support this. Nevertheless, ASV1320, ASV2151, ASV3604
and ASV942 are members of the Rhizobiales order which contains genera that are well
known human pathogens, and ASV5768 and ASV4855 are members of the Chlamydiales
order which also contain pathogenic genera, though these genera were not detected in our
study.

We also encourage further investigations into whether indicator taxa are the cause or
consequence of host growth, or indeed any trait that may be associated with survival, such
as glucocorticoids (Weber et al., 2018) and immune function (Hõrak et al., 1999).
Manipulation of the gut microbiome alongside metabolomics or metagenomics will be the
next step in providing insight into the causal direction and functional mechanisms of
microbial variation and its relationship with host fitness. The latter is particularly important
for identifying whether there is functional overlap between microbial taxa and the
mechanisms by which these taxa affect host traits (Heintz-Buschart & Wilmes, 2018). Given
the temporal importance of host-microbial relationships identified in this study and
elsewhere, we anticipate the maturing microbiome will have important consequences on
long-term fitness effects. More generally, our results highlight the importance of the gut
microbiome in evolutionary processes as it has the potential to mediate rapid adaptation to
environmental change and may also support conservation efforts as an indicator of
individual and population health.

Figure 3. Associations between microbiome metrics and nestling fitness traits. Solid arrows

represent positive associations and open arrows represent negative associations. Dashed

lines indicate no relationship. For visualization purposes, some lines terminate between two

host fitness boxes to indicate that both traits had the same (significant/non-significant)

result. Note that the direction of arrows does not suggest causality from our analyses.

Funding.

This work was supported by Science Foundation Ireland by way of funding to APC

Microbiome Ireland, Cork, Ireland. GLD, SES, MRC and JLQ etc were supported by funding

from the European Research Council under the European Union's Horizon 2020 Programme

(FP7/2007-2013)/ERC Consolidator Grant "Evoecocog" Project No. 617509, awarded to JLQ.

GLD is funded by a Leverhulme Trust Early Career Fellowship (ECF-2018-700) with matched

funding from the Sir Isaac Newton Trust.

Acknowledgements.

Thanks to Jodie Crane and Iván de la Hera, for assistance with fieldwork. Paul Cotter, Fiona

Crispie and Laura Finnegan from the Teagasc Sequencing facility for their role in relation to

the 16S rRNA sequencing. For advice on bioinformatics, we thank Fiona Fouhy, Orla

O'Sullivan and Calum Walsh. We are grateful to James Nichols, Ally Phillimore and Sarah

Knowles for helpful discussions on molecular methodology.

Data accessibility.

Our data will be deposited at Dryad, and sequence data uploaded to a repository such as

NCBI Sequence Read Archive, if the manuscript is published.

Research and Animal Ethics: This study was conducted under licences from the Health

Products Regulatory Authority (AE19130_P017), The National Parks and Wildlife Services

(010/2016) and the British Trust for Ornithology, and permission from Coillte Forestry and

private landowners. The research project received ethical approval from the Animal Welfare

Body at University College Cork, and was in accordance with the ASAB (Association for the

Study of Animal Behaviour) Guidelines for the Treatment of Animals in Behavioural

Research and Teaching.

Author contributions: GLD, CS, RPR and JLQ conceived the ideas and designed the

methodology. GLD and MSR collected the field data. GLD, NW and CNJ carried out the DNA

extraction and library prep. SES analysed the data. GLD and SES wrote the manuscript with

input from all authors. All authors gave final approval for publication.

References:

Alberdi, A., Aizpurua, O., Bohmann, K., Zepeda-Mendoza, M. L., & Gilbert, M. T. P. (2016).
583   Do Vertebrate Gut Metagenomes Confer Rapid Ecological Adaptation? *Trends in*
584   *Ecology & Evolution, 31*(9), 689-699. doi:https://doi.org/10.1016/j.tree.2016.06.008
585   Amato, K. R. (2013). Co-evolution in context: The importance of studying gut microbiomes in
586   wild animals. *Microbiome Science and Medicine, 1*(1). Retrieved from
587   http://dx.doi.org/10.2478/micsm-2013-0002
588   Angelakis, E., & Raoult, D. (2010). The increase of Lactobacillus species in the gut flora of
589   newborn broiler chicks and ducks is associated with weight gain. *PLOS ONE, 5*.
590   Arnold, K. E., Ramsay, S. L., Donaldson, C., & Adam, A. (2007). Parental prey selection affects
591   risk-taking behaviour and spatial learning in avian offspring. *Proceedings of the Royal*
592   *Society B: Biological Sciences, 274*(1625), 2563-2569.
593   doi:doi:10.1098/rspb.2007.0687
594   Bäckhed, F., Ding, H., Wang, T., Hooper, L. V., Koh, G. Y., Nagy, A., . . . Gordon, J. I. (2004).
595   The gut microbiota as an environmental factor that regulates fat storage.
596   *Proceedings of the National Academy of Sciences of the United States of America,*
597   *101*(44), 15718-15723. doi:10.1073/pnas.0407076101

Banerjee, S., Sar, A., Misra, A., Pal, S., Chakraborty, A., & Dam, B. (2018). Increased
productivity in poultry birds by sub-lethal dose of antibiotics is arbitrated by
selective enrichment of gut microbiota, particularly short-chain fatty acid producers.
Microbiology, 164(2), 142-153. doi:https://doi.org/10.1099/mic.0.000597
Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software, 67*(1), 1-48.
Beaumont, M., Goodrich, J. K., Jackson, M. A., Yet, I., Davenport, E. R., Vieira-Silva, S., . . .
Bell, J. T. (2016). Heritable components of the human fecal microbiome are
associated with visceral fat. *Genome Biology, 17*(1), 189. doi:10.1186/s13059-016-
1052-7
Bender, R., & Lange, S. (2001). Adjusting for multiple testing—when and how?
Journal of Clinical Epidemiology, 54(4), 343-349. doi:10.1016/S0895-4356(00)00314-
0
Bian, G., Gloor, G. B., Gong, A., Jia, C., Zhang, W., Hu, J., . . . Li, J. (2017). The Gut Microbiota
of Healthy Aged Chinese Is Similar to That of the Healthy Young. *mSphere, 2*(5).
Retrieved from https://msphere.asm.org/content/2/5/e00327-17
Blomberg, E. J., Sedinger, J. S., Gibson, D., Coates, P. S., & Casazza, M. L. (2014). Carryover
effects and climatic conditions influence the postfledging survival of greater sage-
grouse. *Ecology and Evolution, 4*(23), 4488-4499. doi:10.1002/ece3.1139
Both, C., Visser, M. E., & Verboven, N. (1999). Density–dependent recruitment rates in great
tits: the importance of being heavier. *Proceedings of the Royal Society of London.*
Series B: Biological Sciences, 266(1418), 465-469. doi:10.1098/rspb.1999.0660
Callahan, B. J., McMurdie, P. J., & Holmes, S. P. (2017). Exact sequence variants should
replace operational taxonomic units in marker-gene data analysis. *The ISME Journal,*
11(12), 2639-2643. doi:10.1038/ismej.2017.119
Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.
(2016). DADA2: High-resolution sample inference from Illumina amplicon data.
Nature Methods, 13, 581-581 EP. Retrieved from
https://doi.org/10.1038/nmeth.3869
Chao, A. (1984). Nonparametric Estimation of the Number of Classes in a Population.
Scandinavian Journal of Statistics, 11(4), 265-270. Retrieved from
www.jstor.org/stable/4615964
Coates, M. E., Cole, C. B., Fuller, R., Houghton, S. B., & Yokota, H. (1981). The gut microflora
and the uptake of glucose from the small intestine of the chick. *British Poultry*
Science, 22(3), 289-294. Retrieved from
http://dx.doi.org/10.1080/00071688108447888
Cox, L. M., Yamanishi, S., Sohn, J., Alekseyenko, A. V., Leung, J. M., Cho, I., . . . Blaser, M. J.
(2014). Altering the intestinal microbiota during a critical developmental window has
lasting metabolic consequences. *Cell, 158*(4), 705-721.
doi:10.1016/j.cell.2014.05.052
Crino, O. L., & Breuner, C. W. (2015). Developmental stress: evidence for positive
phenotypic and fitness effects in birds. *Journal of Ornithology, 156*(1), 389-398.
doi:10.1007/s10336-015-1236-z
David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., . . .
Turnbaugh, P. J. (2014). Diet rapidly and reproducibly alters the human gut
microbiome. *Nature, 505*(7484), 559-563. doi:10.1038/nature12820

Davidson, G. L., Raulo, A., & Knowles, S. C. L. (2020). Identifying Microbiome-Mediated
Behaviour in Wild Vertebrates. *Trends in Ecology & Evolution*.
doi:https://doi.org/10.1016/j.tree.2020.06.014
De Caceres, M. (2020). How to use the indicspecies package (ver. 1.7.8).
De Cáceres, M., Legendre, P., & Moretti, M. (2010). Improving indicator species analysis by
combining groups of sites. *Oikos, 119*(10), 1674-1684. doi:10.1111/j.1600-
0706.2010.18334.x
Drissi, F., Raoult, D., & Merhej, V. (2017). Metabolic role of lactobacilli in weight
modification in humans and animals. *Microbial Pathogenesis, 106*, 182-194.
doi:https://doi.org/10.1016/j.micpath.2016.03.006
Fettweis, J. M., Serrano, M. G., Brooks, J. P., Edwards, D. J., Girerd, P. H., Parikh, H. I., . . .
Buck, G. A. (2019). The vaginal microbiome and preterm birth. *Nature Medicine,
25*(6), 1012-1021. doi:10.1038/s41591-019-0450-2
Ford, D. J., & Coates, M. E. (1971). Absorption of glucose and vitamins of the B complex by
germ-free and conventional chicks. *Proc Nutr Soc, 30*(1), 10a-11a.
García-Navas, V., Ferrer, E. S., & Sanz, J. J. (2012). Prey selectivity and parental feeding rates
of Blue Tits Cyanistes caeruleus in relation to nestling age. *Bird Study, 59*(2), 236-242.
doi:10.1080/00063657.2012.662939
Gatesoupe, F. J. (1999). The use of probiotics in aquaculture. *Aquaculture, 180*(1), 147-165.
doi:https://doi.org/10.1016/S0044-8486(99)00187-8
Gomes, B. C., Rodrigues, M. R., Winkelströter, L. K., Nomizo, A., & de Martinis, E. C. (2012).
In vitro evaluation of the probiotic potential of bacteriocin producer Lactobacillus
sakei 1. *J Food Prot, 75*(6), 1083-1089. doi:10.4315/0362-028x.Jfp-11-523
Götmark, F. (2002). Predation by sparrowhawks favours early breeding and small broods in
great tits. *Oecologia, 130*(1), 25-32. doi:10.1007/s004420100769
Hansen, C. H. F., Nielsen, D. S., Kverka, M., Zakostelska, Z., Klimesova, K., Hudcovic, T., . . .
Hansen, A. K. (2012). Patterns of Early Gut Colonization Shape Future Immune
Responses of the Host. *PLOS ONE, 7*(3), e34043. doi:10.1371/journal.pone.0034043
Hartig, F. (2019). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. In.
Heintz-Buschart, A., & Wilmes, P. (2018). Human Gut Microbiome: Function Matters. *Trends
in Microbiology, 26*(7), 563-574. doi:https://doi.org/10.1016/j.tim.2017.11.002
Highet, A. R., Berry, A. M., Bettelheim, K. A., & Goldwater, P. N. (2014). Gut microbiome in
sudden infant death syndrome (SIDS) differs from that in healthy comparison babies
and offers an explanation for the risk factor of prone position. *Int J Med Microbiol,
304*(5-6), 735-741. doi:10.1016/j.ijmm.2014.05.007
Holzapfel, W. H., & Schillinger, U. (2002). Introduction to pre- and probiotics. *Food Research
International, 35*(2), 109-116. doi:https://doi.org/10.1016/S0963-9969(01)00171-5
Hooper, L. V., Littman, D. R., & Macpherson, A. J. (2012). Interactions between the
microbiota and the immune system. *Science, 336*(6086), 1268-1273.
doi:10.1126/science.1223490
Hõrak, P., Lea, T., Ots, I., & Møller, A. P. (1999). Immune Function and Survival of Great Tit
Nestlings in Relation to Growth Conditions. *Oecologia, 121*(3), 316-322. Retrieved
from www.jstor.org/stable/4222472
Jari Oksanen, F. G. B., Michael Friendly, Roeland Kindt, Pierre Legendre, Dan McGlinn, Peter
R. Minchin, R. B. O'Hara, Gavin L. Simpson, Peter Solymos, M. Henry H. Stevens,

Eduard Szoecs and Helene Wagner (2019). vegan: Community Ecology Package
(Version R package version 2.5-6).
Keller, L. F., & Van Noordwijk, A. J. (1994). Effects of Local Environmental-Conditions on
Nestling Growth in the Great Tit Parus-Major. *Ardea, 82*(2), 349-362.
Khan, G., Kangro, H. O., Coates, P. J., & Heath, R. B. (1991). Inhibitory effects of urine on the
polymerase chain reaction for cytomegalovirus DNA. *J Clin Pathol, 44*(5), 360-365.
doi:10.1136/jcp.44.5.360
Kinross, J. M., Darzi, A. W., & Nicholson, J. K. (2011). Gut microbiome-host interactions in
health and disease. *Genome Med, 3*(3), 14. doi:10.1186/gm228
Knowles, S. C. L., Eccles, R. M., & Baltrūnaitė, L. (2019). Species identity dominates over
environment in shaping the microbiota of small mammals. *Ecology Letters, 22*(5),
826-837. doi:10.1111/ele.13240
Koenig, J. E., Spor, A., Scalfone, N., Fricker, A. D., Stombaugh, J., Knight, R., . . . Ley, R. E.
(2010). Succession of microbial consortia in the developing infant gut microbiome.
Proceedings of the National Academy of Sciences, 108(Supplement_1), 4578-4585.
Retrieved from http://dx.doi.org/10.1073/pnas.1000081107
Kohl, K. D., Brun, A., Bordenstein, S. R., Caviedes-Vidal, E., & Karasov, W. H. (2018). Gut
microbes limit growth in house sparrow nestlings (Passer domesticus) but not
through limitations in digestive capacity. (1749-4877 (Electronic)).
Lan, Y., Verstegen, M. W. A., Tamminga, S., & Williams, B. A. (2005). The role of the
commensal gut microbial community in broiler chickens. *World's Poultry Science
Journal, 61*(01), 95-104. Retrieved from http://dx.doi.org/10.1079/WPS200445
Le Chatelier, E., Nielsen, T., Qin, J., Prifti, E., Hildebrand, F., Falony, G., . . . Meta, H. I. T. c.
(2013). Richness of human gut microbiome correlates with metabolic markers.
Nature, 500(7464), 541-546. doi:10.1038/nature12506
Lloyd-Price, J., Abu-Ali, G., & Huttenhower, C. (2016). The healthy human microbiome.
Genome Medicine, 8(1), 51. doi:10.1186/s13073-016-0307-y
Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K., & Knight, R. (2012). Diversity,
stability and resilience of the human gut microbiota. *Nature, 489*(7415), 220-230.
doi:10.1038/nature11550
Martin, T. E. (2015). Age-related mortality explains life history strategies of tropical and
temperate songbirds. *Science, 349*(6251), 966. doi:10.1126/science.aad1173
Maurice, C. F., Cl Knowles, S., Ladau, J., Pollard, K. S., Fenton, A., Pedersen, A. B., &
Turnbaugh, P. J. (2015). Marked seasonal variation in the wild mouse gut microbiota.
The ISME Journal, 9(11), 2423-2434. Retrieved from
http://dx.doi.org/10.1038/ismej.2015.53
McKnight, D. T., Huerlimann, R., Bower, D. S., Schwarzkopf, L., Alford, R. A., & Zenger, K. R.
(2019). Methods for normalizing microbiome data: An ecological perspective.
Methods in Ecology and Evolution, 10(3), 389-400. Retrieved from
https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.13115
McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLOS ONE, 8*(4), e61217-e61217.
Retrieved from http://dx.doi.org/10.1371/journal.pone.0061217
Merila, J. (1996). Genetic Variation in Offspring Condition: An Experiment. *Functional
Ecology, 10*(4), 465-474. doi:10.2307/2389939

Monrós, J. S., Belda, E. J., & Barba, E. (2002). Post-fledging survival of individual great tits:
the effect of hatching date and fledging mass. *Oikos, 99*(3), 481-488.
doi:10.1034/j.1600-0706.2002.11909.x
Motiei, A., Brindefalk, B., Ogonowski, M., El-Shehawy, R., Pastuszek, P., Ek, K., . . .
Gorokhova, E. (2020). Disparate effects of antibiotic-induced microbiome change
and enhanced fitness in Daphnia magna. *PLOS ONE, 15*(1), e0214833.
doi:10.1371/journal.pone.0214833
Noguera, J. C., Aira, M., Pérez-Losada, M., Domínguez, J., & Velando, A. (2018).
Glucocorticoids modulate gastrointestinal microbiome in a wild bird. *Royal Society*
open science, 5(4), 171743-171743. doi:10.1098/rsos.171743
Norris, K. (1993). Heritable variation in a plumage indicator of viability in male great tits
Parus major. *Nature, 362*(6420), 537-539. doi:10.1038/362537a0
O'Hara, A. M., & Shanahan, F. (2006). The gut flora as a forgotten organ. *EMBO reports, 7*(7),
688-693. doi:10.1038/sj.embor.7400731
O'Shea, W., O'Halloran, J., & Quinn, J. L. (2018). Breeding phenology, provisioning
behaviour, and unusual patterns of life history variation across an anthropogenic
heterogeneous landscape. *Oecologia*.
Olszak, T., An, D., Zeissig, S., Vera, M. P., Richter, J., Franke, A., . . . Blumberg, R. S. (2012).
Microbial exposure during early life has persistent effects on natural killer T cell
function. *Science (New York, N.Y.), 336*(6080), 489-493.
doi:10.1126/science.1219328
Patrick, S. C., & Weimerskirch, H. (2014). Personality, foraging and fitness consequences in a
long lived seabird. *PLOS ONE, 9*(2), e87269-e87269.
doi:10.1371/journal.pone.0087269
Patterson, J. A., Orban, J. I., Sutton, A. L., & Richards, G. N. (1997). Selective enrichment of
bifidobacteria in the intestinal tract of broilers by thermally produced kestoses and
effect on broiler performance. *Poultry Science, 76*(3), 497-500.
doi:https://doi.org/10.1093/ps/76.3.497
Pickett, S. R. A., Weber, S. B., McGraw, K. J., Norris, K. J., & Evans, M. R. (2013).
Environmental and Parental Influences on Offspring Health and Growth in Great Tits
(Parus major). *PLOS ONE, 8*(7), e69695. doi:10.1371/journal.pone.0069695
Potti, J., Moreno, J., Yorio, P., Briones, V., García-Borboroglu, P., Villar, S., & Ballesteros, C.
(2002). Bacteria divert resources from growth for magellanic penguin chicks. *Ecology*
Letters, 5(6), 709-714. doi:10.1046/j.1461-0248.2002.00375.x
Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., . . . Glöckner, F. O. (2012).
The SILVA ribosomal RNA gene database project: improved data processing and
web-based tools. *Nucleic Acids Research, 41*(D1), D590-D596. Retrieved from
https://doi.org/10.1093/nar/gks1219
R Core Team. (2019). R: A language and environment for statistical computing. (Version
3.5.1). Vienna, Austria: R Foundation for Statistical Computing.
Ranstam, J. (2019). Hypothesis-generating and confirmatory studies, Bonferroni correction,
and pre-specification of trial endpoints. *Acta Orthopaedica, 90*(4), 297-297.
doi:10.1080/17453674.2019.1612624
Rauw, W. M. (2012). Immune response from a resource allocation perspective. *Frontiers in*
genetics, 3, 267-267. doi:10.3389/fgene.2012.00267

Ringsby, T. H., Sæther, B.-E., & Solberg, E. J. (1998). Factors Affecting Juvenile Survival in
781   House Sparrow Passer domesticus. *Journal of Avian Biology, 29*(3), 241-247.
782   doi:10.2307/3677106

783 Rosshart, S. P., Vassallo, B. G., Angeletti, D., Hutchinson, D. S., Morgan, A. P., Takeda, K., . . .
784   Rehermann, B. (2017). Wild Mouse Gut Microbiota Promotes Host Fitness and
785   Improves Disease Resistance. *Cell, 171*(5), 1015-1028.e1013. Retrieved from
786   http://www.sciencedirect.com/science/article/pii/S0092867417310656

787 Rowland, I., Gibson, G., Heinken, A., Scott, K., Swann, J., Thiele, I., & Tuohy, K. (2018). Gut
788   microbiota functions: metabolism of nutrients and other food components. *Eur J*
789   *Nutr, 57*(1), 1-24. doi:10.1007/s00394-017-1445-8

790 Schwagmeyer, P. L., & Mock, D. W. (2008). Parental provisioning and offspring fitness: size
791   matters. *Animal Behaviour, 75*(1), 291-298.
792   doi:https://doi.org/10.1016/j.anbehav.2007.05.023

793 Shannon, C. E. (1948). A Mathematical Theory of Communication. *Bell System Technical*
794   *Journal, 27*(3), 379-423. doi:10.1002/j.1538-7305.1948.tb01338.x

795 Sherwin, E., Bordenstein, S. R., Quinn, J. L., Dinan, T. G., & Cryan, J. F. (2019). Microbiota and
796   the social brain. *Science, 366*(6465), eaar2016. doi:10.1126/science.aar2016

797 Shutt, J. D., Nicholls, J. A., Trivedi, U. H., Burgess, M. D., Stone, G. N., Hadfield, J. D., &
798   Phillimore, A. B. (2020). Gradients in richness and turnover of a forest passerine's
799   diet prior to breeding: A mixed model approach applied to faecal metabarcoding
800   data. *Molecular Ecology, 29*(6), 1199-1213. doi:10.1111/mec.15394

801 Song, S. J., Sanders, J. G., Delsuc, F., Metcalf, J., Amato, K., Taylor, M. W., . . . Knight, R.
802   (2020). Comparative Analyses of Vertebrate Gut Microbiomes Reveal Convergence
803   between Birds and Bats. *mBio, 11*(1), e02901-02919. doi:10.1128/mBio.02901-19

804 Spencer, K. A., Evans, N. P., & Monaghan, P. (2009). Postnatal Stress in Birds: A Novel Model
805   of Glucocorticoid Programming of the Hypothalamic-Pituitary-Adrenal Axis.
806   *Endocrinology, 150*(4), 1931-1934. doi:10.1210/en.2008-1471

807 Sudo, N., Chida, Y., Aiba, Y., Sonoda, J., Oyama, N., Yu, X.-N., . . . Koga, Y. (2004). Postnatal
808   microbial colonization programs the hypothalamic-pituitary-adrenal system for
809   stress response in mice. *The Journal of Physiology, 558*(1), 263-275. Retrieved from
810   http://dx.doi.org/10.1113/jphysiol.2004.063388

811 Suzuki, T. A. (2017). Links between Natural Variation in the Microbiome and Host Fitness in
812   Wild Mammals. (1557-7023 (Electronic)). doi:10.1093/icb/icx104

813 Teyssier, A., Lens, L., Matthysen, E., & White, J. (2018). Dynamics of Gut Microbiota Diversity
814   During the Early Development of an Avian Host: Evidence From a Cross-Foster
815   Experiment. *Frontiers in Microbiology, 9*. Retrieved from
816   http://dx.doi.org/10.3389/fmicb.2018.01524

817 Therneau, T. M. (2020). coxme: Mixed Effects Cox Models (Version R package version 2.2-
818   16). Retrieved from https://CRAN.R-project.org/package=coxme

819 Torok, V. A., Hughes, R. J., Mikkelsen, L. L., Perez-Maldonado, R., Balding, K., MacAlpine, R., .
820   . . Ophel-Keller, K. (2011). Identification and Characterization of Potential
821   Performance-Related Gut Microbiotas in Broiler Chickens across Various Feeding
822   Trials. *Applied and Environmental Microbiology, 77*(17), 5868-5878. Retrieved from
823   http://dx.doi.org/10.1128/AEM.00165-11

824 Trevelline, B. K., Fontaine, S. S., Hartup, B. K., & Kohl, K. D. (2019). Conservation biology
825   needs a microbial renaissance: a call for the consideration of host-associated

microbiota in wildlife management practices. *Proceedings of the Royal Society B:*
Biological Sciences, 286(1895), 20182448. doi:10.1098/rspb.2018.2448
Turnbaugh, P. J., Ley, R. E., A., M. M., & et al. (2006). An obesity-associated gut microbiome
with increased capacity for energy harvest. *Nature, 444*, 1027-1031.
Van Noordwijk, A. J., McCleery, R. H., & Perrins, C. M. (1995). Selection for the Timing of
Great Tit Breeding in Relation to Caterpillar Growth and Temperature. *Journal of*
Animal Ecology, 64(4), 451-458. doi:10.2307/5648
Vargas-Pellicer, P., Watrobska, C., Knowles, S., Schroeder, J., & Banks-Leite, C. (2019). How
should we store avian faecal samples for microbiota analyses? Comparing efficacy
and cost-effectiveness. *Journal of Microbiological Methods, 165*, 105689.
doi:https://doi.org/10.1016/j.mimet.2019.105689
Vásquez, A., Forsgren, E., Fries, I., Paxton, R. J., Flaberg, E., Szekely, L., & Olofsson, T. C.
(2012). Symbionts as Major Modulators of Insect Health: Lactic Acid Bacteria and
Honeybees. *PLOS ONE, 7*(3), e33188. doi:10.1371/journal.pone.0033188
Videvall, E., Song, S. J., Bensch, H. M., Strandh, M., Engelbrecht, A., Serfontein, N., . . .
Cornwallis, C. K. (2019). Major shifts in gut microbiota during development and its
relationship to growth in ostriches. *Molecular Ecology, 28*(10), 2653-2667.
doi:10.1111/mec.15087
Videvall, E., Strandh, M., Engelbrecht, A., Cloete, S., & Cornwallis, C. K. (2017). Measuring
the gut microbiome in birds: Comparison of faecal and cloacal sampling. *Molecular*
Ecology Resources, 18(3), 424-434. Retrieved from http://dx.doi.org/10.1111/1755-
0998.12744
Vispo, C., & Karasov, W. H. (1997). The Interaction of Avian Gut Microbes and Their Host: An
Elusive Symbiosis. In R. I. Mackie & B. A. White (Eds.), (pp. 116-155): Springer US.
Wang, Q., Garrity Gm Fau - Tiedje, J. M., Tiedje Jm Fau - Cole, J. R., & Cole, J. R. (2007). Naive
Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial
taxonomy. (0099-2240 (Print)).
Warnecke, F., Luginbühl, P., Ivanova, N., Ghassemian, M., Richardson, T. H., Stege, J. T., . . .
Leadbetter, J. R. (2007). Metagenomic and functional analysis of hindgut microbiota
of a wood-feeding higher termite. *Nature, 450*(7169), 560-565.
doi:10.1038/nature06269
Wasielewski, H., Alcock, J., & Aktipis, A. (2016). Resource conflict and cooperation between
human host and gut microbiota: implications for nutrition and health. *Ann N Y Acad*
Sci, 1372(1), 20-28. doi:10.1111/nyas.13118
Weber, B. M., Bowers, E. K., Terrell, K. A., Falcone, J. F., Thompson, C. F., & Sakaluk, S. K.
(2018). Pre- and postnatal effects of experimentally manipulated maternal
corticosterone on growth, stress reactivity and survival of nestling house wrens.
Functional Ecology, 32(8), 1995-2007. doi:10.1111/1365-2435.13126
West, A. G., Waite, D. W., Deines, P., Bourne, D. G., Digby, A., McKenzie, V. J., & Taylor, M.
W. (2019). The microbiome in threatened species conservation. *Biological*
Conservation, 229, 85-98. doi:https://doi.org/10.1016/j.biocon.2018.11.016
Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*: Springer-Verlag New York.
Zaneveld, J. R., McMinds, R., & Vega Thurber, R. (2017). Stress and stability: applying the
Anna Karenina principle to animal microbiomes. *Nature Microbiology, 2*(9), 17121.
doi:10.1038/nmicrobiol.2017.121

Zhu, L., Wu, Q., Dai, J., Zhang, S., & Wei, F. (2011). Evidence of cellulose metabolism by the
giant panda gut microbiome. *Proceedings of the National Academy of Sciences,*
108(43), 17714. doi:10.1073/pnas.1017956108
Zimmer, C., Boogert, N. J., & Spencer, K. A. (2013). Developmental programming:
Cumulative effects of increased pre-hatching corticosterone levels and post-hatching
unpredictable food availability on physiology and behaviour in adulthood. *Hormones*
and Behavior, 64(3), 494-500. doi:https://doi.org/10.1016/j.yhbeh.2013.07.002
878