1   **A time-lagged association between the gut microbiome, nestling weight and nestling**
2   **survival in wild great tits**

4   Gabrielle L. Davidson[a,b,*⧓], Shane E. Somers[a,*], Niamh Wiley[c,d], Crystal N. Johnson[c,d], Michael
5   S. Reichert[a,e], R. Paul Ross[c,d], Catherine Stanton[c,d], John L. Quinn[a]
6
7   *Joint first authors
8   ⧓ Correspondence: gd339@cam.ac.uk
9

a. School of Biological, Earth and Environmental Sciences, Distillery Fields, North Mall,
University College Cork, Cork, Ireland.
b. Department of Psychology, Downing Street, University of Cambridge, Cambridge,
UK.
c. APC Microbiome Ireland, University College Cork, Cork, Ireland.
d. Teagasc Food Research Centre, Moorepark, Fermoy, Ireland.
e. Department of Integrative Biology, Oklahoma State University, USA

Abstract:

1. Natal body mass is a key predictor of viability and fitness in many animals. While
variation in body mass and therefore juvenile viability may be explained by genetic
and environmental factors, emerging evidence points to the gut microbiota as an
important factor influencing host health. The gut microbiota is known to change
during development, but it remains unclear whether the microbiome predicts
fitness, and if it does, at which developmental stage it affects fitness traits.
2. We collected data on two traits associated with fitness in wild nestling great tits
(*Parus major*): weight and survival to fledging. We characterised the gut microbiome
using 16S rRNA sequencing from nestling faeces and investigated temporal
associations between the gut microbiome and fitness traits across development at
day 8 (D8) and day 15 (D15) post-hatching. We also explored whether particular
microbial taxa were 'indicator species' that reflected whether nestlings survived or
not.

3. There was no link between mass and microbial diversity on D8 or D15. However, we detected a time-lagged relationship where weight at D15 was negatively associated with the microbial diversity at D8, controlling for weight at D8, therefore reflecting relative weight gain over the intervening period.

4. Indicator species analysis revealed that specificity values were high and fidelity values were low, suggesting that indicator taxa were primarily detected within either the survived or not survived groups, but not always detected in birds that either survived or died. Therefore these indicator taxa may be sufficient, but not necessary for determining either survival or mortality, perhaps owing to functional overlap in microbiota.

5. We highlight that measuring microbiome-fitness relationships at just one time point may be misleading, especially early in life. Instead, microbial-host fitness effects may be best investigated longitudinally to detect critical development windows for key microbiota and host traits associated with neonatal weight. Our findings should inform future hypothesis testing to pinpoint which features of the gut microbial community impact on host fitness, and when during development this occurs. Such confirmatory research will shed light on population level processes and could have the potential to support conservation.

Introduction:

Understanding the numerous factors that drive fitness has long been the focus of research in ecology and evolution (Merila, 1996; Pickett et al., 2013). This has primarily included behavioural factors, such as foraging (Patrick & Weimerskirch, 2014) and parental care (Schwagmeyer & Mock, 2008), and ecological factors, for example, food availability (Van Noordwijk et al., 1995), predation (Götmark, 2002), and more recently climate change (Blomberg et al., 2014). Proximate factors such as genetics (e.g. Merila (1996), Norris (1993)), stress (Crino & Breuner, 2015; Weber et al., 2018) and natal environment effects (Keller & Van Noordwijk, 1994; Pickett et al., 2013) have also been studied intensively. More recently, a growing body of research is beginning to show the gut microbiome - the

microbial community in the gut and the host environment - as a powerful proximate

mechanism influencing host health and fitness among animals (Kinross et al., 2011; Lloyd-

Price et al., 2016; Rosshart et al., 2017). Emerging evidence has highlighted that the gut

microbiota varies substantially within and between individuals (David et al., 2014; Maurice

et al., 2015), and that microbe-host interactions are likely to be taxonomically widespread

and central to many behavioural, ecological and evolutionary processes (Alberdi et al., 2016;

Davidson et al., 2020; Sherwin et al., 2019). However, the role of the gut microbiome in

determining fitness in natural populations, and whether there are critical developmental

windows whereby it affects future weight and survival, remains poorly understood.

Evidence suggests that the microbiome is associated with many aspects of survival and

fitness. In humans, the maternal vaginal microbiota is associated with pre-term birth

(Fettweis et al., 2019) and sudden infant death syndrome (Fettweis et al., 2019; Highet et

al., 2014), and the gut microbiome is associated with longevity (Biagi et al., 2016).

Laboratory-bred mice, which have substantially different gut microbiota than their wild

counterparts, were more likely to survive an influenza infection if they had received a gut

microbiota transplant from wild donors (Rosshart et al., 2017). Notably, wild mice had

higher abundances of Bacteroidetes and Proteobacteria, and lower abundances of

Firmicutes, suggesting these taxa may play a role in regulating host health. The potential for

the gut microbiome to impact animal fitness in natural systems has gained much attention

in recent years (Amato, 2013; Suzuki, 2017; Trevelline et al., 2019; West et al., 2019).

However, the empirical evidence to support such hypotheses is lacking, perhaps owing to

the logistical constraints associated with collecting longitudinal data on both the gut

microbiome and indicators of fitness such as survival in these systems.

Research on laboratory model species has shown that the microbial community in the gut

can confer specific health benefits to the host (Kinross et al., 2011), and the timing of gut

colonization is critical for normal development (Cox et al., 2014; Hansen et al., 2012). In

particular, gut microbiota are important for the development and regulation of gut

immunity by maintaining intestinal barrier function (Hooper et al., 2012; O'Hara &

Shanahan, 2006) and aiding digestive capabilities through the metabolism of dietary

components (Rowland et al., 2018). Gut bacteria can induce expression of host genes that,

for example, code for proteins in the mucosal layer of the intestines (O'Hara & Shanahan,

2006). Moreover, bacterial genomes can produce enzymes that are absent in the host, yet

are necessary for specific digestive functions such as cellulose digestion in termites

(*Nasutitermes* spp.) and giant pandas (*Ailuropoda melanoleuca*) (Warnecke et al., 2007; Zhu

et al., 2011) and nitrogen metabolism in birds (Vispo & Karasov, 1997). Germ-free mice

devoid of any microbes have depressed immune function and lower nutrient assimilation

efficiency, which can only be improved by introducing intestinal microbes from normal mice

donors (Hansen et al., 2012; Olszak et al., 2012). In many cases, the timing of microbial

colonization is critical and microbiome restoration must occur within an early

developmental window in order to affect host phenotypes that are observed later in life

(Cox et al., 2014; Hansen et al., 2012). Therefore, variation in host microbiota can have

important consequences for digestive capability and body condition, and act as a

mechanism underlying developmental plasticity (Alberdi et al., 2016). Importantly, effects

may be dependent on the timing of microbial colonization if microbial effects on host traits

may have immediate effects (i.e. contemporary associations), or delayed effects (i.e. time-

lagged associations).

In many vertebrate taxa, natal body size (Bowen et al., 2015; Ringsby et al., 1998; Stige et

al., 2019), natal body mass (Both et al., 1999; Monrós et al., 2002), or growth rate (Martin,

2015; Mikaela et al., 2002) can be key predictors of juvenile viability and recruitment into

the breeding population. The association between the gut microbiome and weight has been

studied extensively in clinical studies for application in human obesity (Bäckhed et al., 2004;

Turnbaugh et al., 2006), and in agricultural studies to increase animal productivity (Coates et

al., 1981; Ford & Coates, 1971; Torok et al., 2011). While having a more diverse microbiome

may intuitively suggest a healthier gut microbiome (Lozupone et al., 2012), the effects of

microbial community structure are likely more complex (Zaneveld et al., 2017). In humans,

lower diversity of gut microbiota is associated with high body mass index and high amounts

of visceral fat (Beaumont et al., 2016; Le Chatelier et al., 2013). Specific taxa, such as

Lactobacillus spp. can influence fat storage and weight change, however, the effects of such

bacteria are strain-specific (Drissi et al., 2017). In poultry, beneficial bacteria can be

selectively enriched (e.g. Patterson et al. (1997)), and the use of antibiotics reduce gut

bacterial load, decrease microbial diversity, and improve feed conversion ratios (Banerjee et

al., 2018). Weight gain associated with antibiotic administration and decreased bacterial
diversity has also been shown in house sparrows (*Passer domesticus*) (Kohl et al., 2018),
Daphnia magna (Motiei et al., 2020) and magellanic penguins (*Spheniscus magellanicus*)
(Potti et al., 2002). By contrast, microbial diversity has been positively associated with body
condition and growth rate in wild nestling birds (Teyssier, Lens, et al., 2018).

The great tit (*Parus major*) provides a powerful natural system for addressing microbiome
related effects on host phenotypes. It is an altricial species that readily breeds in nest boxes,
so that the developmental period can be tracked easily in the wild. We investigated the
relationship between natural variation in the gut microbiome and direct and indirect
measures of fitness-related traits in nestling great tits when they were 8 and 15 days old.
The neonatal stage is a critical time for gut microbiome colonization (Koenig et al., 2010)
and the microbial-programming of adult phenotypes (Cox et al., 2014; Hansen et al., 2012;
Sudo et al., 2004). Nestling great tits undergo substantial shifts in gut microbiota between
day 8 and day 15 of development (Teyssier, Lens, et al., 2018), a period when we predicted
the microbiome would have a detectable effect on body weight and the probability of
survival to fledging. We tested whether diversity of the microbiome, or the abundances of
specific microbes, were associated with nestling weight and survival to fledging. Given the
mixed reports of microbial links with host fitness in the existing literature (see above), we
had no a priori predictions regarding the direction of such potential relationships.
Nevertheless, we proposed that if diversity was positively related to measures of host
fitness, then a rich and/or evenly distributed microbial community would be important for
healthy development. By contrast, if the relationship was negative, then the host may
benefit from the presence or absence of specific microbial taxa, which should be observed
in differences in the relative abundance of these specific microbes associated with host
fitness. Moreover, because early developmental windows have been reported to be
important for microbial influences on host traits (Cox et al., 2014; Hansen et al., 2012), we
sampled the gut microbiome from chicks at two developmental stages to test whether
microbiome effects on host fitness traits were contemporary or time-lagged. In other words,
we asked whether the gut microbiome at D8 or D15 was a better predictor of mass and
survival.

Methods:

Field monitoring and microbiome sampling

Birds were sampled from nine nest box populations across Co. Cork, Ireland, five of these were mixed/deciduous habitats and four were coniferous habitats (see O'Shea et al. (2018)). We collected 204 faecal samples from 150 nestling great tits from 54 nests (see below for the number of samples that were successfully sequenced and sample size per developmental stage) for 16S rRNA gene sequencing in order to map the gastrointestinal microbial communities. During April-June 2016, nest boxes were monitored to determine lay dates, hatching dates and nestling survival. Nestlings aged 8 days (+/- 1 day) (D8) and 15 days (D15) were placed into sterile holding bags inside a heated holding case. Coffee filters were used to line the bags in order to soak up uric acid from the faeces, as uric acid has the potential to affect downstream sequencing (Khan et al., 1991). Samples were collected from nestlings that defaecated naturally within 15-20 minutes of being out of the nest before being returned to the nest. Although the aim was to have repeated samples for all individuals, not all birds survived, and not all birds produced samples within this time limit. Faecal sacks were opened using a sterile inoculation loop to release the faecal matter and place in a microcentrifuge tube containing 500uL of 100% ethanol. Samples were stored at -20^{o}C within 8 hours of collection until DNA extraction. D8 birds were weighed, and uniquely individually identified by clipping the tip of one of their nails, avoiding the blood vessel (commonly known as the "quick"). D15 nestlings were weighed and ringed with a unique identifiable metal ring (British Trust for Ornithology) and matched against their unique nail clipping. Nestlings for which we had repeated samples (i.e. for both D8 and D15) are referred to as 'repeat samples'.

DNA Extractions

Briefly, DNA was extracted from the dried faecal contents of all birds using the Qiagen QIAamp DNA Stool Kit, following the "Isolation of DNA from Stool for Pathogen Detection" protocol (June 2012 edition), with modifications described in (Shutt et al., 2020) to accommodate dried avian faeces. A 0.10 - 0.20 g aliquot of each faecal sample was added to the kit, alongside two negative controls which were carried through to sequencing. The negative controls had low sequence reads and were not included in any subsequent analyses. We note that no decontamination methods were used, though they can be

beneficial for samples with low biomass (Eisenhofer et al., 2019). Therefore, we recognise
that our sequence data may contain some Amplicon Sequence Variants (ASVs) from the
external environment, rather than the gut microbiome, although this should not have
systematically biased our results. DNA was stored at -20°C. Full extraction methods are
described in Supporting Information.

Illumina MiSeq sequencing
Full library preparation details are described in Supporting Information and in Davidson et al
2020b). Briefly, the V3-V4 variable region of the 16S rRNA gene was amplified from the DNA
extracts using the 16S metagenomic sequencing library protocol (Illumina). The DNA was
amplified with primers specific to the V3-V4 region of the 16S rRNA gene which also
incorporates the Illumina overhang adaptor. Samples were sequenced on the MiSeq
sequencing platform (Clinical Microbiomics, Denmark), using a 2 x 300 cycle kit, following
standard Illumina sequencing protocols.

Bioinformatics and statistical analysis
The DADA2 pipeline (Callahan et al., 2016) was used to process the raw sequencing data in R
version 3.5 (R Core Team, 2019). Sequence quality was visually inspected. Sequences were
trimmed to remove adapters and lower quality reads (median quality scores below 25-30
threshold) at the extremities of the sequence and filtered to remove sequences with higher
than expected errors. Read errors were estimated before dereplication. Forward and
reverse reads were merged to construct 'contig' sequences, these were used to construct a
sequence table of ASVs, which counts the number of times each unique sequence is
detected. The previous steps were performed for each run separately. Then the separate
sequence tables were merged and chimeras removed using the 'consensus' method.
Taxonomy was assigned to each ASV by RDP's Naive Bayes Classifier (Wang et al., 2007)
against the Silva reference database (version 132) (Quast et al., 2012). This is at 100%
sequence identity in contrast to the lower resolution OTU method which groups sequences
at 97% identity. ASVs allow greater sensitivity and specificity, better discrimination of
ecological patterns than OTU's and are reusable across studies (Callahan et al., 2017).

The DADA2 outputs were assembled into a single Phyloseq object (McMurdie & Holmes,
2013). Sequences identified as mitochondrial or chloroplast were removed. Further filtering
of samples and ASVs took place in R. Sample completeness curves were plotted using vegan
(Oksanen et al., 2019) and helped determine the lower cut-off for sample reads at 10,000
reads. Low read samples (<10000 reads, 9 samples) were removed leaving 195 (Day-8=81,
Day-15=114, repeat samples=41) samples for the analysis. Alpha diversity (both Shannon
and Chao1 diversity) was calculated using the *estimate_richness* function from the
phyloseq package on the filtered dataset. Shannon diversity (Shannon, 1948) measured
richness weighted by abundance (the evenness of a community) and Chao1 (Chao, 1984)
measured richness, specifically estimating taxa abundance and rare taxa missed from under
sampling.

Statistical analysis

Nestling weight: Alpha diversity and phylum-level relative abundance

Linear mixed models (LMMs) and generalised linear mixed models (GLMMs) were used to
examine the association between measures of nestling weight and two measures of alpha
diversity: Shannon's index and Chao1 index. Models were fit using the lme4 package (Bates
et al., 2015) on each data subset. Numeric variables were centered and scaled. Model
diagnostics were assessed using the DHARMa package (Hartig, 2019). Raw weight (g) was
used as the response variable instead of change in weight (delta-weight) as this allowed us
to control for D8 weight in time-lagged models, as well as to differentiate between, for
example, very light D8 nestlings achieving normal D15 weight and heavy D8 nestlings
achieving heavy D15 weight. We also tested whether alpha diversity at D8 was correlated
with alpha diversity at D15 (for repeated samples within individuals) by calculating
Pearson's correlation coefficient with the 'cor.test' function in R.

We examined whether nestling body weight was influenced by the gut microbiome in three
different ways: (1) whether the microbiome at an early developmental stage predicted
future host weight (henceforth 'time-lagged weight'), while controlling for weight at D8, (2)
whether the microbiome at the time of sampling was associated with weight at the same
developmental stage (henceforth 'contemporary weight') at D8, and (3) contemporary
weight at D15. Due to collinearity (alpha diversity) and non-independence (Phylum-level

relative abundance), we performed separate analyses for four different measures of
microbiome as explanatory variables: alpha diversity (Shannon diversity and Chao1 index),
and phylum-level abundance (Proteobacteria and Firmicutes). These two phyla were
selected for our analyses because they were the only two taxa that were prevalent across all
birds (except for three birds).

For the time-lagged weight analyses, we ran a linear mixed model (LMM) using the lmer
function in R with D15 weight as the response variable, and alpha diversity at D8 as a fixed
variable. We included D8 weight as a covariable to control for variation in starting weight.
We also included habitat, lay date of first egg and brood size as fixed factors as these may
influence nestling weight. Site and nest ID were used as nested random effects, as well as
sequence plate as a separate random effect. Only birds that had repeat samples were
included in this analysis (n= 41).

For the D8 contemporary weight analyses (all D8 samples, n= 81), we ran a LMM as
described above, with D8 as the response variable and alpha diversity at D8 as a fixed
factor. We included the same fixed effects and random effects as described above. The D15
contemporary analyses (all D15 samples, n = 114) were the same as described for D8, using
D15 data as opposed to D8 data. D15 weight was squared in the contemporary D15 model
as the model did not converge with raw weight as the response.

Nestling survival: Alpha diversity and phylum-level relative abundance
We tested whether microbiome diversity at D8 predicted nestling survival. Here we define
survival as surviving to fledgling. Birds that were previously alive at D8, but were either dead
in the nest, or absent (e.g. eviction from the parents) at D15 were scored as not surviving.
Nests were checked again approximately 25 days after hatching, when birds were expected
to have fledged. Birds found dead in the nest were scored as not surviving, and those that
were absent were assumed to have fledged because chicks were too large to be evicted,
and we did not see any evidence of predation (woodpeckers are absent in our woodlands,
and predation from stoats always led to the entire nest failing, and was isolated to a single
woodland site not included in the current study). We ran a non-parametric cox proportional
hazards survival model using the Coxme package (Therneau, 2020), and included weight (at

D8), lay date and brood size (at D8) as covariables. We included three ranks of survival times

(i.e. died before D15, died after D15 or fledged) as the ranks of survival times provide

additional temporal information beyond a binary outcome, i.e. died/survived for the model

to estimate mortality risk. The habitat variable was excluded from the survival models as it

caused premature model convergence due to a lack of variation in the variable (i.e. all

coniferous birds survived). Because all but one bird that survived to D15 survived to

fledging, we only performed this analysis for birds that were sampled for gut microbiome at

D8. We repeated this analysis with Proteobacteria abundance as the predictor variable, and

again with Firmicutes abundances as the predictor variables.

Nestling survival: Beta diversity

We investigated whether microbial community structure differed between birds that did or

did not survive to fledgling, using the adonis function (PERMANOVA) from the vegan

package (Oksanen et al., 2019). Beta diversity at D8 was measured using Bray-Curtis

distance (weighted by ASV abundance) and Jaccard (presence/absence of ASVs). We ran

separate analyses for each beta diversity measure in which D8 weight and survival (Yes/No)

were fixed factors. Adonis fits each term sequentially, therefore we used the model formula

beta-diversity~D8-weight + survival, allowing us to estimate the variation in microbiome

community composition associated with survival status after accounting for body weight, as

weight was associated with survival in our population (see results, Table 2). Permutations

were constrained to within nest with the 'strata' option to control for pseudo-replication of

samples from the same nest. Adonis results were very similar with and without this strata

option.

The beta diversity dataset (D8 samples) was filtered by prevalence prior to distance

calculation, where taxa with less than 1 copy in 5% of samples were removed following

Knowles et al. (2019). This resulted in a dataset containing 1254 ASVs. The relative

abundance of each phylum was calculated per sample using 'total sum scaling' (TSS) by

dividing each feature count by the total library size (McKnight et al., 2019). Bray Curtis and

Jaccard distances were calculated from our TSS normalized data. An assumption of the

adonis test is that groups have homogeneity of variance. The dispersion of the groups was

checked for homogeneity of variance using the Betadisper and permutest functions from
the vegan package.

The beta diversity (Bray-Curtis) of D8 nestlings was visualised using a Principle Coordinate
Analysis (PCoA) plot, created using the plot_ordination command in the phyloseq package.
This plot compared individuals that survived to those that did not survive (see Figure S2).

Indicator ASVs for nestling survival

We carried out indicator species analysis to explore whether ASVs were representative of
birds that were alive at D8 (n=81) and survived to fledgling (n=64), or did not survive to
fledgling (n=17). This analysis used the D8 data subset, which was not filtered by prevalence
and contained 29709 ASVs. Indicator analysis identified "indicator species" if they reflected
the state of the environment (i.e. likelihood the bird will survive, or not survive). We used
the multipatt function from the IndicSpecies package (De Caceres & Legendre, 2009), which
assigns a test statistic, or "indicator value" to each ASV for each group, in our case two
groups: Survival-Yes, Survival-No. Taxa with an indicator value of > 0.4 and p <= 0.05 were
considered indicator species. We also report the "specificity value" and the "fidelity value".
The former is the estimate of the probability that an ASV is detected in a particular group.
For example, a value of 1.0 would indicate that the ASV is only detected in that group and
not another group. The latter is the probability estimate of finding the species in sites (i.e.
birds) belonging to the group. For example, a value of 1.0 would indicate that the ASV is
recorded in all birds within that group (De Caceres, 2020). We did not adjust for multiple
comparisons for three reasons. First, our analysis did not include any overlap, and thus
multiple analyses, across different combinations of groups (De Caceres & Legendre, 2009).
Second, the purpose of this analysis was exploratory in order to generate, rather than
confirm hypotheses (Bender & Lange, 2001; Ranstam, 2019). As such, we report ASVs as
potential indicator species, as opposed to reporting a quantitative, total number of taxa
associated with survival/non-survival (De Cáceres et al., 2010). Third, this analysis served to
generate a posteriori predictions regarding the relationship between indicator taxa and
weight gain (see below), although we note that independent studies will be necessary to
confirm these exploratory analyses (Ranstam, 2019). We generated a heatmap of indicator

species for survival and non-survival using the heatmap function from the phyloseq package

(McMurdie & Holmes, 2013).

A posteriori analysis of indicator species

Indicator species from our analysis may reflect whether a bird survives or not. We

hypothesized that indicator species may reflect host survival because such indicator species

may also predict time-lagged, relative weight gain. We performed a posteriori LMMs with

D15 weight as the response variable and D8 relative abundance (of the indicator species) as

a fixed variable. D8 weight, habitat, brood size and lay date were also included as fixed

variables. Site and nest were included as nested random terms (i.e. 1|site/nest). We

restricted this a posteriori analysis to one indicator species that had the highest test statistic

from each group (Survival-Yes: Lactobacillus (ASV46) and Survival-No: Rahnella (ASV38)).

Because there were a high number of zeros in relative abundance of these taxa, and

because our data were proportions and therefore could not be fit using a zero-inflated

(count data) model, we included only birds where the taxa were present and the birds with

0 abundance for that taxa were excluded (ASV46: n=21/41; ASV38: n=8/41).

All graphs were created using ggplot2 (Wickham, 2016), unless otherwise indicated.

Results:

Sequencing results

There were 18,537,795 high quality reads, clustered into 49,655 ASVs before removing

control samples, mitochondria and chloroplast sequences. Mean reads per sample was

88,698 and individual samples ranged from 61 to 579,864 reads. Samples with less than

10,000 reads were discarded before further analysis (n=9). There were 15,183,675 total

high-quality reads, clustered into 47,400 ASVs after removing control samples as well as

mitochondria and chloroplast sequences. Samples ranged from 10,220 to 557,336 reads and

mean reads per sample was 77,865.

The most prominent phyla across all samples (D8 and D15) were as follows (mean

percentage relative abundance ±SE): Firmicutes (43.0%±2.5); Proteobacteria (35.2%±2.2);

Tenericutes (9.7%±1.8); Bacteroidetes (4.2%±0.4) and Actinobacteria (2.2%±0.2). Relative

 abundance of the major phyla at D8 (at least >=10% relative abundance in at least 1 sample)

 per sample are plotted in Figure 1.

Figure 1. Relative abundance of the 8 most prominent phyla (>=10% relative abundance). Each vertical bar corresponds to a nestling at Day 8. "Other" group represents all low abundance phyla as white space.

 Nestling weight

 The time-lagged analysis indicated that weight at D15 was negatively related to alpha

 diversity at D8 (Figure 2a, Table 1, Table 4), controlling for weight at D8 (Table 1). Neither

 habitat, lay date or brood size had detectable influence in this analysis (Table 1). Cross-

 sectional analyses indicated no relationship between contemporary weight and diversity at

 either D8 or D15 (Figure 2b and 2c, Table 4, Table S1). Brood size was negatively related to

 weight at D8. There was no evidence for an effect of habitat on weight in D15 birds. The

 relative abundances of Proteobacteria and Firmicutes did not predict weight in any of these

 analyses.

Figure 2. The effect of Shannon diversity on (a) time-lagged weight (i.e. Day 15 (D15)), (b) contemporary weight Day 8 (D8) and (c) contemporary weight D15. Black dots are individual data points, blue line is regression line with 95% CI (shaded blue). Shannon diversity is centred and scaled.

Table 1. GLMM output from time-lagged (Day-15) weight analyses. The effect of alpha diversity is reported for (a) Shannon diversity and (b) Chao1 diversity. * p < 0.05.

Dependent~Independent variable	Estimate	Std. Error	df	Test statistic	P_estimate
(a) Nestling weight (time-lagged)					
(Intercept)	16.798	0.525	21.052	31.977	<0.001 *
Shannon (Day-8)	-0.593	0.167	23.749	-3.544	0.002 *
Habitat (deciduous)	0.247	0.630	22.081	0.392	0.699
Weight (Day-8)	0.700	0.173	24.672	4.035	<0.001 *
Lay Date (first egg)	0.252	0.296	21.226	0.851	0.404
Brood size (Day-8)	0.168	0.273	21.991	0.616	0.544
(b) Nestling weight (time-lagged)					

(Intercept)	16.961	0.475	20.856	35.681	<0.001 *
Chao1 (Day-8)	-0.525	0.184	34.314	-2.855	0.007 *
Habitat (deciduous)	0.011	0.572	21.903	0.019	0.985
Weight (Day-8)	0.528	0.192	32.990	2.756	0.009 *
Lay Date (first egg)	0.257	0.270	21.399	0.949	0.353
Brood size (Day-8)	0.121	0.250	22.841	0.483	0.634

Correlation tests indicate both Shannon (cor=0.33, t=2.17, df=39, p= 0.036) and Chao1
(cor=0.36, t=2.38, df=39, p=0.022) diversity measures were positively correlated within
pairs.

Nestling Survival
Survival was significantly related to weight at D8; the hazard ratio of 0.67 indicated that for
every unit increase in weight, mortality risk decreased by a factor of 0.67. Survival was not
influenced by lay date or brood size. Nestling survival to fledging was not predicted by
Shannon's diversity or Chao1 index at D8 (Table 2, Table 4). The relative abundance of
Proteobacteria and Firmicutes at D8 also did not predict survival to fledging.

There was no clear evidence for the violation of the assumption of homogeneity of
dispersion for survived/not-survived groups for Jaccard distance (Sum Squares= 0.009,
F=3.295, p=0.081) or Bray-Curtis (Sum squares=0.006, F=1.973, p=0.164). Beta diversity did
not differ between surviving and non-surviving nestlings for Jaccard (Sum Squares=0.563,
F=1.233, p=0.172) or Bray-Curtis (Sum Squares=0.523, F=1.193, p=0.228), Table 4, and see
Supporting Information Figure S2, Table S3.

Table 2. Determinants of mortality risk from Cox proportional hazards model. Positive
coefficients indicate increased risk of mortality. HR = Hazard ratio, which is the
exponentiated coefficients, and gives the ratio of the total number of observed to
expected events. Note: all numeric covariates are centered and scaled. * p < 0.05.

			Standard		
Dependent~Independent variable	Coefficient	HR	error	Test statistic (z)	P estimate

Nestling mortality risk

	Coefficient	HR	error	Test statistic (z)	P estimate
Shannon D8	0.021	1.021	0.327	0.07	0.95
Weight D8	-0.665	0.514	0.282	-2.36	0.018 *
Lay Date (First egg)	0.304	1.356	0.382	0.80	0.43
Brood Size (Day-8)	0.551	1.735	0.363	1.52	0.13

Nestling mortality risk

	Coefficient	HR	error	Test statistic (z)	P estimate
Chao1 D8	0.314	1.368	0.283	1.11	0.27
Weight D8	-0.609	0.544	0.289	-2.11	0.035 *
Lay Date (First egg)	0.297	1.346	0.396	0.75	0.45
Brood Size (Day-8)	0.634	1.885	0.386	1.64	0.1

Indicator species analysis

The indicator species with the highest indicator values associated with survival belonged to

ASV46, ASV310 and ASV150 which all belong to the *Lactobacillaceae* family (Figure 3, Table

4, Table S4). The taxa with the highest indicator values associated with non-survival

belonged to the ASV38, ASV1034, ASV895 which belong to *Enterobacteriaceae*,

Methylopilaceae and *Acidothermaceae* families, respectively. The taxa indicative of non-

survival and survival (p value <=0.05 and test statistic >=0.4) were plotted on a heatmap

(Figure 3). Overall, specificity values were high (mean = 0.94±0.008 standard error) and

fidelity values were low (0.21±0.008), suggesting that indicator species were typically only

detected within a given group (i.e. survive/non-survive), but that not all birds within a group

were recorded to have a given indicator species. All indicator species and test statistics

(specificity values and indicator values) are reported in Supporting Information Table S4.

A posteriori analyses showed that a higher relative abundance of ASV46 (a *Lactobacillus*

spp., and the taxa most strongly associated with survival) at D8 predicted a higher weight at

D15. Relative abundance of ASV38 (a *Rahnella* sp., and the taxa most strongly associated

with non-survival) was not associated with weight at D15 (Table 3, Table 4).

Figure 3. Heatmap of relative abundance of indicator species ASVs for survival. Individual birds are columns with separate panels for individuals that either died or survived. ASVs are in rows where taxa in brackets refer to the ASVs family-level taxonomic assignment, or NA if not resolved to family-level. Taxa associated with survival as opposed to non-survival are in

bold. Lighter colours indicate greater abundance.

Table 3. A posteriori GLMM output from time-lagged weight analyses. The effect of indicator species on weight at D15 are reported for a) *Lactobacillus* sp. (ASV46) and b) *Rahnella* sp. (ASV38). * p <0.05

Dependent/Independent variable	Estimate	Std. Error	df	Test statistic	P_estimate
(a) D15 Weight (time-lagged)					
(Intercept)	17.673	0.739	4.679	23.906	<0.001 *
Abundance **Lactobacillus sp. (ASV46)**	1.042	0.367	8.601	2.839	0.02 *
Habitat (deciduous)	-1.946	1.007	5.126	-1.933	0.11
Weight (Day-8)	0.835	0.340	11.184	2.460	0.031 *
Brood Size (Day-15)	-0.459	0.352	11.573	-1.303	0.218
Lay Date (First egg)	-0.457	0.389	7.245	-1.175	0.277
(b) D15 Weight (time-lagged)					
(Intercept)	16.473	0.656	2.000	25.111	0.002 *
Abundance **Rahnella sp. (ASV38)**	0.040	0.388	2.000	0.103	0.927
Habitat (deciduous)	2.908	1.320	2.000	2.203	0.158
Weight (Day-8)	5.398	1.579	2.000	3.419	0.076
Brood Size (Day-15)	1.690	0.660	2.000	2.558	0.125
Lay Date (First Egg)	-1.069	1.113	2.000	-0.960	0.438

Table 4 Summary of the relationship between microbiota and host performance

Microbiome metric	Survival	Non-survival	Weight (time-lagged)	Weight (contemporary)
Alph diversity	NS	NS	(-)	NS
Beta diversity	NS	NS		
Phylum-level abundance	NS	NS	NS	NS
Inidicator species	(+)	(+)		
Lactobacillus sp. (ASV 46)			(+)	NS
Rahnella sp. (ASV38)			NS	NS

438 Significant relationships are shown as either positive (+), negative (-) or non significant (NS).
439
Discussion:

Overall summary

We demonstrate that the gut microbiota correlates with nestling performance in a
population of wild passerine birds. We found a negative association between weight at D15
and alpha diversity at D8. In other words, individuals with a low microbial diversity at D8
were heavier than expected at D15, given their weight at D8. Although alpha diversity at D8
was correlated with alpha diversity at D15, neither microbial alpha diversity nor phylum-
level abundance predicted contemporary weight. Our exploratory analysis indicated that
eight-day old nestlings with a greater relative abundance of specific *Lactobacillus* taxa were
more likely to fledge, and our a posteriori analysis showed that for at least one of these
taxa, greater relative abundance was specifically associated with increased host weight
(relative to weight at D8). Notably, although we present potential links between gut
microbiota and survival, and a proxy for survival (i.e. weight), the direction of these
relationships or the microbial metrics that predict such relationships are variable within our
own study (Table 4) and contrast with some, but not other studies. A lack of consensus on
what aspects of microbial taxonomic composition predict host traits is prominent in the
microbiome literature (Zaneveld et al., 2017). This may be due to broad statistical
approaches that are correlative in nature, and therefore require manipulative studies to
confirm relationships found here. Nevertheless, we discuss below how methodological
approaches, critical developmental windows, environmental matching and the Anna
Karenina Hypothesis may explain such differential findings across studies. Moreover, we
suggest hypotheses worthy of future confirmatory testing to understand the effects of gut
microbiota on the development of host phenotypes.

We show a negative relationship between alpha diversity and relative weight gain, over one
nesting season. Experimentally reducing diversity of the microbiome through antibiotic
administration has been shown to improve host growth (Banerjee et al., 2018; Kohl et al.,
2018; Lan et al., 2005; Potti et al., 2002), possibly because antibiotics reduce bacterial load,
diverting resources towards growth, as opposed to immune function, in a classic life-history
trade-off (Rauw, 2012). Our study demonstrates that this negative relationship is present in
the absence of microbiome manipulation, although we can only speculate as to the
mechanisms responsible for limiting relative weight gain in nestlings with high gut microbial
diversity. For example, a conventional gut microbiome compared to a germ-free one lowers
absorption capacity of glucose and B-vitamins by increasing the thickness of the gut
epithelium (Ford & Coates, 1971), and non-mutualistic microbes may create greater
competition for nutrients (Wasielewski et al., 2016). While the diversity of the microbial
community may be important for regulating the host's future weight, some taxa may have
disproportionate effects. At least one ASV, *Lactobacillus* sp., with the strongest potential
association with survival, showed a positive, time-lagged association with host weight
(relative to weight at D8). Several *Lactobacillus* spp. have beneficial, probiotic effects on the
host (Angelakis & Raoult, 2010; Banerjee et al., 2018; Gomes et al., 2012; Vásquez et al.,
2012), though this is certainly not the case for all *Lactobacillus* spp. Nevertheless, we can
speculate as to why Lactobacillus may confer benefits to great tit nestlings, and encourage
confirmatory research to test such hypotheses. For example, some strains of *Lactobacillus*
sakei (ASV150) are known to produce antimicrobial peptides which inhibit a known
pathogen (Gomes et al., 2012). Therefore we hypothesise that the presence of *Lactobacillus*
sakei, which was associated with survival, may serve as an anti-pathogen in great tit
nestlings. More generally, *Lactobacillus* spp. produce metabolic byproducts such as short
chain fatty acids that can improve host energy metabolism (LeBlanc et al., 2017) and
modulate immune response (Ratajczak et al., 2019). For example, the SCFA butyrate acts as
an anti-inflammation agent, enhances intestinal barrier function and modulates the
production of mucosal immune products in mammals (see review: Liu et al. (2018)). Future
studies should look to shotgun sequencing to identify specific bacterial strains and genes for
functional profiling and for developing probiotic treatments. This could be followed up by
manipulative studies to identify the causal relationships between ASV and host traits, such

as isolating and culturing ASV46 to test its potential probiotic effects when administered to

nestlings (Davidson et al., 2020; Gatesoupe, 1999; Holzapfel & Schillinger, 2002).

Although a growing number of studies have demonstrated that the microbiome is an

important predictor of host weight, the microbial profiles and the direction of this

relationship vary across species and studies (Kohl et al., 2018; Potti et al., 2002; Teyssier,

Lens, et al., 2018; Videvall et al., 2019). For example, a negative relationship between alpha

diversity and future weight gain was found in juvenile ostriches (Videvall et al., 2019),

whereas a positive relationship between alpha diversity and contemporary body condition

was reported in a different population of great tit nestlings (Teyssier, Lens, et al., 2018). The

contrasting results reported in these studies and our own may be due to variations in the

microbiome sampling method and storage solutions (i.e. cloacal versus faecal, see Videvall

et al. (2017); storing solutions see Vargas-Pellicer et al. (2019)). Moreover, our study and

that of others are limited to a single nesting season, and any general patterns of the gut

microbiota on host traits may differ across years according to local conditions such as food

availability (see below for discussion of Predictive Adaptive Response hypothesis). Therefore

it is likely that differences in local environments (Gillingham et al., 2019; Teyssier, Rouffaer,

et al., 2018; Youngblut et al., 2019) and the temporal resolution of microbial and host traits

dictate when and how microbiota impacts host phenotypes in birds (Videvall et al., 2019).

Time-lagged associations between the gut microbiota and host traits are expected to be

especially important during developmental periods when large shifts in the establishing

microbial community coincide with the programming of host biology (Cox et al., 2014;

Hansen et al., 2012; Sudo et al., 2004), and in systems that are likely to experience abrupt

changes in environmental inputs associated with gut microbiome, such as diet. The latter

may be especially relevant in birds as their microbiomes are tightly linked with

environmental variation, in contrast to mammals where the microbiome is primarily linked

to phylogeny (Song et al., 2020). In ostriches, microbiome diversity predicted future weight

gain, but the direction of this relationship switched after week 1 post-hatching, a time when

ostriches no longer rely on an internal yolk as sustenance (Videvall et al., 2019). In the

current study, alpha diversity at D8 predicted future weight at D15, whereas cross-sectional

analyses were not sufficient to detect such a relationship. This time-lagged association may

be owing to a critical developmental period in which the microbiome impacts on future
phenotypes, a phenomenon that is prominent in clinical microbiome literature (Cox et al.,
2014; Hansen et al., 2012; Sudo et al., 2004). The gut microbiome may also act as a
mechanism for adaptive phenotypic change through its sensitivity to environmental
variability, and its connection to host biological processes (Alberdi et al., 2016). Such a
mechanism would also be in line with the 'Predictive Adaptive Response hypothesis', which
posits that conditions during development cause phenotypic changes that serve to match
individuals to their environments (Bateson et al., 2014; Crino & Breuner, 2015; Weber et al.,
2018). For example, unpredictable food supply during the development stage can influence
adult physiological and behavioral phenotypes (Spencer et al., 2009; Zimmer et al., 2013).
Therefore, microbiome-mediated developmental plasticity may be adaptive in species
where offspring benefit from a gut microbiome that primes the host for dietary shifts.
Dietary shifts are common in tits where the relative frequency of insect taxa can change
throughout the provisioning period (Arnold et al., 2007; García-Navas et al., 2012). Although
Belgian great tit nestlings had the greatest improvement in body condition if they
maintained a stable microbial diversity throughout development (Teyssier, Lens, et al.,
2018), perhaps this may be because the environmental conditions at the time were also
stable. In our study, it is difficult to differentiate between whether the gut microbiome is
responding to dietary shifts as opposed to priming the host. Pinpointing if and when early
life bacterial diversity programs host digestive and/or metabolic efficiency in wild animals,
and the extent to which environmental variation, such as food availability and diet, impacts
the developing microbiome will progress our understanding of microbial-mediated
developmental mechanisms.

The Anna Karenina Hypothesis states that changes in an unhealthy or so-called "dysbiotic"
microbiome can be stochastic rather than deterministic, making them difficult to detect
statistically (Zaneveld et al., 2017). In other words, unhealthy individuals can have greater
variation in their microbial community composition than healthy individuals, and rather
than displaying a specific 'unhealthy' state, the microbiomes of diseased or stressed
individuals can all look very different due to a breakdown in regulation. Although our study
did not examine diseased individuals, as the original Anne Karenina Hypothesis was
intended, this phenomenon may nevertheless explain why nestling survival was not

associated with either alpha or beta diversity. In contrast to the Anne Karenina Hypothesis
and our own study, disease-causing mortality in juvenile ostriches was associated with low
alpha diversity, and beta diversity differed between diseased and non-diseased birds
(Videvall et al., 2020). In the present study, although several indicator species were
identified to be associated with birds that did or did not survive, many of these had low
prevalence across nestlings. This is reflected in the high specificity values and low fidelity
values, suggesting that although the presence of an indicator species in a nestling's gut
microbiome reflects a high probability of survival or non-survival, the same indicator species
is not present across all surviving/non-surviving nestlings in our sample population. This
distinction is important in, for example, conservation practices whereby microbes are used
as markers for population viability (Trevelline et al., 2019; West et al., 2019). Our analysis
suggests that in the case of non-survival, there are no microbes that are universally
indicative of non-survival across birds, but rather the presence of one of several candidate
microbes predicts a higher probability of mortality, which is consistent with the Anna
Karenina Hypothesis. Alternatively, the gut microbiome does not predict survival in great tit
nestlings at all, and the indicator species flagged in our study are an artifact of multiple
testing without a correction due to the exploratory nature of this analysis. Therefore,
whether these same taxa affect survival in other populations and/or other species of birds
should be examined in independent confirmatory studies. For example, we hypothesise that
ASVs reported for non-survival may be pathogenic to great tits, although we found no
evidence in the literature to support this. Nevertheless, ASV1320, ASV2151, ASV3604 and
ASV942 are members of the Rhizobiales order which contains genera that are well known
human pathogens, and ASV5768 and ASV4855 are members of the Chlamydiales order
which also contain pathogenic genera, though these genera were not detected in our study.
583
We also encourage further investigations into whether indicator taxa are the cause or
consequence of host growth, or indeed any trait that may be associated with survival, such
as glucocorticoids (Weber et al., 2018) and immune function (Hõrak et al., 1999).
Manipulation of the gut microbiome alongside metabolomics or metagenomics will be the
next step in providing insight into the causal direction and functional mechanisms of
microbial variation and its relationship with host fitness. The latter is particularly important
for identifying whether there is functional overlap between microbial taxa and the

mechanisms by which these taxa affect host traits (Heintz-Buschart & Wilmes, 2018). Given
the temporal importance of host-microbial relationships identified in this study and
elsewhere, we anticipate the maturing microbiome will have important consequences on
long-term fitness effects. More generally, our results highlight the importance of the gut
microbiome in evolutionary processes as it has the potential to mediate rapid adaptation to
environmental change and may also support conservation efforts as an indicator of
individual and population health.
598

Funding.
This work was supported by Science Foundation Ireland by way of funding to APC
Microbiome Ireland, Cork, Ireland. GLD, SES, MRC and JLQ etc were supported by funding
from the European Research Council under the European Union's Horizon 2020 Programme
(FP7/2007-2013)/ERC Consolidator Grant "Evoecocog" Project No. 617509, awarded to JLQ.
GLD is funded by a Leverhulme Trust Early Career Fellowship (ECF-2018-700) with matched
funding from the Sir Isaac Newton Trust.
606
Acknowledgements.
Thanks to Jodie Crane and Iván de la Hera, for assistance with fieldwork. Paul Cotter, Fiona
Crispie and Laura Finnegan from the Teagasc Sequencing facility for their role in relation to
the 16S rRNA sequencing. For advice on bioinformatics, we thank Fiona Fouhy, Orla
O'Sullivan and Calum Walsh. We are grateful to James Nichols, Ally Phillimore and Sarah
Knowles for helpful discussions on molecular methodology, and to Enrico Pirotta for advice
on statistical analyses. Thank you to Professor Jean-Michel Gaillard, the Associate Editor,
and two anonymous referees for their constructive comments.

615
616 Data accessibility.
617 Our metadata is deposited at Dryad: doi:10.5061/dryad.bk3j9kd9g, and sequence data will
618 be uploaded to a repository such as NCBI Sequence Read Archive, if the manuscript is
619 published.
620
Research and Animal Ethics: This study was conducted under licences from the Health
Products Regulatory Authority (AE19130_P017), The National Parks and Wildlife Services

(010/2016) and the British Trust for Ornithology, and permission from Coillte Forestry and

private landowners. The research project received ethical approval from the Animal Welfare

Body at University College Cork, and was in accordance with the ASAB (Association for the

Study of Animal Behaviour) Guidelines for the Treatment of Animals in Behavioural

Research and Teaching.

Author contributions: GLD, CS, RPR and JLQ conceived the ideas and designed the

methodology. GLD and MSR collected the field data. GLD, NW and CNJ carried out the DNA

extraction and library prep. SES analysed the data. GLD and SES wrote the manuscript with

input from all authors. All authors gave final approval for publication.

References:

Alberdi, A., Aizpurua, O., Bohmann, K., Zepeda-Mendoza, M. L., & Gilbert, M. T. P. (2016).
Do Vertebrate Gut Metagenomes Confer Rapid Ecological Adaptation? *Trends in*
Ecology & Evolution, 31(9), 689-699. doi:https://doi.org/10.1016/j.tree.2016.06.008
Amato, K. R. (2013). Co-evolution in context: The importance of studying gut microbiomes in
wild animals. *Microbiome Science and Medicine, 1*(1). Retrieved from
http://dx.doi.org/10.2478/micsm-2013-0002
Angelakis, E., & Raoult, D. (2010). The increase of Lactobacillus species in the gut flora of
newborn broiler chicks and ducks is associated with weight gain. *PLOS ONE, 5.*
doi:https://doi.org/10.1371/journal.pone.0010463
Arnold, K. E., Ramsay, S. L., Donaldson, C., & Adam, A. (2007). Parental prey selection affects
risk-taking behaviour and spatial learning in avian offspring. *Proceedings of the Royal*
Society B: Biological Sciences, 274(1625), 2563-2569. doi:10.1098/rspb.2007.0687
Bäckhed, F., Ding, H., Wang, T., Hooper, L. V., Koh, G. Y., Nagy, A., . . . Gordon, J. I. (2004).
The gut microbiota as an environmental factor that regulates fat storage.
Proceedings of the National Academy of Sciences of the United States of America,
101(44), 15718-15723. doi:10.1073/pnas.0407076101
Banerjee, S., Sar, A., Misra, A., Pal, S., Chakraborty, A., & Dam, B. (2018). Increased
productivity in poultry birds by sub-lethal dose of antibiotics is arbitrated by
selective enrichment of gut microbiota, particularly short-chain fatty acid producers.
Microbiology, 164(2), 142-153. doi:https://doi.org/10.1099/mic.0.000597
Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software, 67*(1), 1-48.
Bateson, P., Gluckman, P., & Hanson, M. (2014). The biology of developmental plasticity and
the Predictive Adaptive Response hypothesis. *The Journal of Physiology, 592*(11),
2357-2368. doi:https://doi.org/10.1113/jphysiol.2014.271460
Beaumont, M., Goodrich, J. K., Jackson, M. A., Yet, I., Davenport, E. R., Vieira-Silva, S., . . .
Bell, J. T. (2016). Heritable components of the human fecal microbiome are
associated with visceral fat. *Genome Biology, 17*(1), 189. doi:10.1186/s13059-016-
1052-7

Bender, R., & Lange, S. (2001). Adjusting for multiple testing—when and how? *Journal of*
Clinical Epidemiology, 54(4), 343-349. doi:10.1016/S0895-4356(00)00314-0

Biagi, E., Franceschi, C., Rampelli, S., Severgnini, M., Ostan, R., Turroni, S., . . . Candela, M.
(2016). Gut Microbiota and Extreme Longevity. *Current Biology, 26*(11), 1480-1485.
doi:https://doi.org/10.1016/j.cub.2016.04.016

Blomberg, E. J., Sedinger, J. S., Gibson, D., Coates, P. S., & Casazza, M. L. (2014). Carryover
effects and climatic conditions influence the postfledging survival of greater sage-
grouse. *Ecology and Evolution, 4*(23), 4488-4499. doi:10.1002/ece3.1139

Both, C., Visser, M. E., & Verboven, N. (1999). Density–dependent recruitment rates in great
tits: the importance of being heavier. *Proceedings of the Royal Society of London.*
Series B: Biological Sciences, 266(1418), 465-469. doi:10.1098/rspb.1999.0660

Bowen, W. D., den Heyer, C. E., McMillan, J. I., & Iverson, S. J. (2015). Offspring size at
weaning affects survival to recruitment and reproductive performance of
primiparous gray seals. *Ecology and Evolution, 5*(7), 1412-1424.
doi:10.1002/ece3.1450

Callahan, B. J., McMurdie, P. J., & Holmes, S. P. (2017). Exact sequence variants should
replace operational taxonomic units in marker-gene data analysis. *The ISME Journal,*
11(12), 2639-2643. doi:10.1038/ismej.2017.119

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.
(2016). DADA2: High-resolution sample inference from Illumina amplicon data.
Nature Methods, 13, 581-581 EP. Retrieved from
https://doi.org/10.1038/nmeth.3869

Chao, A. (1984). Nonparametric Estimation of the Number of Classes in a Population.
Scandinavian Journal of Statistics, 11(4), 265-270. Retrieved from
www.jstor.org/stable/4615964

Coates, M. E., Cole, C. B., Fuller, R., Houghton, S. B., & Yokota, H. (1981). The gut microflora
and the uptake of glucose from the small intestine of the chick. *British Poultry*
Science, 22(3), 289-294. Retrieved from
http://dx.doi.org/10.1080/00071688108447888

Cox, L. M., Yamanishi, S., Sohn, J., Alekseyenko, A. V., Leung, J. M., Cho, I., . . . Blaser, M. J.
(2014). Altering the intestinal microbiota during a critical developmental window has
lasting metabolic consequences. *Cell, 158*(4), 705-721.
doi:10.1016/j.cell.2014.05.052

Crino, O. L., & Breuner, C. W. (2015). Developmental stress: evidence for positive
phenotypic and fitness effects in birds. *Journal of Ornithology, 156*(1), 389-398.
doi:10.1007/s10336-015-1236-z

David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., . . .
Turnbaugh, P. J. (2014). Diet rapidly and reproducibly alters the human gut
microbiome. *Nature, 505*(7484), 559-563. doi:10.1038/nature12820

Davidson, G. L., Raulo, A., & Knowles, S. C. L. (2020a). Identifying Microbiome-Mediated
Behaviour in Wild Vertebrates. *Trends in Ecology & Evolution*.
doi:https://doi.org/10.1016/j.tree.2020.06.014

Davidson, G.L., Wiley, N., Cooke, A.C., Johson, C.N., Fouhy, Fiona, Reichert, M.S., de la Hera,
I., Crane, J.M.S., Kulahci, I.G., Ross, R.P., Stanton, C., Quinn, J.L. Diet induces parallel
changes to the gut microbiota and problem solving performance in a wild bird.
Scientific Reports 10, 20783 (2020b). https://doi.org/10.1038/s41598-020-77256-y

De Caceres, M. (2020). How to use the indicspecies package (ver. 1.7.8). Retrieved from
https://cran.r-
project.org/web/packages/indicspecies/vignettes/indicspeciesTutorial.pdf
De Caceres, M., & Legendre, P. (2009). Associations between species and groups of sites:
indices and statistical inference. *Ecology*.
De Cáceres, M., Legendre, P., & Moretti, M. (2010). Improving indicator species analysis by
combining groups of sites. *Oikos, 119*(10), 1674-1684. doi:10.1111/j.1600-
0706.2010.18334.x
Drissi, F., Raoult, D., & Merhej, V. (2017). Metabolic role of lactobacilli in weight
modification in humans and animals. *Microbial Pathogenesis, 106*, 182-194.
doi:https://doi.org/10.1016/j.micpath.2016.03.006
Eisenhofer, R., Minich, J. J., Marotz, C., Cooper, A., Knight, R., & Weyrich, L. S. (2019).
Contamination in Low Microbial Biomass Microbiome Studies: Issues and
Recommendations. *Trends in Microbiology, 27*(2), 105-117.
doi:10.1016/j.tim.2018.11.003
Fettweis, J. M., Serrano, M. G., Brooks, J. P., Edwards, D. J., Girerd, P. H., Parikh, H. I., . . .
Buck, G. A. (2019). The vaginal microbiome and preterm birth. *Nature Medicine,
25*(6), 1012-1021. doi:10.1038/s41591-019-0450-2
Ford, D. J., & Coates, M. E. (1971). Absorption of glucose and vitamins of the B complex by
germ-free and conventional chicks. *Proc Nutr Soc, 30*(1), 10a-11a.
García-Navas, V., Ferrer, E. S., & Sanz, J. J. (2012). Prey selectivity and parental feeding rates
of Blue Tits Cyanistes caeruleus in relation to nestling age. *Bird Study, 59*(2), 236-242.
doi:10.1080/00063657.2012.662939
Gatesoupe, F. J. (1999). The use of probiotics in aquaculture. *Aquaculture, 180*(1), 147-165.
doi:https://doi.org/10.1016/S0044-8486(99)00187-8
Gillingham, M. A. F., Béchet, A., Cézilly, F., Wilhelm, K., Rendón-Martos, M., Borghesi, F., . . .
Sommer, S. (2019). Offspring Microbiomes Differ Across Breeding Sites in a
Panmictic Species. *Frontiers in Microbiology, 10*, 35. Retrieved from
https://www.frontiersin.org/article/10.3389/fmicb.2019.00035
Gomes, B. C., Rodrigues, M. R., Winkelströter, L. K., Nomizo, A., & de Martinis, E. C. (2012).
In vitro evaluation of the probiotic potential of bacteriocin producer Lactobacillus
sakei 1. *J Food Prot, 75*(6), 1083-1089. doi:10.4315/0362-028x.Jfp-11-523
Götmark, F. (2002). Predation by sparrowhawks favours early breeding and small broods in
great tits. *Oecologia, 130*(1), 25-32. doi:10.1007/s004420100769
Hansen, C. H. F., Nielsen, D. S., Kverka, M., Zakostelska, Z., Klimesova, K., Hudcovic, T., . . .
Hansen, A. K. (2012). Patterns of Early Gut Colonization Shape Future Immune
Responses of the Host. *PLOS ONE, 7*(3), e34043. doi:10.1371/journal.pone.0034043
Hartig, F. (2019). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. In.
Heintz-Buschart, A., & Wilmes, P. (2018). Human Gut Microbiome: Function Matters. *Trends
in Microbiology, 26*(7), 563-574. doi:https://doi.org/10.1016/j.tim.2017.11.002
Highet, A. R., Berry, A. M., Bettelheim, K. A., & Goldwater, P. N. (2014). Gut microbiome in
sudden infant death syndrome (SIDS) differs from that in healthy comparison babies
and offers an explanation for the risk factor of prone position. *Int J Med Microbiol,
304*(5-6), 735-741. doi:10.1016/j.ijmm.2014.05.007
Holzapfel, W. H., & Schillinger, U. (2002). Introduction to pre- and probiotics. *Food Research
International, 35*(2), 109-116. doi:https://doi.org/10.1016/S0963-9969(01)00171-5

Hooper, L. V., Littman, D. R., & Macpherson, A. J. (2012). Interactions between the
microbiota and the immune system. *Science, 336*(6086), 1268-1273.
doi:10.1126/science.1223490
Hõrak, P., Lea, T., Ots, I., & Møller, A. P. (1999). Immune Function and Survival of Great Tit
Nestlings in Relation to Growth Conditions. *Oecologia, 121*(3), 316-322. Retrieved
from www.jstor.org/stable/4222472
Keller, L. F., & Van Noordwijk, A. J. (1994). Effects of Local Environmental-Conditions on
Nestling Growth in the Great Tit Parus-Major. *Ardea, 82*(2), 349-362.
Khan, G., Kangro, H. O., Coates, P. J., & Heath, R. B. (1991). Inhibitory effects of urine on the
polymerase chain reaction for cytomegalovirus DNA. *J Clin Pathol, 44*(5), 360-365.
doi:10.1136/jcp.44.5.360
Kinross, J. M., Darzi, A. W., & Nicholson, J. K. (2011). Gut microbiome-host interactions in
health and disease. *Genome Med, 3*(3), 14. doi:10.1186/gm228
Knowles, S. C. L., Eccles, R. M., & Baltrūnaitė, L. (2019). Species identity dominates over
environment in shaping the microbiota of small mammals. *Ecology Letters, 22*(5),
826-837. doi:10.1111/ele.13240
Koenig, J. E., Spor, A., Scalfone, N., Fricker, A. D., Stombaugh, J., Knight, R., . . . Ley, R. E.
(2010). Succession of microbial consortia in the developing infant gut microbiome.
Proceedings of the National Academy of Sciences, 108(Supplement_1), 4578-4585.
Retrieved from http://dx.doi.org/10.1073/pnas.1000081107
Kohl, K. D., Brun, A., Bordenstein, S. R., Caviedes-Vidal, E., & Karasov, W. H. (2018). Gut
microbes limit growth in house sparrow nestlings (Passer domesticus) but not
through limitations in digestive capacity. (1749-4877 (Electronic)).
Lan, Y., Verstegen, M. W. A., Tamminga, S., & Williams, B. A. (2005). The role of the
commensal gut microbial community in broiler chickens. *World's Poultry Science
Journal, 61*(01), 95-104. Retrieved from http://dx.doi.org/10.1079/WPS200445
Le Chatelier, E., Nielsen, T., Qin, J., Prifti, E., Hildebrand, F., Falony, G., . . . Meta, H. I. T. c.
(2013). Richness of human gut microbiome correlates with metabolic markers.
Nature, 500(7464), 541-546. doi:10.1038/nature12506
LeBlanc, J. G., Chain, F., Martín, R., Bermúdez-Humarán, L. G., Courau, S., & Langella, P.
(2017). Beneficial effects on host energy metabolism of short-chain fatty acids and
vitamins produced by commensal and probiotic bacteria. *Microbial cell factories,
16*(1), 79-79. doi:10.1186/s12934-017-0691-z
Liu, H., Wang, J., He, T., Becker, S., Zhang, G., Li, D., & Ma, X. (2018). Butyrate: A Double-
Edged Sword for Health? *Advances in Nutrition, 9*(1), 21-29.
doi:10.1093/advances/nmx009
Lloyd-Price, J., Abu-Ali, G., & Huttenhower, C. (2016). The healthy human microbiome.
Genome Medicine, 8(1), 51. doi:10.1186/s13073-016-0307-y
Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K., & Knight, R. (2012). Diversity,
stability and resilience of the human gut microbiota. *Nature, 489*(7415), 220-230.
doi:10.1038/nature11550
Martin, T. E. (2015). Age-related mortality explains life history strategies of tropical and
temperate songbirds. *Science, 349*(6251), 966. doi:10.1126/science.aad1173
Maurice, C. F., Cl Knowles, S., Ladau, J., Pollard, K. S., Fenton, A., Pedersen, A. B., &
Turnbaugh, P. J. (2015). Marked seasonal variation in the wild mouse gut microbiota.
The ISME Journal, 9(11), 2423-2434. Retrieved from
http://dx.doi.org/10.1038/ismej.2015.53

McKnight, D. T., Huerlimann, R., Bower, D. S., Schwarzkopf, L., Alford, R. A., & Zenger, K. R.
(2019). Methods for normalizing microbiome data: An ecological perspective.
Methods in Ecology and Evolution, 10(3), 389-400. Retrieved from
https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.13115
McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLOS ONE, 8*(4), e61217-e61217.
Retrieved from http://dx.doi.org/10.1371/journal.pone.0061217
Merila, J. (1996). Genetic Variation in Offspring Condition: An Experiment. *Functional
Ecology, 10*(4), 465-474. doi:10.2307/2389939
Mikaela, A. J. B., Mark, G. M., Robertson, D. R., & McCormick, M. I. (2002). Larval Growth
Predicts the Recruitment Success of a Coral Reef Fish. *Oecologia, 131*(4), 521-525.
Retrieved from http://www.jstor.org/stable/4223286
Monrós, J. S., Belda, E. J., & Barba, E. (2002). Post-fledging survival of individual great tits:
the effect of hatching date and fledging mass. *Oikos, 99*(3), 481-488.
doi:10.1034/j.1600-0706.2002.11909.x
Motiei, A., Brindefalk, B., Ogonowski, M., El-Shehawy, R., Pastuszek, P., Ek, K., . . .
Gorokhova, E. (2020). Disparate effects of antibiotic-induced microbiome change
and enhanced fitness in Daphnia magna. *PLOS ONE, 15*(1), e0214833.
doi:10.1371/journal.pone.0214833
Norris, K. (1993). Heritable variation in a plumage indicator of viability in male great tits
Parus major. *Nature, 362*(6420), 537-539. doi:10.1038/362537a0
O'Hara, A. M., & Shanahan, F. (2006). The gut flora as a forgotten organ. *EMBO reports, 7*(7),
688-693. doi:10.1038/sj.embor.7400731
O'Shea, W., O'Halloran, J., & Quinn, J. L. (2018). Breeding phenology, provisioning
behaviour, and unusual patterns of life history variation across an anthropogenic
heterogeneous landscape. *Oecologia, 188*, 953–964
Oksanen, J., F. Guillaume Blanchet, Michael Friendly, Roeland Kindt, Pierre Legendre, Dan
McGlinn, . . . Wagner, H. (2019). vegan: Community Ecology Package (Version R
package version 2.5-6).
Olszak, T., An, D., Zeissig, S., Vera, M. P., Richter, J., Franke, A., . . . Blumberg, R. S. (2012).
Microbial exposure during early life has persistent effects on natural killer T cell
function. *Science (New York, N.Y.), 336*(6080), 489-493.
doi:10.1126/science.1219328
Patrick, S. C., & Weimerskirch, H. (2014). Personality, foraging and fitness consequences in a
long lived seabird. *PLOS ONE, 9*(2), e87269-e87269.
doi:10.1371/journal.pone.0087269
Patterson, J. A., Orban, J. I., Sutton, A. L., & Richards, G. N. (1997). Selective enrichment of
bifidobacteria in the intestinal tract of broilers by thermally produced kestoses and
effect on broiler performance. *Poultry Science, 76*(3), 497-500.
doi:https://doi.org/10.1093/ps/76.3.497
Pickett, S. R. A., Weber, S. B., McGraw, K. J., Norris, K. J., & Evans, M. R. (2013).
Environmental and Parental Influences on Offspring Health and Growth in Great Tits
(Parus major). *PLOS ONE, 8*(7), e69695. doi:10.1371/journal.pone.0069695
Potti, J., Moreno, J., Yorio, P., Briones, V., García-Borboroglu, P., Villar, S., & Ballesteros, C.
(2002). Bacteria divert resources from growth for magellanic penguin chicks. *Ecology
Letters, 5*(6), 709-714. doi:10.1046/j.1461-0248.2002.00375.x

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., . . . Glöckner, F. O. (2012).
The SILVA ribosomal RNA gene database project: improved data processing and
web-based tools. *Nucleic Acids Research, 41*(D1), D590-D596. Retrieved from
https://doi.org/10.1093/nar/gks1219
R Core Team. (2019). R: A language and environment for statistical computing. (Version
3.5.1). Vienna, Austria: R Foundation for Statistical Computing.
Ranstam, J. (2019). Hypothesis-generating and confirmatory studies, Bonferroni correction,
and pre-specification of trial endpoints. *Acta Orthopaedica, 90*(4), 297-297.
doi:10.1080/17453674.2019.1612624
Ratajczak, W., Rył, A., Mizerski, A., Walczakiewicz, K., Sipak, O., & Laszczyńska, M. (2019).
Immunomodulatory potential of gut microbiome-derived short-chain fatty acids
(SCFAs). *Acta Biochim Pol, 66*(1), 1-12. doi:10.18388/abp.2018_2648
Rauw, W. M. (2012). Immune response from a resource allocation perspective. *Frontiers in*
genetics, 3, 267-267. doi:10.3389/fgene.2012.00267
Ringsby, T. H., Sæther, B.-E., & Solberg, E. J. (1998). Factors Affecting Juvenile Survival in
House Sparrow Passer domesticus. *Journal of Avian Biology, 29*(3), 241-247.
doi:10.2307/3677106
Rosshart, S. P., Vassallo, B. G., Angeletti, D., Hutchinson, D. S., Morgan, A. P., Takeda, K., . . .
Rehermann, B. (2017). Wild Mouse Gut Microbiota Promotes Host Fitness and
Improves Disease Resistance. *Cell, 171*(5), 1015-1028.e1013. Retrieved from
http://www.sciencedirect.com/science/article/pii/S0092867417310656
Rowland, I., Gibson, G., Heinken, A., Scott, K., Swann, J., Thiele, I., & Tuohy, K. (2018). Gut
microbiota functions: metabolism of nutrients and other food components. *Eur J*
Nutr, 57(1), 1-24. doi:10.1007/s00394-017-1445-8
Schwagmeyer, P. L., & Mock, D. W. (2008). Parental provisioning and offspring fitness: size
matters. *Animal Behaviour, 75*(1), 291-298.
doi:https://doi.org/10.1016/j.anbehav.2007.05.023
Shannon, C. E. (1948). A Mathematical Theory of Communication. *Bell System Technical*
Journal, 27(3), 379-423. doi:10.1002/j.1538-7305.1948.tb01338.x
Sherwin, E., Bordenstein, S. R., Quinn, J. L., Dinan, T. G., & Cryan, J. F. (2019). Microbiota and
the social brain. *Science, 366*(6465), eaar2016. doi:10.1126/science.aar2016
Shutt, J. D., Nicholls, J. A., Trivedi, U. H., Burgess, M. D., Stone, G. N., Hadfield, J. D., &
Phillimore, A. B. (2020). Gradients in richness and turnover of a forest passerine's
diet prior to breeding: A mixed model approach applied to faecal metabarcoding
data. *Molecular Ecology, 29*(6), 1199-1213. doi:10.1111/mec.15394
Song, S. J., Sanders, J. G., Delsuc, F., Metcalf, J., Amato, K., Taylor, M. W., . . . Knight, R.
(2020). Comparative Analyses of Vertebrate Gut Microbiomes Reveal Convergence
between Birds and Bats. *mBio, 11*(1), e02901-02919. doi:10.1128/mBio.02901-19
Spencer, K. A., Evans, N. P., & Monaghan, P. (2009). Postnatal Stress in Birds: A Novel Model
of Glucocorticoid Programming of the Hypothalamic-Pituitary-Adrenal Axis.
Endocrinology, 150(4), 1931-1934. doi:10.1210/en.2008-1471
Stige, L. C., Rogers, L. A., Neuheimer, A. B., Hunsicker, M. E., Yaragina, N. A., Ottersen, G., . . .
Durant, J. M. (2019). Density- and size-dependent mortality in fish early life stages.
Fish and Fisheries, 20(5), 962-976. doi:https://doi.org/10.1111/faf.12391
Sudo, N., Chida, Y., Aiba, Y., Sonoda, J., Oyama, N., Yu, X.-N., . . . Koga, Y. (2004). Postnatal
microbial colonization programs the hypothalamic-pituitary-adrenal system for

stress response in mice. *The Journal of Physiology, 558*(1), 263-275. Retrieved from
http://dx.doi.org/10.1113/jphysiol.2004.063388

Suzuki, T. A. (2017). Links between Natural Variation in the Microbiome and Host Fitness in
Wild Mammals. *Integrative and Comparative Biology, 57*(4), 756-769.
doi:https://doi.org/10.1093/icb/icx104

Teyssier, A., Lens, L., Matthysen, E., & White, J. (2018). Dynamics of Gut Microbiota Diversity
During the Early Development of an Avian Host: Evidence From a Cross-Foster
Experiment. *Frontiers in Microbiology, 9*, 1524.
doi:https://doi.org/10.3389/fmicb.2018.01524

Teyssier, A., Rouffaer, L. O., Hudin, N. S., Strubbe, D., Matthysen, E., Lens, L., & White, J.
(2018). Inside the guts of the city: Urban-induced alterations of the gut microbiota in
a wild passerine. *Science of The Total Environment, 612*, 1276-1286. Retrieved from
http://www.sciencedirect.com/science/article/pii/S0048969717323781

Therneau, T. M. (2020). coxme: Mixed Effects Cox Models (Version R package version 2.2-
16). Retrieved from https://CRAN.R-project.org/package=coxme

Torok, V. A., Hughes, R. J., Mikkelsen, L. L., Perez-Maldonado, R., Balding, K., MacAlpine, R., .
. . Ophel-Keller, K. (2011). Identification and Characterization of Potential
Performance-Related Gut Microbiotas in Broiler Chickens across Various Feeding
Trials. *Applied and Environmental Microbiology, 77*(17), 5868-5878. Retrieved from
http://dx.doi.org/10.1128/AEM.00165-11

Trevelline, B. K., Fontaine, S. S., Hartup, B. K., & Kohl, K. D. (2019). Conservation biology
needs a microbial renaissance: a call for the consideration of host-associated
microbiota in wildlife management practices. *Proceedings of the Royal Society B:*
Biological Sciences, 286(1895), 20182448. doi:10.1098/rspb.2018.2448

Turnbaugh, P. J., Ley, R. E., A., M. M., & et al. (2006). An obesity-associated gut microbiome
with increased capacity for energy harvest. *Nature, 444*, 1027-1031.

Van Noordwijk, A. J., McCleery, R. H., & Perrins, C. M. (1995). Selection for the Timing of
Great Tit Breeding in Relation to Caterpillar Growth and Temperature. *Journal of*
Animal Ecology, 64(4), 451-458. doi:10.2307/5648

Vargas-Pellicer, P., Watrobska, C., Knowles, S., Schroeder, J., & Banks-Leite, C. (2019). How
should we store avian faecal samples for microbiota analyses? Comparing efficacy
and cost-effectiveness. *Journal of Microbiological Methods, 165*, 105689.
doi:https://doi.org/10.1016/j.mimet.2019.105689

Vásquez, A., Forsgren, E., Fries, I., Paxton, R. J., Flaberg, E., Szekely, L., & Olofsson, T. C.
(2012). Symbionts as Major Modulators of Insect Health: Lactic Acid Bacteria and
Honeybees. *PLOS ONE, 7*(3), e33188. doi:10.1371/journal.pone.0033188

Videvall, E., Song, S. J., Bensch, H. M., Strandh, M., Engelbrecht, A., Serfontein, N., . . .
Cornwallis, C. K. (2019). Major shifts in gut microbiota during development and its
relationship to growth in ostriches. *Molecular Ecology, 28*(10), 2653-2667.
doi:10.1111/mec.15087

Videvall, E., Song, S. J., Bensch, H. M., Strandh, M., Engelbrecht, A., Serfontein, N., . . .
Cornwallis, C. K. (2020). Early-life gut dysbiosis linked to juvenile mortality in
ostriches. *Microbiome, 8*(1), 147. doi:10.1186/s40168-020-00925-7

Videvall, E., Strandh, M., Engelbrecht, A., Cloete, S., & Cornwallis, C. K. (2017). Measuring
the gut microbiome in birds: Comparison of faecal and cloacal sampling. *Molecular*
Ecology Resources, 18(3), 424-434. Retrieved from http://dx.doi.org/10.1111/1755-
0998.12744

Vispo, C., & Karasov, W. H. (1997). The Interaction of Avian Gut Microbes and Their Host: An
944   Elusive Symbiosis. In R. I. Mackie & B. A. White (Eds.), (pp. 116-155): Springer US.
945   Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naive Bayesian classifier for rapid
946   assignment of rRNA sequences into the new bacterial taxonomy. *Applied and*
947   *Environmental Microbiology, 73*(16), 5261–5267. doi:http://10.1128/AEM.00062-07
948   Warnecke, F., Luginbühl, P., Ivanova, N., Ghassemian, M., Richardson, T. H., Stege, J. T., . . .
949   Leadbetter, J. R. (2007). Metagenomic and functional analysis of hindgut microbiota
950   of a wood-feeding higher termite. *Nature, 450*(7169), 560-565.
951   doi:10.1038/nature06269
952   Wasielewski, H., Alcock, J., & Aktipis, A. (2016). Resource conflict and cooperation between
953   human host and gut microbiota: implications for nutrition and health. *Annals of the*
954   *New York Academy of Sciences, 1372*(1), 20-28. doi:10.1111/nyas.13118
955   Weber, B. M., Bowers, E. K., Terrell, K. A., Falcone, J. F., Thompson, C. F., & Sakaluk, S. K.
956   (2018). Pre- and postnatal effects of experimentally manipulated maternal
957   corticosterone on growth, stress reactivity and survival of nestling house wrens.
958   *Functional Ecology, 32*(8), 1995-2007. doi:10.1111/1365-2435.13126
959   West, A. G., Waite, D. W., Deines, P., Bourne, D. G., Digby, A., McKenzie, V. J., & Taylor, M.
960   W. (2019). The microbiome in threatened species conservation. *Biological*
961   *Conservation, 229*, 85-98. doi:https://doi.org/10.1016/j.biocon.2018.11.016
962   Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*: Springer-Verlag New York.
963   Youngblut, N. D., Reischer, G. H., Walters, W., Schuster, N., Walzer, C., Stalder, G., . . .
964   Farnleitner, A. H. (2019). Host diet and evolutionary history explain different aspects
965   of gut microbiome diversity among vertebrate clades. *Nature Communications,*
966   *10*(1), 2200. doi:10.1038/s41467-019-10191-3
967   Zaneveld, J. R., McMinds, R., & Vega Thurber, R. (2017). Stress and stability: applying the
968   Anna Karenina principle to animal microbiomes. *Nature Microbiology, 2*(9), 17121.
969   doi:10.1038/nmicrobiol.2017.121
970   Zhu, L., Wu, Q., Dai, J., Zhang, S., & Wei, F. (2011). Evidence of cellulose metabolism by the
971   giant panda gut microbiome. *Proceedings of the National Academy of Sciences,*
972   *108*(43), 17714. doi:10.1073/pnas.1017956108
973   Zimmer, C., Boogert, N. J., & Spencer, K. A. (2013). Developmental programming:
974   Cumulative effects of increased pre-hatching corticosterone levels and post-hatching
975   unpredictable food availability on physiology and behaviour in adulthood. *Hormones*
976   *and Behavior, 64*(3), 494-500. doi:https://doi.org/10.1016/j.yhbeh.2013.07.002
977