Novel *Chlamydiae* and *Amoebophilus* endosymbionts are prevalent in wild isolates of the model social amoebae *Dictyostelium discoideum*

Authors:

Tamara S. Haselkorn[1], Daniela Jimenez[2], Usman Bashir[2], Eleni Sallinger[1], David C. Queller[2], Joan E. Strassmann[2], Susanne DiSalvo[3]

Affiliations:

[1]Department of Biology, University of Central Arkansas, 201 Donaghey Avenue, Conway, AR 72035

[2]Department of Biology, Washington University in St. Louis, One Brookings Drive St. Louis, MO 63130

[3]Department of Biological Sciences, Southern Illinois University Edwardsville, 44 Circle Drive, Edwardsville, IL 62026

* Corresponding author

Department of Biological Sciences, Southern Illinois University Edwardsville, 44 Circle Drive, Edwardsville, IL 62026; sdisalv@siue.edu

Running Title: Prevalent obligate endosymbionts in social amoebae

Significance statement: In a large survey across the eastern US, we identify the presence and prevalence of novel strains of obligate intracellular symbionts from within the *Chlamydiae*, *Amoebophilus*, and *Procabacter* lineages in natural populations of the social amoeba *Dictyostelium discoideum.* We demonstrate that these symbionts do not affect amoeba fitness under standard laboratory conditions. High prevalences in natural populations and some non-random co-infection patterns suggest that these symbionts have environmental context-dependent effects. The presence of these novel bacterial strains that are highly diverged from known amoeba endosymbionts provides an opportunity to study host specificity and host-symbiont interactions in the well-studied *D. discoideum* model organism. It opens the door for exploring the evolutionary history of these widespread obligate symbionts that are associated with a large variety of hosts.

Summary: Amoebae interact with bacteria in diverse and multifaceted ways. Amoeba predation can serve as a selective pressure for the development of bacterial virulence traits. Bacteria may also adapt to life inside amoebae, resulting in symbiotic relationships (pathogenic or mutualistic). Indeed, amoebae are often infected with bacterial endosymbionts. *Acanthamoeba* and *Hartmannella* harbor symbionts from five distinct lineages within the *Proteobacteria*, *Bacteroidetes*, and *Chlamydiae*. Here, we PCR-screened an extensive collection of *Dictyostelium discoideum* wild isolates for the presence of bacterial symbionts. For the first time we found that obligate symbionts are surprisingly common in this highly-studied amoeba species, identified in 42% of screened isolates (N=730). Members of the environmental *Chlamydiae* are particularly prevalent. These strains are novel and phylogenetically distinct. We also found *Amoebophilus* symbionts in 8% of screened isolates (N=730). Under laboratory conditions, antibiotic-cured amoebae behave similarly to their endosymbiont-infected counterparts, suggesting that endosymbionts do not significantly impact host fitness, at least in the laboratory. We found several natural isolates were co-infected with different endosymbionts. The high prevalence and novelty of amoeba endosymbiont clades in the model organism *D. discoideum* opens the door to future research on the significance and mechanisms of amoeba-symbiont interactions.

Introduction:

As voracious microbial predators, free-living amoebae are important shapers of their microbial communities (Clarholm, 1981). This predatory pressure can alter the presence and abundance of specific microbial constituents in the community (Ronn et al., 2002).

Amoeba predation is also postulated to play an important role in microbial evolution, particularly in the evolution of bacterial virulence (Matz and Kjelleberg, 2005; Sun et al., 2018). Bacteria that evolve strategies to avoid or survive amoeba predation would be selected for in amoeba rich environments (Erken et al., 2013). Thus, amoebae may be training grounds for microbial agents to develop virulence traits that allow them to counter or exploit these predators (Barker and Brown, 1994; Cosson and Soldati, 2008; Molmeret et al., 2005). Generalized virulence strategies affect other eukaryotes (such as humans), resulting in the evolution of important bacterial pathogens. Indeed, it has been posited that amoebae may be a missing link in the transition of environmental microbes into pathogens (Taylor-Mulneix et al., 2017).

Amoebae are single-celled eukaryotes that are phylogenetically divergent, but share an amorphous, motile life form. They move via pseudopodia and consume bacteria through phagocytic ingestion and subsequent enzymatic degradation. Some bacteria are susceptible to phagocytosis by amoebae but are able to avoid or withstand subsequent digestion. These are referred to as "amoeba resisting bacteria" and include many important human pathogenic species including *Mycobacteria, Legionella,* and *Chlamydia* (Boamah et al., 2017; Cardenal-Muñoz et al., 2018; Drancourt, 2014; Gomez-Valero and Buchrieser, 2019; Paquet and Charette, 2016; Tosetti et al., 2014). A diverse collection of intracellular bacterial symbionts of amoebae has been found, some of which appear pathogenic, neutral, or beneficial for their amoeba host (DiSalvo et al., 2015; Jeon, 1992; Shu et al., 2018; Taylor et al., 2012).

Bacterial endosymbionts of amoebae were first discovered in the diverse, free-living

amoeba genus *Acanthamoeba* (Horn and Wagner, 2004). These amoebae are found in

soil and freshwater environments and can protect themselves and their bacterial

associates by forming cysts under environmentally unfavorable conditions (Rodríguez-

Zaragoza, 1994). The presence of such endosymbionts was known long before they

could be specifically identified because the inability to isolate obligate associates from

their hosts made characterization challenging. Once molecular methods became

available, it was clear that these bacteria were from specific lineages in the *Chlamydiae*,

Proteobacteria, and *Bacteriodetes* (Horn and Wagner, 2004; Schmitz-Esser et al.,

2008). The potential of amoebae to harbor such bacteria could be of concern to human

health, because environmental *Chlamydiae* have been associated with respiratory

diseases and adverse pregnancy outcomes in humans and cattle, although the extent of

their importance in causing animal illness is unknown (Greub, 2009; Taylor-Brown et al.,

2015).

Amoeba endosymbionts interact with their hosts in varied ways. Some inhabit host-

derived vacuoles, while others persist in the cytoplasm. The most common

endosymbionts are from the order *Chamydiales* within the environmental *Chlamydiae*

group. One symbiont, *Neochlamydia*, can protect its host against the detrimental

consequences of *Legionella* bacterial infection (Ishida et al., 2014; Maita et al., 2018)

and other strains have been shown to improve the growth rate and motility of their

amoeba hosts (Okude et al., 2012). Other known amoeba endosymbionts are from the

less-studied genera *Amoebophilus* and *Procabacter*. Genome sequencing of the

amoeba symbiont *Candidatus Amoebophilus asiaticus* found no evidence that it could provide novel pathways to supplement host nutrition, but other fitness affects have not been tested (Schmitz-Esser et al., 2010).

While many endosymbiont-infected *Acanthamoeba* isolates were singly infected, more recent screening and alternative culturing methods have shown several instances of endosymbiont co-infection (Heinz et al., 2007; Lagkouvardos et al., 2014a). The same bacterial lineages have been found in *Acanthamoeba* sampled globally, and although there is diversity within these lineages, the amount of co-evolution between specific amoebae and bacteria is less clear. Tests of host specificity have shown variable results, with some endosymbiotic bacteria able to infect multiple strains of *Acanthamoeba*, while others have difficulty infecting new hosts (Okude et al., 2012). A high proportion of sampled *Acanthamoeba* (25-80%) are infected with endosymbionts, but the prevalence and variance of infection across populations is unclear due to the challenges of isolating large numbers of amoebae (Schmitz-Esser et al., 2008). Overall, persistent symbionts of *Acanthamoeba* serve as useful models for understanding how amoebae can serve as training grounds for bacterial intracellular adaptation, but the broader distribution of symbiotic bacteria among other amoeba species is unknown.

Another amoeba that has been used as a model to study host-bacteria interactions is the social amoeba, *Dictyostelium discoideum*. This amoeba, distantly related to *Acanthamoeba*, is terrestrial, living and feeding as single cells in forest soils while the bacterial food supply is plentiful. When food becomes scarce, the amoebae begin a

social cycle, which entails thousands of cells aggregating and forming a slug to crawl to the surface of the soil. The cycle culminates in the formation of a fruiting body, made of a ball of spores (sorus) sitting on top of a stalk, which can then be dispersed to more favorable conditions (Kessin, 2001). In many cases, bacteria are cleared from *D. discoideum* cells during the social cycle through innate immune-like mechanisms (Chen et al., 2007). However, vegetative amoebae can be infected with a range of human pathogens such as *Mycobacterium, Bordetella*, and *Legionella* (Bozzaro and Eichinger, 2011; Taylor-Mulneix et al., 2017). *D. discoideum* has thereby been fruitfully utilized to describe the role of host and pathogen factors in mediating infection mechanisms and virulence (Annesley and Fisher, 2009; Bozzaro et al., 2019; Bozzaro and Eichinger, 2011; Cosson and Soldati, 2008; Steinert, 2011; Steinert and Heuner, 2005). Because some of these pathogens survive phagocytosis and even persist in fruiting bodies, it is possible that *D. discoideum* is an environmental reservoir for different types of human pathogenic bacteria (Taylor-Mulneix et al., 2017). Whether these bacterial pathogens commonly infect *Dictyostelium* in the wild is unclear (Amissah et al., 2014; Drancourt, 2014; Salah et al., 2009).

Recently, a variety of bacterial symbionts have been discovered to colonize *D. discoideum* (Brock et al., 2018; DiSalvo et al., 2015; Haselkorn et al., 2018). The *D. discoideum* microbiome is composed of both persistent and transient bacterial associates (Brock et al., 2018). Three different newly named *Paraburkholderia* species, *P. agricolaris*, *P. hayleyella*, and *P. bonniea* are persistent bacterial symbionts that intracellularly infect vegetative amoebae and spore cells (Brock et al., 2020, 2011;

Haselkorn et al., 2018; Khojandi et al., 2019; Shu et al., 2018). While these *Paraburkholderia* themselves do not nourish *D. discoideum*, they confer farming to their hosts by allowing food bacteria to survive in the fruiting body. These food bacteria can then be re-seeded in new environments during spore dispersal, which benefits hosts in food-scarce conditions (Brock et al., 2011; Haselkorn et al., 2018; Khojandi et al., 2019). In natural populations, an average of 25% of *D. discoideum* are infected with *Paraburkholderia* (Haselkorn et al., 2018). Other bacteria, some edible and others not, have the ability to transiently associate with *D. discoideum* even in the absence of *Paraburkholderia* co-infections (Brock et al., 2018). Thus, natural *D. discoideum* hosts a small-scale microbiome and is gaining traction as a simple model system to study the mechanisms of microbiome formation (Dinh et al., 2018; Farinholt et al., 2019). However, all previous sampling of natural *D. discoideum* associates has relied on isolating bacterial colonies from nutrient media plates, which would miss any bacterial species, such as obligate amoeba symbionts, incapable of growing under these conditions.

Since obligate endosymbionts are common in other free-living amoebae, we endeavored to identify them in *D. discoideum.* Here, we screened a frozen stock collection of over 700 natural *D. discoideum* isolates. We first used a 16s rRNA gene amplification strategy with "universal" primers and direct sequencing to detect and identify bacteria within *D. discoideum* spores in a single population. In addition to detecting expected *Paraburkholderia* symbionts, we identified many *Chlamydiae* and *Amoebophilus* sequences*,* and a few *Procabacter* sequences in screened samples. To

determine the extent of *Chlamydiae* and *Amoebophilus* symbionts in our collection we then used symbiont specific PCR screening. From each of these prevalent bacterial lineages, we found genetically distinct novel species and were able to visualize representative endosymbiont cells within spores via electron microscopy. We also used antibiotics to cure natural hosts of *Chlamydiae* and *Amoebophilus* and compared host fitness to uncured counterparts. Finally, we looked at the fitness effects of co-infections since co-infections occur in nature.

Results:

Obligate endosymbionts are common in wild-collected clones of Dictyostelium discoideum

To investigate the presence and prevalence of obligate bacterial symbionts in *D. discoideum*, we screened a frozen collection of 730 natural *D. discoideum* isolates for bacterial DNA. The vast majority of our screened isolates were collected from the Eastern half of the United States (Fig. 1a) and had previously been screened for *Paraburkholderia* (Haselkorn, *et al.* 2018, Table S1). This included 16 populations with seven or more individuals collected. Our largest collection from Virginia consists of about 200 individuals collected in 2000 and in 2014. Across all locations we found over 41% of samples harbored at least one obligate endosymbiont. Specifically, 117 (~16%) were infected with *Amoebophilus* and 198 (~27%) with *Chlamydiae.* Sixteen (2.1%) were co-infected with both *Amoebophilus* and *Chlamydiae* (Fig. 1b). Two isolates (both from Virginia) were infected with an obligate bacterium identified as belonging to the genus *Procabacter*.

For *Procabacter*, *Amoebophilus*, and the three most prevalent *Chamydiae* strains (those

occurring in more than one individual) we sequenced the full length 16S rRNA for our *D.*

discoideum endosymbionts to better determine their evolutionary relationships. In all

cases, the 16S rRNA haplotypes were novel and highly diverged from other sequences

currently in GenBank. For the *Procabacter* and *Amoebophilus* endosymbionts, only a

single haplotype was found for each. The *Procabacter* endosymbiont of *D. discoideum*

is sister to all of the *Procabacter* endosymbionts of *Acanthamoeba* (Fig. 2a). While it is

most closely related to this genus, forming a clade with 100% bootstrap support, the

16S rRNA sequence is only a 93% match. A similar pattern is seen for the

Amoebophilus symbiont of *D. discoideum* (Fig. 2b.). The *Amoebophilus* endosymbiont

that infects *D. discoideum* is sister to the *Amoebophilus asiaticus* endosymbionts of

Acanthamoeba, and is 91% diverged at the 16S rRNA gene.

There are 8 distinct haplotypes of *D. discoideum* endosymbionts in the *Chlamydiae*. The

phylogeny based on the 220bp 16S rRNA fragment-generated screening assay

suggests four distinct groups, although bootstrap support is low (Fig. 2c). Haplotypes 1

and 2 form a well-supported clade with two additional haplotypes, while haplotypes 3, 6,

7, and 8 fall elsewhere on the tree. For the three most prevalent haplotypes (1, 2, and

3), we sequenced the full 16S rRNA gene to reconstruct a more resolved phylogeny

(Fig. 2d). There are currently 9 named *Chlamydiae* families, and the *D. discoideum*

Chamydiae endosymbiont does not fall into any of these clades (Taylor-Brown et al., 2015). Haplotypes 1 and 2 form their own well-supported clade, sister to the *Rhabdochlamydiaceae* family. This highly diverged clade (89% divergent at the sequence level) may represent a new family of *Chamydiae,* as greater than 10% sequence divergence at the full length 16S rRNA gene has been proposed as family level difference (Everett et al., 1999). The phylogenetic position of haplotype 3 is less well-resolved, with weak bootstrap support placing it sister to several *Chlamydiae* families. This highly divergent strain my represent a novel family as well.

Distribution of endosymbionts across populations

While the average overall prevalence of infection of *Chlamydiae* endosymbionts was 27%, it varies among populations, ranging from 0% to 75%, with a variable distribution of haplotypes. The *D. discoideum Chlamydiae* haplotypes 1, 2 and 3 were the most prevalent in natural populations. Haplotype 1 represented 79.8% of all *D. discoideum Chlamydiae* infections, while haplotype 2 was 13.6% and haplotype 3 was 3.5%. Haplotypes 4-8 were represented only once. Haplotype 1 was found in 12 populations, haplotype 2 was found in 8 different populations, and haplotype 3 was found in three populations. The diversity of *Chlamydiae* strains was the highest in the Virginia Mountain Lake collection from 2000, which had six of the eight haplotypes (1-6), and the second highest diversity of *Chlamydiae* haplotypes was found in the Texas Houston Arboretum collection (1, 2, 3 and 8). Interestingly, haplotype 1 was the only haplotype circulating in the most recent 2014 Virginia Mountain Lake collection. While haplotype

diversity was lower in this collection, *Chlamydiae* prevalence (for all haplotypes) in the region increased from 20% in 2000 to 38% in 2014.

The one strain of *Amoebophilus* found to infect *D. discoideum* was distributed across 9 populations, although at low prevalence, often with just a single individual infected in many locations. This endosymbiont was generally less prevalent than *Chlamydiae*, with the exception of the 2000 Virginia Mountain Lake population, where it infected 37% of the population. *Amoebophilus* prevalence in this region, however, decreased to 10% in 2014. *Amoebophilus* was also relatively prevalent in the geographically-distant Texas Houston Arboretum population, infecting 23% of the population.

Obligate endosymbionts can be visualized within reproductive host spore cells

Transmission electron microscopy (TEM) revealed bacterial cells inside developed spores from representative *Amoebophilus, Chlamydiae,* and *Procabacter* host isolates (Fig. 3). Symbiont cells were detectable in the majority of symbiont host spores visualized by TEM. Each symbiont appeared to have gram-negative type cell walls but displayed unique morphologies within spore cells. *Amoebophilus* endosymbionts were rod-like, measured 0.3-0.5μm in width and 0.5-1.3μm in length, and appeared to be distributed throughout the cytoplasm within a host membrane. *Procabacter* endosymbionts were rod shaped, measured 0.25-0.5μm in width and 0.75-1.5μm in length, and in many cases were surrounded by a host membrane. *Chlamydiae* endosymbionts appeared as dense wrinkled spheres measuring 0.3-0.6μm that were distributed throughout the cytoplasm.

Amoebophilus and Chlamydiae *can be eliminated from host isolates without significantly impacting host fitness*

We did not observe any obvious defects in the growth and development of *Chlamydiae* and *Amoebophilus* infected hosts. To better assess the impact of these endosymbionts on their hosts, we measured the reproductive fitness of representative host isolates before and after endosymbiont curing (Fig. 4). We eliminated *Chlamydiae* and *Amoebophilus* from four of their natural hosts through culturing for two social cycles on antibiotic saturated plates. After antibiotic exposure and resumption of development on normal media, we confirmed stable loss of endosymbionts via endosymbiont-specific PCR screening (Fig. 4a). We next cultured $1x10^5$ spores from antibiotic-treated and untreated *D. discoideum* counterparts on nutrient media with *Klebsiella pneumoniae* food bacteria. Following culmination of fruiting body development after a week of incubation, we harvested and measured total spore contents (Fig. 4b). We found no significant differences in spore productivity according to endosymbiont status (one-way analysis of variance, ($F_{(5,42)}=1.106$, $p=0.372$). Thus, symbiont infections in their natural hosts have no significant impact on host spore productivity under these conditions. We could not assess the fitness impact of *Chlamydiae* or *Amoebophilus* in new host amoebae as our attempts to establish infections in new hosts have been thus far unsuccessful.

Amoebophilus and Chlamydiae *infections do not alter the fitness of hosts during exposure to* Paraburkholderia

Since co-infections with unculturable symbionts and *Paraburkholderia* occur in the wild, we wanted to determine whether these co-infections would result in a different fitness outcome than *Paraburkholderia* infections alone. We tested the fitness impact of *Chlamydiae* and *Amoebophilus* during *Paraburkholderia* exposure using the same strategy as above, but with the addition of 5% by volume of the indicated *Paraburkholderia* strain to food bacteria prior to plating (Fig. 5). These *Paraburkholderia* symbionts range from highly detrimental (*P. hayleyella* strain 11), to intermediately detrimental (*P. agricolaris* strains 159 and 70), to neutral (*B. bonniea* strain 859) under the culturing conditions used for this assay (Haselkorn et al., 2018; Khojandi et al., 2019; Shu et al., 2018). We again observed that exposure to distinct *Paraburkholderia* strains significantly influenced spore productivity for both *Amoebophilus* ($F_{(3,60)}=36.87$, $p=<0.001$) and *Chlamydiae* ($F_{(3,60)}=28.52$, $p=<0.001$) originating host lines (Fig. 6). However, as with single unculturable symbiont infections, we found that presence or absence of *Amoebophilus* or *Chlamydiae* had no significant impact on spore productivity during exposure to *P. hayleyella*-11 ($F_{(5,42)}=1.171$, $p=0.339$), *B. agricolaris*-159 ($F_{(5,42)}=0.985$, $p=0.438$), *P. agricolaris*-70 ($F_{(5,42)}=1.095$, $p=0.377$), or *P. bonniea-859* ($F_{(5,42)}=0.661$, $p=0.655$) (Fig. 5). Thus, the fitness impact of *Paraburkholderia* symbiosis does not appear to be altered (for better or worse) by co-infections with obligate endosymbionts.

There are a few population-specific positive and negative associations between the different symbionts

Our earlier culture-based screening of this *D. discoideum* natural collection found that 25% of isolates were infected with *Paraburkholderia* symbionts (Brock et al., 2020; Haselkorn et al., 2018). For this new round of sequence-based screening, we identified multiple instances of *Paraburkholderia-Amoebophilus* and *Paraburkholderia-Chlamydiae* co-infections in addition to the *Amoebophilus-Chlamydiae* co-infections from above. Specifically, 42 isolates (6%) were co-infected with *Paraburkholderia* and *Amoebophilus*, 25 (3.5%) with *Paraburkholderia* and *Chlamydiae*, and 9 (1%) were co-infected with all three symbiont types (Fig. 1b). Co-infection of the different endosymbionts was tested in our three highest sampled populations: Texas-Houston Arboretum, Virginia Mountain Lake 2000, and Virginia Mountain Lake 2014. Pairwise Fisher Exact tests showed a small positive association between *Chlamydiae* and *Paraburkholderia* in the Texas population (p=0.0132, 15 observed with 10 expected) but not the Virginia populations. There was a positive association between *Paraburkholderia* and *Amoebophilus* in the Virginia 2000 collection (p=0.000277, 43 observed with 30 expected) and a negative association between *Chlamydiae* and *Amoebophilus* in the Virginia 2014 collection (p=0.000024, none observed with 10 expected).

Discussion:

 Many microorganisms, particularly those that are obligately host-associated, are often overlooked because they are unculturable in standard laboratory conditions. The presence of obligate bacterial symbionts in free-living amoebae has been known for some time, particularly for *Acanthamoeba* (Fritsche et al., 1993; Hall and Voelz, 1985;

Jeon and Jeon, 1976; Proca-Ciobanu et al., 1975). The list of amoebal endosymbionts and their host species has since expanded but obligate endosymbionts have not been previously described in the social amoeba *D. discoideum* (Corsaro et al., 2010; Horn and Wagner, 2004; Schmitz-Esser et al., 2008; Schulz et al., 2014)*.* Here, we screened a collection of 730 *D. discoideum* isolates using a PCR detection strategy in order to identify bacterial associates that could be missed using conventional culturing techniques. Furthermore, to our knowledge, this is the largest collection of amoebae from natural populations to be screened, facilitated by the relative ease in which social amoeba fruiting bodies can be sampled. Here, we find that obligate symbionts are quite common among *D. discoideum* collected from the wild, with bacterial sequences of unculturable bacteria present in 301 isolates (41.2%). All but two of these unculturable bacterial sequences aligned most closely to *Amoebophilus* or *Chlamydiae* species. The identification of these bacteria in *D. discoideum* reveals that this valuable eukaryotic model organism is a natural vector of environmental bacteria. This opens a new role for *D. discoideum* as a research tool for the investigation of how bacterial endosymbionts evolve to live inside hosts, how they affect the fitness of their amoeba host, and how they interact with other bacteria in the amoeba microbiome.

The intracellular morphology of endosymbionts in *D. discoideum* spores were in some cases similar to previous observations of related endosymbionts in other amoeba. *Amoebophilus* cells in *D. discoideum* spores appeared similar to previous observations of *Amoebophilus* symbionts in *Acanthamoeba sp*. (Horn et al., 2001; Schmitz-Esser et al., 2010). The localization of *Procabacter* endosymbionts within a membrane bound

vacuole has also been observed for Candidatus *Procabacter* sp. OEW1 in *Acanthamoeba* spp. host cells (Heinz et al., 2007). However, this differs from a previous report on *Procabacter* symbionts in *Acanthamoeba* trophozoites and cysts wherein bacterial cells were not found to be enclosed in vacoules (Fritsche et al., 2002). The wrinkled morphology of *Chlamydiae* cells in *D. discoideum* spores is somewhat reminiscent of the spiny structure of *Chlamydia*-like symbionts in *Acanthamoeba, Naegleria,* and *Hartmannella* host species (Casson et al., 2008b; Corsaro and Greub, 2006; Horn et al., 2000).

Chlamydiae-like organisms typically have two or more morphotypes that correspond to distinct developmental stages. Elementary bodies are the infectious and metabolically inert form that can be transmitted extracellularly from cell to cell. Once internalized, elementary bodies differentiate into metabolically active and replicating reticulate bodies within a vacuole (Bayramova et al., 2018). However, in *D. discoideum* host spores, we only visualize one morphology that does not appear to be in an inclusion but is rather distributed throughout the cytosol. It is unclear what morphotype this represents and whether other morphotypes exist in *D. discoideum* prior to spore formation. It is possible that additional *Chlamydiae* developmental stages occur in vegetative amoebae and that only one form is favored within metabolically inert spores.

Out of the 251 amoeba isolates from the 2014 Virginia Mountain Lake collection, we identified only two infected with a bacterium with sequence homology to *Procabacter*, suggesting that *D. discoideum* is not a preferred natural host for this endosymbiont. Yet,

this *D. discoideum Procabacter* lineage possibly represents an important link for understanding the evolution and adaptation of *Procabacter* to amoeba hosts, as the phylogenetic placement of the novel *D. discoideum Procabacter* lineage is sister to the rest of the *Procabacter* species infecting *Acanthamoeba.* Comparative genome sequencing of this clade, as well as future research into the impact of these infections on their hosts and more extensive wild isolate screenings will better inform our understanding of the significance of these infections in *D. discoideum.*

We identified multiple *D. discoideum* infected with an *Amoebophilus* species that most closely matched Ca. *Amoebophilus asiaticus*, an endosymbiont originally discovered in *Acanthamoeba* (Horn et al., 2001). Similar to the *D. discoideum Procabacter* symbionts, the *D. discoideum Amoebophilus* symbiont lineage is sister to the rest of the *Amoebophilus* clade. The next most closely related bacteria to the *Amoebophilus* genus are the insect facultative endosymbionts of the *Cardinium* genus (Santos-Garcia et al., 2014). These bacteria infect up to 7% of all insect species and have diverse effects on their hosts ranging from reproductive parasitism to nutritional supplementation and defense against parasites. Comparative genome sequencing has uncovered potential routes of adaptation from free-living marine bacteria to either an insect or amoeba host (Santos-Garcia et al., 2014). Including this additional *Amoebophilus* species in comparative genome analysis could further elucidate this process and highlight potential amoeba species-specific adaptations. *Acanthamoeba* and *Dictyostelium* are evolutionarily distant hosts with different lifestyles, thus specific bacterial adaptations may be necessary.

The global significance of *Amoebophilus* infections is under active investigation. *Amoebophilus* have been associated with coral microbiomes and bovine digital dermatitis lesions, which may simply be an indicator that their host amoebae are present in these samples (Apprill et al., 2016; Choi et al., 2009; Horn et al., 2001; Huggett and Apprill, 2019; Zinicola et al., 2015). In amoebae, the full extent of this bacterium's affect is unknown. The moderately reduced genome of Ca. *Amoebophilus asiaticus* lacks important metabolic pathways but contains a high number of proteins with eukaryotic domains that are thought to be essential for host cell interactions (Schmitz-Esser et al., 2010). Genes associated with host cell exploitation, such as toxins, antimicrobial factors, and a unique secretion system, suggest that Ca. *Amoebophilus asiaticus* is perhaps more parasitic than mutualistic (Böck et al., 2017; Penz et al., 2010; Schmitz-Esser et al., 2010). We did not observe an obvious fitness cost of hosting *Amoebophilus*, but additional culture conditions and more sensitive assays will help to elucidate its significance for host amoebae.

Chlamydiae endosymbionts were the most prevalent in the screened collection, with approximately 27% of *D. discoideum* isolates infected. Environmental *Chlamydiae* appear to be diverse and ubiquitous and many *Chlamydiae* endosymbionts associate with a wide range of protozoa and other organisms (Corsaro et al., 2013; Corsaro and Venditti, 2006; Coulon et al., 2012). The incredible genetic diversity within the phylum *Chlamydiae* suggests ancient interactions with eukaryotes, with estimates that the association is 700 million years old (Greub and Raoult, 2003). There are currently nine

recognized families within this order. However recent screening of sequence databases has shown even more diverse lineages, with perhaps 181 total families (Lagkouvardos et al., 2014b). This type of screening, however, does not identify the hosts, so insight into adaptation to particular hosts is limited. In this study we have identified eight novel lineages, potentially representing two novel families, one with several closely related haplotypes suggesting ongoing evolution within the *D. discoideum* host. The separation of these novel *D. discoideum Chlamydiae* lineages from the other amoeba lineages suggests a long-standing association and possible co-adaptation. As our attempts to transfer *Amoebophilus* or *Chlamydiae* endosymbionts to new *D. discoideum* hosts were unsuccessful, we could not probe the potential role of host-symbiont adaptation in fitness outcomes. This strengthens the speculation that these symbionts have been associated with their specific host lines for a long period of time but also raises the question of how they were originally acquired. Other lab experiments have shown that some of these *Chlamydiae* strains are easily transferred across different host genera while others cannot be used to successfully re-infect even their original host amoebae (Corsaro et al., 2013; Coulon et al., 2012; Okude et al., 2012). A previous transfer of a *Protochlamydia* strain to new hosts resulted in a decreased growth rate, suggesting mutual co-evolution between the endosymbiont and its original host (Okude et al., 2012).

While we did not find any obvious fitness costs from *Chlamydiae* infection of *D. discoideum* under laboratory conditions, there may be environmental or genotypic context-dependent effects. Studies of the impact of *Chlamydiae* endosymbionts on their

amoeba hosts suggest that the range of outcomes spans the parasite to mutualist continuum (Hayashi et al., 2012). Depending on the temperature, one *Parachlamydia* endosymbiont of *Acanthamoeba* can either be lytic or endosymbiotic (Greub et al., 2003). The *Chlamydiae* endosymbiont S13 does not significantly harm its *Acanthamoeba* host and instead has been shown to protect it from *Legionella pneumophila* infection (Ishida et al., 2014; Maita et al., 2018; Matsuo et al., 2009; Okude et al., 2012). A *Protochlamydia* endosymbiont has been shown to increase the migration rate, actin density, and growth rate of its native host but had an adverse effect when transferred to non-native hosts (Okude et al., 2012). Alternatively, sequencing of a *Rhabochlamydia* genome from ticks shows incomplete metabolic pathways suggesting that this endosymbiont is not providing metabolic benefits and instead may be a specialized intracellular pathogen (Pillonel et al., 2019).

Sampling large numbers of individuals from *D. discoideum* populations allows us to gain insight into the geographic distribution of these endosymbionts. Many of the *Chlamydiae* haplotypes are broadly distributed; haplotype 1 represents 80% of *Chlamydiae* infections and is found in 12 of 16 populations. Our data also suggest increases (*Chlamydiae*) and decreases (*Amoebophilus*) of particular symbionts in specific regions (Virginia Mountain Lake) over time. It is unclear whether the abundance of particular endosymbiont haplotypes represents an epidemic of a new genotype or selective sweeps due to beneficial effects on their hosts under particular environmental conditions, as seen in insect symbionts (Cockburn et al., 2013; Himler et al., 2011). Although the two Virginia collections differ in time, they were also sampled at slightly

different locations, making time and location confounding factors. Further sampling will separate out the effects of season, microhabitat, and time on amoeba endosymbiont prevalence.

At least 11.4% (83 out of 730) of these *D. discoideum* are co-infected with some combination of *Chlamydiae*, *Amoebophilus,* or *Paraburkholderia*. The lack of detectable fitness costs from co-infection with unculturable endosymbionts and *Paraburkholderia* in the laboratory is consistent with random associations found in several populations. In different environments, however, other microbe-microbe interactions may have an effect that favors certain combinations of symbionts, as is seen in aphids (Rock et al., 2018). *Chlamydiae* occur in co-infections less often than would be expected with *Amoebophilus* in one *D. discoideum* population, consistent with the observation that it interacts negatively with the *Legionella* pathogen in *Acanthamoeba*. On the other hand, *Paraburkholderia* and *Amoebophilus* are positively associated in one population. *Paraburkholderia* symbionts of *D. discoideum* have been shown to allow other food bacteria to enter into the amoeba spores (Brock et al 2011, DiSalvo et al 2015), and perhaps a similar positive interaction occurs in this case. This effect may outweigh some of the negative interactions of *Chlamydiae* with other bacteria, as in one population there was positive association with *Chlamydiae* and *Paraburkholderia*. *Paraburkholderia* symbionts are easily transferred to new hosts via co-culturing and can be found both intracellularly and extracellularly in amoeba cells and structures (DiSalvo et al., 2015; Haselkorn et al., 2018; Shu et al., 2018). *Paraburkholderia* infections in new hosts may be detrimental or beneficial depending on symbiont genotype and

environmental context (Haselkorn et al., 2018; Khojandi et al., 2019). Endosymbiont co-infection is not unique to *D. discoideum,* as previous and alternative isolation methods are allowing for identification of co-infections in *Acanthamoeba* (Heinz et al., 2007; Lagkouvardos et al., 2014a).

There is concern that amoeba-resistant microorganisms, such as the obligate endosymbionts described here, may also be resistant to human phagocytic cells (Friedman et al., 2003; Greub et al., 2005; Jacquier et al., 2013; Kebbi-Beghdadi and Greub, 2014). If so, emerging or opportunistic human pathogens could persist and evolve in the wild within their amoeba reservoirs (Guimaraes et al., 2016; Harb et al., 2000; Horn and Wagner, 2004; Molmeret et al., 2005). Adaptations that enable a bacterium to survive within amoeba hosts could correspondingly equip them for intracellular survival in other eukaryotes (Scheid, 2014). Diverse protozoa are the natural hosts of *Legionella pneumophila* while humans are accidental and unfortunate dead-end hosts (Boamah et al., 2017). The prevalence of environmental *Chlamydiae* in *D. discoideum* is potentially concerning given the importance of *Chlamydia* species in human infections (Taylor-Brown et al., 2015). Some *Chlamydiae* symbionts of free-living amoebae can replicate in human cell lines, encode host exploitation factors, and have been associated with healthy and symptomatic animals and humans (Amann et al., 1997; Casson et al., 2008a, 2006; Corsaro and Greub, 2006; Croxatto et al., 2013; Fukumoto et al., 2016; Greub, 2009; Greub et al., 2005; Greub and Raoult, 2002; Heinz et al., 2010; König et al., 2017; Sampo et al., 2014). However, it remains unclear how significant the threat of amoebae vectored *Chlamydiae* are, or will become, for animal

infections (Birtles et al., 1997; Goy and Greub, 2005; Mamolen et al., 1993; Marrie, 2001; Michel et al., 2004). Interestingly, co-infection of endosymbionts within an amoeba host may facilitate gene exchange that could theoretically lead to the evolution of a new pathogen. For instance, genomic analysis of *Legionella* and *Chlamydiae* amoebae endosymbiont isolates suggest that gene exchange has occurred between *Legionellales* and *Chlamydiales*, presumably during co-infection in a common host (Gimenez et al., 2011).

The growing list of bacterial symbionts of free-living amoebae attests to its attractiveness as an intracellular niche for bacterial symbionts. The similarities between amoebae and animal macrophages, as well as the shared intracellular entrance and survival mechanisms employed by endosymbionts and pathogens, further highlight the potential medical and research relevancy of amoebae-symbiosis systems. The discovery of a high prevalence of infection and novel lineages of common amoebae symbiont clades in the model organism *D. discoideum* opens the door to future research on the ecological significance and mechanisms of amoeba-symbiont interactions.

Methods:

Bacterial Strains and Cultural Conditions:

We used *Klebsiella pneumoniae* throughout as our standard *D. discoideum* food supply. For *Paraburkholderia*-unculturable co-infection experiments, we used the *Paraburkholderia* symbionts *P. hayleyella*-11, *P. agricolaris*-70, *P. agricolaris*-159, and *B. bonniea*-859. We grew bacteria on SM/5 agar plates [2 g glucose, 2 g BactoPeptone,

2 g yeast extract, 0.2 g MgCl$_2$, 1.9 g KHPO$_4$, 1 g K2HPO$_5$, and 15 g agar per liter,

premixed from Formedium] at room temperature before resuspending them in KK2

buffer [2.25 g KH2PO$_4$ (Sigma-Aldrich) and 0.67 g K2HPO$_4$ (Fisher Scientific) per liter].

We then adjusted bacterial suspension to a final OD$_{600nm}$ of 1.5. We plated amoebae

spores with 200µl of bacterial suspensions for all culturing conditions.

Symbiont Screening

 Total DNA from *D. discoideum* was harvested by suspending 15-25 sori in a 5% chelex

solution with 10µg proteinase K, incubating samples for 240 minutes at 56°C, 30

minutes at 98°C, and removing DNA supernatant from chelex beads. PCR with

eukaryotic specific primers was run as a DNA extraction control, and symbiont specific

primers were used to detect symbionts in each sample (Table S2) (Corsaro et al., 2002;

Ossewaarde and Meijer, 1999). For all primer sets we used a touchdown PCR protocol

consisting of 94°C for 30 s, 68°C-55°C for 45 x (dropping 1 degree per cycle) for 14

cycles, and 72°C for 1 min, followed by 94°C, 55°C, 72°C for 20 cycles. To confirm

infection we ran PCR samples on a one percent agarose gel with TAE buffer and

imaged on a transilluminator. PCR products were sent for sequencing (Genewiz), and

the resulting Sanger sequences were cleaned using the program Geneious v8

(http://www.geneious.com (Kearse et al., 2012)). We use NCBI Blast to initially identify

the bacterial species/strain based on highest sequence similarity. Sequences have

been deposited to GenBank under accession numbers (MT596490-MT596496).

Amoebae Clone Types and Culture Conditions:

Amoebae were cultured from freezer stocks by resuspending frozen spores onto SM/5 agar plates with 200µl of food *K. pneumoniae* suspensions. Plates were incubated at room temperature under low lights and ~60% humidity for approximately 1 week (or until fruiting body development).

For antibiotic curing and spore quantification, we used four uninfected, *Amoebophilus* infected, and *Chlamydiae* infected host lines respectively. Uninfected representatives were QS864, QS970, QS1003 and QS1069. *Amoebophilus* infected representatives included QS851, QS1002, QS1011, and QS1017. *Chlamydiae* infected representatives included QS866, QS991, QS1023, and QS1089. For all experiments, we initiated experimental conditions from pre-developed fruiting bodies by suspending ~8-15 sorus contents in KK2. We then plated 10^5 spores onto an SM/5 plate with 200µl of bacterial suspensions (*K. pneumoniae* only or with 95% *K. pneumoniae/* 5% *Paraburkholderia* mixtures by volume for co-infection experiments). Plates were then incubated as described above for the time indicated for each experimental condition.

Curing of Symbiont Infections:

We treated uninfected, *Chlamydiae*-infected, and *Amoebophilus* infected representative isolates with rifampacin by plating ~10^2 spores with 400µl *K. pneumoniae* on SM/5 medium containing 30µg/ml rifampacin. Plates were incubated until fruiting body formation. Sori from grown fruiting bodies were then harvested and replated for a second round of antibiotic exposure (using the same strategy as described above). Following fruiting body formation, spores were harvested and transferred to SM/5 with

K. pneumoniae and incubated. Following fruiting body formation on non-antibiotic

plates, we extracted total sorus content DNA and checked for symbiont infection using

the *Chlamydiae*-specific and *Amoebophilus*-specific PCR protocol described above.

Total Spore Productivity Assay:

To determine total spore productivity of antibiotic treated and untreated *D. discoideum*

isolates we plated 10^5 spores on SM/5 agar with 200µl of *K. pneumoniae* and incubated

at room temperature. Five days after plating, total plate contents were collected into

~10ml KK_2 + NP-40 alternative detergent solution. Solutions were then diluted

appropriately (~20x) and spores quantified via hemocytometer counts. We measured

and averaged over at least three replicates. To determine spore productivity after

exposure to *Paraburkholderia*, we conducted the assay as described but plated spores

with 200µl of a 95/5% by volume mixture of *K. pneumoniae/Paraburkholderia* cell

suspensions.

Statistics

Significant positive or negative associations between co-infecting endosymbionts were

determined using a Fisher Exact test (2x2 contingency table) on pairwise comparisons

in our three most highly sampled populations. Statistical analyses for spore productivity

experiments were performed in R 3.6.0. Comparisons were done using one-way

ANOVA's with antibiotic treatment as the independent variable.

Transmission Electron Microscopy

Complete processing and imaging of spores by transmission electron microscopy was performed by Wandy Beatty at the Washington University Molecular Microbiology Imaging Facility. *D. discoideum* isolates imaged included QS864, QS851, QS886, QS1040. Briefly, spores were harvested in KK2 buffer and fixed with 2% paraformaldehyde with 2.5% glutaraldehyde (Polysciences Inc., Warrington, PA) in 100 mM cacodylate buffer (pH 7.2) for 1-3 hrs. at room temperature. We washed samples with cacodylate buffer and postfixed in 1% osmium tetroxide (Polysciences Inc.) for 1 hr before rinsing in dH$_2$O prior to *en bloc* staining with 1% aqueous uranyl acetate (Ted Pella Inc., Redding, CA) for 1 hr. We dehydrated samples in increasing concentrations of ethanol and embedded them in Eponate 12 resin (Ted Pella Inc.). Sections of 95 nm were cut with a Leica Ultracut UCT ultramicrotome (Leica Microsystems Inc., Bannockburn, IL), stained with uranyl acetate and lead citrate, and viewed on a JEOL 1200 EX transmission electron microscope (JEOL USA Inc., Peabody, MA) equipped with an AMT 8 megapixel digital camera (Advanced Microscopy Techniques, Woburn, MA).

Phylogenetic Tree Construction

The *Procabacter* full length 16S rRNA phylogeny was created using the top ten NCBI Blast results for the *Procabacter* endosymbiont of D. *discoideum* sequence and representative species from the phylum *Betaproteobacteria*. The species were chosen using the NCBI taxonomy database and 16S rRNA sequences were obtained from the NCBI nucleotide database. A sequence from *Rickettsia rickettsii* was also obtained from the NCBI nucleotide database to serve as an outgroup. Sequences were imported into Mega7 (Tamura et al, 2013) and aligned with MUSCLE. All sequences were then

trimmed to a uniform 1396 bp. The Kimura 2-parameter model with a discrete Gamma

distribution was selected using the "Find Best DNA/Protein Models" command in

Mega7. A Maximum Likelihood tree was then constructed with 1000 bootstrap

replicates.

The *Amoebophilus* full length 16S rRNA phylogeny was created using the top five NCBI

Blast results for the *Amoebophilus* endosymbiont of *D. discoideum* sequence, a

sequence from the related endosymbiont *Candidatus Cardinium hertigii*, and

representative species from the order *Cytophagales*. The species were chosen using

the phylogeny in Hahnke et al., (2016) as a guide, and full length 16S rRNA sequences

were obtained from the NCBI nucleotide database (Hahnke et al., 2016). All four of

these representatives were isolated in a soil environment. A sequence from

Bacteroides fragilis was also obtained from the NCBI nucleotide database to serve as

an outgroup. Sequences were imported into Mega7, aligned with MUSCLE, and

trimmed to a uniform 1364 bp. The Kimura 2-parameter model with a discrete Gamma

distribution was selected using the "Find Best DNA/Protein Models" command in

Mega7. A Maximum Likelihood tree was then constructed with 1000 bootstrap

replicates.

The *Chlamydiae* short 16S rRNA phylogeny and *Chlamydiae* full length 16S rRNA

phylogeny were each created using 1-2 representative species from the nine families in

the phylum *Chlamydiae* from Taylor-Brown et al., (2015). Full length 16S rRNA

sequences of chosen species were obtained from the NCBI nucleotide database.

Sequences were imported into Mega7 and aligned with MUSCLE. For the short tree, sequences were trimmed to a uniform 218 bp. For the full-length tree, sequences were trimmed to 1354 bp. The Tamura 3-parameter model with a discrete Gamma distribution was selected for the short tree, and the General Time Reversal model with a discrete Gamma distribution and invariant sites was selected for the full-length tree using the "Find Best DNA/Protein Models" command in Mega7. A Maximum Likelihood tree was then constructed for each with 1000 bootstrap replicates.

Acknowledgements: We thank members of the Haselkorn, DiSalvo, and Strassmann-Queller labs for their support and suggestions. We also thank Dierdra Renfroe for her attempts to transfer *Chlamydiae* symbionts to new host amoebae. This material is based upon work supported by the National Science Foundation under grant numbers DEB-1753743 and IOS-1656756 to JS and DQ.

References:

Amann, R., Springer, N., Schönhuber, W., Ludwig, W., Schmid, E.N., Müller, K.D., Michel, R., 1997. Obligate intracellular bacterial parasites of acanthamoebae related to Chlamydia spp. Appl. Environ. Microbiol. 63, 115–121.

Amissah, N.A., Gryseels, S., Tobias, N.J., Ravadgar, B., Suzuki, M., Vandelannoote, K., Durnez, L., Leirs, H., Stinear, T.P., Portaels, F., Ablordey, A., Eddyani, M., 2014. Investigating the Role of Free-living Amoebae as a Reservoir for Mycobacterium ulcerans. PLoS Negl. Trop. Dis. 8, e3148. https://doi.org/10.1371/journal.pntd.0003148

Annesley, S.J., Fisher, P.R., 2009. Dictyostelium discoideum—a model for many reasons. Mol. Cell. Biochem. 329, 73–91. https://doi.org/10.1007/s11010-009-0111-8

Apprill, A., Weber, L.G., Santoro, A.E., 2016. Distinguishing between Microbial Habitats Unravels Ecological Complexity in Coral Microbiomes. mSystems 1, e00143-16, /msys/1/5/e00143-16.atom. https://doi.org/10.1128/mSystems.00143-16

Barker, J., Brown, M.R.W., 1994. Trojan Horses of the microbial world: protozoa and the survival of bacterial pathogens in the environment. Microbiology 140, 1253–1259. https://doi.org/10.1099/00221287-140-6-1253

Bayramova, F., Jacquier, N., Greub, G., 2018. Insight in the biology of Chlamydia-related bacteria. Microbes Infect. 20, 432–440. https://doi.org/10.1016/j.micinf.2017.11.008

Birtles, R., Rowbotham, T., Storey, C., Marrie, T., Raoult, D., 1997. Chlamydia-like obligate parasite of free-living amoebae. The Lancet 349, 925–926. https://doi.org/10.1016/S0140-6736(05)62701-8

Boamah, D.K., Zhou, G., Ensminger, A.W., O'Connor, T.J., 2017. From Many Hosts, One Accidental Pathogen: The Diverse Protozoan Hosts of Legionella. Front. Cell. Infect. Microbiol. 7. https://doi.org/10.3389/fcimb.2017.00477

Böck, D., Medeiros, J.M., Tsao, H.-F., Penz, T., Weiss, G.L., Aistleitner, K., Horn, M., Pilhofer, M., 2017. In situ architecture, function, and evolution of a contractile injection system. Science 357, 713–717. https://doi.org/10.1126/science.aan7904

Bozzaro, S., Buracco, S., Peracino, B., Eichinger, L., 2019. Dictyostelium Host Response to Legionella Infection: Strategies and Assays, in: Buchrieser, C., Hilbi, H. (Eds.), Legionella. Springer New York, New York, NY, pp. 347–370. https://doi.org/10.1007/978-1-4939-9048-1_23

Bozzaro, S., Eichinger, L., 2011. The professional phagocyte Dictyostelium discoideum as a model host for bacterial pathogens. Curr. Drug Targets 12, 942–954. https://doi.org/10.2174/138945011795677782

Brock, D.A., Douglas, T.E., Queller, D.C., Strassmann, J.E., 2011. Primitive agriculture in a social amoeba. Nature 469, 393–396. https://doi.org/10.1038/nature09668

Brock, D.A., Haselkorn, T.S., Garcia, J.R., Bashir, U., Douglas, T.E., Galloway, J., Brodie, F., Queller, D.C., Strassmann, J.E., 2018. Diversity of free-living environmental bacteria and their interactions with a bactivorous amoeba. Front. Cell. Infect. Microbiol. 8, 411–411. https://doi.org/10.3389/fcimb.2018.00411

Brock, D.A., Noh, S., Hubert, A.N.M., Haselkorn, T.S., DiSalvo, S., Suess, M.K., Bradley, A.S., Tavakoli-Nezhad, M., Geist, K.S., Queller, D.C., Strassmann, J.E., 2020. Endosymbiotic adaptations in three new bacterial species associated with *Dictyostelium discoideum* : *Paraburkholderia agricolaris* sp. nov., *Paraburkholderia hayleyella* sp. nov., and *Paraburkholderia bonniea* sp. nov. PeerJ 8, e9151. https://doi.org/10.7717/peerj.9151

Cardenal-Muñoz, E., Barisch, C., Lefrançois, L.H., López-Jiménez, A.T., Soldati, T., 2018. When Dicty Met Myco, a (Not So) Romantic Story about One Amoeba and Its Intracellular Pathogen. Front. Cell. Infect. Microbiol. 7. https://doi.org/10.3389/fcimb.2017.00529

Casson, N., Entenza, J.M., Borel, N., Pospischil, A., Greub, G., 2008a. Murine model of pneumonia caused by Parachlamydia acanthamoebae. Microb. Pathog. 45, 92–97. https://doi.org/10.1016/j.micpath.2008.04.003

Casson, N., Medico, N., Bille, J., Greub, G., 2006. Parachlamydia acanthamoebae enters and multiplies within pneumocytes and lung fibroblasts. Microbes Infect. 8, 1294–1300. https://doi.org/10.1016/j.micinf.2005.12.011

Casson, N., Michel, R., Müller, K.-D., Aubert, J.D., Greub, G., 2008b. *Protochlamydia naegleriophila* as Etiologic Agent of Pneumonia. Emerg. Infect. Dis. 14, 168–172. https://doi.org/10.3201/eid1401.070980

Chen, G., Zhuchenko, O., Kuspa, A., 2007. Immune-like phagocyte activity in the social amoeba. Science 317, 678–681. https://doi.org/10.1126/science.1143991

Choi, S.H., Cho, M.K., Ahn, S.C., Lee, J.E., Lee, J.S., Kim, D.-H., Xuan, Y.-H., Hong, Y.C., Kong, H.H., Chung, D.I., Yu, H.S., 2009. Endosymbionts of Acanthamoeba Isolated from Domestic Tap Water in Korea. Korean J. Parasitol. 47, 337. https://doi.org/10.3347/kjp.2009.47.4.337

Clarholm, M., 1981. Protozoan grazing of bacteria in soil—impact and importance. Microb. Ecol. 7, 343–350. https://doi.org/10.1007/BF02341429

Cockburn, S.N., Haselkorn, T.S., Hamilton, P.T., Landzberg, E., Jaenike, J., Perlman, S.J., 2013. Dynamics of the continent-wide spread of a *Drosophila* defensive symbiont. Ecol. Lett. 16, 609–616. https://doi.org/10.1111/ele.12087

Corsaro, D., Greub, G., 2006. Pathogenic Potential of Novel Chlamydiae and Diagnostic Approaches to Infections Due to These Obligate Intracellular Bacteria. Clin. Microbiol. Rev. 19, 283–297. https://doi.org/10.1128/CMR.19.2.283-297.2006

Corsaro, D., Müller, K.-D., Wingender, J., Michel, R., 2013. "Candidatus Mesochlamydia elodeae" (Chlamydiae: Parachlamydiaceae), a novel chlamydia parasite of free-living amoebae. Parasitol. Res. 112, 829–838. https://doi.org/10.1007/s00436-012-3213-2

Corsaro, D., Pages, G.S., Catalan, V., Loret, J.-F., Greub, G., 2010. Biodiversity of amoebae and amoeba-associated bacteria in water treatment plants. Int. J. Hyg. Environ. Health 213, 158–166. https://doi.org/10.1016/j.ijheh.2010.03.002

Corsaro, D., Venditti, D., 2006. Diversity of the Parachlamydiae in the Environment. Crit. Rev. Microbiol. 32, 185–199. https://doi.org/10.1080/10408410601023102

Corsaro, D., Venditti, D., Valassina, M., 2002. New parachlamydial 16S rDNA phylotypes detected in human clinical samples. Res. Microbiol. 153, 563–567. https://doi.org/10.1016/S0923-2508(02)01369-4

Cosson, P., Soldati, T., 2008. Eat, kill or die: when amoeba meets bacteria. Curr. Opin. Microbiol. 11, 271–276. https://doi.org/10.1016/j.mib.2008.05.005

Coulon, C., Eterpi, M., Greub, G., Collignon, A., McDonnell, G., Thomas, V., 2012. Amoebal host range, host-free survival and disinfection susceptibility of environmental *Chlamydiae* as compared to *Chlamydia trachomatis*. FEMS Immunol. Med. Microbiol. 64, 364–373. https://doi.org/10.1111/j.1574-695X.2011.00919.x

Croxatto, A., Murset, V., Chassot, B., Greub, G., 2013. Early expression of the type III secretion system of *Parachlamydia acanthamoebae* during a replicative cycle within its natural host cell *Acanthamoeba castellanii*. Pathog. Dis. 69, 159–175. https://doi.org/10.1111/2049-632X.12065

Dinh, C., Farinholt, T., Hirose, S., Zhuchenko, O., Kuspa, A., 2018. Lectins modulate the microbiota of social amoebae. Science 361, 402–406. https://doi.org/10.1126/science.aat2058

DiSalvo, S., Haselkorn, T.S., Bashir, U., Jimenez, D., Brock, D.A., Queller, D.C., Strassmann, J.E., 2015. Burkholderia bacteria infectiously induce the proto-farming symbiosis of Dictyostelium amoebae and food bacteria. Proc. Natl. Acad. Sci. U. S. A. 112, E5029–E5037. https://doi.org/10.1073/pnas.1511878112

Drancourt, M., 2014. Looking in amoebae as a source of mycobacteria. Microb. Pathog. 77, 119–124. https://doi.org/10.1016/j.micpath.2014.07.001

Erken, M., Lutz, C., McDougald, D., 2013. The Rise of Pathogens: Predation as a Factor Driving the Evolution of Human Pathogens in the Environment. Microb. Ecol. 65, 860–868. https://doi.org/10.1007/s00248-013-0189-0

Everett, K.D.E., Bush, R.M., Andersen, A.A., 1999. Emended description of the order Chlamydiales, proposal of Parachlamydiaceae fam. nov. and Simkaniaceae fam. nov., each containing one monotypic genus, revised taxonomy of the family Chlamydiaceae, including a new genus and five new species, and standards for the identification of organisms. Int. J. Syst. Evol. Microbiol. 49, 415–440. https://doi.org/10.1099/00207713-49-2-415

Farinholt, T., Dinh, C., Kuspa, A., 2019. Microbiome management in the social amoeba Dictyostelium discoideum compared to humans. Int. J. Dev. Biol. 63, 447–450. https://doi.org/10.1387/ijdb.190240ak

Friedman, M.G., Dvoskin, B., Kahane, S., 2003. Infections with the chlamydia-like microorganism Simkania negevensis, a possible emerging pathogen. Microbes Infect. 5, 1013–1021. https://doi.org/10.1016/S1286-4579(03)00188-6

Fritsche, T.R., Gautom, R.K., Seyedirashti, S., Bergeron, D.L., Lindquist, T.D., 1993. Occurrence of bacterial endosymbionts in Acanthamoeba spp. isolated from corneal and environmental specimens and contact lenses. J. Clin. Microbiol. 31, 1122–1126.

Fritsche, T.R., Linner, T., Wagner, M., Gautom, R.K., Harzenetter, M.D., Horn, M., 2002. Obligate bacterial endosymbionts of Acanthamoeba spp. related to the beta-Proteobacteria: proposal of "Candidatus Procabacter acanthamoebae" gen. nov., sp. nov. Int. J. Syst. Evol. Microbiol. 52, 599–605. https://doi.org/10.1099/00207713-52-2-599

Fukumoto, T., Matsuo, J., Okubo, T., Nakamura, S., Miyamoto, K., Oka, K., Takahashi, M., Akizawa, K., Shibuya, H., Shimizu, C., Yamaguchi, H., 2016. Acanthamoeba containing endosymbiotic chlamydia isolated from hospital environments and its potential role in inflammatory exacerbation. BMC Microbiol. 16, 292. https://doi.org/10.1186/s12866-016-0906-1

Gimenez, G., Bertelli, C., Moliner, C., Robert, C., Raoult, D., Fournier, P.-E., Greub, G., 2011. Insight into cross-talk between intra-amoebal pathogens. BMC Genomics 12, 542. https://doi.org/10.1186/1471-2164-12-542

Gomez-Valero, L., Buchrieser, C., 2019. Intracellular parasitism, the driving force of evolution of Legionella pneumophila and the genus Legionella. Genes Immun. 20, 394–402. https://doi.org/10.1038/s41435-019-0074-z

Goy, G., Greub, G., 2005. [The pathogenicity of amoebae-resisting chlamydiales]. Rev. Med. Suisse 1, 1916–1920.

Greub, G., 2009. Parachlamydia acanthamoebae, an emerging agent of pneumonia. Clin. Microbiol. Infect. 15, 18–28. https://doi.org/10.1111/j.1469-0691.2008.02633.x

Greub, G., Mege, J.-L., Gorvel, J.-P., Raoult, D., Méresse, S., 2005. Intracellular trafficking of Parachlamydia acanthamoebae: Intracellular trafficking of Parachlamydia. Cell. Microbiol. 7, 581–589. https://doi.org/10.1111/j.1462-5822.2004.00488.x

Greub, G., Raoult, D., 2003. History of the ADP/ATP-Translocase-Encoding Gene, a Parasitism Gene Transferred from a Chlamydiales Ancestor to Plants 1 Billion Years Ago. Appl. Environ. Microbiol. 69, 5530–5535. https://doi.org/10.1128/AEM.69.9.5530-5535.2003

Greub, G., Raoult, D., 2002. *Parachlamydiaceae* : Potential Emerging Pathogens. Emerg. Infect. Dis. 8, 626–630. https://doi.org/10.3201/eid0806.010210

Greub, G., Scola, B., Raoult, D., 2003. *Parachlamydia acanthamoeba* Is Endosymbiotic or Lytic for *Acanthamoeba polyphaga* Depending on the Incubation Temperature. Ann. N. Y. Acad. Sci. 990, 628–634. https://doi.org/10.1111/j.1749-6632.2003.tb07437.x

Guimaraes, A.J., Gomes, K.X., Cortines, J.R., Peralta, J.M., Peralta, R.H.S., 2016. Acanthamoeba spp. as a universal host for pathogenic microorganisms: One bridge from environment to host virulence. Microbiol. Res. 193, 30–38. https://doi.org/10.1016/j.micres.2016.08.001

Hahnke, R.L., Meier-Kolthoff, J.P., García-López, M., Mukherjee, S., Huntemann, M., Ivanova, N.N., Woyke, T., Kyrpides, N.C., Klenk, H.-P., Göker, M., 2016. Genome-Based Taxonomic Classification of Bacteroidetes. Front. Microbiol. 7, 2003. https://doi.org/10.3389/fmicb.2016.02003

Hall, J., Voelz, H., 1985. Bacterial Endosymbionts of Acanthamoeba sp. J. Parasitol. 71, 89. https://doi.org/10.2307/3281982

Harb, O.S., Gao, L.-Y., Kwaik, Y.A., 2000. From protozoa to mammalian cells: a new paradigm in the life cycle of intracellular bacterial pathogens. Minireview. Environ. Microbiol. 2, 251–265. https://doi.org/10.1046/j.1462-2920.2000.00112.x

Haselkorn, T.S., DiSalvo, S., Miller, J.W., Bashir, U., Brock, D.A., Queller, D.C., Strassmann, J.E., 2018. The specificity of Burkholderia symbionts in the social amoeba farming symbiosis: prevalence, species, genetic and phenotypic diversity. Mol. Ecol. 0. https://doi.org/10.1111/mec.14982

Hayashi, Y., Yimin, Matsuo, J., Nakamura, S., Kunichika, M., Yoshida, M., Takahashi, K., Yamaguchi, H., 2012. A domino-like chlamydial attachment process: concurrent Parachlamydia acanthamoebae attachment to amoebae is required for several amoebal released molecules and serine protease activity. Microbiology 158, 1607–1614. https://doi.org/10.1099/mic.0.057190-0

Heinz, E., Kolarov, I., Kästner, C., Toenshoff, E.R., Wagner, M., Horn, M., 2007. An Acanthamoeba sp. containing two phylogenetically different bacterial endosymbionts. Environ. Microbiol. 9, 1604–1609. https://doi.org/10.1111/j.1462-2920.2007.01268.x

Heinz, E., Rockey, D.D., Montanaro, J., Aistleitner, K., Wagner, M., Horn, M., 2010. Inclusion Membrane Proteins of Protochlamydia amoebophila UWE25 Reveal a Conserved Mechanism for Host Cell Interaction among the Chlamydiae. J. Bacteriol. 192, 5093–5102. https://doi.org/10.1128/JB.00605-10

Himler, A.G., Adachi-Hagimori, T., Bergen, J.E., Kozuch, A., Kelly, S.E., Tabashnik, B.E., Chiel, E., Duckworth, V.E., Dennehy, T.J., Zchori-Fein, E., Hunter, M.S., 2011. Rapid Spread of a Bacterial Symbiont in an Invasive Whitefly Is Driven by Fitness Benefits and Female Bias. Science 332, 254–256. https://doi.org/10.1126/science.1199410

Horn, M., Harzenetter, M.D., Linner, T., Schmid, E.N., Muller, K.-D., Michel, R., Wagner, M., 2001. Members of the Cytophaga-Flavobacterium-Bacteroides phylum as intracellular bacteria of acanthamoebae: proposal of "Candidatus Amoebophilus asiaticus." Environ. Microbiol. 3, 440–449. https://doi.org/10.1046/j.1462-2920.2001.00210.x

Horn, M., Wagner, M., 2004. Bacterial Endosymbionts of Free-living Amoebae1. J. Eukaryot. Microbiol. 51, 509–514. https://doi.org/10.1111/j.1550-7408.2004.tb00278.x

Horn, M., Wagner, M., Müller, K.-D., Schmid, E.N., Fritsche, T.R., Schleifer, K.-H., Michel, R., 2000. Neochlamydia hartmannellae gen. nov., sp. nov. (Parachlamydiaceae), an endoparasite of the amoeba Hartmannella vermiformis. Microbiology, 146, 1231–1239.

Huggett, M.J., Apprill, A., 2019. Coral microbiome database: Integration of sequences reveals high diversity and relatedness of coral-associated microbes. Environ. Microbiol. Rep. 11, 372–385. https://doi.org/10.1111/1758-2229.12686

Ishida, K., Sekizuka, T., Hayashida, K., Matsuo, J., Takeuchi, F., Kuroda, M., Nakamura, S., Yamazaki, T., Yoshida, M., Takahashi, K., Nagai, H., Sugimoto, C., Yamaguchi, H., 2014. Amoebal Endosymbiont Neochlamydia Genome Sequence Illuminates the Bacterial Role in the Defense of the Host Amoebae against Legionella pneumophila. PLoS ONE 9, e95166. https://doi.org/10.1371/journal.pone.0095166

Jacquier, N., Aeby, S., Lienard, J., Greub, G., 2013. Discovery of New Intracellular Pathogens by Amoebal Coculture and Amoebal Enrichment Approaches. J. Vis. Exp. 51055. https://doi.org/10.3791/51055

Jeon, K.W., 1992. Prokaryotic Symbionts of Amoebae and Flagellates, in: Balows, A., Trüper, H.G., Dworkin, M., Harder, W., Schleifer, K.-H. (Eds.), The Prokaryotes. Springer New York, New York, NY, pp. 3855–3864. https://doi.org/10.1007/978-1-4757-2191-1_51

Jeon, K.W., Jeon, M.S., 1976. Endosymbiosis in amoebae: Recently established endosymbionts have become required cytoplasmic components. J. Cell. Physiol. 89, 337–344. https://doi.org/10.1002/jcp.1040890216

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., Drummond, A., 2012. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28, 1647–1649. https://doi.org/10.1093/bioinformatics/bts199

Kebbi-Beghdadi, C., Greub, G., 2014. Importance of amoebae as a tool to isolate amoeba-resisting microorganisms and for their ecology and evolution: the *C hlamydia* paradigm: Discovering new pathogens. Environ. Microbiol. Rep. 6, 309–324. https://doi.org/10.1111/1758-2229.12155

Kessin, R.H., 2001. Dictyostelium: evolution, cell biology, and the development of multicellularity, Developmental and cell biology series. Cambridge University Press, Cambridge, UK ; New York.

Khojandi, N., Haselkorn, T.S., Eschbach, M.N., Naser, R.A., DiSalvo, S., 2019. Intracellular Burkholderia Symbionts induce extracellular secondary infections; driving diverse host outcomes that vary by genotype and environment. ISME J. https://doi.org/10.1038/s41396-019-0419-7

König, L., Siegl, A., Penz, T., Haider, S., Wentrup, C., Polzin, J., Mann, E., Schmitz-Esser, S., Domman, D., Horn, M., 2017. Biphasic Metabolism and Host Interaction of a Chlamydial Symbiont. mSystems 2, e00202-16, /msys/2/3/e00202-16.atom. https://doi.org/10.1128/mSystems.00202-16

Lagkouvardos, I., Shen, J., Horn, M., 2014a. Improved axenization method reveals complexity of symbiotic associations between bacteria and acanthamoebae: An improved method for the axenization of amoebae. Environ. Microbiol. Rep. 6, 383–388. https://doi.org/10.1111/1758-2229.12162

Lagkouvardos, I., Weinmaier, T., Lauro, F.M., Cavicchioli, R., Rattei, T., Horn, M., 2014b. Integrating metagenomic and amplicon databases to resolve the phylogenetic and ecological diversity of the Chlamydiae. ISME J. 8, 115–125. https://doi.org/10.1038/ismej.2013.142

Maita, C., Matsushita, M., Miyoshi, M., Okubo, T., Nakamura, S., Matsuo, J., Takemura, M., Miyake, M., Nagai, H., Yamaguchi, H., 2018. Amoebal endosymbiont Neochlamydia protects host amoebae against Legionella pneumophila infection by preventing Legionella entry. Microbes Infect. 20, 236–244. https://doi.org/10.1016/j.micinf.2017.12.012

Mamolen, M., Lewis, D.M., Blanchet, M.A., Satink, F.J., Vogt, R.L., 1993. Investigation of an outbreak of "Humidifier fever" in a print shop. Am. J. Ind. Med. 23, 483–490. https://doi.org/10.1002/ajim.4700230311

Marrie, T., 2001. Legionella-Like and Other Amoebal Pathogens as Agents of Community-Acquired Pneumonia. Emerg. Infect. Dis. 7, 1026–1029. https://doi.org/10.3201/eid0706.010619

Matsuo, J., Kawaguchi, K., Nakamura, S., Hayashi, Y., Yoshida, M., Takahashi, K., Mizutani, Y., Yao, T., Yamaguchi, H., 2009. Survival and transfer ability of phylogenetically diverse bacterial endosymbionts in environmental Acanthamoeba isolates: Survival and transfer of bacterial endosymbionts in amoebae. Environ. Microbiol. Rep. 2, 524–533. https://doi.org/10.1111/j.1758-2229.2009.00094.x

Matz, C., Kjelleberg, S., 2005. Off the hook – how bacteria survive protozoan grazing. Trends Microbiol. 13, 302–307. https://doi.org/10.1016/j.tim.2005.05.009

Michel, R., Steinert, M., Zöller, L., Hauroeder, B., Henning, K., 2004. Free-living Amoebae May Serve as Hosts for the Chlamydia-like Bacterium Waddlia chondrophila Isolated from an Aborted Bovine Foetus. Acta Protozool. 43, 37–42.

Molmeret, M., Horn, M., Wagner, M., Santic, M., Abu Kwaik, Y., 2005. Amoebae as training grounds for intracellular bacterial pathogens. Appl. Environ. Microbiol. 71, 20–28. https://doi.org/10.1128/AEM.71.1.20-28.2005

Okude, M., Matsuo, J., Nakamura, S., Kawaguchi, K., Hayashi, Y., Sakai, H., Yoshida, M., Takahashi, K., Yamaguchi, H., 2012. Environmental Chlamydiae Alter the Growth Speed and Motility of Host Acanthamoebae. Microbes Environ. 27, 423–429. https://doi.org/10.1264/jsme2.ME11353

Ossewaarde, J.M., Meijer, A., 1999. Molecular evidence for the existence of additional members of the order Chlamydiales. Microbiology 145, 411–417. https://doi.org/10.1099/13500872-145-2-411

Paquet, V.E., Charette, S.J., 2016. Amoeba-resisting bacteria found in multilamellar bodies secreted by *Dictyostelium discoideum:* social amoebae can also package bacteria. FEMS Microbiol. Ecol. 92, fiw025. https://doi.org/10.1093/femsec/fiw025

Penz, T., Horn, M., Schmitz-Esser, S., 2010. The genome of the amoeba symbiont "Candidatus *Amoebophilus asiaticus* " encodes an afp-like prophage possibly used for protein secretion. Virulence 1, 541–545. https://doi.org/10.4161/viru.1.6.13800

Pillonel, T., Bertelli, C., Aeby, S., de Barsy, M., Jacquier, N., Kebbi-Beghdadi, C., Mueller, L., Vouga, M., Greub, G., 2019. Sequencing the Obligate Intracellular Rhabdochlamydia

helvetica within Its Tick Host Ixodes ricinus to Investigate Their Symbiotic Relationship. Genome Biol. Evol. 11, 1334–1344. https://doi.org/10.1093/gbe/evz072

Proca-Ciobanu, M., Lupascu, Gh., Petrovici, Al., Ionescu, M.D., 1975. Electron microscopic study of a pathogenic Acanthamoeba castellani strain: The presence of bacterial endosymbionts. Int. J. Parasitol. 5, 49–56. https://doi.org/10.1016/0020-7519(75)90097-1

Rock, D.I., Smith, A.H., Joffe, J., Albertus, A., Wong, N., O'Connor, M., Oliver, K.M., Russell, J.A., 2018. Context-dependent vertical transmission shapes strong endosymbiont community structure in the pea aphid, *Acyrthosiphon pisum*. Mol. Ecol. 27, 2039–2056. https://doi.org/10.1111/mec.14449

Rodríguez-Zaragoza, S., 1994. Ecology of Free-Living Amoebae. Crit. Rev. Microbiol. 20, 225–241. https://doi.org/10.3109/10408419409114556

Ronn, R., McCaig, A.E., Griffiths, B.S., Prosser, J.I., 2002. Impact of Protozoan Grazing on Bacterial Community Structure in Soil Microcosms. Appl. Environ. Microbiol. 68, 6094–6105. https://doi.org/10.1128/AEM.68.12.6094-6105.2002

Salah, I.B., Ghigo, E., Drancourt, M., 2009. Free-living amoebae, a training field for macrophage resistance of mycobacteria. Clin. Microbiol. Infect. 15, 894–905. https://doi.org/10.1111/j.1469-0691.2009.03011.x

Sampo, A., Matsuo, J., Yamane, C., Yagita, K., Nakamura, S., Shouji, N., Hayashi, Y., Yamazaki, T., Yoshida, M., Kobayashi, M., Ishida, K., Yamaguchi, H., 2014. High-temperature adapted primitive *Protochlamydia* found in *Acanthamoeba* isolated from a hot spring can grow in immortalized human epithelial HEp-2 cells: High-temperature adapted primitive *Protochlamydia*. Environ. Microbiol. 16, 486–497. https://doi.org/10.1111/1462-2920.12266

Santos-Garcia, D., Rollat-Farnier, P.-A., Beitia, F., Zchori-Fein, E., Vavre, F., Mouton, L., Moya, A., Latorre, A., Silva, F.J., 2014. The Genome of Cardinium cBtQ1 Provides Insights into Genome Reduction, Symbiont Motility, and Its Settlement in Bemisia tabaci. Genome Biol. Evol. 6, 1013–1030. https://doi.org/10.1093/gbe/evu077

Scheid, P., 2014. Relevance of free-living amoebae as hosts for phylogenetically diverse microorganisms. Parasitol. Res. 113, 2407–2414. https://doi.org/10.1007/s00436-014-3932-7

Schmitz-Esser, S., Tischler, P., Arnold, R., Montanaro, J., Wagner, M., Rattei, T., Horn, M., 2010. The Genome of the Amoeba Symbiont "Candidatus Amoebophilus asiaticus" Reveals Common Mechanisms for Host Cell Interaction among Amoeba-Associated Bacteria. J. Bacteriol. 192, 1045–1057. https://doi.org/10.1128/JB.01379-09

Schmitz-Esser, S., Toenshoff, E.R., Haider, S., Heinz, E., Hoenninger, V.M., Wagner, M., Horn, M., 2008. Diversity of Bacterial Endosymbionts of Environmental Acanthamoeba Isolates. Appl. Environ. Microbiol. 74, 5822–5831. https://doi.org/10.1128/AEM.01093-08

Schulz, F., Lagkouvardos, I., Wascher, F., Aistleitner, K., Kostanjšek, R., Horn, M., 2014. Life in an unusual intracellular niche: a bacterial symbiont infecting the nucleus of amoebae. ISME J. 8, 1634–1644. https://doi.org/10.1038/ismej.2014.5

Shu, L., Brock, D.A., Geist, K.S., Miller, J.W., Queller, D.C., Strassmann, J.E., DiSalvo, S., 2018. Symbiont location, host fitness, and possible coadaptation in a symbiosis between social amoebae and bacteria. eLife 7. https://doi.org/10.7554/eLife.42660

Steinert, M., 2011. Pathogen–host interactions in Dictyostelium, Legionella, Mycobacterium and other pathogens. Semin. Cell Dev. Biol. 22, 70–76. https://doi.org/10.1016/j.semcdb.2010.11.003

Steinert, M., Heuner, K., 2005. Dictyostelium as host model for pathogenesis. Cell. Microbiol. 7, 307–314. https://doi.org/10.1111/j.1462-5822.2005.00493.x

Sun, S., Noorian, P., McDougald, D., 2018. Dual Role of Mechanisms Involved in Resistance to Predation by Protozoa and Virulence to Humans. Front. Microbiol. 9, 1017. https://doi.org/10.3389/fmicb.2018.01017

Taylor, M., Mediannikov, O., Raoult, D., Greub, G., 2012. Endosymbiotic bacteria associated with nematodes, ticks and amoebae. FEMS Immunol. Med. Microbiol. 64, 21–31. https://doi.org/10.1111/j.1574-695X.2011.00916.x

Taylor-Brown, A., Vaughan, L., Greub, G., Timms, P., Polkinghorne, A., 2015. Twenty years of research into Chlamydia-like organisms: a revolution in our understanding of the biology and pathogenicity of members of the phylum Chlamydiae. Pathog. Dis. 73, 1–15. https://doi.org/10.1093/femspd/ftu009

Taylor-Mulneix, D.L., Hamidou Soumana, I., Linz, B., Harvill, E.T., 2017. Evolution of Bordetellae from Environmental Microbes to Human Respiratory Pathogens: Amoebae as a Missing Link. Front. Cell. Infect. Microbiol. 7. https://doi.org/10.3389/fcimb.2017.00510

Tosetti, N., Croxatto, A., Greub, G., 2014. Amoebae as a tool to isolate new bacterial species, to discover new virulence factors and to study the host–pathogen interactions. Microb. Pathog. 77, 125–130. https://doi.org/10.1016/j.micpath.2014.07.009

Zinicola, M., Lima, F., Lima, S., Machado, V., Gomez, M., Döpfer, D., Guard, C., Bicalho, R., 2015. Altered Microbiomes in Bovine Digital Dermatitis Lesions, and the Gut as a Pathogen Reservoir. PLOS ONE 10, e0120504. https://doi.org/10.1371/journal.pone.0120504

Figures

Figure 1. Unculturable symbiont prevalence in natural *D. discoideum* populations. (a) Pie charts indicate *Chlamydiae*, *Amoebophilus*, and *Chlamydiae-Amoebophilus* co-infection patterns in respective populations of *D. discoideum* isolates from across the Eastern United States. *Procabacter*, due to its low prevalence, is not included. Numbers next to pie charts indicate the number of isolates for each population. (b) Of all symbiont positive amoebae isolates, several are co-infected with more than one symbiont species, represented by the Venn diagram indicating the number of isolates infected with *Paraburkholderia, Amoebophilus,* and *Chlamydiae*.

Figure 2a. *Procabacter* full length 16S rRNA phylogeny

Figure 2b. *Amoebophilus* full length 16S rRNA phylogeny

Figure 2c *Chlamydiae* short 16S rRNA phylogeny with all haplotypes

Figure 2d. *Chlamydiae* Full length 16SrRNA phylogeny

Figure 2. Unculturable endosymbionts of *Dictyostelium* are related to unculturable endosymbionts of other amoebae. 16S rRNA phylogenies of *Procobacter* (a) *Amoebophilus* (b) and *Chlamydiae* (c & d) endosymbionts constructed using the Maximum Likelihood method in Mega7, with 1000 bootstrap replicates. Family names, where applicable are delineated in bold italics.

QS864 Uninfected

1µm

QS851 *Amoebophilus* Infected

500nm

QS886 *Chlamydiae* Infected

QS1040 *Procabacter* Infected

Figure 3. Endosymbionts have distinct morphologies in infected spores.
Transmission electron micrographs of spores from the indicated representative uninfected and infected *D. discoideum* isolates. Images on the right show close-ups of intracellular bacteria.

Figure 4. Elimination of *Amoebophilus* and *Chlamydiae* from natural hosts with antibiotics does not impact host reproductive fitness. Four uninfected, *Amoebophilus*-infected, and *Chlamydiae*-infected representative host isolates were treated with rifampicin to eliminate symbionts (a) and reproductive fitness (b) was assessed. Symbiont presence is indicated by amplification with *Amoebophilus*-specific or *Chlamydiae*-specific PCR and DNA gel electrophoresis (with Eukaryote specific gene amplification serving as an internal control) (a). Reproductive fitness was assessed by quantifying total spore productivity of uninfected, *Amoebophilus* infected, or *Chlamydiae* infected *D. discoideum* cultures with or without antibiotic pre-treatment after completion of one social cycle (b). Individual *D. discoideum* isolates are represented by point color.

Figure 5. *Amoebophilus* and *Chlamydiae* infections do not impact the reproductive fitness of hosts co-infected with *Paraburkholderia* symbionts. Reproductive fitness was quantified for four untreated and antibiotic-cured *Amoebophilus*-infected (a) and *Chlamydiae*-infected isolates (b) following social cycle completion after exposure to the indicated *Paraburkholderia* symbiont strains. Individual *D. discoideum* isolates are represented by point color.

Supplemental Files

Table S1. Table of Screened *D. discoideum* Natural Isolates. Table lists each D. discoideum isolate by identification number and location of collection. PCR amplification of Dictyostelium, Paraburkholderia, Amoebophilus, or Chlamydiae specific DNA is indicated with a 1 (for positive amplification) or 0 (for negative amplification).

Table S2. PCR Primers Used for this Study. PCR primer names, sequences, and associated references.