Novel Chlamydiae and *Amoebophilus* endosymbionts are prevalent in wild

isolates of the model social amoeba *Dictyostelium discoideum*

Authors:

Tamara S. Haselkorn[1*], Daniela Jimenez[2], Usman Bashir[2], Eleni Sallinger[1], David C.

Queller[2], Joan E. Strassmann[2], Susanne DiSalvo[3*]

Affiliations:

[1]Department of Biology, University of Central Arkansas, 201 Donaghey Avenue,

Conway, AR 72035

[2]Department of Biology, Washington University in St. Louis, One Brookings Drive St.

Louis, MO 63130

[3]Department of Biological Sciences, Southern Illinois University Edwardsville, 44 Circle

Drive, Edwardsville, IL 62026

* Corresponding authors

Department of Biological Sciences, Southern Illinois University Edwardsville, 44 Circle

Drive, Edwardsville, IL 62026; sdisalv@siue.edu

Department of Biology, University of Central Arkansas, 201 Donaghey Avenue,

Conway, AR 72035; thaselkorn@uca.edu

Running Title: Prevalent obligate endosymbionts in a social amoeba

Summary: Amoebae interact with bacteria in diverse and multifaceted ways. Amoeba

predation can serve as a selective pressure for the development of bacterial virulence

traits. Bacteria may also adapt to life inside amoebae, resulting in symbiotic relationships

(pathogenic or mutualistic). Indeed, particular lineages of obligate bacterial endosymbionts

have been found in different amoebae. Here, we screened an extensive collection of

Dictyostelium discoideum wild isolates for the presence of such bacterial symbionts using

PCR primers that identify these endosymbionts. This is the first report of obligate

symbionts in this highly-studied amoeba species. They are surprisingly common, identified

in 42% of screened isolates (N=730). Members of the Chlamydiae phylum are particularly

prevalent, occurring in 27% of the host strains. They are novel and phylogenetically

distinct. We also found *Amoebophilus* symbionts in 8% of screened isolates (N=730).

Antibiotic-cured amoebae behave similarly to their endosymbiont-infected counterparts,

suggesting that endosymbionts do not significantly impact host fitness, at least in the

laboratory. We found several natural isolates were co-infected with multiple

endosymbionts, with no obvious fitness effects of co-infection under laboratory conditions.

The high prevalence and novelty of amoeba endosymbiont clades in the model organism

D. discoideum opens the door to future research on the significance and mechanisms of

amoeba-symbiont interactions.

Introduction:

As voracious microbial predators, free-living amoebae are important shapers of their

microbial communities (Clarholm, 1981). This predatory pressure can alter the presence

and abundance of specific microbial constituents in the community (Ronn et al., 2002).

Amoeba predation is also postulated to play an important role in microbial evolution,

particularly in the evolution of bacterial virulence (Matz and Kjelleberg, 2005; Sun et al.,

2018). Bacteria that evolve strategies to avoid or survive amoeba predation would be

selected for in amoeba rich environments (Erken et al., 2013). Bacteria that are able to

enter amoeba cells (via phagocytosis or other entry mechanisms) but avoid subsequent

digestion gain access to an attractive intracellular niche. A diverse collection of

intracellular bacterial symbionts of amoebae has been found, some of which appear

pathogenic, neutral, or beneficial for their amoeba host (DiSalvo et al., 2015; Jeon,

1992; König et al., 2019; Maita et al., 2018; Shu et al., 2018; Taylor et al., 2012).   Some

of these symbionts are important human pathogenic species such as *Mycobacteria,*

Legionella, and *Chlamydia* (Boamah et al., 2017; Cardenal-Muñoz et al., 2018;

Drancourt, 2014; Gomez-Valero and Buchrieser, 2019; Paquet and Charette, 2016;

Tosetti et al., 2014). .

Bacterial endosymbionts of amoebae were first discovered in the diverse, free-living

amoeba genus *Acanthamoeba* (Horn and Wagner, 2004). These amoebae are found in

soil and freshwater environments and can protect themselves and their bacterial

associates by forming cysts under environmentally unfavorable conditions (Rodríguez-

Zaragoza, 1994). The presence of such endosymbionts was known long before they

could be specifically identified because the inability to isolate obligate associates from

their hosts made characterization challenging. Once molecular methods became

available, it was clear that these bacteria were from specific lineages in the Chlamydiae,

Proteobacteria, *Bacteriodetes,* and *Dependentiae* (Delafont et al., 2015; Horn and

Wagner, 2004; König et al., 2019; Maita et al., 2018; Samba-Louaka et al., 2019;

Schmitz-Esser et al., 2008). These endosymbionts interact with their amoeba hosts in

varied ways. Some inhabit host-derived vacuoles, while others persist in the cytoplasm.

A *Neochlamydia* symbiont can protect its host against the detrimental consequences of

Legionella bacterial infection (Ishida et al., 2014; König et al., 2019; Maita et al., 2018)

and other strains have been shown to improve the growth rate and motility of their hosts

(Okude et al., 2012). Other known amoeba endosymbionts are from the less-studied

genera *Amoebophilus* and *Procabacter*. Genome sequencing of the amoeba symbiont

Candidatus Amoebophilus asiaticus found no evidence that it could provide novel

pathways to supplement host nutrition, but other fitness affects have not been tested

(Schmitz-Esser et al., 2010).

Overall, a high proportion of sampled *Acanthamoeba* (25-80%) are infected with

endosymbionts and these serve as useful models for understanding how amoebae can

serve as training grounds for bacterial intracellular adaptation. The broader distribution

of symbiotic bacteria among other amoeba species and their prevalence in natural host

populations, however, is unclear (Schmitz-Esser et al., 2008). The potential of amoebae

to harbor such bacteria could be of concern to human health. For instance,

environmental Chlamydiae have been associated with respiratory diseases and adverse

pregnancy outcomes in humans and cattle, although the extent of their importance in

causing animal illness is unknown (Borel et al., 2018; Greub, 2009; Taylor-Brown et al.,

2015; Taylor-Brown and Polkinghorne, 2017).

Another amoeba that has been used as a model to study host-bacteria interactions is

the social amoeba, *Dictyostelium discoideum*. This amoeba, distantly related to

Acanthamoeba, is terrestrial, living and feeding as single cells in forest soils while the

bacterial food supply is plentiful. When food becomes scarce, the amoebae begin a

social cycle, which entails thousands of cells aggregating and forming a slug to crawl to

the surface of the soil. The cycle culminates in the formation of a fruiting body, made of

a ball of spores (sorus) sitting on top of a stalk, which can then be dispersed to more

favorable conditions (Kessin, 2001). In many cases, bacteria are cleared from *D.*

discoideum cells during the social cycle through innate immune-like mechanisms (Chen

et al., 2007). However, vegetative amoebae can be infected with a range of human

pathogens such as *Mycobacterium, Bordetella*, and *Legionella* (Bozzaro and Eichinger,

2011; Taylor-Mulneix et al., 2017). As a model system *D. discoideum* has been fruitfully

utilized to describe the role of host and pathogen factors in mediating infection

mechanisms and virulence (Annesley and Fisher, 2009; Bozzaro et al., 2019; Bozzaro

and Eichinger, 2011; Cosson and Soldati, 2008; Steinert, 2011; Steinert and Heuner,

2005).

Recently, a variety of bacterial symbionts have been discovered to colonize *D.*

discoideum (Brock et al., 2018; DiSalvo et al., 2015; Haselkorn et al., 2018). The *D.*

discoideum microbiome is composed of both persistent and transient bacterial

associates (Brock et al., 2018). Three different newly named *Paraburkholderia* species,

P. agricolaris, *P. hayleyella*, and *P. bonniea* are persistent bacterial symbionts that

intracellularly infect vegetative amoebae and spore cells (Brock et al., 2020, 2011;

Haselkorn et al., 2018; Khojandi et al., 2019; Shu et al., 2018). While these

Paraburkholderia themselves do not nourish *D. discoideum*, they confer farming to their

hosts by allowing food bacteria to survive in the fruiting body. These food bacteria can

then be re-seeded in new environments during spore dispersal, which benefits hosts in

food-scarce conditions (Brock et al., 2011; Haselkorn et al., 2018; Khojandi et al., 2019).

In natural populations, an average of 25% of *D. discoideum* are infected with

Paraburkholderia (Haselkorn et al., 2018). Other bacteria, some edible and others not,

have the ability to transiently associate with *D. discoideum* even in the absence of

Paraburkholderia co-infections (Brock et al., 2018). Thus, natural *D. discoideum* hosts a

small-scale microbiome and is gaining traction as a simple model system to study the

mechanisms of microbiome formation (Dinh et al., 2018; Farinholt et al., 2019).

However, all previous sampling of natural *D. discoideum* associates has relied on

isolating bacterial colonies from nutrient media plates, which would miss any bacterial

species, such as obligate amoeba symbionts, incapable of growing under these

conditions.

Since obligate endosymbionts are common in other free-living amoebae, we

endeavored to identify them in *D. discoideum.* Here, we screened a frozen stock

collection of 730 natural *D. discoideum* isolates. We first used a 16S rRNA gene

amplification strategy with "universal" primers and direct sequencing to detect and

identify bacteria within *D. discoideum* spores in a single population. In addition to

detecting expected *Paraburkholderia* symbionts, we identified many Chlamydiae and

Amoebophilus sequences*,* and a few *Procabacter* sequences in screened samples. To

determine the extent of Chlamydiae and *Amoebophilus* symbionts in our collection we

then used symbiont specific PCR screening. From each of these prevalent bacterial

lineages, we found genetically distinct novel species and were able to visualize

representative endosymbiont cells within spores via electron microscopy. We also used

antibiotics to cure natural hosts of Chlamydiae and *Amoebophilus* and compared host

fitness to uncured counterparts. Finally, we looked at the fitness effects of co-infections

since co-infections occur in nature.

Results:

<u>*Obligate endosymbionts are common in wild-collected clones of* Dictyostelium</u>

<u>discoideum</u>

To investigate the presence and prevalence of obligate bacterial symbionts in *D.*

discoideum, we screened a frozen collection of 730 natural *D. discoideum* isolates for

bacterial DNA. The vast majority of our screened isolates were collected from the

Eastern half of the United States (Fig. 1a) and had previously been screened for

Paraburkholderia (Haselkorn, *et al.* 2018, Table S1). This included 16 populations with

seven or more individuals collected. Our largest collection from Virginia consists of

about 200 individuals collected in 2000 and in 2014. Across all locations we found over

41% of samples harbored at least one obligate endosymbiont. Specifically, 117 (~16%)

were infected with *Amoebophilus* and 198 (~27%) with Chlamydiae*.* Sixteen (2.1%)

were co-infected with both *Amoebophilus* and Chlamydiae (Fig. 1b). Two isolates (both

from Virginia) were infected with an obligate bacterium identified as belonging to the

genus *Procabacter*.

The Procabacter*,* Amoebophilus*, and Chlamydiae endosymbionts of* D. discoideum *are*

novel and phylogenetically distinct from known symbionts.

For *Procabacter*, *Amoebophilus*, and the three most prevalent Chamydiae strains (those

occurring in more than one individual) we sequenced the full length 16S rRNA for our *D.*

discoideum endosymbionts to better determine their evolutionary relationships. In all

cases, the 16S rRNA haplotypes were novel and highly diverged, ranging from 6-12%

sequence difference from other sequences currently in GenBank or SILVA (Table S2).

For the *Procabacter* and *Amoebophilus* endosymbionts, only a single haplotype was

found for each. The *Procabacter* endosymbiont of *D. discoideum* is sister to all of the

Procabacter endosymbionts of *Acanthamoeba* (Fig. 2a). While it is most closely related

to this genus, forming a clade with 100% bootstrap support, the 16S rRNA sequence is

only a 93% match. A similar pattern is seen for the *Amoebophilus* symbiont of *D.*

discoideum (Fig. 2b.). This symbiont groups most closely with the putative

Amoebophilus symbiont from genome assembly of the recently sequenced

Dictyostelium polycephalum. This Dictyostelid symbiont clade is sister to the

Amoebophilus asiaticus endosymbionts of *Acanthamoeba*, and is 6% diverged at the

16S rRNA gene (Table S2).

There are 8 distinct and novel haplotypes of *D. discoideum* endosymbionts in the

Chlamydiae. For the three most prevalent haplotypes (1, 2, and 3), we sequenced the

full 16S rRNA gene to reconstruct a more resolved phylogeny which included their

nearest neighbors from the SILVA database and representatives from many

Chlamydiae families (Fig. 2c). There are currently 6 formally named and 8 proposed

Chlamydiae families (Pillonel et al., 2018; Taylor-Brown et al., 2015), and several other

recently discovered novel clades from marine sediments (Dharamshi et al., 2020). The

D. discoideum Chlamydiae endosymbionts do not appear to be closely related (<97%

sequence match) to any of these groups. Haplotypes 1 and 2 and 4 form their own well-

supported clade, grouping with the *Rhabdochlamydiaceae* and *Ca.*

Arenachlamydiaceae, sister to the *Simkaniaceae*. Many of the internal nodes on this

phylogeny have low bootstrap support, making more detailed phylogenetic inferences

impossible. This *D. discoideum* endosymbiont clade, however, does have only a 91%

identity to its closest phylogenetic neighbor and may represent a new genus or family of

Chamydiae*.* Greater than 10% sequence divergence at the full length 16S rRNA gene

has been proposed as family level difference (Everett et al., 1999), although full genome

sequences may be necessary to make this assertion. Detailed searches for this most

common Chlamydiae haplotype 1 in the IMNGS database returned 13 matches (>99%

sequence identity), mostly as "uncultured bacteria" from soil metagenomic studies

(Table S3). One such study was in the Northeast United States (Massachusetts), in a

region close to where we sampled *D. discoideum*, and it is likely that amoeba

endosymbionts are being sequenced as part of these metagenomes.

Our other Chlamydiae haplotypes are scattered throughout the phylogeny. Haplotypes 3

and 5 form a well-supported clade and may be sister to *Rhabdochlamydiaceae*/ *Ca.*

Arenachlamydiaceae/ *Simkaniaceae* families. These haplotypes have a greater than

11% nucleotide difference from its nearest neighbors on the phylogeny (and in the

SILVA and NCBI databases, Table S2) and may represent a novel family as well.

Haplotypes 6, 7 and 8 fall in different places in the phylogeny and do not group closely

with any other taxa, although these haplotypes are based on only 220bp of sequence

information so their taxonomic identity is less clear. Finally, while the newly discovered

marine Chlamydiae lineages (Dharamshi et al., 2020) were not included in this

phylogeny, when we aligned our Chlamydiae haplotypes with these sequences, the

closest match had only an 89% sequence identity.

Distribution of endosymbionts across populations

While the average overall prevalence of infection of Chlamydiae endosymbionts was

27%, it varied among populations, ranging from 0% to 75% (Table S1). The *D.*

discoideum Chlamydiae haplotypes 1, 2 and 3 were the most prevalent in natural

populations. Haplotype 1 represented 79.8% of all *D. discoideum* Chlamydiae

infections, while haplotype 2 was 13.6% and haplotype 3 was 3.5%. Haplotypes 4-8

were represented only once. Haplotype 1 was found in 12 populations, haplotype 2 was

found in 8 different populations, and haplotype 3 was found in three populations. The

diversity of Chlamydiae strains was the highest in the Virginia Mountain Lake collection

from 2000, which had six of the eight haplotypes (1-6), and the second highest diversity

of Chlamydiae haplotypes was found in the Texas Houston Arboretum collection (1, 2, 3

and 8). Interestingly, haplotype 1 was the only haplotype circulating in the most recent

2014 Virginia Mountain Lake collection. While haplotype diversity was lower in this

collection, Chlamydiae prevalence (for all haplotypes) in the region increased from 20%

in 2000 to 38% in 2014.

The one strain of *Amoebophilus* found to infect *D. discoideum* was distributed across 9

populations, although at low prevalence, often with just a single individual infected in

many locations. This endosymbiont was generally less prevalent than Chlamydiae, with

the exception of the 2000 Virginia Mountain Lake population, where it infected 37% of

the population. *Amoebophilus* prevalence in this region, however, decreased to 10% in

2014. *Amoebophilus* was also relatively prevalent in the geographically-distant Texas

Houston Arboretum population, infecting 23% of the population.

Obligate endosymbionts can be visualized within reproductive host spore cells

Transmission electron microscopy (TEM) revealed bacterial cells inside developed

spores from representative *Amoebophilus,* Chlamydiae*, and *Procabacter* host isolates

(Fig. 3). Symbiont cells were detectable in the majority of symbiont host spores

visualized by TEM. Each symbiont appeared to have gram-negative type cell walls but

displayed unique morphologies within spore cells. *Amoebophilus* endosymbionts were

rod-like, measured 0.3-0.5µm in width and 0.5-1.3µm in length, and appeared to be

distributed throughout the cytoplasm within a host membrane. *Procabacter*

endosymbionts were rod shaped, measured 0.25-0.5µm in width and 0.75-1.5µm in

length, and in many cases were surrounded by a host membrane. Chlamydiae

endosymbionts appeared as dense wrinkled spheres measuring 0.3-0.6µm that were

distributed throughout the cytoplasm.

Amoebophilus *and Chlamydiae can be eliminated from host isolates without significantly impacting host fitness*

We did not observe any obvious defects in the growth and development of Chlamydiae and *Amoebophilus* infected hosts. To better assess the impact of these endosymbionts on their hosts, we measured the reproductive fitness of representative host isolates before and after endosymbiont curing (Fig. 4). We eliminated Chlamydiae and *Amoebophilus* from four of their natural hosts through culturing for two social cycles on antibiotic saturated plates. After antibiotic exposure and resumption of development on normal media, we confirmed stable loss of endosymbionts via endosymbiont-specific PCR screening (Fig. 4a). We next cultured $1x10^5$ spores from antibiotic-treated and untreated *D. discoideum* counterparts on nutrient media with *Klebsiella pneumoniae* food bacteria. Following culmination of fruiting body development after a week of incubation, we harvested and measured total spore contents (Fig. 4b). We found no significant differences in spore productivity according to endosymbiont status (one-way analysis of variance, $(F(5,42)=1.106, p=0.372)$. Thus, symbiont infections in these natural host isolates have no significant impact on host spore productivity under these conditions. We could not assess the fitness impact of Chlamydiae or *Amoebophilus* in new host amoebae as our attempts to establish infections in new hosts have been thus far unsuccessful.

Amoebophilus *and Chlamydiae infections do not alter the fitness of hosts during exposure to* Paraburkholderia

Since co-infections with unculturable symbionts and *Paraburkholderia* occur in the wild,

we wanted to determine whether these co-infections would result in a different fitness

outcome than *Paraburkholderia* infections alone. We tested the fitness impact of

Chlamydiae and *Amoebophilus* during *Paraburkholderia* exposure using the same

strategy as above, but with the addition of 5% by volume of the indicated

Paraburkholderia strain to food bacteria prior to plating (Fig. 5). These *Paraburkholderia*

symbionts range from highly detrimental (*P. hayleyella* strain 11), to intermediately

detrimental (*P. agricolaris* strains 159 and 70), to neutral (*B. bonniea* strain 859) under

the culturing conditions used for this assay (Haselkorn et al., 2018; Khojandi et al.,

2019; Shu et al., 2018). We again observed that exposure to distinct *Paraburkholderia*

strains significantly influenced spore productivity for both *Amoebophilus* ($F_{(3,60)}=36.87$,

$p=<0.001$) and Chlamydiae ($F_{(3,60)}=28.52$, $p=<0.001$) originating host lines (Fig. 6).

However, as with single unculturable symbiont infections, we found that presence or

absence of *Amoebophilus* or Chlamydiae had no significant impact on spore

productivity during exposure to *P. hayleyella*-11 ($F_{(5,42)}=1.171$, $p=0.339$), *B.*

agricolaris-159 ($F_{(5,42)}=0.985$, $p=0.438$), *P. agricolaris*-70 ($F_{(5,42)}=1.095$, $p=0.377$), or

P. bonniea-859 ($F_{(5,42)}=0.661$, $p=0.655$) (Fig. 5). Thus, under these conditions the

fitness impact of *Paraburkholderia* symbiosis on host spore productivity does not appear

to be altered (for better or worse) by co-infections with obligate endosymbionts.

Detection of positive associations between Chlamydiae and Paraburkholderia *and*

Amoebophilus *and* Paraburkholderia *and a negative association between* Amoebophilus

and Chlamydiae

Our earlier culture-based screening of this *D. discoideum* natural collection found that

25% of isolates were infected with *Paraburkholderia* symbionts (Brock et al., 2020;

Haselkorn et al., 2018). For this new round of sequence-based screening, we identified

multiple instances of *Paraburkholderia-Amoebophilus* and *Paraburkholderia-*

Chlamydiae co-infections in addition to the *Amoebophilus*-Chlamydiae co-infections

from above. Specifically, 42 isolates (6%) were co-infected with *Paraburkholderia* and

Amoebophilus, 25 (3.5%) with *Paraburkholderia* and Chlamydiae, and 9 (1%) were co-

infected with all three symbiont types (Fig. 1b). Co-infection prevalence of the different

endosymbionts were tested in our three highest sampled populations: Texas-Houston

Arboretum, Virginia Mountain Lake 2000, and Virginia Mountain Lake 2014. Pairwise

Fisher Exact tests showed a small positive association between Chlamydiae and

Paraburkholderia in the Texas population (p=0.0132, 15 observed with 10 expected) but

not the Virginia populations. There was a positive association between

Paraburkholderia and *Amoebophilus* in the Virginia 2000 collection (p=0.000277, 43

observed with 30 expected) and a negative association between Chlamydiae and

Amoebophilus in the Virginia 2014 collection (p=0.000024, none observed with 10

expected).

Discussion:

Many microorganisms, particularly those that are obligately host-associated, are often

overlooked because they are unculturable in standard laboratory conditions. The

presence of obligate bacterial symbionts in free-living amoebae has been known for

some time, particularly for *Acanthamoeba* (Fritsche et al., 1993; Hall and Voelz, 1985;

Jeon and Jeon, 1976; Proca-Ciobanu et al., 1975). The list of amoebal endosymbionts

and their host species has since expanded but obligate endosymbionts have not been

previously described in the social amoeba *D. discoideum* (Corsaro et al., 2010; Delafont

et al., 2015; Horn and Wagner, 2004; Samba-Louaka et al., 2019; Schmitz-Esser et al.,

2008; Schulz et al., 2014)*. Here, we screened a collection of 730 *D. discoideum* isolates

using a PCR detection strategy in order to identify bacterial associates that could be

missed using conventional culturing techniques. Furthermore, to our knowledge, this is

the largest collection of amoebae from natural populations to be screened, facilitated by

the relative ease in which social amoeba fruiting bodies can be sampled. Here, we find

that obligate symbionts are quite common among *D. discoideum* collected from the wild,

with bacterial sequences of unculturable bacteria present in 301 isolates (41.2%). All

but two of these unculturable bacterial sequences aligned most closely to *Amoebophilus*

or *Chlamydiae* species. The identification of these bacteria in *D. discoideum* reveals

that this valuable eukaryotic model organism is a natural vector of environmental

bacteria. This opens a new role for *D. discoideum* as a research tool for the

investigation of how bacterial endosymbionts evolve to live inside hosts, how they affect

the fitness of their amoeba host, and how they interact with other bacteria in the

amoeba microbiome.

The intracellular morphology of endosymbionts in *D. discoideum* spores were in some

cases similar to previous observations of related endosymbionts in other amoebae.

Amoebophilus cells in *D. discoideum* spores appeared similar to previous observations

of *Amoebophilus* symbionts in *Acanthamoeba sp*. (Horn et al., 2001; Schmitz-Esser et

al., 2010). The localization of *Procabacter* endosymbionts within a membrane bound

vacuole has also been observed for Candidatus *Procabacter* sp. OEW1 in

Acanthamoeba spp. host cells (Heinz et al., 2007). However, this differs from a previous

report on *Procabacter* symbionts in *Acanthamoeba* trophozoites and cysts wherein

bacterial cells were not found to be enclosed in vacuoles (Fritsche et al., 2002). The

wrinkled morphology of Chlamydiae cells in *D. discoideum* spores is somewhat

reminiscent of the spiny structure of *Chlamydia*-like symbionts in *Acanthamoeba,*

Naegleria, and *Hartmannella* host species (Casson et al., 2008; Corsaro and Greub,

2006; Horn et al., 2000). However, these morphological observations may be artifacts of

(or exaggerated by) sample preparation prior to TEM and may be better resolved using

cryo-electron tomography (Huang et al., 2010; Santarella-Mellwig et al., 2013).

Chlamydiae-like organisms typically have two or more morphotypes that correspond to

distinct developmental stages. Elementary bodies are the infectious form that can be

transmitted extracellularly from cell to cell. Once internalized, elementary bodies

differentiate into reticulate bodies that typically replicate within inclusion vesicles,

although some environmental Chlamydiae can be found directly in the cytosol

(Bayramova et al., 2018; Benamar et al., 2017; Bou Khalil et al., 2017; Collingro et al.,

2020; Horn, 2008; Nylund et al., 2018). However, in *D. discoideum* host spores, we only

visualize one morphology that does not appear to be in an inclusion but is rather

distributed throughout the cytosol. It is unclear what morphotype this represents and

whether other morphotypes exist in *D. discoideum* prior to spore formation. It is possible

that additional Chlamydiae developmental stages occur in vegetative amoebae and that

only one form is favored within metabolically inert host spores.

Out of the 251 amoeba isolates from the 2014 Virginia Mountain Lake collection, we

identified only two infected with a bacterium with sequence homology to *Procabacter*,

suggesting that *D. discoideum* is not a preferred natural host for this endosymbiont. Yet,

this *D. discoideum Procabacter* lineage possibly represents an important link for

understanding the evolution and adaptation of *Procabacter* to amoeba hosts, as the

phylogenetic placement of the novel *D. discoideum Procabacter* lineage is sister to the

rest of the *Procabacter* species infecting *Acanthamoeba.* Comparative genome

sequencing of this clade, as well as future research into the impact of these infections

on their hosts and more extensive wild isolate screenings will better inform our

understanding of the significance of these infections in *D. discoideum.*

We identified multiple *D. discoideum* infected with an *Amoebophilus* species that most

closely matched Ca. *Amoebophilus asiaticus*, an endosymbiont originally discovered in

Acanthamoeba (Horn et al., 2001). Similar to the *D. discoideum Procabacter* symbionts,

the *D. discoideum Amoebophilus* symbiont lineage is sister to the rest of the

Amoebophilus clade. The fact that it groups most closely with a putative *Amoebophilus*

symbiont discovered in the genome assembly of *Coremiostelium polycephalum,*

formerly called *Dictyostelium polycephalum* (Sheikh et al., 2018) suggests that there

may be some host specificity of *Amoebophilus* among social amoebae, although

sampling of additional amoeba species would be necessary to elucidate this pattern.

The next most closely related bacteria to the *Amoebophilus* genus are the insect

facultative endosymbionts of the *Cardinium* genus (Santos-Garcia et al., 2014). These

bacteria infect up to 7% of all insect species and have diverse effects on their hosts

ranging from reproductive parasitism to nutritional supplementation and defense against

parasites. Comparative genome sequencing has uncovered potential routes of

adaptation from free-living marine bacteria to either an insect or amoeba host (Santos-

Garcia et al., 2014). Including this additional *Amoebophilus* species in comparative

genome analysis could further elucidate this process and highlight potential amoeba

species-specific adaptations. *Acanthamoeba* and *Dictyostelium* are evolutionarily

distant hosts with different lifestyles, thus specific bacterial adaptations may have

evolved.

The global significance of *Amoebophilus* infections is under active investigation.

Amoebophilus have been associated with coral microbiomes and bovine digital

dermatitis lesions, which may simply be an indicator that their host amoebae are

present in these samples (Apprill et al., 2016; Choi et al., 2009; Horn et al., 2001;

Huggett and Apprill, 2019; Zinicola et al., 2015). In amoebae, the full extent of this

bacterium's affect is unknown. The moderately reduced genome of Ca. *Amoebophilus*

asiaticus lacks important metabolic pathways but contains a high number of proteins

with eukaryotic domains that are thought to be essential for host cell interactions

(Schmitz-Esser et al., 2010). Genes associated with host cell exploitation, such as

toxins, antimicrobial factors, and a unique secretion system, suggest that Ca.

Amoebophilus asiaticus is more parasitic than mutualistic (Böck et al., 2017; Penz et al.,

2010; Schmitz-Esser et al., 2010). We did not observe an obvious fitness cost of hosting

Amoebophilus via our spore productivity assays. However, this measures just one

aspect of host fitness in defined laboratory conditions and it remains possible that these

symbionts may impact host fitness in important ways at different life stages or under

different environmental conditions.

Chlamydiae endosymbionts were the most prevalent in the screened collection, with

approximately 27% of *D. discoideum* isolates infected. Environmental Chlamydiae

appear to be diverse and ubiquitous and many Chlamydiae endosymbionts associate

with a wide range of protozoa and other organisms (Corsaro et al., 2013; Corsaro and

Venditti, 2006; Coulon et al., 2012). The incredible genetic diversity within the phylum

Chlamydiae suggests ancient interactions with eukaryotes, with estimates that the

association is 700 million years old (Greub and Raoult, 2003). The diversity of

Chlamydiae, as estimated from various 16S rRNA genome databases and sequence

read archives, ranges anywhere from 1157-1483 different families (Collingro et al.,

2020). This type of screening, however, does not identify the hosts, so insight into

adaptation to particular hosts is limited. In this study we have identified eight novel

lineages, potentially representing two novel families, both with closely related

haplotypes suggesting ongoing evolution within the *D. discoideum* host. The separation

of these novel *D. discoideum* Chlamydiae lineages from the other amoeba lineages

suggests a long-standing association and possible co-adaptation. As our attempts to

transfer *Amoebophilus* or Chlamydiae endosymbionts to new *D. discoideum* hosts were

unsuccessful, we could not probe the potential role of host-symbiont adaptation in

fitness outcomes. This strengthens the speculation that these symbionts have been

associated with their specific host lines for a long period of time but also raises the

question of how they were originally acquired. Other lab experiments have shown that

some of these Chlamydiae strains are easily transferred across different host genera

while others cannot be used to successfully re-infect even their original host amoebae

(Corsaro et al., 2013; Coulon et al., 2012; Okude et al., 2012). A previous transfer of a

Protochlamydia strain to new hosts resulted in a decreased growth rate, suggesting

mutual co-evolution between the endosymbiont and its original host (Okude et al.,

2012).

Similar to our results for *Amoebophilus*-infected hosts, we did not find any obvious

fitness costs from Chlamydiae infection of *D. discoideum* under standard lab conditions

for a subset of *D. discoideum* isolates. This does not, however, rule out the possibility

that these symbionts may impart significant impacts on their hosts which may be life-

stage, environmental, or genotypically context-dependent (Taylor-Brown et al., 2015).

Studies of the impact of *Chlamydiae* endosymbionts on their amoeba hosts suggest that

the range of outcomes spans the parasite to mutualist continuum (Hayashi et al., 2012).

The high prevalence and widespread distribution of particular haplotypes, like haplotype

1, may suggest ecological relevance. It is unclear whether the abundance of particular

endosymbiont haplotypes represents an epidemic of a new genotype or selective

sweeps due to beneficial effects on their hosts under particular environmental

conditions, as seen in insect symbionts (Cockburn et al., 2013; Himler et al., 2011).

Depending on the temperature, one *Parachlamydia* endosymbiont of *Acanthamoeba*

can either be lytic or endosymbiotic (Greub et al., 2003). The Chlamydiae endosymbiont

S13 does not significantly harm its *Acanthamoeba* host and instead has been shown to

protect it from *Legionella pneumophila* infection (Ishida et al., 2014; König et al., 2019;

Maita et al., 2018; Matsuo et al., 2009; Okude et al., 2012). A *Protochlamydia*

endosymbiont has been shown to increase the migration rate, actin density, and growth

rate of its native host but had an adverse effect when transferred to non-native hosts

(Okude et al., 2012). Overall, future investigations into the impact of these symbionts on

hosts using diverse fitness assays and a range of culture conditions will help to

elucidate their significance in infected host amoebae.

At least 11.4% (83 out of 730) of these *D. discoideum* are co-infected with some

combination of *Chlamydiae*, *Amoebophilus,* or *Paraburkholderia*. The lack of detectable

fitness costs from co-infection with unculturable endosymbionts and *Paraburkholderia* in

the laboratory is consistent with random associations found in several populations. In

different environments, however, other microbe-microbe interactions may have an effect

that favors certain combinations of symbionts, as is seen in aphids (Rock et al., 2018).

Chlamydiae occur in co-infections less often than would be expected with

Amoebophilus in one *D. discoideum* population, consistent with the observation that

Chlamydiae interact negatively with the *Legionella* pathogen in *Acanthamoeba*. On the

other hand, *Paraburkholderia* and *Amoebophilus* are positively associated in one

population. *Paraburkholderia* symbionts of *D. discoideum* have been shown to allow

other food bacteria to enter into the amoeba spores (Brock et al 2011, DiSalvo et al

2015), and perhaps a similar positive interaction occurs in this case. This effect may

outweigh some of the negative interactions of Chlamydiae with other bacteria, as in one

population there was positive association with Chlamydiae and *Paraburkholderia*.

Paraburkholderia symbionts are easily transferred to new hosts via co-culturing and can

be found both intracellularly and extracellularly in amoeba cells and structures (DiSalvo

et al., 2015; Haselkorn et al., 2018; Shu et al., 2018). *Paraburkholderia* infections in new

hosts may be detrimental or beneficial depending on symbiont genotype and

environmental context (Haselkorn et al., 2018; Khojandi et al., 2019). Endosymbiont co-

infection is not unique to *D. discoideum,* as previous and alternative isolation methods

are allowing for identification of co-infections in *Acanthamoeba* (Heinz et al., 2007;

Lagkouvardos et al., 2014).

The growing list of bacterial symbionts of free-living amoebae attests to their attractiveness

as an intracellular niche for bacterial symbionts. The similarities between amoebae and

animal macrophages, as well as the shared intracellular entrance and survival

mechanisms employed by endosymbionts and pathogens, further highlight the potential

medical and research relevancy of amoebae-symbiosis systems. The discovery of a high

prevalence of infection and novel lineages of common amoebae symbiont clades in the

model organism *D. discoideum* opens the door to future research on the ecological

significance and mechanisms of amoeba-symbiont interactions.

Methods:

Bacterial Strains and Cultural Conditions:

We used *Klebsiella pneumoniae* throughout as our standard *D. discoideum* food supply.

For *Paraburkholderia*-unculturable co-infection experiments, we used the

Paraburkholderia symbionts *P. hayleyella*-11, *P. agricolaris*-70, *P. agricolaris*-159, and

B. bonniea-859. We grew bacteria on SM/5 agar plates [2 g glucose, 2 g BactoPeptone,

2 g yeast extract, 0.2 g $MgCl_2$, 1.9 g $KHPO_4$, 1 g $K2HPO_5$, and 15 g agar per liter,

premixed from Formedium] at room temperature before resuspending them in KK2

buffer [2.25 g $KH2PO_4$ (Sigma-Aldrich) and 0.67 g $K2HPO_4$ (Fisher Scientific) per liter].

We then adjusted bacterial suspension to a final OD_{600nm} of 1.5. We plated amoebae

spores with 200μl of bacterial suspensions for all culturing conditions.

<u>Symbiont Screening</u>

Total DNA from *D. discoideum* was harvested by suspending 15-25 sori in a 5% chelex

solution with 10μg proteinase K, incubating samples for 240 minutes at 56°C, 30

minutes at 98°C, and removing DNA supernatant from chelex beads. PCR with

eukaryotic specific primers was run as a DNA extraction control, and symbiont specific

primers were used to detect symbionts in each sample (Table S2) (Corsaro et al., 2002;

Ossewaarde and Meijer, 1999). For all primer sets we used a touchdown PCR protocol

consisting of 94°C for 30 s, 68°C-55°C for 45 x (dropping 1 degree per cycle) for 14

cycles, and 72°C for 1 min, followed by 94°C, 55°C, 72°C for 20 cycles. To confirm

infection we ran PCR samples on a one percent agarose gel with TAE buffer and

imaged on a transilluminator. PCR products were sent for sequencing (Genewiz), and

the resulting Sanger sequences were cleaned using the program Geneious v8

(http://www.geneious.com (Kearse et al., 2012)). We use NCBI Blast to initially identify

the bacterial species/strain based on highest sequence similarity. Sequences have

been deposited to GenBank under accession numbers (MT596490-MT596496).

<u>Amoebae Clone Types and Culture Conditions:</u>

Amoebae were cultured from freezer stocks by resuspending frozen spores onto SM/5

agar plates with 200µl of food *K. pneumoniae* suspensions. Plates were incubated at

room temperature under low lights and ~60% humidity for approximately 1 week (or until

fruiting body development).

For antibiotic curing and spore quantification, we used four uninfected, four

Amoebophilus infected, and four Chlamydiae infected host lines. Uninfected

representatives were QS864, QS970, QS1003 and QS1069 (highlighted in yellow in

Table S1). *Amoebophilus* infected representatives included QS851, QS1002, QS1011,

and QS1017 (highlighted in green in Table S1). Chlamydiae infected representatives

included QS866, QS991, QS1023, and QS1089 (highlighted in blue in Table S1). For all

experiments, we initiated experimental conditions from pre-developed fruiting bodies by

suspending ~8-15 sorus contents in KK2. We then plated 10^5 spores onto an SM/5 plate

with 200µl of bacterial suspensions (*K. pneumoniae* only or with 95% *K. pneumoniae/*

5% *Paraburkholderia* mixtures by volume for co-infection experiments). Plates were

then incubated as described above for the time indicated for each experimental

condition.

<u>Curing of Symbiont Infections:</u>

We treated uninfected, Chlamydiae-infected, and *Amoebophilus* infected representative

isolates with rifampacin by plating ~10^2 spores with 400µl *K. pneumoniae* on SM/5

medium containing 30µg/ml rifampacin. Plates were incubated until fruiting body

formation. Sori from grown fruiting bodies were then harvested and replated for a

second round of antibiotic exposure (using the same strategy as described above).

Following fruiting body formation, spores were harvested and transferred to SM/5 with

K. pneumoniae and incubated. Following fruiting body formation on non-antibiotic

plates, we extracted total sorus content DNA and checked for symbiont infection using

the Chlamydiae-specific and *Amoebophilus*-specific PCR protocol described above.

<u>Total Spore Productivity Assay:</u>

To determine total spore productivity of antibiotic treated and untreated *D. discoideum*

isolates we plated 10^5 spores on SM/5 agar with 200µl of *K. pneumoniae* and incubated

at room temperature. Five days after plating, total plate contents were collected into

~10ml KK$_2$ + NP-40 alternative detergent solution. Solutions were then diluted

appropriately (~20x) and spores quantified via hemocytometer counts. We measured

and averaged over at least three replicates. To determine spore productivity after

exposure to *Paraburkholderia*, we conducted the assay as described but plated spores

with 200µl of a 95/5% by volume mixture of *K. pneumoniae/Paraburkholderia* cell

suspensions.

<u>Statistics</u>

Significant positive or negative associations between co-infecting endosymbionts were

determined using a Fisher Exact test (2x2 contingency table) on pairwise comparisons

in our three most highly sampled populations. Statistical analyses for spore productivity

experiments were performed in R 3.6.0. Comparisons were done using one-way

ANOVA's with antibiotic treatment as the independent variable.

<u>Transmission Electron Microscopy</u>

Complete processing and imaging of spores by transmission electron microscopy was

performed by Wandy Beatty at the Washington University Molecular Microbiology

Imaging Facility. *D. discoideum* isolates imaged included QS864, QS851, QS886,

QS1040. Briefly, spores were harvested in KK2 buffer and fixed with 2%

paraformaldehyde with 2.5% glutaraldehyde (Polysciences Inc., Warrington, PA) in 100

mM cacodylate buffer (pH 7.2) for 1-3 hrs. at room temperature. We washed samples

with cacodylate buffer and postfixed in 1% osmium tetroxide (Polysciences Inc.) for 1 hr

before rinsing in dH_2O prior to *en bloc* staining with 1% aqueous uranyl acetate (Ted

Pella Inc., Redding, CA) for 1 hr. We dehydrated samples in increasing concentrations

of ethanol and embedded them in Eponate 12 resin (Ted Pella Inc.). Sections of 95 nm

were cut with a Leica Ultracut UCT ultramicrotome (Leica Microsystems Inc.,

Bannockburn, IL), stained with uranyl acetate and lead citrate, and viewed on a JEOL

1200 EX transmission electron microscope (JEOL USA Inc., Peabody, MA) equipped

with an AMT 8 megapixel digital camera (Advanced Microscopy Techniques, Woburn,

MA).

<u>Phylogenetic Tree Construction</u>

The *Procabacter* full length 16S rRNA phylogeny was created using the top ten NCBI

Blast results for the *Procabacter* endosymbiont of D. *discoideum* sequence and

representative species from the phylum *Betaproteobacteria*. The species were chosen

using the NCBI taxonomy database and 16S rRNA sequences were obtained from the

NCBI nucleotide database as well as the three top hits to the SILVA database. A

sequence from *Rickettsia rickettsii* was also obtained from the NCBI nucleotide

database to serve as an outgroup. Sequences were imported into Mega7 (Tamura et

al, 2013) and aligned with MUSCLE. All sequences were then trimmed to a uniform

1396 bp. The Kimura 2-parameter model with a discrete Gamma distribution was

selected using the "Find Best DNA/Protein Models" command in Mega7. A Maximum

Likelihood tree was then constructed with 1000 bootstrap replicates.

The *Amoebophilus* full length 16S rRNA phylogeny was created using the top five NCBI

Blast results for the *Amoebophilus* endosymbiont of *D. discoideum* sequence, a

sequence from the recently sequenced *Dictyostelium polycephalum* genome assembly,

a sequence from the related endosymbiont *Candidatus Cardinium hertigii*, and

representative species from the order *Cytophagales*. The species were chosen using

the phylogeny in Hahnke et al., (2016) as a guide, and full length 16S rRNA sequences

were obtained from the NCBI nucleotide database (Hahnke et al., 2016) as well as the

three top hits from the SILVA database. A sequence from *Bacteroides fragilis* was also

obtained from the NCBI nucleotide database to serve as an outgroup. Sequences were

imported into Mega7, aligned with MUSCLE, and trimmed to a uniform 1364 bp. The

Kimura 2-parameter model with a discrete Gamma distribution was selected using the

"Find Best DNA/Protein Models" command in Mega7. A Maximum Likelihood tree was

then constructed with 1000 bootstrap replicates.

The Chlamydiae full length 16S rRNA phylogeny was created using 1-2 representative

species from the nine families in the phylum Chlamydiae from Taylor-Brown et al.,

(2015), 1 from *Ca.* Arenachlamydiaceae (Pillonel et al. 2018), and the three top hits for

each haplotype from the SILVA database. Full length 16S rRNA sequences of chosen

species were obtained from both the NCBI nucleotide and SILVA databases.

Sequences were imported into Mega7 and aligned with MUSCLE, and then trimmed to

1354 bp. The Tamura 3-parameter model with a discrete Gamma distribution was

selected for the short tree, and the General Time Reversal model with a discrete

Gamma distribution and invariant sites was selected for the full-length tree using the

"Find Best DNA/Protein Models" command in Mega7. A Maximum Likelihood tree was

then constructed for each with 1000 bootstrap replicates.

<u>Verification of symbiont strain novelty</u>

In order to verify that, to the best of our knowledge, these *D. discoideum* Chlamydiae

lineages are novel, we searched the SILVA database (in addition to the NCBI Blast

database) for nearest neighbors. A table of the sequence identities to the SILVA

database for all of our full length 16S rRNA endosymbiont haplotypes is included as

Supplemental Table 2. Two recent papers (Pillonel et al., 2018 and Dharamshi et al.,

2020) have newly discovered Chlamydiae lineages so we compared our 16S rRNA

sequences to theirs where possible. The phylogenetic analyses from the Pillonel et al.

paper were done from metagenomic sequences, which often lack identified 16S rRNA

genes, and thus it was not always possible to compare directly. We did use BLAST to

query our sequences against these metagenomes in NCBI and in no case did we find a

match with >85% sequence identity, if a match was found at all. For the Dharamshi et

al. paper we separately aligned our sequences (full length Chlamydiae haplotypes 1-3)

with their publicly available 16S rRNA sequences. We ran a Neighbor-joining

phylogenetic analysis in MEGA 7 and computed a pairwise distance matrix. Finally, we

searched the IMNGS database (Lagkouvardos et al. 2016), which is designed to query

sequence read archives from NCBI for 16S rRNA sequences, for matches to our most

common *D. discoideum* Chlamydiae haplotype.

Acknowledgements: We thank members of the Haselkorn, DiSalvo, and Strassmann-

Queller labs for their support and suggestions, in particular Debbie Brock. We also

thank the many people in the group that collected the samples, in particular Debbie

Brock, Tom Platt and Angelo Fortunato. We also thank Dierdra Renfroe for her attempts

to transfer *Chlamydiae* symbionts to new host amoebae. This material is based upon

work supported by the National Science Foundation under grant numbers DEB-

1753743 and IOS-1656756 to JES and DCQ.

References:
Annesley, S.J., Fisher, P.R., 2009. Dictyostelium discoideum—a model for many reasons. Mol.
Cell. Biochem. 329, 73–91. https://doi.org/10.1007/s11010-009-0111-8
Apprill, A., Weber, L.G., Santoro, A.E., 2016. Distinguishing between Microbial Habitats Unravels
Ecological Complexity in Coral Microbiomes. mSystems 1, e00143-16,
/msys/1/5/e00143-16.atom. https://doi.org/10.1128/mSystems.00143-16
Bayramova, F., Jacquier, N., Greub, G., 2018. Insight in the biology of Chlamydia-related
bacteria. Microbes Infect. 20, 432–440. https://doi.org/10.1016/j.micinf.2017.11.008

Benamar, S., Bou Khalil, J.Y., Blanc-Tailleur, C., Bilen, M., Barrassi, L., La Scola, B., 2017. Developmental Cycle and Genome Analysis of Protochlamydia massiliensis sp. nov. a New Species in the Parachlamydiacae Family. Front. Cell. Infect. Microbiol. 7, 385. https://doi.org/10.3389/fcimb.2017.00385

Boamah, D.K., Zhou, G., Ensminger, A.W., O'Connor, T.J., 2017. From Many Hosts, One Accidental Pathogen: The Diverse Protozoan Hosts of Legionella. Front. Cell. Infect. Microbiol. 7. https://doi.org/10.3389/fcimb.2017.00477

Böck, D., Medeiros, J.M., Tsao, H.-F., Penz, T., Weiss, G.L., Aistleitner, K., Horn, M., Pilhofer, M., 2017. In situ architecture, function, and evolution of a contractile injection system. Science 357, 713–717. https://doi.org/10.1126/science.aan7904

Borel, N., Polkinghorne, A., Pospischil, A., 2018. A Review on Chlamydial Diseases in Animals: Still a Challenge for Pathologists? Vet. Pathol. 55, 374–390. https://doi.org/10.1177/0300985817751218

Bou Khalil, J.Y., Benamar, S., Di Pinto, F., Blanc-Tailleur, C., Raoult, D., La Scola, B., 2017. Protochlamydia phocaeensis sp. nov., a new Chlamydiales species with host dependent replication cycle. Microbes Infect. 19, 343–350. https://doi.org/10.1016/j.micinf.2017.02.003

Bozzaro, S., Buracco, S., Peracino, B., Eichinger, L., 2019. Dictyostelium Host Response to Legionella Infection: Strategies and Assays, in: Buchrieser, C., Hilbi, H. (Eds.), Legionella. Springer New York, New York, NY, pp. 347–370. https://doi.org/10.1007/978-1-4939-9048-1_23

Bozzaro, S., Eichinger, L., 2011. The professional phagocyte Dictyostelium discoideum as a model host for bacterial pathogens. Curr. Drug Targets 12, 942–954. https://doi.org/10.2174/138945011795677782

Brock, D.A., Douglas, T.E., Queller, D.C., Strassmann, J.E., 2011. Primitive agriculture in a social amoeba. Nature 469, 393–396. https://doi.org/10.1038/nature09668

Brock, D.A., Haselkorn, T.S., Garcia, J.R., Bashir, U., Douglas, T.E., Galloway, J., Brodie, F., Queller, D.C., Strassmann, J.E., 2018. Diversity of free-living environmental bacteria and their interactions with a bactivorous amoeba. Front. Cell. Infect. Microbiol. 8, 411–411. https://doi.org/10.3389/fcimb.2018.00411

Brock, D.A., Noh, S., Hubert, A.N.M., Haselkorn, T.S., DiSalvo, S., Suess, M.K., Bradley, A.S., Tavakoli-Nezhad, M., Geist, K.S., Queller, D.C., Strassmann, J.E., 2020. Endosymbiotic adaptations in three new bacterial species associated with *Dictyostelium discoideum* : *Paraburkholderia agricolaris* sp. nov., *Paraburkholderia hayleyella* sp. nov., and *Paraburkholderia bonniea* sp. nov. PeerJ 8, e9151. https://doi.org/10.7717/peerj.9151

Cardenal-Muñoz, E., Barisch, C., Lefrançois, L.H., López-Jiménez, A.T., Soldati, T., 2018. When Dicty Met Myco, a (Not So) Romantic Story about One Amoeba and Its Intracellular Pathogen. Front. Cell. Infect. Microbiol. 7. https://doi.org/10.3389/fcimb.2017.00529

Casson, N., Michel, R., Müller, K.-D., Aubert, J.D., Greub, G., 2008. *Protochlamydia naegleriophila* as Etiologic Agent of Pneumonia. Emerg. Infect. Dis. 14, 168–172. https://doi.org/10.3201/eid1401.070980

Chen, G., Zhuchenko, O., Kuspa, A., 2007. Immune-like phagocyte activity in the social amoeba. Science 317, 678–681. https://doi.org/10.1126/science.1143991

Choi, S.H., Cho, M.K., Ahn, S.C., Lee, J.E., Lee, J.S., Kim, D.-H., Xuan, Y.-H., Hong, Y.C., Kong, H.H.,
Chung, D.I., Yu, H.S., 2009. Endosymbionts of Acanthamoeba Isolated from Domestic
Tap Water in Korea. Korean J. Parasitol. 47, 337.
https://doi.org/10.3347/kjp.2009.47.4.337
Clarholm, M., 1981. Protozoan grazing of bacteria in soil—impact and importance. Microb. Ecol.
7, 343–350. https://doi.org/10.1007/BF02341429
Cockburn, S.N., Haselkorn, T.S., Hamilton, P.T., Landzberg, E., Jaenike, J., Perlman, S.J., 2013.
Dynamics of the continent-wide spread of a *Drosophila* defensive symbiont. Ecol. Lett.
16, 609–616. https://doi.org/10.1111/ele.12087
Collingro, A., Köstlbacher, S., Horn, M., 2020. Chlamydiae in the Environment. Trends Microbiol.
28, 877–888. https://doi.org/10.1016/j.tim.2020.05.020
Corsaro, D., Greub, G., 2006. Pathogenic Potential of Novel Chlamydiae and Diagnostic
Approaches to Infections Due to These Obligate Intracellular Bacteria. Clin. Microbiol.
Rev. 19, 283–297. https://doi.org/10.1128/CMR.19.2.283-297.2006
Corsaro, D., Müller, K.-D., Wingender, J., Michel, R., 2013. "Candidatus Mesochlamydia
elodeae" (Chlamydiae: Parachlamydiaceae), a novel chlamydia parasite of free-living
amoebae. Parasitol. Res. 112, 829–838. https://doi.org/10.1007/s00436-012-3213-2
Corsaro, D., Pages, G.S., Catalan, V., Loret, J.-F., Greub, G., 2010. Biodiversity of amoebae and
amoeba-associated bacteria in water treatment plants. Int. J. Hyg. Environ. Health 213,
158–166. https://doi.org/10.1016/j.ijheh.2010.03.002
Corsaro, D., Venditti, D., 2006. Diversity of the Parachlamydiae in the Environment. Crit. Rev.
Microbiol. 32, 185–199. https://doi.org/10.1080/10408410601023102
Corsaro, D., Venditti, D., Valassina, M., 2002. New parachlamydial 16S rDNA phylotypes
detected in human clinical samples. Res. Microbiol. 153, 563–567.
https://doi.org/10.1016/S0923-2508(02)01369-4
Cosson, P., Soldati, T., 2008. Eat, kill or die: when amoeba meets bacteria. Curr. Opin. Microbiol.
11, 271–276. https://doi.org/10.1016/j.mib.2008.05.005
Coulon, C., Eterpi, M., Greub, G., Collignon, A., McDonnell, G., Thomas, V., 2012. Amoebal host
range, host-free survival and disinfection susceptibility of environmental *Chlamydiae* as
compared to *Chlamydia trachomatis*. FEMS Immunol. Med. Microbiol. 64, 364–373.
https://doi.org/10.1111/j.1574-695X.2011.00919.x
Delafont, V., Samba-Louaka, A., Bouchon, D., Moulin, L., Héchard, Y., 2015. Shedding light on
microbial dark matter: a TM6 bacterium as natural endosymbiont of a free-living
amoeba: TM6 bacterium as endosymbiont of amoeba. Environ. Microbiol. Rep. 7, 970–
978. https://doi.org/10.1111/1758-2229.12343
Dharamshi, J.E., Tamarit, D., Eme, L., Stairs, C.W., Martijn, J., Homa, F., Jørgensen, S.L., Spang,
A., Ettema, T.J.G., 2020. Marine Sediments Illuminate Chlamydiae Diversity and
Evolution. Curr. Biol. 30, 1032-1048.e7. https://doi.org/10.1016/j.cub.2020.02.016
Dinh, C., Farinholt, T., Hirose, S., Zhuchenko, O., Kuspa, A., 2018. Lectins modulate the
microbiota of social amoebae. Science 361, 402–406.
https://doi.org/10.1126/science.aat2058
DiSalvo, S., Haselkorn, T.S., Bashir, U., Jimenez, D., Brock, D.A., Queller, D.C., Strassmann, J.E.,
2015. Burkholderia bacteria infectiously induce the proto-farming symbiosis of

757   Dictyostelium amoebae and food bacteria. Proc. Natl. Acad. Sci. U. S. A. 112, E5029–
758   E5037. https://doi.org/10.1073/pnas.1511878112
759 Drancourt, M., 2014. Looking in amoebae as a source of mycobacteria. Microb. Pathog. 77,
760   119–124. https://doi.org/10.1016/j.micpath.2014.07.001
761 Erken, M., Lutz, C., McDougald, D., 2013. The Rise of Pathogens: Predation as a Factor Driving
762   the Evolution of Human Pathogens in the Environment. Microb. Ecol. 65, 860–868.
763   https://doi.org/10.1007/s00248-013-0189-0
764 Everett, K.D.E., Bush, R.M., Andersen, A.A., 1999. Emended description of the order
765   Chlamydiales, proposal of Parachlamydiaceae fam. nov. and Simkaniaceae fam. nov.,
766   each containing one monotypic genus, revised taxonomy of the family Chlamydiaceae,
767   including a new genus and five new species, and standards for the identification of
768   organisms. Int. J. Syst. Evol. Microbiol. 49, 415–440. https://doi.org/10.1099/00207713-
769   49-2-415
770 Farinholt, T., Dinh, C., Kuspa, A., 2019. Microbiome management in the social amoeba
771   Dictyostelium discoideum compared to humans. Int. J. Dev. Biol. 63, 447–450.
772   https://doi.org/10.1387/ijdb.190240ak
773 Fritsche, T.R., Gautom, R.K., Seyedirashti, S., Bergeron, D.L., Lindquist, T.D., 1993. Occurrence of
774   bacterial endosymbionts in Acanthamoeba spp. isolated from corneal and
775   environmental specimens and contact lenses. J. Clin. Microbiol. 31, 1122–1126.
776 Fritsche, T.R., Linner, T., Wagner, M., Gautom, R.K., Harzenetter, M.D., Horn, M., 2002. Obligate
777   bacterial endosymbionts of Acanthamoeba spp. related to the beta-Proteobacteria:
778   proposal of "Candidatus Procabacter acanthamoebae" gen. nov., sp. nov. Int. J. Syst.
779   Evol. Microbiol. 52, 599–605. https://doi.org/10.1099/00207713-52-2-599
780 Gomez-Valero, L., Buchrieser, C., 2019. Intracellular parasitism, the driving force of evolution of
781   Legionella pneumophila and the genus Legionella. Genes Immun. 20, 394–402.
782   https://doi.org/10.1038/s41435-019-0074-z
783 Greub, G., 2009. Parachlamydia acanthamoebae, an emerging agent of pneumonia. Clin.
784   Microbiol. Infect. 15, 18–28. https://doi.org/10.1111/j.1469-0691.2008.02633.x
785 Greub, G., Raoult, D., 2003. History of the ADP/ATP-Translocase-Encoding Gene, a Parasitism
786   Gene Transferred from a Chlamydiales Ancestor to Plants 1 Billion Years Ago. Appl.
787   Environ. Microbiol. 69, 5530–5535. https://doi.org/10.1128/AEM.69.9.5530-5535.2003
788 Greub, G., Scola, B., Raoult, D., 2003. *Parachlamydia acanthamoeba* Is Endosymbiotic or Lytic
789   for *Acanthamoeba polyphaga* Depending on the Incubation Temperature. Ann. N. Y.
790   Acad. Sci. 990, 628–634. https://doi.org/10.1111/j.1749-6632.2003.tb07437.x
791 Hahnke, R.L., Meier-Kolthoff, J.P., García-López, M., Mukherjee, S., Huntemann, M., Ivanova,
792   N.N., Woyke, T., Kyrpides, N.C., Klenk, H.-P., Göker, M., 2016. Genome-Based Taxonomic
793   Classification of Bacteroidetes. Front. Microbiol. 7, 2003.
794   https://doi.org/10.3389/fmicb.2016.02003
795 Hall, J., Voelz, H., 1985. Bacterial Endosymbionts of Acanthamoeba sp. J. Parasitol. 71, 89.
796   https://doi.org/10.2307/3281982
797 Haselkorn, T.S., DiSalvo, S., Miller, J.W., Bashir, U., Brock, D.A., Queller, D.C., Strassmann, J.E.,
798   2018. The specificity of Burkholderia symbionts in the social amoeba farming symbiosis:
799   prevalence, species, genetic and phenotypic diversity. Mol. Ecol. 0.
800   https://doi.org/10.1111/mec.14982

Hayashi, Y., Yimin, Matsuo, J., Nakamura, S., Kunichika, M., Yoshida, M., Takahashi, K., Yamaguchi, H., 2012. A domino-like chlamydial attachment process: concurrent Parachlamydia acanthamoebae attachment to amoebae is required for several amoebal released molecules and serine protease activity. Microbiology 158, 1607–1614. https://doi.org/10.1099/mic.0.057190-0

Heinz, E., Kolarov, I., Kästner, C., Toenshoff, E.R., Wagner, M., Horn, M., 2007. An Acanthamoeba sp. containing two phylogenetically different bacterial endosymbionts. Environ. Microbiol. 9, 1604–1609. https://doi.org/10.1111/j.1462-2920.2007.01268.x

Himler, A.G., Adachi-Hagimori, T., Bergen, J.E., Kozuch, A., Kelly, S.E., Tabashnik, B.E., Chiel, E., Duckworth, V.E., Dennehy, T.J., Zchori-Fein, E., Hunter, M.S., 2011. Rapid Spread of a Bacterial Symbiont in an Invasive Whitefly Is Driven by Fitness Benefits and Female Bias. Science 332, 254–256. https://doi.org/10.1126/science.1199410

Horn, M., 2008. *Chlamydiae* as Symbionts in Eukaryotes. Annu. Rev. Microbiol. 62, 113–131. https://doi.org/10.1146/annurev.micro.62.081307.162818

Horn, M., Harzenetter, M.D., Linner, T., Schmid, E.N., Muller, K.-D., Michel, R., Wagner, M., 2001. Members of the Cytophaga-Flavobacterium-Bacteroides phylum as intracellular bacteria of acanthamoebae: proposal of "Candidatus Amoebophilus asiaticus." Environ. Microbiol. 3, 440–449. https://doi.org/10.1046/j.1462-2920.2001.00210.x

Horn, M., Wagner, M., 2004. Bacterial Endosymbionts of Free-living Amoebae1. J. Eukaryot. Microbiol. 51, 509–514. https://doi.org/10.1111/j.1550-7408.2004.tb00278.x

Horn, M., Wagner, M., Müller, K.-D., Schmid, E.N., Fritsche, T.R., Schleifer, K.-H., Michel, R., 2000. Neochlamydia hartmannellae gen. nov., sp. nov. (Parachlamydiaceae), an endoparasite of the amoeba Hartmannella vermiformis. Microbiology, 146, 1231–1239.

Huang, Z., Chen, M., Li, K., Dong, X., Han, J., Zhang, Q., 2010. Cryo-electron tomography of Chlamydia trachomatis gives a clue to the mechanism of outer membrane changes. J. Electron Microsc. (Tokyo) 59, 237–241. https://doi.org/10.1093/jmicro/dfp057

Huggett, M.J., Apprill, A., 2019. Coral microbiome database: Integration of sequences reveals high diversity and relatedness of coral-associated microbes. Environ. Microbiol. Rep. 11, 372–385. https://doi.org/10.1111/1758-2229.12686

Ishida, K., Sekizuka, T., Hayashida, K., Matsuo, J., Takeuchi, F., Kuroda, M., Nakamura, S., Yamazaki, T., Yoshida, M., Takahashi, K., Nagai, H., Sugimoto, C., Yamaguchi, H., 2014. Amoebal Endosymbiont Neochlamydia Genome Sequence Illuminates the Bacterial Role in the Defense of the Host Amoebae against Legionella pneumophila. PLoS ONE 9, e95166. https://doi.org/10.1371/journal.pone.0095166

Jeon, K.W., 1992. Prokaryotic Symbionts of Amoebae and Flagellates, in: Balows, A., Trüper, H.G., Dworkin, M., Harder, W., Schleifer, K.-H. (Eds.), The Prokaryotes. Springer New York, New York, NY, pp. 3855–3864. https://doi.org/10.1007/978-1-4757-2191-1_51

Jeon, K.W., Jeon, M.S., 1976. Endosymbiosis in amoebae: Recently established endosymbionts have become required cytoplasmic components. J. Cell. Physiol. 89, 337–344. https://doi.org/10.1002/jcp.1040890216

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., Drummond, A., 2012. Geneious Basic: An integrated and extendable desktop software platform for the

organization and analysis of sequence data. Bioinformatics 28, 1647–1649.
https://doi.org/10.1093/bioinformatics/bts199

Kessin, R.H., 2001. Dictyostelium: evolution, cell biology, and the development of
multicellularity, Developmental and cell biology series. Cambridge University Press,
Cambridge, UK ; New York.

Khojandi, N., Haselkorn, T.S., Eschbach, M.N., Naser, R.A., DiSalvo, S., 2019. Intracellular
Burkholderia Symbionts induce extracellular secondary infections; driving diverse host
outcomes that vary by genotype and environment. ISME J.
https://doi.org/10.1038/s41396-019-0419-7

König, L., Wentrup, C., Schulz, F., Wascher, F., Escola, S., Swanson, M.S., Buchrieser, C., Horn,
M., 2019. Symbiont-Mediated Defense against Legionella pneumophila in Amoebae.
mBio 10. https://doi.org/10.1128/mBio.00333-19

Lagkouvardos, I., Shen, J., Horn, M., 2014. Improved axenization method reveals complexity of
symbiotic associations between bacteria and acanthamoebae: An improved method for
the axenization of amoebae. Environ. Microbiol. Rep. 6, 383–388.
https://doi.org/10.1111/1758-2229.12162

Maita, C., Matsushita, M., Miyoshi, M., Okubo, T., Nakamura, S., Matsuo, J., Takemura, M.,
Miyake, M., Nagai, H., Yamaguchi, H., 2018. Amoebal endosymbiont Neochlamydia
protects host amoebae against Legionella pneumophila infection by preventing
Legionella entry. Microbes Infect. 20, 236–244.
https://doi.org/10.1016/j.micinf.2017.12.012

Matsuo, J., Kawaguchi, K., Nakamura, S., Hayashi, Y., Yoshida, M., Takahashi, K., Mizutani, Y.,
Yao, T., Yamaguchi, H., 2009. Survival and transfer ability of phylogenetically diverse
bacterial endosymbionts in environmental Acanthamoeba isolates: Survival and transfer
of bacterial endosymbionts in amoebae. Environ. Microbiol. Rep. 2, 524–533.
https://doi.org/10.1111/j.1758-2229.2009.00094.x

Matz, C., Kjelleberg, S., 2005. Off the hook – how bacteria survive protozoan grazing. Trends
Microbiol. 13, 302–307. https://doi.org/10.1016/j.tim.2005.05.009

Nylund, A., Pistone, D., Trösse, C., Blindheim, S., Andersen, L., Plarre, H., 2018. Genotyping of
Candidatus Syngnamydia salmonis (chlamydiales; Simkaniaceae) co-cultured in
Paramoeba perurans (amoebozoa; Paramoebidae). Arch. Microbiol. 200, 859–867.
https://doi.org/10.1007/s00203-018-1488-0

Okude, M., Matsuo, J., Nakamura, S., Kawaguchi, K., Hayashi, Y., Sakai, H., Yoshida, M.,
Takahashi, K., Yamaguchi, H., 2012. Environmental Chlamydiae Alter the Growth Speed
and Motility of Host Acanthamoebae. Microbes Environ. 27, 423–429.
https://doi.org/10.1264/jsme2.ME11353

Ossewaarde, J.M., Meijer, A., 1999. Molecular evidence for the existence of additional
members of the order Chlamydiales. Microbiology 145, 411–417.
https://doi.org/10.1099/13500872-145-2-411

Paquet, V.E., Charette, S.J., 2016. Amoeba-resisting bacteria found in multilamellar bodies
secreted by *Dictyostelium discoideum:* social amoebae can also package bacteria. FEMS
Microbiol. Ecol. 92, fiw025. https://doi.org/10.1093/femsec/fiw025

Penz, T., Horn, M., Schmitz-Esser, S., 2010. The genome of the amoeba symbiont "Candidatus
Amoebophilus asiaticus " encodes an afp-like prophage possibly used for protein
secretion. Virulence 1, 541–545. https://doi.org/10.4161/viru.1.6.13800
Pillonel, T., Bertelli, C., Aeby, S., de Barsy, M., Jacquier, N., Kebbi-Beghdadi, C., Mueller, L.,
Vouga, M., Greub, G., 2019. Sequencing the Obligate Intracellular Rhabdochlamydia
helvetica within Its Tick Host Ixodes ricinus to Investigate Their Symbiotic Relationship.
Genome Biol. Evol. 11, 1334–1344. https://doi.org/10.1093/gbe/evz072
Pillonel, T., Bertelli, C., Greub, G., 2018. Environmental Metagenomic Assemblies Reveal Seven
New Highly Divergent Chlamydial Lineages and Hallmarks of a Conserved Intracellular
Lifestyle. Front. Microbiol. 9, 79. https://doi.org/10.3389/fmicb.2018.00079
Proca-Ciobanu, M., Lupascu, Gh., Petrovici, Al., Ionescu, M.D., 1975. Electron microscopic study
of a pathogenic Acanthamoeba castellani strain: The presence of bacterial
endosymbionts. Int. J. Parasitol. 5, 49–56. https://doi.org/10.1016/0020-
7519(75)90097-1
Rock, D.I., Smith, A.H., Joffe, J., Albertus, A., Wong, N., O'Connor, M., Oliver, K.M., Russell, J.A.,
2018. Context-dependent vertical transmission shapes strong endosymbiont community
structure in the pea aphid, *Acyrthosiphon pisum*. Mol. Ecol. 27, 2039–2056.
https://doi.org/10.1111/mec.14449
Rodríguez-Zaragoza, S., 1994. Ecology of Free-Living Amoebae. Crit. Rev. Microbiol. 20, 225–
241. https://doi.org/10.3109/10408419409114556
Ronn, R., McCaig, A.E., Griffiths, B.S., Prosser, J.I., 2002. Impact of Protozoan Grazing on
Bacterial Community Structure in Soil Microcosms. Appl. Environ. Microbiol. 68, 6094–
6105. https://doi.org/10.1128/AEM.68.12.6094-6105.2002
Samba-Louaka, A., Delafont, V., Rodier, M.-H., Cateau, E., Héchard, Y., 2019. Free-living
amoebae and squatters in the wild: ecological and molecular features. FEMS Microbiol.
Rev. 43, 415–434. https://doi.org/10.1093/femsre/fuz011
Santarella-Mellwig, R., Pruggnaller, S., Roos, N., Mattaj, I.W., Devos, D.P., 2013. Three-
Dimensional Reconstruction of Bacteria with a Complex Endomembrane System. PLoS
Biol. 11, e1001565. https://doi.org/10.1371/journal.pbio.1001565
Santos-Garcia, D., Rollat-Farnier, P.-A., Beitia, F., Zchori-Fein, E., Vavre, F., Mouton, L., Moya, A.,
Latorre, A., Silva, F.J., 2014. The Genome of Cardinium cBtQ1 Provides Insights into
Genome Reduction, Symbiont Motility, and Its Settlement in Bemisia tabaci. Genome
Biol. Evol. 6, 1013–1030. https://doi.org/10.1093/gbe/evu077
Schmitz-Esser, S., Tischler, P., Arnold, R., Montanaro, J., Wagner, M., Rattei, T., Horn, M., 2010.
The Genome of the Amoeba Symbiont "Candidatus Amoebophilus asiaticus" Reveals
Common Mechanisms for Host Cell Interaction among Amoeba-Associated Bacteria. J.
Bacteriol. 192, 1045–1057. https://doi.org/10.1128/JB.01379-09
Schmitz-Esser, S., Toenshoff, E.R., Haider, S., Heinz, E., Hoenninger, V.M., Wagner, M., Horn, M.,
2008. Diversity of Bacterial Endosymbionts of Environmental Acanthamoeba Isolates.
Appl. Environ. Microbiol. 74, 5822–5831. https://doi.org/10.1128/AEM.01093-08
Schulz, F., Lagkouvardos, I., Wascher, F., Aistleitner, K., Kostanjšek, R., Horn, M., 2014. Life in an
unusual intracellular niche: a bacterial symbiont infecting the nucleus of amoebae. ISME
J. 8, 1634–1644. https://doi.org/10.1038/ismej.2014.5

Sheikh, S., Thulin, M., Cavender, J.C., Escalante, R., Kawakami, S., Lado, C., Landolt, J.C.,
Nanjundiah, V., Queller, D.C., Strassmann, J.E., Spiegel, F.W., Stephenson, S.L., Vadell,
E.M., Baldauf, S.L., 2018. A New Classification of the Dictyostelids. Protist 169, 1–28.
https://doi.org/10.1016/j.protis.2017.11.001
Shu, L., Brock, D.A., Geist, K.S., Miller, J.W., Queller, D.C., Strassmann, J.E., DiSalvo, S., 2018.
Symbiont location, host fitness, and possible coadaptation in a symbiosis between social
amoebae and bacteria. eLife 7. https://doi.org/10.7554/eLife.42660
Steinert, M., 2011. Pathogen–host interactions in Dictyostelium, Legionella, Mycobacterium
and other pathogens. Semin. Cell Dev. Biol. 22, 70–76.
https://doi.org/10.1016/j.semcdb.2010.11.003
Steinert, M., Heuner, K., 2005. Dictyostelium as host model for pathogenesis. Cell. Microbiol. 7,
307–314. https://doi.org/10.1111/j.1462-5822.2005.00493.x
Sun, S., Noorian, P., McDougald, D., 2018. Dual Role of Mechanisms Involved in Resistance to
Predation by Protozoa and Virulence to Humans. Front. Microbiol. 9, 1017.
https://doi.org/10.3389/fmicb.2018.01017
Taylor, M., Mediannikov, O., Raoult, D., Greub, G., 2012. Endosymbiotic bacteria associated
with nematodes, ticks and amoebae. FEMS Immunol. Med. Microbiol. 64, 21–31.
https://doi.org/10.1111/j.1574-695X.2011.00916.x
Taylor-Brown, A., Polkinghorne, A., 2017. New and emerging chlamydial infections of creatures
great and small. New Microbes New Infect. 18, 28–33.
https://doi.org/10.1016/j.nmni.2017.04.004
Taylor-Brown, A., Vaughan, L., Greub, G., Timms, P., Polkinghorne, A., 2015. Twenty years of
research into Chlamydia-like organisms: a revolution in our understanding of the biology
and pathogenicity of members of the phylum Chlamydiae. Pathog. Dis. 73, 1–15.
https://doi.org/10.1093/femspd/ftu009
Taylor-Mulneix, D.L., Hamidou Soumana, I., Linz, B., Harvill, E.T., 2017. Evolution of Bordetellae
from Environmental Microbes to Human Respiratory Pathogens: Amoebae as a Missing
Link. Front. Cell. Infect. Microbiol. 7. https://doi.org/10.3389/fcimb.2017.00510
Tosetti, N., Croxatto, A., Greub, G., 2014. Amoebae as a tool to isolate new bacterial species, to
discover new virulence factors and to study the host–pathogen interactions. Microb.
Pathog. 77, 125–130. https://doi.org/10.1016/j.micpath.2014.07.009
Zinicola, M., Lima, F., Lima, S., Machado, V., Gomez, M., Döpfer, D., Guard, C., Bicalho, R., 2015.
Altered Microbiomes in Bovine Digital Dermatitis Lesions, and the Gut as a Pathogen
Reservoir. PLOS ONE 10, e0120504. https://doi.org/10.1371/journal.pone.0120504
963
964

 Table and Figure Legends:

Figure 1. Unculturable symbiont prevalence in natural *D. discoideum* populations.

(a) Pie charts indicate Chlamydiae, *Amoebophilus*, and Chlamydiae-*Amoebophilus* co-

infection patterns in respective populations of *D. discoideum* isolates from across the

Eastern United States. *Procabacter*, due to its low prevalence, is not included. Numbers

next to pie charts indicate the number of isolates for each population. (b) Of all symbiont

972   positive amoebae isolates, several are co-infected with more than one symbiont

973   species, represented by the Venn diagram indicating the number of isolates infected

974   with *Paraburkholderia, Amoebophilus,* and Chlamydiae.

995   a.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011 b.

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

c.

0.050

Figure 2. Unculturable endosymbionts of *Dictyostelium* are related to

unculturable endosymbionts of other amoebae. 16S rRNA phylogenies of

Procobacter (a) *Amoebophilus* (b) and Chlamydiae (c & d) endosymbionts constructed

using the Maximum Likelihood method in Mega7, with 1000 bootstrap replicates. Family

names, where applicable are delineated in bold italics.

QS864 Uninfected

1μm

QS851 *Amoebophilus* Infected

500nm

QS886 *Chlamydiae* Infected

QS1040 *Procabacter* Infected

1033

1034 **Figure 3. Unculturable symbionts have distinct morphologies in infected spores.**

1035 Transmission electron micrographs of spores from the indicated representative

1036 uninfected and infected *D. discoideum* isolates. Images on the right show close-ups of

1037 intracellular bacteria.

Figure 4. Elimination of *Amoebophilus* and Chlamydiae from natural hosts with

antibiotics does not impact host reproductive fitness. Four uninfected,

Amoebophilus-infected, and Chlamydiae-infected representative host isolates were

treated with rifampicin to eliminate symbionts (a) and reproductive fitness (b) was

assessed. Symbiont presence is indicated by amplification with *Amoebophilus*-specific

or Chlamydiae-specific PCR and DNA gel electrophoresis (with Eukaryote specific gene

amplification serving as an internal control) (a). Reproductive fitness was assessed by

quantifying total spore productivity of *D. discoideum* cultures after completion of one

social cycle (b). Individual *D. discoideum* isolates are represented by point color.

Figure 5. *Amoebophilus* **and Chlamydiae infections do not impact the**

reproductive fitness of hosts co-infected with *Paraburkholderia* **symbionts.**

Reproductive fitness was quantified for four untreated and antibiotic-cured

Amoebophilus-infected (a) and Chlamydiae-infected isolates (b) following social cycle

completion after exposure to the indicated *Paraburkholderia* symbiont strains. Individual

D. discoideum isolates are represented by point color.

1055 **Table S1. Table of Screened *D. discoideum* Natural Isolates.** Table lists each *D.*

1056 *discoideum* isolate by identification number and location of collection. PCR amplification

1057 of *Dictyostelium*, *Paraburkholderia*, *Amoebophilus*, or Chlamydiae specific DNA is

1058 indicated with a 1 (for positive amplification) or 0 (for negative amplification). Samples

1059 used for fitness assays are highlighted in yellow.

1060

1061 **Table S2. SILVA Database Classification of Symbiont Haplotypes.** Percent

1062 sequence identity to closest hit in the database for the five full length 16S rRNA

1063 sequence haplotypes and their least common ancestor taxonomic classification.

1064

1065 **Table S3. IMNGS database >99% matches for Chlamydiae Endosymbiont of *D.***

1066 ***discoideum* Haplotype 1.** Includes metagenome sample type and collection location.

1067

1068 **Table S4. PCR Primers Used for this Study.** PCR primer names, sequences, and

1069 associated references.

1070