Consideration of macrophage phenotypes under biomaterials mediated host defense process

Shengan Rung[a 1], Chenyu Chu[a 1], Chen Hu[a], Li Liu[b], Yili Qu[a*], Yi Man[a*]

[a] *State Key Laboratory of Oral Diseases & National Clinical Research Center for Oral Diseases & Department of Oral Implantology, West China Hospital of Stomatology, Sichuan University, Chengdu 610041, China.*

[b] *State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, Collaborative Innovation Center for Biotherapy, Chengdu, Sichuan 610041, China*

1 These authors contributed equally to this work.

Corresponding author:

Yi Man, DDS, PhD

Chair and Professor

State Key Laboratory of Oral Diseases & National Clinical Research Center for

Oral Diseases & Department of Oral Implantology, West China Hospital of

Stomatology, Sichuan University

Tel: 0086-28-85503571

Email: Manyi780203@126.com

Yili Qu, DDS, PhD

Associate Professor

State Key Laboratory of Oral Diseases, West China Hospital of Stomatology,

Sichuan

University, 14#, 3rd section, Renmin South Road

Chengdu 610041, China

Tel: 0086-28-85503571

Email: qqyili@126.com

Key words: macrophages, CAMP, candida albicans, biomaterials, cutaneous

Abstract

Culturing macrophage in vitro is an important means to understand its reaction towards co-culture substances. However, the source of macrophages can be either purchased from specific cell line or extracted from bone marrow as differentiated macrophages. In order to assure the precision, when adopting extracted primary cell, screening in advance will be prominent before any processes to avoid results reserving that of non-macrophages. Here, we perform single-cell sequencing on open wound of skin in mice and focusing on the secreted proteins and surface markers related to traditional macrophage phenotypes (M1/M2) to ensure the importance of screening. The expression of CAMP when macrophages fight against candida albicans is another target to see its relationship with current classification. And results showed that identifying its phenotype without screening macrophages will far from the exact situation, and the expression of CAMP cannot be carried out by the traditional M1 and M2 macrophage. Thus, determining phenotype of macrophages based on function would be a promising way.

Introduction

Macrophage is one of the most important defenders and regulators of the immune response, which will be activated by the immunological recognition of pathogens and

foreign bodies. The timely shifting from M1 towards the M2 phenotype of

Macrophage is usually concerned with tissue repair. In contrast, Gao et al. mentioned

that an opposite strategy was needed for tackling Candida albicans with the demand

effectively to improve the survival rate after being infected [1]. To evaluate the

nano-trinity treatment effect on phenotypes of macrophage and its expression, flow,

western blot (WB), and ELISA kits were adopted. However, the controversies in this

study not only appeared in the selected markers and the grouping process which seem

lacking comprehensive consideration, but also emerged in the glancing description to

the interaction between specific phenotypes of macrophages and Candida. In this

study, we discussed the related topic with single cell RNA sequencing data,

investigated the contradictory results and analysis the underlying way of classification

of phenotype during biomaterials mediated host defense against Candida albicans.

Method and materials

The original data is obtained from previous database[2] and processed through Seurat3

(https://satijalab.org/seurat/) and Loupe Browser v4.0 (10xgenomics, USA) for

subsequent analysis.

Results

Macrophage-specific markers should be detected before determining its

phenotype

In order to confirm the specificity of the macrophage phenotypic markers (Fig. 1A, B)

and proteins (Fig. 1C-G) secreted by specific phenotypes, we analyzed the expression

of each cluster in open wound of skin. And it was found that none of the detected

markers are completely specific; most of the targeting gene have close relationship

with neutrophils (Fig. 1H) or fibroblasts (Fig. 1I). That is, through those markers to

determine macrophage phenotype from an obvious potentially complex environment

without screening the specific marker of macrophages like F4/80 in advance, the

result is prone to false positives from other non- macrophage cells.

The expression of Camp is independent of traditional M1/M2 phenotypic

markers

Comparing the gene expression of macrophages and neutrophils, Arg1 could be

expressed by both cells in open wound of skin; this phenomenon can also be seen in

Nos2 (Fig. 2A, B and D). In addition, we found that the expression of Camp has

nothing to do with traditional macrophage phenotypic markers (M1-Nos2/M2-Arg1),

for neither $Arg1^+Camp^+$ nor $Nos2^+Camp^+$ cells exist (Fig. 2E and 2F). Moreover,

Camp could only be seen in neutrophil cluster with relative low expression (Fig. 2C).

Discussion

First, the phenotype of both J774A.1 macrophage cell line and bone marrow-derived

macrophages (BMDMs) were detected by simultaneously incubating with F4/80 and

CD206 (M2 phenotype) or CD80 (M1 phenotype), respectively. This indifferent

treatment would accompany inevitable error to BMDMs, which would cause the data

of flow cytometry included CD80$^+$ and CD206$^+$ of non-macrophage cells and

increased false-positive results. For example, in the murine open wound of skin,

CD80$^+$ cells are not restrained to cluster of monomolecular macrophages (Fig. 1A),

while CD206, which has always been recognized as a representative marker of M2

macrophage, is not specifically expressed by macrophages (Fig. 1B). Moreover,

through the heat map of our single-cell sequencing result, neutrophils also express

Mrc1 that encodes CD206 (Fig. 1H and 1I). That is, BMDMs, unlike J774A.1

macrophage, is difficult to avoid the presence of multiple types of cells apart from

macrophages. As extracted *in vivo*, BMDMs is essentially different from the cell line.

And merely CD80 or CD206 could not represented for a certain phenotype; more

markers should be adopted. Also, the F4 / 80 staining results showed that the positive

population had no obvious clustering that was difficult to demonstrate the existence of

an F4 / 80^{high} population in BMDMs. To resolve this problem, conducting flow

cytometry for phenotype indication should begin with extracting CD11b$^+$F4 / 80$^+$

population from the entire BMDMs to determine macrophage, then analyzing those

phenotypic M1/M2 specific markers. Besides, according to the previous study,

macrophages with both M1 and M2 markers must also be considered, therefore it

would be better to stained CD80 and CD206 at the same time based on F4 / 80^{high}

population (CD80$^+$CD206$^-$ represented for M1 phenotype; CD80$^-$CD206$^+$ represented

for M2 phenotype). As for, the secretion of iNOS, Arg1, TNF-α, IL-6, and IL-10 to

determine macrophage polarization also neglected whether they are specific for

macrophage (Fig. 1C-G). Especially iNOS (encoded by Nos2), which was different

from Arg1, our results showed that it could be secreted by neutrophil and fibroblast

(Fig. 1H and 1I). Therefore, without quality control, WB and ELISA results of

BMDMs must include the expression levels of other bone marrow-derived cells. Such

protein quantitative results cannot correspond to flow results. In particular, separating

macrophages from BMDMs before processing any assays, could better reflect the real

situation. Last but not least, Candida albicans could reduce the production of NO, not

by inhibiting the expression of NOS2 mRNA, but lowering the activity of NOS2

enzyme and decreasing the expression of NOS2 protein [3]. Whereas other researchers

suggested that it is due to the promotion of Arg1 through chitin from candida albicans

in the competition for the L-arginine substrate. With the help of Arg1 inhibitor, the

synthesis of NO could be restored while the survival rate of the host being improved [4].

It may seem that weakening macrophages polarization towards M2 phenotype when

tackling such pathogens would be a promising solution, but the host response under

the attack candida albicans is not that simple. Indeed, there are corresponding studies

aiming at both macrophages of murine and human that have an M1-to-M2 switch

after candida albicans infection [5, 6], but whether this polarization trend is candida's

strategy to improve its survival rate or the result of host self-protection is still

unknown. In addition, as recommended by Gao et al., increasing M1 macrophages can

promote the expression of NO, ROS and enhance phagocytosis to fight candida

albicans [1], however, candida albicans may end up resulting in the death of

macrophages by transferring its metabolic way on glucose that macrophage could

rival it [7]. Thus, to confirm that macrophages can reduce the damage of Candida by

enhancing M1 macrophage polarization, longer-term observation is needed. Cramp,

an antimicrobial peptide, is another effective weapon of the host against candida

albicans, which could be produced by Neutrophils, macrophages and epithelial cells [8].

According to our single-cell sequencing results, Camp (encoding Cramp) was

expressed in the open wound of skin that did not exclude candida albicans infection

(Fig. 2A-D), but neither Arg1 or Nos2 are correlated to this gene (Fig. 2E and 2F),

showing that those conventional classification markers of macrophage could not serve

as an indicator for the production of Cramp in murine. And compared to macrophages,

neutrophils even expressed more Camp. Other than antimicrobial peptides, Cabezón,

Virginia, et al. had also clarified multiple proteins acting between macrophages and

Candida, which is related to the induction of candida albicans' apoptosis, confirming

macrophages contribution [9]. However, whether the M1 and M2 macrophage phenotypes are the key to regulating Candida infection, and how the specific phenotype of macrophages prolongs the host survival time needs further study. Base on the information mentioned above, by means of functionally relevant markers may be a more reasonable option for understanding the role of macrophages in fungal infections. The removal of pathogens by macrophages is indispensable, which functions closely related to the phenotype. However, in the grouping process, it should be considered whether the selected object composed of macrophages only. If there were other types of cells, they should be excluded before starting a phenotypic assay. The gating for the baseline and selection of markers should according to the composition of the tested samples. The phenotypes and corresponding antibacterial effects also interested us a lot. It would be our pleasure to have further communication on the perspective mentioned above and the author's considerations for experimental design.

Acknowledgement

This work was supported by Research and Develop Program, West China Hospital of Stomatology Sichuan University (No.LCYJ2019-19); The Fundamental Research Funds for The Central Universities (No. 2082604401239)

Statement of conflict of interest

There are no conflicts of interest related to this manuscript.

Reference

1.   Gao, Q.; Zhang, J.; Chen, C.; Chen, M.; Sun, P.; Du, W.; Zhang, S.; Liu, Y.; Zhang, R.; Bai, M*., et al.* In Situ Mannosylated Nanotrinity-Mediated Macrophage Remodeling Combats Candida Albicans Infection. *ACS nano* **2020,** 14(4), 3980-3990.

2.   Hu, C.; Chu, C.; Liu, L.; Qu, Y.; Man, Y. Dissecting the Microenvironment around Biosynthetic Scaffolds in Murine Skin Wound Healing. *bioRxiv* **2020**.

3.   Schröppel, K.; Kryk, M.; Herrmann, M.; Leberer, E.; Röllinghoff, M.; Bogdan, C. Suppression of Type 2 No-Synthase Activity in Macrophages by Candida Albicans. *International journal of medical microbiology : IJMM* **2001,** 290(8), 659-68.

4.   Wagener, J.; Maccallum, D. M.; Brown, G. D.; Gow, N. A. Candida Albicans Chitin Increases Arginase-1 Activity in Human Macrophages, with an Impact on Macrophage Antimicrobial Functions. *mBio* **2017,** 8(1).

5.   Alvarez-Rueda, N.; Rouges, C.; Touahri, A.; Misme-Aucouturier, B.; Albassier, M.; Pape, P. L. In Vitro Immune Responses of Human Pbmcs against Candida Albicans Reveals Fungal and Leucocyte Phenotypes Associated with Fungal Persistence. *Scientific reports* **2020,** 10(1), 6211.

6.   Reales-Calderón, J. A.; Aguilera-Montilla, N.; Corbí Á, L.; Molero, G.; Gil, C. Proteomic Characterization of Human Proinflammatory M1 and Anti-Inflammatory M2 Macrophages and Their Response to Candida Albicans. *Proteomics* **2014,** 14(12), 1503-18.

7.   Tucey, T. M.; Verma, J.; Harrison, P. F.; Snelgrove, S. L.; Lo, T. L.; Scherer, A. K.; Barugahare, A. A.; Powell, D. R.; Wheeler, R. T.; Hickey, M. J*., et al.* Glucose Homeostasis Is Important for Immune Cell Viability During Candida Challenge and Host Survival of Systemic Fungal Infection. *Cell Metab* **2018,** 27(5), 988-1006.e7.

8.   López-García, B.; Lee, P. H.; Yamasaki, K.; Gallo, R. L. Anti-Fungal Activity of Cathelicidins and Their Potential Role in Candida Albicans Skin Infection. *The Journal of investigative dermatology* **2005,** 125(1), 108-15.

9.   Cabezón, V.; Vialás, V.; Gil-Bona, A.; Reales-Calderón, J. A.; Martínez-Gomariz, M.; Gutiérrez-Blázquez, D.; Monteoliva, L.; Molero, G.; Ramsdale, M.; Gil, C. Apoptosis of Candida Albicans During the Interaction with Murine Macrophages: Proteomics and Cell-Death Marker Monitoring. *Journal of proteome research* **2016,** 15(5), 1418-34.

Figure 1. Single-cell sequencing result of skin open wound. Expression of gene (A)

CD80, (B) Mrc1, (C) Nos2, (D) Arg1, (E) Tnf, (F) Il6, (G) Il10 and heat map of

monomolecular macrophage versus (H) neutrophil and (I) fibroblast cluster

Figure 2. (A) Single-cell sequencing result of monomolecular macrophage (red) and neutrophil (blue). The expression of (B) Arg1, (C) Camp, (D) Nos2, (E) Arg1+Camp+ and (F) Nos2+Camp+.