Social relationships among captive male pygmy slow lorises (*Nycticebus pygmaeus*): Is

forming male same-sex pairs a feasible management strategy?

Running head: social relationships among captive male pygmy slow lorises

Yumi Yamanashi[1,2], Kei Nemoto[3], Josue Alejandro[4]

1. Center for Research and Education of Wildlife, Kyoto City Zoo

2. Wildlife Research Center, Kyoto University

3. Japan Monkey Center

4. Primate Research Institute, Kyoto University

Correspondent author

Yumi Yamanashi

Email: yumi.yamanashi.kycz@gmail.com

Physical address: Okazaki kouen, Okazaki hosshoji-cho, Sakyo-ku, Kyoto City, Kyoto,

Japan

Tel: +81-75-771-0210

Abstract

Little is known about the social behavior of pygmy slow lorises, in particular,

the social relationships of same-sex individuals have rarely been investigated. The Slow

Loris Conservation Center was built at the Japan Monkey Center to enhance the welfare

of confiscated slow lorises, promote their conservation, improve public education and

perform scientific research on the species. In the course of improving housing conditions,

several same-sex pairs of pygmy slow lorises were formed. We monitored their behaviors

and fecal glucocorticoid metabolite (FGM) levels to understand whether male same-sex

pairings could be a feasible management strategy. The subjects were 10 male and 6 female

lorises for comparison, all of whom were over five years old. We successfully formed five

pairs of male lorises after eight formation attempts. Male pairs initially showed some

aggressive behaviors; however, the rate decreased approximately 10 days after

introduction. All of the male pairs eventually exhibited extensive affiliative social

behaviors, including allogrooming and social play, during the dark (active) phase, and

sleep site sharing during the light (inactive) phase. The rate of sleep site sharing during

the light phase was higher than expected, suggesting that the pairs preferred to stay near

each other. There was no evidence of increased stress after a long period of male–male

social housing. Female same-sex pairs and male-female pairs demonstrated a high level

of affiliative behaviors right after introduction. These results highlight the flexibility and

high sociability of this species and indicate that such same-sex pairings are a feasible

option for their social management.

Keywords

Pygmy slow loris, sociality, animal welfare, conservation, social management

Introduction

Slow lorises (*Nycticebus spp.*) are in danger of extinction. The IUCN Red List currently classifies them into nine species, among which the pygmy slow loris (*Nycticebus pygmaeus*) is categorised as Endangered (IUCN, 2020). Although their international commercial trade has been prohibited as a general rule since 2007 by their inclusion in CITES Appendix I, one of the causes of their loss remains their frequent illegal trade (Musing, Suzuki, & Nekaris, 2015; Nekaris, 2007). Thus, an important conservation goal for this species is to reduce their use in the illegal pet trade through, among other strategies, public education (Norconk et al., 2020). Slow lorises are confiscated in both range and non-range countries, including Japan (Fuller, Eggen, Wirdateti, & Nekaris, 2018; Kitade & Naruse, 2020; Musing et al., 2015). Once confiscated, lorises are typically transferred to sanctuaries, rehabilitation centers or zoos (Fuller et al., 2018). Sanctuaries and rescue centers can often become strained past their capacity due to the unscheduled arrival of the confiscated animals (Moore, Wihermanto, & Nekaris, 2014), which can result in suboptimal captive conditions, mainly due to the lack of available space. Therefore, it is important to develop a practical strategy to house these individuals to ensure their well-being.

Little is known about the social behavior of wild pygmy slow lorises due to the inherent difficulty of nocturnal observation. The species was historically considered solitary, but studies have revealed that they are not asocial. Some slow loris species have been found to have spatial overlaps in home ranges between adult males and females (Wiens & Zitzmann, 2003), and friendly interactions between individuals have been reported (Wiens, 2002). Javan slow loris spend as much as 65% of their time in spatial proximity, including body contact for an average of 18% of their time (Nekaris, 2014).

Studies have also demonstrated that male slow lorises are often found with scars, which

has led some researchers to conclude that slow loris venom is used for intraspecific

competition (Nekaris, Moore, Rode, & Fry, 2013; Nekaris et al., 2020). A recent study

investigating the ranging patterns of pygmy slow loris in Cambodia found that the home

ranges of some individuals overlapped with others to a small degree. Adult pygmy slow

lorises display primarily solitary sleeping patterns, except for adult females with infants,

who always sleep with their young (Starr & Nekaris, 2020). These solitary sleeping

patterns are consistent with greater slow loris behavior (Wiens & Zitzmann, 2003), but

different from Javan slow lorises, who often share sleep sites with conspecifics (Nekaris,

2014). Activity budgets differ across species and studies, with the proportions of social

behaviors ranging from 0%–18% (Nekaris, 2014; Nekaris & Bearder, 2007; Rode-

Margono, Nijman, Wirdateti., & Nekaris, 2014). Overall, the current available evidence

suggests that slow lorises are more social than previously thought, and form some groups

(Wiens & Zitzmann, 2003), including dispersed family groups (Javan slow loris and

greater slow loris), and polygynous groups (pygmy slow loris) (Poindexter & Nekaris,

2020).

Slow lorises display social behaviors in captivity, not only between mother and

infant (Fitch-Snyder & Ehrlich, 2003), but also between adults (Ehrlich & Alan, 1977). A

previous study reported that two polygynous groups of greater slow lorises participated

in various social behaviors, including allogrooming and social play (Ehrlich & Alan,

1977). Researchers believe that such stimulation from conspecifics can be a form of social

enrichment (Fitch-Snyder, Schulze, & Larson, 2001). Despite these data, one survey

focusing on the husbandry practices of North American zoos for lorisid primates revealed

that approximately 55% of slow lorises are housed alone (Fuller, Kuhar, Dennis, & Lukas,

2013). Conspecific aggression during animal introductions was listed as one reason for

the single housing. Although aggression can be a source of injury and stress in primates

(Honess & Marin, 2006; Tennenhouse, Putman, Boisseau, & Brown, 2017; Yamanashi et

al., 2016), and can compromise welfare if it continues over a long period or if the level

of aggression is severe, moderate aggression is a natural behavior (Bloomsmith, 2001)

that often occurs during the introduction phases of gregarious primates, and often

decreases once their social relationships are established (Kutsukake et al., 2018). A well-

established, cohesive social environment can offer ever-changing stimulation to animals

in captivity. Successful social management can help to maximize the positive benefits of

social groups and minimize negative effects, such as stress and aggression, but discussion

on the practical social management of slow lorises is still limited.

Ideally, social structures in captivity should mimic the social structures seen in

the wild for a particular species (Price & Stoinski, 2007); however, social structures in

wild primates can vary in group size and composition depending on resource availability,

predator presence, and other factors. In captivity, such environmental constraints differ

(e.g. less resource competition and absent predators), which could allow for increased

social group flexibility in captivity (Price & Stoinski, 2007). For example, same-sex

groups are sometimes formed to control breeding or to manage surplus animals (Fàbregas

& Guillén-Salazar, 2007; Sha, Alagappasamy, Chandran, Cho, & Guha, 2013; Stoinski,

Kuhar, Lukas, & Maple, 2004; Stoinski, Lukas, Kuhar, & Maple, 2004; Yamanashi,

Nogami, Teramoto, Morimura, & Hirata, 2018), even in species where such social

compositions are non-existent in the wild (e.g. chimpanzees). Similarly, wild orangutans

are semi-solitary species (van Schaik, 1999), but are sometimes kept in social groups in

zoos (Amrein, Heistermann, & Weingrill, 2014; Weingrill, Willems, Zimmermann,

Steinmetz, & Heistermann, 2011). Such flexible groups enable us to maximize space, and

avoid suboptimal housing. However, the degree of social flexibility is species-dependent,

and needs to be carefully investigated for each species. In slow lorises, same-sex pairs

have also been formed in some captive settings (Fuller et al., 2013; Moore, Cabana, &

Nekaris, 2015). In one study conducted at a rehabilitation center in Indonesia, confiscated

slow lorises housed with conspecifics, particularly in same-sex pairs, displayed fewer

stereotypic behaviors (Moore et al., 2015). However, the social relationships between

same-sex slow lorises have rarely been investigated either in the wild or in captivity, and

it is not well understood whether same-sex housing is a viable social management option

for *Nycticebus spp.*

The purpose of this study was to investigate the social behaviors of male pairs

of pygmy slow lorises, in order to consider whether same-sex pairing could be a feasible

management strategy for this species. If slow lorises are motivated to interact with

conspecifics, and benefit from being together, then we would consider that same-sex

pairings are a feasible option. We particularly focused on male–male pairings in this study,

because males often exhibit competitive tendencies (Van Schaik, 1994), which often

makes their social management more complicated than female groups (Sha et al., 2013;

Stoinski, Lukas, et al., 2004). In addition, we also measured the fecal glucocorticoid

metabolite (FGM) levels as a measure of physiological stress. Glucocorticoids (GCs), the

steroid hormones secreted by the adrenal glands of vertebrates, are frequently used as an

indicator of physiological and psychological stress, because they are often increased when

organisms face stressors (Selye, 1941; Squires, 2010). GCs generally increase available

energy, and their acute increase is generally considered adaptive in stressful situations.

However, prolonged exposure to GCs can result in a number of maladaptive

consequences, such as neuronal cell death, insulin resistance, muscle and bone atrophy,

poor wound healing, hypertension, and even immune system collapse to the point of death

(Reeder & Kramer, 2005). Therefore, prolonged GC activation can be considered

maladaptive from the view of animal welfare. For the purposes of this study, we explored

five variables. First, we investigated the types of social relationships observed between

adult males. Second, we determined how often sleep sites were shared, in order to

understand whether the lorises preferred to stay together during their light phases (i.e.

when the individuals are typically resting or sleeping). To exclude the potential influences

of environmental factors, such as locational preference or temperature, we also recorded

temperatures, and calculated the chance rate of sleep site sharing. Third, we observed the

process that occurred during the formation of the social relationships, and its effects on

the individuals' physiological stress levels. Fourth, we compared the social relationship

formation process across pairs of different sex combinations. Fifth, we considered the

stability of each social relationship by comparing social behaviors observed during two

different study periods.

If the pygmy slow lorises were motivated to interact with conspecifics, then we

hypothesized that male pygmy slow lorises would display affiliative social behaviors, and

prefer to share sleep sites during their light phases. In this case, physiological stress levels

may decrease after social relationships have been formed, or they may remain the same.

If slow lorises were not motivated to interact with conspecifics, then we hypothesized that

they would choose to keep distance between each other. In the worst case, they would

continue fighting without showing any sign of affiliation. Physiological stress levels in

this case would then likely increase over time as a cost of the social environment.

Methods

Subjects

The subjects of this research were 16 pygmy slow lorises living at the Japan Monkey Center (JMC), in Inuyama, Aichi, Japan (10 males and 6 males). Fifteen of the individuals were confiscated between 2006 and 2007, and the final male was born at JMC. The confiscated slow lorises came from the customs office of airports in Japan. Specific information on each subject is given in Supplementary Table 1. Prior to the establishment of the Slow Loris Conservation Center (SLCC) at JMC, the subjects were housed individually in small cages due to space limitations (0.1–0.3 m³), except for one father-son pair. Once the center was established, the lorises were introduced to larger enclosures, as described below. This study was carried out in accordance with the recommendations in the "Guide for Animal Research Ethics" of the Japan Monkey Center, and the legal requirements of the country, and ASP Principles for Ethical Treatment of Non-Human Primates. The study protocol was approved by the institutional committee of the Japan Monkey Center (#2015001, #2016001, #2017003).

Slow loris conservation center

The SLCC was established to improve the living conditions of confiscated pygmy slow lorises, and to serve as a place for environmental education, and slow loris research. The first slow lorises started to live there in 2015, and its facility and function gradually improved after that. The facility had previously been used for nocturnal primates, and was renovated by keepers and researchers to become the SLCC. Animal welfare is the first priority at SLCC, thus the renovations were carried out to match the needs of slow lorises. The facility uses a reverse-lighting cycle. Lighting levels were

measured on 19 December 2019, and were found to be between 0.00 and 0.18 lux during the dark phase, depending on the distance from the light in each room. We used red-filmed light in each room during dark phase observations, in order to minimize the effects of lighting on the slow lorises (Ariana, Marco, & Nekaris, 2020; Fuller et al., 2016; Starr, Nekaris, & Leung, 2012). Enclosure size varied from 2.0 to 16.0 m^3. For the male–male pair experiments, we used rooms that were about 8.0 m^3. Branches were installed to facilitate locomotion, and nest boxes and hiding places were added to each enclosure (Fitch-Snyder, Schulze, & Streicher, 2008). The number of nest boxes was equal to, or more than, the number of individuals in the enclosure. The animals also used the branches and the floor for resting and sleeping. Diets mainly comprised of gum, insects and vegetables were provided in multiple locations in each enclosure (Cabana & Nekaris, 2015; Cabana & Plowman, 2014). The mean temperature was 25.8 ± 2.08 ℃ (measured between January and December 2017). Organized tours with a limited number of zoo visitors were held several times a month. Participants of the organized tour could explore the center using red light with educational staff, who described the ecology and conservation of slow lorises and the activities of JMC. Before arriving at the SLCC, most lorises lived alone except for the father-son pair (Yashi-Hiiragi). They were kept alone at least since 2013. The details of the social management in the past were not well-recorded, but some individuals had social housing experience. Based on husbandry records, some individuals confiscated on the same date were kept in the same cages for a while after their arrival, but they were separated because of aggression or death of some individuals. Although there is a possibility that some individuals were housed together for a few months up to a few years when they were young, Yanagi and Shuro were never housed together before the current study.

Social housing that restricted breeding was necessary to accommodate all the slow lorises

housed at the facility when it was first established.

Social introduction

Each pair was formed in an enclosure that were unfamiliar to both individuals.

When necessary, we set up acclimation phases by placing small cages in close proximity

for several days or by allowing the individuals to spend time in adjacent cages. In the

beginning of this study, we conducted social introduction on a trial-and-error basis before

establishing the described acclimation protocol. We judged whether the pairing could

continue based on the behaviors exhibited. If the individuals immediately displayed

aggressive behavior, and little to no affiliative behaviors were observed, we decided to

stop the pairing and change the combination. If some affiliative behaviors were observed

with few or no aggressive behaviors, or if no social behaviors were observed at all, then

we continued the pairing. With this protocol, we formed five male–male pairs from eight

formation attempts with nine individuals (one male formed two pairs with different

individuals during different periods of the study, Supplementary Table 2). A tenth male

always behaved aggressively; therefore, he was kept solitary at all times, and was not

included further in the study. During the study of male-male pairing, they only had a

visual access to females. The males lived in adjacent cages where they could visually,

olfactorily, auditorily, and, possibly, tactilely socialize with other pairs. The females lived

in cages a short distance away, with glass partitions in-between (Supplementary Figure

1).

Data collection

The behaviors were recorded using infrared cameras (Zcam-CV, Bohankobo).

The cameras were attached to the ceiling and recorded each enclosure. In most cases, it

was difficult to identify the individuals from the recorded videos due to the video's

resolution, therefore, we recorded aggressive and non-aggressive interactions for the

lorises without identifying each individual. An ethogram is provided in Table 1. We

recorded aggressive behaviors using all-occurrence sampling methods (Martin & Bateson,

2007). We counted the bouts of aggressive behaviors. We included contact aggression in

the category of aggressive interactions, as it is often difficult to differentiate non-contact

aggression (i.e. chasing) from non-aggressive behaviors (i.e. following behaviors to

inspect conspecifics) on the recorded videos. We recorded affiliative social behaviors that

involved direct social contact (allogrooming, social play, co-feeding, nest box sharing)

every 5 minutes during the dark phase. We categorized the affiliative behaviors based on

(Fitch-Snyder & Ehrlich, 2003). We often observed that the two lorises entered a nest box,

and categorized such behaviors as a form of affiliative social behavior. We sometimes

observed the animals grooming each other or playing socially inside of the nest box, but

could not observe all the behaviors that occurred inside the boxes. In addition to

monitoring dark phase behaviors, we determined whether the lorises used the same sleep

sites during their light phase. We primarily did so using recorded videos, but also used

long-term data from Tsuge–Poplar, which were obtained by keepers (including the second

author (KN)) during the light phase.

Behavioral data recording

Behavioral data were collected between July 2016 and January 2020. Five types

of behavioral collections were made. We used the data obtained in different periods for

each purpose.

1. Description of social behavior in five pairs of male slow lorises

Behaviors were recorded from the 5 pairs of male slow lorises (9 individuals) for

a total of 150 hours between November 2016 and August 2017. We recorded their

behaviors 5 hours a day from the start of the dark phase for 5 days, 3–4 months after the

start of the social housing for each pair. We recorded the behavior for another 5 hours for

5 days and 1 year after the start of social housing for the Olive–Kunugi pair because the

pair took a longer time to exhibit affiliative behaviors.

2. Did they choose to stay in the same sleeping site during light phase?

Subjects were 5 pairs of male slow lorises (N = 9). A total of 450 days of sleeping

site data were obtained between August 2016 and October 2019. Sleeping occurred

primarily in their nest boxes, but the ground and branches were sometimes also used. We

considered that slow lorises shared a sleeping site when they slept in close proximity,

within reaching distance. Room temperature was recorded daily by keepers. To check

whether sleep sites were shared by chance, we used sleeping site data collected over a

month for each pair during the period when they formed affiliative relationships (four

pairs, except for the father–son pair, for which data were collected for 16 days only

because recordings were only available for this time period). To determine whether a

relationship between sleep site sharing and temperature existed, we used data from three

of the pairs. We selected the three pairs based on our ability to collect sleep site sharing

data and room temperature data simultaneously for these three pairs during a period when

they performed affiliative behaviors with almost no aggressive behaviors in their light

phase. We collected data from the Tsuge–Poplar pair for 281 days, the Nagi–Poplar pair

for 56 days and the Olive–Kunugi pair for 66 days.

3. Process of social group formation and changes in fecal glucocorticoid metabolites in

two pairs of male pygmy slow lorises

For this part of the experiment, we selected two pairs of male slow lorises as

subjects (N = 4). We recorded their behaviors 5 hours a day from the day after introduction

every day for 20 days, with additional sporadic observations after this period, for a total

of 260 hours between August 2016 and August 2017. We started each observation from

the start of the dark phase. Since the social relationships were established over different

time periods for the different pairs, the timing of the observations varied.

4. Sex differences in forming social relationships

The subjects were two pairs of male–male, female–female, and male–female

pairs (N = 10). Two males experienced two types of pairings during different periods of

the study. Total observation time was 144 hours between August 2016 and January 2020.

We observed each female–female and male–female pair for 12 days for 2 hours from the

start of the dark phase each day. Observations were made for the first 6 days after social

housing began, and then for another 6 days one month after the start of social housing.

Any injury incidents that occurred were recorded by keepers between August 2016 and

January 2020.

5. Stability of social relationship

Subjects were 3 pairs of male lorises (N = 6). Behaviors were compared between

two periods; after 3–4 months, and after 8–12 months from the start of the social housing.

We recorded their behaviors 3 hours each day from the start of the dark phase, for six

days per observation period. The total observation time was 108 hours between November

2016 and August 2017.

Fecal glucocorticoid metabolite (FGM) analysis

The feces from two pairs of male lorises were collected by keepers (N = 99) at

16:00 to 17:30 in the evenings (beginning of light phase). Fecal samples were collected

1–15 days before the start of social housing (11–14 days of sample collections for each

pair) and during 0–31 days after the start of social housing (20–32 days of sample

collections for each pair), between 15 July and 15 September, 2016. We could not identify

the individual feces once the pairs were housed together. We cannot exclude the

possibility of urine contamination, but the amount of slow loris urine is very small.

Therefore, compared with large-bodied primates, urine contamination is less likely to

occur. Before starting the social housings, individual feces were processed separately, and

the average value was used for the analyses. From one day after social housing, all the

feces collected were mixed and processed together. Collected feces were stored in freezers

(−20℃) until use.

We extracted FGM based on the methodologies developed by Kinoshita et al.

(2011). Feces were placed in a vacuum oven for about 18 hours at 60℃. The dried

samples were pulverized using a hammer and contaminants were removed as much as

possible using mesh. Approximately 0.02 g of the powdered samples were placed in a

conical tube and 3 ml of 80% methanol was added to each tube. The tubes were shaken

for 30 mins at ambient temperature, and then centrifuged for 10 mins at 3000 rpm. The

supernatant was then placed into new 2.0 ml tubes, and dried in a vacuum oven at $60°C$.

The dried samples were reconstituted with the assay buffer before an assay. Cortisol levels

were measured using enzyme immunoassay kits (Salimetric salivary cortisol kits: #1-

3002). Parallelism was confirmed, and the recovery rates were 85.6% on average. Inter-

assay variabilities were 3.92% (high) and 12.4% (low), respectively, and intra-assay

variabilities were 4.41% on average. Data with CV values that were higher than 10% were

removed from the analysis.

Statistical analysis

We used R 3.6.1 to perform all the statistical tests described below (R

Development Core Team, 2019). The data that support the findings of this study are

openly available in Open Science Framework (Yamanashi et al., 2020).

Did they choose to stay in the same sleeping site during light phase?

We performed two analyses, in order to determine whether the pairs chose to

share sleep sites during the light phase. First, we compared the rate of sleep site sharing

with the rate that could have occurred by chance. The chance level was calculated based

on Cohen's kappa calculation (Martin & Bateson, 2007). For example, suppose the two

individuals, A and B, used two locations, x and y, for sleeping, and that data was obtained

for 30 days. Individual A used sleeping location x a total of 11 times, and y a total of 19

times. Individual B used sleeping location x a total of 20 times, and y a total of 10 times.

The chance proportion of sharing the location x can be calculated by: $\frac{11}{30} \times \frac{20}{30}$ and that of

sharing the location y can be calculated by: $\frac{19}{30} \times \frac{10}{30}$. The chance proportion of sharing

the same sleeping site can then be calculated as $\left(\frac{11}{30} \times \frac{20}{30}\right) + \left(\frac{19}{30} \times \frac{10}{30}\right)$. We calculated the

chance proportion for each pair. Then, we checked whether the observed proportion of

sharing the sleep sites was higher than the chance proportion calculated as above by

binomial test. We used the function 'pbinom' in R.

We also checked the relationship between room temperature and sleep site

sharing using Man–Whitney's U tests for comparing temperatures when sleep sites were

shared, and when they did not. We used the function 'wilcox.exact' of the package

'exactRankTests' to perform the analyses (Hothorn, Hornik, van de Wiel, & Zeileis, 2006).

Changes in FGM levels

Fecal samples could not be assigned to specific individuals once the pairs were

housed together. We compared the FGM levels among three conditions: 15 days before

social housing ('before'), 15 days right after social housing began ('soon after', 0–14

days) and days 15–31 ('later') after social housing began for each pair. For statistical

analyses, we calculated the average values from each individual each day before social

housing. We used the values obtained from FGM measurements of mixed samples after

social housings. Among the four subjects, Nagi lived in a small cage and Poplar, Olive

and Kunugi each lived in one of the enclosures (8.0 m^3) during the 'before' period. We

used Kruskal–Wallis tests to compare the FGM levels across the three conditions. If we

statistically significant differences were determined, then we used Man–Whitney U tests

for multiple comparisons with the Bonferroni correction.

Sex differences in forming social relationships

The effect of sex on the formation process of social relationships was compared

using a generalized linear model. We used the function 'glm' with Poisson distribution in

R. Independent variables included sex (male–male, male–female, female–female),

introduction phase (0 months or 1 month), pairs and the interaction of sex and introduction

phase. Dependent variables included the number of times we observed aggressive and the

number of sampling points we observed affiliative behaviors. We also looked for

differences in the rate of affiliative behaviors one month after the start of social housing,

by constructing a model that included the independent variables of sex and pair. We

compared Akaike Information Criteria (AIC) among models composed of various

combinations of the aforementioned factors to identify the models that produced the

lowest AIC values. We used the dredge function in the package 'MuMIn' to choose the

best models (Barton & Barton, 2019). The AIC tables, including the models containing

$\triangle$AIC < 2 from the final model and the parameter estimates for the best models, are given

in the supplementary materials.

Stability of social relationships

We used a Mann–Whitney U test to compare the rates of social behaviors

between two periods for each pair. We used a proportion test to compare the probability

of sharing the sleeping sites.

Results

1. Social behaviors observed in the five male–male pairs

All five pairs performed some affiliative behaviors. On average, the pairs

continued for 595 ± 334 days (mean of four pairs), except for the father-son pair, who had

lived together from the birth to the death of the son (Supplementary Table 2).

Allogrooming, nest box sharing during dark phase and sleep site sharing during light

phase were observed in all pairs (Table 2, Figure 1). Social play was observed in three of

the pairs (Nagi-Poplar, Shuro-Yanagi, Yashi-Hiiragi: Supplementary video). Unlike other

pairs, one pair (Olive–Kunugi) did not show any affiliative behaviors during the

observations made three months after the start of the social housing; however, they did

show affiliative behaviors similar to the other pairs after one year had passed.

2. Did they choose to stay in the same sleeping site during light phase?

All pairs were observed to remain at the same sleeping site during the light phase.

They used multiple locations for sleeping together. The proportion of sleep site sharing

was above the calculated chance levels for all the pairs (Table 3).

There was no statistically significant relationship between temperature and sleep

site sharing for two of the pairs. The proportion of sleep site sharing was high for all the

three pairs (Tsuge–Poplar: 96.4%; Nagi–Poplar: 67.9%; Olive–Kunugi: 84.8%). Nagi–

Poplar had an average temperature of (mean ± SD): 26.2℃ ± 0.98℃ when they shared a

sleeping site, and an average temperature of 26.3℃ ± 0.73℃ when they slept alone

(Man–Whitney's U test: W = 328, P = 0.81). Olive–Kunugi had an average temperature

of 25.4℃ ± 0.75℃ when they shared a sleeping site, and an average temperature of

25.7℃ ± 0.75℃ when they did not (Man–Whitney's U test: W = 218, P = 0.25). In the

case of Tsuge–Poplar, the average temperature was relatively high when they shared sleep

sites with an average temperature 24.7℃ ± 0.73℃, and an average temperature of

24.2℃ ± 0.67℃ when they slept alone (Man–Whitney's U test: W = 817, P = 0.029). It

should be noted that they did not share sleep sites for 10 days out of 281, and that the

temperatures when they did not share sleep sites were within the range of those when they

shared.

3. Process of social group formation and changes in FGM in two pairs of male slow lorises

The male–male pairs showed some forms of aggression in the beginning of the

study that ceased after about 10 days. One pair (Nagi–Poplar) gradually increased their

affiliative behaviors after 14 days. It took more time for another pair (Olive–Kunugi) to

exhibit comparable levels of affiliative social behaviors (Figure 2). Sleep site sharing

during the light phase was observed after affiliative behaviors were observed during the

dark phase. Aggressive behaviors were observed most often after intense following of

another individual, during which the follower grabbed the other male and a short jostling

occurred.

Changes in FGM levels are shown in Figure 3. In the case of Nagi–Poplar, there

was a significant increase in FGM levels soon after the start of the experiment (Man–

Whitney's U test: 'before' vs 'soon after': $W = 22$, $P < 0.001$), but levels decreased to

lower than those observed before the social housing began (Man–Whitney's U test: 'soon

after' vs 'later': $W = 227$, $P < 0.001$; 'before' vs 'later': $W = 242$, $P < 0.001$). There was

no statistically significant change in FGM levels for Olive–Kunugi (Kruskal–Wallis test:

$X^2 = 2.5$, $df = 2$, $P = 0.28$). Before social housing, the FGM levels of Nagi were higher

than those of Poplar ($W = 164$, $P = 0.0091$).

4. Sex differences in forming social relationships

The sex of each animal did make a difference in social behavior during the

beginning phase of the social housing (Figure 4). Only male–male pairs showed

aggressive behaviors, while the other types of pairs did not show any aggressive behaviors

(Figure 4-a). The final model of aggressive behaviors included sex, introduction phase

and the interaction of sex and introduction phase (Supplementary Tables 3 and 4).

Parameter estimates of the final model indicated that aggression decreased one month

after the start of social housing, compared with the beginning phase. Female–female and

male–female pairs showed affiliative behaviors from the beginning, while male–male

pairs showed such behaviors later (Figure 4-b). The final model of affiliative behaviors

included sex, introduction phase, pair and the interaction of phase and sex

(Supplementary Tables 5 and 6). All types of pairs showed similar affiliative behaviors

by the end of the experiment. The final model explaining the factors associated with the

rate of affiliative behaviors one month after the start of the social housings only included

the pair factor, not sex (Supplementary Tables 7 and 8). Some female–female and male–

female pairs showed sleep sites share from the beginning, while male–male pairs showed

such behaviors later (Table 4).

Injuries were rarely observed, with a total of six incidents requiring veterinary

interventions between 2016 and 2019. Four individuals out of 16 received some form of

injury (three males and one female). Most injuries were treated without separation, but

the pairs in two cases were separated because the wounds did not heal well. The potential

effects of social grooming on slow recovery were suspected (Supplementary table 2).

5. Stability of social relationship

Social relationship stability differed across the pairs. One pair showed similar

levels of social behaviors between the two observation periods, another pair increased the

amount of affiliative behaviors and one pair decreased the amount of affiliative behaviors

and increased aggressive behaviors (Table 5). During the periods when aggressive

behaviors were observed, without any affiliative behaviors, sleep sites were not shared

during the light phase.

Discussion

All five pairs of male slow lorises formed affiliative relationships by the end of

the first observation period. Allogrooming, social play, nest box sharing during the dark

phase, and sleep site sharing during the light phase were the main behaviors observed.

Some pairs ate together at the same feeding platform in close proximity at times, even if

there were other places to forage. The rate of affiliative social behaviors was high in the

father-son pair, but some pairs also showed comparable levels of such behaviors. One

pair (Olive–Kunugi) did not show any aggressive or affiliative behaviors at times when

the other pairs would have; however, even this pair showed affiliative behaviors without

any aggressive behaviors by the end of the first observation period. This is surprising

considering reports of the wild lorises who rarely interact with same-sex adults, but they

can express social flexibility (Starr & Nekaris, 2020). In an environment without

resource competition or mating pressure, males can coexist, even into adulthood. It is

imperative that enclosures have adequate space and options to separate or hide from the

group in captivity (Global Federation of Animal Sanctuaries, 2013), especially during

the introductory phase. Additionally, it is necessary to closely monitor their

relationships considering the fact that the quality of relationship of the Yanagi–Shuro

pair deteriorated later.

All pairs shared sleep sites during the light phase. Sleep site sharing is often

observed in various wild, nocturnal strepsirrhines (Bearder et al., 2003; Nekaris &

Bearder, 2007; Rode et al., 2013), as researchers often analyze grouping tendencies by

observing daytime sleep sites in the wild (Bearder et al., 2003). Although it is rare for multiple males to share sleep sites, it has been reported in the slender loris and in giant mouse lemurs (Nekaris, 2003; Rode et al., 2013). This result is not consistent with those observed in the solitary sleeping patterns of wild pygmy slow lorises (Starr & Nekaris, 2020). Given their small body size, sleeping together may have a thermoregulatory advantage at lower temperatures, but room temperature in this experiment was set within the range recommended by the husbandry manual (Fitch-Snyder et al., 2001), and there was no evidence of increased sharing at lower room temperatures. On the contrary, one pair (Tsuge–Poplar) tended to increase their sharing rate when the temperature increased, which was not expected. Consideration has to be made for the fact that they rarely slept alone, and for the temperature range, which leads us to conclude that the trend was not strong. Furthermore, all the pairs used multiple sleep sites together, and the rate of sleep site sharing was above the chance level for all pairs. These results suggest that the pygmy slow lorises in this study preferred to stay together with other individuals during the light phase. In addition, relationships with both aggressive and affiliative behaviors during the dark phase suggest that sleep site sharing could only be observed during periods when their relationship was more affiliative. Therefore, sleep site sharing is a simple and good indicator of their relationship status.

The two pairs of males that showed aggression at the beginning of the experiment, had stopped displaying aggressive behaviors around day 10. Most aggressive behaviors did not appear to be serious, and veterinary interventions were rare. We did not record any deaths that were suspected to be related to the wounds incurred through aggressions, but sometimes, recovery from injuries took time. There are previous studies that have reported a few cases of death that may have been related to trauma or infection secondary

to bite wounds (Fuller, Lukas, Kuhar, & Dennis, 2014). In particular, slow lorises are

venomous primates (Nekaris et al., 2013), which results in a need for close monitoring

whenever adult pairs are formed to avoid serious injury or death.

Results of the changes in FGM levels for Nagi–Poplar indicates that their FGM

levels increased during their phase of heightened aggression, and decreased in the later

phase. The final FGM levels were even lower than those in the period when they were

housed alone in the small cage or an enclosure. This result suggests that forming pairs in

a more enriched enclosure is beneficial to reducing the level of physiological stress. This

interpretation of these benefits was supported by the fact that the FGM levels of Nagi

before social housing were higher than those of Nagi collected in the same period.

However, the results from Olive–Kunugi were not consistent with that of Nagi–Poplar.

There was no significant change in the level of FGM for Olive–Kunugi, which is counter

intuitive. This might be related to the fact that Olive and Kunugi already lived in an

enriched environment prior to the start of this study. Therefore, in addition to the

heightened aggression, changes in the physical environment may have affected the

increase in FGM levels of the Nagi-Poplar pair. Nevertheless, combined with the

behavioral results and FGM levels, there was no evidence of increased stress over a long

period of social housing. Instead, forming same-sex pairs increased the opportunity for

affiliative social behaviors.

There were sex-dependent differences in the formation process of social groups.

Aggressive behaviors were observed in the male–male pairs at the start of the experiment.

The male–female and female–female pairs showed affiliative behaviors from the

beginning. This low level of aggressive behaviors between females is interesting,

considering the fact that both adult males and females in the wild do not overlap their

home range with same-sex individuals (Starr & Nekaris, 2020). The details in the behaviors seen in female–female pairings will be reported in another paper (Alejandro et al. in prep). Injury risk was relatively high in the male–male pairs, but an injury was observed in a female-female pair once. Nevertheless, all types of pairs showed affiliative behaviors by the end of the experiment, without any serious incidence of aggression.

All-male groups have been formed in several captive primate species, including gorillas (Stoinski, Kuhar, et al., 2004), chimpanzees (Morimura, Idani, & Matsuzawa, 2010; Yamanashi, Teramoto, Morimura, Nogami, & Hirata, 2018), proboscis monkeys (Sha et al., 2013) and white crowned mangabeys (Fàbregas & Guillén-Salazar, 2007). All of these species are gregarious, and some of these species also have naturally occurring all-male groups in the wild, the type of species that typically form polygynous groups [e.g. gorillas (Gatti, Levréro, Ménard, & Gautier‐Hion, 2004) and proboscis monkeys (Murai, 2004)]. When mating competition is removed from the environment, as in all-male captive groups, coexistence of multiple adult males is more feasible. Fundamentally, male bonding is less easily occurred compared with females because males tend to compete for mating, which cannot be shared easily, in addition to food resources (Van Schaik, 1994). In some species, particularly male philopatric species, social bonding among males might be advantageous for mating success (Van Schaik, 1994). Pygmy slow lorises are not such species, rather they are territorial, and rarely overlap their home ranges with other same-sex adult individuals (Starr & Nekaris, 2020; Wiens & Zitzmann, 2003). Additionally, it is normal for other species to exhibit aggression, even after they establish their social relationships (Leeds, Boyer, Ross, & Lukas, 2015), which has led to the recommendation for some species to form social groups at an early age (Sha et al., 2013; Stoinski, Lukas, et al., 2004). Surprisingly, in the case of male pygmy slow lorises,

aggressive behaviors were very rare once they formed affiliative relationships, even when

they were introduced after sexual maturation. Familiarity could not explain these

relationships, except for the father-son pair as they were housed solitary at least since

2013. We are still not sure what the proximate and ultimate mechanisms of this flexible

trait are, but the results of this study indicate that this species has a high motivation to

interact with conspecifics if conditions are appropriate.

We are not sure if the results of this study can be generalized to other *Nycticebus*

species. Although similarity exists across different species of lorises, pygmy slow lorises

are unique in terms of their body size (smallest) and high rate of twin births (Fitch-Snyder

et al., 2001). It is noteworthy that the chemical compounds secreted from the brachial

glands of pygmy slow lorises were less than half of those of Bengal slow lorises (Hagey,

Fry, & Fitch-Snyder, 2007). Because the secretion from brachial glands is the main

component of the venom, bites from pygmy slow lorises might be less venomous and

thus, less risky for social housings. Nevertheless, recent findings on high sociability of

other slow lorises species show the importance of social management of *Nycticebus spp*.

Therefore, more studies on social behaviors of both wild and captive *Nycticebus spp*. are

needed to find out better social management strategies. It might be also important to study

other nocturnal strepsirhines that were also often misunderstood as solitary because of the

lack of understanding of their sociality.

There were some limitations to our study pertaining to the understanding of

social behaviors, and optimal social management regimes. First, we could not record

vocalizations and social behaviors that did not involve direct social contact. Studies have

reported that vocalizations often accompany social behaviors (Daschbach, Schein, &

Haines, 1981; Zimmermann, 1985). A recent study in Javan slow loris reported that the

lorises used ultrasonic vocalizations for communication (Geerah, O'Hagan, Wirdateti, &

Nekaris, 2019), and reported that lorises emitted vocalizations in the context of following

or leading each other, and entering a sleep site. In this study, we sometimes observed that

the pygmy slow lorises looked into a nest box where another individual was already inside,

or kept in social proximity for a while in front of the nest box before entering the box

together. It would be interesting to explore the details of pygmy slow loris communication

prior to social behavior initiation in future studies. A second limitation is the fact that the

judgement criteria used to test pair compatibility during the first introduction trial was

obscure. We introduced two unfamiliar individuals based on published suggestions to

minimize aggression during the introduction of a male and female pair (Fitch-Snyder et

al., 2001). We judged their compatibility based on behaviors displayed during their first

encounters. One male always showed aggressive tendencies to all individuals introduced,

thus, we did not form any long-lasting pair with him. We are not sure whether his

aggression was a personality trait or the result of some past experience. Since all other

pairs managed to form affiliative relationships by the end of the study, it is possible that

this individual may have successfully formed an affiliative relationship if given more time

or tried with a different individual.

In conclusion, male slow lorises appeared to be motivated to interact with

conspecifics. All pairs formed affiliative relationship by the end of the study, and they all

preferred to sleep together during their light phases. Furthermore, there was no evidence

of increasing physiological stress levels over a long period of time. These results suggest

that living with conspecifics increases the behavioral options of pygmy slow lorises, and

enables facilities to more efficiently use their available space and improve animal well-

being. We believe that providing opportunities to interact with conspecifics is one of the

most important parts of captive care for this species. Forming same-sex pairs might be a

feasible strategy for their social management. Although adult male–male pairings might

not be the first choice considering the risk of aggression during the introductory phase,

such flexibility in social housing should be considered to reduce the number of individuals

housed under suboptimal conditions in sanctuaries, zoos and rehabilitation centers.

Acknowledgement

This study was financially supported by captive care grant of the International

Primatological Society, Ishizue of Kyoto University, and Japan Society for Promotion of

Science (17K17828, JSPS-LGP-U04, JSPS core-to-core CCSN). It is also supported by

the Collaborative Research Program of Wildlife Research Center, Kyoto University. We

thank Tetsuro Matsuzawa, Genichi Idani, Naoto Kimura, Koshiro Watanuki, Mari

Hirosawa, Rui Hirokawa, Makiko Uchikoshi, Yuko Tawa, Yuta Shintaku and staff of the

Japan Monkey Center for their support for this project. We also thank Kei Matsushima,

Shiro Koshima, Kodzue Kinoshita, Satoshi Hirata, Takashi Hayakawa, Shinichi Kioka

for their valuable advice and kind cooperation for this study. Thanks are also due to Anna

Nekaris for her valuable advice for this project. We thank the editor and two anonymous

reviewers for their constructive comments. We have no conflict of interest to declare.

Reference

Amrein, M., Heistermann, M., & Weingrill, T. (2014). The effect of fission–fusion zoo housing

on hormonal and behavioral indicators of stress in Bornean orangutans (*Pongo*

pygmaeus). *International Journal of Primatology, 35*(2), 509-528. doi:10.1007/s10764-

014-9765-5

Ariana, W., Marco, C., & Nekaris, K. A. I. (2020). Red light for nocturnal observations. In

Evolution, Ecology and Conservation of Lorises and Pottos (Cambridge Studies in

Biological and Evolutionary Anthropology) (pp. 492-494). Cambridge, UK: Cambridge

University Press.

Barton, K., & Barton, M. K. (2019). Package 'MuMIn'. *R package version, 1*(6).

Bearder, S. K., Ambrose, L., Harcourt, C., Honess, P., Perkin, A., Pimley, E., . . . Svoboda, N.

(2003). Species-typical patterns of infant contact, sleeping site use and social cohesion

among nocturnal primates in Africa. *Folia Primatologica, 74*(5-6), 337-354.

Bloomsmith, M. A., Baker, K.C. (2001). Social Management of Captive Chimpanzees. In L.

Brent (Ed.), *The care and management of captive chimpanzees* (pp. 205-242). San

Antonio: American Society of Primatologists.

Cabana, F., & Nekaris, K. A. I. (2015). Diets high in fruits and low in gum exudates promote

the occurrence and development of dental disease in pygmy slow loris (*Nycticebus*

pygmaeus). *Zoo Biology, 34*, 547-553. doi:10.1002/zoo.21245

Cabana, F., & Plowman, A. (2014). Pygmy slow loris *Nycticebus pygmaeus*— natural diet

replication in captivity. *Endangered Species Research, 23*, 197-204.

Daschbach, N. J., Schein, M. W., & Haines, D. E. (1981). Vocalizations of the slow

loris,*Nycticebus coucang* (Primates, *Lorisidae*). *International Journal of Primatology,*

2(1), 71-80. doi:10.1007/BF02692301

Ehrlich, A., & Alan, M. (1977). Social and Individual Behaviors in Captive Slow Lorises.

Behaviour, 60(3/4), 195-220.

Fàbregas, M., & Guillén-Salazar, F. (2007). Social compatibility in a newly formed all-male

group of white crowned mangabeys (*Cercocebus atys lunulatus*). *Zoo Biology, 26*(1),

63-69. doi:10.1002/zoo.20117

Fitch-Snyder, H., & Ehrlich, A. (2003). Mother-infant interactions in slow lorises (*Nycticebus*

bengalensis) and pygmy lorises (*Nycticebus pygmaeus*). *Folia Primatologica, 74*(5-6),

259-271.

Fitch-Snyder, H., Schulze, H., & Larson, L. (2001). Management of lorises in captivity. *A*

husbandry manual for Asian lorisines.

Fitch-Snyder, H., Schulze, H., & Streicher, U. (2008). Enclosure design for captive slow and

pygmy lorises. *Primates of the oriental night. LIPI, Jakarta*, 123-135.

Fuller, G., Eggen, W. F., Wirdateti, W., & Nekaris, K. A. I. (2018). Welfare impacts of the
illegal wildlife trade in a cohort of confiscated greater slow lorises, Nycticebus
coucang. *Journal of Applied Animal Welfare Science, 21*(3), 224-238.
doi:10.1080/10888705.2017.1393338

Fuller, G., Kuhar, C. W., Dennis, P. M., & Lukas, K. E. (2013). A survey of husbandry practices
for lorisid primates in North American zoos and related facilities. *Zoo Biology, 32*(1),
88-100. doi:10.1002/zoo.21049

Fuller, G., Lukas, K. E., Kuhar, C., & Dennis, P. M. (2014). A retrospective review of mortality
in lorises and pottos in North American zoos, 1980-2010. *Endangered Species*
Research, 23(3), 205-217.

Fuller, G., Raghanti, M. A., Dennis, P. M., Kuhar, C. W., Willis, M. A., Schook, M. W., & Lukas,
K. E. (2016). A comparison of nocturnal primate behavior in exhibits illuminated with
red and blue light. *Applied Animal Behaviour Science, 184*, 126-134.
doi:http://dx.doi.org/10.1016/j.applanim.2016.08.011

Gatti, S., Levréro, F., Ménard, N., & Gautier‐Hion, A. (2004). Population and group
structure of western lowland gorillas (*Gorilla gorilla gorilla*) at Lokoue, Republic of
Congo. *American Journal of Primatology: Official Journal of the American Society of*
Primatologists, 63(3), 111-123.

Geerah, D. R., O'Hagan, R. P., Wirdateti, W., & Nekaris, K. A. I. (2019). The use of ultrasonic
communication to maintain social cohesion in the Javan slow loris (*Nycticebus*
javanicus). *Folia Primatologica, 90*(5), 392-403. doi:10.1159/000495077

Global Fereration of Animal Sanctuaries. (2013). Retrieved from

Hagey, L. R., Fry, B. G., & Fitch-Snyder, H. (2007). Talking Defensively, a Dual Use for the
Brachial Gland Exudate of Slow and Pygmy Lorises. In S. L. Gursky & K. A. I.
Nekaris (Eds.), *Primate Anti-Predator Strategies* (pp. 253-272). Boston, MA:
Springer US.

Honess, P. E., & Marin, C. M. (2006). Behavioural and physiological aspects of stress and
aggression in nonhuman primates. *Neuroscience & Biobehavioral Reviews, 30*(3),
390-412.

Hothorn, T., Hornik, K., van de Wiel, M., & Zeileis, A. (2006). A Lego system for conditional
inference. *The American Statistician, 60*(3), 257-263. doi:10.1198/000313006X118430

IUCN. (2020). The IUCN Red List of Threatened Species. Version 2020-2. Retrieved from
https://www.iucnredlist.org

Kinoshita, K., Inada, S., Seki, K., Sasaki, A., Hama, N., & Kusunoki, H. (2011). Long-term
monitoring of fecal steroid hormones in female snow leopards (*Panthera uncia*)
during pregnancy or pseudopregnancy. *PLOS ONE, 6*(5), e19314.

doi:10.1371/journal.pone.0019314

Kitade, T., & Naruse, Y. (2020). Crossing the red line: Japan's exotic pet trade. Retrieved from

https://www.traffic.org/publications/reports/crossing-the-red-line/

Kutsukake, N., Teramoto, M., Honma, S., Mori, Y., Ikeda, K., Yamamoto, R., . . . Hasegawa,

T. (2018). Behavioural and hormonal changes during group formation by male

chimpanzees. *Behaviour*, 1-21. doi:https://doi.org/10.1163/1568539X-00003532

Leeds, A., Boyer, D., Ross, S. R., & Lukas, K. E. (2015). The effects of group type and young

silverbacks on wounding rates in western lowland gorilla (*Gorilla gorilla gorilla*)

groups in North American zoos. *Zoo Biology, 34*(4), 296-304.

doi:doi:10.1002/zoo.21218

Martin, P., & Bateson, P. (2007). *Measuring behaviours: an introductory guide* (3rd ed.).

Cambridge: Cambridge University Press.

Moore, R. S., Cabana, F., & Nekaris, K. A. I. (2015). Factors influencing stereotypic

behaviours of animals rescued from Asian animal markets: A slow loris case study.

Applied Animal Behaviour Science, 166, 131-136.

doi:https://doi.org/10.1016/j.applanim.2015.02.014

Moore, R. S., Wihermanto, & Nekaris, K. A. I. (2014). Compassionate conservation,

rehabilitation and translocation of Indonesian slow lorises. *Endangered Species*

Research, 26(2), 93-102.

Morimura, N., Idani, G. i., & Matsuzawa, T. (2010). The first chimpanzee sanctuary in Japan:

an attempt to care for the "surplus" of biomedical research. *American Journal of*

Primatology, 73(3), 226-232. doi:10.1002/ajp.20887

Murai, T. (2004). Social behaviors of all-male proboscis monkeys when joined by females.

Ecological Research, 19(4), 451-454. doi:10.1111/j.1440-1703.2004.00656.x

Musing, L., Suzuki, K., & Nekaris, K. A., I. (2015). Crossing international borders: the trade

of slow lorises, Nycticebus spp., as pets in Japan. *Asian Primates*.

Nekaris, K. A. I. (2003). Spacing system of the Mysore slender loris (*Loris lydekkerianus*

lydekkerianus). *American Journal of Physical Anthropology, 121*(1), 86-96.

doi:10.1002/ajpa.10189

Nekaris, K. A. I. (2007). Unexpected diversity of slow lorises (*Nycticebus spp.*) within the

Javan pet trade: implications for slow loris taxonomy. *Contributions to zoology*

(Amsterdam, Netherlands : 1995), 76(3), 187.

Nekaris, K. A. I. (2014). Extreme primates: ecology and evolution of Asian lorises.

Evolutionary Anthropology: Issues, News, and Reviews, 23(5), 177-187.

Nekaris, K. A. I., & Bearder, S. (2007). The Lorisiform Primates of Asia and Mainland Africa:

Diversity Shrouded in Darkness. In *Primate in perspective*: Oxford University Press.

Nekaris, K. A. I., & Bearder, S. K. (2007). The Lorisiform primates of Asia and mainland
Africa. *Primates in Perspective. Oxford University Press, New York*, 24-45.

Nekaris, K. A. I., Campera, M., Nijman, V., Birot, H., Rode-Margono, E. J., Fry, B. G., . . .
Imron, M. A. (2020). Slow lorises use venom as a weapon in intraspecific competition.
Current Biology, 30(20), R1252-R1253. doi:https://doi.org/10.1016/j.cub.2020.08.084

Nekaris, K. A. I., Moore, E., Rode, J., & Fry, B. (2013). Mad, bad and dangerous to know: the
biochemistry, ecology and evolution of slow loris venom. *Journal of Venomous*
Animals and Toxins including Tropical Diseases, 19-21.

Norconk, M. A., Atsalis, S., Tully, G., Santillan, A. M., Waters, S., Knott, C. D., . . . Stiles, D.
(2020). Reducing the primate pet trade: Actions for primatologists. *Am J Primatol,*
82(1), e23079. doi:10.1002/ajp.23079

Poindexter, S., & Nekaris, K. A. I. (2020). The evolution of social organisation in lorisiformes.
In K. A. I. Nekaris & A. M. Burrows (Eds.), *Evolution, Ecology and Conservation of*
Lorises and Pottos. Cambridge: Cambridge University Press.

Price, E. E., & Stoinski, T. S. (2007). Group size: Determinants in the wild and implications
for the captive housing of wild mammals in zoos. *Applied Animal Behaviour Science,*
103(3–4), 255-264. doi:http://dx.doi.org/10.1016/j.applanim.2006.05.021

R Development Core Team. (2019). R: A language and environment for statistical computing.
(3-900051-07-0). http://www.R-project.org/

Reeder, D. M., & Kramer, K. M. (2005). Stress in free-ranging mammals: integrating
physiology, ecology, and natural history. *Journal of Mammalogy, 86*(2), 225-235.
doi:10.1644/bhe-003.1

Rode-Margono, E. J., Nijman, V., Wirdateti., & Nekaris, K. A. I. (2014). Ethology of the
critically endangered Javan slow loris *Nycticebus javanicus* E. Geoffroy Saint-Hilaire
in West Java. *Asian Primates, 4*, 27-41.

Rode, E. J., Nekaris, K. A. I., Markolf, M., Schliehe-Diecks, S., Seiler, M., Radespiel, U., . . .
Nijman, V. (2013). Social organisation of the northern giant mouse lemur Mirza zaza
in Sahamalaza, north western Madagascar, inferred from nest group composition and
genetic relatedness. *Contributions to Zoology, 82*(2), 71.
doi:https://doi.org/10.1163/18759866-08202001

Selye, H. (1941). Pharmacological classification of steroid hormones. *Nature, 148*, 84-85.

Sha, J. C. M., Alagappasamy, S., Chandran, S., Cho, K. M., & Guha, B. (2013). Establishment
of a captive all-male group of proboscis monkey (*Nasalis larvatus*) at the Singapore
zoo. *Zoo Biology, 32*(3), 281-290. doi:10.1002/zoo.21020

Squires, J. E. (2010). *Applied animal endocrinology: 2nd edtion*. Oxfordshire: CAB
International.

Starr, C., & Nekaris, K. A. I. (2020). Ranging Patterns of the Pygmy Slow Loris (*Nycticebus*
pygmaeus) in a Mixed Deciduous Forest in Eastern Cambodia. In K. A. I. Nekaris &
A. M. Burrows (Eds.), *Evolution, Ecology and Conservation of Lorises and Pottos*.
Cambridge: Cambridge University Press.

Starr, C., Nekaris, K. A. I., & Leung, L. (2012). Hiding from the moonlight: luminosity and
temperature affect activity of Asian nocturnal primates in a highly seasonal forest.
PloS One.

Stoinski, T. S., Kuhar, C. W., Lukas, K. E., & Maple, T. L. (2004). Social dynamics of captive
western lowland gorillas living in all-male groups. *Behaviour, 141*(2), 169-195.
doi:10.1163/156853904322890807

Stoinski, T. S., Lukas, K. E., Kuhar, C. W., & Maple, T. L. (2004). Factors influencing the
formation and maintenance of all-male gorilla groups in captivity. *Zoo Biology, 23*(3),
189-203. doi:doi:10.1002/zoo.20005

Tennenhouse, E. M., Putman, S., Boisseau, N. P., & Brown, J. L. (2017). Relationships
between steroid hormones in hair and social behaviour in ring-tailed lemurs (*Lemur*
catta). *Primates, 58*(1), 199-209. doi:10.1007/s10329-016-0566-7

van Schaik, C. P. (1994). Male bonds: afilliative relationships among nonhuman primate
males. *Behaviour, 130*(3-4), 309-337. doi:10.1163/156853994X00587

van Schaik, C. P. (1999). The socioecology of fission-fusion sociality in Orangutans. *Primates,*
40(1), 69-86. doi:10.1007/bf02557703

Weingrill, T., Willems, E. P., Zimmermann, N., Steinmetz, H., & Heistermann, M. (2011).
Species-specific patterns in fecal glucocorticoid and androgen levels in zoo-living
orangutans (*Pongo spp.*). *General and Comparative Endocrinology, 172*(3).

Wiens, F. (2002). Behavior and ecology of wild slow lorises (*Nycticebus coucang*): social
organization, infant care system, and diet.

Wiens, F., & Zitzmann, A. (2003). Social structure of the solitary slow loris Nycticebus
coucang (*Lorisidae*). *Journal of Zoology, 261*(1), 35-46.
doi:10.1017/s0952836903003947

Yamanashi, Y. (2020). Social relationships among captive male pygmy slow lorises
(*Nyctycebus pygmaeus*): Is forming male same-sex pairs a feasible management
strategy? Retrieved from: https://osf.io/bdtas

Yamanashi, Y., Nogami, E., Teramoto, M., Morimura, N., & Hirata, S. (2018). Adult-adult
social play in captive chimpanzees: Is it indicative of positive animal welfare? *Applied*
Animal Behaviour Science, 199, 75-83.
doi:https://doi.org/10.1016/j.applanim.2017.10.006

Yamanashi, Y., Teramoto, M., Morimura, N., Hirata, S., Inoue-Murayama, M., & Idani, G.

816   (2016). Effects of relocation and individual and environmental factors on the long-

817   term stress levels in captive chimpanzees (*Pan troglodytes*): monitoring hair cortisol

818   and behaviors. *Plos One, 11*(7), e0160029. doi:10.1371/journal.pone.0160029

819 Yamanashi, Y., Teramoto, M., Morimura, N., Nogami, E., & Hirata, S. (2018). Social

820   relationship and hair cortisol level in captive male chimpanzees (*Pan troglodytes*).

821   *Primates, 59*(2), 145-152. doi:10.1007/s10329-017-0641-8

822 Zimmermann, E. (1985). Vocalizations and associated behaviours in adult slow loris

823   (*Nycticebus coucang*). *Folia Primatologica, 44*(1), 52-64. doi:10.1159/000156197

824   Table 1. Ethogram used for the study

Category	Behaviors	Definition
Affiliative behaviors	Social grooming	Lick fur and apply toothcomb while grasping other animal with one or both hands.
	Play	Mock fight (mutual biting, hiting, classping but no other signs of threat; low intensity, prolonged, neither animal tries to get away)
	Co-feeding	Eat foods from the same dish in proximity
	Sharing a nest box	Enter into a same nest box during dark phase
	Sharing a sleeping site	Enter the same nest box or sleep in a close proximity, within reaching distance, during light phase
Aggressive behaviors	Attack	Bite, hit hard with hands or push with apparent force; lunge and bite at the same time, butt with head with apparent force
	Fight	Mutual hitting and biting with apparent force; wrestle vigorously

829　Table 2. The rate of social behaviors among five pairs of male slow lorises (5 hours for 5

830　days/column)

831　We recorded their behaviors 3–4 months after the start of the social housing for each pair

832　and 1 year after the start of social housing for the Olive–Kunugi.

	Yashi–Hiiragi	Shuro - Yanagi	Nagi–Poplar	Olive–Kunugi		Tsuge–Poplar
Sampling period (months)	Father-son	4	4	3	12	3
Aggression	0	0	0	0	0	0
Affiliation	0.38 ± 0.043	0.14 ± 0.059	0.31 ± 0.14	0	0.27 ± 0.15	0.11 ± 0.084
Sleep site share(days)	5	5	3	0	4	5

835   Table 3. The proportion of sleep site sharing in the five male pairs

	Number of sleeping site used	Rate of sleeping site share	Rate of sleeping site share expected by chance	P
Nagi-Poplar	3	0.68	0.38	<0.001
Shuro-Yanagi	2	1.00	0.85	<0.001
Olive-Kunugi	3	0.81	0.68	0.018
Tsuge-Poplar	4	0.90	0.46	<0.001
Yashi-Hiiragi	3	0.94	0.64	<0.001

839   Table 4. Sex differences in the proportion of sleep site sharing

840   The data were obtained on the first 6 days (0 month), then again for 6 days one month

841   after the start of the social housing (1 month). The data of final phase of male-male pairs

842   derived from the study 1 (5 pairs). The data of female-female and male-female pairs were

843   the average of data obtained one month after the start of the social housing (2 pairs each).

Phase	Pair	Male-Male	Female-Female	Male-Female
0 month	Pair 1	0.00	0.17	0.00
	Pair 2	0.00	0.50	1.00
1 month	Pair 1	0.67	0.50	1.00
	Pair 2	0.00	0.00	1.00
Final	Average of the pairs	0.92	0.25	1.00

845 Table 5. Changes in social behaviors in the three male pairs between two periods

	Shuro – Yanagi			Nagi-Poplar			Olive-Kunugi		
Sampling period (months)	4	8		4	8		3	12	
Aggression	0	1.2 ± 1.3	W = 6, P = 0.061	0	0	W =18, P =1	0.17 ± 0.37	0	W = 21, P =1
Affiliation	0.15 ± 0.048	0	W =36, P = 0.0022	0.28 ± 0.13	0.26 ± 0.064	W = 19, P = 0.98	0	0.29 ± 0.19	W = 0, P = 0.0022
Sleeping site share(days)	6	0	X^2=3.4, P = 0.066	4	5	X^2=0, P = 1	0	4	X^2=8.3, P < 0.001

846
847

Figure 1. A photo of two males (Shuro and Yanagi) sharing a nest box

Figure 2. Changes in the social behavior from the start of the social housing in two pairs.

Figure 2-a. Changes in aggressive behaviors for Nagi–Poplar.

Figure 2-b. Changes in affiliative behaviors for Nagi–Poplar.

Figure 2-c. Changes in aggressive behaviors for Olive–Kunugi

Figure 2-d. Changes in affiliative behaviors for Olive–Kunugi

857
858

859   Figure 3. Changes in FGM levels before and after social housing.

860   Figure 3-a. Nagi–Poplar

861   Figure 3-b. Olive–Kunugi

862
863

Figure 4. Sex differences in social behaviors during the beginning phase of social housing.

FF represents data from the two female-female pairs, MF represents data from the male–

female pairs, and MM represents data from the male–male pairs. The data were obtained

on the first 6 days (0M), then again for 6 days one month after the start of the social

housing (1M). The data of MM final derived from the study 1.

Figure 4-a. Sex differences in aggressive behaviors

Figure 4-b. Sex differences in affiliative behaviors

Fig. 4-a

Fig. 4-b

872   Supplementary Table 1. Subject information

Name	Sex	Date of arrival	Origin	Pairing
Yanagi	M	2006/12/7	Confiscation (Narita)	Shuro
Tsubaki	F	2007/8/2	Confiscation (Osaka)	Tsuge
Mizuki	F	2007/8/2	Confiscation (Osaka)	Nanten
Momiji	F	2007/8/2	Confiscation (Osaka)	Poplar
Olive	M	2007/8/28	Confiscation (Osaka)	Kunugi
Nagi	M	2007/8/28	Confiscation (Osaka)	Poplar
Kunugi	M	2007/8/28	Confiscation (Osaka)	Olive
Nanten	F	2007/8/28	Confiscation (Osaka)	Mizuki
Hiiragi	M	2007/8/28	Confiscation (Osaka)	Yashi
Aoki	M	2007/8/28	Confiscation (Osaka)	Unsuccessful
Shuro	M	2007/8/28	Confiscation (Osaka)	Yanagi
Kaede	F	2007/8/28	Confiscation (Osaka)	Mimoza
Poplar	M	2007/8/28	Confiscation (Osaka)	Nagi, Tsuge, Momiji
Tsuge	M	2007/8/28	Confiscation (Osaka)	Poplar, Tsubaki
Mimoza	F	2007/8/28	Confiscation (Osaka)	Kaede
Yashi	M	2010.03.28	Born at the JMC	Hiiragi

875   Supplementary Table 2. Duration of each pair

Pair	Type of pair	Start	End	Reason of ending the pair
Nagi - Poplar	MM	2016/8/1	2017/3/31	To treat an injury of Poplar
Olive - Kunugi	MM	2016/8/14	2019/1/21	To form a male-female pair
Yashi - Hiiragi	MM	2016/9/16	2018/5/30	Yashi passed away from sickness.
Shuro - Yanagi	MM	2016/11/29	2017/12/12	Shuro passed away from sickness.
Tsuge - Poplar	MM	2017/5/26	2019/10/14	To form a male-female pair
Kaede - Mimoza	FF	2017/1/7	2017/7/2	To treat an injury of Mimoza
Nanten - Mizuki	FF	2017/9/12	2018/11/5	To form a male-female pair
Poplar - Momiji	MF	2019/10/14	Ongoing	
Tsuge - Tsubaki	MF	2019/11/27	Ongoing	

877 Supplementary Table 3. AIC table for explaining the factors associated with occurrence
878 of aggressive behaviors

AIC	Independent variables
70.0	Sex + Phase + Sex : Phase
70.6	Sex + Phase + Pair + Sex : Phase
71.4	Sex + Phase
72.0	Phase + Pair
72.0	Sex + Phase + Pair

880 Supplementary Table 4. Parameter estimates for the best-fit model to explain the factors
881 associated with occurrence of aggressive behaviors

	Factor	estimate	SE	z	p
	Intercept	0.51	0.22	2.3	0.022
Sex	Female- female	-21	4428	-0.0050	1.0
	Male - female	-21	4428	-0.0050	1.0
Phase	1 month	-3.0	1.0	-2.9	0.0035
Sex Phase	Female-female : 1month	21	4487	0.0050	1.0
	Male-female : 1month	3.0	6345	0	1.0

883 Supplementary Table 5. AIC table for explaining the factors associated with the rate of
884 affiliative behaviors

AIC	Independent variables
329	Sex + Phase + Pair + Sex : Phase

886 Supplementary Table 6. Parameter estimates for the best-fit model to explain the factors
887 associated with the rate of affiliative behaviors

	Factor	estimate	SE	z	p
	Intercept	-20	1436	-0.014	0.99
Sex	Female-female	22	1436	0.016	0.99
	Male-female	23	1436	0.016	0.99
Phase	1 month	2.7	0.60	4.4	<0.001
Pair	Mizuki-Nanten	-0.11	0.15	-0.74	0.46
	Nagi-Poplar	19	1436	0.013	0.99
	Olive-Kunugi	NA	NA	NA	NA

	Poplar-Momiji	-1.6	0.15	-11	<0.001
	Tsuge-Tsubaki	NA	NA	NA	NA
Sex : Phase	Female-female : 1month	-3.0	0.62	-4.9	<0.001
	Male-female : 1month	-2.5	0.61	-4.1	<0.001

891 Supplementary Table 7. AIC table for explaining the factors associated with the rate of

892 affiliative behaviors one month after the start of the social housings

AIC	Independent variables
156	Pair
156	Sex + Pair

894 Supplementary Table 8. Parameter estimates for the best-fit model to explain the factors

895 associated with the rate of affiliative behaviors one month after the start of the social

896 housings

	Factor	estimate	SE	z	p
	Intercept	2.1	0.14	14	<0.001
Pair	Mizuki-Nanten	-0.50	0.2	-2.1	0.032
	Nagi-Poplar	-0.11	0.21	-0.52	0.60
	Olive-Kunugi	-20	2334	-0.0090	0.99
	Poplar-Momiji	-0.50	0.24	-2.1	0.032
	Tsuge-Tsubaki	1.1	0.17	6.3	<0.001

900 Supplementary Figure 1. A schematic of the Slow Loris Conservation Centre (SLCC).
901 Male-male pairings were conducted in cages A-E, while male-female and female-female
902 pairings were conducted in all types of cages. During the study periods of male-male
903 pairings, females lived in cages F- J. G-H and I-J can either be used as a single cage or
904 two adjacent cages.
905

906
907
908 Supplementary video
909 https://youtu.be/3gv1qbnppBo