Wiring diagram of the oxytocin system in the mouse brain

Seoyoung Son[1], Steffy B. Manjila[1], Kyra T. Newmaster[1], Yuan-ting Wu[1], Daniel J. Vanselow[2],
Matt Ciarletta[1], Keith C. Cheng[2], and Yongsoo Kim[1]

1 Department of Neural and Behavioral Sciences, Penn State University, Hershey, PA, USA

2 Department of Pathology, Penn State University, Hershey, PA, USA

Corresponding author:

Yongsoo Kim

Email: yuk17@psu.edu

Keyword: Oxytocin, Oxytocin receptor, Brain mapping, synaptic input, axonal output,
anatomical connectivity, serial two-photon tomography, connectome

Abstract

In the brain, oxytocin (OT) neurons make direct connections with discreet regions to regulate social behavior and diverse physiological responses. Obtaining an integrated neuroanatomical understanding of pleiotropic OT functions requires comprehensive wiring diagram of OT neurons. Here, we have created a whole-brain map of distribution and anatomical connections of hypothalamic OT neurons, and their relationship with OT receptor (OTR) expression. We used our brain-wide quantitative mapping at cellular resolution combined with a 2D flatmap to provide an intuitive understanding of the spatial arrangements of OT neurons. Then, we utilized knock-in *Ot-Cre* mice injected with Cre dependent retrograde monosynaptic rabies viruses and anterograde adeno associated virus to interrogate input-output patterns. We find that brain regions with cognitive functions such as the thalamus are reciprocally connected, while areas associated with physiological functions such as the hindbrain receive unidirectional outputs. Lastly, comparison between OT output and OTR expression showed no significant quantitative correlation, suggesting that OT transmission mostly occurs through indirect pathways. In summary, our OT wiring diagram provides structural and quantitative insights of distinct behavioral functions of OT neurons in the brain.

Significance Statement

Oxytocin (OT) neurons in the brain play an important role in socio-physiological responses. Impairment of OT signaling has been implicated in many neurodevelopmental disorders. To understand diverse OT functions in the context of discreet neural circuits, it is imperative to understand the anatomical arrangement of OT neurons across the whole brain in significant detail. Here, we have established a comprehensive brain-wide wiring diagram of OT neurons. Our anatomical and connectivity map of OT neurons includes brain-wide cell distribution, synaptic inputs, axonal outputs, and their relationships with the oxytocin receptor expression. This whole brain structural perspective of the OT system provides a foundation for understanding the diversity of neural circuits modulated by OT and will guide future circuit-based OT functional studies.

Introduction

Oxytocin (OT) is a highly conserved neuropeptide, playing key roles in regulating social behavior and other physiological functions (1–4). Impairment in OT signaling has been heavily implicated in many neurodevelopmental disorders including autism (5, 6). Correcting OT signaling is being pursued as a potential therapy to alleviate social behavioral symptoms in many brain disorders (7), but our limited neurobiological understanding of OT signaling in the brain hampers the development of targeted approaches aimed at altering specific OT functions without affecting other biological pathways. A comprehensive anatomical understanding of OT neurons would enable integrated neural circuit specific studies to decipher the neural substrate of distinct OT functions.

The majority of OT producing neurons are located in the paraventricular nucleus of the hypothalamus (PVH) and the supraoptic nucleus (SO) while fewer OT neurons reside in the bed nucleus of the stria terminalis (BST), the medial amygdala (MEA), and the accessory nuclei (AC) (8). OT neurons receive input from distinct brain regions [e.g., the thalamus (TH)] to integrate sensory input and internal information to release OT in a context dependent manner in order to modulate specific downstream circuitry (9, 10). The actions of OT are mainly mediated by a single subtype of the OT receptor (OTR) expressed in distinct brain regions as well as peripheral tissues (11–13). In addition to the well-known peripheral release of OT as a hormone via the posterior pituitary, OT neurons send direct projections to discreet brain areas that frequently express OTR, thereby modulating circuit specific functions. For example, OT signaling is linked with the medial prefrontal cortex for social cognition (14, 15), CA2 of the hippocampus for social memory (16, 17), the central amygdala for fear modulations (18, 19), the parabrachial nucleus (PB) for fluid intake (20), and the spinal cord for pain perception (21, 22). Despite these prior studies, we still lack a quantitative and comprehensive wiring diagram of the OT neurons in a standard 3D reference brain. Moreover, quantitatively characterizing the wiring relationships between OT neurons and OTR expression patterns across the whole brain is made difficult due to the need to simultaneously examine microscopic structures (e.g., cell bodies and their axonal projections) in a macroscopic structure, namely the whole brain.

Here, we use a high-resolution quantitative brain mapping method in combination with cell type specific transgenic mice and viral tools to establish the wiring diagram of OT neurons in the mouse brain. We devised a 2D hypothalamic flatmap to quantify OT distribution, and analyzed synaptic inputs and long-range output patterns of hypothalamic OT neurons. We find reciprocally and unidirectionally connected brain regions with OT neurons, linked with distinct cognitive and physiological functions, respectively. Moreover, systemic comparisons between the OT projectome and OTR expression reveal potential brain regions with direct synaptic or non-synaptic OT signaling. The imaging data are mapped onto a standard reference brain to facilitate cross-examination (23) and freely available via user-friendly web visualizations at https://kimlab.io/brain-map/ot_wiring/.

Results

Quantitative expression of oxytocin neurons in the mouse brain

Although the location of OT neurons in the mouse hypothalamus (HY) is relatively well-established (8), data showing quantitative brain-wide OT distribution in complex 3D structures remains elusive. To examine the anatomical distribution of OT neurons across the whole brain, we used heterozygote OT knock-in mice with Cre recombinase (*Ot-Cre*) crossed with Ai14 reporter mice (OT:Ai14) (24). We imaged the entire mouse brain at cellular resolution using serial two-photon tomography (STPT) and performed quantitative mapping using previously established computational methods (25) (n=8 brains, 3 males, 3 virgin females, and 2 lactating females, Fig. 1A-B, Movie S1). There was no significant difference between male, virgin female, and lactating female (Table S1), which is consistent with an earlier study (18). The majority of OT neurons (~42%, 792 out of total 1896 cells) were expressed in the PVH regions (PVH, descending division of PVH, anterior, intermediate, and preoptic part of periventricular hypothalamic nucleus, and subparaventricular zone) followed by the SO, the tuberal nucleus (TU), the MEA, and the BST (Table S1). The rest of the OT neurons are expressed in 10 different brain regions (Table S1). The Cre-driven reporter system used in the present study permanently labels all Cre positive cells including developmentally transient expression (11). To distinguish neurons actively expressing OT in adult from developmentally labeled cells, we performed immunohistochemistry using an OT antibody in OT:Ai14 mice. We confirmed that almost all OT immuno positive neurons (97%, 1733 out of 1790 cells, n=4) were labeled by tdTomato from OT:Ai14 mice (Fig. S1). In contrast, 76% of tdTomato labeled cells were OT immuno positive (1733 out of 2277 cells) in the PVH. Smaller portions of tdTomato cells in the SO (44%, 654 out of 1508 cells) and the MEA (8%, 31 out of 375 cells) retain active OT expression (Fig. S1). This suggests that OT neurons in different brain regions undergo differential rates of developmental down-regulation in the adult brain (26). To further visualize the spatial expression pattern of OT neurons, we created a 2D hypothalamic flatmap. Evenly spaced bins provide a flattened 2D spatial unit to quantify and to display signals from the 3D brain (Fig. 1C-D). The 2D hypothalamic flatmap was delineated with Allen Common Coordinate Framework (CCF) and Franklin-Paxinos atlas based anatomical labels (23, 27, 28) (Fig 1C). The regional boundaries of the two labeling systems generally agreed with each other in the major OT expressing regions (e.g., the PVH and the SO) despite noticeable discrepancies in the ventricle hypothalamic area (e.g., the TU) (Fig 1C-D) (28). The density heatmap on the hypothalamic flatmap provides an intuitive and quantitative display of the regionally heterogeneous distribution of OT neurons (Fig. 1D). We also provide detailed distribution patterns of the OT neurons within the PVH using a 2D PVH flatmap (Fig. S2). Together, these data provide the quantitative expression pattern of OT neurons throughout the whole brain.

Figure 1. Distribution of oxytocin neurons.
(A) Signals from OT:Ai14 across representative coronal planes of the mouse brain. (top) STPT images registered onto the Allen CCF reference brain, (bottom) averaged signals (n=8, green dots) overlaid on the reference brain. White and red arrowheads indicate the PVH and SO, the separately. Scale bar = 1mm. See also Movie S1, Table S1. (B) 3D distribution of oxytocin neurons. A/P: anterior/posterior, L/M: Lateral/medial, D/V: dorsal/ventral. (C) 2D hypothalamic flatmaps. Small inserts with coronal sections illustrate how bins (while areas with numbers) were generated at different coronal planes. Anatomical labels in the flatmap are delineated based on Allen mouse brain atlas (left) and Franklin-Paxinos atlas (right). The X-axis is for bin numbers and Y-axis is for the bregma A/P axis. (D) Heatmap of oxytocin neuronal density in the overlay of Allen and Franklin-Paxinos labels in solid and dotted lines, respectively. The full name of abbreviations can be found in Table S1.

Monosynaptic inputs to oxytocin neurons

Previous studies suggest that the PVH area receives input from specific brain areas such as the
dorsomedial hypothalamus area, the medial preoptic area (MPO), the subparafascicular area, and
the posterior intralaminar nuclei to integrate external stimuli and internal state (29–31). To map
brain-wide trans-synaptic inputs in a cell type specific manner, we injected monosynaptic
retrograde rabies viruses into the PVH and the SO of the *Ot-Cre* knock-in mice (32). We used
our qBrain mapping method to quantify input neurons throughout the whole brain (25). To
capture total input to each anatomical area, we combined input signals from multiple
independent injections targeting a specific brain region and overlaid them onto a reference brain

while displaying input neurons with pseudo-colored green for the PVH (n=6 brains) and red for
the SO (n=4 brains) (Fig. 2A, Movie S2). Our analysis reveals that more than 50 different
structures provide inputs to OT neurons (Fig. 2B). Both the PVH and the SO receive major
inputs from other hypothalamic brain regions. Overall, PVH OT neurons take inputs from more
regions than the SO neurons. Notably, PVH and SO OT neurons have different regional input
preferences (Fig. 2B). For example, thalamic regions mainly provide inputs to the PVH neurons
(Fig. 2A, blue arrowhead), while the striatum (STR) areas (e.g., nucleus of accumbens) provide
more input to SO neurons (Fig. 2A, yellow arrowhead). Moreover, the ventral anterior-lateral
complex of the thalamus, piriform cortex, and anterior amygdalar area only provide inputs to the
SO neurons. Both the PVH and SO received inputs from the midbrain reticular nucleus (MRN)
and the periaqueductal gray (PAG) in the midbrain (MB). The motor-related superior colliculus
provides inputs only to the PVH neurons, while the substantia nigra reticular part provides input
only to the SO neurons. Notably, PVH OT neurons are strongly connected with the thalamic
brain area, which communicates with the cortex and is a key structure for the rapid integration of
new learning, working memory, and adaptive decision-making (25), while the SO receives more
input from the striatum-like amygdala, which is centered on emotion-based cognitive function
(26). This input pattern is overall in agreement with prior reports that utilize traditional
retrograde tracers or viral tools (10, 30). The anterior two-thirds of the PVH is known to contain
the majority of magnocellular neuroendocrine neurons, while the posterior one-third of the PVH
contains the descending preautonomic neurons (8). To investigate potentially segregating input
patterns into different PVH subregions, we performed a series of injections with a small volume
(150 nL) of the rabies virus into the anterior-posterior PVH area. However, we did not observe a
topographically distinct input pattern into different PVH subregions (Fig. S3). Collectively, we
conclude that OT neurons in the PVH received stronger and broader inputs than the SO with
distinctive regional input preferences (Fig. 2C).

Figure 2. Monosynaptic input map of oxytocin neurons in the PVH and the SO.
(A) Inputs into the PVH (green) and SO (red) OT neurons in representative coronal planes from the Allen CCF. The signals were overlaid on the reference brain. See also Movie S2. Blue and yellow arrowheads indicate unique inputs from the thalamus and the ventral striatum into the PVH and the SO, respectively. Scale bar = 1mm. (B) Number of cells across anatomical regions providing synaptic inputs into OT neurons in the PVH (green bar) and SO (red bar). The input regions from both PVH and SO were represented as gray. Neighboring brain regions to the PVH and SO with potential signal contamination are not included in the graph, indicated with arrowhead. For example, red arrowhead in the medial amygdala means that input data from the SO was excluded. (C) Schematic input map in the mouse brain. Anatomical brain regions were segmented with different color code and arrows represent input brain regions (green: PVH and red: SO neurons). The full name of abbreviations can be found in Table S2.

Anterograde projections of oxytocin neurons

OT neurons can secrete OT through axonal and dendritic processes in coordinated or
independent compartment-specific manner (33, 34). To visualize the entire central projection
patterns of OT neurons, we injected Cre-dependent adeno associated virus 2 (AAV2-CAG-Flex-
EGFP) in multiple subregions of the PVH and the SO of *Ot-Cre* knock-in mice. Long-range
projection signals from multiple injections were registered onto a reference brain and merged to
represent efferent output from each anatomical area (Fig 3A, Movie S3). First, we observed that
OT neurons in the PVH project to the HY, the TH, the STR, and the pallidum (PAL), as well as
the posterior region including the MB and the hindbrain (HB) (Fig 3A-B). In contrast, the SO OT
neurons project to very limited areas (e.g., pons) in the brain (Fig 3A-B). Quantitatively, OT
neurons in the PVH project to over 99 brain regions including sparse projection to forebrain
areas (N = 3 males and 3 females, Fig 3B, Table S3). OT neurons in the SO project mainly to
hypothalamic regions and sparsely to posterior brain regions including the PB and the superior
olivary complex (SOC) (Fig. 3A-B and Table S3). To examine potential projection topology
within the PVH, we targeted PVH subregions by injecting a small volume (40~150 nL) of the
AAV virus injection along different A/P locations. We did not observe a strong regional
projectome difference within the PVH (Fig. S4).

Comparison of input-output connection of oxytocin neurons

Next, we systematically compared brain regions connected with OT neurons in the PVH and the
SO either unidirectionally or bidirectionally (Fig. 3D-E). In the PVH, selected hypothalamic
areas (e.g. the TU, the lateral hypothalamic area, and the arcuate hypothalamic nucleus) and the
MB (e.g. the PAG) show reciprocal input-output connection. Moreover, the thalamic area (e.g.
the nucleus of reuniens, the parataenial nucleus, the ventral medial nucleus of the thalamus, and
the reticular nucleus of the thalamus) has strong input to the PVH OT neurons while the MRN
shows a stronger output pattern (Fig. 3D). In the SO, OT neurons have dominant input from the
PVH and reciprocal connection with the TU (Fig. 3E). To further understand the input-output
relationship quantitatively, we calculated relative input-output strength within the brain regions
bidirectionally connected with OT neurons and plotted a log scale of relative input-output ratio
(Fig. 3F-G). For example, log-ratio '0' means that the relative strength of input and the output to
that specific region is similar. In PVH OT neurons, many thalamic and hypothalamic regions
show input preference and cerebral nuclei have balanced input-output log-ratios while MB
regions have output preference (Fig. 3F). In contrast, the SO OT neurons show stronger input
preference largely due to their limited projection areas (Fig. 3G). These input-output patterns
indicate that after integrating cognitive and physiological information from relatively closer
regions (e.g., the TH, the HY), PVH OT neurons broadly project to other brain regions from the
forebrain to the brain stem to exert fast and site-specific neuronal modulation (Fig. 3F). In
contrast, OT neurons in the SO mainly receive emotion-based cognitive information from
cerebral nuclei (e.g., the PAL) and social-sexual information from hypothalamic area while
mainly providing limited output to the HY and posterior pituitary to regulate metabolism and
autonomic body functions (Fig. 3G).

Figure 3. Projection map of Oxytocin neurons in the PVH and the SO.
(A) Projection outputs from the PVH (green) and SO (red) oxytocin neurons across representative coronal planes from the Allen CCF. Scale bar = 1mm. See also Movie S3. (B) Area normalized projection (percentage of the area with the signal/total area) of brain regions with more than 0.1% of the area filled with projection signal originated from the PVH (green), the SO (red), or both (grey). Neighboring brain regions to the PVH and SO with potential signal contamination are not included in the graph, indicated with arrowhead. (C) Schematic output map in the mouse brain. Anatomical brain regions were segmented with different color code and arrows represent output projection (green: PVH and red: SO OT neurons). (D-E) Scatter plots of brain regions with synaptic input and axonal output from the PVH (D) and the SO OT neurons (E). (F-G) Comparison of the relative output and input strength of the PVH (F) and the SO OT neurons (G). The full name of abbreviations can be found in Table S3.

Comparison of oxytocin output and oxytocin receptor expression

The actions of OT are transduced by a single subtype of the OTR, a seven-transmembrane protein (1). Previous studies suggested a mismatch between presynaptic OT projection and postsynaptic OTR expression (3, 34), yet quantitative brain-wide comparison has not been conducted. Previously, we validated that OTR-Venus mice represent the endogenous OTR expression faithfully (11). We imaged a cohort of adult OTR-Venus mice using serial two-photon tomography and mapped OTR expression in the whole adult brain (n=14, green in Fig.

4A). Then, we compared the combined projection pattern of *Ot-Cre* mice in the PVH and the SO
(Fig. 3A-C) and OTR-Venus (+) cell density across the entire brain via image registration onto
the reference brain (Fig. 4, Table S4, Movie S4). Overall, the OTR (pseudo colored as green)
showed high expression in the cortical area with minimal OT projection, while many midbrain
and hindbrain regions (e.g., nucleus of the lateral lemniscus) have strong OT projection (pseudo
colored as red) with little OTR expression (Fig 4A-C; Movie S4). Correlation analysis between
OT and OTR expression across the whole brain did not result in any significant correlation (Fig
4D). Nevertheless, a few brain areas contained both OT projection and OTR expression
including the lateral septal nucleus, the central amygdalar nucleus (CEA), the paraventricular
nucleus of the thalamus (PVT), the PAG, the SOC, and the facial motor nucleus (Fig. 4B).
Moreover, a significant OT-OTR correlation was observed within the thalamic and medullary
areas. (Figs. 4D-4F). Thus, our results highlight quantitative and spatial discrepancies between
OT and OTR in the mouse brain, suggesting largely indirect OT transmission in the brain.

Figure 4. Comparison between the oxytocin output and oxytocin receptor expression.
(A) Composite images of representative OT neuronal projection outputs (red: combined from both the PVH and the SO) and OTR expression (green) across the mouse brain. See also Movie S4. (B) Representative images to show the co-expression of oxytocinergic fibers and oxytocin receptors. White lines represent anatomical borders from the Allen CCF. Scale bars in the column 1 = 1 mm and the column 2-4 = 500 µm. (C) Quantitative comparison of relative OT projection pattern and OTR expression. Note that cerebral cortex has very small OT/OTR ratio while the hindbrain and the midbrain shows higher ratio. (D) Correlation between OT projection and oxytocin receptor density (Spearman nonparametric correlation, *: p<0.05). Note the significant correlation in the thalamus and the medulla. (E-F) Correlation of brain regions with axonal projection and OTR expression in the thalamus (E) and the medulla (F). (G) Schematic summary of synaptic input and axonal output connectivity of OT neurons and correlated OTR in the mouse brain (green: PVH and red: SO OT neurons). All abbreviations for brain regions can be found in Table S4.

Discussion

The wiring diagram of the brain is a structural foundation to decipher neural circuits underlying brain function. Here, we present a comprehensive anatomical and connectivity map of the hypothalamic OT neurons and their relationship with postsynaptic OTR expression in the whole mouse brain. We find that OT neurons broadly project to brain regions from the forebrain to the brain stem, while the majority of inputs to OT neurons comes from the thalamus and the hypothalamus. Moreover, we confirm the quantitative mismatch between OT projection and OTR distribution in the brain.

OT neurons are mostly located in hypothalamic nuclei with a complex 3D shape (8). To examine OT expression intuitively and quantitatively, we devised a 2D flatmap for OT containing hypothalamic regions from an Allen CCF based reference brain while incorporating anatomical labels from the Allen Institute and Franklin-Paxinos (23, 27). This approach allows for the interpretation of OT anatomical location from two independently created and commonly used atlases (28) and provides an alternative coordinate system to understand anatomical connectivity. Our transgenic labeling approach using OT:Ai14 mice revealed that OT neurons in PVH largely maintain their expression throughout adulthood whereas other brain regions (e.g., SO, MEA) showed significant down-regulation of OT expression. Considering the significant role OT signaling plays in shaping the brain development including neural connectivity (35), this transient OT expression in non-PVH areas may provide extra OT signaling to promote early brain development.

Our OT input-output wiring diagram provides brain-wide insights about connectivity-based brain function, which can be divided into two groups (Fig. 4G). The first group contains the brain regions with unidirectional OT output. This pattern is mainly observed in the midbrain, pons, and medulla which are known to be associated with physiological function including body metabolism, sleep, and motor sensory control. For example, the nucleus of the solitary tract (NTS) in the medulla is a part of the neural circuit for food intake and energy expenditure (36). OT neurons in the PVH directly communicate with the NTS to modulate visceral afferent transmission (37). The second group contains brain regions with reciprocal input-output connections with OT neurons including the hypothalamus, thalamus, and striatum that regulate cognitive brain functions such as sexual behavior, memory, attention, fear, reward, and pain. For instance, reciprocally connected regions in the hypothalamic area are highly related to the parental and sexual behaviors such as the MPO (38). Although OT neurons can be further divided into magno- and parvo-cellular neurons, a recent study in rats showed that both cell types in the PVH receive overall similar inputs despite small differences (10). Moreover, we did not observe topographically distinct subregions within the PVH based on input-output maps. This is consistent with a previous retrograde mapping study from the spinal cord, showing that spinal cord projecting OT neurons are distributed across the entire A/P axis (8).

Our data also provides a brain-wide perspective on the spatial relationships between OT central projections and oxytocin receptor expression (Fig. 4G). Although the spatial discrepancy of OT-OTR was noted before, recent studies showed that most OTR expressing areas contain at least sparse OT projection (12, 18). Nonetheless, our analysis showed that there is overall no significant quantitative correlation between OT and OTR across entire brain regions. For example, the cerebral cortex area contains abundant OTR with little to no OT axons. However, OT can still mediate sensory stimuli in the cortex to modify mouse behavior (39). Previous

studies suggest that OTR neurons in the isocortex may receive OT signals indirectly from
ventricular pathways via cerebral spinal fluid with delayed and long-lasting effects (40, 41). In
addition, OT axonal ends in the CEA exert a direct effect on fear suppression with delayed
electrophysiological responses (~seconds) in the CEA, suggesting that OT is being released in a
non-synaptic way (18, 42). In contrast, we also found few areas with high levels of both OTR
and OT projection such as the PVT which is involved in maternal behavior (43) and the
ventrolateral periaqueductal gray that is implicated in for social defeat stress (44). OT signaling
in these areas is likely to occur as the spatially and temporally precise synaptic transmission.
Another noteworthy OT-OTR discrepancy is brain regions with abundant OT projection without
OTR expression such as sensory related hindbrain areas. Although OTR is a main OT receptor,
OT can bind to another receptor to exert its effect. For example, OT can elicit TRPV1 activity in
the spinal cord to modulate nociception (45). Non-canonical pathways like this are an under
explored mechanism of OT modulation in the central nervous system that requires further study.

In summary, our study provides an anatomical foundation to understand diverse functions based
on OT neurons in the brain. We deposit all high-resolution imaging data in publicly accessible
databases and our website to facilitate data mining. We envision that this OT wiring diagram
with quantitative expression data will guide future studies to understand circuit-based
mechanisms of OT function and its changes in various brain disorders such as autism.

289
290

Material and Method

<u>Animals</u>

All animal care and experimental procedures are approved by the Penn State University Institutional Animal Care Use Committee (IACUC). *Ot-Cre* mice (24) were originally produced in the Gloria B. Choi lab at the Massachusetts Institute of Technology and imported to the Penn State University (Kim Lab). To generate OT:Ai14 mice, *Ot-Cre* mice were crossed with Ai14 mice, expressing tdTomato following Cre-mediated recombination (Jax: 007914, C57Bl/6 J background). Mice received food and water ad libitum and were housed under constant temperature and light conditions (12 hrs light and 12 hrs dark cycle).

<u>Stereotaxic surgery and virus injections</u>

Ot-Cre mice (8-11 weeks old, males and females) were anesthetized with isoflurane (controlled with Somnosuite, Kent Scientific) and mounted on a stereotaxic instrument (Angle Two, Leica) with a heating pad placed underneath. All injections were performed with pulled micropipettes (VWR, Cat# 53432-706). Through the small opening of the micropipette, virus was delivered at a rate of 75-100 nL per minute. The speed and volume of injection were monitored along with the calibration marks on the micropipette (1 mm =100 nL). To target the PVH, coordinates are anteroposterior (AP) from the Bregma: -0.58 mm; mediolateral (ML): 0.27 mm; dorsoventral (DV): -4.75 mm. Anterior PVH and posterior PVH injection coordinates are -0.35 mm (AP), 0.3 mm (ML), and -4.5 mm (DV) and -0.94 mm (AP), 0.39 mm (ML), and -4.55 mm (DV), respectively. Coordinates for the SO are -0.66 mm (AP), 1.3 mm (ML), and -5.8 mm (DV). For anterograde tracing, 50-500 nL of AAV2-CAG-Flex-EGFP virus (titer 3.7×10^{12} vg/ml, purchased from UNC vector core) was injected into the PVH (500 nL for maximum coverage, 50-150 nL for PVH subregion) and 150 nL of the virus was injected into the SO. Mice were euthanized three weeks later with Ketamine (100 mg/kg) and Xylazine (10 mg/kg) mixture. For monosynaptic retrograde labeling, 50-500 nL of rAAV1-svnp-DIO-STPEPB (titer 3.9×10^{12} cg/ml, purchased from UNC vector core, a gift from Ian Wickersham (46), Addgene plasmid # 52473 ; http://n2t.net/addgene:52473 ; RRID:Addgene_52473) was injected into the PVH, followed by the same quantity of EnvA G-deleted Rabies-mcherry virus (titer: 8.12×10^{8} transduction unit (TU) /ml, purchased from the Salk Institute Viral Vector Core, a gift from Edward Callaway (47), Addgene plasmid # 32636 ; http://n2t.net/addgene:32636 ; RRID:Addgene_32636) three weeks later into the same location. Mice were euthanized 7-8 days later with Ketamine (100 mg/kg) and Xylazine (10 mg/kg) mixture.

<u>STPT imaging and related data analysis</u>

Transgenic or virus injected mice were transcardially perfused with 4% paraformaldehyde (PFA) in 0.1M phosphate buffer (PB, pH 7.4) after 0.9% saline. Brains were dissected out and post-fixed in 4% PFA overnight at 4°C. Fixed brains were stored in 0.05 M phosphate buffer at 4°C until imaged. To image the entire brain, a serial two-photon tomography (TissueCyte 1000; Tissuevision) was used as previously described (25, 48). Briefly, the brain was embedded in 4% oxidized agarose and cross-linked with 0.2% sodium borohydride solution. The brain was imaged as 12 x 16 x 280 tiles with 1 x 1 μm^2 *x,y* pixel resolution in every 50 μm *z*-section. We used 910 nm wavelength for two-photon excitation to excite both green (e.g., eGFP) and red

signals (e.g., tdTomato). Signals were separated with 560 nm dichroic mirror and two band path
filters (607/70-25 for red and 520/35- 25 for green). Imaging tiles in each channel were stitched
with custom-built software (25).

For quantitative projection data analysis, we used our previously published pipeline (49). Briefly,
both signal and background channels were z-normalized. Then, the background channel images
were subtracted from the signal channel images to increase signal-to-noise ratio. Then, projection
signals were converted to a binary map by applying an optimized threshold (8x standard
deviation) to detect signals while minimizing noise from background autofluorescence. Then,
binarized signals in each pixel were counted in 20 x 20 (x,y) pixel unit (voxel) and the value was
assigned the corresponding voxel across the brain, which is defined as "projection strength".
Thus, range of the projection strength in a given voxel is between 0 and 400. Projection strength
of each area is calculated by summing up all projection strength within an anatomically defined
area. Autofluorescence of brains was used to register each brain to the Allen CCF using Elastix
(50), then, the projection signals were transformed to the reference brain. "Area normalized
projection" represents normalized occupancy of projection signals in the ROI by dividing the
projection strength with a total number of voxels in each ROI. For example, if total voxel count
for one ROI was 20,000 and our projection strength showed 2,000 in the ROI, it will be
(2,000/20,000)*100 = 10%. For cell counting analysis, we used a machine-learning algorithm to
detect fluorescently labeled cells (25). The cell density in 2D (counts/mm^2) was calculated by
dividing cell number with ROI area. 2D counting numbers were also converted into the 3D
counting using our previously calculated 3D conversion factor (1.4 for tdTomato) (25). To
measure the volume of anatomical ROI, the reference Allen CCF was reversely registered onto
individual brains using the Elastix. "Cell density (counts/mm^3)" was calculated by dividing
detected cell numbers in 3D with the anatomical ROI volume. The cell counting analysis was
applied to OT:Ai14 and OTR-Venus cell distribution and inputs to the OT neurons.

To compare relative quantity of OT input-output (Fig. 3F-G), bidirectionally connected areas
were first selected. Within the selected areas, relative cell density or output data in each region
was calculated by dividing each data by summed density or output data from all chosen areas,
respectively. Then, log10 (relative input/relative output) was used to quantify input or output
preference of each area. The same approach was used to calculate relative abundance between
OT output and OTR expression in Fig. 4C.

To create a representative connectivity map for each anatomical area, we microinjected OT
neurons with a large quantity of virus into subregions within an anatomically defined area. Then,
the maximum signal of several (4-6) brains from each ROI registered in the reference was used
for final analysis to cover entire target anatomical areas for OT input and output data in Figures
2-4, and Movies S2-S4.

To determine the correlation between OT area normalized projection and OTR density, we first
tested for the normality of the data using the D'Agostino-Pearson normality test. Based on the
normality test result, we performed Spearman nonparametric correlation test. GraphPad Prism 8
was used for all statistical analysis and graphs.

2D hypothalamic and PVH Flatmap

To generate the hypothalamic flatmap, we adapted the previously used method (25) and applied
it to the hypothalamic region. First, we created a binary image in the hypothalamic area based on
the oxytocin expression. Second, a zero line was placed to generate evenly spaced bins along the
dorsal to the ventral direction of the PVH and laterally extended to include TU and MEA at
different coronal plains. To capture signals on the flatmap, bins were registered into the reference
brain and the cell number in each bin was quantified as described before in the STPT data
analyses section. Lastly, the mean number of the OT neurons in 8 OT:Ai14 brains were plotted
in each flatmap using a custom-build matlab code. For the PVH flatmap, we followed the same
procedure to generate a hypothalamic flatmap except for bin generation. Instead of delineating
bins in a binary image, we assigned bin numbers in the PVH subregion of Franklin-Paxinos atlas
(27) along the dorsal to the ventral direction.

Immunohistochemistry, microscopic image, and cell counting

For immunohistochemistry, fixed brains were either embedded in 3% agarose or frozen after
sinking in 30% sucrose in 0.2 M Phosphate buffer. Embedded or frozen brains were then cut on a
vibratome (Leica vt1000s) or a microtome (Leica SM2010 R) at 50 μm thickness. Sections were
stored at −20°C in a cryoprotectant solution (30% sucrose and 30% glycerol in 0.1 M PB) until
immunostaining. For oxytocin staining, sections were washed three times in 1x PBS. After 1
hour incubation in blocking solution (10% donkey serum and 0.1 % Triton X-100), slices were
incubated with oxytocin primary antibody (ImmunoStar Cat# 20068, RRID:AB_572258, 1:1000)
in blocking solution for overnight at 4 °C. Sections were then washed three times with 1x PBS
and further incubated in secondary antibodies (Thermo Fisher Scientific Cat# A-21206,
RRID:AB_2535792, 1:500) for 1 hour at room temperature. After washing three times, slices
were mounted onto slides and coverslipped with vectashield mounting media (Vector
laboratories, H-1500-10). For microscopic imaging, a BZ-X700 fluorescence microscope
(Keyence) and a confocal microscope (Zeiss 510) were used. A low magnification objective lens
(4x) was used to image with a large enough view to define brain anterior-posterior location from
bregma and higher magnification objective lenses (10x ~ 40x) were used to image sections
depending on the cell density. Images were delineated manually based on the Franklin-Paxinos
atlas and fluorescently tagged cells were manually quantified using the cell counter plug-in in
FIJI (ImageJ, NIH).

Acknowledgments

This publication was made possible by an NIH grant R01MH116176 to Y.K. and an NIH grant R24OD18559 to K.C. Its contents are solely the responsibility of the authors and do not necessarily represent the views of the funding agency. We thank Dr. Gloria Choi for kindly sharing OT-Cre transgenic mice and Rebecca Betty for assistance in editing the manuscript. We acknowledge use of computational resources in the High Performance Computing cluster at the Penn State College of Medicine.

Contributions

Conceptualization, Y.K.; Data Collection and analysis, S.S., ; Data Collection, S.M., K.N., M.C., Computer Coding, Y.W.; Web visualization, D.J.V., K.C.; Manuscript preparation: S.S., Y.K with help from the other authors.

Competing Interests

The authors declare no competing interests.

Data Sharing Plan

High-resolution serial two-photon tomography images can be found at https://kimlab.io/brain-map/ot_wiring/

Flatmaps and custom-built codes are available at https://kimlab.io/ and can be used without any restriction.

References

1. B. Jurek, I. D. Neumann, The Oxytocin Receptor: From Intracellular Signaling to Behavior. *Physiol. Rev.* **98**, 1805–1908 (2018).

2. V. Grinevich, R. Stoop, Interplay between Oxytocin and Sensory Systems in the Orchestration of Socio-Emotional Behaviors. *Neuron* **99**, 887–904 (2018).

3. H.-J. Lee, A. H. Macbeth, J. H. Pagani, W. S. Young, Oxytocin: the great facilitator of life. *Prog. Neurobiol.* **88**, 127–151 (2009).

4. D. S. Quintana, A. J. Guastella, An Allostatic Theory of Oxytocin. *Trends Cogn. Sci.* (2020) https:/doi.org/10.1016/j.tics.2020.03.008.

5. S. M. Francis, *et al.*, Oxytocin and vasopressin systems in genetic syndromes and neurodevelopmental disorders. *Brain Res.* **1580**, 199–218 (2014).

6. K. T. Rajamani, S. Wagner, V. Grinevich, H. Harony-Nicolas, Oxytocin as a Modulator of Synaptic Plasticity: Implications for Neurodevelopmental Disorders. *Front. Synaptic Neurosci.* **10** (2018).

7. A. J. Guastella, I. B. Hickie, Oxytocin Treatment, Circuitry and Autism: A Critical Review of the Literature Placing Oxytocin into the Autism Context. *Biol. Psychiatry* (2015).

8. J. Biag, *et al.*, Cyto- and chemoarchitecture of the hypothalamic paraventricular nucleus in the C57BL/6J male mouse: A study of immunostaining and multiple fluorescent tract tracing. *J. Comp. Neurol.* **520**, 6–33 (2012).

9. V. Grinevich, I. D. Neumann, Brain oxytocin: how puzzle stones from animal studies translate into psychiatry. *Mol. Psychiatry*, 1–15 (2020).

10. Y. Tang, *et al.*, Social touch promotes interfemale communication via activation of parvocellular oxytocin neurons. *Nat. Neurosci.* (2020) https:/doi.org/10.1038/s41593-020-0674-y.

11. K. T. Newmaster, *et al.*, Quantitative cellular-resolution map of the oxytocin receptor in postnatally developing mouse brains. *Nat. Commun.* **11**, 1885 (2020).

12. V. Grinevich, H. S. Knobloch-Bollmann, M. Eliava, M. Busnelli, B. Chini, Assembling the Puzzle: Pathways of Oxytocin Signaling in the Brain. *Biol. Psychiatry* **79**, 155–164 (2016).

13. G. Gimpl, F. Fahrenholz, The oxytocin receptor system: structure, function, and regulation. *Physiol. Rev.* **81**, 629–683 (2001).

14. K. Li, M. Nakajima, I. Ibañez-Tallon, N. Heintz, A Cortical Circuit for Sexually Dimorphic Oxytocin-Dependent Anxiety Behaviors. *Cell* **167**, 60-72.e11 (2016).

15. S. Sabihi, S. M. Dong, N. E. Durosko, B. Leuner, Oxytocin in the medial prefrontal cortex regulates maternal care, maternal aggression and anxiety during the postpartum period. *Front. Behav. Neurosci.* **8**, 258 (2014).

16. N. N. Tirko, *et al.*, Oxytocin Transforms Firing Mode of CA2 Hippocampal Neurons.

Neuron **100**, 593-608.e3 (2018).

17.   T. Raam, K. M. McAvoy, A. Besnard, A. H. Veenema, A. Sahay, Hippocampal oxytocin
receptors are necessary for discrimination of social stimuli. *Nat. Commun.* **8**, 2001 (2017).

18.   H. S. Knobloch, *et al.*, Evoked axonal oxytocin release in the central amygdala attenuates
fear response. *Neuron* **73**, 553–566 (2012).

19.   V. Ferretti, *et al.*, Oxytocin Signaling in the Central Amygdala Modulates Emotion
Discrimination in Mice. *Curr. Biol.* **29**, 1938--1953.e6 (2019).

20.   P. J. Ryan, S. I. Ross, C. A. Campos, V. A. Derkach, R. D. Palmiter, Oxytocin-receptor-
expressing neurons in the parabrachial nucleus regulate fluid intake. *Nat. Neurosci.* **20**,
1722–1733 (2017).

21.   M. Eliava, *et al.*, A New Population of Parvocellular Oxytocin Neurons Controlling
Magnocellular Neuron Activity and Inflammatory Pain Processing. *Neuron* **89**, 1291–
1304 (2016).

22.   S. Boll, A. C. Almeida de Minas, A. Raftogianni, S. C. Herpertz, V. Grinevich, Oxytocin
and Pain Perception: From Animal Models to Human Research. *Neuroscience* **387**, 149–
161 (2018).

23.   Q. Wang, *et al.*, The Allen Mouse Brain Common Coordinate Framework: A 3D
Reference Atlas. *Cell* **181**, 936-953.e20 (2020).

24.   H. K. Choe, *et al.*, Oxytocin Mediates Entrainment of Sensory Stimuli to Social Cues of
Opposing Valence. *Neuron* **87**, 152–163 (2015).

25.   Y. Kim, *et al.*, Brain-wide Maps Reveal Stereotyped Cell-Type-Based Cortical
Architecture and Subcortical Sexual Dimorphism. *Cell* **171**, 456-469.e22 (2017).

26.   M. P. Madrigal Verdú, S. Jurado, Specification of oxytocinergic and vasopressinergic
circuits in the developing mouse brain. *bioRxiv*, 2020.08.07.241364 (2020).

27.   G. Franklin, Keith and Paxinos, *The Mouse Brain in Stereotaxic Coordinates* (Academic
Press, 2008).

28.   U. Chon, D. J. Vanselow, K. C. Cheng, Y. Kim, Enhanced and unified anatomical labeling
for a common mouse brain atlas. *Nat. Commun.* **10**, 5067 (2019).

29.   S. Campeau, S. J. J. Watson, Connections of some auditory-responsive posterior thalamic
nuclei putatively involved in activation of the hypothalamo-pituitary-adrenocortical axis
in response to audiogenic stress in rats: an anterograde and retrograde tract tracing study
combined with . *J. Comp. Neurol.* **423**, 474–491 (2000).

30.   M. Cservenák, *et al.*, A Thalamo-Hypothalamic Pathway That Activates Oxytocin
Neurons in Social Contexts in Female Rats. *Endocrinology* **158**, 335–348 (2017).

31.   C. Boudaba, K. Szabó, J. G. Tasker, Physiological mapping of local inhibitory inputs to
the hypothalamic paraventricular nucleus. *J. Neurosci.* **16**, 7151–7160 (1996).

32.   I. R. Wickersham, *et al.*, Monosynaptic restriction of transsynaptic tracing from single,
genetically targeted neurons. *Neuron* **53**, 639–647 (2007).

33.   M. Ludwig, G. Leng, Dendritic peptide release and peptide-dependent behaviours. *Nat.*

Rev. Neurosci. **7**, 126–136 (2006).

34.   B. Jurek, I. D. Neumann, The Oxytocin Receptor: From Intracellular Signaling to
Behavior. *Physiol. Rev.* **98**, 1805–1908 (2018).

35.   T. V Miller, H. K. Caldwell, Oxytocin during Development: Possible Organizational
Effects on Behavior. *Front. Endocrinol. (Lausanne).* **6**, 76 (2015).

36.   R. A. Travagli, L. Anselmi, Vagal neurocircuitry and its influence on gastric motility. *Nat.*
Rev. Gastroenterol. Hepatol. **13**, 389–401 (2016).

37.   J. H. Peters, *et al.*, Oxytocin enhances cranial visceral afferent synaptic transmission to the
solitary tract nucleus. *J. Neurosci.* **28**, 11731–11740 (2008).

38.   M. Gil, R. Bhatt, K. B. Picotte, E. M. Hull, Oxytocin in the medial preoptic area facilitates
male sexual behavior in the rat. *Horm. Behav.* **59**, 435–443 (2011).

39.   B. J. Marlin, M. Mitre, J. A. D'amour, M. V Chao, R. C. Froemke, Oxytocin enables
maternal behaviour by balancing cortical inhibition. *Nature* **520**, 499–504 (2015).

40.   J. J. Zheng, *et al.*, Oxytocin mediates early experience-dependent cross-modal plasticity in
the sensory cortices. *Nat. Neurosci.* **17**, 391–399 (2014).

41.   W. B. Mens, A. Witter, T. B. van Wimersma Greidanus, Penetration of neurohypophyseal
hormones from plasma into cerebrospinal fluid (CSF): half-times of disappearance of
these neuropeptides from CSF. *Brain Res.* **262**, 143–149 (1983).

42.   H. S. Knobloch, V. Grinevich, Evolution of oxytocin pathways in the brain of vertebrates.
Front. Behav. Neurosci. **8**, 31 (2014).

43.   A. Watarai, *et al.*, The blockade of oxytocin receptors in the paraventricular thalamus
reduces maternal crouching behavior over pups in lactating mice. *Neurosci. Lett.* **720**,
134761 (2020).

44.   N. Nasanbuyan, *et al.*, Oxytocin-Oxytocin Receptor Systems Facilitate Social Defeat
Posture in Male Mice. *Endocrinology* **159**, 763–775 (2018).

45.   Y. Nersesyan, *et al.*, Oxytocin Modulates Nociception as an Agonist of Pain-Sensing
TRPV1. *Cell Rep.* **21**, 1681–1691 (2017).

46.   K. Kohara, *et al.*, Cell type-specific genetic and optogenetic tools reveal hippocampal
CA2 circuits. *Nat. Neurosci.* **17**, 269–279 (2014).

47.   F. Osakada, *et al.*, New rabies virus variants for monitoring and manipulating activity and
gene expression in defined neural circuits. *Neuron* **71**, 617–631 (2011).

48.   T. Ragan, *et al.*, Serial two-photon tomography for automated ex vivo mouse brain
imaging. *Nat. Methods* **9**, 255–258 (2012).

49.   M. Jeong, *et al.*, Comparative three-dimensional connectome map of motor cortical
projections in the mouse brain. *Sci. Rep.* **6**, 20072 (2016).

50.   S. Klein, M. Staring, K. Murphy, M. A. Viergever, J. P. W. Pluim, elastix: a toolbox for
intensity-based medical image registration. *IEEE Trans. Med. Imaging* **29**, 196–205
(2010).

Supplementary Information
553
554   **Wiring diagram of the oxytocin system in the mouse brain**
555
556   Son et al.,
557
558

560
561
562 Figure S1. Fluorescent images across 5 levels of the PVH, SO, and MEA. Genetically expressed
563 oxytocin neurons (OT:Ai14) are red and oxytocin immuno staining cellar are labeled with green
564 fluorescent marker are green. Scale bar = 50 µm.
565

566
567
568
569 Figure S2. (A) 2D PVH flatmap. Small inserts with coronal sections illustrate the binning
570 system. Bin numbers were assigned on the subregion of PVH at different coronal planes in
571 Franklin-Paxinos labels. (B) Heatmap of oxytocin neuronal density in the 2D PVH flatmap. The
572 number in each brain region represents the percentage of developmentally expressed neurons.
573 The X-axis is for bin numbers and Y-axis is for the bregma A/P axis.
574
575

577
578
579 Fig S3. PVH subregion infection. Left: infection coverage on the 2D hypothalamic flatmap.
580 Right: scatter plot between the virus injection coverage and the mean bregma A/P value of input
581 brain regions to the PVH.
582

583
584 Fig S4. PVH subregional topology. Left: infection area of five different injection on the 2D
585 hypothalamic flatmap. Right: scatter plot of the mean Bregma A/P value of output brain regions.
586
587
588

Supplementary Tables

Table S1. Oxytocin neuronal distribution in the mouse brain

Table S2. Number of cells in brain areas providing monosynaptic input to oxytocin neurons

Table S3. Projection percentage of brain regions with oxytocin neuronal projection

Table S4. Comparison between oxytocin projection and oxytocin receptor expression

Supplementary Movies

Movie S1. Oxytocin neuronal expression in the whole brain

Movie S2. Monosynaptic input to oxytocin neurons in the PVH and the SO

Movie S3. Brain-wide projection of oxytocin neurons in the PVH and the SO

Movie S4. Comparison between oxytocin receptor expression and projection of hypothalamic oxytocin neurons.