Cross-compartment signal propagation in the Mitotic Exit Network.

Xiaoxue Zhou[1,*], Wenxue Li[2], Yansheng Liu[2], Angelika Amon[1,*]

[1]David H. Koch Institute for Integrative Cancer Research, Howard Hughes Medical Institute, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

[2]Yale Cancer Biology Institute, Department of Pharmacology, Yale University, West Haven, CT 06516, USA

[*]Correspondence to: xiaoxue@mit.edu and angelika@mit.edu

ABSTRACT (150 words)

In budding yeast, the Mitotic Exit Network (MEN), a GTPase signaling cascade integrates spatial and temporal cues to promote exit from mitosis. This signal integration requires transmission of a signal generated on the cytoplasmic face of spindle pole bodies (SPBs; yeast equivalent of centrosomes) to the nucleolus, where the MEN effector protein Cdc14 resides. Here, we show that the MEN activating signal at SPBs is relayed to Cdc14 in the nucleolus through the dynamic localization of its terminal kinase complex Dbf2-Mob1. Cdc15, the protein kinase that activates Dbf2-Mob1 at SPBs, also regulates its nuclear access. Once in the nucleus, priming phosphorylation of Cfi1/Net1, the nucleolar anchor of Cdc14, by the Polo-like kinase Cdc5 targets Dbf2-Mob1 to the nucleolus. Nucleolar Dbf2-Mob1 then phosphorylates Cfi1/Net1 and Cdc14, activating Cdc14. The kinase-primed transmission of the MEN signal from the cytoplasm to the nucleolus exemplifies how signaling cascades can bridge distant inputs and responses.

INTRODUCTION

In cellular signaling, the sensing of signals (i.e. binding of signaling molecules at cell surface) and the response (i.e. transcription in the nucleus) often occur in different cellular compartments. Determining how signals are transmitted across compartments is thus essential for understanding signal transmission. The Mitotic Exit Network (MEN), a Ras-like GTPase kinase signaling cascade and budding yeast homolog of the Hippo pathway (Hergovich and Hemmings, 2012), represents such an example for signaling across cellular compartments. The MEN-activating signal is sensed and processed at the cytoplasmic face of spindle pole bodies (SPBs; yeast functional equivalent of the centrosomes), whereas the MEN effector protein Cdc14 resides in the nucleolus (Figure 1A). Because budding yeast undergo a closed mitosis without disassembling the nuclear envelope and nucleolus, the MEN must transmit a signal generated at the cytoplasmic face of SPBs, across the nuclear envelope and into the nucleolus to activate its effector Cdc14. The molecular mechanisms governing this cross-compartment signaling process remain largely unknown.

The central function of the MEN is to couple the final cell cycle transition, exit from mitosis (when the mitotic spindle is disassembled, chromosomes decondense and cytokinesis ensues), to nuclear/spindle position. In many organisms such as budding yeast, fission yeast as well as some plant species, the site of cytokinesis/division plane (i.e. the bud neck) is determined prior to mitosis (Guertin et al., 2002). Thus, the mitotic spindle must be positioned accordingly to ensure accurate genome partitioning between the daughter cells. In addition, these organisms have evolved surveillance mechanisms to monitor spindle position and regulate cell cycle progression in response. This surveillance mechanism is best understood in budding yeast where spindle position controls the activity of the MEN.

The MEN senses spindle position through a Ras-like GTPase Tem1. Tem1 is activated when a SPB enters the bud (Bardin et al., 2000; Pereira et al., 2000). Together with the Polo-like kinase Cdc5, Tem1 activates its effector, the Hippo-like protein kinase Cdc15, presumably by recruiting Cdc15 to the SPBs (Rock and Amon, 2011) (Figure 1A). Cdc15 then activates the LATS/NDR kinase Dbf2-Mob1 via a two-step process (Rock et al., 2013). Cdc15 first phosphorylates the MEN scaffold Nud1, a core component of the SPB outer plaque. This creates a docking site for Dbf2-Mob1 on Nud1 and facilitates subsequent phosphorylation of Dbf2-Mob1 by Cdc15 which activates Dbf2-Mob1 (Mah et al., 2001; Rock et al., 2013).

In addition to sensing spindle position, the MEN also integrates cues of cell cycle

progression through the downstream kinases Cdc15 and Dbf2-Mob1 (Campbell et al., 2019).

Two cell cycle events are sensed by the MEN: (1) Activity of the Polo-like kinase Cdc5:

activation of Cdc15 depends on Cdc5 activity (Rock and Amon, 2011) which occurs only in

mitosis (Cheng et al., 1998); (2) Initiation of anaphase: cyclin-dependent kinases (CDKs)

phosphorylate Cdc15 and Mob1 thereby inhibiting their activity (Jaspersen and Morgan, 2000;

König et al., 2010). At the onset of anaphase, CDK activity declines due to cyclin degradation,

which lifts this inhibition. This loss of inhibition by CDKs creates a state whereby the MEN is

poised for activation.

Once activated, the MEN promotes exit from mitosis by activating the phosphatase

Cdc14. As an antagonist of CDKs, Cdc14 reverses CDK-dependent phosphorylation and

promotes mitotic CDK inactivation thereby returning the cell to a G1 state (reviewed in

Stegmeier and Amon, 2004). Given the central role of Cdc14 in promoting mitotic exit, it is not

surprising that the phosphatase is tightly regulated. Cdc14 is sequestered in the nucleolus by its

inhibitor Cfi1/Net1 from G1 until the onset of anaphase (Shou et al., 1999; Visintin et al., 1999).

During anaphase, Cdc14 is released from its inhibitor, spreads throughout the nucleus and

cytoplasm to dephosphorylate its targets.

Two pathways, the Cdc fourteen early anaphase release (FEAR) network and the MEN

promote the dissociation of Cdc14 from its inhibitor during anaphase. Upon anaphase entry,

Cdc14 is transiently released from the nucleolus by the FEAR network (reviewed in Rock and

Amon, 2009). The FEAR network promotes dissociation of Cdc14 from Cfi1/Net1 through

facilitating phosphorylation of Cfi1/Net1 by mitotic CDKs (Azzam, 2004; Queralt et al., 2006).

This transient release, although not essential for exit from mitosis, is crucial for the timely

execution of several anaphase events such as segregation of the nucleolus (D'Amours et al.,

2004; Sullivan et al., 2004; Torres-Rosell et al., 2004) and MEN activation by counteracting

CDK inhibition of MEN kinases (Campbell et al., 2019; Jaspersen and Morgan, 2000; König et

al., 2010). In late anaphase, the activated MEN drives a more sustained and complete release of

Cdc14 from the nucleolus which ultimately results in exit from mitosis. In the absence of MEN

activity, Cdc14, after a transient FEAR network-mediated release from the nucleolus during

early anaphase, is re-sequestered in the nucleolus and cells arrest in late anaphase.

Despite our extensive knowledge of the MEN and Cdc14, how the MEN promotes the

sustained release of Cdc14 from its inhibitor in the nucleolus is not well understood. One

contributing mechanism involves the phosphorylation of the nuclear localization signal (NLS)

sequence in the C-terminus of Cdc14 by the MEN kinase Dbf2-Mob1 (Mohl et al., 2009).

Inactivation of the NLS by the MEN promotes redistribution of Cdc14 via its nuclear export

signal (NES) from the nucleus to the cytoplasm. However, retention of Cdc14 in the cytoplasm is

not required for mitotic exit (Bembenek et al., 2005; Kuilman et al., 2015; Mohl et al., 2009).

Furthermore, a Cdc14 mutant lacking the Dbf2 phosphorylation sites within its NLS is still

released from the nucleolus in late anaphase (Mohl et al., 2009). These results suggest that MEN-

mediated cytoplasmic retention of Cdc14 is not the main mechanism whereby the MEN activates

Cdc14. Rather, the MEN must also disrupt the interaction between Cdc14 and its inhibitor

Cfi1/Net1 in the nucleolus.

How does the MEN, activated at the outer plaque of SPBs in the cytosol, liberate Cdc14

from its inhibitor in the nucleolus? Here, we demonstrate that the terminal kinase in the MEN,

Dbf2-Mob1, serves as the molecular messenger traveling between the SPBs and the nucleolus to

release Cdc14 from its inhibitor. We show that Dbf2-Mob1, normally kept out of the nucleus by

Crm1, gains access to the nucleus following activation by Cdc15. We further demonstrate that

Dbf2-Mob1 utilizes a nucleolar docking site created by the Polo-like kinase Cdc5 in order to

phosphorylate Cfi1/Net1, resulting in Cdc14 liberation. These findings define the molecular

mechanisms of cross-compartment signal transmission in the MEN and provide a novel

paradigm for how signaling can occur across organelle boundaries.

RESULTS

Dbf2-Mob1 dynamically associates with SPBs.

When the MEN is activated in anaphase, Dbf2-Mob1 is recruited to the outer plaque of SPBs by

binding to Cdc15-phosphorylated Nud1 (Rock et al., 2013). However, immobilizing Dbf2-Mob1

at SPBs by fusing Mob1 to Nud1 disrupts MEN activity (Rock et al., 2013), suggesting that

Dbf2-Mob1 is likely needed away from the SPBs for the MEN to function. Additionally, a small

fraction of Dbf2-Mob1 was found to enter mitotic nuclei (Stoepel et al., 2005). We thus

hypothesized that the MEN liberates Cdc14 from its nucleolar inhibitor through the dynamic

shuttling of Dbf2-Mob1 between the outer plaque of the SPB and the nucleolus. We reasoned

that as a messenger between the SPB and nucleolus, Dbf2-Mob1 needs to be mobile at the SPB.

To test this, we performed fluorescence recovery after photobleaching (FRAP) analysis on eGFP

tagged Mob1 in anaphase cells (Figure 1B). We observed a rapid recovery of fluorescence with a

half-recovery time of 4.6 ± 0.7 s (mean $\pm$ std, $n = 6$ cells) after photobleaching of Mob1-eGFP

fluorescence either at the daughter (dSPB) or the mother (mSPB) SPB. This fast turnover rate

(~1/500 of the total duration for Dbf2-Mob1's SPB localization in anaphase) indicates that Dbf2-

Mob1 is highly mobile at SPBs.

Dbf2-Mob1 transiently localizes to the nucleolus during anaphase.

We next investigated whether Dbf2-Mob1 localizes to the nucleolus by live-cell fluorescence

microscopy. Although subtle, we observed transient localization of Mob1-eGFP to the nucleolus

in some cells as judged by co-localization with the nucleolar protein Cfi1/Net1. Importantly, this

nucleolar localization was only observed in late anaphase cells after nucleolar segregation, when

the MEN is normally active (Figure 1C).

It was reported previously that two N-terminally truncated Mob1 mutant proteins,

Mob1Δ78 and Mob1Δ132 (Mob1 missing the first 78 and 132 amino acids, respectively),

localize more prominently to the nucleus (Stoepel et al., 2005). We found that they also

displayed increased nucleolar localization (Figure 1D-F). We hypothesize that the N-terminus of

Mob1 harbors auto-inhibitory sequences that prevent access of the protein to the nucleolus.

Hence, deleting these sequences ought to cause hyperactivation of Dbf2-Mob1. Indeed, we found

that N-terminal truncation mutants of Mob1 partially suppressed temperature sensitive alleles of

upstream MEN components (*cdc15-2* and *tem1-3*; Figure S1B). This suppression was not a result

of elevated Mob1 protein levels because overexpression of Mob1 from the *GPD* promoter did

not suppress the growth defect of *cdc15-2* or *tem1-3* mutants (Figure S1A-B), nor did it increase

Mob1's nucleolar localization (Figure S1C). We conclude that N-terminal truncations result in

enhanced nucleolar localization and hyperactivation of Mob1.

To further characterize the cellular localization of Dbf2-Mob1, we quantified the relative

enrichment of full-length and truncated GFP-Mob1 at SPBs and in the nucleolus during the cell

cycle (Figure 1E). Full-length Mob1 localized to SPBs and the nucleolus during anaphase.

Mob1Δ78's localization to SPBs was similar to that of full-length Mob1 while Mob1Δ132's

dissociation from SPBs was slightly delayed. The nucleolar localization of Mob1 and truncated

Mob1 (Figure 1E) correlated with MEN activation, as judged by Mob1 association with SPBs,

translocation of the MEN activity reporter NLS$_{Cdc14}$ (Campbell et al., 2019) into the cytoplasm,

and Cdc14 release from the nucleolus (Figure S2). Consistent with earlier observations, the

Mob1 truncations displayed significantly greater nucleolar enrichment relative to full-length

Mob1 in anaphase (~30% and 120% increase on average for Mob1Δ78 and Mob1Δ132

respectively, Figure 1F). Mob1Δ78 localization to the nucleolus was, like full-length Mob1,

restricted to anaphase but accumulated in the nucleolus to higher levels. In contrast, Mob1Δ132

displayed both greater and earlier nucleolar enrichment, evident already in metaphase. We

conclude that Dbf2-Mob1 localizes to the nucleolus during anaphase when the MEN is active. N-

terminal truncation mutants of Mob1 exhibit enhanced nucleolar localization and are

hypermorphic. Given that the nucleolar localization of full-length Mob1 is quite subtle, we used

the Mob1 truncation mutants as tools to study Dbf2-Mob1's nucleolar localization.

Dbf2-Mob1 localizes to the nucleolus through interacting with Cfi1/Net1.

To validate the nucleolar localization of Mob1 we observed by microscopy and to identify the

potential nucleolar receptor for Dbf2-Mob1, we performed TurboID proximity-based labeling

(Branon et al., 2018). We fused the promiscuous biotin ligase TurboID to the MEN components

Mob1, Dbf2, Tem1 and Cdc15, and identified their protein interactors by streptavidin pull-down

followed by mass spectrometry (MS) (Figure S3A). In this experiment, we identified the

nucleolar protein Cfi1/Net1 as the top hit for Mob1- and Dbf2-TurboID labeling (Figure 2A,

S3B, Table S3). Biotinylation of Cfi1/Net1 by Mob1-TurboID was further confirmed by the

detection of a biotinylated peptide of Cfi1/Net1 (Figure S3C). Importantly, Cfi1/Net1 was only

detected in the labeling experiments where Mob1 or Dbf2 were tagged with TurboID but not

when Tem1 or Cdc15 were used as baits (Figure 2A). In contrast, Nud1, the MEN scaffold

protein at SPBs, was detected in the TurboID labeling experiments for all MEN proteins (Figure

2A).

We validated these MS findings using streptavidin gel-shift assays (Fairhead and

Howarth, 2015; Housley et al., 2014). To monitor whether a target protein was biotinylated by

the TurboID-tagged bait protein *in vivo*, we treated the denatured cell lysates with excess

streptavidin prior to immunoblotting. Biotinylated form(s) of the target protein will migrate more

slowly in SDS-PAGE due to binding of streptavidin, with each added biotin molecule causing a

theoretical size increase of up to 53 kD, the size of a streptavidin tetramer (Figure S3D). Using

this assay, we observed a slower migrating form of Nud1 in cell lysates from cells expressing all

TurboID-tagged MEN proteins (Figure 2B). Slower migrating forms of Cfi1/Net1 were only

observed in lysates obtained from cells expressing Mob1-TurboID (Figure 2B).

To determine whether Cfi1/Net1 was the sole receptor for Mob1 in the nucleolus, we

characterized the localization of Mob1Δ78 and Mob1Δ132 in cells lacking *CFI1/NET1*. While

still localized to SPBs, Mob1Δ78 and Mob1Δ132 no longer accumulated in the nucleolus during

anaphase in *cfi1/net1Δ* cells (Figure 2C-D). Furthermore, when we overexpressed *CFI1/NET1*

from the strong galactose-inducible *GAL1-10* promoter, nucleolar localization of both full-length

and N-terminal truncation mutants of Mob1 was increased by at least 50% (full-length) up to

300% (truncations) (Figure 2E-F). Interestingly, this increase in nucleolar localization was

accompanied by a decrease in Mob1's SPB localization (Figure 2F). We conclude that Dbf2-

Mob1 localization in the nucleolus during anaphase is mediated by interactions with Cfi1/Net1.

Nucleolar localization of Dbf2-Mob1 depends on MEN activation.

The localization pattern of Dbf2-Mob1 leads to the model where Dbf2-Mob1 is activated by

Cdc15 at SPBs. Active Dbf2-Mob1 then binds to and phosphorylates Cfi1/Net1, promoting the

dissociation of Cdc14 from its inhibitor to carry out mitotic exit. This model predicts that the

nucleolar localization of Dbf2-Mob1 depends on MEN activity. To test this prediction, we

employed an analog-sensitive allele of *CDC15*, *cdc15-as1* (Bishop et al., 2000; D'Aquino et al.,

2005)*.* As expected, inhibition of *cdc15-as1* (through addition of the analog 1-NA-PP1)

prevented localization of Mob1 to SPBs and translocation of the MEN activity reporter NLS_{Cdc14}

into the cytoplasm (Figure 3A). Nucleolar localization of full-length Mob1 was also significantly

reduced (Figure 3A). Nucleolar localization of the N-terminally truncated Mob1Δ78 and

Mob1Δ132 mutants, on the other hand, was only moderately reduced (Figure 3A), which is

consistent with the finding that these alleles partially suppress the temperature sensitive growth

defect of *cdc15-2* cells (Figure S1B). We conclude that Mob1 localization to the nucleolus

depends on MEN activity. The N-terminal hyperactive truncation mutations in Mob1 are less

reliant on upstream MEN kinases for their nucleolar localization.

Cdc15 regulates nuclear access of Dbf2-Mob1.

How does Cdc15 cause Dbf2-Mob1 to localize to the nucleolus? To reach the nucleolus, Dbf2-

Mob1 must first enter the nucleus. Considering that the size of the complex (102 kD) is above

the passive diffusion limit of the nuclear envelope (~40-60 kD) (Knockenhauer and Schwartz,

2016), we hypothesized that nuclear access of Dbf2-Mob1 is regulated. We explored this

possibility using the PhyB-PIF based light-inducible organelle targeting system (Yang et al.,

2013) (Figure S4A). We fused Mob1-eGFP to the PIF protein, which binds PhyB upon exposure

to red light (650 nm). Using this system, we successfully recruited Mob1-eGFP-PIF to various

subcellular locations such as SPBs or the outer mitochondrial membrane (Figure S4B).

To gauge Dbf2-Mob1's nuclear access in different cell cycle stages, we created a trap for

nuclear Dbf2-Mob1 using nucleolar-anchored PhyB (PhyB-Sik1, Figure 3B). If Dbf2-Mob1 was

shuttling between the nucleus and cytoplasm, PhyB-Sik1 would capture nuclear Mob1-eGFP-PIF

when activated by light. Interestingly, in pre-anaphase cells, we were not able to capture a

notable amount of nuclear Dbf2-Mob1 after 2 minutes of red-light activation. In contrast, in

anaphase cells in which the MEN is active, Dbf2-Mob1 was readily recruited to the nucleolus

within 2 minutes (Figure 3C). We obtained similar results with a Dbf2-PIF fusion (Figure S4C).

Furthermore, when we activated PhyB every 15 minutes for 2 minutes to recruit Mob1 to the

nucleolus in cells progressing through the cell cycle, Mob1 was only recruited to the nucleolus

by light during anaphase when the protein was also present at SPBs (Figure S4D). By

comparison, we were not able to recruit the upstream kinase Cdc15 to the nucleolus in any cell

cycle stage (Figure S4C). Given that (1) Dbf2-Mob1 protein levels do not fluctuate considerably

during the cell cycle (Visintin and Amon, 2001), that (2) the interaction between other PIF

fusions and PhyB-Sik1 are not cell cycle regulated (Yang et al., 2013) and that (3) Mob1 can be

recruited to cytoplasmic targets throughout the cell cycle (Figure S4B), we conclude that nuclear

access of Dbf2-Mob1 is cell cycle regulated.

Because nuclear access of Dbf2-Mob1 correlates with MEN activation, we next tested

whether it was regulated by the MEN by quantifying the relative enrichment of Mob1 in the

nucleolus as a function of PhyB activation time (exposure to 650 nm light) in *CDC15* or *cdc15-2*

cells. In cells with wild-type *CDC15*, light-induced nucleolar recruitment of Mob1 was higher in

anaphase than pre-anaphase cells. In contrast, in *cdc15-2* cells this difference was abolished

(Figure 3D). We conclude that in addition to activating Dbf2's kinase activity (Mah et al., 2001),

Cdc15 regulates Dbf2-Mob1's nuclear access. Consistent with this notion, we find that

Mob1Δ78, which partially suppresses the temperature sensitivity of the *cdc15-2* allele, displayed

increased nuclear access in all cell cycle stages (Figure S4E).

Dbf2-Mob1 is exported from the nucleus by Crm1.

Dbf2-Mob1 is a substrate of the nuclear exportin Crm1 *in vitro* (Kırlı et al., 2015). To

determine whether Crm1 plays a role in controlling Dbf2-Mob1 localization *in vivo*, we

quantified the nucleolar localization of full-length and truncated Mob1 in cells carrying an allele

of *CRM1* (*crm1T539C*) that is sensitive to the nuclear export inhibitor leptomycin B (LMB)

(Neville, 1999). Treatment of *crm1T539C* cells with LMB led to an increase in nucleolar

localization of both full-length and N-terminally truncated Mob1 (Figure S5A), suggesting that

Crm1 controls nuclear export of Dbf2-Mob1.

Crm1 recognizes substrates with a leucine-rich nuclear export signal (NES). To test

whether there was a functional NES in Dbf2 or Mob1, we overexpressed Dbf2 or Mob1 from the

galactose-inducible *GAL1-10* promotor. Overexpressed Mob1 was enriched in the nucleus

(Figure S5B) similar to what we observed for *pGPD-GFP-MOB1* (Figure S1C). In contrast,

when we overexpressed Mob1 together with Dbf2, Mob1 was no longer nuclear enriched,

suggesting that Dbf2 not Mob1 harbors a NES. Consistently, when overexpressed on its own,

Dbf2 exhibited diffuse localization but inhibition of *crm1T539C* with LMB led to accumulation

of Dbf2 in the nucleus (Figure S5C).

Sequence analysis identified a putative NES sequence in the N-terminus of Dbf2

beginning with L12 (Figure S5D). Incidentally, we noticed that this leucine was mutated to

methionine in a previously isolated hyperactive allele of *DBF2* (*DBF2-HyA*)(Geymonat et al.,

2009). Dbf2-HyA, when overexpressed, was nuclear enriched (Figure S5E). Inspired by this

observation, we overexpressed *DBF2* mutants where L12 had been mutated to either methionine

(*dbf2-L12M*) or alanine (*dbf2-L12A*). Both mutants accumulated in the nucleus (Figure S5E).

Furthermore, we found that the first 23 amino acids of Dbf2 were sufficient to drive nuclear

export of eGFP (Figure S5E, G). To test whether the NES in Dbf2 influences Dbf2-Mob1's

nucleolar localization in anaphase under normal expression level, we characterized Mob1's

cellular localization in *dbf2-L12A* and observed an increase in nucleolar localization by at least

40% for both full-length and N-terminally truncated Mob1(Figure 3E).

A previous phospho-proteomics study reported that S17 and S20 within the NES of Dbf2

are phosphorylated in anaphase-arrested cells (Holt et al., 2009). We found that mutating S17

and S20 to phospho-mimetic residues (S17,20D or S17,20E) disrupted the NES while mutating

these residues to alanine (S17,20A) retained the NES activity of Dbf2 (Figure S5F, G). We

propose that phosphorylation of S17 and S20 is regulated, possibly by Cdc15, to control nuclear

access of Dbf2-Mob1. This is consistent with the observation that *dbf2-S17,20A* exacerbated the

temperature sensitivity of *cdc15-2* (Figure S6A). In addition, nucleolar localization of Mob1,

particularly of Mob1Δ78, was reduced in cells harboring the *dbf2-S17,20A* allele compared to

cells with wild-type *DBF2* (Figure S6B). However, cells carrying the *dbf2-S17,20A* allele, while

exhibiting reduced nuclear access of Mob1 during all cell cycle stages, still showed differential

nuclear access between pre-anaphase and anaphase, as is observed in *DBF2* cells (Figure S6C).

This observation suggests that additional regulatory mechanism(s) control Dbf2-Mob1's nuclear

access. In contrast, cells harboring the *dbf2-S17,20D* allele exhibited increased nuclear access of

Mob1 during all cell cycle stages (Figure S6C), confirming that Dbf2-Mob1 is normally kept out

of the nucleus through Dbf2's NES. The NES sequence in Dbf2 is well conserved among

Saccharomycetes (Figure S5D) suggesting that regulated nuclear access of Dbf2-Mob1 is

conserved at least across this class of fungi.

Nucleolar localization of Dbf2-Mob1 is regulated by Cdc5.

The analysis of nuclear access and nucleolar localization of the N-terminal truncations of Mob1

indicated that the MEN is not the only pathway controlling Dbf2-Mob1's nucleolar localization.

The truncation mutants localize to the nucleolus in a manner largely independent of the MEN,

yet their nucleolar localization is still restricted to metaphase and anaphase (Figure 3A). This

restriction of nucleolar localization is not due to limited nuclear access. Truncated Mob1 mutants

have increased nuclear access prior to anaphase (Figure S4E). These data indicate that nucleolar

localization or the interaction between Dbf2-Mob1 with Cfi1/Net1 is regulated by additional

factors.

An obvious candidate for this additional regulator is the Polo-like kinase Cdc5, which is

active throughout mitosis and plays multiple essential roles in mitotic exit (Lee et al., 2005). As

part of both the FEAR network (Rock and Amon, 2009; Stegmeier et al., 2002) and the MEN,

Cdc5 is indispensable for Cdc14's nucleolar release. However, the exact role(s) of Cdc5 during

this process is not fully understood. To determine whether Cdc5 regulated binding of Dbf2-

Mob1 to Cfi1/Net1, we examined the consequences of inhibiting Cdc5's kinase activity on

nucleolar localization of Dbf2-Mob1 using an analog-sensitive allele of *CDC5* (*cdc5-as1*).

Consistent with the known functions of Cdc5 in MEN activation, we observed loss of Mob1's

SPB localization and Dbf2-Mob1's kinase activity as monitored by translocation of the NLS_{Cdc14}

reporter into the cytoplasm when Cdc5 was inhibited (Figure 4A). Nucleolar localization of

Mob1, Mob1Δ78 and Mob1Δ132 was also lost in cells lacking Cdc5 activity (Figure 4A). This is

in direct contrast to Cdc15 inhibition, where the nucleolar localization of N-terminal Mob1

truncation mutants particularly Mob1Δ132 was only partially reduced (Figure 3A). These results

suggested that Cdc5 regulates Dbf2-Mob1's nucleolar localization independently of its role in

activating Cdc15.

To directly determine whether *CDC5* regulated Dbf2-Mob1's nucleolar localization

independently of the MEN, we took advantage of a hyperactive *CDC15* allele, *GAL-CDC15(1-*

750) (Bardin et al., 2003), which is active even in the absence of *CDC5* (Rock and Amon, 2011).

When *GAL-CDC15(1-750)* was expressed, Mob1's SPB localization and Dbf2-Mob1's kinase

activity was no longer restricted to anaphase but rather was high throughout the cell cycle as a

result of MEN hyper-activation (Figure 4B). Interestingly, *GAL-CDC15(1-750)* did not abolish

cell-cycle regulation of Mob1's nucleolar localization but rather advanced it to early anaphase

and metaphase (Figure 4B). In cells expressing *GAL-CDC15(1-750),* inactivation of *CDC5* still

abolished nucleolar localization of both full-length and the hyperactive N-terminally truncated

Mob1 while Mob1 binding to SPBs was unaffected (Figure 4B). These results demonstrate that

nucleolar localization of Dbf2-Mob1 directly depends on Cdc5 independently of its role in MEN

activation.

Could *CDC5* regulate the nucleolar localization of Dbf2-Mob1 through its role in the

FEAR network? To test this, we quantified Mob1's nucleolar localization in cells lacking the

FEAR network component *SLK19* (*slk19Δ*) or carrying a temperature sensitive allele of the

FEAR network effector *CDC14* (*cdc14-3*). MEN activation (as determined by Mob1 localization

to the dSPB and nuclear release of the NLS_{Cdc14} reporter) and as a result mitotic exit were

considerably delayed and more variable in *slk19Δ* cells (Figure S7A). Consistent with a delay in

MEN activation, nucleolar localization of Mob1 and Mob1Δ78 but not Mob1Δ132 was also

delayed. Importantly, maximum enrichment of Mob1 in the nucleolus was not reduced in *slk19Δ*

cells for all three forms of Mob1 (Figure S7A). We observed similar results in *cdc14-3* mutants

(Figure S7B). We conclude that Cdc5 regulates Dbf2-Mob1's nucleolar localization through

mechanisms in addition to its role in the MEN and the FEAR network.

Cdc5 and Dbf2-Mob1 phosphorylate Cfi1/Net1 at distinct sites.

Our results suggest a model where Cdc5 promotes the interaction between Dbf2-Mob1 and its

nucleolar receptor Cfi1/Net1, likely through phosphorylating Cfi1/Net1. This interaction then

facilitates phosphorylation of Cfi1/Net1 by Dbf2-Mob1 to bring about the release of Cdc14 from

Cfi1/Net1. Cfi1/Net1 is a highly phosphorylated protein with 64 known phosphorylation sites *in*

vivo (Holt et al., 2009; Swaney et al., 2013). About one fifth of these sites were identified as

CDK targets (Holt et al., 2009) including six key CDK sites whose phosphorylation is controlled

by the FEAR network (Azzam, 2004). To map sites in Cfi1/Net1 that are phosphorylated in a

CDC5 or MEN-dependent manner, we performed phosphoproteomics analyses on wild-type

anaphase cells and cells in which Cdc5 or Cdc15 were inhibited using the *cdc5-as1* and *cdc15-*

as1 alleles, respectively (Figure S8A). This analysis identified 44 of the 64 previously known

sites in Cfi1/Net1 and 18 new sites (Table S4). To achieve complete or close to complete

coverage of Cfi1/Net1's phosphorylation sites in anaphase, we also performed

immunoprecipitation-mass spectrometry (IP-MS) for Cfi1/Net1 in anaphase enriched cultures

and identified 9 additional sites (Table S4) resulting in an astonishing total of 91 phosphorylation

sites in Cfi1/Net1. These phosphorylation sites appear to cluster in regions of disorder as

predicted by the PONDR score (Romero et al., 1997) (Figure 5A).

By comparing the peptide signals between wild-type, *cdc15-as1* and *cdc5-as1* cells in our

quantitative phosphoproteomics dataset, we identified phosphorylation sites that depended on

Cdc15 or Cdc5 activity or both (Figure S8B-E). Among them, we found 11 *CDC15*-dependent

and 22 *CDC5*-dependent sites in Cfi1/Net1. Six of the *CDC15*-dependent sites fit Dbf2-Mob1's

preferred sequence motif RXXS* (* represents the phosphorylation site) (Mah et al., 2005),

supporting our model that Dbf2-Mob1 phosphorylates Cfi1/Net1. Given that Cdc5 activates

Cdc15, sites that depended on *CDC15* ought to also depend on *CDC5*. This was indeed the case

for 10 of the 11 *CDC15*-dependent phosphorylation sites in Cfi1/Net1.

To identify sites that only depended on *CDC5* but not *CDC15*, we subtracted *CDC15*-

dependent sites from *CDC5*-dependent sites yielding 12 sites (denoted as *CDC5*-only, Figure

5A). Based on the distribution of all 91 phosphorylation sites and degree of disorder, we divided

Cfi1/Net1 into four zones: residues 31-69 (z1, 7 sites), 160-615 (z2, 53 sites), 676-840 (z3, 12

sites) and 1017-1166 (z4, 19 sites) (Figure 5A). *CDC15*-dependent phosphorylation sites were

concentrated in zone 2, whereas *CDC5*-dependent sites were also found in zones 1 and 3. In

contrast to Cfi1/Net1, all the *CDC5*-dependent phosphorylation sites in Dbf2 and Mob1 were

also *CDC15*-dependent (Figure S8F). These data indicate that Cdc5 and Dbf2-Mob1 directly

phosphorylate Cfi1/Net1. Dbf2-Mob1 on the other hand is a direct substrate of Cdc15 but not

Cdc5.

Cdc5 promotes Dbf2-Mob1's nucleolar localization by phosphorylating Cfi1/Net1.

Having identified the phosphorylation sites within Cfi1/Net1 we next asked whether they were

important for the interaction between Cfi1/Net1 and Dbf2-Mob1. We generated a *CFI1* allele in

which all 91 phosphorylation sites were mutated to alanine (*cfi1-91A*). Cells harboring this allele

as the sole source of *CFI1/NET1* were viable and progressed through anaphase with only a slight

delay in mitotic exit as judged by the timing of Mob1's dissociation from the SPBs (Figure 5B).

Interestingly, Mob1Δ132, which showed the most pronounced nucleolar localization among all

Mob1 alleles analyzed, still localized to the SPBs during anaphase in *cfi1-91A* cells, but failed to

accumulate in the nucleolus (Figure 5B-C, S9A). We conclude that Cfi1/Net1 phosphorylation is

required for interacting with Dbf2-Mob1.

Next, we tested whether *CDC5*-dependent phosphorylation of Cfi1/Net1 regulated Dbf2-

Mob1's nucleolar localization. We focused our analysis on zone 2 of Cfi1/Net1 because previous

studies had shown that the first 621 amino acids of the protein are sufficient to confer Cdc14

regulation (Azzam, 2004). We further note that mutating phosphorylation sites in zone 1 and 3 of

Cfi1/Net1 did not affect the nucleolar localization of Mob1Δ132 (Figure 5B-C, S9A). We

generated a *CFI1/NET1* allele with mutated phosphorylation sites in zone 2 that were

phosphorylated in a *CDC5*-dependent but *CDC15*-independent manner (henceforth *cfi1-*

Cdc5only(z2)). For comparison we generated a *CFI1/NET1* allele in which we only mutated sites

that were phosphorylated in a *CDC15*- dependent manner (henceforth *cfi1-Cdc15(z2)*). Analysis

of Mob1Δ132 localization in these mutants revealed that, similar to inhibition of Cdc5, *cfi1-*

Cdc5only(z2) abolished the nucleolar localization of Mob1Δ132, while *cfi1-Cdc15(z2)* slightly

increased nucleolar localization of the protein (Figure 5D-E, S9B). To validate these findings, we

performed TurboID labeling experiments followed by Streptavidin gel-shift assays to probe the

interaction between TurboID tagged full-length Mob1 and 13Myc tagged Cfi1/Net1. The slower

migrating form corresponding to biotinylated Cfi1/Net1 was absent in cells expressing *cfi1-*

Cdc5only(z2) but present in cells expressing *CFI* or *cfi1-Cdc15(z2)* (Figure 5F). We conclude

that phosphorylation of Cfi1/Net1 by Cdc5 is required for Dbf2-Mob1 binding to the protein.

Three lines of evidence indicate that Cdc5 directly phosphorylates Cfi1/Net1. First,

consistent with the finding that Cdc5 is already active in metaphase, 10 out of 12 *CDC5*-only

sites identified in anaphase cells are already phosphorylated in cells arrested in metaphase using

the microtubule depolymerizing drug nocodazole (Figure S10) with 6 of them also determined to

be *CDC5*-dependent in metaphase (Figure S10E). Second, we found considerable overlap

between *CDC5*-dependent phosphorylation sites *in vivo* and sites identified *in vitro* (Loughrey

Chen et al., 2002; Shou et al., 2002). Four out of five *in vivo* Cdc5only sites in region 1-341 were

previously found to be phosphorylated by Cdc5 *in vitro* (Table S4). Third, Cdc5 and Cfi1/Net1

interact with each other *in vivo* as determined by TurboID labeling (Figure S11A). The

biotinylation of Cfi1/Net1 by Cdc5-TurboID was further confirmed by the detection of

biotinylated peptides in Cfi1/Net1 as well as the streptavidin gel-shift assay (Figure S11B-C).

We conclude that Cdc5 phosphorylates Cfi1/Net1 at the onset of metaphase which serves as a

priming event for Dbf2-Mob1 binding to Cfi1/Net1 in anaphase. We note that these findings also

explain why Mob1Δ132's nucleolar localization is already evident in metaphase (Figure 4B).

Dbf2-Mob1 promotes the release of Cdc14 from the nucleolus through Cdc5-mediated priming

of Cfi1/Net1.

To determine whether the *CDC5* and MEN-dependent phosphorylation sites in Cfi1/Net1

regulate the interaction between Cfi1/Net1 and Cdc14 as our model predicted, we examined the

consequences of disrupting these phosphorylation sites on the release of Cdc14 from the

nucleolus. We first determined which region of Cfi1/Net1 was mediating phospho-regulation of

this interaction. We mutated all phosphorylation sites in the individual zones as well as in

combination and analyzed the effects on Cdc14 release from the nucleolus (Figure S12A-B).

This analysis revealed that only phosphorylation in zone 2 controlled Cdc14 release from

Cfi1/Net1 (Figure S12A-B). It is important to note that mutating the phosphorylation sites in

zone 2 also affected the ability of Cfi1/Net1 to bind Cdc14, as judged by the lower degree of

Cdc14 nucleolar sequestration prior to anaphase and localization of Cdc14 to the dSPB prior to

anaphase (Figure S12C). This finding indicates that the same residues involved in regulating the

interaction between Cfi1/Net1 and Cdc14 are also important for forming the complex in the first

place and mutating them to alanine weakens this interaction. Alternatively, mutating so many

residues at once (53 sites in zone 2) could change the structure of Cfi1/Net1 and thus disrupt

binding to Cdc14. Nevertheless, the increased level of free Cdc14 in the cell with *cfi1-91A* and

the zone 2-phosphomutant explains why such severe defect in Cdc14 release from the nucleolus

did not cause a significant anaphase delay as assayed by the kinetics of Cdc14 re-sequestration.

Next, we examined kinase-specific phospho-mutants in Cfi1/Net1. There are three known

kinases that phosphorylate Cfi1/Net1 to regulate binding to Cdc14: mitotic CDKs (mainly Clb2-

Cdk1), Cdc5 and Dbf2-Mob1. CDK phosphorylation of Cfi1/Net1 during early anaphase

underlies FEAR network-dependent release of Cdc14 from its inhibitor (Azzam, 2004). In FEAR

network mutants such as *slk19Δ*, Cdc14 release from the nucleolus and anaphase progression are

delayed and are accompanied by increased cell-to-cell variability (Figure 6A). As reported

previously (Azzam, 2004), cells harboring a *CFI1/NET1* allele with 6 CDK sites mutated to

alanine, *cfi1/net1-6Cdk*, caused Cdc14 release defects similar to those of FEAR network mutants

(Figure 6B).

Inactivation of the MEN using the *cdc15-as1* allele led to the previously described

pattern of Cdc14 localization, where Cdc14 is initially released from the nucleolus by the FEAR

network during early anaphase but is then re-sequestered in the nucleolus during later stages of

anaphase (Stegmeier et al., 2002) (Figure 6A). Mutating the *CDC15*-dependent phosphorylation

sites in zone 2 (*cfi1-Cdc15(z2)*) resulted in a significant reduction of Cdc14 release from the

nucleolus but only recapitulated ~50% of the effect of inactivating *CDC15* (compare Figure 6A

and B). As previously reported (Visintin et al., 2008), inhibition of the analog sensitive *cdc5-as1*

allele caused defects in both FEAR network and MEN-mediated release of Cdc14 from the

nucleolus (Figure 6A). Mutating the *CDC5*-only phosphorylation sites in zone 2 (*cfi1-*

Cdc5only(z2)) resulted in a similar reduction of Cdc14 release from the nucleolus (Figure 6B).

Finally, combining *cfi1-Cdc5only(z2)* with *cfi1-Cdc15(z2)* (*cfi1-Cdc15&Cdc5(z2)*) caused an

even greater defect in Cdc14 release from the nucleolus than either mutant alone (Figure 6B,

S13). These results confirmed our model where Cdc5, in addition to activating the MEN, directly

phosphorylates Cfi1/Net1 to target Dbf2-Mob1 to Cfi1/Net1. Phosphorylation of Cfi1/Net1 by

Dbf2-Mob1 then promotes the dissociation of Cdc14 from Cfi1/Net1.

Phosphorylation of Cfi1/Net1 by Cdc5 and Dbf2-Mob1 promotes mitotic exit.

Preventing the dissociation of Cdc14 from its inhibitor during anaphase ought to interfere

with mitotic exit. Indeed, we observed a delay in all mutants analyzed (Figure 6C, S14). Both

cfi1-Cdc15(z2) and *cfi1-Cdc5only(z2)* mutant cells exited mitosis with an average delay of ~6

minutes (~25% increase, Figure 6C, S14C). In addition, consistent with Cdc5's role in the FEAR

network, we observed a significant delay in nucleolar segregation in *cfi1-Cdc5only(z2)* mutant

cells (Figure 6C, S14C). Surprisingly, *cfi1-Cdc15&Cdc5(z2)* double mutant cells which had the

most severe defect in Cdc14 release from the nucleolus exhibited a similar delay in mitotic exit

as the *cfi1-Cdc15(z2)* and *cfi1-Cdc5only(z2)* single mutants and a less severe defect in nucleolar

segregation compared to *cfi1-Cdc5only(z2)* (Figure 6C). This relatively short delay in mitotic

exit is likely due to the fact that Cdc14 was not tightly sequestered in this mutant prior to

anaphase. We observed elevated levels of Cdc14 at dSPB in both *cfi1-Cdc15(z2)* and *cfi1-*

Cdc15&Cdc5(z2) cells prior to anaphase (Figure 6D, S13). It appears that mutating *CDC15-*

dependent sites in Cfi1/Net1 interferes with its ability to bind Cdc14. Nonetheless, the majority

(>60%) of Cdc14 was not released during anaphase in cells harboring *cfi1-Cdc15&Cdc5*.

One possible explanation for the apparently discrepancy between the severity of the

Cdc14 release defect and the more subtle delay in mitotic exit is that in addition to Cfi1/Net1,

MEN also promotes cytoplasmic retention of Cdc14 by phosphorylating its NLS (Mohl et al.,

2009). To determine the contribution of this MEN function to promoting mitotic exit in

Cfi1/Net1 mutants, we constructed a *CDC14* mutant (*cdc14-3A*) in which all three potential

Dbf2-Mob1 phosphorylation sites capable of driving translocation of the NLS_{Cdc14} reporter into

the cytoplasm were mutated to alanine (Figure S15A). Interestingly, cells expressing *cdc14-3A*

instead of *CDC14* exited from mitosis 3 minutes faster (Figure S15B) indicating that cytoplasmic

retention of Cdc14 delays rather than promotes mitotic exit.

Another possibility is that some MEN-dependent phosphorylation sites in Cfi1/Net1

eluded our MS analysis. However, given that Cdc14 is a potent phosphatase, we favor the idea

that, incomplete sequestration of Cdc14 prior to anaphase and the residual release of Cdc14 from

the nucleolus during anaphase combined with a functional FEAR network-mediated release of

Cdc14 from the nucleolus during early anaphase is sufficient to promote exit from mitosis with

only a modest delay in the *cfi1-Cdc15&Cdc5* mutant. Consistent with this idea, we found that the

cfi1-Cdc5only(z2) mutant, which does not display defects in sequestering Cdc14 prior to

anaphase, was synthetic lethal with the FEAR network mutants *slk19Δ* and *spo12Δ* (Figure 6E).

We conclude that *CDC5* and MEN-dependent phosphorylation of Cfi1/Net1 controls the

protein's binding to Cdc14 and exit from mitosis.

DISCUSSION

As a model system for mitotic exit control and cellular signaling in eukaryotes, the MEN has

been studied extensively for decades. Yet, how the MEN activates its effector Cdc14 to promote

exit from mitosis has remained an enigma. Furthermore, the spatial aspect of signal transmission

in the MEN, namely how a signal generated at the outer plaque of the SPBs in the cytosol

reaches its target in the nucleolus, was largely unexplored. Our study provides mechanistic

insights into these questions and together with prior observations lead to a model for Cdc14

regulation and mitotic exit in budding yeast (Figure 7).

The role of the Polo-like kinase Cdc5 in regulating Cdc14 activation.

The Polo-like kinase Cdc5 is essential for Cdc14 activation and is part of both the FEAR

network and the MEN. However, the exact role(s) of Cdc5 in regulating Cdc14's nucleolar

release has remained elusive due to its multiple functions in the MEN and the FEAR network.

Using an allele that bypasses *CDC5*'s role in MEN activation, we revealed a novel function of

Cdc5 as a priming kinase that targets Dbf2-Mob1 to its substrate Cfi1/Net1.

It was hypothesized that phosphorylation of Cfi1/Net1 by Cdc5 weakens the interaction

between Cfi1/Net1 and Cdc14. Additional phosphorylation by mitotic CDKs or Dbf2-Mob1

were thought to further reduce the interaction resulting in the dissociation of Cdc14 from its

inhibitor. Our results suggest an alternative model. Instead of, or at least in addition to,

weakening the Cfi1/Net1-Cdc14 interaction, Cdc5 phosphorylation targets Dbf2-Mob1 to

Cfi1/Net1. Mutating the Cdc5-only phosphorylation sites in Cfi1/Net1 abolished nucleolar

enrichment of Dbf2-Mob1 and resulted in a reduction in MEN-mediated dissociation of Cdc14

from Cfi1/Net1. Interestingly, this mutation also caused severe defects in the FEAR network

mediated release of Cdc14 from Cfi1/Net1. We thus propose that Cdc5 priming phosphorylation

is required not only for Dbf2-Mob1 to phosphorylate Cfi1/Net1 but also for mitotic CDKs

(Figure 7). Consistent with this hypothesis, we found that most Cdc5 phosphorylation sites in

Cfi1/Net1 are already phosphorylated by Cdc5 in metaphase (Figure S10E). This observation

indicates that the docking site(s) on Cfi1/Net1 for Dbf2-Mob1 (and mitotic CDKs) is already

present in metaphase prior to the activation of the FEAR network and the MEN (Figure 7). This

model is further supported by the observation that inhibition of Cdc5 eliminates CDK mediated

phosphorylation of T212 in Cfi1/Net1 (Azzam, 2004). Cdc5 activity is regulated by the DNA

damage checkpoint (Cheng et al., 1998; Sanchez, 1999). We speculate that making FEAR

network and MEN-mediated release of Cdc14 from the nucleolus dependent on Cdc5's priming

activity ensures that DNA damage has been repaired and the checkpoint silenced prior to exit

from mitosis.

The FEAR network and the MEN regulate Cdc14 binding to Cfi1/Net1 by different

mechanisms.

While both mitotic CDKs and Dbf2-Mob1 appear to require *CDC5*-dependent priming

phosphorylation of Cfi1/Net1, the mechanism whereby mitotic CDKs and Dbf2-Mob1 disrupt

the interaction between Cfi1/Net1 and Cdc14 is quite different. Mitotic CDKs phosphorylate

Cfi1/Net1 mainly on S166, T212, S252, T297 and T304; Dbf2-Mob1 targets sites S259, S295,

S362, S439 and S497 (Figure 5A, Table S4). Given that mitotic CDKs and Dbf2-Mob1 target

different sites, we propose that increasing the acidity of aa160–500 within Cfi1/Net1 rather than

site-specific phosphorylation disrupts the interaction between Cdc14 and its inhibitor. Cfi1/Net1

is an integral part of the nucleolus, which has recently been described as a phase-separated

structure (Feric et al., 2016; Shin and Brangwynne, 2017). Overall phosphorylation rather than

phosphorylation of specific sites has been shown to disrupt interactions within such structures

(Carpenter et al., 2018; Owen and Shewmaker, 2019). Perhaps extraction of Cdc14 from the

nucleolar phase requires a similar mechanism.

The MEN as a model for cross-compartment signaling.

The MEN, most closely related to the Hippo pathway, employs most, if not all, of the principles

governing classic receptor tyrosine signaling logic to convey a signal generated at SPBs to the

MEN effector Cdc14 in the nucleolus: (1) scaffold assisted signaling (at the SPB), (2) signal

transmission across organelle boundaries - from the cytoplasm to the nucleus, and (3) activation

of the effector in a sub-compartment (the nucleolus). As such, we believe that the molecular

mechanisms governing MEN activity are broadly applicable to intracellular signal transmission

in general.

(1) Dynamic scaffold assisted signaling.

Scaffold assisted assembly of signaling complexes is a widespread phenomenon in eukaryotic

signal transduction cascades (Good et al., 2011). We find that in the MEN, assembly of Cdc15-

(Dbf2-Mob1) signaling complex on the scaffold Nud1 is highly dynamic and this dynamicity is

crucial for effector activation. We propose that this dynamicity also serves to amplify the signal.

Cdc15 is the limiting enzyme of the pathway: it is the least abundant component of the MEN and

hyperactivating Cdc15 increases Dbf2-Mob1's kinase activity by > 40-fold (Rock and Amon,

2011). We further hypothesize that the relatively low affinity of Mob1 for phosphorylated-Nud1

(K_d = 2.4 µM; note that for an optimal Mob1 binding phosphopeptide the K_d is 174 nM) (Rock et

al., 2013) is selected for to facilitate the fast turnover rate of Dbf2-Mob1 at SPBs and thus to

promote release of the kinase and signal amplification. In this model, the binding

affinity/kinetics of kinases to their signaling scaffolds is an important parameter that cells fine-

tune to generate desirable signaling properties of scaffold-assisted signaling pathways.

(2) Regulated compartment access.

Most signals, be they generated outside or within the cell, ultimately, result in a nuclear

response. As such, signals have to be propagated from the cytoplasm into the nucleus. Our

studies have led to the discovery that in the MEN, this nuclear access is cell cycle regulated.

Prior to anaphase, Dbf2-Mob1 is actively exported out of the nucleus by Crm1 through the

conserved NES within the N-terminus of Dbf2. Upon MEN activation, nuclear partitioning of

Dbf2-Mob1 increases, likely a result of both increased nuclear import through modification of

Mob1 and decreased nuclear export through modification of Dbf2.

Disrupting the NES resulted in an increase in nuclear/nucleolar localization of Dbf2-

Mob1 in all cell cycle stages, interestingly, including anaphase (Figure 3E). This suggests that

only a small fraction of Dbf2-Mob1 is activated by Cdc15 at any given time during anaphase due

to limited levels of Cdc15 as discussed earlier. In addition, active Dbf2-Mob1 is likely needed in
the cytosol to phosphorylate substrates other than Cfi1/Net1 such as those involved in
cytokinesis. We speculate that fine-turning the balance of nuclear versus cytosolic Dbf2-Mob1,
possibly through maintaining a dynamic shuttling of active Dbf-Mob1 between the nucleus and
cytoplasm, is important for the timing of late cell cycle events. The dynamic shuttling of Dbf2-
Mob1 in combination with the relatively small fraction of active Dbf2-Mob1 would also explain
the absence of visible nuclear translocation of Dbf2-Mob1 upon activation. Interestingly,
mammalian Dbf2, known as LATS which are thought to mainly function in the cytosol (Yu and
Guan, 2013), has been found to localize to the nucleus (Britschgi et al., 2017; Li et al., 2014),
indicating nuclear shuttling of kinases might play a role in Hippo signaling as well.
579
(3) Substrate targeting by priming phosphorylation.
Upon entry into the nucleus, Dbf2-Mob1 specifically functions in the nucleolus to promote the
dissociation of Cdc14 from its inhibitor Cfi1/Net1. Priming phosphorylation by Cdc5 on
Cfi1/Net1 ensures that Dbf2-Mob1 executes this function effectively. Dbf2-Mob1 binds to
Cdc15-phosphorylated Nud1 through Mob1's phosphoserine-threonine binding domain (Rock et
al., 2013). We propose that a similar mechanism mediates the interaction between Dbf2-Mob1
and Cdc5-phosphorylated Cfi1/Net1. This hypothesis is supported by the finding that during
cytokinesis, phosphorylation of Dbf2-Mob1's substrate Hof1 by Cdc5 facilitates the binding of
Hof1 to Mob1 (Meitinger et al., 2011; Rock et al., 2013). Based on our observation that a quarter
of potential Dbf2-Mob1 substrates are also targets of Cdc5 (Figure S8E) we further speculate
that priming phosphorylation by Cdc5 is a general mechanism for targeting Dbf2-Mob1 to its
substrates.
It is worth noting that Cdc5's consensus motif with an acidic residue at the -2 position of
the pS/T (Kettenbach et al., 2011; Nakajima et al., 2003) does not fit well with the optimal Mob1
binding motif which prefers Y/F at the -2 position of the pS/T (Rock et al., 2013). This is likely
beneficial as a low affinity between Dbf2-Mob1 and its substrates would increase substrate
turnover and prevent sequestration of active Dbf2-Mob1. This low affinity would also explain
the weak and transient localization of Dbf2-Mob1 in the nucleolus that evaded detection
previously. Mob1's phosphoserine-threonine binding domain is well conserved from yeast to

humans (Rock et al., 2013) suggesting that substrate targeting through priming phosphorylation

might occur for LATS-MOB1 in the Hippo pathway as well.

Priming phosphorylation is a widely used mechanism to ensure effective kinase action at

a particular site in the cell. Perhaps the best studied example for priming is the Polo-like kinases

whose polo-box domain directs the kinase to specific subcellular structures and substrates that

have been previously phosphorylated by a priming kinase such as CDKs (Elia et al., 2003;

Lowery et al., 2005). Although MAP kinases are mainly directed to their substrates through

specialized docking motifs (Bardwell, 2006; Cargnello and Roux, 2011) without priming, it has

been demonstrated that successive phosphorylation through priming could contribute to the

sensing of MAPK signal duration and strength (Murphy et al., 2002). These examples together

with our findings for Dbf2-Mob1 underscore the importance of priming phosphorylation as a

conserved paradigm for achieving specificity and efficiency in cellular signal transduction.

ACKNOWLEDGEMENTS

We thank F. Luca (Philadelphia, USA), R. Deshaies (Caltech, USA), E. Schiebel (ZMBH,

Germany), E. Unal (Berkeley, USA) and J. Haber (Waltham, USA) for strains and reagents; R.

Schiavoni and the Biopolymers & Proteomics Facility at Koch Institute Swanson Biotechnology

Center for processing samples for the TurboID-MS experiments; members of our laboratory for

their critical reading of the manuscript. This work was supported by National Institute of General

Medical Science (GM 118066 to A.A.) and the Eunice Kennedy Shriver National Institute of

Child Health and Human Development (HD085866 to A.A.). X.Z. was supported by a Helen

Hay Whitney postdoctoral fellowship. A.A. is also investigator of the Howard Hughes Medical

Institute and the Paul F. Glenn Center for Biology of Aging Research at MIT.

AUTHOR CONTRIBUTIONS

X.Z. and A.A. conceived the study. X.Z. performed all experiments. W.L. and Y.L. provided the

phosphoproteomic analysis. X.Z. and A.A. wrote the manuscript with input from all coauthors.

COMPETING INTERESTS

The authors declare no competing interests.

    FIGURES

Figure 1. Dbf2-Mob1 transiently localizes to the nucleolus in late anaphase.

(**A**) Major components of the MEN and their subcellular localization.

(**B**) FRAP analysis of Mob1-eGFP (A39695). Red circles indicate the area of photo-bleaching.

Cells were grown and imaged at room temperature in SC medium + 2% glucose. Graph to the

right represents average measurements of double normalized fluorescence intensities ($n = 6$ cells)

after correcting for photo-bleaching during acquisition. Red curve is the average fit and shaded

area represents standard deviation (SD) of the fits. Half recovery time $t_{1/2}$ ± SD is indicated.

(**C**) Localization of Mob1 during the cell cycle. A40257 (with Mob1-eGFP, Cfi1-mCherry and

Spc42-mCherry) cells were grown at room temperature in SC medium + 2% glucose and imaged

every minute for 2 hours. Arrows highlight the nucleolar localization.

(**D**) Nucleolar localization of full-length (A39931) and N-terminally truncated (A39933 and

A39935) Mob1. Cells were grown at room temperature in SC medium + 2% glucose and imaged

every 3 minutes for 4 hours. Arrows highlight the nucleolar localization.

(**E**) Enrichment of Mob1 (A41211, n = 62 cells), Mob1Δ78 (A41212, n = 60 cells) and

Mob1Δ132 (A41213, n = 48 cells) at the dSPB (left) and in the nucleolus (right) as a function of

cell cycle progression. Cells were grown at 25°C in SC medium + 2% glucose and imaged every

3 minutes for 4 hours. Single cell traces were aligned based on anaphase onset, as defined as

spindle length > 3 μm (measured based on SPB marker Spc42-mCherry), and averaged. Solid

lines represent the average, shaded areas represent 95% confidence intervals.

(**F**) Maximum enrichment of full-length Mob1 (WT) and truncated Mob1 (Mob1Δ78 and

Mob1Δ132) in the nucleolus in anaphase of cells in (E). Solid lines represent the median. ****P

< 0.0001 by two-sided Wilcoxon rank sum test.

Figure 2. Dbf2-Mob1 localizes to the nucleolus through interacting with Cfi1/Net1.

(**A**) Results of proximity-based biotinylation with TurboID for MEN proteins (A41367, A41370, A41368, A41369 and A2588). Cells were grown at room temperature in YEP + 2% glucose + 50 µM biotin.

(**B**) Streptavidin gel-shift assays to probe the interactions of TurboID-labeled MEN proteins with the MEN scaffold Nud1 (left, A11869, A41381, A41382 and A41380) or Cfi1/Net1 (right, A1638, A41406, A41407 and A41372). Cells were grown at room temperature in YEP + 2% glucose and lysates were treated with streptavidin and immunoblotted as indicated. Red arrows highlight biotinylated proteins.

(**C** and **D**) Representative images (C) and quantification (D) of Mob1Δ78 localization in wild-type *CFI1/NET1* (A41344, *n* = 106 cells) or *cfi1/net1Δ* (A41347, *n* = 18 cells) cells and Mob1Δ132 localization in *CFI1/NET1* (A41345, *n* = 95 cells) or *cfi1/net1Δ* (A41348, *n* = 18

cells) cells. Cells were grown at 25°C in SC medium + 2% glucose and imaged every 5 minutes

for 4 hours. Arrows highlight nucleolar localization.

(**E** and **F**) Representative images (E) and quantification (F) of Mob1 localization in wild-type

(A41343, *n* = 110 cells) or *GAL-CFI1/NET1* expressing cells (A41340, *n* = 71 cells), Mob1Δ78

localization in wild-type (A41344, *n* = 103 cells) or *GAL-CFI1/NET1* expressing cells (A41341,

n = 68 cells), and Mob1Δ132 localization in wild-type (A41345, *n* = 71 cells) or cells expressing

GAL-CFI1/NET1 (A41342, *n* = 53 cells). # denotes that the image was linearly contrast adjusted

to avoid over-saturation for Mob1Δ78 and Mob1Δ132. Cells were first grown at room

temperature in SC medium + 2% raffinose. Cells were then mounted onto agarose pads made

with SC medium + 1% raffinose + 1% galactose and imaged every 5 minutes for 5 hours at

25°C. Arrows highlight nucleolar localization. Solid lines represent the average of single cell

traces aligned to anaphase onset while shaded areas represent 95% confidence intervals. For

maximum enrichment, each dot represents a single cell. The solid lines represent the median.

****$P < 0.0001$ by two-sided Wilcoxon rank sum test.

Figure 3. *CDC15* regulates nuclear access of Dbf2-Mob1.

(**A**) Enrichment of Mob1 at the dSPB, in the nucleolus, and Dbf2-Mob1's kinase activity were
determined in cells going through anaphase in *CDC15* (A41211, A41212, and A41213; *n* = 74,
94, and 55 cells respectively) or *cdc15-as1* (A41214, A41215, and A41216; *n* = 37, 63, and 30
cells respectively) cells. Cells were grown at 25°C in SC medium + 2% glucose and 10 μM 1-
NA-PP1 and imaged every 3 minutes for 4 hours.

(**B**) Probing Dbf2-Mob1's nuclear access by recruiting Mob1 to the nucleolus with the PhyB-PIF
optogenetics system. By anchoring PhyB to the nucleolus, diffuse nuclear Dbf2-Mob1, if
present, can be visualized by recruiting Mob1-eGFP-PIF to the nucleolus.

(**C**) Recruiting Mob1 to the nucleolus at different cell cycle stages. A40260 cells were grown at
25°C in SC medium + 2% glucose, imaged after a 2-hour incubation with 12.5 μM PCB in the

dark. Red dot denotes the frame where 650 nm light was applied to activate PhyB. Yellow

arrows highlight the light-induced recruitment.

(**D**) Recruitment of Mob1 to the nucleolus in *CDC15* (A41360) or *cdc15-2* (A41361) cells.

Quantifications of Mob1's enrichment in the nucleolus as a function of PhyB activation time in

CDC15 (A41360, $n = 27$ and 16 cells for pre-anaphase and anaphase respectively) or *cdc15-2*

(A41361, $n = 14$ and 36 cells for pre-anaphase and anaphase respectively) cells. Cells were

grown at room temperature in SC medium + 2% glucose, incubated with 12.5 µM PCB for 2

hours in the dark, and shifted to 34 °C for 50 minutes before imaging.

(**E**) Nucleolar enrichment of wild-type and truncated Mob1 in wild-type and *dbf2-L12A* cells

(A41394, A41395 and A41396; $n = 32$, 28 and 31 cells respectively). Wild-type traces for

comparison were the same as in Fig. 1E. Cells were grown similarly as in Fig. 1E.

For graphs in (A) and (E), solid lines represent the average of single cell traces aligned to

anaphase onset. Shaded areas represent 95% confidence intervals. For maximum enrichment,

each dot represents a single cell. Solid lines represent the median. ****$P < 0.0001$; ***$P < 0.001$

by two-sided Wilcoxon rank sum test.

Figure 4. Nucleolar localization of Dbf2-Mob1 is regulated by *CDC5* independently of *CDC15*.

(**A**) Enrichment of Mob1 at the dSPB, in the nucleolus, and Dbf2-Mob1's kinase activity in cells wild-type for *CDC5* (A41211, A41212, and A41213; n = 49, 60 and 47 cells respectively) or harboring a *cdc5-as1* allele (A41334, A41335, and A41336; n = 23, 30 and 28 cells

respectively). Cells were grown at 25°C in SC medium + 2% glucose and 5 μM CMK and

imaged every 3 minutes for 4 hours.

(**B**) Cells harboring *GAL-CDC15(1-750)* and *cdc5-as1* either containing *eGFP-MOB1* (A41211,

A41334, A41376 and A41337; n = 44, 41, 58 and 61 cells respectively), or *eGFP-MOB1Δ78*

(A41212, A41335, A41377, and A41338; n = 54, 30, 12 and 22 cells respectively), or *eGFP-*

MOB1Δ132 (A41213, A41336, A41378 and A41339; n = 62, 28, 26 and 22 cells respectively)

were analyzed to determine Mob1 localization. Localization to the mSPB instead of dSPB was

quantified here because cells expressing *GAL-CDC15(1-750)* often exit from mitosis in the

mother (without movement of a SPB into the bud). For cells exited with two SPBs in the mother

cell, maximum intensities of the two SPBs were used. Cells were grown at 25°C in SC medium +

1% raffinose, 1% galactose and 5 μM CMK and imaged every 5 minutes for 5 hours. Solid lines

represent the average of single cell traces aligned to anaphase onset. Shaded areas represent 95%

confidence intervals. For maximum enrichment, each dot represents a single cell. Solid lines

represent the median. ****P < 0.0001; **P < 0.01; *P < 0.05 by two-sided Wilcoxon rank sum

test.

Figure 5. Cdc5 promotes Dbf2-Mob1's nucleolar localization by phosphorylating Cfi1/Net1.

(**A**) Distribution of all, CDK sites, *CDC15*- and *CDC5*-dependent phosphorylation sites (Table S4) and disordered regions in Cfi1/Net1. For CDK sites, open circles represent sites identified and mutated in Azzam et al. 2004 (*cfi1/net1-6Cdk*) and filled circles represent sites identified in Holt et al. 2009. Dashed boxes denote the four zones.

(**B-E**) Localization of Mob1Δ132 in *CFI1/NET1* (A41411, *n* = 67 cells for B-C and 66 cells for

D-E), *cfi1-91A* mutants (A41412 and A41413, *n* = 36 and 35 cells), *cfi1-Cdc15(z2)* (A41593, *n* =

34 cells) or *cfi1-Cdc5only(z2)* (A41594, *n* = 69 cells). Cells were grown at 25°C in SC medium +

2% glucose and imaged every 3 minutes for 4 hours. Solid lines represent the average of single

cell traces aligned to anaphase onset. Shaded areas represent 95% confidence intervals. For

maximum enrichment, each dot represents a single cell. Solid lines represent the median. *****P

< 0.0001 by two-sided Wilcoxon rank sum test.

(**F**) Streptavidin gel-shift assays to probe the interactions between TurboID-tagged Mob1 and

different *CFI1/NET1* alleles (from left to right: A2587, A41596, A41597, A41598, A41379,

A41611, A41612, A41613). Cells were grown at room temperature in YEP + 2% glucose and

lysates were treated with streptavidin and immunoblotted as indicated. Red arrows highlight

biotinylated proteins.

Figure 6. MEN and Cdc5 promote release of Cdc14 from the nucleolus by phosphorylating Cfi1/Net1.

(**A**) Cdc14 nucleolar release kinetics in wild-type (A41387, n = 134, 123 and 96 cells for each condition), *slk19Δ* (A41410, n = 86 cells), *cdc15-as1* (A41408, n = 38 cells) or *cdc5-as1* mutant (A41409, n = 61 cells). Cells were grown at 25°C in SC medium + 2% glucose with corresponding inhibitors and imaged every 5 minutes for 5 hours. Release of Cdc14 from the nucleolus was quantified as the ratio of fluorescence intensity of Cdc14-eGFP to Cfi1/Net1-mScarlet in the nucleolus (I_{Cdc14}/I_{Cfi1}). Relative degree of Cdc14 release from the nucleolus was calculated as the normalized minimal Cdc14 level in the nucleolus: $(I_{Cdc14}(t_{min})/I_{Cfi1}(t_{min}))/$ $(I_{Cdc14}(t_{-20})/I_{Cfi1}(t_{-20}))$ where t_{min} represents the frame with minimal Cdc14 level in the nucleolus and t_{-20} represent 20 min before movement of the SPB into bud.

(**B**) Cdc14 nucleolar release kinetics in cells harboring wild-type *CFI1/NET1* (A41387, n = 102 and 114 cells) or *CFI1/NET1* phospho-mutants for CDK sites (A41420, n = 95 cells), Cdc15 sites (A41587, n = 104 cells), Cdc5 sites (A441588, n = 86 cells), and Cdc15&Cdc5 sites

(A41589, *n* = 131 cells). Cells were grown at 25°C in SC medium + 2% glucose and imaged

every 5 minutes for 5 hours.

(**C**) Distribution of anaphase duration and relative delay of nucleolar segregation for different

CFI1/NET1 phospho-mutants (A41436, A41590, A41591 and A41592; *n* = 76, 85, 99, 92 cells

respectively) measured using the SPB marker Spc42-eGFP and the nucleolar marker Cfi1/Net1-

mScarlet- I (see Figure S14 for details). Cells were grown at 25°C in SC medium + 2% glucose

and imaged every 3 minutes for 4 hours.

(**D**) Intensities of Cdc14-eGFP at dSPBs in different *CFI1/NET1* phospho-mutant cells (A41387,

A41587, A441588 and A41589; *n* = 80, 82, 77 and 89 cells respectively). Cells were grown and

imaged as in (B).

(**E**) Genetic interactions between different *CFI1/NET1* phospho-mutants and *slk19Δ* (A41645,

A41646, A41647, A41648, A41649) or *spo12Δ* (A41650, A41651, A41652, A41653, A41654)

analyzed by plasmid shuffling (see Methods for details). 5-fold serial dilutions were spotted onto

plates with or without 5'-fluoroorotic acid (5-FOA) and incubated at 25°C for 2 to 3 days. The

presence of 5-FOA selects cells that are viable after losing the *CFI1(URA3/CEN)* plasmid.

For all graphs, single cell traces were aligned to the frame where the dSPB entered the bud and

averaged. Solid lines represent the average. Shaded areas represent 95% confidence intervals.

For distributions, each dot represents a single cell. Solid lines represent the median for (A-B) and

the mean for (C). ****$P < 0.0001$; ***$P < 0.001$; **$P < 0.01$; *$P < 0.05$ by two-sided Wilcoxon

rank sum test.

Figure 7. Phosphorylation of Cfi1/Net1 by Cdc5 and Dbf2-Mob1 promotes mitotic exit.

A model for Cdc14 activation. In metaphase, Cdc5 phosphorylates Cfi1/Net1 in the nucleolus to prepare for Cdc14 release/activation in anaphase. Upon anaphase onset, the FEAR network promotes phosphorylation of Cfi1/Net1 by Clb2-Cdk1 which results in transient release of Cdc14 from the nucleolus. In the meantime, the MEN kinase Cdc15 is activated integrating inputs from both spindle position (via Tem1) and cell cycle progression (via Cdc5 and CDK activities). Activated (SPB-localized) Cdc15 phosphorylates the SPB outer plaque protein Nud1 which creates a dynamic docking site for the MEN terminal kinase complex Dbf2-Mob1 and facilitates phosphorylation and activation of Dbf2-Mob1 by Cdc15. Activated Dbf2-Mob1 gains access to the nucleus and is targeted to the nucleolus by interacting with Cdc5-primed Cfi1/Net1. Nucleolar Dbf2-Mob1 then phosphorylates Cfi1/Net1, keeping Cdc14 dissociated from its nucleolar inhibitor to trigger exit from mitosis. In addition, active Dbf2-Mob1 in the nucleolus and/or cytoplasm phosphorylates Cdc14 at its NLS resulting in cytoplasmic retention of Cdc14 to facilitate cytokinesis.

MATERIALS AND METHODS

Construction of Yeast strains and plasmids

All *Saccharomyces cerevisiae* yeast strains used in this study are derivatives of W303 (A2587) and are listed in Table S1. All plasmids used in this study are listed in Table S2. Yeast were cultured in standard YEP media (1% yeast extract, 2% peptone) with 2% D-glucose, or in standard Synthetic Complete media (SC) with either 2% D-glucose or 2% raffinose prior to induction of *GAL1/10* promoters with 1% raffinose + 1% galactose as indicated in the figure legends. Cells were cultured at 25°C unless noted otherwise.

C-terminal fusions were constructed using standard PCR-based methods. N-terminal truncation of *MOB1* mutants were made by first constructing the truncations with N-terminal eGFP fusion in plasmids (pA2828-2830) based on pRS305H-*MOB1* (König et al., 2010). PCR products of the fusion with 500bp upstream promoter region and 300bp downstream of the *MOB1* ORF as well as the hphMX6 marker were transformed and integrated at the *MOB1* locus. Correct integration and mutations were confirmed by PCR and sequencing. The NLS$_{Cdc14}$ reporter was made by Gibson assembly. NLS$_{Cdc14}$ (Cdc14aa450-551) from pA2725 (Mohl et al., 2009) and synthesized yeast codon optimized miRFP670 (Shcherbakova and Verkhusha, 2013) (denoted as ymiRFP670) were assembled into the p404TEF1 vector (pA2726) which yielded pA2786 for integration at the *trp1* locus. The PhyB constructs were integrated at the *leu2* locus as single integrants (Yang et al., 2013). The eGFP-PIF C-terminal tagging plasmid (pA2821) was made by replacing the mCitrine in pA2721 with eGFP and correcting the frame shift mutation (missing a G at codon#2) in the *natMx6* coding sequence in the original plasmid acquired from Addgene. The TurboID tagging plasmid (pA2859) was constructed by inserting the TurboID-V5 sequence from pA2847 (Branon et al., 2018) into the vector backbone of pA2821 between the linker and natMx6.

Point mutations and truncations in the *GAL-DBF2* plasmids were made using Q5 site-directed mutagenesis (NEB). Single point mutation in *DBF2* (*dbf2-L12A*) was introduced at the endogenous locus using Cas9-mediated gene editing based on previously published protocols (Anand et al., 2017) with modifications. Briefly, BplI cut plasmid pA2911 containing Cas9 and gRNA missing the 20 bp target specific complementary sequence was transformed into A2587 together with (1) a ~250 bp synthesized double stranded DNA (IDT) containing the 20 bp complementary sequence for *DBF2* (TGCTCATATTGCCTGCCAAG) sandwiched with

homologous sequences to the gRNA sequence in plasmid pA2911 for gap repair, and (2) an

80mer single strand DNA harboring *dbf2-L12A* mutation as the repair template. Successfully

edited clones were checked by sequencing and cured of the Cas9 plasmid. Phospho-mutants of

DBF2 (*dbf2-S17,20A* and *dbf2-S17,20D*) were first constructed in plasmids (pA2948, pA2953

and pA2954) by Q5 site-directed mutagenesis. The *DBF2* ORF as well as the *URA3* marker were

then PCR amplified and transformed into A32629 to integrate at the *DBF2* locus as the sole copy

of *DBF2*. Correct integration and mutations were confirmed by PCR and sequencing.

To generate *CFI1/NET1* phospho-mutants, a fusion of wild-type *CFI1/NET1* and

mScarlet-I with ~300 bp upstream and downstream regions of *CFI1/NET1* were first Gibson

assembled into a single-integration vector backbone to generate plasmid pA2898. *CFI1/NET1*

phospho-mutants were synthesized (Twist Bioscience) in two segments (aa25-624 and aa617-

1173) and Gibson assembled into the vector backbone of pA2898 to replace wild-type

CFI1/NET1. These *CFI1/NET1* constructs were integrated into strain A41362 harboring a

heterozygous deletion of *CFI1/NET1* and haploid strains with *cfi1Δ* and different phospho-mutant

forms of *CFI1/NET1* integrated as a single copy at the *leu2* locus as the sole source of

CFI1/NET1 were obtained through tetrad dissection.

Phospho-mutants in the NLS$_{Cdc14}$ reporter were made by first introducing the point

mutations into the integration plasmid (pA2735) with Q5 site-directed mutagenesis and then

transformed and integrated at the *ura3* locus. The triple mutants of *CDC14* (S531,537,546A) was

introduced at the endogenous locus of *CDC14-eGFP* (A34515) using Cas9-mediated gene

editing as described above. The 20 bp complementary sequence for the gRNA used to target

CDC14 was GGAGAGTAACGTCAGGGAGA.

FRAP analysis

FRAP analysis was performed on a DeltaVision-OMX Super-Resolution Microscope (Applied

Precision, GE Healthcare Bio-Sciences) using a 60x oil objective and a 488 nm laser adjusted to

bleach an area of approximately 0.5 μm in radius. Two prebleach images were acquired followed

by a laser pulse (100% intensity) of 0.02 s duration and postbleach images were acquired at 1

s/frame for 30 s. Images at each time points were maximum projections of 7 *z* stacks with 0.5 μm

spacing. Images were analyzed with a custom MATLAB script. After subtracting the

background, fluorescence intensities in the cytosol, $I_{cytosol}(t)$, and at the SPB, $I_{SPB}(t)$ were

measured after segmenting the cell and SPBs. Photobleaching was corrected by normalizing I_{SPB}

(t) with $I_{cytosol}(t)$, $I_{SPB\_norm}(t) = I_{SPB}(t) / I_{cytosol}(t)$. Double normalization for FRAP was calculated

to scale the photobleaching effect between 0 and 1: $I_{SPB\_FRAP}(t) = [I_{SPB\_norm}(t) - I_{SPB\_norm}(0)]$ /

$[I_{SPB\_norm}(pre) - I_{SPB\_norm}(0)]$, where $t = 0$ is the time point (frame) right after photobleaching and

$t = pre$ is the time point (frame) right before photobleaching. The double normalized FRAP

curves were then fitted to a single exponential curve: $y = y_{max}\left(1 - e^{\left(-ln2/t_{1/2}\right)t}\right)$, where y_{max}

is the fraction recovered while $t_{1/2}$ is the half-recovery time.

Microscopy and image analysis

For live-cell microscopy, cells were imaged on agarose pads (2% agarose in SC medium + 2%

glucose, unless otherwise noted) affixed to a glass slide and covered with a coverslip. Imaging

was performed on a DeltaVision Elite platform (GE Healthcare Bio-Sciences) with an InsightSSI

solid state light source, an UltimateFocus hardware autofocus system and a model IX-71,

Olympus microscope controlled by SoftWoRx software. A 60x Plan APO 1.42NA objective and

CoolSNAP HQ2 camera were used for image acquisition. For each time point, 7 *z* sections with

1 μm spacing were collect for each channel and were deconvolved. Maximum projections of the

deconvolved *z* stack were used for fluorescence quantification.

Image analysis was performed with custom scripts in MATLAB. First, yeast cells were

segmented and tracked through time using the bright-field image stacks and a previously

reported algorism (Ricicova et al., 2013). A few modifications were made for the tracking

process. Images were first aligned to correct for drift in the *xy* plane and cell segmentations in the

last frame of the time-lapse series were used as the reference for tracking. Next, fluorescence

images of cell cycle markers (such as Spc42 and or Cfi1/Net1) were segmented and tracked

based on cell segmentation. Appearance of a cell cycle marker in a cell during the acquisition

period was used to identify buds (daughter cells) and cell division events. Tracking of the cell

cycle markers that migrated into the buds were used to identify the corresponding mother cells.

Finally, for each division event identified, localization of Mob1 (or Cdc14) at regions defined by

the cell cycle markers was quantified.

For Mob1 localization at the SPBs (I_{SPB}), maximum intensity of Mob1 at SPBs (dilated

from the SPB area based on segmentation of Spc42) was used given that the size of SPBs (~100

nm) is within the diffraction limit of light microscopy. For Mob1 localization in the nucleolus

($I_{nucleolus}$), the median intensity of Mob1 in the nucleolus as segmented by Cfi1/Net1 was used.

To calculate the relative enrichment of Mob1 at the SPB or in the nucleolus, Mob1 intensities at

these sites as defined above were normalized to the median intensity of Mob1 in the cytosol

which was defined as the cell area with cell cycle marker area subtracted ($I_{cytosol}$):

$I_{SPB\ or\ nucleolus}/I_{cytosol} - 1$. For the Dbf2-Mob1 kinase activity reporter, NLS_{Cdc14}, its

translocation from the nucleus into the cytoplasm was quantified with the coefficient of variation

(CV) of the pixel intensities within the dividing cell. CV is defined as the standard deviation

divided by the mean.

To quantify Cdc14 release from the nucleolus, the ratio of Cdc14 intensity to Cfi1/Net1

intensity (I_{Cdc14}/I_{Cfi1}) was calculated for each pixel within the nucleolus as segmented using

Cfi1/Net1 and averaged.

Single cell traces were aligned based on the timing of anaphase onset or the movement of

a SPB into the bud as indicated in figures, and averaged. 95% confidence intervals were

calculated as $\mu \pm 1.96 * \sigma/\sqrt{n}$, where μ and σ denote the mean and standard deviation

respectively and n is the number of cells measured.

Immunoblot analysis

Log-phase cultures of cells grown in YEP + 2% glucose were harvested and treated with

5% TCA at 4°C overnight. TCA treated cell pellets were washed with acetone, air dried, and

resuspended in lysis buffer (10 mM Tris, 1 mM EDTA, 2.75 mM DTT, pH = 8). Cells were

lysed by bead-beating using a Multivortexer (max speed, 20 minutes) and glass beads at 4°C and

followed by boiling in SDS PAGE protein loading buffer for 5 minutes. Lysates were clarified

by centrifugation and were resolved on a 15-well NuPAGE 4-12% Bis-Tris protein gel (Thermo

Fisher Scientific) prior to transfer onto nitrocellulose membranes. GFP-Mob1 and variants were

detected using an anti-GFP antibody (Clontech, JL-8) at a 1:1000 dilution. Nud1-13myc and

Cfi1/Net1-3myc were detected using an anti-Myc antibody (abcam, 9E10) at a 1:500 dilution.

Mob1-V5-TurboID was detected using an anti-V5 antibody (Invitrogen) at a 1:2000 dilution.

Kar2 was detected using a rabbit anti-Kar2 antiserum at a 1:200,000 dilution. Secondary

antibodies were used at a 1 :10,000 dilution. Blots were imaged using the ECL Plus system (GE

Healthcare).

TurboID-MS and streptavidin gel-shift assay

To identify biotinylated proteins as a result of interaction or physical proximity with TurboID

tagged bait protein by mass spectrometry (MS), log-phase cells with TurboID tagged bait and

untagged control cells were grown in YEP + 2% glucose + 50 µM biotin at room temperature for

3.5 h (~ two doublings). ~40 OD of cells were harvested for each sample and were treated with

5% trichloroacetic acid (TCA) at 4°C for a minimum of 10 minutes. TCA treated cells were

pelleted, washed with 50 mM Tris (pH = 7.5) and acetone, and dried. Dried cell pellets were

resuspended in lysis buffer (50 mM Tris, pH 7.5, 1 mM EDTA, 5 mM DTT, 1 mM PMSF, and

complete mini protease inhibitor cocktail by Roche) and were lysed by bead-beating with chilled

MiniBeadbeater (Biospec) and glass beads for 5 minutes followed by boiling in 1% SDS for 5

minutes. Lysates were diluted with RIPA buffer (50 mM Tris, pH 7.5, 150 mM NaCl, 0.1% SDS,

0.5% sodium deoxycholate, 1% NP40) and clarified by centrifugation. Protein concentration of

the lysates was measured by Bradford assay. 350 µl MyOne streptavidin C1 dynabeads (Thermo

Fisher Scientific) were washed twice with RIPA buffer, incubated with clarified lysates

containing ~ 3 mg of total protein for each sample with rotation for 1 hour at room temperature,

then moved to 4°C and incubated overnight with rotation. On the second day, the supernatants

(flow through) were removed and the beads were washed twice with 1 ml of 0.1 M Na_2CO_3,

once with 1 ml of 2 M urea in 10 mM Tris (pH = 7.5), and twice with 1 ml RIPA buffer. Bound

proteins were eluted by boiling the beads in 30 µl 3x protein loading buffer supplemented with 2

mM biotin. Small aliquots of samples were saved along the process to monitor for the

enrichment for biotinylated proteins by immunoblotting for V5 (included in the TurboID tagged

bait) and biotin.

Eluted proteins were resolved on a 10-well NuPAGE 4-12% Bis-Tris protein gel (Thermo

Fisher Scientific), stained with Coomassie (Imperial Protein Stain, Thermo Fisher Scientific) and

entire gel lanes were excised and cut into 1-mm pieces. Proteins were reduced with 20 mM

dithiothreitol (Sigma) for 1 hour at 56°C and then alkylated with 60 mM iodoacetamide (Sigma)

for 1 hour at 25°C in the dark. Proteins were then digested with 12.5 ng/uL modified trypsin

(Promega) in 50 µL 100 mM ammonium bicarbonate, pH 8.9 at 25°C overnight. Peptides were

extracted by incubating the gel pieces with 5% formic acid in 50% acetonitrile then 100 mM

ammonium bicarbonate, repeated twice followed by incubating the gel pieces with 100%

acetonitrile then 100 mM ammonium bicarbonate, repeated twice. Each fraction was collected,

combined, and reduced to near dryness in a vacuum centrifuge. Peptides were desalted using

C18 SpinTips (Protea, Morgantown, WV).

Peptides were separated by reverse phase HPLC (Thermo Easy nLC1000) using a

precolumn (made in house, 6 cm of 10 µm C18) and a self-pack 5 µm tip analytical column (12

cm of 5 µm C18, New Objective) over a 140-minute gradient before nanoelectrospray using a

QExactive HF-X mass spectrometer (Thermo). Raw mass spectral data files (.raw) were searched

using Proteome Discoverer (Thermo) and Mascot version 2.4.1 (Matrix Science). Mascot search

parameters were: 10 ppm mass tolerance for precursor ions; 15 mmu for fragment ion mass

tolerance; 2 missed cleavages of trypsin; fixed modification was carbamidomethylation of

cysteine; variable modifications were methionine oxidation and lysine biotinylation. Only

peptides with a Mascot score greater than or equal to 25 and an isolation interference less than or

equal to 30 were included in the data analysis.

There are several endogenous biotinylated proteins in yeast. To identify specific proteins

enriched in the TurboID labeling experiments (hits), total peptides identified for each protein in

cells with TurboID tagged bait were compared with the unlabeled control cells to calculate the

ratio of enrichment. A threshold of 10 standard deviations above the average ratio of enrichment

for all proteins in each sample was used to identify hits (see Figure S3B).

For the streptavidin gel-shift assays, cell lysates were prepared as for typical

immunoblotting experiments. Prior to loading samples onto protein gels, 20 ul of lysates for each

sample was incubated with 2 µl of 10 mg/ml streptavidin for 10 min at room temperature with

rotation. Treated samples were then resolved by SDS-PAGE gel at 4°C and processed for

immunoblotting as described above.

PhyB-PIF based optogenetics

The PhyB-PIF based optogenetics experiments were performed based on previous reports (Jost

and Weiner, 2015; Yang et al., 2013). Cells grown to log phase in 1x SC medium + 2% glucose

were incubated with 31.25 µM (0.5 µl of 12.5 mM stock in 200 ul culture) Phycocyanobilin

(PCB, Santa Cruz Biotechnology) for 2 hours at room temperature in the dark. Cells were then

pelleted and resuspended in fresh medium without PCB and were mounted onto an agarose pad

with 2x SC medium + 2% glucose for imaging. To apply the light, we attached one 650 nm and

one 750 nm light-emitting diode (LED, Light-speed Technologies) onto the microscope

condenser. Light was controlled manually during the experiments. For the continuous exposure

of 650 nm light during time-course experiments, the light was briefly turned off for each image

acquisition. To quantify the relative enrichment of target protein in the PhyB-anchored region,

pixel intensities of the target protein (Mob1-eGFP-PIF, I_{PIF}) and PhyB (PhyB-mCherry, I_{PhyB})

inside the cell were fitted to a line ($I_{PIF} = \alpha + \beta I_{PhyB}$). The slope of the fitted line (β) was used

to assess the extent of co-localization or enrichment of the target protein in the PhyB-anchored

region. This method is robust against both photobleaching of the target protein during the time-

course and difference in the shape and size of the anchored region (the nucleolus) at different cell

cycle stages.

Phosphoproteomics

To map phosphorylation events that depend on *CDC5* or *CDC15* (MEN) activity, we used the

analog-sensitive alleles of *CDC5* (*cdc5-as1*) or *CDC15* (*cdc15-as1*) and compared

phosphopeptides with and without inhibiting the kinase in anaphase (and metaphase for Cdc5)

enriched cultures. To enrich for anaphase cells, we first synchronized cells in G1 with α-factor (5

μg/ml for 2.5 hours). α-factor arrested cultures were then washed and released into YEP + 2%

glucose + inhibitor (5 μM CMK for *CDC5/cdc5-as1* or 10 μM 1-NAPP1 for *CDC5/cdc5-as1*) at

room temperature to progress to anaphase (~ 100 minutes). Synchronization and cell cycle stage

were assessed by monitoring budding and spindle length at various time points. Spindles were

visualized by immunofluorescence using an anti-tubulin antibody (Abcam ab6161). Cultures

collected for the anaphase experiment harbored 74% (*WT* + 1-NA-PP1), 95% (*cdc15-as1*, 1-NA-

PP1), 69% (*WT* + CMK), and 94% (*cdc5-as1* + CMK) cells with anaphase spindles.

Phosphopeptide preparation was based on previous methods (Holt et al., 2009; Villén and

Gygi, 2008). ~25 ODs of cells were harvested for each replicate (6 replicates per samples for the

anaphase experiment and 3 replicates per sample for the metaphase experiment) by

centrifugation (5 min at 3000 rpm). Cell pellets were washed with cold 50 mM Tris (pH = 7.5)

and resuspended in cold lysis buffer containing 8 M urea in 50 mM Tris (pH = 7.5), 75 mM

NaCl, 50 mM NaF, 50 mM β-glycerophosphate, 1 mM sodium orthovanadate, 10 mM sodium

pyrophosphate and protease inhibitor cocktail (complete mini, EDTA-free, Roche). Cells were

lysed under denaturing conditions by bead-beating with a chilled MiniBeadbeater (Biospec, two

rounds of 3 cycles of 90 s each) and glass beads. The protein extract was then separated from the

beads and insoluble material by centrifugation. Protein concentration of the lysate was

determined by BCA protein assay (Pierce).

The protein extraction was ultrasonically lysed at 4°C for 2 minutes with six rounds using

a VialTweeter device (Hielscher-Ultrasound Technology). The proteins extracted were then

processed with a precipitation-based digestion protocol using trypsin (Promega) at a ratio of 1:20

as described previously(Collins et al., 2017). About 500 μg of resultant peptides from each

sample were subjected for phosphopeptide enrichment, using the High-Select™ Fe-NTA kit

(Thermo Scientific, A32992) according to the kit instruction, as described previously(Gao et al.,

2019). About 1.5 μg of phosphopeptides enriched from each sample were used for DIA-MS

measurement on an Orbitrap Fusion Lumos Tribrid mass spectrometer (Thermo Scientific)

platform coupled to a nanoelectrospray ion source, as described previously (Mehnert et al.,

2019). DIA-MS data analyses were performed using Spectronaut v13 (Bruderer et al., 2017)

using default setting, by searching against a database of 6632 yeast open reading frames(Hebert

et al., 2014) and the Swiss-prot yeast proteome database. In particular, the PTM localization

option in Spectronaut v13 was enabled to locate phosphorylation sites (Bekker-Jensen et al.,

2020; Collins et al., 2017), with the probability score cutoff >0.75(7), resulting Class-I peptides

to be identified and quantified. Both peptide and protein FDR cutoff (Q value) were controlled at

1%, and the label-free protein quantification of proteome and phospho-proteome was performed

using the default settings in Spectronaut.

To identify peptides whose phosphorylation depended on *CDC5* or *CDC15*, we

calculated the ratio as well as the *p*-value of peptide intensities in *WT* versus *as1* samples. The *p*-

values were calculated using a two-sided Student's *t*-test and were adjusted for false discovery

rate (FDR) of multiple hypothesis testing using the linear step-up procedure (Benjamini-

Hochberg procedure). A threshold of ratio $R > 2$ and $p_{adj} < 0.05$ was used to identify hits.

Additional hits were also included for peptides that were not defected in the *as1* samples but

were detected in at least 5 out of 7 replicates (or 4 out of 6 replicates in the metaphase

experiment) of the *WT* samples. Peptides that were only detected in one replicate of the *as1*

samples but were detected in at least 5 out of 7 replicates (or 4 out of 6 replicates in the

metaphase experiment) of the *WT* samples with a ratio > 2 were also included. After we have

identified hits for peptides, we mapped the peptides and phosphorylation sites in those peptides

to proteins in the yeast proteome. For phosphorylation sites that were detected in multiple

peptides, peptides with a single phosphorylation were given priority. We marked a site as a

strong hit only when all single phosphorylation peptides for the specific site fit our selection

criteria. If a site was only detected in multi-phosphorylation peptides that fit our selection

criteria, we designated that site as a weak hit.

1054

Plasmid shuffling

To assess the genetic interactions between different *CFI1/NET1* phospho-mutants and FEAR

mutants (*slk19Δ* or *spo12Δ*), we constructed strains in the background of *slk19Δ* or *spo12Δ* with

cfi1/net1Δ carrying a *URA3*-based *CEN* plasmid expressing wild-type *CFI1/NET1* (pA2858) and

different *CFI1/NET1* phospho-mutants integrated at the *leu2* locus (A41645-A41654). Growing

these strains on plates with 5'-fluoroorotic acid (5-FOA) selects cells that are viable after losing

the *URA3* plasmid.

1062

1063   ***Data and Code Availability***

1064   The mass spectrometry phosphoproteomics data have been deposited to the ProteomeXchange

1065   Consortium via the PRIDE (Perez-Riverol et al., 2019) partner repository. Custom MATLAB

1066   scripts are available at https://github.com/snow-zhou/Dbf2-Mob1.

REFERENCES

Anand, R., Memisoglu, G., and Haber, J. (2017). Cas9-mediated gene editing in Saccharomyces cerevisiae. Protoc. Exch.

Azzam, R. (2004). Phosphorylation by Cyclin B-Cdk Underlies Release of Mitotic Exit Activator Cdc14 from the Nucleolus. Science *305*, 516–519.

Bardin, A.J., Visintin, R., and Amon, A. (2000). A Mechanism for Coupling Exit from Mitosis to Partitioning of the Nucleus. Cell *102*, 21–31.

Bardin, A.J., Boselli, M.G., and Amon, A. (2003). Mitotic Exit Regulation through Distinct Domains within the Protein Kinase Cdc15. Mol. Cell. Biol. *23*, 5018–5030.

Bardwell, L. (2006). Mechanisms of MAPK signalling specificity. Biochem. Soc. Trans. *34*, 837–841.

Bekker-Jensen, D.B., Bernhardt, O.M., Hogrebe, A., Martinez-Val, A., Verbeke, L., Gandhi, T., Kelstrup, C.D., Reiter, L., and Olsen, J.V. (2020). Rapid and site-specific deep phosphoproteome profiling by data-independent acquisition without the need for spectral libraries. Nat. Commun. *11*, 787.

Bembenek, J., Kang, J., Kurischko, C., Li, B., Raab, J.R., Belanger, K.D., Luca, F.C., and Yu, H. (2005). Crm1-Mediated Nuclear Export of Cdc14 is Required for the Completion of Cytokinesis in Budding Yeast. Cell Cycle *4*, 961–971.

Bishop, A.C., Ubersax, J.A., Petsch, D.T., Matheos, D.P., Gray, N.S., Blethrow, J., Shimizu, E., Tsien, J.Z., Schultz, P.G., Rose, M.D., et al. (2000). A chemical switch for inhibitor-sensitive alleles of any protein kinase. Nature *407*, 395–401.

Branon, T.C., Bosch, J.A., Sanchez, A.D., Udeshi, N.D., Svinkina, T., Carr, S.A., Feldman, J.L., Perrimon, N., and Ting, A.Y. (2018). Efficient proximity labeling in living cells and organisms with TurboID. Nat. Biotechnol. *36*, 880–887.

Britschgi, A., Duss, S., Kim, S., Couto, J.P., Brinkhaus, H., Koren, S., De Silva, D., Mertz, K.D., Kaup, D., Varga, Z., et al. (2017). The Hippo kinases LATS1 and 2 control human breast cell fate via crosstalk with ERα. Nature *541*, 541–545.

Bruderer, R., Bernhardt, O.M., Gandhi, T., Xuan, Y., Sondermann, J., Schmidt, M., Gomez-Varela, D., and Reiter, L. (2017). Optimization of Experimental Parameters in Data-Independent Mass Spectrometry Significantly Increases Depth and Reproducibility of Results. Mol. Cell. Proteomics MCP *16*, 2296–2309.

Campbell, I.W., Zhou, X., and Amon, A. (2019). The Mitotic Exit Network integrates temporal and spatial signals by distributing regulation across multiple components. ELife *8*.

Cargnello, M., and Roux, P.P. (2011). Activation and Function of the MAPKs and Their Substrates, the MAPK-Activated Protein Kinases. Microbiol. Mol. Biol. Rev. *75*, 50–83.

Carpenter, K., Bell, R.B., Yunus, J., Amon, A., and Berchowitz, L.E. (2018). Phosphorylation-Mediated Clearance of Amyloid-like Assemblies in Meiosis. Dev. Cell *45*, 392-405.e6.

Cheng, L., Hunke, L., and Hardy, C.F.J. (1998). Cell Cycle Regulation of the Saccharomyces cerevisiae Polo-Like Kinase Cdc5p. Mol. Cell. Biol. *18*, 7360–7370.

Collins, B.C., Hunter, C.L., Liu, Y., Schilling, B., Rosenberger, G., Bader, S.L., Chan, D.W., Gibson, B.W., Gingras, A.-C., Held, J.M., et al. (2017). Multi-laboratory assessment of reproducibility, qualitative and quantitative performance of SWATH-mass spectrometry. Nat. Commun. *8*, 291.

D'Amours, D., Stegmeier, F., and Amon, A. (2004). Cdc14 and Condensin Control the Dissolution of Cohesin-Independent Chromosome Linkages at Repeated DNA. Cell *117*, 455–469.

D'Aquino, K.E., Monje-Casas, F., Paulson, J., Reiser, V., Charles, G.M., Lai, L., Shokat, K.M., and Amon, A. (2005). The Protein Kinase Kin4 Inhibits Exit from Mitosis in Response to Spindle Position Defects. Mol. Cell *19*, 223–234.

Elia, A.E.H., Rellos, P., Haire, L.F., Chao, J.W., Ivins, F.J., Hoepker, K., Mohammad, D., Cantley, L.C., Smerdon, S.J., and Yaffe, M.B. (2003). The Molecular Basis for Phosphodependent Substrate Targeting and Regulation of Plks by the Polo-Box Domain. Cell *115*, 83–95.

Fairhead, M., and Howarth, M. (2015). Site-specific biotinylation of purified proteins using BirA. Methods Mol. Biol. Clifton NJ *1266*, 171–184.

Feric, M., Vaidya, N., Harmon, T.S., Mitrea, D.M., Zhu, L., Richardson, T.M., Kriwacki, R.W., Pappu, R.V., and Brangwynne, C.P. (2016). Coexisting Liquid Phases Underlie Nucleolar Subcompartments. Cell *165*, 1686–1697.

Gao, Q., Zhu, H., Dong, L., Shi, W., Chen, R., Song, Z., Huang, C., Li, J., Dong, X., Zhou, Y., et al. (2019). Integrated Proteogenomic Characterization of HBV-Related Hepatocellular Carcinoma. Cell *179*, 1240.

Geymonat, M., Spanos, A., de Bettignies, G., and Sedgwick, S.G. (2009). Lte1 contributes to Bfa1 localization rather than stimulating nucleotide exchange by Tem1. J. Cell Biol. *187*, 497–511.

Good, M.C., Zalatan, J.G., and Lim, W.A. (2011). Scaffold proteins: hubs for controlling the flow of cellular information. Science *332*, 680–686.

Guertin, D.A., Trautmann, S., and McCollum, D. (2002). Cytokinesis in Eukaryotes. Microbiol. Mol. Biol. Rev. *66*, 155–178.

Hebert, A.S., Richards, A.L., Bailey, D.J., Ulbrich, A., Coughlin, E.E., Westphall, M.S., and Coon, J.J. (2014). The one hour yeast proteome. Mol. Cell. Proteomics MCP *13*, 339–347.

Hergovich, A., and Hemmings, B.A. (2012). Hippo signalling in the G2/M cell cycle phase: Lessons learned from the yeast MEN and SIN pathways. Semin. Cell Dev. Biol. *23*, 794–802.

Holt, L.J., Tuch, B.B., Villen, J., Johnson, A.D., Gygi, S.P., and Morgan, D.O. (2009). Global Analysis of Cdk1 Substrate Phosphorylation Sites Provides Insights into Evolution. Science *325*, 1682–1686.

Housley, M.P., Reischauer, S., Dieu, M., Raes, M., Stainier, D.Y.R., and Vanhollebeke, B. (2014). Translational profiling through biotinylation of tagged ribosomes in zebrafish. Development *141*, 3988–3993.

Jaspersen, S.L., and Morgan, D.O. (2000). Cdc14 activates Cdc15 to promote mitotic exit in budding yeast. Curr. Biol. *10*, 615–618.

Jost, A.P.-T., and Weiner, O.D. (2015). Probing Yeast Polarity with Acute, Reversible, Optogenetic Inhibition of Protein Function. ACS Synth. Biol. *4*, 1077–1085.

Kettenbach, A.N., Schweppe, D.K., Faherty, B.K., Pechenick, D., Pletnev, A.A., and Gerber, S.A. (2011). Quantitative Phosphoproteomics Identifies Substrates and Functional Modules of Aurora and Polo-Like Kinase Activities in Mitotic Cells. Sci. Signal. *4*, rs5–rs5.

Kırlı, K., Karaca, S., Dehne, H.J., Samwer, M., Pan, K.T., Lenz, C., Urlaub, H., and Görlich, D. (2015). A deep proteomics perspective on CRM1-mediated nuclear export and nucleocytoplasmic partitioning. ELife *4*, e11466.

Knockenhauer, K.E., and Schwartz, T.U. (2016). The Nuclear Pore Complex as a Flexible and Dynamic Gate. Cell *164*, 1162–1171.

König, C., Maekawa, H., and Schiebel, E. (2010). Mutual regulation of cyclin-dependent kinase and the mitotic exit network. J. Cell Biol. *188*, 351–368.

Kuilman, T., Maiolica, A., Godfrey, M., Scheidel, N., Aebersold, R., and Uhlmann, F. (2015). Identification of Cdk targets that control cytokinesis. EMBO J. *34*, 81–96.

Lee, K.S., Park, J.-E., Asano, S., and Park, C.J. (2005). Yeast polo-like kinases: functionally conserved multitask mitotic regulators. Oncogene *24*, 217–229.

Li, W., Cooper, J., Zhou, L., Yang, C., Erdjument-Bromage, H., Zagzag, D., Snuderl, M., Ladanyi, M., Hanemann, C.O., Zhou, P., et al. (2014). Merlin/NF2 loss-driven tumorigenesis linked to CRL4(DCAF1)-mediated inhibition of the hippo pathway kinases Lats1 and 2 in the nucleus. Cancer Cell *26*, 48–60.

Loughrey Chen, S., Huddleston, M.J., Shou, W., Deshaies, R.J., Annan, R.S., and Carr, S.A. (2002). Mass Spectrometry-based Methods for Phosphorylation Site Mapping of Hyperphosphorylated Proteins Applied to Net1, a Regulator of Exit from Mitosis in Yeast. Mol. Cell. Proteomics *1*, 186–196.

Lowery, D.M., Lim, D., and Yaffe, M.B. (2005). Structure and function of Polo-like kinases. Oncogene *24*, 248–259.

Mah, A.S., Jang, J., and Deshaies, R.J. (2001). Protein kinase Cdc15 activates the Dbf2-Mob1 kinase complex. Proc. Natl. Acad. Sci. *98*, 7325–7330.

Mah, A.S., Elia, A.E.H., Devgan, G., Ptacek, J., Schutkowski, M., Snyder, M., Yaffe, M.B., and Deshaies, R.J. (2005). Substrate specificity analysis of protein kinase complex Dbf2-Mob1 by peptide library and proteome array screening. BMC Biochem. *6*, 22.

Mehnert, M., Li, W., Wu, C., Salovska, B., and Liu, Y. (2019). Combining Rapid Data Independent Acquisition and CRISPR Gene Deletion for Studying Potential Protein Functions: A Case of HMGN1. Proteomics *19*, e1800438.

Meitinger, F., Boehm, M.E., Hofmann, A., Hub, B., Zentgraf, H., Lehmann, W.D., and Pereira, G. (2011). Phosphorylation-dependent regulation of the F-BAR protein Hof1 during cytokinesis. Genes Dev. *25*, 875–888.

Mohl, D.A., Huddleston, M.J., Collingwood, T.S., Annan, R.S., and Deshaies, R.J. (2009). Dbf2–Mob1 drives relocalization of protein phosphatase Cdc14 to the cytoplasm during exit from mitosis. J. Cell Biol. *184*, 527–539.

Murphy, L.O., Smith, S., Chen, R.-H., Fingar, D.C., and Blenis, J. (2002). Molecular interpretation of ERK signal duration by immediate early gene products. Nat. Cell Biol. *4*, 556–564.

Nakajima, H., Toyoshima-Morimoto, F., Taniguchi, E., and Nishida, E. (2003). Identification of a Consensus Motif for Plk (Polo-like Kinase) Phosphorylation Reveals Myt1 as a Plk1 Substrate*. J. Biol. Chem. *278*, 25277–25280.

Neville, M. (1999). The NES-Crm1p export pathway is not a major mRNA export route in Saccharomyces cerevisiae. EMBO J. *18*, 3746–3756.

Owen, I., and Shewmaker, F. (2019). The Role of Post-Translational Modifications in the Phase Transitions of Intrinsically Disordered Proteins. Int. J. Mol. Sci. *20*, 5501.

Pereira, G., Höfken, T., Grindlay, J., Manson, C., and Schiebel, E. (2000). The Bub2p Spindle Checkpoint Links Nuclear Migration with Mitotic Exit. Mol. Cell *6*, 1–10.

Perez-Riverol, Y., Csordas, A., Bai, J., Bernal-Llinares, M., Hewapathirana, S., Kundu, D.J., Inuganti, A., Griss, J., Mayer, G., Eisenacher, M., et al. (2019). The PRIDE database and related tools and resources in 2019: improving support for quantification data. Nucleic Acids Res. *47*, D442–D450.

Queralt, E., Lehane, C., Novak, B., and Uhlmann, F. (2006). Downregulation of PP2ACdc55 Phosphatase by Separase Initiates Mitotic Exit in Budding Yeast. Cell *125*, 719–732.

Ricicova, M., Hamidi, M., Quiring, A., Niemisto, A., Emberly, E., and Hansen, C.L. (2013). Dissecting genealogy and cell cycle as sources of cell-to-cell variability in MAPK signaling using high-throughput lineage tracking. Proc. Natl. Acad. Sci. *110*, 11403–11408.

Rock, J.M., and Amon, A. (2009). The FEAR network. Curr. Biol. CB *19*, R1063-1068.

Rock, J.M., and Amon, A. (2011). Cdc15 integrates Tem1 GTPase-mediated spatial signals with Polo kinase-mediated temporal cues to activate mitotic exit. Genes Dev. *25*, 1943–1954.

Rock, J.M., Lim, D., Stach, L., Ogrodowicz, R.W., Keck, J.M., Jones, M.H., Wong, C.C.L., Yates, J.R., Winey, M., Smerdon, S.J., et al. (2013). Activation of the Yeast Hippo Pathway by Phosphorylation-Dependent Assembly of Signaling Complexes. Science *340*, 871–875.

Romero, P., Obradovic, Z., Kissinger, C., Villafranca, J.E., and Dunker, A.K. (1997). Identifying disordered regions in proteins from amino acid sequence. In Proceedings of International Conference on Neural Networks (ICNN'97), (Houston, TX, USA: IEEE), pp. 90–95.

Sanchez, Y. (1999). Control of the DNA Damage Checkpoint by Chk1 and Rad53 Protein Kinases Through Distinct Mechanisms. Science *286*, 1166–1171.

Shcherbakova, D.M., and Verkhusha, V.V. (2013). Near-infrared fluorescent proteins for multicolor in vivo imaging. Nat. Methods *10*, 751–754.

Shin, Y., and Brangwynne, C.P. (2017). Liquid phase condensation in cell physiology and disease. Science *357*, eaaf4382.

Shou, W., Seol, J.H., Shevchenko, A., Baskerville, C., Moazed, D., Chen, Z.W.S., Jang, J., Shevchenko, A., Charbonneau, H., and Deshaies, R.J. (1999). Exit from Mitosis Is Triggered by Tem1-Dependent Release of the Protein Phosphatase Cdc14 from Nucleolar RENT Complex. Cell *97*, 233–244.

Shou, W., Azzam, R., Chen, S.L., Huddleston, M.J., Baskerville, C., Charbonneau, H., Annan, R.S., Carr, S.A., and Deshaies, R.J. (2002). Cdc5 influences phosphorylation of Net1 and disassembly of the RENT complex. BMC Mol. Biol. 14.

Stegmeier, F., and Amon, A. (2004). Closing Mitosis: The Functions of the Cdc14 Phosphatase and Its Regulation. Annu. Rev. Genet. *38*, 203–232.

Stegmeier, F., Visintin, R., and Amon, A. (2002). Separase, Polo Kinase, the Kinetochore Protein Slk19, and Spo12 Function in a Network that Controls Cdc14 Localization during Early Anaphase. Cell *108*, 207–220.

Stoepel, J., Ottey, M.A., Kurischko, C., Hieter, P., and Luca, F.C. (2005). The Mitotic Exit Network Mob1p-Dbf2p Kinase Complex Localizes to the Nucleus and Regulates Passenger Protein Localization□D □V. Mol. Biol. Cell *16*, 15.

Sullivan, M., Higuchi, T., Katis, V.L., and Uhlmann, F. (2004). Cdc14 Phosphatase Induces rDNA Condensation and Resolves Cohesin-Independent Cohesion during Budding Yeast Anaphase. Cell *117*, 471–482.

Swaney, D.L., Beltrao, P., Starita, L., Guo, A., Rush, J., Fields, S., Krogan, N.J., and Villén, J. (2013). Global analysis of phosphorylation and ubiquitylation cross-talk in protein degradation. Nat. Methods *10*, 676–682.

Torres-Rosell, J., Machín, F., Jarmuz, A., and Aragón, L. (2004). Nucleolar Segregation Lags Behind the Rest of the Genome and Requires Cdc14p Activation by the FEAR Network. Cell Cycle *3*, 494–500.

Villén, J., and Gygi, S.P. (2008). The SCX/IMAC enrichment approach for global phosphorylation analysis by mass spectrometry. Nat. Protoc. *3*, 1630–1638.

Visintin, R., and Amon, A. (2001). Regulation of the mitotic exit protein kinases Cdc15 and Dbf2. Mol. Biol. Cell *12*, 2961–2974.

Visintin, C., Tomson, B.N., Rahal, R., Paulson, J., Cohen, M., Taunton, J., Amon, A., and Visintin, R. (2008). APC/C-Cdh1-mediated degradation of the Polo kinase Cdc5 promotes the return of Cdc14 into the nucleolus. Genes Dev. *22*, 79–90.

Visintin, R., Hwang, E.S., and Amon, A. (1999). Cfi1 prevents premature exit from mitosis by anchoring Cdc14 phosphatase in the nucleolus. *398*, 6.

Yang, X., Jost, A.P.-T., Weiner, O.D., and Tang, C. (2013). A light-inducible organelle-targeting system for dynamically activating and inactivating signaling in budding yeast. Mol. Biol. Cell *24*, 2419–2430.

Yu, F.-X., and Guan, K.-L. (2013). The Hippo pathway: regulators and regulations. Genes Dev. *27*, 355–371.

SUPPLEMENTARY DATA

5 µm

Figure S1. N-terminally truncated Mob1 is hyperactive, Related to Figure 1

(**A**) Immunoblot (top) and quantification (bottom) of untagged (A2587), full-length (A41351) and truncated GFP-Mob1 (A41352, A41353) as well as full-length GFP-Mob1 expressed from the *pGPD/TDH3* promoter (A41350).

(**B**) 5-fold serial dilutions of *cdc15-2* (A41424, A41425, A41426, A41427), *tem1-3* (A41429, A41430, A41431, A41428) and *cdc5-1* (A41432, A41433, A41434, A41435) harboring the indicated *MOB1* constructs in YEP + 2% glucose at the indicated temperatures.

(**C**) Localization of GFP-Mob1 expressed under the control of *pGPD* promoter (A41595) during the cell cycle. Cells were grown and imaged as in Figure 1E. Increased nuclear but not nucleolar localization of Mob1 was observed.

Figure S2. Mob1's nucleolar localization correlates with MEN activation and Cdc14 release from the nucleolus, Related to Figure 1

(**A**) Illustration of Mob1's SPB localization and translocation of NLS$_{Cdc14}$ reporter in response to MEN activation.

(**B**) Localization of Mob1Δ132 and the MEN activity reporter NLS$_{Cdc14}$ during the cell cycle. A41213 (*GFP-MOB1Δ132, CFI1/NET1-mCherry, SPC42-mCherry* and *NLS$_{Cdc14}$-ymiRFP670*) cells were grown as in Figure 1E. Red square indicates anaphase.

(**C**) Relative timing of nucleolar localization of Mob1 (black) and SPB localization of Mob1 (green) in cells harboring the indicated *MOB1* alleles.

(**D**) Relative timing of nucleolar localization of Mob1 (black) and release of the NLS_{Cdc14} reporter from the nucleus (red) in cells harboring the indicated *MOB1* alleles.

(**E**) Localization of Mob1 and Cdc14 during the cell cycle. A40314 (*MOB1-eGFP*, *SPC42-mCherry* and *CDC14-mCherry*) cells were grown at room temperature in SC medium + 2% glucose and imaged every 2 minutes for 4 hours. The red square indicates the frames where Mob1 localizes to the nucleolus and Cdc14 is fully released from the nucleolus.

Figure S3. TurboID identified interaction between Dbf2-Mob1 and Cfi1/Net1, Related to Figure 2

(**A**) Overview of TurboID proximity-based biotinylation to identify interaction partners of a target protein.

(**B**) Number of total peptides detected for MS-identified proteins in cells with TurboID tagged baits relative to untagged control cells (source data for Figure 2A). Shaded red dots represent proteins (hits) identified as interaction partners of the bait protein. Solid yellow lines denote the mean ratio of total peptide detected for all proteins. Dashed lines denote the threshold used for identifying hits (10 standard derivations above the mean).

(**C**) Biotinaylated peptide of Cfi1/Net1 detected in cells harboring Mob1-TurboID.

(**D**) Streptavidin gel-shift assay to probe the interaction between Mob1 and Cfi1/Net1. Lysates of A41379, A1638, and A41372 cells with or without streptavidin treatment were immunoblotted as indicated. Red arrows highlight biotinylated proteins.

Figure S4. Examining nuclear access of Dbf2-Mob1 by optogenetics, Related to Figure 3

(**A**) The PhyB-PIF based light-inducible organelle targeting system. When exposed to 650 nm light, anchored PhyB interacts with PIF and thus targets protein of interest (POI) to designated subcellular regions. This interaction is reversed with exposure to 750 nm light.

(**B**) Recruiting Mob1 to SPBs (A40346) or the mitochondrial surface (A40354) with the PhyB-PIF optogenetics system. Cells were grown at room temperature in SC medium + 2% glucose, incubated with 12.5 μM PCB for 2 hours in the dark prior to imaging. Red dots indicate application of red light (650 nm) for 5 minutes to activate PhyB.

(**C**) Recruiting Dbf2 (A40262) or Cdc15 (A40258) to the nucleolus with PhyB-Sik1. Cells were grown similarly to (B). Yellow arrows highlight the light-induced recruitment.

(**D**) Probing Mob1's nuclear access during the cell cycle with PhyB-Sik1. A40260 cells were grown similarly to (B) and imaged every 5 minutes while the red light (650 nm) was applied for 2 minutes every 15 minutes. Red dots denote frames where 650 nm light was applied to activate PhyB prior to imaging as indicated in the cartoon above. Yellow arrows highlight the light-induced recruitment.

(**E**) Quantifications of enrichment of Mob1 in the nucleolus as a function of PhyB activation time for wild-type Mob1 (A41360; $n = 22$, 14, and 21 cells for G1-S, metaphase and anaphase respectively), Mob1Δ78 (A41366; $n = 11$, 6, and 2 cells for G1-S, metaphase and anaphase respectively) at 25°C and for Mob1Δ78 in *CDC15* (A41366; $n = 25$, 4, and 9 cells for G1-S, metaphase and anaphase respectively) or *cdc15-2* cells (A41365; $n = 39$, 9, and 10 cells for G1-S, metaphase and anaphase respectively) at 34°C. Cells were grown similarly to (B).

Figure S5. Identification of a functional NES in Dbf2, Related to Figure 3

(**A**) Mob1's cellular localization when nuclear export is inhibited. Left, cells (A39893, A41349) were grown at room temperature in SC medium + 2% glucose and leptomycin B. Red box indicates anaphase and yellow arrows highlight increased nuclear and nucleolar localization. Right, comparison of nucleolar enrichment of Mob1 in *CRM1* (A41211, A41212 and A41213; *n* = 12, 18 and 11 cells respectively) and *crm1T539C* cells (A41373, A41374 and A41375; *n* = 13,

9 and 16 cells respectively). Cells were grown at 25°C in SC medium + 2% glucose with 100 ng/ml LMB and imaged every 3 minutes for 4 hours.

(**B**) Localization of overexpressed Mob1-eGFP from the *GAL1-10* promoter with (A41363) or without (A41364) co-overexpressing Dbf2 from the same promoter. Cells were grown in SC medium + 2% raffinose and mounted onto agarose pads made with SC medium + 1% raffinose and 1% galactose.

(**C**) Localization of overexpressed Dbf2-eGFP with (A41383) or without (A41384) inhibiting Crm1 with LMB. Cells were grown in SC medium + 2% raffinose and mounted onto agarose pad made with SC medium + 1% raffinose, 1% galactose and 100 ng/ml LMB.

(**D**) Alignment of Dbf2 homologs in *Saccharomycetes* and *S. pombe*.

(**E-G**) Localization of various Dbf2-eGFP mutants expressed from the *GAL1-10* promoter (A41388, A41389, A41386, A41390, A41391, A41392, A41393, A41440, A41441 and A41442; $n = 97, 98, 123, 135, 120, 88, 58, 76$ and 79 cells respectively). Cells were grown as in (B) and analyzed after 5 hours of growth in galactose containing medium.

Figure S6. Phosphorylation of Dbf2's NES partially regulates Dbf2-Mob1's nuclear access, Related to Figure 3

(**A**) 5-fold serial dilutions of *cdc15-2* cells harboring wild-type *DBF2* (A41624), *dbf2-S17/20A* (A41625) or *dbf2-S17/20D* (A41626) in YEP + 2% glucose at the indicated temperatures.

(**B**) Nucleolar enrichment of wild-type and truncated Mob1 in wild-type, *dbf2-S17/20A* or *dbf2-S17/20D* cells (A41614, A41617, A41620, A41615, A41618, A41621, A41616, A41619, A41622; *n* = 32, 28, 33,39, 43, 37, 31, 35 and 37 cells respectively). Cells were grown and imaged as in Figure 1E. Solid lines represent the average, shaded areas represent 95% confidence intervals. For maximum enrichment, each dot represents a single cell. Solid lines represent the median. ****P < 0.0001; *P < 0.05 by two-sided Wilcoxon rank sum test.

(**C**) Quantifications of enrichment of Mob1 in the nucleolus as a function of PhyB activation time for wild-type *DBF2* (A41608; *n* = 42 and 24 cells for pre-anaphase and anaphase, respectively), *dbf2-S17/20A* (A41609; *n* = 55 and 41 cells for pre-anaphase and anaphase, respectively) and *dbf2-S17/20D* (A41610; *n* = 58 and 28 cells for pre-anaphase and anaphase, respectively). Cells were grown as in Figure S4B.

Figure S7. Nucleolar localization of Dbf2-Mob1 does not depend on the FEAR network, Related to Figure 4

(**A**) Enrichment of Mob1 at the dSPB, in the nucleolus, and Dbf2-Mob1's kinase activity in wild-type (A41211, A41212 and A41213; n = 12, 18, 11) or *slk19Δ* (A41357, A41358, A41359; n = 26, 29, 22 cells) cells. Cells were grown at 25°C in SC medium + 2% glucose and imaged every 3 minutes for 4 hours.

(**B**) Enrichment of Mob1 at the dSPB, in the nucleolus, and Dbf2-Mob1's kinase activity in wild-type (A41212 and A41213; n = 58 and 66 cells) or *cdc14-3* (A41355 and A41356; n = 15 and 25 cells) cells. Cells were grown at 34°C in SC medium + 2% glucose and imaged every 3 minutes for 4 hours. For all graphs, solid lines represent the average of single cell traces aligned to anaphase onset. Shaded areas represent 95% confidence intervals. For maximum enrichment, each dot represents a single cell. The solid lines represent the median. *P < 0.05; *ns*, not significant (P > 0.05) by two-sided Wilcoxon rank sum test.

Figure S8. Phosphoproteomics identifies *CDC15* (MEN) and *CDC5* dependent phosphorylation in anaphase, Related to Figure 5

(**A**) Overview of the sample preparation for phosphoproteomics analysis to map *CDC15* (MEN) and *CDC5* dependent phosphorylation in anaphase. Paired wild-type (*WT*, A2587) and *cdc15-as1* (A10991) or paired wild-type and *cdc5-as1* (A40903) cultures were synchronized in G1 with

α-factor (5 µg/ml). After 150 minutes cells were released into fresh medium with the corresponding inhibitors in the absence of pheromone at room temperature. After ~100 minutes cells were harvested to extract proteins and processed for data-independent acquisition mass spectrometry (DIA-MS) analysis with 7 technical replicates for each sample. Immuno-fluorescence (IF) using an anti-tubulin antibody was performed on the collected cells to determine the percentage of cells with anaphase spindle (~70% for wild-type cells and ~95% for cdc5-as1 and cdc15-as1 cells).

(**B**) Volcano plots of -$\log_{10}$ transformed FDR adjusted *P*-value versus $\log_2$(fold change or ratio) of intensities measured for peptides identified in anaphase cells with *WT* or analog-sensitive (*as1*) alleles of *CDC15* or *CDC5*. Yellow dashed lines indicate the cutoff ($R > 2$ and $p_{adj} < 0.05$) used to identify peptides whose phosphorylation depends on the corresponding kinase as marked with red shaded dots.

(**C**) Correlations of peptide intensity in *WT* and *as1* samples for Cdc15 and Cdc5 inhibition in anaphase. Data points on the axis represent peptides that were only detected in one sample but not the other. Red shaded dots denote hits for *CDC15* or *CDC5*-dependent phosphopeptides identified based on the cutoff described in (B) and for peptides that were detected in at least 5 out of 7 replicates in *WT* samples but were missing in *as1* samples (thus no fold change could be calculated).

(**D-E**) Summary of phosphorylation sites (D) and phospho-proteins (E) determined as *CDC15*- or *CDC5*-dependent. Potential Dbf2-Mob1 targets were identified as *CDC15*-dependent and to fit the Dbf2 phosphorylation consensus motif RXXS*, where * denotes the site of phosphorylation. Cdc5-only sites are sites that are *CDC5*-dependent but not *CDC15*-dependent.

(**F**) *CDC15*- and *CDC5*-dependent sites in Mob1 and Dbf2. Light gray sites represent sites that were detected but were not determined as either *CDC15*- or *CDC5*-dependent.

Figure S9. Phosphorylation of Cfi1/Net1 modulates Dbf2-Mob1's nucleolar localization, Related to Figure 5

(**A-B**) Representative images of Mob1Δ132 localization in *CFI1/NET1* (A41411), *cfi1-91A* (A41412 and A41413), *cfi1-Cdc15(z2)* (A41593) or *cfi1-Cdc5only(z2)* (A41594) cells.

Figure S10. Identification of *CDC5*-dependent phosphorylation sites in metaphase, Related to Figure 5

(**A**) Overview of the sample preparation for phosphoproteomics analysis to map *CDC5*-dependent phosphorylation sites in metaphase. Same procedures were followed as in Figure S8A except that following release of cells from the G1 arrest, cells were resuspended in medium containing the inhibitor CMK (5 μM) and nocodazole (15 μg/ml).

(**B**) Volcano plot and correlation of peptide intensities with and without inhibition of Cdc5 (*cdc5-as1* or *WT*) in metaphase. Yellow dashed lines indicate the cutoff ($R > 2$ and $P_{adj} < 0.05$, or detected in at least 4 out of 6 replicates in *WT* sample but missing in *cdc5as1* sample) used to identify peptides whose phosphorylation depends on *CDC5* as marked with red shaded dots.

(**C-D**) Summary and comparison of phosphorylation sites (C) and phospho-proteins (D) detected in metaphase and anaphase.

(**E**) *CDC5*-only and *CDC15*-dependent sites in Cfi1/Net1 and Cdc14 mapped in metaphase and anaphase. Light gray sites in metaphase/anaphase represent phosphorylation sites that were detected but were not determined as *CDC5*-dependent.

A

# of peptides detected		Bait (TurboID tagged)	
		Cdc5	Untagged
	Cfi1	164	6
Hits	Boi2	21	0
	Mps3	20	0
	Csa1	17	0

B

                        618
Cfi1 IVPQDSDSSSFPkSDLFK
                      ↑
                    biotin

                              623
Cfi1 IVPQDSDSSSFPKSDLFkMIEGDDTDLPQWFK
                               ↑
                             biotin

C

+ Streptavidin

Cdc5-TurboID-V5   -   +
Cfi1-3myc   +   +

α-myc

α-V5

Kar2

Figure S11. Cdc5 interacts with Cfi1/Net1 *in vivo*, Related to Figure 5

(**A**) Results of TurboID proximity-based biotinylation for Cdc5 (A41385 and A2588).

(**B**) Biotinaylated peptides of Cfi1/Net1 detected in cells with Cdc5-TurboID.

(**C**) Streptavidin gel-shift assay to probe the interaction between Cdc5-TurboID and Cfi1/Net1 (A1638 and A41418). Cells were grown at room temperature in YPED. Lysates were treated with streptavidin and immunoblotted as indicated. Red arrows highlight biotinylated proteins.

Figure S12. Phosphorylation in zone 2 of Cfi1/Net1 regulates Cdc14 release from the nucleolus, Related to Figure 6

(**A-B**) Kinetics of Cdc14 release from the nucleolus in cells harboring wild-type *CFI1/NET1* (A41387, *n* = 103 cells for A, *n* = 130 cells for B) or different *CFI1/NET1* phospho-null mutants (A41398, A41399, A41400, A41401, A41404, A41405, A41397, A41402 and A41403; *n* = 95, 103, 146, 59, 102, 113, 128, 114 and 102 cells respectively). Cells were grown at 25°C in SC medium + 2% glucose and imaged every 5 minutes for 5 hours. Release of Cdc14 from the nucleolus was quantified as the ratio of fluorescence intensity of Cdc14-eGFP to Cfi1/Net1-mScarlet in the nucleolus (I_{Cdc14}/I_{Cfi1}). Relative degree of Cdc14 release from the nucleolus was calculated as the normalized minimal Cdc14 level in the nucleolus: $(I_{Cdc14}(t_{min})/I_{Cfi1}(t_{min}))/$ $(I_{Cdc14}(t_{-20})/I_{Cfi1}(t_{-20}))$ where t_{min} represents the frame with minimal Cdc14 levels in the nucleolus and t_{-20} represent 20 min before movement of the SPB into bud. Each dot represents a single cell and the solid lines represent the median. ****$P < 0.0001$; **$P < 0.01$; *$P < 0.05$ by two-sided Wilcoxon rank sum test.

(**C**) Representative images showing Cdc14 nucleolar release kinetics for cells harboring wild-type *CFI1/NET1* (A41387) or *CFI1/NET1* phospho-mutants with all 91 sites (A41404) or only sites in zone 2 (A41399) mutated to alanine. Yellow arrows highlight localization of Cdc14 at dSPB prior to anaphase.

Figure S13. The MEN and *CDC5* promote release of Cdc14 from the nucleolus by phosphorylating Cfi1/Net1, Related to Figure 6

Representative images showing Cdc14 nucleolar release kinetics for cells harboring wild-type *CFI1/NET1* (A41387) or *CFI1/NET1* phospho-mutants for Cdc15 sites (A41587), Cdc5 sites (A441588), and Cdc15&Cdc5 sites (A41589). Red box highlights anaphase when Cdc14 is fully released in WT cells. Yellow arrows highlight localization of Cdc14 at dSPB prior to anaphase.

Figure S14. Cfi1/Net1 phospho-mutants delay mitotic exit and nucleolar segregation, Related to Figure 6

(**A**) Tracking anaphase progression and nucleolar segregation using Spc42-eGFP and Cfi1/Net1-mScarlet-I as markers, respectively. A41436 cells were grown at 25°C in SC medium + 2% glucose and imaged every 3 minutes for 4 hours.

(**B**) Quantification of spindle elongation (spindle length was estimated by measuring the distance between the two SPBs) and nucleolar segregation (nucleoli length was estimated by measuring

the length of the major axis of the nucleolar mass) for *CFI1/NET1* mutants using the SPB marker Spc42-eGFP and the nucleolar marker Cfi1/Net1-mScarlet-I (A41436, A41590, A41591 and A41592; same dataset as in Fig. 6C). Single cell traces were aligned to anaphase onset (spindle length > 3 μm) and averaged. Solid lines represent the average. Shaded areas represent 95% confidence intervals.

(**C**) Cumulative density of anaphase duration (left) and delay of nucleolar segregation (right) for cells in (B). Anaphase duration was defined as the time from anaphase onset (spindle length > 3 μm) to mitotic exit (spindle breakdown, determined as relaxation of the distance between SPBs). Delay of nucleolar segregation was defined as the time of nucleolar segregation (clear separation of two nucleolar masses) relative to anaphase onset.

A

B

Figure S15. Phosphorylation of the Cdc14 NLS delays mitotic exit, Related to Figure 6

(**A**) Profiles (left) and relative degrees (right) of nuclear release of wild-type (A41584, $n = 69$ cells) and mutant NLS_{Cdc14} reporters where all three potential Dbf2-Mob1 (MEN) target sites (A41585, $n = 59$ cells) or only two out of three sites (A41586 and A41623, $n = 77$ and 55 cells) were mutated to alanine. Cells were grown at 25°C in SC medium + 2% glucose and imaged every 5 minutes for 4 hours. Solid lines (right) represent the median.

(**B**) Distribution of anaphase duration for different *CFI1/NET1* phospho-mutants in combination with *CDC14* phospho-mutants measured using the SPB marker Spc42-eGFP (A41603, A41604, A41605, A41606 and A41607; $n = 59$, 62, 50, 67 and 70 cells, respectively). Cells were grown at 25°C in SC medium + 2% glucose and imaged every 3 minutes for 4 hours. Anaphase duration was defined as the time from anaphase onset (spindle length > 3 µm) to mitotic exit (spindle breakdown). Solid lines represent the mean. ****$P < 0.0001$; ***$P < 0.001$ by two-sided Wilcoxon rank sum test.

Table S1. Yeast strains used in this study

Strain	Relevant Genotype
A2587	*MATa, ade2-1, leu2-3, ura3, trp1-1, his3-11,15, can1-100, GAL, psi+ (wild- type W303)*
A2588	*MATalpha*
A1638	*CFI1-3MYC*
A10991	*MATa, cdc15::cdc15-as1(L99G)::URA3*
A11869	*NUD1-13MYC::KanMx6*
A32629	*MATa, dbf2::HIS3, CDC14::TAB6-1::TRP1*
A34515	*MATalpha, CDC14-eGFP::HIS3*
A39695	*MATa, ADE2, MOB1-eGFP::KanMx6, SPC42-mCherry:NatMx6, ura3::pRS306-mCherry-TUB1::URA3*
A39893	*MATa, ADE2, MOB1-eGFP::KanMX6, ura3::pADH1-NLS$_{Cdc14}$-TagRFP::URA3*
A39931	*MATa, GFP-MOB1-URA3::mob1::HIS3, CFI1-mCherry::KAN*
A39933	*MATa, GFP-mob1aa79-314-URA3::mob1::HIS3, CFI1-mCherry::KAN*
A39935	*MATa, GFP-mob1aa133-314-URA3::mob1::HIS3, CFI1-mCherry::KAN*
A40257	*MATa, ADE2, MOB1-eGFP::KanMX6, CFI1-mCherry::KAN, SPC42-mCherry:NatMx6, trp1::pTEF-NLS$_{Cdc14}$-ymiRFP670::TRP1*
A40258	*MATa, ADE2, leu2::PhyB-mCherry-SIK1::LEU2, CDC15-eGFP-PIF::NatMX6*
A40260	*MATa, ADE2, leu2::PhyB-mCherry-SIK1::LEU2, MOB1-eGFP-PIF::NatMX6*
A40262	*MATa, ADE2, leu2::PhyB-mCherry-SIK1::LEU2, DBF2-eGFP-PIF::NatMX6*
A40346	*MATa, ADE2, leu2::PhyB-mCherry-Spc72::LEU2, MOB1-eGFP-PIF::NatMX6*
A40354	*MATa, ADE2, leu2::PhyB-mCherry-Tom7::LEU2, MOB1-eGFP-PIF::NatMX6*
A40903	*MATa, cdc5::cdc5-as1(L158G)*
A41211	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN*
A41212	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN*
A41213	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN*
A41214	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, cdc15::cdc15-as1(L99G)::URA3*
A41215	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, cdc15::cdc15-as1(L99G)::URA3*
A41216	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, cdc15::cdc15-as1(L99G)::URA3*
A41334	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, cdc5::cdc5-as1 (L158G)*
A41335	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, cdc5::cdc5-as1 (L158G)*
A41336	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, cdc5::cdc5-as1 (L158G)*
A41337	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, ura3::GAL-CDC15(aa1-750)-3HA::URA3, cdc5::cdc5-as1 (L158G)*

A41338	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, ura3::GAL-CDC15(aa1-750)-3HA::URA3, cdc5::cdc5-as1 (L158G)*
A41339	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, ura3::GAL-CDC15(aa1-750)-3HA::URA3, cdc5::cdc5-as1 (L158G)*
A41340	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, FOB1-2xmScarlet-I::LEU2, GAL-3HA-CFI1::URA3 (YIplac211)*
A41341	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, FOB1-2xmScarlet-I::LEU2, GAL-3HA-CFI1::URA3 (YIplac211)*
A41342	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, FOB1-2xmScarlet-I::LEU2, GAL-3HA-CFI1::URA3 (YIplac211)*
A41343	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, FOB1-2xmScarlet-I::LEU2*
A41344	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, FOB1-2xmScarlet-I::LEU2*
A41345	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, FOB1-2xmScarlet-I::LEU2*
A41346	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, FOB1-2xmScarlet-I::LEU2, cfi1::URA3*
A41347	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, FOB1-2xmScarlet-I::LEU2, cfi1::URA3*
A41348	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, FOB1-2xmScarlet-I::LEU2, cfi1::URA3*
A41349	*MATalpha, ADE2, crm1::KAN, pDC-crm1T539C(LEU2/CEN), MOB1-eGFP::KanMX6, ura3::pADH1-NLS$_{Cdc14}$-TagRFP::URA3*
A41350	*MATa, ADE2, mob1::pGPD-yeGFP-MOB1::NatMX6*
A41351	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1*
A41352	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1*
A41353	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1*
A41354	*MATalpha, ADE2, mob1::mob1aa79-314::hphNT1*
A41355	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, cdc14-3*
A41356	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, cdc14-3*
A41357	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, slk19::HIS3*
A41358	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, slk19::HIS3*
A41359	*MATa, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, slk19::HIS3*
A41360	*MATa, ADE2, leu2::PhyB-mCherry-SIK1::LEU2, SPC42-mCherry:NatMx6, MOB1-eGFP-PIF::NatMX6*
A41361	*MATa, leu2::PhyB-mCherry-SIK1::LEU2, SPC42-mCherry:NatMx6, MOB1-eGFP-PIF::NatMX6, cdc15-2*
A41362	*MATa/alpha, CFI1/cfi1::KanMx6*
A41363	*MATa, ADE2, leu2::GAL-MOB1-eGFP::LEU2, trp1::GAL-DBF2::TRP1*

A41364	*MATa, ADE2, leu2::GAL-MOB1-eGFP::LEU2*
A41365	*MATa, leu2::PhyB-mCherry-SIK1::LEU2, SPC42-mCherry:NatMx6, mob1::Mob1aa79-314-eGFP-PIF::NatMX6::hphNT1, cdc15-2*
A41366	*MATa, ADE2, leu2::PhyB-mCherry-SIK1::LEU2, SPC42-mCherry:NatMx6, mob1::mob1aa79-314-eGFP-PIF::NatMX6::hphNT1*
A41367	*MATalpha, MOB1-TurboID-V5::NatMx6*
A41368	*MATalpha, TEM1-TurboID-V5::NatMx6*
A41369	*MATalpha, CDC15-TurboID-V5::NatMx6*
A41370	*MATalpha, DBF2-TurboID-V5::NatMx6*
A41371	*MATalpha, ADE2, mob1::mob1aa79-314-TurboID-V5::NatMx6::hphNT1*
A41372	*MATa, CFI1-3MYC, mob1::Mob1-TurboID-V5::NatMx6*
A41373	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, crm1::KAN, pDC-crm1T539C(LEU2/CEN)*
A41374	*MATalpha, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, crm1::KAN, pDC-crm1T539C(LEU2/CEN)*
A41375	*MATalpha, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, crm1::KAN, pDC-crm1T539C(LEU2/CEN)*
A41376	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, ura3::GAL-CDC15(aa1-750)-3HA::URA3*
A41377	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, ura3::GAL-CDC15(aa1-750)-3HA::URA3*
A41378	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, ura3::GAL-CDC15(aa1-750)-3HA::URA3*
A41379	*MATa, MOB1-TurboID-V5::NatMx6*
A41380	*MATa, NUD1-13MYC::KanMX6, MOB1-TurboID-V5::NatMx6*
A41381	*MATa, NUD1-13MYC::KanMX6, TEM1-TurboID-V5::NatMx6*
A41382	*MATa, NUD1-13MYC::KanMX6, CDC15-TurboID-V5::NatMx6*
A41383	*MATa, trp1::GAL-DBF2-eGFP::His3MX6::TRP1, crm1::KAN, pDC-crm1T539C(LEU2/CEN)*
A41384	*MATa, trp1::GAL-DBF2-eGFP::His3MX6::TRP1*
A41385	*MATalpha, CDC5-TurboID-V5::NatMx6*
A41386	*MATa, trp1::GAL-dbf2-HyA-eGFP::TRP1*
A41387	*MATa, cfi1::KanMx6, leu2::CFI1-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41388	*MATa, trp1::GAL-eGFP::TRP1, ura3::pRS306-pCTS1-2xmCherry-SV40NLS::URA3*
A41389	*MATa, trp1::GAL-DBF2-eGFP::TRP1, ura3::pRS306-pCTS1-2xmCherry-SV40NLS::URA3*
A41390	*MATa, trp1::GAL-dbf2-L12M-eGFP::TRP1, ura3::pRS306-pCTS1-2xmCherry-SV40NLS::URA3*
A41391	*MATa, trp1::GAL-dbf2-L12A-eGFP::TRP1, ura3::pRS306-pCTS1-2xmCherry-SV40NLS::URA3*
A41392	*MATa, trp1::GAL-dbf2aa1-23-eGFP::TRP1, ura3::pRS306-pCTS1-2xmCherry-SV40NLS::URA3*

A41393	*MATa, trp1::GAL-dbf2-12del-eGFP::TRP1, ura3::pRS306-pCTS1-2xmCherry-SV40NLS::URA3*
A41394	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::dbf2-L12A*
A41395	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::dbf2-L12A*
A41396	*MATalpha, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::dbf2-L12A*
A41397	*MATa, cfi1::KanMx6, leu2::cfi1-91A(z3&4)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41398	*MATa, cfi1::KanMx6, leu2::cfi1-91A(z1)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41399	*MATa, cfi1::KanMx6, leu2::cfi1-91A(z2)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41400	*MATa, cfi1::KanMx6, leu2::cfi1-91A(z3)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41401	*MATa, cfi1::KanMx6, leu2::cfi1-91A(z4)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41402	*MATa, cfi1::KanMx6, leu2::cfi1-91A(z2&4)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41403	*MATa, cfi1::KanMx6, leu2::cfi1-91A(z1&3)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41404	*MATa, cfi1::KanMx6, leu2::cfi1-91A-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41405	*MATa, cfi1::KanMx6, leu2::cfi1-91A(z1&2)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41406	*MATa, CFI1-3MYC, TEM1-TurboID-V5::NatMx6*
A41407	*MATa, CFI1-3MYC, CDC15-TurboID-V5::NatMx6*
A41408	*MATa, cfi1::KanMx6, leu2::CFI1-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3, cdc15::cdc15-as1(L99G)::URA3*
A41409	*MATacfi1::KanMx6, leu2::CFI1-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3, cdc5::cdc5-as1(L158G)*
A41410	*MATa, cfi1::KanMx6, leu2::CFI1-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3, slk19::HIS3*
A41411	*MATa, ADE2, cfi1::KanMx6, leu2::CFI1-mScarlet-I::LEU2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1*
A41412	*MATa, ADE2, cfi1::KanMx6, leu2::cfi1-91A-mScarlet-I::LEU2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1*
A41413	*MATa, ADE2, cfi1::KanMx6, leu2::cfi1-91A(z1&3)-mScarlet-I::LEU2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1*
A41418	*MATa, CFI1-3MYC, CDC5-TurboID-V5::NatMx6*
A41420	*MATa, cfi1::KanMx6, leu2::cfi1-6Cdk-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CDC14-eGFP::HIS3*
A41424	*MATa, cdc15-2, mob1::eGFP-MOB1::hphNT1*
A41425	*MATa, cdc15-2, mob1::eGFP-mob1aa79-314::hphNT1*
A41426	*MATa, cdc15-2, mob1::eGFP-mob1aa133-314::hphNT1*

A41427   *MATa, cdc15-2, mob1::PGPD-yeGFP-MOB1::NatMX6*

A41428   *MATa, tem1-3, mob1::pGPD-yeGFP-MOB1::NatMX6*

A41429   *MATa, tem1-3, mob1::eGFP-MOB1::hphNT1*

A41430   *MATa, tem1-3, mob1::eGFP-mob1aa79-314::hphNT1*

A41431   *MATa, tem1-3, mob1::eGFP-mob1aa133-314::hphNT1*

A41432   *MATa, cdc5-1, mob1::eGFP-MOB1::hphNT1*

A41433   *MATa, cdc5-1, mob1::eGFP-mob1aa79-314::hphNT1*

A41434   *MATa, cdc5-1, mob1::eGFP-mob1aa133-314::hphNT1*

A41435   *MATa, cdc5-1, mob1::pGPD-yeGFP-MOB1::NatMX6*

A41436   *MATa, cfi1::KanMx6, leu2::CFI1-mScarlet-I::LEU2, SPC42-GFP::TRP1*

A41440   *MATa, trp1::GAL-dbf2(S17A,S20A)-eGFP::TRP1, ura3::pRS306-pCTS1-2xmCherry-SV40NLS::URA3*

A41441   *MATa, trp1::GAL-dbf2(S17D,S20D)-eGFP::TRP1, ura3::pRS306-pCTS1-2xmCherry-SV40NLS::URA3*

A41442   *MATa, trp1::GAL-dbf2(S17E,S20E)-eGFP::TRP1, ura3::pRS306-pCTS1-2xmCherry-SV40NLS::URA3*

A41584   *MATalpha, ADE2, ura3::pADH1-NLS$_{Cdc14}$-GFP::URA3, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1*

A41585   *MATalpha, ADE2, ura3::pADH1-NLS$_{Cdc14}$(S531A,S537A,S546A)-GFP::URA3, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1*

A41586   *MATalpha, ADE2, ura3::pADH1-NLS$_{Cdc14}$(S531A,S537A)-GFP::URA3, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1*

A41587   *MATa, cfi1::KanMx6, leu2::cfi1-Cdc15(z2)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670, CDC14-eGFP::HIS3*

A41588   *MATa, cfi1::KanMx6, leu2::cfi1-Cdc5only(z2)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670, CDC14-eGFP::HIS3*

A41589   *MATa, cfi1::KanMx6, leu2::cfi1-Cdc15&5(z2)-mScarlet-I::LEU2, SPC42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670, CDC14-eGFP::HIS3*

A41590   *MATa, cfi1::KanMx6, leu2::cfi1-Cdc15(z2)-mScarlet-I::LEU2, SPC42-GFP::TRP1*

A41591   *MATa, cfi1::KanMx6, leu2::cfi1-Cdc5only(z2)-mScarlet-I::LEU2, SPC42-GFP::TRP1*

A41592   *MATa, cfi1::KanMx6, leu2::cfi1-Cdc15&5(z2)-mScarlet-I::LEU2, SPC42-GFP::TRP1*

A41593   *MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, cfi1::KanMx6, leu2::cfi1-Cdc15(z2)-mScarlet-I::LEU2*

A41594   *MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, cfi1::KanMx6, leu2::cfi1-Cdc5only(z2)-mScarlet-I::LEU2*

A41595   *MATa, ADE2, mob1::pGPD-yeGFP-MOB1::NatMX6, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN*

A41596   *MATa, cfi1::KanMx6, leu2::CFI1-13MYC::TRP1::LEU2*

A41597   *MATa, cfi1::KanMx6, leu2:: cfi1-Cdc15(z2)-13MYC::TRP1::LEU2*

A41598   *MATa, cfi1::KanMx6, leu2:: cfi1-Cdc5only(z2)-13MYC::TRP1::LEU2*

A41603   *MATa, cfi1::KanMx6, leu2::CFI1-mScarlet-I::LEU2, SPC42-GFP::TRP1, CDC14-eGFP::HIS3*

A41604   *MATa, cfi1::KanMx6, leu2::CFI1-mScarlet-I::LEU2, SPC42-GFP::TRP1, cdc14::cdc14(S531A,S537A,S546A)-eGFP::HIS3*

A41605   *MATa, cfi1::KanMx6, leu2::cfi1-Cdc15(z2)-mScarlet-I::LEU2, SPC42-GFP::TRP1, cdc14::cdc14(S531A,S537A,S546A)-eGFP::HIS3*

A41606	*MATa, cfi1::KanMx6, leu2::cfi1-Cdc5only(z2)-mScarlet-I::LEU2, SPC42-GFP::TRP1, cdc14::cdc14(S531A,S537A,S546A)-eGFP::HIS3*
A41607	*MATa, cfi1::KanMx6, leu2::cfi1-Cdc15&5(z2)-mScarlet-I::LEU2, SPC42-GFP::TRP1, cdc14::cdc14(S531A,S537A,S546A)-eGFP::HIS3*
A41608	*MATa, leu2::PhyB-mCherry-SIK1::LEU2, MOB1-eGFP-PIF::NatMX6, SPC42-mCherry:NatMx6, dbf2::3MYC-DBF2::URA3*
A41609	*MATa, leu2::PhyB-mCherry-SIK1::LEU2, MOB1-eGFP-PIF::NatMX6, SPC42-mCherry:NatMx6, dbf2::3MYC-dbf2(S17A,S20A)::URA3*
A41610	*MATa, leu2::PhyB-mCherry-SIK1::LEU2, MOB1-eGFP-PIF::NatMX6, SPC42-mCherry:NatMx6, dbf2::3MYC-dbf2(S17D,S20D)::URA3*
A41611	*MATa, cfi1::KanMx6, leu2::CFI1-13MYC::TRP1::LEU2, MOB1-TurboID-V5::NatMx6*
A41612	*MATa, cfi1::KanMx6, leu2::cfi1-Cdc15(z2)-13MYC::TRP1::LEU2, MOB1-TurboID-V5::NatMx6*
A41613	*MATa, cfi1::KanMx6, leu2::cfi1-Cdc5only(z2)-13MYC::TRP1::LEU2, MOB1-TurboID-V5::NatMx6*
A41614	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::3MYC-DBF2::URA3*
A41615	*MATa, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::3MYC-DBF2::URA3*
A41616	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::3MYC-DBF2::URA3*
A41617	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::3MYC-dbf2(S17A,S20A)::URA3*
A41618	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::3MYC-dbf2(S17A,S20A)::URA3*
A41619	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::3MYC-dbf2(S17A,S20A)::URA3*
A41620	*MATa, ADE2, mob1::eGFP-MOB1::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::3MYC-dbf2(S17D,S20D)::URA3*
A41621	*MATa, ADE2, mob1::eGFP-mob1aa79-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::3MYC-dbf2(S17D,S20D)::URA3*
A41622	*MATa, ADE2, mob1::eGFP-mob1aa133-314::hphNT1, SPC42-mCherry:NatMx6, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1, CFI1-mCherry::KAN, dbf2::3MYC-dbf2(S17D,S20D)::URA3*
A41623	*MATalpha, ADE2, ura3::pADH1-NLS$_{Cdc14}$(S537A,S546A)-GFP::URA3, Spc42-mScarlet-I::LEU2, trp1::pTEF1-NLS$_{Cdc14}$-ymiRFP670::TRP1*
A41624	*MATalpha, cdc15-2, dbf2::3MYC-DBF2::URA3*
A41625	*MATalpha, cdc15-2, dbf2::3MYC-dbf2(S17A,S20A)::URA3*
A41626	*MATalpha, cdc15-2, dbf2::3MYC-dbf2(S17D,S20D)::URA3*
A41645	*MATa, cfi1::KanMx6, leu2::CFI1-mScarlet-I::LEU2, slk19::HIS3, pRS316-Cfi1*
A41646	*MATa, cfi1::KanMx6, leu2::cfi1-Cdc15(z2)-mScarlet-I::LEU2, slk19::HIS3, pRS316-CFI1*
A41647	*MATa, cfi1::KanMx6, leu2::cfi1-Cdc5only(z2)-mScarlet-I::LEU2, slk19::HIS3, pRS316-CFI1*

A41648	*MATa, cfi1::KanMx6, leu2::cfi1-Cdc15&Cdc5(z2)-mScarlet-I::LEU2, slk19::HIS3, pRS316-CFI1*
A41649	*MATa, cfi1::KanMx6, leu2::cfi1-6Cdk-mScarlet-I::LEU2, slk19::HIS3, pRS316-CFI1*
A41650	*MATa, cfi1::KanMx6, leu2::CFI1-mScarlet-I::LEU2, spo12::HIS3, pRS316-CFI1*
A41651	*MATa, cfi1::KanMx6, leu2::cfi1-Cdc15(z2)-mScarlet-I::LEU2, spo12::HIS3, pRS316-CFI1*
A41652	*MATa, cfi1::KanMx6, leu2::cfi1-Cdc5only(z2)-mScarlet-I::LEU2, spo12::HIS3, pRS316-CFI1*
A41653	*MATa, cfi1::KanMx6, leu2::cfi1-Cdc15&Cdc5(z2)-mScarlet-I::LEU2, spo12::HIS3, pRS316-CFI1*
A41654	*MATa, cfi1::KanMx6, leu2::cfi1-6Cdk-mScarlet-I::LEU2, spo12::HIS3, pRS316-CFI1*

Table S2. Plasmids used in this study

Plasmid	Description	Source
pA2721	linker-mCitrine-PIF-NatMx6	C. Tang (Addgene #51576)
pA2723	PhyB-mCherry-Spc72	C. Tang (Addgene #51582)
pA2724	PhyB-mCherry-Tom7	O. Weiner (Addgene #66571)
pA2808	PhyB-mCherry-SIK1	C. Tang (Addgene #51577)
pA2727	p404TEF1	N. Buchler & F. Cross (Addgene #15972)
pA2725	Gal-NLS$_{Cdc14}$-GFP, URA3	R. Deshaies
pA2735	p406-pADH1-NLS$_{Cdc14}$-GFP	This study
pA2786	p404-pTEF1-NLS$_{Cdc14}$-ymiRFP670	This study
pA2821	linker-eGFP-PIF-NatMx6	This study
pA2621	pRS305H-Mob1	E. Schiebel
pA2824	pRS305H-Mob1aa79-314	This study
pA2828	pRS305H-eGFP-Mob1	This study
pA2829	pRS305H-eGFP-Mob1aa79-314	This study
pA2830	pRS305H-eGFP-Mob1aa133-314	This study
pA2840	YIplac211-GAL-Cdc15(1-750)	This study
pA2852	YIplac128-Gal-Mob1-eGFP	This study
pA2854	YIplac211-Gal-3HA-Cfi1	This study
pA2847	pRS415-TurboID-V5	A. Ting (Addgene #107167)
pA2858	pRS316-Cfi1	This study
pA2859	linker-TurboID-V5-NatMx6	This study
pA2868	pFA6a-Vinnylinker-mScarlet-I-cgLEU2	E. Unal
pA2869	pFA6a-Vinnylinker-2xmScarlet-I-cgLEU2	E. Unal
pA2890	YIplac204-Gal-eGFP	This study
pA2891	YIplac204-Gal-Dbf2-eGFP	This study
pA2892	YIplac204-Gal-Dbf2-HyA-eGFP	This study
pA2908	YIplac204-Gal-dbf2-L12M-eGFP	This study
pA2926	YIplac204-Gal-dbf2-L12A-eGFP	This study
pA2927	YIplac204-Gal-dbf2aa1-23-eGFP	This study
pA2928	YIplac204-Gal-dbf2-12del-eGFP	This study
pA2941	YIplac204-Gal-dbf2-S17,20A-eGFP	This study
pA2942	YIplac204-Gal-dbf2-S17,20D-eGFP	This study
pA2943	YIplac204-Gal-dbf2-S17,20E-eGFP	This study
pA2898	pRS305s-Cfi1-mScarlet	This study
pA2911	bRA90 pPGK1-Cas9-LEU	J. Haber
pA2948	pFA6a-3myc-Dbf2-CaURA3	This study
pA2953	pFA6a-3myc-dbf2-S17,20A-CaURA3	This study
pA2954	pFA6a-3myc-dbf2-S17,20D-CaURA3	This study
pA2956	pRS305s-Cfi1-91A-mScarlet	This study
pA2957	pRS305s-Cfi1-91A(z1)-mScarlet	This study
pA2958	pRS305s-Cfi1-91A(z2)-mScarlet	This study
pA2959	pRS305s-Cfi1-91A(z3)-mScarlet	This study
pA2960	pRS305s-Cfi1-91A(z4)-mScarlet	This study
pA2961	pRS305s-Cfi1-91A(z1&2)-mScarlet	This study
pA2962	pRS305s-Cfi1-91A(z3&4)-mScarlet	This study
pA2963	pRS305s-Cfi1-91A(z1&3)-mScarlet	This study
pA2964	pRS305s-Cfi1-91A(z2&4)-mScarlet	This study
pA2965	pRS305s-Cfi1-Cdc15(z2)-mScarlet	This study
pA2966	pRS305s-Cfi1-Cdc5only(z2)-mScarlet	This study
pA2967	pRS305s-Cfi1-Cdc15&5(z2)-mScarlet	This study
pA2968	pRS305s-Cfi1-6Cdk-mScarlet	This study
pA2969	p406-pADH1-NLS$_{Cdc14}$(S531,537A)-GFP	This study
pA2970	p406-pADH1-NLS$_{Cdc14}$(S537,546A)-GFP	This study
pA2971	p406-pADH1-NLS$_{Cdc14}$(S531,537,546A)-GFP	This study

Table S3. Summary of TurboID labeling experiments

Proteins identified in TurboID labeling experiments for MEN proteins (Figure 2A, S3). Proteins listed in Figure 2A are highlighted in green.

Bait (TurboID tagged)	Hits	#Total/Unique peptides in tagged sample	#Total/Unique peptides in untagged control
Mob1	Net1/Cfi1	84/37	0
	Mob1	12/6	0
	Dbf20	8/5	0
	Nud1	5/4	0
	Dbf2/Dbf20	3/2	0
	Ctr9	3/3	0
	Rpn11	3/3	0
	Bni4	3/3	0
	Vac7	3/3	0
	Yck2	2/2	0
	Ski2	2/2	0
	Ski3	2/2	0
	Bud3	2/2	0
	Hog1	2/2	0
	Cnm67	2/2	0
	Sxm1	2/2	0
	Gsh2	2/2	0
	Aim21	2/2	0
	Sac6	2/2	0
	Alg9	2/2	0
	Ufd2	2/2	0
	Arp8	2/2	0
	Nup157	2/2	0
	Sec16	2/2	0
	Tif35	2/2	0
	Gin4	8/8	1
	Dbf2	5/3	1
	Scp160	2/2	1
Mob1Δ78	Net1/Cfi1	70/38	0
	Mob1	8/3	0
	Dbf20	7/5	0
	Dbf2/Dbf20	6/3	0
	Dhh1	3/3	0
	Pre8	3/2	0
	Ndi1	3/3	0
	Rsm23	2/2	0
	Ski2	2/2	0
	Bud14	2/2	0
	Npa3	2/2	0
	Hmf1	2/2	0
	Frt1	2/2	0
	YMR027W	2/2	0
	Tps2	2/2	0

Dbf2	Net1/Cfi1	45/29	0
	Spa2	5/5	0
	Mob1	4/4	0
	Ski2	4/4	0
	Bud14	4/3	0
	Ssm4	3/3	0
	Pop2	3/2	0
	Nud1	2/2	0
	Dbf2/Dbf20	2/2	0
	Aim21	2/1	0
	Sec3	2/2	0
	Utp6	2/1	0
	Tan1	2/2	0
	Map1	2/1	0
	Apl1	2/2	0
	Rrp6	2/2	0
	Hxt1	2/2	0
	Nup2	2/2	0
	Dbf2	7/6	1
Tem1	Prs1	5/5	0
	Cdc28	5/4	0
	Ski2	4/4	0
	Rpn11	4/4	0
	Hog1	4/4	0
	Mtr4	4/4	0
	Bfa1	4/4	0
	Utp6	3/2	0
	Paa1	3/2	0
	Ski3	3/3	0
	Gsh2	3/3	0
	Yck1	3/3	0
	Atp2	3/3	0
	Tyr1	3/2	0
	Ndi1	3/3	0
	Cat2	3/3	0
	Nup133	3/3	0
	Utp9	3/3	0
	Ipi3	3/3	0
	YMR027W	3/3	0
	Ssm4	2/2	0
	Nud1	2/2	0
	Sec3	2/2	0
	Map1	2/2	0
	Rho5	2/2	0
	Tma108	2/2	0
	Dhh1	2/2	0
	Pre10	2/2	0
	Msc7	2/2	0
	Lsm12	2/2	0

	Ald4	2/2	0
	Mot1	2/2	0
	Vma6	2/2	0
	Nop13	2/2	0
	Tps1	2/2	0
	Tim44	2/2	0
	Trm3	2/2	0
	Cms1	2/2	0
	Mmn1	2/2	0
	Bdh1	2/2	0
	YGR283C	2/2	0
	Scp160	5/5	1
	Gin4	3/3	1
	Dbf2	2/2	1
Cdc15	Prs1	8/5	0
	Gsh2	5/3	0
	Utp6	4/2	0
	Paa1	4/2	0
	Nup133	4/3	0
	Vma6	4/3	0
	Msn5	4/4	0
	Mtr4	3/2	0
	Ndi1	3/3	0
	Utp9	3/3	0
	Mcm3	3/2	0
	Urk1	3/3	0
	Rrp45	3/2	0
	Arp8	3/3	0
	Syp1	3/3	0
	Cdc28	2/2	0
	Ski2	2/2	0
	Hog1	2/2	0
	Yck1	2/2	0
	Tyr1	2/2	0
	YMR027W	2/2	0
	Nud1	2/2	0
	Msc7	2/2	0
	Tps1	2/2	0
	Tim44	2/2	0
	Yck2	2/2	0
	Rvs167	2/2	0
	Cct5	2/2	0
	Mcm5	2/2	0
	Snu114	2/2	0
	Dbp7	2/2	0
	Nup82	2/2	0
	Cnm67	2/2	0
	Dop1	2/2	0
	Doa1	2/2	0

	Pho91	2/2	0
	Scp160	9/7	1
	Dbf2	4/3	1
Cdc5	Net1/Cfi1	164/63	6/6
	Boi2	21/14	0
	Mps3	20/13	0
	YPR174C/Csa1	17/7	0
	Pfy1	16/4	0
	Prs1	11/6	0
	Ycp4	10/6	0
	Mtr4	9/9	0
	Nip7	8/5	0
	Arf2	7/2	0
	YMR144W	7/5	0
	Gsh2	7/7	0
	Rpn13	6/3	0
	Rpb9	6/3	0
	Rbp8	5/3	0
	Cdc11	5/3	0
	Nic96	5/3	0
	Vtc1	5/3	0
	Ipi3	5/4	0
	Tof2	5/4	0
	Utp6	5/3	0
	Dtd1	5/4	0

Table S4. Phosphorylation of Cfi1/Net1

Summary of phosphorylation sites in Cfi1/Net1. SGD data was retrieved on 2019-02-06, the numbers represent number of studies reported for each site. CDK sites were extracted from Holt et al. 2009 (denoted as +); green highlight, sites mutated to alanine in *cfi1/net1-6CDK* (Azzam et al. 2004). PP, phospho-proteomics with DIA-MS; ana, anaphase culture; meta, metaphase culture; +, detected or determined as *CDC15/CDC5*-dependent; w, weakly dependent (see methods for criteria); *, fit Dbf2's phosphorylation motif of RXXS; ^, phosphorylated by Cdc5 *in vitro* (Chen et al. 2002); yellow highlight, mutated to alanine in *cfi1-Cdc15(z2)*; purple highlight, mutated to alanine in *cfi1-Cdc5only(z2)*; A, mutated to alanine in *cfi1-91A*; red highlight, mutated to alanine in *cfi1-Cdc15&5(z2)*.

Sites	SGD	CDK	IP-MS (ana)	PP (ana)	PP (meta)	Cdc15 (ana)	Cdc5 (ana)	Cdc5 (meta)	*cfi1-91A*
S31			+						A
S43	1			+			+		A
S48	1		+	+	+		+^		A
Y51					+				
S56				+					A
S60	1		+	+	+		+^	+	A
T62		(green)			+				
S64	1		+	+			+^		A
S69	1				+				A
S160	1		+						A
S166	4	+ (green)	+	+	+				A
S169			+	+					A
S178	2			+		+	+		A
S179	1								A
S180	1								A
S207	1								A
T212	2	(green)	+	+		+	+		A
S228	3			+	+				A
S231	5		+	+	+				A
S242	1			+	+		+^	+	A
T248				+	+				A
S252	4	+ (green)	+	+	+				A
S259	2		+	+		+*	+		A
S269	1		+	+		+	+		A
S270	1		+	+		+	+		A
S278				+		+			A
T282			+						A
T288			+						A
S295	2		+	+	+	+*	+		A

T297	2	+	+	+	+				A
S301				+					A
T304	1		+	+	+				A
T311			+	+	+				A
S317				+					
T356				+					A
T357			+	+	+				A
S362	2		+	+		+*	+		A
S384	2			+	+		+		A
S385	1		+	+	+				A
S388				+	+				A
S412	1								A
S415	1								A
S433	1								A
S437	3			+	+		+	+	A
S439	3		+	+	+	+*	+		A
S440	2		+	+	+		+		A
S447	2	+	+	+	+				A
S452	1	+	+	+	+				A
S496			+		+				A
S497	5		+	+	+	+*	+		A
T500				+					A
S511			+						A
S557				+	+				A
S558				+	+				A
S561			+						A
S567	1								A
T584				+					A
T602				+	+				A
T603				+	+		+	+	A
S611	3		+	+	+				A
S613	2				+				A
S614				+					A
S615	2		+		+				A
T676	4	+	+	+	+				A
S679	1								A
S734	1			+	+		+		A

S744	1		+	+	+		+	+	A
S747	1			+			+	+	A
S782			+		+				A
S785	1								A
T787					+				
S803	1				+				A
S823	1								A
S830	2	+							A
T838	1								A
T839			+						A
S840	2		+	+	+				A
T1017	2		+	+	+				A
S1024	1								A
S1025			+						A
S1026	3		+	+		+*	+		A
S1032	5	+	+	+	+				A
S1034	2								A
T1042	3	+	+	+	+				A
T1049	3		+	+					A
S1055	2		+	+	+				A
S1056	2	+	+	+	+			+	A
S1059	2			+	+				A
S1066	2		+	+	+				A
S1069				+					A
S1070			+		+				A
S1082	5		+	+	+				A
S1084	2		+						A
S1085	2		+						A
S1091				+					A
S1166	1								A

Supporting databases (Excel spreadsheets)

Database S1. Summary of phospho-proteomics results for inhibiting Cdc15 and Cdc5 in anaphase cells and Cdc5 in metaphase cells.

Database S2. Complete list of phosphopeptides for each phosphorylation site detected in the DIA-MS experiments.